

A discrepancy version of the Hajnal-Szemerédi theorem

Balogh, Jozsef; Csaba, Bela; Pluhar, Andras; Treglown, Andrew

DOI:

[10.1017/S0963548320000516](https://doi.org/10.1017/S0963548320000516)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Balogh, J, Csaba, B, Pluhar, A & Treglown, A 2021, 'A discrepancy version of the Hajnal-Szemerédi theorem', *Combinatorics, Probability and Computing*, vol. 30, no. 3, pp. 444-459.
<https://doi.org/10.1017/S0963548320000516>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

A DISCREPANCY VERSION OF THE HAJNAL–SZEMERÉDI THEOREM

JÓZSEF BALOGH, BÉLA CSABA, ANDRÁS PLUHÁR AND ANDREW TREGLOWN

ABSTRACT. A perfect K_r -tiling in a graph G is a collection of vertex-disjoint copies of the clique K_r in G covering every vertex of G . The famous Hajnal–Szemerédi theorem determines the minimum degree threshold for forcing a perfect K_r -tiling in a graph G . The notion of discrepancy appears in many branches of mathematics. In the graph setting, one assigns the edges of a graph G labels from $\{-1, 1\}$, and one seeks substructures F of G that have ‘high’ discrepancy (i.e. the sum of the labels of the edges in F is far from 0). In this paper we determine the minimum degree threshold for a graph to contain a perfect K_r -tiling of high discrepancy.

1. INTRODUCTION

1.1. Discrepancy of graphs. Classical discrepancy theory, or the study of irregularities of distribution, concerns with the following question: given some space, how evenly can one distribute a set of n points in it (where here evenness is measured with respect to certain subsets)? Perhaps the first significant result in the area is by Hermann Weyl on the criterion for a sequence to be uniformly distributed in the unit interval. In the other direction, answering a question by van der Corput, van Aardenne-Ehrenfest proved that some irregularity of a point sequence in the unit interval is inevitable. Since then discrepancy theory has become a widely studied area, with lots of ramifications and applications in ergodic theory, number theory, statistics, geometry, computer science, etc. For more details see the monograph by Beck and Chen [4], the book by Chazelle [7] and the book chapter by Alexander and Beck [1].

In this paper we study the discrepancy of graphs; a topic that lies in the wider framework of *hypergraph discrepancy theory* (see e.g. [3, 7]). Before we can rigorously discuss this topic we must introduce some definitions.

Definition 1.1. *Suppose G is a graph and $f : E(G) \rightarrow \{-1, 1\}$. We say a subgraph G' of a graph G has discrepancy t (with respect to f) if $\sum_{e \in E(G')} f(e) = t$ and absolute discrepancy t (with respect to f) if $\left| \sum_{e \in E(G')} f(e) \right| = t$.*

If G and G' are n -vertex graphs, then we say that G contains a copy of G' of *high discrepancy* (with respect to f) if there is a copy of G' in G with absolute discrepancy $\Omega(n)$. Note that this concept also has a natural rephrasing in terms of Ramsey theory: given any 2-colouring of the edges of G , one seeks a copy of G' in G whose edge set contains significantly more edges from one colour class than the other.

JB: Department of Mathematics, University of Illinois at Urbana–Champaign, IL, USA, and Moscow Institute of Physics and Technology, Dolgoprudny, Russian Federation, jobal@illinois.edu. Research of this author is partially supported by NSF Grants DMS-1500121, DMS-1764123, Arnold O. Beckman Research Award (UIUC) Campus Research Board 18132 and the Langan Scholar Fund (UIUC).

BC: Bolyai Institute, University of Szeged, Hungary, bcsaba@math.u-szeged.hu. Research of this author was supported in part by the grant TUDFO/47138-1/2019-ITM of the Ministry for Innovation and Technology, Hungary, and by NKFIH grant KH.18 129597.

AP: Department of Computer Science, University of Szeged, Hungary, pluhar@inf.u-szeged.hu.

AT: University of Birmingham, United Kingdom, a.c.treglown@bham.ac.uk.

A natural question in graph discrepancy is to seek a fixed spanning subgraph H of a graph G of high discrepancy (or at least discrepancy ‘far’ away from zero). The first result of this type was obtained by Erdős, Füredi, Loeb and Sós [10]: they proved that, for some constant $c > 0$, given any labelling $f : E(K_n) \rightarrow \{-1, 1\}$ of K_n and any fixed spanning tree T_n with maximum degree Δ , K_n contains a copy of T_n of absolute discrepancy at least $c(n - 1 - \Delta)$. Note that in [10] this result was phrased in the equivalent Ramsey setting.

In a previous paper [3], Jing and the first three authors of this paper investigated the graph discrepancy problem of spanning trees, paths and Hamilton cycles for various classes of graphs G . For example, the following result determines the minimum degree threshold for forcing a Hamilton cycle of high discrepancy.

Theorem 1.2 (Balogh, Csaba, Jing and Pluhár [3]). *Let $0 < c < 1/4$ and $n \in \mathbb{N}$ be sufficiently large. If G is an n -vertex graph with*

$$\delta(G) \geq (3/4 + c)n$$

and $f : E(G) \rightarrow \{-1, 1\}$, then there is a Hamilton cycle in G with absolute discrepancy at least $cn/32$ (with respect to f). Moreover, if 4 divides n , there is an n -vertex graph with $\delta(G) = 3n/4$ and an edge labelling $f : E(G) \rightarrow \{-1, 1\}$ for which every Hamilton cycle has discrepancy 0 (with respect to f).

One can view such results about discrepancy as a measure of how robustly a graph contains a spanning structure. Indeed, Theorem 1.2 implies that every n -vertex graph G with $\delta(G) > (3/4 + o(1))n$ contains a Hamilton cycle that spans an ‘unbalanced’ collection of edges for any partition $A \cup B$ of $E(G)$. (See [22] for a survey on other measures of graph robustness.)

After submitting this paper, a multicolour extension of Theorem 1.2 was proven where the underlying graph is the random graph, see [11].

1.2. Perfect tilings in graphs. An H -tiling in a graph G is a collection of vertex-disjoint copies of H contained in G . An H -tiling is *perfect* if it covers all the vertices of G . Perfect H -tilings are also often referred to as H -factors, *perfect H -packings* or *perfect H -matchings*. H -tilings can be viewed as generalisations of both the notion of a matching (which corresponds to the case when H is a single edge) and the Turán problem (i.e. a copy of H in G is simply an H -tiling of size one).

Except for the case when H contains no component of size at least 3, the decision problem of whether a graph contains a perfect H -tiling is NP-complete (see [13]). Thus, there has been substantial efforts to obtain sufficient conditions that force a graph to contain a perfect H -tiling. In particular, a cornerstone result in extremal graph theory is the Hajnal–Szemerédi theorem [12], which characterises the minimum degree threshold that ensures a graph contains a perfect K_r -tiling.

Theorem 1.3 (Hajnal and Szemerédi [12]). *Every graph G whose order n is divisible by r and whose minimum degree satisfies $\delta(G) \geq (1 - 1/r)n$ contains a perfect K_r -tiling. Moreover, there are n -vertex graphs G with $\delta(G) = (1 - 1/r)n - 1$ that do not contain a perfect K_r -tiling.*

There has also been much interest in the minimum degree threshold that ensures a perfect H -tiling for an arbitrary graph H . After earlier work on this topic (see e.g. [2, 18]), Kühn and Osthus [20, 21] determined, up to an additive constant, the minimum degree that forces a perfect H -tiling for *any* fixed graph H . Furthermore, there are now many different generalisations of the Hajnal–Szemerédi theorem. In particular, Kierstead and Kostochka [15] proved an *Ore-type* analogue, Keevash and Mycroft [14] proved a version for r -partite graphs, whilst there are now several generalisations of Theorem 1.3 in the setting of *directed graphs* (see e.g. [8, 9]).

1.3. Our main result. In this paper we prove the following discrepancy version of the Hajnal–Szemerédi theorem.

Theorem 1.4. *Suppose $r \geq 3$ is an integer and let $\eta > 0$. Then there exists $n_0 \in \mathbb{N}$ and $\gamma > 0$ such that the following holds. Let G be a graph on $n \geq n_0$ vertices where r divides n and where*

$$\delta(G) \geq \left(1 - \frac{1}{r+1} + \eta\right)n.$$

Given any function $f : E(G) \rightarrow \{-1, 1\}$ there exists a perfect K_r -tiling \mathcal{T} in G so that

$$\left| \sum_{e \in E(\mathcal{T})} f(e) \right| \geq \gamma n.$$

Comparing Theorem 1.4 with Theorem 1.3 we see that having minimum degree just above that which forces a perfect K_{r+1} -tiling in fact ensures a perfect K_r -tiling of high discrepancy. Moreover, the minimum degree condition in Theorem 1.4 is essentially best-possible for all values of $r \geq 3$. Interestingly, whilst the underlying extremal graph is the same for all $r \geq 3$ (the $(r+1)$ -partite Turán graph), the precise labelling of the edges we take is rather different depending on the value of r modulo 4. In Section 3 we construct extremal labellings in the cases when $r \equiv 1, 2 \pmod{4}$. In the case when $r \equiv 0, 3 \pmod{4}$ the extremal labelling is easy to describe: let K be the complete graph K_{r+1} with precisely half of its edges labelled with 1, the remaining edges with -1 (the choice of r ensures this is possible). Then for any $n \in \mathbb{N}$ divisible by $r(r+1)$ consider the blow-up G of K with $n/(r+1)$ vertices in each class, and where the labellings of each edge in G are induced by the labelling of $E(K)$. It is easy to see that every perfect K_r -tiling in G has discrepancy precisely 0, whilst $\delta(G) = (1 - 1/(r+1))n$. Moreover, in the case when $r(r+1)$ does not divide n , the same construction G is such that every perfect K_r -tiling has absolute discrepancy $O_r(1)$.

Note that the $r = 2$ case (i.e. perfect matchings) is covered by Theorem 1.2. Indeed, it is easy to see that since the hypothesis of Theorem 1.2 forces a Hamilton cycle of high discrepancy, this ensures a perfect matching of high discrepancy. Moreover, consider the 4-partite Turán graph G on n vertices (where 4 divides n). Label all edges incident to one of the vertex classes of G with -1 . All remaining edges are labelled 1. Then every perfect matching in G has discrepancy 0. Thus, perhaps surprisingly, this observation and Theorem 1.4 imply that the minimum degree threshold for forcing a perfect K_3 -tiling of high discrepancy is the same as the analogous threshold for perfect matchings.

The paper is organised as follows. In the next section we introduce some notation and definitions. In Section 3 we give the extremal examples for Theorem 1.4 in the cases when $r \equiv 1, 2 \pmod{4}$. We introduce a number of tools that will be used in the proof of Theorem 1.4 in Section 4. In Section 5 we give an outline of the proof of Theorem 1.4 before giving the full proof in Section 6. Finally, in Section 7 we present a number of open questions.

2. NOTATION AND DEFINITIONS

Let G be a graph. We write $V(G)$ for the vertex set of G , $E(G)$ for the edge set of G and define $|G| := |V(G)|$ and $e(G) := |E(G)|$. Given a subset $X \subseteq V(G)$, we write $G[X]$ for the subgraph of G induced by X and $G \setminus X$ for the subgraph of G induced by $V(G) \setminus X$. The degree of x is denoted by $d_G(x)$ and its neighbourhood by $N_G(x)$. Given a vertex $x \in V(G)$ and a set $Y \subseteq V(G)$ we write $d_G(x, Y)$ to denote the number of edges xy where $y \in Y$. Given a subgraph F of G we write $d_G(x, F) := d_G(x, V(F))$. Given disjoint vertex classes $X, Y \subseteq V(G)$, we write $G[X, Y]$ for the bipartite graph with vertex classes X and Y whose edge set consists of all those edges in G with one endpoint in X and the other in Y ; we write $e_G(X, Y)$ for the number of edges in $G[X, Y]$.

Suppose G is a graph and $f : E(G) \rightarrow \{-1, 1\}$. We say that $e \in E(G)$ is a *1-edge* if $f(e) = 1$ and a *(-1)-edge* if $f(e) = -1$. The *(-1)-neighbourhood* $N_G^-(x)$ of a vertex $x \in V(G)$ is the set of all

vertices $y \in V(G)$ so that xy is a (-1) -edge in G ; the 1-neighbourhood $N_G^+(x)$ of a vertex $x \in V(G)$ is the set of all vertices $y \in V(G)$ so that xy is a 1-edge in G .

The following notion of a K_r -template is crucial for the proof of Theorem 1.4.

Definition 2.1. *Let F be a graph. A K_r -template of F of size s is a collection $\{H_1, \dots, H_s\}$ of not necessarily distinct copies of K_r in F for which there is some $s' \in \mathbb{N}$ so that every vertex $x \in V(F)$ lies in precisely s' of the H_i . (In fact, note we must have $s' = sr/|F|$.) Suppose $f : E(F) \rightarrow \{-1, 1\}$ and $\mathcal{K} := \{H_1, \dots, H_s\}$ is a K_r -template of F . We say that \mathcal{K} has discrepancy t if*

$$\sum_{i=1}^s \sum_{e \in E(H_i)} f(e) = t.$$

The following special labelled copies of K_r appear in the proof of Theorem 1.4.

Definition 2.2. *We write K_r^+ for a copy of K_r whose edges are each assigned 1; define K_r^- to be a copy of K_r whose edges are each assigned -1 . The $(K_r, +)$ -star is a copy of K_r whose 1-edges induce a copy of $K_{1,r-1}$. We call the root of this $K_{1,r-1}$ the head of the $(K_r, +)$ -star. We define the $(K_r, -)$ -star and its head analogously.*

We write $0 < \alpha \ll \beta \ll \gamma$ to mean that we can choose the constants α, β, γ from right to left. More precisely, there are increasing functions f and g such that, given γ , whenever we choose some $\beta \leq f(\gamma)$ and $\alpha \leq g(\beta)$, all calculations needed in our proof are valid. Hierarchies of other lengths are defined in the obvious way. Throughout the paper we omit floors and ceilings whenever this does not affect the argument.

3. THE EXTREMAL EXAMPLES

After its statement, we introduced an extremal example for Theorem 1.4 in the case when $r \equiv 0, 3 \pmod{4}$. In this section we first describe an extremal example for the case when $r \equiv 1 \pmod{4}$ and then give a construction for the $r \equiv 2 \pmod{4}$ case.

Proposition 3.1. *Let $m \in \mathbb{N}$, $r := 4m + 1$ and $n \in \mathbb{N}$ be divisible by $2r(r + 1)$. Let G be the complete balanced $(r + 1)$ -partite graph on n vertices (and so $\delta(G) = (1 - 1/(r + 1))n$). There is a function $f : E(G) \rightarrow \{-1, 1\}$ so that for every perfect K_r -tiling \mathcal{T} in G , \mathcal{T} has discrepancy zero (i.e. $\sum_{e \in E(\mathcal{T})} f(e) = 0$).*

Proof. Let V_1, \dots, V_{r+1} denote the vertex classes of G ; so $|V_i| = n/(r + 1)$ for all $i \in [r + 1]$. Consider a copy K of K_r on vertex set $[r]$. Since $r = 4m + 1$ we can assign labels from $\{-1, 1\}$ to each edge of K so that the (-1) -edges induce a spanning $2m$ -regular subgraph of K ; the 1-edges induce a spanning $2m$ -regular subgraph of K . Let X, Y be a partition of V_{r+1} so that $|X| = |Y|$.

We now define $f : E(G) \rightarrow \{-1, 1\}$ as follows. The labelling of K induces a labelling of the edges in $G' := G \setminus V_{r+1}$. That is, if $xy \in E(G)$ and $x \in V_i, y \in V_j$ where $1 \leq i < j \leq r$, then $f(xy) = 1$ if ij is a 1-edge in K ; $f(xy) = -1$ if ij is a (-1) -edge in K . Every vertex in X sends 1-edges to each vertex in $V(G')$; every vertex in Y sends (-1) -edges to each vertex in $V(G')$.

There are precisely three types of copy of K_r in G : *Type 1* K_r have every vertex in $V(G')$; *Type 2* K_r have one vertex in X , the remaining vertices in $V(G')$; *Type 3* K_r have one vertex in Y , the remaining vertices in $V(G')$. Note that a type 1 copy of K_r has discrepancy 0, a type 2 copy of K_r has discrepancy $r - 1$, and a type 3 copy of K_r has discrepancy $-r + 1$. Given any perfect K_r -tiling \mathcal{T} in G , \mathcal{T} must contain precisely the same number of type 2 and type 3 copies of K_r . Thus, \mathcal{T} has discrepancy 0, as desired. \square

A similar function $f : E(G) \rightarrow \{-1, 1\}$ to that in Proposition 3.1 now resolves the case when $r \equiv 2 \pmod{4}$.

Proposition 3.2. *Let $m \in \mathbb{N}$, $r := 4m + 2$ and $n \in \mathbb{N}$ be divisible by $2r(r + 1)$. Let G be the complete balanced $(r + 1)$ -partite graph on n vertices (and so $\delta(G) = (1 - 1/(r + 1))n$). There is a function $f : E(G) \rightarrow \{-1, 1\}$ so that for every perfect K_r -tiling \mathcal{T} in G , \mathcal{T} has discrepancy zero (i.e. $\sum_{e \in E(\mathcal{T})} f(e) = 0$).*

Proof. Let V_1, \dots, V_{r+1} denote the vertex classes of G ; so $|V_i| = n/(r + 1)$ for all $i \in [r + 1]$. Consider a copy K of K_r on vertex set $[r]$ whose edges are assigned labels from $\{-1, 1\}$ so that there is precisely one more 1-edge than (-1) -edge. Let X, Y be a partition of V_{r+1} so that $|X| = \frac{(r-1)n}{2r(r+1)}$ and $|Y| = \frac{n}{2r}$.

We now define $f : E(G) \rightarrow \{-1, 1\}$ as follows. As in Proposition 3.1, the labelling of K induces a labelling of the edges in $G' := G \setminus V_{r+1}$. Every vertex in X sends 1-edges to each vertex in $V(G')$; every vertex in Y sends (-1) -edges to each vertex in $V(G')$.

As before, there are precisely three types of copy of K_r in G : *Type 1* K_r have every vertex in $V(G')$; *Type 2* K_r have one vertex in X , the remaining vertices in $V(G')$; *Type 3* K_r have one vertex in Y , the remaining vertices in $V(G')$. Consider any perfect K_r -tiling \mathcal{T} in G . Our aim is to show that \mathcal{T} has discrepancy 0.

Note that \mathcal{T} contains precisely $\frac{n}{r} - \frac{n}{r+1} = \frac{n}{r(r+1)}$ copies of K_r of type 1; each of these K_r s has discrepancy 1. Consider the subtiling \mathcal{T}' of \mathcal{T} induced by the type 2 and type 3 copies of K_r . Let \mathcal{T}'' be the K_{r-1} -tiling in G' obtained from \mathcal{T}' by removing all those vertices from $V_{r+1} = X \cup Y$. Note that \mathcal{T}'' covers precisely $\frac{n}{r+1} - \frac{n}{r(r+1)} = \frac{(r-1)n}{r(r+1)}$ vertices in V_i for each $i \in [r]$. In total \mathcal{T}'' consists of $n/(r + 1)$ copies of K_{r-1} . Moreover, for each pair (i, j) with $1 \leq i < j \leq r$, a precisely $\frac{r-2}{r}$ -proportion of the copies of K_{r-1} in \mathcal{T}'' contain an edge xy with $x \in V_i, y \in V_j$. Together, this implies that \mathcal{T}'' has discrepancy

$$\frac{r-2}{r} \times \frac{n}{r+1}.$$

Recalling that each edge incident to X is a 1-edge and each edge incident to Y is a (-1) -edge, we conclude that \mathcal{T} has discrepancy

$$\frac{n}{r(r+1)} + \left(\frac{r-2}{r} \times \frac{n}{r+1} \right) + |X|(r-1) - |Y|(r-1) = \frac{n}{r(r+1)} + \frac{(r-2)n}{r(r+1)} - \frac{(r-1)n}{r(r+1)} = 0,$$

as required. \square

4. USEFUL RESULTS

4.1. The regularity lemma. In the proof of our main result we will use a discrepancy variant of Szemerédi's regularity lemma [23]. Before stating this result, we introduce some notation. The *density* of a bipartite graph G with vertex classes A and B is defined to be

$$d(A, B) := \frac{e(A, B)}{|A||B|}.$$

Given any $\varepsilon, d > 0$, we say that G is (ε, d) -regular if $d(A, B) \geq d$ and, for all sets $X \subseteq A$ and $Y \subseteq B$ with $|X| \geq \varepsilon|A|$ and $|Y| \geq \varepsilon|B|$, we have $|d(A, B) - d(X, Y)| < \varepsilon$. We say that G is (ε, d) -superregular if all sets $X \subseteq A$ and $Y \subseteq B$ with $|X| \geq \varepsilon|A|$ and $|Y| \geq \varepsilon|B|$ satisfy that $d(X, Y) > d$, that $d_G(a) > d|B|$ for all $a \in A$ and that $d_G(b) > d|A|$ for all $b \in B$.

Suppose G is a graph with edge labelling $f : E(G) \rightarrow \{-1, 1\}$. Given disjoint $X, Y \subseteq V(G)$ we write $G_+[X, Y]$ (or $(X, Y)_G^+$) for the bipartite graph with vertex classes X, Y whose edge set

consists of all those 1-edges between X and Y in G . We define $G_-[X, Y]$ and $(X, Y)_G^-$ analogously (now with respect to (-1) -edges).

We will apply the following variant of Szemerédi's regularity lemma that can be easily deduced from the multicoloured version e.g. given in [19].

Lemma 4.1. *For every $\varepsilon > 0$ and $\ell_0 \in \mathbb{N}$ there exists $L_0 = L_0(\varepsilon, \ell_0)$ so that the following holds. Let $d \in [0, 1]$ and G be a graph on $n \geq L_0$ vertices with edge labelling $f : E(G) \rightarrow \{-1, 1\}$. Then there exists a partition V_0, V_1, \dots, V_ℓ of $V(G)$ and a spanning subgraph G' of G , such that the following conditions hold:*

- (i) $\ell_0 \leq \ell \leq L_0$;
- (ii) $d_{G'}(x) \geq d_G(x) - (2d + \varepsilon)n$ for every $x \in V(G)$;
- (iii) the subgraph $G'[V_i]$ is empty for all $1 \leq i \leq \ell$;
- (iv) $|V_0| \leq \varepsilon n$;
- (v) $|V_1| = |V_2| = \dots = |V_\ell|$;
- (vi) for all $1 \leq i < j \leq \ell$ either $(V_i, V_j)_{G'}^+$ is an (ε, d) -regular pair or $G'_+[V_i, V_j]$ is empty;
- (vii) for all $1 \leq i < j \leq \ell$ either $(V_i, V_j)_{G'}^-$ is an (ε, d) -regular pair or $G'_-[V_i, V_j]$ is empty.

□

We call V_1, \dots, V_ℓ *clusters*, V_0 the *exceptional set* and the vertices in V_0 *exceptional vertices*. We refer to G' as the *pure graph*. The *reduced graph* R of G with parameters ε , d and ℓ_0 is the graph whose vertices are V_1, \dots, V_ℓ and in which $V_i V_j$ is an edge precisely when at least one of $(V_i, V_j)_{G'}^+$ and $(V_i, V_j)_{G'}^-$ is (ε, d) -regular. Associated with the reduced graph R is an edge labelling $f_R : E(R) \rightarrow \{-1, 1\}$ where $f_R(V_i V_j) := 1$ if $(V_i, V_j)_{G'}^+$ is (ε, d) -regular and $f_R(V_i V_j) := -1$ otherwise. (So if both $(V_i, V_j)_{G'}^+$ and $(V_i, V_j)_{G'}^-$ is (ε, d) -regular, then f_R only 'records' the former property.)

We will use the following well-known property of the reduced graph.

Fact 4.2. *Given a constant $c > 0$, let G be an n -vertex graph with $\delta(G) \geq cn$ that we have applied Lemma 4.1 to (with parameters ε , d and ℓ_0). Let R be the corresponding reduced graph. Then $\delta(R) \geq (c - 2d - 2\varepsilon)|R|$.*

□

The following well-known result allows us to use subgraphs of a reduced graph as 'templates' for embedding in the original graph G .

Lemma 4.3 (Key lemma [19]). *Suppose that $0 < \varepsilon < d$, that $q, t \in \mathbb{N}$ and that R is a graph with $V(R) = \{v_1, \dots, v_k\}$. We construct a graph G as follows: Replace every vertex $v_i \in V(R)$ with a set V_i of q vertices and replace each edge of R with an (ε, d) -regular pair. For each $v_i \in V(R)$, let U_i denote the set of t vertices in $R(t)$ corresponding to v_i . Let H be a subgraph of $R(t)$ with maximum degree Δ and set $h := |H|$. Set $\delta := d - \varepsilon$ and $\varepsilon_0 := \delta^\Delta / (2 + \Delta)$. If $\varepsilon \leq \varepsilon_0$ and $t - 1 \leq \varepsilon_0 q$ then there are at least*

$$(\varepsilon_0 q)^h \text{ labelled copies of } H \text{ in } G$$

so that if $x \in V(H)$ lies in U_i in $R(t)$, then x is embedded into V_i in G .

The following fundamental result of Komlós, Sárközy and Szemerédi [16], known as the *blow-up lemma*, essentially says that (ε, d) -superregular pairs behave like complete bipartite graphs with respect to containing bounded degree subgraphs.

Lemma 4.4 (Blow-up lemma [16]). *Given a graph F on vertices $\{1, \dots, f\}$ and $d, \Delta > 0$, there exists an $\varepsilon_0 = \varepsilon_0(d, \Delta, f) > 0$ such that the following holds. Given $L_1, \dots, L_f \in \mathbb{N}$ and $\varepsilon \leq \varepsilon_0$, let F^* be the graph obtained from F by replacing each vertex $i \in F$ with a set V_i of L_i new vertices and joining all vertices in V_i to all vertices in V_j whenever ij is an edge of F . Let G be a spanning*

subgraph of F^* such that for every edge $ij \in F$ the pair $(V_i, V_j)_G$ is (ε, d) -superregular. Then G contains a copy of every subgraph H of F^* with $\Delta(H) \leq \Delta$.

4.2. An absorbing lemma. We will apply the following well-known *absorbing lemma* (which e.g. is a special case of [24, Theorem 4.1]). Given a graph G we say a set $S \subseteq V(G)$ is a K_r -*absorbing set* for $Q \subseteq V(G)$, if both $G[S]$ and $G[S \cup Q]$ contain perfect K_r -tilings.

Lemma 4.5. *Let $0 < 1/n \ll \nu \ll \eta \ll 1/r$ where $n, r \in \mathbb{N}$ and $r \geq 2$. Suppose that G is a graph on n vertices with $\delta(G) \geq (1 - 1/r + \eta)n$. Then $V(G)$ contains a set M so that $|M| \leq \nu n$ and M is a K_r -absorbing set for every $W \subseteq V(G) \setminus M$ such that $|W| \in r\mathbb{N}$ and $|W| \leq \nu^3 n$.*

5. OVERVIEW OF THE PROOF OF THEOREM 1.4

In the proof of Theorem 1.4 we will apply the regularity lemma to obtain the reduced graph R of G with an associated edge labelling $f_R : E(R) \rightarrow \{-1, 1\}$. Since the reduced graph R ‘inherits’ the minimum degree condition on G (see Fact 4.2), the Hajnal–Szemerédi theorem implies that R contains a perfect K_{r+1} -tiling \mathcal{T} .

In Claim 6.1 we establish the following crucial property: (a) if \mathcal{T} has high absolute discrepancy (with respect to f_R), then we can use this structure in R as a framework to build a perfect K_r -tiling in G with high absolute discrepancy (with respect to f). To build this tiling in G we make use of the absorbing method.

We then establish another vital property of R : (b) if R has a ‘small’ subgraph F so that F has two K_r -templates with different discrepancies (with respect to f_R), then we can use this to again build a perfect K_r -tiling in G with high absolute discrepancy (see Claim 6.2).

We may therefore assume neither (a) nor (b) holds. This in turn forces the cliques of size at most $r + 2$ in R to have some very rigid structure. In particular, we deduce that every copy of K_{r+1} in R (therefore in our tiling \mathcal{T}) is one of the following: a K_{r+1}^+ ; a K_{r+1}^- ; a $(K_{r+1}, +)$ -star; a $(K_{r+1}, -)$ -star (see Claim 6.5).

After this, we then argue that in fact almost all of the tiles in \mathcal{T} are copies of $(K_{r+1}, +)$ -stars and $(K_{r+1}, -)$ -stars. Finally, we prove that there are two tiles K, K' in \mathcal{T} for which (b) must hold with respect to $F := R[K \cup K']$, and so we do have a perfect K_r -tiling in G with high absolute discrepancy.

6. PROOF OF THEOREM 1.4

It suffices to prove the theorem in the case when $\eta \ll 1/r$. Define additional constants $\gamma, \varepsilon, d, \nu > 0$ and $n_0, \ell_0, L_0 \in \mathbb{N}$ so that

$$(1) \quad 0 < 1/n_0 \ll \gamma \ll 1/L_0 \leq 1/\ell_0 \ll \varepsilon \ll d \ll \nu \ll \eta \ll 1/r.$$

Here L_0 is the constant obtained from Lemma 4.1 on input ε, ℓ_0 .

Let G be a graph on $n \geq n_0$ vertices as in the statement of the theorem. Fix an arbitrary edge labelling $f : E(G) \rightarrow \{-1, 1\}$.

By Lemma 4.5 we obtain a set of vertices $Abs \subseteq V(G)$ where $|Abs| \leq \nu n$ and where both $G[Abs]$ and $G[Abs \cup W]$ contain perfect K_r -tilings for any set $W \subseteq V(G) \setminus Abs$ of size at most $\nu^3 n$ where r divides $|W|$. Let $G_1 := G \setminus Abs$. Thus,

$$(2) \quad \delta(G_1) \geq \left(1 - \frac{1}{r+1} + \frac{3\eta}{4}\right)n.$$

6.1. Applying the regularity lemma. Apply the regularity lemma (Lemma 4.1) to G_1 with parameters ε, d, ℓ_0 . We thus obtain clusters V_1, \dots, V_ℓ of size m (where $\ell_0 \leq \ell \leq L_0$) an exceptional set V_0 (of size at most εn) and a pure graph G'_1 of G_1 . We may assume that $r+1$ divides ℓ . (If not, we can achieve this by deleting at most r of the clusters, and move the vertices in these clusters to the exceptional set V_0 .) Further we obtain the reduced graph R of G_1 with an edge labelling $f_R : E(R) \rightarrow \{-1, 1\}$ ‘inherited’ from f (as defined in Section 4.1). Note that (2) and Fact 4.2 imply that

$$(3) \quad \delta(R) \geq \left(1 - \frac{1}{r+1} + \frac{\eta}{2}\right) \ell.$$

The following two claims will be used several times in our proof. The first implies that to obtain our desired perfect K_r -tiling in G it suffices to find a perfect K_{r+1} -tiling in R of high absolute discrepancy.

Claim 6.1. *Suppose that R contains a perfect K_{r+1} -tiling \mathcal{T}_R with absolute discrepancy $t \geq \eta^2 \ell$ (with respect to f_R). Then G contains a perfect K_r -tiling with absolute discrepancy at least γn (with respect to f).*

Proof. Consider any copy H of K_{r+1} in \mathcal{T}_R . Suppose that H has discrepancy $t_H \in \mathbb{Z}$ (with respect to f_R). The vertices W_1, \dots, W_{r+1} in H are clusters in G . Write G_H for the $(r+1)$ -partite graph $G'_1[W_1 \cup \dots \cup W_{r+1}]$. Through repeated applications of the key lemma (Lemma 4.3) we obtain that there is a K_r -tiling \mathcal{T}_H in G_H so that:

- (i) All but precisely $\varepsilon^{1/2}m$ vertices in W_i are covered by \mathcal{T}_H for each $i \in [r+1]$;
- (ii) Given any edge $xy \in E(\mathcal{T}_H)$, if $x \in W_i$ and $y \in W_j$ then $f(xy) = f_R(W_i W_j)$;
- (iii) Each copy of K_r in \mathcal{T}_H contains at most one vertex from every cluster W_i . Furthermore, given any $1 \leq i < j \leq r+1$, a $\frac{r-1}{r+1}$ -proportion of the K_r in \mathcal{T}_H contain an edge from $G'_1[W_i, W_j]$.

Note that (ii) follows from the definition of f_R ; (iii) simply states that we embed copies of K_r in G_H in a balanced way, alternating which cluster W_i is ‘uncovered by a copy of K_r ’. Since H has discrepancy t_H , (ii) and (iii) imply that \mathcal{T}_H has discrepancy

$$\frac{r-1}{r+1} \times |\mathcal{T}_H| \times t_H = \frac{(r-1)}{r} (1 - \varepsilon^{1/2}) m t_H$$

(with respect to f).

Consider the K_r -tiling \mathcal{T}' in G'_1 obtained by taking the union of the \mathcal{T}_H for each H in \mathcal{T}_R . By (i), \mathcal{T}' contains all but $|V_0| + \varepsilon^{1/2}m\ell \leq 2\varepsilon^{1/2}n$ of the vertices in G_1 . Noting that $\sum_{H \in \mathcal{T}_R} t_H \in \{t, -t\}$, we deduce that \mathcal{T}' has absolute discrepancy

$$\frac{(r-1)}{r} (1 - \varepsilon^{1/2}) m t \geq \frac{2}{3} (1 - \varepsilon^{1/2}) \eta^2 m \ell \geq \eta^2 n / 2$$

(with respect to f). Let W be the set of vertices in G_1 uncovered by \mathcal{T}' ; so $|W| \leq 2\varepsilon^{1/2}n \leq \nu^3 n$. Thus, $G[Abs \cup W]$ has a perfect K_r -tiling \mathcal{T}'' . As $|Abs \cup W| \leq 2\nu n$, $\mathcal{T}' \cup \mathcal{T}''$ is a perfect K_r -tiling in G with absolute discrepancy at least $\eta^2 n / 2 - \binom{r}{2} 2\nu n \geq \gamma n$, as desired. \square

The next claim gives us a useful condition that guarantees our desired perfect K_r -tiling in G ; it will be used repeatedly through the proof.

Claim 6.2. *Let F be a subgraph of R on p vertices where $r+1 \leq p \leq 2r+2$. Given some $s \leq r^{100}$, suppose that F has two K_r -templates $\mathcal{K} = \{H_1, \dots, H_s\}$ and $\mathcal{K}' = \{H'_1, \dots, H'_s\}$, both of size s . If \mathcal{K} and \mathcal{K}' have different discrepancies (with respect to f_R), then G contains a perfect K_r -tiling with absolute discrepancy at least γn .*

Proof. Let W_1, \dots, W_p denote the clusters of G'_1 that correspond to the vertices of F . So if $W_i W_j \in E(F)$ and $f_R(W_i W_j) = 1$ then $(W_i, W_j)_{G'_1}^+$ is (ε, d) -regular; otherwise if $W_i W_j \in E(F)$ and $f_R(W_i W_j) = -1$ then $(W_i, W_j)_{G'_1}^-$ is (ε, d) -regular. A well-known property of regular pairs implies that we can delete $\varepsilon^{1/2}m$ vertices from each of these clusters to obtain subclusters W'_1, \dots, W'_p with the following properties: if $W_i W_j \in E(F)$ and $f_R(W_i W_j) = 1$ then $(W'_i, W'_j)_{G'_1}^+$ is $(2\varepsilon, d/2)$ -superregular; if $W_i W_j \in E(F)$ and $f_R(W_i W_j) = -1$ then $(W'_i, W'_j)_{G'_1}^-$ is $(2\varepsilon, d/2)$ -superregular. Write $m' := (1 - \varepsilon^{1/2})m$; so $|W'_i| = m'$ for all $i \in [p]$.

Let F^* be the p -partite graph with vertex classes W'_1, \dots, W'_p , and where for each $i \neq j$, there are all possible edges between W'_i and W'_j precisely if $W_i W_j \in E(F)$; that is, F^* is a blow-up of F . Define $f_{F^*} : E(F^*) \rightarrow \{-1, 1\}$ so that $f_{F^*}(xy) = 1$ if $x \in W'_i, y \in W'_j$ and $f_R(W_i W_j) = 1$; $f_{F^*}(xy) = -1$ if $x \in W'_i, y \in W'_j$ and $f_R(W_i W_j) = -1$.

Write t for the discrepancy of \mathcal{K} and t' for the discrepancy of \mathcal{K}' ; by the assumption in the claim, $t \neq t'$. Note that we can use \mathcal{K} as a ‘framework’ to find a perfect K_r -tiling \mathcal{T} in F^* as follows: consider any H_k in \mathcal{K} and let W_{i_1}, \dots, W_{i_r} be the vertices of H_k ; in \mathcal{T} there are $m'p/sr$ copies of K_r corresponding to H_k which contain precisely one vertex from each of $W'_{i_1}, \dots, W'_{i_r}$. Thus, \mathcal{T} has discrepancy $m'pt/sr$ (with respect to f_{F^*}).

Similarly, we can use \mathcal{K}' as a framework to find a perfect K_r -tiling \mathcal{T}' in F^* of discrepancy $m'pt'/sr$ (with respect to f_{F^*}).

Now applying the blow-up lemma, this ensures $G_0 := G'_1[W'_1 \cup \dots \cup W'_p]$ contains two perfect K_r -tilings \mathcal{T}_1 and \mathcal{T}_2 with discrepancy $m'pt/sr$ and $m'pt'/sr$ respectively (with respect to f). Note that

$$|m'pt/sr - m'pt'/sr| \geq (1 - \varepsilon^{1/2}) \frac{m}{s} \geq \frac{n}{2L_0 r^{100}} \stackrel{(1)}{\geq} 2\gamma n.$$

Further, $G \setminus G_0$ comfortably satisfies

$$\delta(G \setminus G_0) \geq (1 - 1/r)n,$$

so contains a perfect K_r -tiling \mathcal{T}_3 by the Hajnal–Szemerédi theorem. Therefore, both $\mathcal{T}_1 \cup \mathcal{T}_3$ and $\mathcal{T}_2 \cup \mathcal{T}_3$ are perfect K_r -tilings in G , whose discrepancies differ by at least $2\gamma n$; thus, one of these perfect K_r -tilings has absolute discrepancy at least γn , as desired. \square

From now on we may assume that the hypotheses of Claims 6.1 and 6.2 fail; this will eventually lead to a contradiction, thereby proving the theorem.

6.2. Properties of cliques in R . The minimum degree condition on R ensures the following easy observation.

Claim 6.3. *Let $1 \leq k \leq r + 1$. Every copy of K_k in R lies in a copy of K_{r+2} .*

We now use Claim 6.2 to prove that the copies of K_{r+2} in R have a limited number of possible edge labellings.

Claim 6.4. *Every copy K of K_{r+2} in R is one of the following: a K_{r+2}^+ ; a K_{r+2}^- ; a $(K_{r+2}, +)$ -star; a $(K_{r+2}, -)$ -star.*

Proof. Consider an arbitrary Hamilton cycle C in K . We obtain a K_r -template \mathcal{K}_C of K of size $r + 2$ by going around the Hamilton cycle as follows: take each copy of K_r whose vertices are r consecutive vertices along C and add it to \mathcal{K}_C .

Consider any two Hamilton cycles $C = W_1 \dots W_i W_{i+1} W_{i+2} W_{i+3} \dots W_{r+2}$ and C' obtained from C by reordering $W_i W_{i+1} W_{i+2} W_{i+3}$ as $W_i W_{i+2} W_{i+1} W_{i+3}$ (i.e. we just swap the order of W_{i+1} and

W_{i+2}). Since we are assuming the hypothesis of Claim 6.2 does not hold, we must have that \mathcal{K}_C and $\mathcal{K}_{C'}$ have the same discrepancy with respect to f_R .

This implies that $f_R(W_i W_{i+1}) + f_R(W_{i+2} W_{i+3}) = f_R(W_i W_{i+2}) + f_R(W_{i+1} W_{i+3})$. (The left hand side considers the contribution to the discrepancy of \mathcal{K}_C not ‘present’ in the discrepancy of $\mathcal{K}_{C'}$; the right hand side considers the contribution to the discrepancy of $\mathcal{K}_{C'}$ not ‘present’ in the discrepancy of \mathcal{K}_C .)

The choice of the Hamilton cycle C in K was arbitrary. So this implies that

$$(4) \quad f_R(ab) + f_R(cd) = f_R(ac) + f_R(bd) \text{ for all distinct } a, b, c, d \in V(K).$$

Consider any $a \in V(K)$. Suppose $|N_K^-(a)| \geq 3$. Given any distinct $b, c, d \in N_K^-(a)$, (4) implies that $f_R(bd) = f_R(cd)$. This implies that the edges in $N_K^-(a)$ are either all 1-edges or all (-1) -edges. A similar argument holds if $|N_K^+(a)| \geq 3$.

In particular, this implies that if one of $N_K^-(a)$ and $N_K^+(a)$ is empty then K is one of the following: a K_{r+2}^+ ; a K_{r+2}^- ; a $(K_{r+2}, +)$ -star; a $(K_{r+2}, -)$ -star. We may therefore assume that both $N_K^-(a)$ and $N_K^+(a)$ are non-empty, and without loss of generality assume that $|N_K^+(a)| \geq 2$.

Choose any distinct $c, d \in N_K^+(a)$ and $b \in N_K^-(a)$. Noting that ac is a 1-edge and ab is a (-1) -edge, (4) implies cd is a 1-edge and bd is a (-1) -edge. The choice of $c, d \in N_K^+(a)$ and $b \in N_K^-(a)$ was arbitrary so this implies all edges between $N_K^+(a)$ and $N_K^-(a)$ are (-1) -edges.

If $|N_K^-(a)| = 1$ we are immediately done now: indeed, we have just argued that $b \in N_K^-(a)$ sends out (-1) -edges to everything else; and as $|N_K^+(a)| \geq 3$ in this case, all edges in $N_K^+(a)$ are $+1$ -edges. That is, K is a $(K_{r+2}, -)$ -star.

Thus, we now may additionally assume $|N_K^-(a)| \geq 2$. Choose any distinct $c', d' \in N_K^-(a)$ and $b' \in N_K^+(a)$. Then (4) implies that $b'd'$ is a 1-edge. This is a contradiction, as we already proved that all edges between $N_K^+(a)$ and $N_K^-(a)$ are (-1) -edges. Thus this case does not occur, and we are done. \square

Combining Claims 6.3 and 6.4 we obtain the following.

Claim 6.5. *Let $1 \leq k \leq r + 2$. Every copy of K_k in R is one of the following: a K_k^+ ; a K_k^- ; a $(K_k, +)$ -star; a $(K_k, -)$ -star.*

6.3. Using a perfect K_{r+1} -tiling in R . Note that (3) and Theorem 1.3 imply that R contains a perfect K_{r+1} -tiling \mathcal{T} . By Claim 6.5, there are only four types of K_{r+1} in \mathcal{T} . Let A denote the set of K_{r+1}^+ in \mathcal{T} ; let B denote the set of K_{r+1}^- in \mathcal{T} ; let C denote the set of $(K_{r+1}, +)$ -stars in \mathcal{T} ; let D denote the set of $(K_{r+1}, -)$ -stars in \mathcal{T} . Without loss of generality we may assume that

$$(5) \quad |B| + |C| \geq |A| + |D|.$$

6.3.1. *Assume that A is non-empty.*

Claim 6.6. *Consider any vertex $V_a \in V(A)$ and any copy $K \in B$ of K_{r+1}^- . Then we may assume $d_R(V_a, K) \leq r - 2$ if r is even; $d_R(V_a, K) \leq r - 1$ if r is odd.*

Proof. Write K_A for the clique in \mathcal{T} that contains V_a . Let $F := R[K_A \cup K]$.

First consider the case when r is even, and suppose V_a sends $r - 1$ edges to K in R . Suppose i of these edges are 1-edges (and so $r - 1 - i$ of them are (-1) -edges). Let $X, Y \in V(K)$ be the vertices in K that are not incident to one of these $r - 1$ edges. We will prove that F satisfies the hypothesis of Claim 6.2.

Write \mathcal{K}_A for the set of all copies of K_r in K_A , and \mathcal{K} for the set of all copies of K_r in K ; so $|\mathcal{K}| = |\mathcal{K}_A| = r + 1$.

Define \mathcal{K}_1 to be the K_r -template for F of size $2r(r + 1)$ that contains precisely r copies of each of the cliques in $\mathcal{K}_A \cup \mathcal{K}$. Note that indeed \mathcal{K}_1 is a K_r -template for F as each vertex $V \in V(F)$ is

contained in precisely r^2 of the cliques in \mathcal{K}_1 . Since $K_A \in A$, $K \in B$, and \mathcal{K}_1 contains the same number of copies of cliques from \mathcal{K}_A and \mathcal{K} , \mathcal{K}_1 has discrepancy 0 (with respect to f_R).

We define another K_r -template \mathcal{K}_2 for F of size $2r(r+1)$ as follows:

- (i) for the clique $H \in \mathcal{K}_A$ that does not contain V_A , add $2r-1$ copies of H to \mathcal{K}_2 ;
- (ii) for each clique $H \in \mathcal{K}_A$ that contains V_A , add $r-1$ copies of H to \mathcal{K}_2 ;
- (iii) add to \mathcal{K}_2 r copies of the clique in F that contains V_A and the $r-1$ vertices in $V(K) \setminus \{X, Y\}$;
- (iv) add $r+1$ copies of each clique $H \in \mathcal{K}$ that contains both X and Y ;
- (v) add one copy of each clique $H \in \mathcal{K}$ that avoids one of X and Y .

To prove that \mathcal{K}_2 is a K_r -template for F of size $2r(r+1)$ it suffices to prove that every vertex $V \in V(F)$ lies in precisely r^2 of the cliques in \mathcal{K}_2 : if $V \in V(K_A) \setminus \{V_A\}$ then (i) and (ii) give that V lies in $(2r-1) + (r-1)(r-1) = r^2$ such cliques; (ii) and (iii) imply V_A lies in $(r-1)r + r = r^2$ such cliques; if $V \in V(K) \setminus \{X, Y\}$ then (iii)–(v) imply that V lies in $r + (r+1)(r-2) + 1 \cdot 2 = r^2$ such cliques; if $V \in \{X, Y\}$ then (iv) and (v) imply that V lies in $(r+1)(r-1) + 1 \cdot 1 = r^2$ such cliques.

To compute the discrepancy of \mathcal{K}_2 note that, compared to \mathcal{K}_1 it has: one fewer clique from \mathcal{K}_A ; $r-1$ fewer cliques from \mathcal{K} ; an additional r cliques (from (iii)) that each have discrepancy $2i - \binom{r}{2}$. As \mathcal{K}_1 has discrepancy 0 this implies that \mathcal{K}_2 has discrepancy

$$-\binom{r}{2} + (r-1)\binom{r}{2} + r\left(2i - \binom{r}{2}\right) = 2ir - r(r-1) \neq 0$$

as $i \neq (r-1)/2$ (recall we assumed that r is even). So F satisfies the hypothesis of Claim 6.2.

Now suppose r is odd and V_a sends at least r edges to K in R . We can fix $r-1$ such edges so that $i \neq (r-1)/2$ of them are 1-edges and $r-1-i$ of them are (-1) -edges. Now arguing precisely as before we conclude F satisfies the hypothesis of Claim 6.2 as desired. \square

Claim 6.7. *Consider any $V_a \in V(A)$ and any $K \in C$. Then we may assume $d_R(V_a, K) \leq r-2$ if r is even; $d_R(V_a, K) \leq r-1$ if r is odd.*

Proof. Write K_A for the clique in \mathcal{T} that contains V_a . Let $F := R[K_A \cup K]$. Write \mathcal{K}_A for the set of all copies of K_r in K_A , and \mathcal{K} for the set of all copies of K_r in K ; so $|\mathcal{K}| = |\mathcal{K}_A| = r+1$.

The proof proceeds similarly to the previous claim. If r is even, suppose V_a sends $r-1$ edges to K in R ; if r is odd suppose V_a sends r edges to K in R . If all these edges avoid the head¹ V_H of K then we can argue precisely as in Claim 6.6 to obtain two K_r -templates \mathcal{K}_1 and \mathcal{K}_2 of F , both with the same size, but different discrepancy. Note that how we construct \mathcal{K}_1 and \mathcal{K}_2 is identical to the proof of Claim 6.6, though the discrepancies will differ from that claim since now $K \in C$.

Next suppose r is even and V_a sends $r-1$ edges to K in R , one of the endpoints being the head V_H . Suppose i of these edges are 1-edges and $r-1-i$ of them are (-1) -edges. Let $X, Y \in V(K)$ be the vertices in K that are not endpoints of such edges. Again, we choose \mathcal{K}_1 and \mathcal{K}_2 as in Claim 6.6.

That is, we define \mathcal{K}_1 to be the K_r -template for F of size $2r(r+1)$ that contains precisely r copies of each of the cliques in $\mathcal{K}_A \cup \mathcal{K}$. We define \mathcal{K}_2 as follows:

- (i) for the clique $H \in \mathcal{K}_A$ that does not contain V_A , add $2r-1$ copies of H to \mathcal{K}_2 ;
- (ii) for each clique $H \in \mathcal{K}_A$ that contains V_A , add $r-1$ copies of H to \mathcal{K}_2 ;
- (iii) add to \mathcal{K}_2 r copies of the clique in F that contains V_A and the $r-1$ vertices in $V(K) \setminus \{X, Y\}$;
- (iv) add $r+1$ copies of each clique $H \in \mathcal{K}$ that contains both X and Y ;
- (v) add one copy of each clique $H \in \mathcal{K}$ that avoids one of X and Y .

The same argument as in Claim 6.6 implies both \mathcal{K}_1 and \mathcal{K}_2 are K_r -templates for F of size $2r(r+1)$.

To complete the proof we have to again show the discrepancies of \mathcal{K}_1 and \mathcal{K}_2 are different. Note that (i) and (ii) imply that \mathcal{K}_2 has one fewer copy of K_r^+ from \mathcal{K}_A compared to \mathcal{K}_1 ; compared to \mathcal{K}_1 ,

¹ K is a copy of a $(K_k, +)$ -star; the head of such a star was defined in Definition 2.2.

\mathcal{K}_2 has an additional r cliques arising from (iii); from (iv) and (v) we conclude that \mathcal{K}_2 has r fewer $(K_r, +)$ -stars from \mathcal{K} compared to \mathcal{K}_1 ; by (iv) \mathcal{K}_2 has one more copy of a K_r^- from \mathcal{K} compared to \mathcal{K}_1 . Thus, the difference in discrepancy between \mathcal{K}_1 and \mathcal{K}_2 is precisely

$$-\binom{r}{2} + r \left(2i + 2(r-2) - \binom{r}{2} \right) - r \left(-\binom{r}{2} + 2(r-1) \right) - \binom{r}{2} = 2ri - r^2 - r.$$

As r is even, this term is non-zero (since $i \neq (r+1)/2$ in this case). Therefore, \mathcal{K}_1 and \mathcal{K}_2 are K_r -templates for F of different discrepancies; that is, the hypothesis of Claim 6.2 holds.

Next suppose $r \geq 5$ is odd, and V_a has at least r neighbours in K , including the head V_H . We can choose $r-1$ such neighbours, including V_H , so that i of the corresponding edges incident to V_a are 1-edges (and $r-1-i$ of them are (-1) -edges), where vitally, $i \neq (r+1)/2$. In particular, here we are using that $(r+1)/2 < r-1$ to guarantee that we can choose i as desired. Then arguing as in the previous case we obtain two K_r -templates for F of different discrepancies. This argument also resolves the case when $r=3$ unless all the edges from V_a to K are 1-edges; in which case we would be forced to ‘choose’ $i=2=(r+1)/2$. However, in this case we have that V_a sends two 1-edges to vertices in $V(K) \setminus \{V_H\}$. In this case can argue precisely as in Claim 6.6 to obtain two K_r -templates for F of different discrepancies. This completes the proof of the claim. \square

By the last two claims we have that each $V_a \in V(A)$ has average degree of at most $r-1$ into each $K \in B \cup C$. Trivially V_a has average degree of at most $r+1$ into each $K \in A \cup D$. So by (5), each $V_a \in V(A)$ has average degree at most r into each $K \in A \cup B \cup C \cup D$. This is a contradiction as R has minimum degree $\delta(R) \geq (1 - 1/(r+1) + \eta/2)\ell$. Thus we conclude that A is empty.

Further this implies B is small. Indeed, if $|B| \geq \eta^2\ell$ then (5) implies that the perfect K_{r+1} -tiling \mathcal{T} of R has absolute discrepancy at least $\eta^2\ell$. Thus the hypothesis of Claim 6.1 holds, contradicting our assumption.

Therefore assume $A = \emptyset$ and $|B| \leq \eta^2\ell$. We now split into cases.

6.3.2. Case 1: $r \geq 4$. Note that in this case we have $|D| - \eta^2\ell \leq |C| \leq |D| + \eta^2\ell$. Indeed, otherwise (5) implies that the perfect K_{r+1} -tiling \mathcal{T} of R has absolute discrepancy at least $\eta^2\ell$.

Together with the fact that $\delta(R) \geq (1 - 1/(r+1) + \eta/2)\ell$, this implies that every $V_c \in V(C)$ has at least $(1 - 2/(r+1) + \eta/3)|C|$ neighbours in D . Thus, this immediately implies the following.

Claim 6.8. *Given any $V_c \in V(C)$ there is some $K \in D$ such that $d_R(V_c, K) \geq r$.* \square

Fix $V_c \in V(C)$ to be the head of some tile K_C in \mathcal{T} . So V_c sends at least $r-1$ edges to $K \setminus \{V_H\}$ where V_H is the head of K . Fix $r-1$ of these edges. Call the endpoints of these edges in K *good*. Write X for the vertex in $K \setminus \{V_H\}$ that is not good. Write \mathcal{K}_C for the set of all copies of K_r in K_C , and \mathcal{K} for the set of all copies of K_r in K ; so $|\mathcal{K}| = |\mathcal{K}_C| = r+1$.

Set $F := R[K_C \cup K]$. Define \mathcal{K}_1 to be the K_r -template for F of size $2r(r+1)$ that contains precisely r copies of each of the cliques in $\mathcal{K}_C \cup \mathcal{K}$. Note that indeed \mathcal{K}_1 is a K_r -template for F as each vertex $V \in V(F)$ is contained in precisely r^2 of the cliques in \mathcal{K}_1 .

We define another K_r -template \mathcal{K}_2 for F of size $2r(r+1)$ as follows:

- (i) for the clique $H \in \mathcal{K}_C$ that does not contain V_c , add $2r-1$ copies of H to \mathcal{K}_2 ;
- (ii) for each clique $H \in \mathcal{K}_C$ that contains V_c , add $r-1$ copies of H to \mathcal{K}_2 ;
- (iii) add to \mathcal{K}_2 r copies of the clique in F that contains V_c and the good vertices;
- (iv) for each clique $H \in \mathcal{K}$ that contains both X and V_H , add $r+1$ copies of H to \mathcal{K}_2 ;
- (v) add one copy of the clique $H \in \mathcal{K}$ that avoids X ;
- (vi) add one copy of the clique $H \in \mathcal{K}$ that avoids V_H .

To prove that \mathcal{K}_2 is a K_r -template for F of size $2r(r+1)$ it suffices to prove that every vertex $V \in V(F)$ lies in precisely r^2 of the cliques in \mathcal{K}_2 : if $V \in V(K_C) \setminus \{V_c\}$ then (i) and (ii) give that V

lies in $(2r-1) + (r-1)(r-1) = r^2$ such cliques; (ii) and (iii) imply V_c lies in $(r-1)r + r = r^2$ such cliques; if $V \in V(K) \setminus \{X, V_H\}$ then (iii)–(vi) imply that V lies in $r + (r+1)(r-2) + 1 + 1 = r^2$ such cliques; if $V = X$ then (iv) and (vi) imply that V lies in $(r+1)(r-1) + 1 = r^2$ such cliques; if $V = V_H$ then (iv) and (v) imply that V lies in $(r+1)(r-1) + 1 = r^2$ such cliques.

We will now complete this case by showing that \mathcal{K}_1 and \mathcal{K}_2 have different discrepancies with respect to f_R ; that is, the hypothesis of Claim 6.2 holds, as desired.

Write i for the number of (-1) -edges in F with one endpoint V_c , the other a good vertex. So there are $r-1-i$ 1-edges between V_c and the good vertices. Note that (i) implies that \mathcal{K}_2 has $r-1$ more copies of K_r^- from \mathcal{K}_C compared to \mathcal{K}_1 ; compared to \mathcal{K}_1 , (ii) implies that \mathcal{K}_2 has r fewer copies of $(K_r, +)$ -stars from \mathcal{K}_C ; the r cliques from (iii) are contained in \mathcal{K}_2 but not \mathcal{K}_1 ; from (iv) and (v) we conclude that \mathcal{K}_2 has the same number of $(K_r, -)$ -stars as \mathcal{K}_1 ; by (vi) \mathcal{K}_2 has $r-1$ fewer copies of K_r^+ from \mathcal{K} compared to \mathcal{K}_1 . Thus, the difference in discrepancy between \mathcal{K}_1 and \mathcal{K}_2 is precisely

$$-(r-1)\binom{r}{2} - r\left(-\binom{r}{2} + 2(r-1)\right) + r\left(\binom{r}{2} - 2i\right) - (r-1)\binom{r}{2} = -r(r-1) - 2ri < 0.$$

Therefore, \mathcal{K}_1 and \mathcal{K}_2 are K_r -templates for F of different discrepancies; that is, the hypothesis of Claim 6.2 holds, as required.

6.3.3. Case 2: $r = 3$. As $\delta(R) \geq (3/4 + \eta/2)\ell$ and $|B| \leq \eta^2\ell$ we obtain the following.

Claim 6.9. *Given any $V_c \in V(C)$ there is some $K \in C \cup D$ such that $d_R(V_c, K) = 4$.* \square

Fix $V_c \in V(C)$ to be the head of some tile K_C in \mathcal{T} . Write \mathcal{K}_C for the set of all copies of K_3 in K_C , and \mathcal{K} for the set of all copies of K_3 in K ; so $|\mathcal{K}| = |\mathcal{K}_C| = 4$. Set $F := R[K_C \cup K]$.

Subcase 2a: $K \in D$.

Note that V_c together with K forms a copy of K_5 in R . As $K \in D$, Claim 6.4 tells us that the edge between V_c and the head V_H of K is a (-1) -edge; all other edges between V_c and K are 1-edges.

Define \mathcal{K}_1 to be the K_3 -template for F of size 24 that contains precisely 3 copies of each of the cliques in $\mathcal{K}_C \cup \mathcal{K}$. Note that indeed \mathcal{K}_1 is a K_r -template for F as each vertex $V \in V(F)$ is contained in precisely 9 of the cliques in \mathcal{K}_1 .

We define another K_3 -template \mathcal{K}_2 for F of size 24 as follows:

- (i) for the clique $H \in \mathcal{K}_C$ that does not contain V_c , add 5 copies of H to \mathcal{K}_2 ;
- (ii) for each clique $H \in \mathcal{K}_C$ that contains V_c , add 2 copies of H to \mathcal{K}_2 ;
- (iii) add to \mathcal{K}_2 one copy of each clique in F that contains V_c and precisely two of the vertices in $V(K) \setminus \{V_H\}$;
- (iv) for each clique $H \in \mathcal{K}$ that contains V_H , add 3 copies of H to \mathcal{K}_2 ;
- (v) add one copy of the clique $H \in \mathcal{K}$ that avoids V_H .

It is easy to check that every $V \in V(F)$ lies in precisely 9 cliques in \mathcal{K}_2 ; so indeed \mathcal{K}_2 is K_3 -template for F of size 24. Further, \mathcal{K}_1 has discrepancy 0, \mathcal{K}_2 has discrepancy -6 . Thus, the hypothesis of Claim 6.2 holds, as desired.

Subcase 2b: $K \in C$.

Note that V_c together with K forms a copy of K_5 in R . As $K \in C$, Claim 6.4 tells us that the edge between V_c and the head V_H of K is a 1-edge; all other edges between V_c and K are (-1) -edges.

We define \mathcal{K}_1 and \mathcal{K}_2 precisely as in Subcase 2a. That is, define \mathcal{K}_1 to be the K_3 -template for F of size 24 that contains precisely 3 copies of each of the cliques in $\mathcal{K}_C \cup \mathcal{K}$. Define \mathcal{K}_2 as follows:

- (i) for the clique $H \in \mathcal{K}_C$ that does not contain V_c , add 5 copies of H to \mathcal{K}_2 ;
- (ii) for each clique $H \in \mathcal{K}_C$ that contains V_c , add 2 copies of H to \mathcal{K}_2 ;

- (iii) add to \mathcal{K}_2 one copy of each clique in F that contains V_c and precisely two of the vertices in $V(K) \setminus \{V_H\}$;
- (iv) for each clique $H \in \mathcal{K}$ that contains V_H , add 3 copies of H to \mathcal{K}_2 ;
- (v) add one copy of the clique $H \in \mathcal{K}$ that avoids V_H .

In this subcase, \mathcal{K}_1 has discrepancy 0, \mathcal{K}_2 has discrepancy -12 . Thus, the hypothesis of Claim 6.2 holds, as desired. This completes the proof of Theorem 1.4. \square

7. OPEN PROBLEMS

The r th power of a Hamilton cycle C is obtained from C by adding an edge between every pair of vertices of distance at most r on C . The Pósa–Seymour conjecture states that every n -vertex graph G with minimum degree $\delta(G) \geq (1 - 1/(r+1))n$ contains the r th power of a Hamilton cycle. Komlós, Sárközy and Szemerédi [17] proved this conjecture for sufficiently large n .

It is natural to seek a discrepancy analogue of the Pósa–Seymour conjecture. We believe that the hypothesis of Theorem 1.4 additionally ensures that the host graph G contains the $(r-1)$ th power of a Hamilton cycle with high discrepancy. Furthermore, the minimum degree in such a result should be best-possible (in the same sense Theorem 1.4 is best-possible). We believe the proof of such a result can be obtained via the connecting–absorbing method, and using Theorem 1.4 as a black-box (applied to the reduced graph of the host graph G); this would be a suitable project for a strong Master’s student. Note that such a result (combined with Theorem 1.2) would show that $\delta(G) = (3/4 + o(1))n$ is the threshold for a graph G to contain *both* a Hamilton cycle of high discrepancy and the square of a Hamilton cycle of high discrepancy.

It is also natural to seek an extension of Theorem 1.4 to perfect H -tilings for *any* graph H .

Question 7.1. *Given any graph H , what is the minimum degree threshold that forces a perfect H -tiling of high discrepancy in a graph G (with respect to any edge labelling $f : E(G) \rightarrow \{-1, 1\}$)?*

A famous conjecture of Bollobás and Eldridge [5], and Catlin [6] asserts that every n -vertex graph G with $\delta(G) \geq (rn - 1)/(r + 1)$ contains every n -vertex graph H with $\Delta(H) = r$.

Question 7.2. *Given any $\eta > 0$ and $r \geq 2$, does there exist an $n_0 \in \mathbb{N}$ so that the following holds for all $n \geq n_0$? Let G, H be n -vertex graphs, and assume that*

$$\delta(G) \geq (1 - 1/(r + 2) + \eta)n,$$

where $r := \Delta(H)$. Then G contains a copy of H of high discrepancy (with respect to any edge labelling $f : E(G) \rightarrow \{-1, 1\}$).

Note that the Bollobás–Eldridge–Catlin conjecture has still not been fully resolved. So it seems extremely challenging to answer Question 7.2 in general. However, our main result (Theorem 1.4) resolves Question 7.2 in the affirmative when H is a perfect K_r -tiling. It would be interesting to resolve Question 7.2 in cases for which the Bollobás–Eldridge–Catlin conjecture is known to be true (in particular the case when $r \leq 3$).

8. ACKNOWLEDGEMENTS

The first and fourth authors are grateful to the BRIDGE strategic alliance between the University of Birmingham and the University of Illinois at Urbana-Champaign. In particular, much of the research in this paper was carried out whilst the fourth author was visiting UIUC. The authors are also grateful to the referee for their helpful and careful review.

REFERENCES

- [1] J.R. Alexander, J. Beck, and W.W.L. Chen, Geometric discrepancy theory and uniform distribution. In *Handbook of Discrete and Computational Geometry*. 1997.
- [2] N. Alon and R. Yuster, H -factors in dense graphs, *J. Combin. Theory B* **66** (1996), 269–282.
- [3] J. Balogh, B. Csaba, Y. Jing and A. Pluhár, On the discrepancies of graphs, *Electron. J. Combin.* **27** (2020) P2.12.
- [4] J. Beck and W.W.L. Chen, Irregularities of Distribution. Vol. **89** of *Cambridge Tracts in Math.*, Cambridge University Press, 1987.
- [5] B. Bollobás and S.E. Eldridge, Packing of graphs and applications to computational complexity, *J. Combin. Theory Ser. B* **25** (1978), 105–124.
- [6] P.A. Catlin, Embedding subgraphs and coloring graphs under extremal degree conditions, Ph.D. Thesis, Ohio State Univ., Columbus (1976).
- [7] B. Chazelle, *The Discrepancy Method*, Cambridge University Press, 2000.
- [8] A. Czygrinow, L. DeBiasio, H.A. Kierstead and T. Molla, An extension of the Hajnal–Szemerédi theorem to directed graphs, *Combin. Probab. Comput.* **24** (2015), 754–773.
- [9] A. Czygrinow, L. DeBiasio, T. Molla and A. Treglown, Tiling directed graphs with tournaments, *Forum Math. Sigma* **6** (2018) e2.
- [10] P. Erdős, Z. Füredi, M. Loeb and V.T. Sós, Discrepancy of Trees, *Stud Sci Math*, **30** (1995), 47–57.
- [11] L. Gishboliner, M. Krivelevich and P. Michaeli, Colour-biased Hamilton cycles in random graphs, arXiv:2007.12111.
- [12] A. Hajnal and E. Szemerédi, Proof of a conjecture of Erdős, *Combinatorial Theory and its Applications vol. II* **4** (1970), 601–623.
- [13] P. Hell and D.G. Kirkpatrick, On the complexity of general graph factor problems, *SIAM J. Computing* **12** (1983), 601–609.
- [14] P. Keevash and R. Mycroft, A multipartite Hajnal–Szemerédi theorem, *J. Combin. Theory Ser. B* **114** (2015), 187–236.
- [15] H.A. Kierstead and A.V. Kostochka, An Ore-type Theorem on Equitable Coloring, *J. Combin. Theory Ser. B* **98** (2008), 226–234.
- [16] J. Komlós, G.N. Sárközy and E. Szemerédi, Blow-up lemma, *Combinatorica* **17**(1) (1997), 109–123.
- [17] J. Komlós, G.N. Sárközy and E. Szemerédi, Proof of the Seymour conjecture for large graphs, *Ann. Combin.* **2** (1998), 43–60.
- [18] J. Komlós, G.N. Sárközy and E. Szemerédi, Proof of the Alon–Yuster conjecture, *Discrete Math.* **235** (2001), 255–269.
- [19] J. Komlós and M. Simonovits, Szemerédi’s Regularity Lemma and its applications in graph theory, *Combinatorics, Paul Erdős is Eighty* (Volume 2), Keszthely (Hungary), 1993, (D. Miklós, V.T. Sós, T. Szőnyi eds.), Bolyai Math. Stud., Budapest (1996), 295–352.
- [20] D. Kühn and D. Osthus, Critical chromatic number and the complexity of perfect packings in graphs, *17th ACM-SIAM Symposium on Discrete Algorithms* (SODA 2006), 851–859.
- [21] D. Kühn and D. Osthus, The minimum degree threshold for perfect graph packings, *Combinatorica* **29** (2009), 65–107.
- [22] B. Sudakov, Robustness of graph properties, *Surveys in Combinatorics 2017*, Cambridge University Press, 2017, 372–408.
- [23] E. Szemerédi, Regular partitions of graphs, *Problèmes Combinatoires et Théorie des Graphes Colloques Internationaux CNRS* **260** (1978), 399–401.
- [24] A. Treglown, A degree sequence Hajnal–Szemerédi theorem, *J. Combin. Theory Ser. B* **118** (2016), 13–43.