

Single-electron transfer in frustrated Lewis pair chemistry

Holtrop, Flip; Jupp, Andrew; Kooij, Bastiaan; Leest, Nicolaas P.; de Bruin, Bas; Slootweg, J. Chris

DOI:

[10.1002/anie.202009717](https://doi.org/10.1002/anie.202009717)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Holtrop, F, Jupp, A, Kooij, B, Leest, NP, de Bruin, B & Slootweg, JC 2020, 'Single-electron transfer in frustrated Lewis pair chemistry', *Angewandte Chemie (International Edition)*, vol. 59, no. 49, pp. 22210-22216.
<https://doi.org/10.1002/anie.202009717>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



Frustrated Lewis Pairs Hot Paper

How to cite:

International Edition: doi.org/10.1002/anie.202009717

German Edition: doi.org/10.1002/ange.202009717

Single-Electron Transfer in Frustrated Lewis Pair Chemistry

Flip Holtrop, Andrew R. Jupp, Bastiaan J. Kooij, Nicolaas P. van Leest, Bas de Bruin, and J. Chris Slootweg*


Abstract: Frustrated Lewis pairs (FLPs) are well known for their ability to activate small molecules. Recent reports of radical formation within such systems indicate single-electron transfer (SET) could play an important role in their chemistry. Herein, we investigate radical formation upon reacting FLP systems with dihydrogen, triphenyltin hydride, or tetrachloro-1,4-benzoquinone (TCQ) both experimentally and computationally to determine the nature of the single-electron transfer (SET) events; that is, being direct SET to $B(C_6F_5)_3$ or not. The reactions of H_2 and Ph_3SnH with archetypal P/B FLP systems do not proceed via a radical mechanism. In contrast, reaction with TCQ proceeds via SET, which is only feasible by Lewis acid coordination to the substrate. Furthermore, SET from the Lewis base to the Lewis acid–substrate adduct may be prevalent in other reported examples of radical FLP chemistry, which provides important design principles for radical main-group chemistry.

Introduction


Frustrated Lewis pairs (FLPs) combine a Lewis acidic electron-pair acceptor and a Lewis basic electron-pair donor to activate small molecules, most notably H_2 and CO_2 , granting access to fascinating main group chemistry and catalysis.^[1] It is generally accepted that the FLP components cooperatively interact with the substrate to facilitate heterolytic bond cleavage;^[2] however, recent reports suggest that radicals may play an important role too and, in some cases, provide alternative homolytic pathways.^[2b,c,3] Stephan et al. reported the detection of a weak radical signal by electron paramagnetic resonance (EPR) spectroscopy for the archetypal FLP $PMes_3/B(C_6F_5)_3$ ($Mes = 2,4,6$ -triphenylmethyl), and after switching the Lewis acid to $Al(C_6F_5)_3$ found a similar, yet much clearer, EPR signal that could be unambiguously attributed to the phosphine radical cation ($PMes_3^{\cdot+}$).^[2c] Furthermore, reaction of $PMes_3/B(C_6F_5)_3$ with

Ph_3SnH resulted in formation of $[Mes_3PH][HB(C_6F_5)_3]$ and $Ph_3SnSnPh_3$, instead of the $[Mes_3P-SnPh_3][HB(C_6F_5)_3]$ product that would be expected for heterolytic cleavage of the $Sn-H$ bond.^[2c] In the case of $PtBu_3/B(C_6F_5)_3$, for which no radicals were detected in the reaction mixture, indeed $[tBu_3P-SnPh_3][HB(C_6F_5)_3]$ was obtained, corresponding to nucleophilic substitution at tin (Scheme 1a). As the difference in products was proposed to be caused by a change in mechanism (homolytic vs. heterolytic), this led to the use of Ph_3SnH as a probe for determining the mechanistic nature of FLP reactions.^[2c,4] Furthermore, Stephan et al. determined that reacting $PMes_3/B(C_6F_5)_3$ with tetrachloro-1,4-benzoquinone


a) Stephan et al.




b) Stephan et al.




c) Melen et al.



d) Ooi et al.



e) Slootweg et al.



Scheme 1. a) Different pathways proposed by Stephan et al. for reactions of FLPs with Ph_3SnH . b) Reactivity observed by Stephan et al. for $Mes_3P/B(C_6F_5)_3$ with tetrachloro-1,4-quinone (TCQ). c) Reactivity observed by Melen et al. ($Ar^F = Ph, p-F-Ph$ or fluorene; $Ar =$ variety of aryl groups, see Ref. [6]). d) Reactivity observed by Ooi et al. utilizing catalytic $B(C_6F_5)_3$ (10 mol%) ($R = Me$ or Br); e) Light dependence for radical ion pair generation from archetypal FLP systems observed by Slootweg et al. (For P: $R = Mes$ or tBu , for N: $R = Ph$ or $p-Me-Ph$).

[*] F. Holtrop, Dr. A. R. Jupp, B. J. Kooij, N. P. van Leest, Prof. Dr. B. de Bruin, Prof. Dr. J. C. Slootweg
Van't Hoff Institute for Molecular Sciences, University of Amsterdam
PO Box 94157, 1090 GD Amsterdam (The Netherlands)
E-mail: j.c.slootweg@uva.nl

Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:
<https://doi.org/10.1002/anie.202009717>.


© 2020 The Authors. Published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution Non-Commercial NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial, and no modifications or adaptations are made.

(TCQ) leads to radical formation after observation of PMes_3^+ by EPR spectroscopy (Scheme 1 b).^[2c,5] Furthermore, Melen et al. recently reported that the $\text{PMes}_3/\text{B}(\text{C}_6\text{F}_5)_3$ pair can be utilized to facilitate C–C bond formation by coupling diarylmethyl groups to styrenes via a mechanism involving single-electron transfer (SET; Scheme 1 c).^[6] Ooi et al. also achieved C–C bond formation using an amine/ $\text{B}(\text{C}_6\text{F}_5)_3$ system to couple methylvinylketone to the amine employing catalytic amounts of borane (Scheme 1 d).^[7] They also showed that the reaction requires light and proceeds via radical species which they postulated to be the result of photo-induced SET from the amine directly to $\text{B}(\text{C}_6\text{F}_5)_3$ yielding the corresponding radical ion pair [amine $^{\cdot+}$, $\text{B}(\text{C}_6\text{F}_5)_3^{\cdot-}$]. Subsequent addition of this species to the substrate was proposed, which then led to product formation.

Recently, we demonstrated the generality of SET for FLP type donor–acceptor systems^[8,9] and showed that for common P/B FLPs ($\text{PMes}_3/\text{B}(\text{C}_6\text{F}_5)_3$ and $\text{PtBu}_3/\text{B}(\text{C}_6\text{F}_5)_3$) and analogous N/B systems visible light is required to induce SET to generate the corresponding transient radical ion pairs (Scheme 1 e).^[10] This light dependence provides an excellent probe for determining whether an FLP reaction proceeds via a radical mechanism or via concerted, polar pathways, as carrying out the reaction in the absence of light precludes the formation of the radical ion pair. The work presented herein focuses on applying this notion to investigate the reaction of archetypal FLPs with the substrates H_2 , Ph_3SnH and TCQ. In addition, we analyzed the nature of the initial single-electron transfer event that is responsible for the radical chemistry observed by Melen et al. and Ooi et al. For all cases, we determine whether the boron Lewis acid is directly involved in SET, or plays a facilitating role by enhancing the oxidizing power of the substrate.^[11]

Results and Discussion

First, we assessed the influence of light on the reaction of PMes_3 and $\text{B}(\text{C}_6\text{F}_5)_3$ with H_2 (1 atm), which is known to generate the corresponding phosphonium borate $[\text{Mes}_3\text{PH}][\text{HB}(\text{C}_6\text{F}_5)_3]$.^[12] We previously showed that this combination of donor (PMes_3) and acceptor ($\text{B}(\text{C}_6\text{F}_5)_3$) forms a violet charge-transfer complex in solution from which the corresponding radical ion pair $[\text{PMes}_3^{\cdot+}, \text{B}(\text{C}_6\text{F}_5)_3^{\cdot-}]$ is generated by irradiating this electron donor–acceptor (EDA) complex with visible light (534 nm).^[10] Thus, if formation of this radical ion pair is a significantly contributing factor in hydrogen splitting, the reaction should exhibit a change in reaction rate depending on the absence or presence of light. Comparison of reaction samples kept in the dark or irradiated (534 nm, 2.2 W LEDs; Scheme 2) whilst stirring for 2.5 hours showed near-identical conversions to the phosphonium borate $[\text{Mes}_3\text{PH}][\text{HB}(\text{C}_6\text{F}_5)_3]$, and again after 4 hours, as determined by ^{31}P -NMR spectroscopy (Supporting Information, Figures S1, S2).^[13] These data show that the reaction is not light dependent and therefore the formation of the radical ion pair does not significantly influence the reaction kinetics. This finding suggests that the photo-stationary concentration of the radical ion pair $[\text{PMes}_3^{\cdot+}, \text{B}(\text{C}_6\text{F}_5)_3^{\cdot-}]$ is too low and/or its




Scheme 2. Reactivity of $\text{PMes}_3/\text{B}(\text{C}_6\text{F}_5)_3$ with H_2 for which no light dependence was observed.

lifetime is too short to significantly influence the reaction rate. Indeed, this charge-separated state lies much higher in energy (54.4 kcal mol $^{-1}$) than the neutral donor–acceptor pair $[\text{PMes}_3, \text{B}(\text{C}_6\text{F}_5)_3]$ and undergoes rapid back-electron transfer (lifetime = 237 ps) as determined by transient absorption spectroscopy to regenerate the FLP,^[10] thus preventing build-up of a concentration of radicals large enough to influence the reaction kinetics. This leads to the conclusion that the splitting of dihydrogen by PMes_3 and $\text{B}(\text{C}_6\text{F}_5)_3$ proceeds via a two-electron, heterolytic pathway, even when the reaction is performed in ambient light.^[2c–g]

Next, we probed the reaction between $\text{PMes}_3/\text{B}(\text{C}_6\text{F}_5)_3$ and Ph_3SnH (2 equiv) to analyse whether light affects the formation of phosphonium borate $[\text{Mes}_3\text{PH}][\text{HB}(\text{C}_6\text{F}_5)_3]$ and $\text{Ph}_3\text{Sn–SnPh}_3$. We found that the reaction proceeds rapidly in both darkness and ambient light and, in both cases, within minutes full conversion to $[\text{Mes}_3\text{PH}][\text{HB}(\text{C}_6\text{F}_5)_3]$ and $\text{Ph}_3\text{Sn–SnPh}_3$ was observed by multi-nuclear NMR spectroscopy ($\delta^{31}\text{P} = -28.6$, $\delta^{11}\text{B} = -26.1$, $\delta^{119}\text{Sn} = -131.7$; Supporting Information, Figure S5–S9; Scheme 1 a). This suggests that also in this case radicals are not responsible for the reaction outcome.

But how is $[\text{Mes}_3\text{PH}][\text{HB}(\text{C}_6\text{F}_5)_3]$ formed when using Ph_3SnH instead of H_2 ? For this, changing the phosphine to PtBu_3 provided insight. Addition of 1 equiv of Ph_3SnH to $\text{PtBu}_3/\text{B}(\text{C}_6\text{F}_5)_3$ in $\text{C}_6\text{H}_5\text{Cl}$ at room temperature instantly led to heterolytic cleavage of the Sn–H bond and the formation of $[\text{tBu}_3\text{P–SnPh}_3][\text{HB}(\text{C}_6\text{F}_5)_3]$ ($\delta^{31}\text{P} = 65.8$, $^1J_{\text{P–Sn}} = 90$ Hz; Scheme 3; Supporting Information, Figure S10), supporting the observations by Stephan et al.^[2c] We noted, however, that when more Ph_3SnH (up to 2.5 equiv) was used, the reaction continued and after 20 hours $[\text{tBu}_3\text{PH}][\text{HB}(\text{C}_6\text{F}_5)_3]$ ($\delta^{31}\text{P} = 58.1$, $^1J_{\text{P–H}} = 410$ Hz; Supporting Information, Figure S12) as well as $\text{Ph}_3\text{Sn–SnPh}_3$ (Scheme 3; Supporting Information, Figure S14) was observed.^[13] We also noted that this reaction proceeds equally in the absence of light, in ambient light, or under direct irradiation of the charge-transfer band of $[\text{PtBu}_3, \text{B}(\text{C}_6\text{F}_5)_3]$ (400 nm, 2.2 W LED). These findings show that for both phosphines R_3P (R = Mes and *t*Bu) a polar, heterolytic



Scheme 3. Reactivity of $\text{PtBu}_3/\text{B}(\text{C}_6\text{F}_5)_3$ with Ph_3SnH .

mechanism is operative and that the initial product $[R_3P-SnPh_3][HB(C_6F_5)_3]$ can convert into $[R_3PH][HB(C_6F_5)_3]$ in the presence of Ph_3SnH .

To elucidate the heterolytic splitting of Ph_3SnH in more detail, we first combined it with $B(C_6F_5)_3$ in calculations and found the formation of an adduct with a bridging hydride $[Ph_3Sn-H-B(C_6F_5)_3]$ ($\Delta E = -21.3$, $\Delta G^\circ_{298K} = -1.9$ kcal mol $^{-1}$; Figure 1, left), which is analogous to the key, transient intermediate in the $B(C_6F_5)_3$ -catalyzed hydrosilylation.^[14] We also observed the $[Ph_3Sn-H-B(C_6F_5)_3]$ adduct in C_6H_5Cl solution by ^{19}F -NMR spectroscopy that shows a decrease in resonance difference between the *meta*- and *para*-fluorines ($\Delta\delta$ 18.2 to 13.7 ppm), which is indicative of a transition from a trigonal planar borane to a more tetrahedral geometry.^[15] Furthermore, ^{119}Sn -NMR spectroscopy supports this notion, the clear downfield shift indicates a more electron deficient Sn nucleus ($\delta = 165$ to 130 ppm; Supporting Information, Figure S15, S16).^[16] These observations evidence activation of the tin hydride by $B(C_6F_5)_3$, making it more susceptible to nucleophilic attack by a phosphine in an S_N2 fashion to produce the initial $[R_3P-SnPh_3][HB(C_6F_5)_3]$ species.

To investigate the subsequent reaction steps and determine the influence of the P-substituent (Mes vs. *t*Bu), we again employed computational chemistry ($\omega B97X-D/def2-TZVP$; $[HB(C_6F_5)_3]^-$ anion omitted),^[15] which highlighted a formal metathesis reaction of the $[R_3P-SnPh_3]^+$ cation with Ph_3Sn-H , reminiscent of reactions between tin hydrides and neutral stannyl phosphines.^[17] When using $PMes_3$ (Figure 2, in green), $[Mes_3P-SnPh_3]^+$ undergoes a facile reaction with Ph_3SnH to afford the bridging hydride $[Ph_3Sn-H-SnPh_3]^+$ ($\Delta E^\ddagger_{TS1} \approx 9$, $\Delta E = -4.9$ kcal mol $^{-1}$),^[15] akin to the tin hydride- $B(C_6F_5)_3$ adduct (Figure 1). Subsequent deprotonation by the liberated phosphine, which induces Sn-Sn bond formation ($\Delta E^\ddagger_{TS2} = 19.9$, $\Delta E = -13.1$ kcal mol $^{-1}$), affords $[Mes_3PH]^+$ and $Ph_3Sn-SnPh_3$. This reaction profile supports the notion that $[Mes_3P-SnPh_3][HB(C_6F_5)_3]$ is a transient, unobserved intermediate in the formation of $[Mes_3PH][HB(C_6F_5)_3]$.^[2c]




Figure 1. Computed structure for the adducts of Ph_3SnH with $B(C_6F_5)_3$ (left) and Ph_3Sn^+ (right) featuring a bridging hydride (DFT: $\omega B97X-D/def2-TZVP$). Selected bond lengths and angles: $Ph_3Sn-H-B(C_6F_5)_3$: Sn-H 1.83 Å, B-H 1.37 Å, Sn-H-B 180°. $[Ph_3Sn-H-SnPh_3]^+$: Both Sn-H 1.87 Å; Sn-H-B 147°.





Figure 2. Proposed reaction mechanism based on DFT calculations at the $\omega B97X-D/def2-TZVP$ level of theory. $R = tBu$ (blue, dashed) or Mes (green, dotted). $[HB(C_6F_5)_3]^-$ anion has been omitted for clarity. Energies in kcal mol $^{-1}$.

Changing the phosphine to $PrBu_3$ has a significant impact. First, the formation of the bridging hydride $[Ph_3Sn-H-SnPh_3]^+$ is now endothermic ($\Delta E^\ddagger_{TS1} = 14.1$, $\Delta E = 9.5$ kcal mol $^{-1}$; Figure 2, in blue) and the subsequent deprotonation faces a sizeable barrier ($\Delta E^\ddagger_{TS2} = 30.9$, $\Delta E = -0.2$ kcal mol $^{-1}$). The near thermoneutral reaction profile and high barrier accounts for the slow and modest formation of $[tBu_3PH][HB(C_6F_5)_3]$ and explains why the $[tBu_3P-SnPh_3][HB(C_6F_5)_3]$ intermediate can be isolated after short reaction times and immediate work-up.^[2c]

To support the intermediacy of the bridging $[Ph_3Sn-H-SnPh_3]^+$ cation, we combined $[Ph_3C][B(C_6F_5)_4]$ with 2 equiv of Ph_3SnH in C_6H_5Cl at $-35^\circ C$ in order to access this species by hydride abstraction (Scheme 4). Indeed, after 1 hour, the characteristic yellow color of the trityl cation disappeared, and an expected downfield shift of the aromatic 1H nuclei of the Ph_3Sn species in combination with a broadening of the hydride peak at 6.91 ppm was observed by 1H NMR spectroscopy (Supporting Information, Figures S20, S21).^[15] In addition, the spectrum evidenced formation of triphenylmethane ($\delta^1H = 5.55$).^[18] As predicted by DFT, subsequent addition of $PMes_3$ led to formation of $[Mes_3PH][B(C_6F_5)_4]$ by



Scheme 4. Hydride abstraction from Ph_3SnH using $[Ph_3C][B(C_6F_5)_4]$ and subsequent reaction with $PMes_3$.

deprotonation, as observed by ^1H - and ^{31}P -NMR spectroscopy (Supporting Information, Figure S23, S24), and the formation of $\text{Ph}_3\text{Sn-SnPh}_3$, evidenced by ^{119}Sn NMR spectroscopy (Supporting Information, Figure S25). Addition of PtBu_3 instead of PMes_3 afforded both $[\text{tBu}_3\text{PH}][\text{B}(\text{C}_6\text{F}_5)_4]$ and $[\text{tBu}_3\text{P-SnPh}_3][\text{B}(\text{C}_6\text{F}_5)_4]$ according to ^{31}P -NMR spectroscopy (approx. 4:5 ratio; Supporting Information, Figure S26), since both the forward and reverse pathways (in blue, Figure 2) have accessible barriers (21.4 and 4.6 kcal mol^{-1} , respectively); this yields the thermodynamically controlled product distribution. These findings demonstrate that after heterolytic Sn-H bond cleavage to form $[\text{R}_3\text{P-SnPh}_3][\text{HB}(\text{C}_6\text{F}_5)_3]$, subsequent bond metathesis leads to formation of $[\text{R}_3\text{PH}][\text{HB}(\text{C}_6\text{F}_5)_3]$ via a bridging hydride intermediate and highlights that the complete reaction of $\text{PMes}_3/\text{B}(\text{C}_6\text{F}_5)_3$ and $\text{PtBu}_3/\text{B}(\text{C}_6\text{F}_5)_3$ with Ph_3SnH is accessible via heterolytic polar pathways.

Next, we set out to analyze the reaction of $\text{PMes}_3/\text{B}(\text{C}_6\text{F}_5)_3$ with tetrachloro-1,4-benzoquinone (TCQ) for which Stephan et al. detected radical formation (PMes_3^+) by EPR spectroscopy.^[2c] They postulated that this proceeds via SET from PMes_3 to $\text{B}(\text{C}_6\text{F}_5)_3$ to form the corresponding radical ion pair $[\text{PMes}_3^+, \text{B}(\text{C}_6\text{F}_5)_3^-]$, after which 2 equiv of $\text{B}(\text{C}_6\text{F}_5)_3^-$ react with the quinone to form dianion **7**, while 1 equiv of $\text{B}(\text{C}_6\text{F}_5)_3^-$ affords the neutral adduct $\text{Mes}_3\text{P-TCQ-B}(\text{C}_6\text{F}_5)_3 **4** (Scheme 5).^[2c] We performed this reaction in the absence of light and found that the reaction still proceeds rapidly, forming a deep purple solution for which EPR spectroscopy confirmed the formation of PMes_3^+ (two-line signal simulated with $g_{\text{iso}} = 2.0050$, $A_{\text{iso}} = 670.00 \text{ MHz}$),^[2c,19] but also showed for the first time a featureless signal ($g_{\text{iso}} = 2.0058$) that we attribute to a TCQ centered radical anion, most likely $\text{TCQ-B}(\text{C}_6\text{F}_5)_3^-$ (Figure 3). Furthermore, we noted an unknown smaller third signal, which was also reported by Müller and Klare et al. when combining PMes_3 and the strongly accepting silyl and trityl cations.^[20,21]$

So, how is it possible that radicals are formed in the dark? Clearly, a strong electron acceptor is required to oxidise





Figure 3. Experimental EPR spectrum (bottom) for reaction of PMes_3 , $\text{B}(\text{C}_6\text{F}_5)_3$ and TCQ (2:2:1) and simulated spectra for PMes_3^+ , $\text{TCQ-B}(\text{C}_6\text{F}_5)_3^-$ and the third smaller signal. See the Supporting Information for experimental and simulation parameters. HFI = hyperfine interaction.

PMes_3 ($\text{IE}_\text{D} = 5.25 \text{ eV}$; $\text{SCRf}^{[15]}/\omega\text{B97X-D/6-311+G(d,p)}$, solvent = chlorobenzene) and neither $\text{B}(\text{C}_6\text{F}_5)_3$ nor TCQ are suitable ($E_\text{A} = 3.31$ and 4.45 eV , respectively)^[15] to accommodate the needed thermal SET. Yet, $\text{B}(\text{C}_6\text{F}_5)_3$ can coordinate to one of the carbonyl moieties of TCQ, affording the corresponding Lewis adduct $\text{TCQ-B}(\text{C}_6\text{F}_5)_3$, which has an increased electron affinity ($E_\text{A} = 5.57 \text{ eV}$) and therefore should be capable of oxidizing PMes_3 .^[11] Note that such interactions between a Lewis acid and a carbonyl moiety are typically exploited in photo-redox catalysis to facilitate SET events.^[22] As the carbonyl moieties of TCQ are electron poor, the interaction with $\text{B}(\text{C}_6\text{F}_5)_3$ is weak ($\Delta E = -4.6$, $\Delta G^\circ_{298\text{K}} =$



Scheme 5. a) Orbitals involved in the SET between PMes_3 and the $\text{TCQ-B}(\text{C}_6\text{F}_5)_3$ adduct. b) Reactivity, featuring all possible pathways for the reaction of TCQ, $\text{B}(\text{C}_6\text{F}_5)_3$, and PMes_3 .




10.7 kcal mol⁻¹) leading to an equilibrium featuring low concentrations of the TCQ-B(C₆F₅)₃ adduct, which supports the notion of Stephan et al. that no interaction between B(C₆F₅)₃ and TCQ is observable by NMR spectroscopy.^[2c] However, in presence of PMes₃, the transient TCQ-B(C₆F₅)₃ adduct will undergo SET from the PMes₃ HOMO to the TCQ-B(C₆F₅)₃ LUMO (Scheme 5, left) generating the radical ion pair [PMes₃⁺, TCQ-B(C₆F₅)₃⁻] **3**, which drives the equilibrium towards the TCQ-B(C₆F₅)₃ adduct (Scheme 5b, blue). Subsequent radical coupling of PMes₃⁺ and TCQ-B(C₆F₅)₃⁻ via a computed 8 kcal mol⁻¹ (ΔG°_{298K}) barrier leads to the formation of Mes₃P-TCQ-B(C₆F₅)₃ **4** as observed experimentally by Stephan et al. (Scheme 5b, black).^[2c] This mechanism highlights that, rather than directly participating in SET, B(C₆F₅)₃ is facilitating the process by increasing the electron affinity of the quinone acceptor.

As TCQ features two carbonyl moieties, coordination of two B(C₆F₅)₃ molecules can also occur prior to SET (**2**→**5** $\Delta E = -16.0$, $\Delta G^\circ_{298K} = -2.9$ kcal mol⁻¹; coordination after SET is unlikely: **3**→**6** $\Delta E = 2.7$, $\Delta G^\circ_{298K} = 18.8$ kcal mol⁻¹), yielding radical ion pair [PMes₃⁺, (C₆F₅)₃B-TCQ-B(C₆F₅)₃⁻] **6** (Scheme 5b, green). The radical anion of **6** has a high electron affinity ($E_A = 6.11$ eV), which allows another SET from a second equiv of PMes₃ to generate dianion **7**.^[2c] To complete the picture, dianion **7** is in equilibrium with TCQ-B(C₆F₅)₃⁻ radical anion **3** ($\Delta E = 5.2$, $\Delta G^\circ_{298K} = 2.3$ kcal mol⁻¹, Scheme 5b, purple) that can, as noted earlier, undergo radical


coupling with PMes₃⁺ to form **4** ($\Delta E = -48.9$, $\Delta G^\circ_{298K} = -20.4$ kcal mol⁻¹; Scheme 5b, black).

Changing the phosphine to PtBu₃ was shown by Stephan et al. to only yield the *t*Bu₃P-TCQ-B(C₆F₅)₃ adduct, akin to **4**, without detectable radicals or dianion **7**, which could indicate a different mechanism. The ionization energy of *t*Bu₃P ($IE_D = 5.54$ eV), however, suggests that SET from the phosphine to the TCQ-B(C₆F₅)₃ adduct ($E_A = 5.57$ eV) is still feasible. In this case, though, the subsequent radical coupling is barrierless,^[23] which leads to the immediate formation of *t*Bu₃P-TCQ-B(C₆F₅)₃ ($\Delta E = -56.5$, $\Delta G^\circ_{298K} = -20.4$ kcal mol⁻¹) and prevents detection of radical species or subsequent reactivity to form dianion **7**. This shows that, similar to our findings for Ph₃SnH, changing from PMes₃ to PtBu₃ does not alter the mechanism, but merely the energy levels along the reaction path leading to observation of radical intermediates for PMes₃, but not in case of PtBu₃.

Since the groups of Melen and Ooi recently reported FLP type reactions featuring radical formation when using B(C₆F₅)₃ and carbonyl containing substrates,^[6,7] we postulated that also in these cases Lewis acid coordination to the substrate could increase its electron acceptor capacity and promote SET (Scheme 6). Indeed, for methylvinylketone (MVK), the substrate utilized by Ooi et al. (Scheme 1d),^[7] we found that B(C₆F₅)₃ forms an adduct ($\Delta E = -16.3$, $\Delta G^\circ_{298K} = 0.9$ kcal mol⁻¹; SCRF/ ω B97X-D/6-311 + G(d,p), solvent = dichloroethane) and increases its electron affinity from 1.43 to 2.73 eV (Scheme 7, left). This decreases the energy gap between the ground state amine donor ($IE_D = 5.11$ eV; R = Me; Scheme 1d) and methylvinylketone acceptor pair to the corresponding radical ion pair [amine⁺, MVK-B(C₆F₅)₃⁻] from 3.68 to 2.38 eV, which results in visible light induced (1.5–3.1 eV, 400–800 nm) SET becoming feasible. Indeed, Ooi et al. used 400 nm light to promote this reaction. In case of substrate **9**, used by Melen et al. (Scheme 1c),^[6] we found a similar result. Binding of B(C₆F₅)₃ ($\Delta E = -19.1$, $\Delta G^\circ_{298K} = -0.2$ kcal mol⁻¹; SCRF/ ω B97X-D/6-311 + G(d,p), solvent = THF) increases the electron affinity from 2.31 to 3.56 eV (Scheme 7, right) bringing the energy required for SET (with PMes₃ as donor) down from 2.89 to 1.74 eV. This reduced energy gap (40.0 kcal mol⁻¹) is still sizeable and suggests that, also in this case, the SET is photo-induced and thus performing this reaction in broad daylight (or using a high-power



Scheme 6. Lewis acid coordination to a carbonyl moiety facilitating SET. LB = Lewis base.



Scheme 7. Change in electron affinity when B(C₆F₅)₃ coordinates and the resulting LUMO for two different B(C₆F₅)₃-coordinated substrates.

LED) will be beneficial. These results, in combination with the transient nature of the highly reactive $B(C_6F_5)_3^-$ species in solution,^[3a,24] make it highly plausible that also for these systems, $B(C_6F_5)_3$ is facilitating SET through binding to the substrate and increasing its electron affinity, instead of participating directly in SET.

Conclusion

Although the archetypal $PMes_3/B(C_6F_5)_3$ and $PrBu_3/B(C_6F_5)_3$ FLP systems can form high energy radical ion pairs via photo-induced single-electron transfer, we found that this pathway is not a major contributor in the reaction with H_2 or Ph_3SnH , and that in both cases the reactions occur via polar, heterolytic mechanisms. Furthermore, we discovered that the SET reactivity observed for FLP systems with substrates featuring carbonyl moieties is not the result of SET from the Lewis base directly to the borane Lewis acid. Instead, adduct formation between the Lewis acid and substrate activates the substrate for SET, after which electron donor-acceptor complex formation with the Lewis base provides the corresponding radical ion pair, via either thermal or photoinduced SET, depending on the energy required. To promote radical reactivity in cases based on photoinduced SET, it is thus important to locate the CT-band arising after Lewis acid coordination to determine the optimal wavelength for irradiation of reaction mixtures. These important mechanistic insights are of fundamental importance for both efficient usage of current radical FLP systems as well as the design of novel radical FLP systems and new examples of main-group redox catalysis,^[25] which we are currently exploring in our laboratories.

Acknowledgements

This work was supported by the Council for Chemical Sciences of The Netherlands Organization for Scientific Research (NWO/CW) by a VIDI grant (J.C.S.), NWA Idea Generator grant (J.C.S.) and a VENI grant (A.R.J.).

Conflict of interest

The authors declare no conflict of interest.

Keywords: frustrated Lewis pairs · radicals · reactivity · single-electron transfer · substrate coordination

- [1] a) G. C. Welch, R. R. San Juan, J. D. Masuda, D. W. Stephan, *Science* **2006**, *314*, 1124–1126; b) J. Paradies, *Coord. Chem. Rev.* **2019**, *380*, 170–183; c) G. Skara, F. De Vleeschouwer, P. Geerlings, F. De Proft, B. Pinter, *Sci. Rep.* **2017**, *7*, 16024; d) D. W. Stephan, *Science* **2016**, *354*, aaf7229; e) D. W. Stephan, *J. Am. Chem. Soc.* **2015**, *137*, 10018–10032; f) D. W. Stephan, G. Erker, *Angew. Chem. Int. Ed.* **2015**, *54*, 6400–6441; *Angew. Chem.* **2015**, *127*, 6498–6541; g) A. R. Jupp, D. W. Stephan, *Trends Chem.* **2019**, *1*, 35–48.

- [2] a) J. Paradies, *Eur. J. Org. Chem.* **2019**, 283–294; b) L. Liu, L. L. Cao, D. Zhu, J. Zhou, D. W. Stephan, *Chem. Commun.* **2018**, 7431–7434; c) L. Liu, L. L. Cao, Y. Shao, G. Ménard, D. W. Stephan, *Chem* **2017**, *3*, 259–267; d) L. Rocchigiani, G. Ciancaleoni, C. Zuccaccia, A. Macchioni, *J. Am. Chem. Soc.* **2014**, *136*, 112–115; e) T. A. Rokob, I. Bakó, A. Stirling, A. Hamza, I. Pápai, *J. Am. Chem. Soc.* **2013**, *135*, 4425–4437; f) S. Grimme, H. Kruse, L. Goerigk, G. Erker, *Angew. Chem. Int. Ed.* **2010**, *49*, 1402–1405; *Angew. Chem.* **2010**, *122*, 1444–1447; g) J. Daru, I. Bakó, A. Stirling, I. Pápai, *ACS Catal.* **2019**, *9*, 6049–6057.
- [3] a) E. J. Lawrence, V. S. Oganessian, G. G. Wildgoose, A. E. Ashley, *Dalton Trans.* **2013**, 782–789; b) W. E. Piers, A. J. V. Marwitz, L. G. Mercier, *Inorg. Chem.* **2011**, *50*, 12252–12262; c) Z. Dong, H. H. Cramer, M. Schmidtman, L. A. Paul, I. Siewert, T. Müller, *J. Am. Chem. Soc.* **2018**, *140*, 15419–15424; d) L. L. Liu, D. W. Stephan, *Chem. Soc. Rev.* **2019**, *48*, 3454–3463; e) Z. Dong, C. Pezzato, A. Sienkiewicz, R. Scopelliti, F. Fadaei-Tirani, K. Severin, *Chem. Sci.* **2020**, *11*, 7615–7618.
- [4] J. N. Bentley, E. Pradhan, T. Zeng, C. B. Caputo, *Dalton Trans.* **2020**, <https://doi.org/10.1039/d0dt00506a>.
- [5] Note triphenylphosphine forms an adduct with TCQ; see: F. Ramirez, S. Dershowitz, *J. Am. Chem. Soc.* **1956**, *78*, 5614–5622.
- [6] Y. Soltani, A. Dasgupta, T. A. Gazis, D. M. C. Ould, E. Richards, B. Slater, K. Stefkova, V. Y. Vladimirov, L. C. Wilkins, D. Wilcox, R. L. Melen, *Cell Rep. Phys. Chem.* **2020**, *1*, 10016.
- [7] Y. Aramaki, N. Imaizumi, M. Hotta, J. Kumagai, T. Ooi, *Chem. Sci.* **2020**, *11*, 4305–4311.
- [8] a) R. S. Mulliken, *J. Am. Chem. Soc.* **1952**, *74*, 811–824; b) R. Foster, *J. Phys. Chem.* **1980**, *84*, 2135–2141; c) S. V. Rosokha, J. K. Kochi, *Acc. Chem. Res.* **2008**, *41*, 641–653; d) for an ethylene/ I_2 EDA complex, see: A. Kalume, L. George, A. D. Powell, R. Dawes, S. A. Reid, *J. Phys. Chem. A* **2014**, *118*, 6838–6845.
- [9] For selected examples of electron donor-acceptor complexes in photocatalysis, see: a) E. Arceo, I. D. Jurberg, A. Álvarez-Fernández, P. Melchiorre, *Nat. Chem.* **2013**, *5*, 750–756; b) Y. Zhu, L. Zhang, S. Luo, *J. Am. Chem. Soc.* **2014**, *136*, 14642–14645; c) M. Nappi, G. Bergonzini, P. Melchiorre, *Angew. Chem. Int. Ed.* **2014**, *53*, 4921–4925; *Angew. Chem.* **2014**, *126*, 5021–5025; d) Ł. Woźniak, J. J. Murphy, P. Melchiorre, *J. Am. Chem. Soc.* **2015**, *137*, 5678–5681; e) S. R. Kandukuri, A. Bahamonde, I. Chatterjee, I. D. Jurberg, E. C. Escudero-Adan, P. Melchiorre, *Angew. Chem. Int. Ed.* **2015**, *54*, 1485–1489; *Angew. Chem.* **2015**, *127*, 1505–1509; f) F. Sandfort, F. Strieth-Kalthoff, F. J. R. Klauck, M. J. James, F. Glorius, *Chem. Eur. J.* **2018**, *24*, 17210–17214; g) J. Wu, P. S. Grant, X. Li, A. Noble, V. K. Aggarwal, *Angew. Chem. Int. Ed.* **2019**, *58*, 5697–5701; *Angew. Chem.* **2019**, *131*, 5753–5757; h) for a recent review, see: G. E. M. Crisenza, D. Mazzarella, P. Melchiorre, *J. Am. Chem. Soc.* **2020**, *142*, 5461–5476.
- [10] F. Holtrop, A. R. Jupp, N. P. Van Leest, M. Paradiz Dominguez, R. M. Williams, A. M. Brouwer, B. De Bruin, A. W. Ehlers, J. C. Sloodteweg, *Chem. Eur. J.* **2020**, *26*, 9005–9011.
- [11] For increasing the electron-acceptor capacity of quinones by hydrogen-bond donors, see: a) A. K. Turek, D. J. Hardee, A. M. Ullman, D. G. Nocera, E. N. Jacobsen, *Angew. Chem. Int. Ed.* **2016**, *55*, 539–544; *Angew. Chem.* **2016**, *128*, 549–554; for the $B(C_6F_5)_3$ coordination to and thereby redox-activation of S-nitrosothiols, see: b) V. Hosseiniinasab, A. C. McQuilken, A. Bakhoda, J. A. Bertke, Q. K. Timerghazin, T. H. Warren, *Angew. Chem. Int. Ed.* **2020**, *59*, 10854–10858; *Angew. Chem.* **2020**, *132*, 10946–10950; for the reduction of disulfides by ferrocene enabled by Lewis acids like $B(C_6F_5)_3$, see: c) L. L. Liu, L. L. Cao, Y. Shao, D. W. Stephan, *J. Am. Chem. Soc.* **2017**, *139*, 10062–10071.
- [12] G. C. Welch, D. W. Stephan, *J. Am. Chem. Soc.* **2007**, *129*, 1880–1881.



- [13] Note that the conversion values are based on the relative integrals of the $t\text{Bu}_3\text{P}$ and $[t\text{Bu}_3\text{PH}]^+$ resonances in each case. These values do not accurately correspond to conversion because the spectra were not obtained in a quantitative manner; however, the key result is that the values are the same regardless of whether the samples were irradiated or not.
- [14] a) A. Y. Houghton, J. Hurmalainen, A. Mansikkamäki, W. E. Piers, H. M. Tuononen, *Nat. Chem.* **2014**, *6*, 983–988; b) S. Rendler, M. Oestreich, *Angew. Chem. Int. Ed.* **2008**, *47*, 5997–6000; *Angew. Chem.* **2008**, *120*, 6086–6089; c) M. Mewald, M. Oestreich, *Chem. Eur. J.* **2012**, *18*, 14079–14084.
- [15] a) A. D. Horton, J. de With, *Organometallics* **1997**, *16*, 5424–5436; b) J. M. Blackwell, W. E. Piers, M. Parvez, *Org. Lett.* **2000**, *2*, 695–698; c) T. Beringhelli, D. Donghi, D. Maggioni, G. D'Alfonso, *Coord. Chem. Rev.* **2008**, *252*, 2292–2313; d) W. E. Piers, *Adv. Organomet. Chem.* **2005**, *52*, 1–76.
- [16] See the Supporting Information.
- [17] W. P. Neumann, B. Schneider, R. Sommer, *Justus Liebigs Ann. Chem.* **1966**, 692, 1–11.
- [18] M. Horn, L. H. Schappele, G. Lang-Wittkowski, H. Mayr, A. R. Ofial, *Chem. Eur. J.* **2013**, *19*, 249–263.
- [19] a) A. V. Il'yasov, Y. M. Kargin, E. V. Nikitin, A. A. Vafina, G. V. Romanov, O. V. Parakin, A. A. Kazakova, A. N. Pudovik, *Phosphorus Sulfur Relat. Elem.* **1980**, *8*, 259–262; b) M. Culcasi, Y. Berchadsky, G. Gronchi, P. Tordo, *J. Org. Chem.* **1991**, *56*, 3537–3542.
- [20] A. Merk, H. Großekappenberg, M. Schmidtman, M.-P. Luecke, C. Lorent, M. Driess, M. Oestreich, H. F. T. Klare, T. Müller, *Angew. Chem. Int. Ed.* **2018**, *57*, 15267–15271; *Angew. Chem.* **2018**, *130*, 15487–15492.
- [21] The third signal corresponds to either a radical species which shows hyperfine coupling with two ^{31}P nuclei, or a species featuring two uncoupled (that is, non-interacting) unpaired electrons, for which one shows hyperfine coupling to a ^{31}P nucleus. One possible assignment would be $\text{Mes}_3\text{P}-\text{PMes}_3^+$, although this species has no stable minimum on the potential energy surface; see also: S. Tojo, S. Yasui, M. Fujitsuka, T. Majima, *J. Org. Chem.* **2006**, *71*, 8227–8232.
- [22] a) J. Du, K. L. Skubi, D. M. Schultz, T. P. Yoon, *Science* **2014**, *344*, 392–396; b) T. P. Yoon, *Acc. Chem. Res.* **2016**, *49*, 2307–2315; c) K. N. Lee, M.-Y. Ngai, *Chem. Commun.* **2017**, 53, 13093–13112; d) E. Speckmeier, P. J. W. Fuchs, K. Zeitler, *Chem. Sci.* **2018**, *9*, 7096–7103.
- [23] The computed barrier ($\Delta G^\circ_{298\text{K}} = -0.4\text{ kcal mol}^{-1}$) is less than the error margin and can thus be considered approximately equal to zero.
- [24] R. J. Kwaan, C. J. Harlan, J. R. Norton, *Organometallics* **2001**, *20*, 3818–3820.
- [25] a) R. C. McAtee, E. J. McClain, C. R. J. Stephenson, *Trends Chem.* **2019**, *1*, 111–125; b) N. A. Romero, D. A. Nicewicz, *Chem. Rev.* **2016**, *116*, 10075–10166; c) M.-C. Fu, R. Shang, B. Zhao, B. Wang, Y. Fu, *Science* **2019**, *363*, 1429–1434; d) Z. Deng, J.-H. Lin, J.-C. Xiao, *Nat. Commun.* **2016**, *7*, 10337.

Manuscript received: July 14, 2020

Accepted manuscript online: August 25, 2020

Version of record online: ■ ■ ■ ■ ■ ■ ■ ■ ■ ■

Research Articles



Frustrated Lewis Pairs

F. Holtrop, A. R. Jupp, B. J. Kooij,
N. P. van Leest, B. de Bruin,
J. C. Slootweg* ————— ■■■-■■■

Single-Electron Transfer in Frustrated
Lewis Pair Chemistry



The radical reactivity of archetypal P/B frustrated Lewis pair (FLP) systems is probed through utilization of the light dependency of direct single-electron transfer from the phosphine to the borane to determine that reactivity with H_2 does not proceed via a radical pathway. For radical FLP reactions with carbonyl-containing substrates, direct SET is excluded in favor of borane coordination to the carbonyl, after which SET occurs.