

# Cardiac contractility modulation increases action potential duration dispersion and decreases ventricular fibrillation threshold via $\beta$ 1-adrenoceptor activation in the crystalloid perfused normal rabbit heart

Winter, James; Brack, Kieran E; Coote, John H; Ng, G André

DOI:

[10.1016/j.ijcard.2013.12.184](https://doi.org/10.1016/j.ijcard.2013.12.184)

License:

Creative Commons: Attribution (CC BY)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Winter, J, Brack, KE, Coote, JH & Ng, GA 2014, 'Cardiac contractility modulation increases action potential duration dispersion and decreases ventricular fibrillation threshold via  $\beta$ 1-adrenoceptor activation in the crystalloid perfused normal rabbit heart', *International Journal of Cardiology*, vol. 172, no. 1, pp. 144-154. <https://doi.org/10.1016/j.ijcard.2013.12.184>

[Link to publication on Research at Birmingham portal](#)

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.


When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

Download date: 17. Apr. 2024



# Cardiac contractility modulation increases action potential duration dispersion and decreases ventricular fibrillation threshold via $\beta_1$ -adrenoceptor activation in the crystalloid perfused normal rabbit heart ☆☆



James Winter <sup>a,1</sup>, Kieran E. Brack <sup>a,1</sup>, John H. Coote <sup>b,1</sup>, G. André Ng <sup>a,c,d,\*</sup>,<sup>1</sup>

<sup>a</sup> Cardiology Group, Department of Cardiovascular Sciences, University of Leicester, UK

<sup>b</sup> School of Clinical and Experimental Medicine, University of Birmingham, UK

<sup>c</sup> NIHR Leicester Cardiovascular Biomedical Research Unit, Leicester, UK

<sup>d</sup> University Hospitals of Leicester NHS Trust, Leicester, UK

## ARTICLE INFO

### Article history:

Received 11 July 2012

Received in revised form 19 June 2013

Accepted 31 December 2013

Available online 8 January 2014

### Keywords:

Cardiac contractility modulation

Non-excitatory stimulation (NES)

Ventricular arrhythmia

Ventricular fibrillation

Action potential duration dispersion

## ABSTRACT

**Background/objectives:** Cardiac contractility modulation (CCM) is a new treatment being developed for heart failure (HF) involving application of electrical current during the absolute refractory period. We have previously shown that CCM increases ventricular force through  $\beta_1$ -adrenoceptor activation in the whole heart, a potential pro-arrhythmic mechanism. This study aimed to investigate the effect of CCM on ventricular fibrillation susceptibility.

**Methods:** Experiments were conducted in isolated New Zealand white rabbit hearts (2.0–2.5 kg,  $n = 25$ ). The effects of CCM ( $\pm 20$  mA, 10 ms phase duration) on the left ventricular basal and apical monophasic action potential duration (MAPD) were assessed during constant pacing (200 bpm). Ventricular fibrillation threshold (VFT) was defined as the minimum current required to induce sustained VF with rapid pacing ( $30 \times 30$  ms). Protocols were repeated during perfusion of the  $\beta_1$ -adrenoceptor antagonist metoprolol (1.8  $\mu$ M). In separate hearts, the dynamic and spatial electrophysiological effects of CCM were assessed using optical mapping with di-4-ANEPPS.

**Results:** CCM significantly shortened MAPD close to the stimulation site (Basal:  $102 \pm 5$  [CCM] vs.  $131 \pm 6$  [Control] ms,  $P < 0.001$ ). VFT was reduced during CCM ( $2.6 \pm 0.6$  [CCM] vs.  $6.1 \pm 0.8$  [Control] mA,  $P < 0.01$ ) and was correlated ( $r^2 = 0.40$ ,  $P < 0.01$ ) with increased MAPD dispersion ( $26 \pm 4$  [CCM] vs.  $5 \pm 1$  [Control] ms,  $P < 0.01$ ) ( $n = 8$ ). Optical mapping revealed greater spread of CCM induced MAPD shortening during basal vs. apical stimulation. CCM effects were abolished by metoprolol and exogenous acetylcholine. No evidence for direct electrotonic modulation of APD was found, with APD adaptation occurring secondary to adrenergic stimulation.

**Conclusions:** CCM decreases VFT in a manner associated with increased MAPD dispersion in the crystalloid perfused normal rabbit heart.

© 2014 The Authors. Published by Elsevier Ireland Ltd. Open access under [CC BY license](http://creativecommons.org/licenses/by/4.0/).

**Abbreviations:** ACh, acetylcholine; APD, action potential duration; CCM, cardiac contractility modulation; LV, left ventricle; MAPD<sub>90</sub>, monophasic action potential duration at 90% repolarization; NE, norepinephrine; VFT, ventricular fibrillation threshold.

☆☆ Funding: This work was supported by grants from the British Heart Foundation [PG/08/039, FS/12/2/29300] and Garfield Weston Trust [PMS/MMS-08/09-3023]. The study is part of the research portfolio supported by the Leicester NIHR Biomedical Research Unit in Cardiovascular Disease.

\* Corresponding author at: Department of Cardiovascular Sciences, Cardiology group, University of Leicester, Glenfield Hospital, Leicester LE39QP, UK. Tel.: +44 1162502438; fax: +44 1162875792.

E-mail address: [gan1@le.ac.uk](mailto:gan1@le.ac.uk) (G.A. Ng).

<sup>1</sup> All authors take responsibility for all aspects of the reliability and freedom from bias of the data presented and their discussed interpretation.

## 1. Introduction

Cardiac contractility modulation (CCM) is an electrical device therapy being developed for the treatment of heart failure (HF), which remains a significant clinical burden despite decades of research and the development of useful drug therapy [1]. During CCM, electrical signals are applied to the ventricular myocardium to increase force and are timed to coincide with the absolute refractory period. Experimental and clinical studies demonstrate that CCM improves contractile performance, increases patient exercise capacity and improves quality of life (reviewed by Winter et al. [2]). Several studies have highlighted that CCM reverses the molecular remodelling associated with heart failure opposing the down-regulation of several calcium handling proteins (e.g. phospholamban, sarco-endoplasmic reticulum ATPase (SERCA)),

increased expression of detrimental signalling factors (e.g. atrial natriuretic peptide) and fibrosis, which may contribute to its long term beneficial action [2,3].

We have previously demonstrated that the acute effects of CCM are mediated via stimulated norepinephrine (NE) release resulting in cardiac  $\beta$ 1-adrenoceptor activation and shortening of locally recorded monophasic action potential duration (MAPD); in the crystalloid perfused normal rabbit heart [4]. The adrenergic dependence of CCM raises interesting questions regarding its neuro-cardiac action and the potential pro-arrhythmic nature of induced electrophysiological changes.

Efferent and afferent sympathetic and parasympathetic nerves richly innervate the cardiac ventricle and it has long been known that afferent nerves relay sensory information from the myocardium to the intra-cardiac and intra-thoracic ganglia, spinal cord and the brain for integration to regulate cardiac function [5]. It is recognised that local reflexes, acting through the cardiac and intra-thoracic ganglia, can modulate cardiac function on a beat-to-beat basis *without* involvement of the central nervous system [6]. It is perceivable that CCM captures both afferent and efferent nerve fibres, and that the former could engage intrathoracic ganglia or reflex activated sympathetic preganglionic neurones in the spinal cord. Furthermore, CCM may release acetylcholine (ACh) from parasympathetic nerves within the ventricle. ACh is known to antagonise NE synaptic release and antagonise post-synaptic adrenergic signalling and so it is necessary to establish whether ACh has a role in acute action of CCM (i.e. does ACh modulate the adrenergic component of CCM?).

It has long been acknowledged that adrenergic activation is pro-arrhythmic. We have previously shown that sympathetic nerve stimulation increases the susceptibility of the heart to ventricular fibrillation (VF) [7]. Given the adrenergic dependence of CCM and the pro-arrhythmic effect of positive inotropic drugs in HF, it is important to determine the effects of CCM on ventricular arrhythmia susceptibility [8].

The aims of this study were to investigate: 1) the role of pre and post-ganglionic signalling in CCM, 2) any role for ACh in modulating the acute ventricular response to CCM and 3) the effects of CCM on ventricular electrophysiology and arrhythmia susceptibility.

## 2. Methods

### 2.1. Animal welfare and ethical publishing declaration

Experiments were conducted on Adult male New Zealand White rabbit hearts (2.5–3.5 kg,  $n = 22$ ), using the non-innervated and innervated heart preparations. All procedures were undertaken in accordance with ethical guidelines set out by the UK Animals (Scientific Procedures) Act 1986 and conformed to the Guide for the Care and Use of Laboratory Animals Published by the US National Institutes of Health (NIH Publication No. 85-23, revised 2010).

The author(s) of this manuscript have certified that they comply with the Principles of Ethical Publishing in the International Journal of Cardiology.

### 2.2. Isolation of the non-innervated heart preparation

Adult male New Zealand White rabbit hearts were isolated as previously described [4]. In brief, animals were pre-medicated with ketamine (Ketaset, 10 mg/kg, Fort Dodge, UK), medetomidine hydrochloride (Sedator, 0.2 mg/kg, Dechra, UK) and butorphanol (Torbugesic, 0.05 mg/kg, Fort Dodge, UK) (i.m.). Following stable sedation, animals were sacrificed with an overdose of pentobarbitone sodium (Sagatal, Rhone Merieux, UK; 111 mg/kg body weight, i.v.) containing Heparin (1000 IU, Multiparin, UK) delivered via the marginal ear vein. The hearts were rapidly excised, placed into ice cold tyrode solution to reduce metabolic rate and retrogradely perfused through the ascending aorta in conditions of constant flow Langendorff mode (40 ml/min) using a Gilson Minipuls 3 peristaltic pump (Anachem, UK).

### 2.3. Isolation of heart with intact autonomic innervation

The innervated Langendorff perfused isolated heart preparation was utilised as previously described [9]. An illustration of the preparation can be found in Fig. 1B. In brief, following pre-medication, anaesthesia was maintained with i.v. propofol (5 mg as required, Rapinovel, Schering-Plough Animal Health, UK) for the remainder of surgery. Following tracheotomy animals were ventilated with room air using a small animal ventilator (Harvard Apparatus Ltd, UK; 60 breaths/min). The left and right vagus nerves were isolated, and

the blood vessels leading to and from the ribcage were ligated and dissected. Animals were sacrificed with an overdose of pentobarbitone sodium containing heparin (as above). The anterior portion of the ribcage was removed, and the descending aorta was cannulated. The pericardium was cut, and ice-cold Tyrode solution was applied to the surface of the heart. The preparation, extending from C1 to T12 vertebrae, was dissected from surrounding tissues and perfused through the descending aorta (100 ml/min) [9].

### 2.4. Solutions

Hearts were perfused with Tyrode solution of the following composition (mM):  $\text{Na}^+$  138.0,  $\text{K}^+$  4.0,  $\text{Ca}^{2+}$  1.8,  $\text{Mg}^{2+}$  1.0,  $\text{HCO}_3^-$  24.0,  $\text{H}_2\text{PO}_4^-$  0.4,  $\text{Cl}^-$  121.0, glucose 11.0 and acetate 20.0. The solution was continuously bubbled with 95%  $\text{O}_2$ –5%  $\text{CO}_2$  to maintain a constant pH of 7.4. Temperature was maintained at 37 °C. A 3 F polypropylene catheter (Portex, UK) was inserted at the left ventricular (LV) apex for drainage of Thebesian venous effluent.

### 2.5. Functional parameters

Intra-ventricular LV pressure (LVP) was measured with a fluid-filled latex balloon connected to a pressure transducer (MTLO380, ADInstruments Ltd, UK) inserted into the LV via the left atrium. LV end diastolic pressure was maintained between 0 and 5 mm Hg. Aortic perfusion pressure (PP) was monitored with a second pressure transducer connected in series to the aortic cannula.

### 2.6. Cardiac electrical recording and pacing

MAP-TIP recording catheters (73-0150, Harvard Apparatus, UK), with a tip diameter of 1.5 mm electrode spacing of 4 mm, were used to record action potentials from the epicardial surface of the LV using a DC-coupled high input impedance differential amplifier (Joint Biomedical Workshop, University of Leicester, UK). MAPs were recorded from the LV epicardium at apical and/or basal sites depending on the experimental protocol.

Pacing electrode locations are illustrated on Fig. 1A. Endocardial pacing electrodes were inserted through the pulmonary trunk or left atrial appendage for RV and LV apical endocardial pacing, respectively (electrode diameter = 2 mm, spacing = 4 mm). For epicardial pacing, a pair of platinum hook electrodes (spacing = 4 mm) were inserted into the basal anterior LV free wall. Pacing stimuli delivered using a constant current electrical stimulator (DS7A, Digitimer Ltd, UK).

### 2.7. CCM signal generation and delivery

Square wave electrical pulses were generated using a Neurolog modular system (Digitimer, UK) with signals delivered using a constant current stimulator (Model A385, World Precision Instruments, UK). CCM stimuli were triggered using the pacing stimulus as previously described [4]. CCM signals were applied to the epicardial surface of the LV through a pair of platinum hook electrodes (inter-electrode distance ~ 1 cm) as a biphasic waveform with equal positive and negative phase amplitudes (stimulus amplitude = 20 mA and phase duration = 10 ms). We have previously demonstrated that these parameters induce optimal responses [4]. CCM signals were timed to coincide with the plateau phase using a locally recorded MAP (see Fig. 1.) and were delivered 2–3 min, allowing for a 30 second period of steady state. A 10–15 minute rest period between stimulations was used throughout the experiment for parameters to return to baseline. Hearts were paced during all CCM protocols at a 300 ms cycle length.

#### 2.7.1. Protocols

**2.7.1.1. Role of the intracardiac/intrathoracic ganglia.** The effects of CCM on ventricular and electrophysiological performance were assessed in the *innervated Langendorff heart* in the absence and presence of the nicotinic ACh channel antagonist hexamethonium (0.5 mM) [10]. Ganglionic blockade was confirmed by inhibition of bradycardia with right vagus nerve stimulation (5 Hz, 5 V). CCM was applied in basal regions of the LV.


**2.7.1.2. Role of ACh.** The effects of CCM on ventricular and electrophysiological performance were assessed in the *non-innervated Langendorff heart* preparation in the absence (Control) and presence of;

- A. The muscarinic ACh receptor antagonist atropine (0.1 mM) [10].
- B. Exogenous ACh (1  $\mu\text{M}$ ) [13] with and without atropine (0.1  $\mu\text{M}$ ) [11].

2–3 stimulations were conducted at baseline and during perfusion. The order of perfusion was randomised from heart to heart to reduce any bias. CCM was applied in basal regions of the LV.

#### 2.7.1.3. VFT, ERP and APD dispersion

**2.7.1.3.1. VFT.** The susceptibility of the heart to VF was studied by measuring the ventricular fibrillation threshold (VFT) in the *non-innervated Langendorff heart*. VFT was obtained by RV endocardial, LV endocardial or LV epicardial pacing using a train of 30-stimuli (CL = 30 ms, 2 ms pulse width) spanning the refractory period following a 20-beat S1 drive train (CL = 300 ms, 2 ms pulse width) and determined by progressively increasing the pacing current (0.5 mA steps), with 2-second rest period before the next pacing train if no VF was induced (see Fig. 2). VFT was defined as the minimum current required to induce sustained VF. VFT was assessed in the absence and



**Fig. 1.** CCM signal delivery and the isolated innervated rabbit heart. A) Illustration of heart with pacing, recording and CCM stimulation sites. A square wave bipolar signal is applied directly to the surface of the left ventricle (LV) during the absolute refractory period. Signals were applied to either basal or apical regions of the LV epicardium depending upon the experimental protocol. Hearts were paced at a cycle length of 300 ms. Monophasic action potentials (MAP) were recorded in both basal and apical regions. B) Illustration of isolated innervated Langendorff rabbit heart preparation. Key: LA = left atrium, LVP = left ventricular pressure, PP = perfusion pressure, RA<sub>EG</sub> = right atrial electrogram, RA = right atrium, RVNS = right vagus nerve stimulation, RV = right ventricle.

subsequently in the presence of CCM at basal and apical regions. CCM was applied for a minimum period of 60 s during constant pacing (CL = 300 ms) before commencing the VFT protocol. During VFT protocol, CCM was applied during the S1 beats but not during the rapid pacing period of the VF protocol. VF was terminated using a 2–3 ml bolus injection of 50 mM KCL. Hearts were allowed to beat intrinsically between protocols, allowing for a 10-minute recovery period before initiation of pacing. A 5-minute period was allowed for the stabilisation of MAPD before commencement of each protocol. In separate hearts, the effects of CCM on VFT were repeated during perfusion of the  $\beta_1$ -adrenoceptor antagonist Metoprolol (1.8  $\mu$ M) [4,9]. Baseline, Basal CCM and Apical CCM data reflects the average of 3 VFT repeats at each steady state during the experimental protocols, in each heart. The order of pacing was randomised in each experiment.


**2.7.1.3.2. Effective refractory period.** The effective refractory period (ERP) was estimated using an extra-stimulus protocol consisting of a 20-beat drive train (S1) at a constant CL (300 ms) followed by an extra-stimulus (S2) at progressively shorter CLs (200 ms, 5 ms steps). ERP was measured in RV endocardial, LV endocardial and LV epicardial regions during basal and apical CCM. ERP was defined as the longest S1-S2 interval that failed to capture. Pacing stimuli were applied at twice the diastolic threshold current, with a pulse width of 2 ms.

**2.7.1.3.3. MAPD dispersion.** Ventricular MAPD dispersion was measured as the difference between in MAPD the two sites using contact electrodes (*non-innervated hearts*). These data were obtained during the period of constant pacing (CL = 300 ms), prior to the assessment of VFT, as described above, at baseline and during CCM delivery in basal and apical regions.

**2.7.1.4. AP dynamics and spatial effects of CCM (optical mapping).** APs were recorded with optical mapping of the anterior LV surface of the non-innervated rabbit heart (18 × 18 mm). Optical action potentials (AP) were recorded following a bolus injection of di-4 ANEPPS (40  $\mu$ l, 1 mg/ml in DMSO, Invitrogen, USA). Hearts were illuminated episcopically with an LED light source (535 nm). Emitted light was filtered through a 630 nm long-pass filter and collected using a Hamamatsu 16 × 16 element photodiode array (Cairn Research, Faversham, UK). Hearts were mechanically uncoupled with blebbistatin (5  $\mu$ M). Optical studies were conducted in the non-innervated rabbit heart.

**2.7.1.4.1. APD dynamics.** Experiments were performed to assess the dynamics to APD change during CCM. Optical APs were recorded continuously at stable baseline (300 ms CL) and during applications of biphasic CCM signals ( $\pm 20$  mA, 20 ms) to the LV free wall. The effects of CCM on APD was assessed at regions within 2 mm of the stimulating electrode and in APs with a clear deflection associated with CCM stimulation.





**Fig. 2.** Ventricular fibrillation threshold protocol. VFT was obtained by right ventricular pacing using a train of 30 stimuli (CL = 30 ms) spanning the refractory period following a 20-beat drive train (CL = 300 ms) and determined by progressively increasing the pacing current in 0.5 mA steps, with a 2-second rest period before the next pacing train if no VF was induced. VFT was defined as the minimum current required inducing sustained VF. VFT was assessed in the absence and presence of CCM applied separately in basal and apical regions. CCM was applied for a 60-second period during constant pacing (CL = 300 ms) before commencement of the protocol, and throughout the S1 drive train. CCM was not applied during the 2-second rest period between consecutive S1 drive trains or during the rapid pacing period.

**2.7.1.4.2. Spatial effects.** CCM ( $\pm 20$  mA, 20 ms) was applied in basal and apical regions during constant pacing (CL = 300 ms). The degree of spread of APD shortening during CCM was assessed from the site of CCM signal delivery (Base  $\rightarrow$  Apex or Apex  $\rightarrow$  Base). Studies were conducted in the non-innervated rabbit heart.

### 2.8. Signal measurements and analysis

Functional parameters and MAPs were recorded with a PowerLab 8 s system and digitised at 2 kHz using Chart and Scope software (ADInstruments Ltd). Optical AP signals were digitised at 2 kHz and recorded on a custom designed computer system (National Instruments, USA) using QRecord software (Dr Francis Burton, University of Glasgow, UK). LV contractile performance was assessed from an average of 20 cardiac cycles during baseline and subsequently during the steady state CCM response. Contact MAPD was measured at 90% repolarisation (MAPD<sub>90</sub>) averaged over 20-cardiac cycles recorded during steady state before and immediately on cessation of CCM stimulation due to signal interference from the current used during CCM. Optical APD<sub>90</sub> was assessed before and during the steady state response to CCM signal delivery as there was no signal interference from the current used during CCM on the recorded APs. Both MAPD<sub>90</sub> and optical APD<sub>90</sub> calculations were performed using custom written analysis software (NewMap and Optiq respectively, Dr. F. Burton).

### 2.9. Statistical methods

Statistical comparisons were made using Student's paired t-tests, one- or two-way ANOVA where appropriate with Bonferroni post-hoc test.  $P < 0.05$  was considered significant. All statistically significant differences reported in this manuscript represent a sample size to give a minimum statistical power of 0.94, at an alpha level of 0.05. Data are presented as mean values  $\pm$  standard error of the mean.

## 3. Results

### 3.1. Role of intracardiac/intrathoracic ganglia

Data illustrating the effects of hexamethonium perfusion on the acute ventricular response to CCM, in the *innervated Langendorff heart*, are summarised in Fig. 3. Ganglionic blockade was confirmed by the abolition of right vagus nerve stimulation induced bradycardia (Fig. 3A&B). In control, right vagus nerve stimulation reduced heart rate by  $76 \pm 9$  bpm and was abolished during perfusion with 0.5 mM hexamethonium ( $\Delta = -1 \pm 1$  bpm,  $n = 3$ ). During hexamethonium perfusion, CCM stimulated enhancement of LV pressure, maximal rate of change of pressure and shortening of MAPD<sub>90</sub> (Fig. 3C–E) were comparable to control ( $n = 5$ ).

### 3.2. Role of ACh

#### 3.2.1. Atropine

Perfusion of atropine had no effect on the increase in LVP ( $\Delta 3.9 \pm 0.7$  [Atropine] vs.  $\Delta 4.2 \pm 0.6$  [Control] mm Hg),  $dP/dt_{\max}$  ( $\Delta 186 \pm 47$  [Atropine] vs.  $\Delta 175 \pm 27$  [Control] mm Hg/s) or MAPD<sub>90</sub> shortening ( $\Delta -27 \pm 9$  [Atropine] vs.  $\Delta -28 \pm 7$  [Control] ms) associated with CCM ( $n = 5$ ), indicating no role for endogenous ACh release in the acute ventricular effects of CCM.

#### 3.2.2. Exogenous ACh

Raw and mean data illustrating the effect of exogenous ACh perfusion on the acute effects of CCM are presented in Fig. 4. Exogenous ACh inhibited the increase in peak LV pressure (Fig. 4B), maximal rate of change of pressure ( $\Delta 126 \pm 16$  vs.  $\Delta 30 \pm 4$  mm Hg/s,  $P < 0.05$ ) and shortening of MAPD<sub>90</sub> (Fig. 4C) associated with CCM ( $n = 4$ ). The inhibition of CCM with ACh was reversed during combined perfusion with 0.1  $\mu$ M atropine (Fig. 4,  $n = 4$ ). Exogenous ACh perfusion caused coronary vasodilation, evidenced by a reduction in PP ( $54.4 \pm 10.9$  vs.  $50.9 \pm 10.6$  mm Hg,  $P < 0.05$ ), but had no significant effect on LVP or MAPD<sub>90</sub> in baseline conditions.


### 3.3. VFT, ERP and MAPD dispersion

#### 3.3.1. VFT

Mean data on the effect of CCM on VFT are illustrated in Fig. 5A. CCM was associated with a significant reduction in VFT during RV (Fig. 5Ai) and LV (Fig. 5Aii) endocardial pacing, indicating increased susceptibility to VF. Similarly CCM applied at basal regions, during LV epicardial pacing, reduced VFT with a borderline significant trend associated with apical CCM (Fig. 5Aiii,  $P = 0.06$ ). The magnitude change in VFT with apical and basal CCM was similar in all conditions, with the exception of LV epicardial stimulation where there was a larger reduction in VFT during basal vs. apical CCM (Fig. 5Aiii). Perfusion of metoprolol abolished the reduction in VFT during basal ( $5.35 \pm 0.8$  [Basal CCM + Metoprolol] vs.  $5.4 \pm 0.7$  [Metoprolol] mA) and apical CCM ( $5.42$  [Apical CCM + Metoprolol] vs.  $5.4 \pm 0.7$  [Metoprolol] mA) (RV endocardial pacing,  $n = 6$ ).

#### 3.3.2. ERP


Changes in ERP were dependent upon the relative location of the pacing site and site of CCM stimulation. ERP was shortened during




**Fig. 3.** Role of ganglionic transmission in the acute ventricular effects of CCM. A) Raw data illustrating instantaneous left ventricular pressure (LVP) and heart rate (HR) during right vagus nerve stimulation (RVNS, 5Hz, 5 V) in the absence (left) and presence of 0.5 mM hexamethonium (right). B) Mean data representing the bradycardia effect of RVNS in the absence and presence of hexamethonium ( $n = 3$ ). C–E) Mean data demonstrating the change in LVP (C), rate of pressure production ( $dPdt_{max}$ , D) and basal  $MAPD_{90}$  (E) during basal CCM in the absence and presence of hexamethonium ( $n = 5$ ). Key: BL = baseline, SS = steady state, CCM = cardiac contractility modulation. Baseline vs. stimulation (VNS or CCM); \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ .

LV epicardial pacing when CCM signals were applied to basal (close to) but not apical regions (far from) (Fig. 5Diii). Similarly there was a trend for ERP shortening, during LV endocardial pacing, with CCM applied in

apical (close) but not basal regions (far) (Fig. 5Dii,  $P = 0.07$ ). No effect of CCM on ERP was seen during RV endocardial pacing with either basal or apical stimulation (Fig. 5Di).



**Fig. 4.** Effect of exogenous acetylcholine on the acute ventricular response to CCM. A) Raw data illustrating left ventricular pressure (LVP) and peak LVP in the absence and presence of exogenous acetylcholine (ACh, 1  $\mu$ M) and ACh + atropine (0.1  $\mu$ M) perfusion associated with basal CCM. B & C) Mean data demonstrating the change in peak LVP and basal monophasic action potential duration ( $MAPD_{90}$ ) in the absence and presence of ACh and ACh + atropine ( $n = 4$ ). Difference in CCM response between experimental conditions; \* $P < 0.05$ , \*\* $P < 0.01$ .



**Fig. 5.** The effect of CCM on the ventricular fibrillation threshold and ventricular dispersion of monophasic action potential duration. A) Mean data showing ventricular fibrillation threshold (VFT) at baseline (BL) and during CCM at basal and apical sites during RV endocardial, LV endocardial and LV epicardial pacing. B) Mean data illustrating monophasic action potential duration (MAPD<sub>90</sub>) at BL and during CCM. C) Maximum–minimum MAPD<sub>90</sub> at BL and with CCM. D) Effective refractory period (ERP) at baseline and during CCM E) The correlation between Max–Min MAPD<sub>90</sub> vs. VFT (regression fit represents combined datasets, with the exception of (Eiii) in which basal CCM data points are excluded (dotted line)). Comparison to baseline; \*P < 0.05, \*\*P < 0.01, \*\*\*P < 0.001, base vs. apex; #P < 0.05, ##P < 0.01 (VFT, MAPD; n = 8, ERP; n = 4, same hearts).

**3.3.3. MAPD dispersion**

Mean data on the effect of CCM on MAPD<sub>90</sub> and max-min dispersion are presented in Fig. 5B & C. Un-stimulated MAPD<sub>90</sub> was similar

between basal and apical regions during RV and LV endocardial pacing but longer in basal, vs. apical, regions during LV epicardial pacing (Fig. 5B). CCM caused significant shortening of MAPD<sub>90</sub> close to the

site of stimulation but not at distant sites (Fig. 5B) resulting in a significant increase of max-min MAPD<sub>90</sub> (Fig. 5C) (n = 8). The increase in dispersion of repolarisation was similar during apical and basal CCM and occurred in all pacing protocols (Fig. 5C). The scatterplots presented in Fig. 5E show that VFT was inversely correlated with max-min MAPD<sub>90</sub> during RV endocardial (5Ei) and LV endocardial (5Eii), but not LV epicardial (5Eiii), pacing. It is apparent that the reduction in VFT with basal CCM, during LV epicardial pacing, may be the product of increased MAPD<sub>90</sub> dispersion and shortening of ERP at the site of CCM stimulation. Analysis of LV epicardial pacing, when basal CCM data are removed from the dataset, reveals a trend of correlation between VFT and max-min MAPD<sub>90</sub> dispersion (Fig. 5Eiii), similar to that seen with LV and RV endocardial pacing.

Perfusion of metoprolol abolished the shortening of MAPD<sub>90</sub> during basal (131 ± 4 [Metoprolol + Basal CCM] vs. 128 ± 2 [Metoprolol] ms) and apical CCM (132 ± 4 [Metoprolol + Apical CCM] vs. 129 ± 5 [Metoprolol] ms). Max-min MAPD<sub>90</sub> dispersion during CCM stimulation with metoprolol was similar to baseline values (4.7 ± 2.0 [Metoprolol + Basal CCM] vs. 4.2 ± 1.7 [Metoprolol] ms) (RV endocardial pacing, n = 6).


### 3.4. AP dynamics and spatial electrophysiological effects of CCM

#### 3.4.1. Optical vs. contact

Optical action potentials (n = 4) recorded at baseline under unstimulated conditions were significantly longer than those recorded with contact electrodes (RV endocardial pacing, n = 8) in both basal (159 ± 6 [Optical] vs. 131 ± 6 [Contact] ms, P < 0.01) and apical (169 ± 6 [Optical] vs. 133 ± 6 [Contact] ms, P < 0.01) regions. Optical recordings were associated with a significant apico-basal dispersion of APD<sub>90</sub> (169 ± 6 [Apex] vs. 159 ± 6 [Base] ms, P < 0.01) not present in contact electrode recordings (133 ± 6 [Apex] vs. 131 ± 6 [Base]) and attributable to the adverse electrophysiological effects of blebbistatin.

#### 3.4.2. AP Dynamics

Raw and mean data demonstrating time dependent adaptation of ventricular APD during LV free wall CCM are presented in Fig. 6. APs were recorded close to the site of CCM stimulation (within 2 mm). Despite a clear deflection attributable to the CCM stimulus (Fig. 6), there was no instantaneous effect on APD<sub>90</sub>. Rather, APD shortened



**Fig. 6.** Temporal changes in action potential duration during local CCM stimulation. A) Example action potentials demonstrating the effects of CCM on action potential duration (APD<sub>90</sub>) on the first beat, and after 10–30 s of stimulation (optical). B) Mean data demonstrating the time course of APD<sub>90</sub> changes during local CCM stimulation. CCM vs. baseline; \*\*\*P < 0.001 (n = 3).

over a period of seconds with a maximal response at 30 s. Responses recovered to baseline 2–3 min after cessation of stimulation (data not shown).

#### 3.4.3. Spatial effects of CCM

Raw and mean data demonstrating the effect of CCM on optically recorded APs are illustrated in Fig. 7A. In accordance with data obtained from contact electrodes, CCM caused APD shortening close to the stimulation site, but not at recording sites distant to CCM stimulation. Raw data illustrating the spatial spread of optical AP shortening during high amplitude basal CCM is shown in Fig. 7 and shows that the degree of APD shortening decays relatively uniformly away from the stimulation site. The range of APD shortening appears greater during basal CCM as defined by distance from site of stimulation when APD shortening fails to reach statistical significance (Fig. 7C–D, n = 4).

## 4. Discussion

To the best of our knowledge, this is the first study to demonstrate that CCM (20 mA, 20 ms) increases susceptibility to ventricular fibrillation, an effect that is dependent on β<sub>1</sub>-adrenoceptor activation and is associated with an increase in ventricular APD dispersion. In addition, we have shown that the ventricular effects of CCM are abolished during the perfusion of exogenous ACh but unaffected during muscarinic receptor antagonism and ganglionic blockade, suggesting that local intracardiac/spinal reflex pathways are not involved. These data support the notion that the acute effects of CCM are mediated through local release of NE from post-ganglionic sympathetic nerves.

#### 4.1. Recruitment of intracardiac/intrathoracic ganglia


Intracardiac and intrathoracic ganglia have been shown to be involved in the beat-to-beat control of ventricular performance independently from the central nervous system [6]. The electrical current used during CCM should be sufficient to activate ventricular afferent fibres, as suggested by experiments using electrical stimulation to identify sympathetic afferents [12], and thus a proportion of the CCM response may occur through ganglionic activation and recruitment of additional efferent sympathetic inputs. We investigated this hypothesis using the ganglionic neurotransmission inhibitor hexamethonium. Although hexamethonium abolished the bradycardic response to vagal nerve stimulation it had no effect on the acute ventricular responses to CCM. These data demonstrate that the effects of CCM occur solely through the electrical stimulation of post-ganglionic sympathetic fibres that innervate the ventricle.

#### 4.2. The role of acetylcholine

We confirm our previous finding that CCM increases ventricular contractility and shortens MAPD through β<sub>1</sub>-adrenoceptor activation following NE release [4]. These data are supported by studies demonstrating the adrenergic dependence of CCM in the goat and dog [13]. In the present study we investigated whether ACh, released from parasympathetic nerves with the ventricle, plays any role in the acute action of CCM.

It is well known that the vagus nerve antagonises the action of the sympathetic nervous system. Binding of NE to the β-adrenoceptors increases the activity of soluble adenylyl cyclase, catalysing the formation of cAMP, which activates protein kinase A to phosphorylate several protein targets increasing contractile force. ACh binds to muscarinic receptors reducing the release of NE from sympathetic nerves and inhibits the formation of cAMP within cardiac myocytes [14,15]. It is reasonable to assume that exogenous ACh perfusion prevents the acute actions of CCM through a combination of these mechanisms. In much the same way that CCM stimulates sympathetic nerves, parasympathetic nerves can also be stimulated. Despite substantial functional





**Fig. 7.** The effects of CCM on action potential duration across the anterior ventricular surface. A) Raw data demonstrating optically action potentials recorded at base and apex regions at baseline (BL) and during basal and apical CCM. B) The spatial spread of APD shortening during basal CCM from a representative experiment. D & E) Mean data demonstrating the effective range of APD shortening from the site of CCM delivery. Dotted lines denote the statistical cut off for each data set. (n = 4).

and histological evidence of parasympathetic innervation of the ventricle, atropine failed to modify the ventricular CCM response, suggesting that there was insufficient ACh release from tentative parasympathetic nerve stimulation to modulate the adrenergic component of CCM.

#### 4.3. Calcium vs. hormone hypothesis

CCM was initially proposed as a technique to increase cardiac contractility through direct modulation of calcium entry during the action potential (calcium/electrotonic hypothesis). However, our data demonstrate that application of CCM signals to the ventricle stimulates the release of NE from sympathetic nerves [4], which acts upon the myocardium through classical  $\beta$ -adrenoceptor signalling pathways. This hormone hypothesis is supported by the results of the present study, in that both metoprolol and ACh abolished the ventricular responses to CCM. Moreover,

the positive inotropic response and shortening of MAPD during CCM are in keeping with adrenergic activation. Although we did not perform experiments with calcium channel blockers, the lack of response to CCM during both metoprolol and ACh perfusion indicates that its effects on calcium, as seen in other studies [16], are secondary to adrenergic activation.

Initial experiments describing an electrotonic modulation of action potential duration by CCM were performed in the isolated rabbit papillary muscle and utilised unipolar field stimulation. These studies demonstrate that cathodic and anodic electrical stimulation have opposing influences on force and APD, presumably due to an accumulation of electrical charge at the stimulating electrode and resultant modulation of membrane potential and L-type calcium current. However, the electrotonic hypothesis predicts that opposing effects would be exerted at the anodal and cathode electrodes and that the net change in contractility during bipolar CCM could be zero (if all regions respond

comparably). Furthermore, CCM is applied in a biphasic manner and should exert equal depolarising and hyperpolarising influences at each electrode. The electrotonic hypothesis does not provide adequate explanation for the effects of CCM in whole heart and the lack of any immediate contractile effects in whole heart studies provides support for the hormone hypothesis [4,16]. Moreover, we have previously shown that CCM does not exert any effect on ventricular force during  $\beta$ -adrenoceptor blockade [4]. In the present study we show, using optical mapping, that biphasic CCM signals do not prolong the action potential but that APD shortens over a period of seconds, in keeping with the dynamics of adrenergic stimulation [17].

#### 4.4. Regional electrophysiological response to CCM

We have previously shown that the CCM induced APD shortening was partially mediated by activation of the slow delayed rectifying potassium current ( $I_{Ks}$ ) [4]. Our findings are in keeping with the classical effects of sympathetic activation on ventricular electrophysiology and are supported by other studies reporting the adrenergic dependence of CCM [13]. In the present study we confirm localised APD shortening during CCM, close to the stimulation site but not in regions that are distant. The mechanisms whereby CCM exerts a greater functional effect during basal CCM and apparently “captures” a greater region is not known although it is plausible that this reflects either the pattern of sympathetic innervation, ion channel distribution and/or the effect of coronary circulation. We have previously reported a greater expression of tyrosine hydroxylase, a surrogate marker of sympathetic nerve distribution, at basal LV myocardium of the rabbit [18]. Other investigators have reported a graded distribution of catecholamines and sympathetic nerve fibres from basal to apical regions [19,20]. It is reasonable to hypothesise that CCM captures a larger number of sympathetic nerves when applied to the basal myocardium. Although we have not measured NE release in the present study we would expect that to see a greater NE release during basal CCM than with apical CCM based on the physiological parameters. We have previously shown that NE release during CCM is dependent upon the current amplitude applied to the basal LV, likely reflecting “capture” of a greater area of the LV and thus sympathetic nerves [4]. The expression of KCNQ1, the pore forming protein subunit responsible for the adrenergic sensitive slow delayed rectifier potassium current, is also higher at the base than the apex [18] which may also contribute to the greater degree of APD shortening with CCM at the base. It is also possible that the greater distance range of CCM induced shortening during basal stimulation reflects the downstream coronary circulation transporting NE away from its release site deeper and further away from the stimulation site. Because of the location, NE release in apical regions will be distributed to a smaller proportion of the myocardium via coronary circulation.

#### 4.5. CCM and ventricular arrhythmias

In the present study we show that CCM signals (20 mA, 20 ms), applied to the LV epicardial surface of the normal rabbit heart, promote localised action potential shortening and give rise of non-uniformities of APD across the ventricular surface. We demonstrate that CCM increases apico-basal APD dispersion in a manner correlated with a downward shift in VFT and dependent upon  $\beta_1$ -adrenoceptor stimulation (being abolished by metoprolol). Furthermore, ERP is substantially shortened at the site of CCM stimulation and appears to contribute to the pro-arrhythmic phenotype, causing a greater reduction in VFT when compared to the effects of MAPD<sub>90</sub> dispersion alone. It is probable that the localised effects of CCM contribute to electrical instability during rapid pacing and that this underpins an increase in VF susceptibility. It is well known that spatial repolarisation heterogeneity provides an ideal substrate for the generation of ventricular arrhythmias [21]. Dispersion of APD is a known pro-arrhythmic mechanism underlying circus movement re-entry around refractory regions [22] and computational studies

demonstrate a key role of APD dispersion in the genesis and propagation of VF [23].

##### 4.5.1. Clinical implications

Our data demonstrate that CCM increases the susceptibility of the crystalloid perfused rabbit heart to experimentally induced VF, however it is recognised that the present study utilises healthy non-disease animal hearts and as such we are cautious of offering direct correlations with clinical application of CCM therapy in human heart failure. Further study in animal models of heart failure, failing human tissue and prospective clinical studies would be necessary to confirm our results. It is clear that CCM signals of  $\pm 20$  mA and 20 ms applied to the rabbit heart may not be directly comparable to the application of similar or smaller amplitude signals in the larger canine or human heart and these values were selected from our previous experience and not as a comparison to clinical CCM. Nevertheless, the observed effects on action potential dispersion, regional repolarisation and VFT may be of scientific and clinical interest.

Anecdotal evidence from clinical studies of CCM suggests that there is no increase in the number of serious adverse arrhythmic events with CCM. However these trials are relatively small in size, short in duration and do not provide any record on the occurrence of arrhythmic events i.e. number of premature ventricular beats, non-sustained VT events, other than those presenting in clinic i.e. sustained VT and VF [2]. Several factors including the intermittent nature of chronic CCM therapy (4–6 h per day), the interaction of CCM with long-term beneficial remodelling, and the use of  $\beta$ -blockers may influence the occurrence of arrhythmias in vivo. Exercise regimes, like CCM, are associated with intermittent sympathetic activation and are widely accepted to improve patient quality of life without affecting mortality and hospitalisation rates (during a 4 year follow up period) [24]. It is attractive to suggest that CCM, which is commonly applied intermittently for 4–6 h per day, mimics the effects of daily exercise; however exercise will have many additional effects other than an increased sympathetic tone.

Whilst exercise regimes and CCM are both adrenergic in nature, it should be noted that clinical studies of CCM have been carried out in patients on  $\beta$ -blockers. Although the degree of  $\beta$ -blockade cannot be accurately assessed in each patient, there is the possibility that the level of  $\beta$ -blockade present could counteract at least in part the adverse effects of CCM on arrhythmogenesis. Complete  $\beta$ -adrenoceptor blockade is unlikely to be achieved clinically due to the association with significant side effects with potentially high doses. Rather the goal of  $\beta$ -blockade therapy in heart failure is to reduce sympathetic over-activation. CCM has been shown to be effective (acutely and chronically) in patients receiving  $\beta$ -blockade drugs [2]. It cannot be ruled out that the long-term beneficial effects of CCM are not mediated by  $\beta_1$ -adrenoceptor signalling. It is equally plausible that the long-term beneficial effects of CCM in heart failure, for example upon molecular remodelling, offer some protection from associated arrhythmias negating the negative influence of acute CCM therapy. It is clear that more detailed study on the long-term mechanism of action of CCM is required, with a special focus on the effects of long term CCM on spontaneous ventricular arrhythmias and arrhythmias susceptibility.

#### 4.6. CCM and ventricular relaxation

As with our previous study, CCM increased LV pressure development and contractility but had little effect on ventricular relaxation. This finding is in accordance with other reports [25]. We have previously proposed [4] that the localised effects of CCM and the proportion of sympathetic fibres stimulated during CCM may be insufficient to significantly alter global relaxation. Relaxation is dependent upon both calcium reuptake into the sarcoplasmic reticulum and the passive properties of the myocardium and as such may not be a sensitive to the localised effects of CCM. We cannot discount that there may be regional

effects of CCM on relaxation that are not reflected in the more global measure reported in our study.

#### 4.7. Ventricular fibrillation threshold

The susceptibility of the heart to VF was studied by measuring the VFT, which is a quantifiable and reproducible measure reflecting electrical stability [7]. Despite controversy regarding its ability to predict the effects of some anti-arrhythmic drugs, VFT has been shown to be a powerful predictor of altered arrhythmia vulnerability in relation to the autonomic nervous system [7,26].

#### 4.8. Study limitations

##### 4.8.1. Site of CCM delivery different to clinical use and comparisons with larger species

We use epicardial LV sites for CCM delivery to ascertain the effects and mechanisms underlying CCM. Importantly, this method is in accordance with previous mechanistic studies in both the canine and ferret [33,34]. Clinically, the site of CCM delivery chosen is the RV septum [2], as it is convenient for and akin to standard cardiac device lead placement. Whilst our study differs in this regard, other investigators have reported that the effects of CCM are abolished by  $\beta$ -blockade during stimulation of the RV septum of the goat [13], in accord with our previous work [4]. The localised electrophysiological effects of CCM suggest that application in any region may promote heterogeneities of repolarisation and provide a substrate for arrhythmia.

Histological studies suggest that there exists a transmural dispersion of nerve distribution through the hearts, with less sympathetic innervation but more parasympathetic innervation of the ventricular endocardium [20]. We cannot discount that epicardial CCM captures more sympathetic fibres than the endocardial approach used clinically and that this will influence the change in VFT with CCM.

Although the spacing between our electrodes is proportionally similar to that used in larger species (e.g. dogs and humans) we cannot exclude that a 20 mA stimulus might stimulate a greater proportion of the smaller rabbit heart in relation to larger hearts. Despite this, the degree of inotropic enhancement is similar to that reported in other studies [16,25]. The parameters of our CCM signal were selected on the basis of our previous work characterising functional and electrophysiological responses with varying signal amplitude, delay and duration [4].

##### 4.8.2. Electromechanical uncoupling agent

During optical mapping we use the electromechanical uncoupler blebbistatin to prevent cardiac movement during imaging. We have recently reported that blebbistatin has a direct effect on ventricular electrophysiological parameters prolonging APD and increasing resistance to VF induction [27]. Due to the pronounced electrophysiological effects of blebbistatin, data on the effects of CCM on VFT were not collected in optical studies, rather these studies were used solely to provide confirmatory data on the regional influence of CCM, to complement our previously published findings [4] and add additional insight into the spatial distribution and dynamics of CCM effects. The mechanisms of action of blebbistatin's electrophysiological effects are not clear and require further investigation.

## 5. Conclusions

To our knowledge, this study represents the first direct investigation of the effects of CCM on ventricular arrhythmia vulnerability. Our data show that CCM ( $\pm 20$  mA, 20 ms) increases the vulnerability to ventricular fibrillation through a  $\beta 1$ -adrenoceptor dependent mechanism and is associated with APD shortening and an increase in APD dispersion (in the crystalloid perfused rabbit heart). In conjunction with our previous study, we conclude that the acute effects of CCM are mediated via

NE release from efferent postganglionic sympathetic nerves as the response to CCM can be abolished by exogenous ACh perfusion.

## Disclosures

None.

## Acknowledgements

All authors contributed to the design, planning and implementation of the experiments described in this manuscript. Data was collected and analysed by JW and KEB. All authors contributed to the drafting and preparation of the manuscript.

## References

- [1] Hjalmarson A, Goldstein S, Fagerberg B, et al. Effects of controlled-release metoprolol on total mortality, hospitalizations, and well-being in patients with heart failure: the metoprolol CR/XL randomized intervention trial in congestive heart failure (MERIT-HF). *JAMA* 2000;283:1295–302.
- [2] Winter J, Brack KE, Ng GA. Cardiac contractility modulation in the treatment of heart failure: initial results and unanswered questions. *Eur J Heart Fail* 2010;13:700–10.
- [3] Imai M, Rastogi S, Gupta RC, et al. Therapy with cardiac contractility modulation electrical signals improves left ventricular function and remodeling in dogs with chronic heart failure. *J Am Coll Cardiol* 2007;49:2120–8.
- [4] Winter J, Brack KE, Ng GA. The acute inotropic effects of cardiac contractility modulation (CCM) are associated with action potential duration shortening and mediated by beta1-adrenoceptor signalling. *JMCC* 2010;51:252–62.
- [5] Brown AM. Excitation of afferent cardiac sympathetic nerve fibres during myocardial ischaemia. *J Physiol* 1967;190:35–53.
- [6] Armour JA. Instant to instant reflex cardiac regulation. *Cardiology* 1976;61:309–28.
- [7] Ng GA, Brack KE, Patel VH, Coote JH. Autonomic modulation of electrical restitution, alternans and ventricular fibrillation initiation in the isolated heart. *Cardiovasc Res* 2007;73:750–60.
- [8] Packer M, Carver JR, Rodeheffer RJ, et al. Effect of oral milrinone on mortality in severe chronic heart failure. The PROMISE Study Research Group. *N Engl J Med* 1991;325:1468–75.
- [9] Ng GA, Brack KE, Coote JH. Effects of direct sympathetic and vagus nerve stimulation on the physiology of the whole heart—a novel model of isolated Langendorff perfused rabbit heart with intact dual autonomic innervation. *Exp Physiol* 2001;86:319–29.
- [10] Brack KE, Coote JH, Ng GA. Vagus nerve stimulation protects against ventricular fibrillation independent of muscarinic receptor activation. *Cardiovasc Res* 2011;91:437–46.
- [11] Brack KE, Patel VH, Mantravadi R, Coote JH, Ng GA. Direct evidence of nitric oxide release from neuronal nitric oxide synthase activation in the left ventricle as a result of cervical vagus nerve stimulation. *J Physiol* 2009;587:3045–54.
- [12] Fu L-W, Longhurst JC. Interactions between histamine and bradykinin in stimulation of ischaemically sensitive cardiac afferents in felines. *J Physiol* 2005;565:1007–17.
- [13] Cornelussen RN, Splett V, Klepfer RN, Stegemann B, Kornet L, Prinzen FW. Electrical modalities beyond pacing for the treatment of heart failure. *Heart Fail Rev* 2011;16:315–25.
- [14] Levy MN, Blattberg B. Effect of vagal stimulation on the overflow of norepinephrine into the coronary sinus during cardiac sympathetic nerve stimulation in the dog. *Circ Res* 1976;38:81–4.
- [15] Vincent NH, Ellis S. Inhibitory effect of acetylcholine on glycogenolysis in the isolated guinea-pig heart. *J Pharmacol Exp Ther* 1963;139:60–8.
- [16] Mohri S, Shimizu J, Mika Y, et al. Electric currents applied during refractory period enhance contractility and systolic calcium in the ferret heart. *Am J Physiol Heart Circ Physiol* 2003;284:H1119–23.
- [17] Winter J, Tanko AS, Brack KE, Ng GA. Differential cardiac responses to unilateral sympathetic nerve stimulation in the isolated innervated rabbit heart. *Auton Neurosci* 2012;166:4–14.
- [18] Mantravadi R, Gabris B, Liu T, et al. Autonomic nerve stimulation reverses ventricular repolarization sequence in rabbit hearts. *Circ Res* 2007;100:e72–80.
- [19] Angelakos ET. Regional distribution of catecholamines in the dog heart. *Circ Res* 1965;16:39–44.
- [20] Kawano H, Okada R, Yano K. Histological study on the distribution of autonomic nerves in the human heart. *Heart Vessels* 2003;18:32–9.
- [21] Han J, Moe GK. Nonuniform recovery of excitability in ventricular muscle. *Circ Res* 1964;14:44–60.
- [22] Kuo C, Munakata K, Reddy C, Surawicz B. Characteristics and possible mechanism of ventricular arrhythmia dependent on the dispersion of action potential durations. *Circulation* 1983;67:1356–67.
- [23] Clayton RH, Holden AV. Dispersion of cardiac action potential duration and the initiation of re-entry: a computational study. *Biomed Eng Online* 2005;4:11.

- [24] Flynn KE, Pina IL, Whellan DJ, et al. Effects of exercise training on health status in patients with chronic heart failure: HF-ACTION randomized controlled trial. *JAMA* 2009;301:1451–9.
- [25] Mohri S, He KL, Dickstein M, et al. Cardiac contractility modulation by electric currents applied during the refractory period. *Am J Physiol Heart Circ Physiol* 2002;282:H1642–7.
- [26] Quesada A, Angus J, Chorro J, et al. Changes in canine ventricular fibrillation threshold induced by verapamil, flecanide and bretylium. *Eur Heart J* 1993;14:712–6.
- [27] Brack KE, Narang R, Winter J, Ng GA. The mechanical uncoupler blebbistatin is associated with significant electrophysiological effects in the isolated rabbit heart. *Exp Physiol* 2013;98:1009–27.