

## Different types of physical activity are positively associated with indicators of mental health and psychological wellbeing in rheumatoid arthritis during COVID-19

Brady, Sophia M.; Fenton, Sally A. M.; Metsios, George S.; Bosworth, Ailsa; Duda, Joan L.; Kitas, George D.; Veldhuijzen van Zanten, Jet J. C. S.

DOI:

[10.1007/s00296-020-04751-w](https://doi.org/10.1007/s00296-020-04751-w)

License:

Creative Commons: Attribution (CC BY)

### Document Version

Publisher's PDF, also known as Version of record

### Citation for published version (Harvard):

Brady, SM, Fenton, SAM, Metsios, GS, Bosworth, A, Duda, JL, Kitas, GD & Veldhuijzen van Zanten, JJCS 2020, 'Different types of physical activity are positively associated with indicators of mental health and psychological wellbeing in rheumatoid arthritis during COVID-19', *Rheumatology International*, vol. 2020, pp. 1-10. <https://doi.org/10.1007/s00296-020-04751-w>

[Link to publication on Research at Birmingham portal](#)

### Publisher Rights Statement:

Brady, S.M., Fenton, S.A.M., Metsios, G.S. et al. Different types of physical activity are positively associated with indicators of mental health and psychological wellbeing in rheumatoid arthritis during COVID-19. *Rheumatol Int* (2020). <https://doi.org/10.1007/s00296-020-04751-w>

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.



# Different types of physical activity are positively associated with indicators of mental health and psychological wellbeing in rheumatoid arthritis during COVID-19

Sophia M. Brady<sup>1,2,3</sup> · Sally A. M. Fenton<sup>1,2,3</sup> · George S. Metsios<sup>2,4</sup> · Ailsa Bosworth<sup>5</sup> · Joan L. Duda<sup>1,3</sup> · George D. Kitas<sup>1,2</sup> · Jet J. C. S. Veldhuijzen van Zanten<sup>1,2,3</sup>

Received: 10 October 2020 / Accepted: 4 November 2020

© The Author(s) 2020

## Abstract

Nationwide lockdowns during SARS-CoV-2 (COVID-19) can compromise mental health and psychological wellbeing and limit opportunities for physical activity (PA), particularly in clinical populations, such as people with rheumatoid arthritis (RA), who are considered at risk for COVID-19 complications. This study aimed to investigate associations between PA and sedentary time (ST) with indicators of mental health and wellbeing in RA during COVID-19 lockdown, and examine the moderation effects of self-isolating. 345 RA patients completed an online questionnaire measuring PA (NIH-AARP Diet and Health Study Questionnaire), ST (International Physical Activity Questionnaire-Short Form), pain (McGill Pain Questionnaire and Visual Analogue Scale), fatigue (Multidimensional Fatigue Inventory), depressive and anxious symptoms (Hospital Anxiety and Depression Scale), and vitality (Subjective Vitality Scale) during the United Kingdom COVID-19 lockdown. Associations between PA and ST with mental health and wellbeing were examined using hierarchical multiple linear regressions. Light PA (LPA) was significantly negatively associated with mental fatigue ( $\beta = -.11$ ), depressive symptoms ( $\beta = -.14$ ), and positively with vitality ( $\beta = .13$ ). Walking was negatively related to physical fatigue ( $\beta = -.11$ ) and depressive symptoms ( $\beta = -.12$ ) and positively with vitality ( $\beta = .15$ ). Exercise was negatively associated with physical ( $\beta = -.19$ ) and general ( $\beta = -.12$ ) fatigue and depressive symptoms ( $\beta = -.09$ ). ST was positively associated with physical fatigue ( $\beta = .19$ ). Moderation analyses showed that LPA was related to lower mental fatigue and better vitality in people not self-isolating, and walking with lower physical fatigue in people self-isolating. These findings show the importance of encouraging PA for people with RA during a lockdown period for mental health and wellbeing.

**Keywords** Physical activity · Sedentary behaviour · Mental health · COVID-19 · Self-isolation · Rheumatoid arthritis

## Introduction

SARS-CoV-2 (COVID-19) has been declared a pandemic by the World Health Organisation (WHO) [1]. Unprecedented nationwide restrictions were put in place to limit the spread of the virus. In the United Kingdom (UK), the general population was instructed to only leave their home for basic necessities (i.e. food shopping and medical treatment),

essential work that could not be carried out at home and once a day for exercise. People considered at increased risk of serious complications following infection were advised to “self-isolate” (i.e. limit outside contact and not leave their homes). One such at risk population is Rheumatoid Arthritis (RA), which is an autoimmune disease characterised by inflammation, pain, fatigue and poor mental health, and commonly managed by immunosuppressive therapies [2, 3]. Whilst restrictions were deemed necessary to contain the spread of the virus, they can negatively impact mental health and psychological wellbeing [4], as well as behaviours which can support mental health and wellbeing such as physical activity (PA) and sedentary behaviour [5].

High levels of anxiety, depression and stress have been reported in the general population during COVID-19 restrictions [6–9]. In people living with rheumatic diseases, a

**Electronic supplementary material** The online version of this article (<https://doi.org/10.1007/s00296-020-04751-w>) contains supplementary material, which is available to authorized users.

✉ Jet J. C. S. Veldhuijzen van Zanten  
veldhujj@bham.ac.uk

Extended author information available on the last page of the article

study conducted in the United States has reported difficulty managing negative emotions, perceived increased risk of being infected and reduced access to healthcare and medications during this pandemic [10]. Such COVID-19-related concerns have been associated with poor mental health and compromised psychological wellbeing in the general population [8, 9]. As the mental health impact of COVID-19 is even greater in those self-isolating [4, 11], people living with RA, a population already at risk of compromised mental health [12], may be at even greater risk of adverse psychological impact [5]. Indeed, negative consequences of COVID-19 on mental health has been reported in 73% participants with rheumatic diseases in one study [13]. Therefore, identifying factors that could positively impact mental health during the COVID-19 pandemic is critical for establishing effective management to attenuate the negative impact of this pandemic on wellbeing.

In the general population, PA is positively associated with indicators of mental health and psychological wellbeing and reductions in PA are associated with negative mental health during COVID-19 [11]. In RA, PA is related to reduced anxiety, depression, fatigue, pain and increased vitality [3, 14–17]. However, the COVID-19 lockdown has restricted opportunities for PA and not surprisingly, lower PA and increased screen time have been reported [11, 18]. This may pose a significant risk for the mental health and psychological wellbeing of those, who are already at risk of low levels of PA and high levels of sedentary behaviour (i.e. waking activities involving sitting/lying and energy expenditure  $\leq 1.5$  metabolic equivalents), such as people with RA [19]. Indeed, during COVID-19, people with rheumatic diseases reported challenges to being active [10] and reductions in PA and increased sedentary time (ST) have been reported in RA [20], which could further worsen mental health and wellbeing [21].

Emerging evidence during COVID-19 emphasises the benefits of PA for mental health in the general population [8, 22, 23]. However, associations between PA, mental health and psychological wellbeing during COVID-19 have not been investigated in RA. The aims of this study were to (1) explore the associations between PA and ST with indicators of mental health and wellbeing in RA during COVID-19 and (2) examine the impact of self-isolation on these associations.

## Methods

### Participants

People with RA were recruited through social media of the National Rheumatoid Arthritis Society (NRAS) in the UK. Inclusion criteria were a self-reported clinical diagnosis of

RA and aged  $\geq 18$  years. Ethical approval was obtained from the University of Birmingham ethics committee (ERN\_20-0475). Participants were given an information sheet, provided informed consent and completed the online questionnaire between April 8 and April 30 2020, when the most stringent lockdown restrictions applied in the UK.

### Patient and public involvement

This online questionnaire was developed in collaboration with NRAS.

### Measures

#### Physical activity and sedentary behaviour

PA was assessed using the National Institutes of Health-American Association of Retired Persons (NIH-AARP) Diet and Health Study questionnaire to record participation in different types of PA [24]. Participants were asked to indicate how much time they had spent during the last 7 days in 3 PA types: (1) light PA (LPA) (e.g. cooking, laundry), (2) walking and (3) exercise (e.g. tennis, cycling). Items were scored using a categorical scale with response options from “none” to “ $\geq 10$  h”. Sedentary behaviour was assessed using the International Physical Activity Questionnaire-Short Form (IPAQ-SF), which comprises 2 questions. Participants are asked “during the last 7 days, how much time did you spend sitting on a”.... (1) weekday and (2) weekend day, to calculate overall weekly ST. Both questionnaires are reliable and valid measures of PA and ST in older adults [24, 25].

#### Pain

Pain experienced during the past week was assessed with the McGill Pain Questionnaire (MPQ), comprising 15 pain descriptors (e.g. “throbbing” and “tender”) on a scale from 0 (none) to 3 (severe) [26]. Participants were also asked to rate last week’s pain on a visual analogue scale (VAS), from 0 (no pain) to 10 (worst imaginable pain) [27]. The MPQ demonstrated good internal reliability in this study (Cronbach  $\alpha = .92$ ) and the MPQ and VAS have been validated in RA [28].

#### Fatigue

Fatigue during the past week was assessed using the Multidimensional Fatigue Inventory (MFI) reflecting physical, mental and general fatigue (4 items each), rated on a scale from 1 (no, that’s not true) to 5 (yes, that’s true) [29]. MFI is a validated fatigue measure used in RA, with good internal

reliability in this study (physical:  $\alpha = .78$ , mental:  $\alpha = .86$ , general:  $\alpha = .73$ ) [30].

### Anxious and depressive symptoms

Anxious (7 items) and depressive (7 items) symptoms during the past week were measured using the Hospital Anxiety and Depression Scale (HADS) [31]. Participants were asked to indicate their agreement (on a scale from 0 to 3) with each statement (e.g. “I feel tense or ‘wound-up’”). The HADS has previously shown good validity [32] and good internal reliability (anxious symptoms:  $\alpha = .87$ , depressive symptoms:  $\alpha = .83$ ) in this study.

### Subjective vitality

Vitality, a measure of positive wellbeing, experienced during the past week was measured using the Subjective Vitality Scale (SVS) [33]. Participants were asked to rate 6 statements (e.g. “I’ve been feeling energised”) on a scale from 1 (not at all true) to 7 (very true). The SVS demonstrates good reliability in this study ( $\alpha = .88$ ) and has been validated in RA [17].

### Functional disability

Participants’ functional disability was determined using the Stanford Health Assessment Questionnaire disability index (HAQ-DI), comprising 8 subscales each reflecting an activity of daily living (i.e. dressing, rising, eating, walking, hygiene, grip, reach and activities) [34]. Participants were asked to rate their ability to perform specific activities (e.g. “open car doors”) on a scale from “without any difficulty” to “unable to do” and indicate if they used aids/devices for each activity. An overall disability index score is calculated as an average of the 8 subscales. A higher score represents higher functional disability.

### General COVID-19 concern

Concerns regarding COVID-19 were measured as the extent to which participants were apprehensive about (1) testing positive for COVID-19, (2) a family member testing positive for COVID-19 and (3) not being able to receive arthritis-related medical care. Each item was scored on a scale from 1 (not concerned at all) to 5 (very concerned) and the average was calculated.

### COVID-19 living

COVID-19 Living reflected living circumstances during COVID-19, which included “self-isolating at home” (i.e. not leaving the house due to medical recommendation (i.e.

shielding) or personal concern) and “not self-isolating” (i.e. leaving the house for basic necessities, exercise and/or work).

### Data reduction and statistical analysis

The data were analysed using IBM SPSS Version 26 and checked for normality using the Kolmogorov–Smirnov test. Independent variables were PA types (LPA, walking and exercise) and ST. Dependent variables were indicators of mental health and psychological wellbeing (i.e. pain (MPQ and VAS), physical, mental and general fatigue, anxious symptoms, depressive symptoms and vitality). Covariates were age, gender, functional disability, living alone/with others, education and general COVID-19 concern, all with known associations with dependent variables in RA. The moderator variable was COVID-19 Living.

408 participants provided complete data for PA and ST. Participants were excluded due to implausible ST ( $> 18$ hours/day [35],  $n = 26$ ), missing covariate ( $n = 8$ ) or moderator data ( $n = 6$ ). For the dependent variables, the missing data were imputed using the expectation maximisation method where participants were missing one item of a questionnaire (MPQ:  $n = 24$ ; MFI:  $n = 15$ ; HADS:  $n = 1$ ; SVS:  $n = 1$ ). Participants with  $> 1$  missing value per questionnaire were excluded ( $n = 23$ ). The final sample size for all statistical analysis was 345 participants.

Differences between those self-isolating vs not self-isolating were assessed with Mann–Whitney or  $\chi^2$  tests, as appropriate. To address the primary aim, hierarchical linear regression analyses were conducted to examine associations between PA and ST with indicators of mental health and psychological wellbeing, while adjusting for potential covariates. In these hierarchical regression analyses, we explored the following sequential models:

Regression Model 1 examined the associations between the covariates (age, gender, functional disability, living alone/with others, education and general COVID-19 concern) with the indicators of mental health and psychological wellbeing as dependent variables (pain, physical, mental and general fatigue, anxious symptoms, depressive symptoms and vitality). Separate regression analyses were conducted for each indicator of mental health and wellbeing. For each regression analysis, the  $F$  value and  $p$  value are reported to reflect statistical significance, the  $R^2$  value is presented to reflect the variance in the indicator of mental health and psychological wellbeing explained by all covariates combined and standardised beta values ( $\beta$  values) are presented to reflect the direction and strength of the association between each covariate and indicator of mental health and psychological wellbeing.

Regression Model 2 explored the associations between LPA, walking, exercise or ST with each indicator of mental

health and psychological wellbeing indicator, while adjusting for the covariates included in Model 1. In other words, Regression Model 2 expanded the analyses conducted in Model 1 (with only covariates as the predictors of mental health and psychological wellbeing) to include LPA, walking, exercise or ST (independent variable) as a predictor of the indicators of mental health and psychological wellbeing (dependent variable). Separate analyses were conducted for each combination of independent and dependent variable, with the covariates included in all regression models. For each regression analysis,  $\Delta R^2$  was calculated to reflect the additional variance in the dependent variable explained by including the independent variable to the model with covariates only.  $F$  and  $p$  values are reported to reflect statistical significance of adding the independent variable to the model and  $\beta$  values reflect the direction and strength of the association between each independent (LPA, walking, exercise or ST) and dependent variable (indicator of mental health and psychological wellbeing).

Regression Model 3 explored whether the associations between the activity-related independent variables (i.e. LPA, walking or exercise) with the indicators of mental health and psychological wellbeing were independent of ST, and vice versa, whether the associations between ST and indicators of mental health and psychological wellbeing were independent of the levels of activity (LPA, walking and exercise). More specifically, where regression Model 2 revealed significant associations between LPA, walking or exercise with a specific dependent variable, Model 3 included both the significant PA type as well as ST as predictors for that dependent variable, while also adjusting for covariates. For each regression analysis,  $\Delta R^2$  was calculated to reflect the additional variance explained in the dependent variable as compared to Model 2.  $F$  and  $p$  values are reported to reflect statistical significance of adding the independent variable to the model and  $\beta$  values reflected the direction and strength of the association between each independent (LPA, walking, exercise or ST) and dependent variable (indicator of mental health and psychological wellbeing).

Finally, to explore the impact of COVID-19 living situation (i.e. self-isolating at home vs not self-isolating) on all associations between independent (LPA, walking, exercise, ST) and dependent variables (indicators of mental health and wellbeing), moderation analyses were conducted using the PROCESS model [36]. In all moderation analyses, age, gender, living situation, education, general COVID-19 concern and functional disability were included as covariates.

As the majority of variables were not normally distributed, bootstrapping was employed in regression analyses. Bootstrapping is a nonparametric re-sampling procedure that does not impose the assumption of normal distribution on the data [37]. Significance was interpreted based on the bootstrap-generated 95% bias-corrected confidence intervals

(CIs) (5000 samples). CIs also provide more information than  $p$ -values, showing the possible variability of effect size and therefore are more appropriate for determining significance of bootstrapped data [37, 38] and standardised beta coefficients ( $\beta$ ) were used to interpret the strength of associations.

## Results

Participant characteristics are reported in Table 1. The sample predominantly comprised white females, with moderate functional disability. Mann – Whitney and Chi-square tests revealed a longer disease duration, lower levels of LPA and walking, more pain, physical fatigue, functional disability and COVID-19 concern in self-isolating participants (all  $p$ 's < .05, see Table 1).

### Regression analyses

Model 1: The associations between covariates with each mental health and psychological wellbeing indicator are summarised in Table 2. As is evident from Table 2, all regression models were statistically significant. Examination of the  $\beta$  values of the covariates showed that functional disability was most strongly and consistently associated with all indicators of mental health and psychological wellbeing, with higher functional disability related to more pain, fatigue and depressive and anxious symptoms and lower vitality.

Model 2 expanded on Model 1 with the covariates only, by adding the independent variables (LPA, walking, exercise, ST) as predictors of the indicators of mental health and psychological wellbeing in separate analyses. Table 3 presents the summary findings of Model 2 regression analyses, by focusing on the additional amount of variance explained by each independent variable and the associated  $\beta$ -coefficient. More detailed information on these models, including all covariates, unstandardised beta values ( $B$ ) and the 95% CI (used to assess significance of the model), is reported in Supplementary Tables S1–S4. LPA was significantly negatively associated with mental fatigue and depressive symptoms and positively with vitality. Walking was negatively related to physical fatigue and depressive symptoms and positively with vitality. Exercise was negatively associated with physical and general fatigue and depressive symptoms. ST was positively linked to physical fatigue. No other significant associations were detected. In all these analyses, the  $\beta$ -coefficients for the covariates remained broadly similar to those reported in Table 2. More detailed information about the full regression models is presented in Supplementary Tables S1–S4.

Model 3 explored whether the associations reported between the activity-related independent variables with

**Table 1** Descriptive statistics with *p* values for total sample and sample stratified by COVID-19 Living status

	Self-isolating at home ( <i>n</i> = 230)	Not self-isolating ( <i>n</i> = 115)	All participants ( <i>n</i> = 345)	<i>p</i> value
<b>Demographic Information</b>				
Age (years)	51.53 ± 11.82	51.37 ± 11.58	51.48 ± 11.73	.805
Gender ( <i>n</i> = female (%))	214 (93.0)	107 (93.0)	321 (93.0)	1.000
Ethnicity ( <i>n</i> = white (%))	224 (97.4)	110 (96.5)	334 (96.8)	.640
RA duration (years from diagnosis)	11.51 ± 10.51	8.52 ± 8.08*	10.52 ± 9.87	.012
<b>Independent variables</b>				
LPA (minutes/week)	150 ± 420	300 ± 420*	300 ± 420	.033
Walking (minutes/week)	60 ± 150	240 ± 420*	90 ± 300	< .001
Exercise (minutes/week)	0 ± 30	0 ± 30	0 ± 30	.820
Sedentary time (minutes/week)	3360 ± 1823	3360 ± 1980	3360 ± 1680	.088
<b>Outcomes</b>				
Pain (MPQ)	15.34 ± 9.30	13.12 ± 9.02*	14.60 ± 9.26	.024
Pain (VAS rating)	4.57 ± 2.62	4.10 ± 2.58	4.41 ± 2.62	.095
Physical fatigue	15.25 ± 4.06	14.06 ± 4.15*	14.85 ± 4.12	.007
Mental fatigue	12.04 ± 5.00	11.94 ± 5.01	12.01 ± 5.00	.860
General fatigue	15.88 ± 3.63	15.31 ± 4.21	15.69 ± 3.83	.383
Anxious symptoms	9.39 ± 4.58	8.34 ± 4.51	9.04 ± 4.58	.055
Depressive symptoms	7.39 ± 3.94	6.86 ± 4.04	7.21 ± 3.98	.146
Subjective Vitality	2.56 ± 1.20	2.74 ± 1.32	2.62 ± 1.24	.266
<b>Covariates</b>				
Functional disability (HAQ-DI)	1.33 ± 0.77	0.99 ± 0.69*	1.22 ± 0.76	< .001
Living situation ( <i>n</i> = living alone (%))	43 (18.7)	17 (14.8)	60 (17.4)	.366
Education ( <i>n</i> = higher education (%))	119 (51.7)	61 (53.0)	180 (52.2)	.819
General COVID-19 Concern	3.91 ± 0.90	3.62 ± 0.90*	3.81 ± 0.91	.003

Values are reported as means ± SD, except for PA and ST variables which show medians ± IQR

Living situation was characterised as living with others (i.e. partner, family) or living alone. Education was characterised as higher education (university degree, doctorate) or secondary education (GCSE/O-level, A-level/GCE)

RA rheumatoid arthritis, LPA light physical activity, MPQ McGill Pain Questionnaire, VAS Visual Analogue Scale, HAQ-DI Health Assessment Questionnaire-Disability Index, SD standard deviation, IQR interquartile range

\*Significantly different from self-isolating at home with *p* < .05. Differences were examined using Mann–Whitney *U* and  $\chi^2$  tests, as appropriate

the dependent variables were independent of levels of ST and vice versa. When adding ST as an additional predictor, all significant associations between activity-related independent variables with the indicators of mental health and psychological wellbeing observed in Model 2 remained significant, with the exception of the association between walking with physical fatigue, which no longer remained significant when ST was added to the model ( $\beta = -.08$ ,  $B = -0.001$ , 95% CI =  $-3.1 \cdot 10^{-3}$ ,  $1.5 \cdot 10^{-4}$ ). In both walking and exercise models, ST was significantly associated with physical fatigue (walking model:  $\beta = .18$ ,  $B < 0.001$ , 95% CI =  $2.8 \cdot 10^{-4}$ ,  $8.3 \cdot 10^{-4}$ ; exercise model:  $\beta = .18$ ,  $B < 0.001$ , 95% CI =  $2.9 \cdot 10^{-4}$ ,  $8.2 \cdot 10^{-4}$ ).

## Moderation analyses

Moderation analyses revealed that COVID-19 living situation only moderated the associations between LPA with

mental fatigue and vitality and walking with physical fatigue. More LPA was significantly associated with lower mental fatigue and better vitality in those who were not self-isolating (mental fatigue model:  $\beta = -.26$ , 95% CI =  $-.42$ ,  $-.09$ ; vitality model:  $\beta = .31$ , 95% CI =  $.15$ ,  $.47$ ), but not in those who were self-isolating (mental fatigue model:  $\beta = -.04$ , 95% CI =  $-.16$ ,  $.10$ ; vitality model:  $\beta = .03$ , 95% CI =  $-.09$ ,  $.16$ ). Walking was associated with lower physical fatigue in people who were self-isolating ( $\beta = -.22$ , 95% CI =  $-.35$ ,  $-.08$ ), but not in those not self-isolating ( $\beta = -.02$ , 95% CI =  $-.15$ ,  $.12$ ). For more detailed information of moderation analysis results, please see Supplementary Table S5.

**Table 2** Model 1 Regression Analyses for all covariates with each indicator of mental health and psychological wellbeing (dependent variable)

	Pain (MPQ)	Pain (VAS rating)	Physical Fatigue	Mental Fatigue	General Fatigue	Anxious Symptoms	Depressive Symptoms	Vitality
	$R^2 = .383$	$R^2 = .330$	$R^2 = .348$	$R^2 = .140$	$R^2 = .260$	$R^2 = .207$	$R^2 = .219$	$R^2 = .176$
	$F = 34.91$	$F = 27.73$	$F = 30.09$	$F = 9.15$	$F = 19.78$	$F = 14.67$	$F = 15.77$	$F = 12.05$
	$p < .001$	$p < .001$	$p < .001$	$p < .001$	$p < .001$	$p < .001$	$p < .001$	$p < .001$
	$\beta$	$\beta$	$\beta$	$\beta$	$\beta$	$\beta$	$\beta$	$\beta$
Age	-.11*	-.08	-.11*	-.27*	-.21*	-.25*	-.16*	.18*
Gender	-.01	-.02	.05	-.06	-.03	-.01	.01	-.03
Education	-.03	.01	.05	-.04	.03	-.10*	-.04	.06
Living Situation	.02	.04	.01	-.10*	.01	.02	-.09	.07
Concern	.15*	.08	.09	.10	.04	.32*	.15*	-.08
Functional Disability	.55*	.55*	.55*	.23*	.46*	.14*	.38*	-.34*

Model 1: Regressions included all covariates (age, gender, education, living situation, general COVID-19 concern and functional disability) as predictors for each indicator of mental health and wellbeing in separate analyses

$R^2$  represents the variance explained in the dependent variable by all covariates together. Statistical information about each model is presented by the  $F$ -value and the  $p$ -value, with  $\beta$  representing the standardised beta coefficient of each covariate

Concern general COVID-19 concern, MPQ McGill Pain Questionnaire, VAS Visual Analogue Scale, CI confidence interval

\*Significant associations between covariates with indicators of mental health and wellbeing derived using bootstrapped 95% CI

**Table 3** Summary Model 2 Regression Analyses for LPA, Walking, Exercise and Sedentary Time with dependent variables adjusting for covariates

	Pain (MPQ)		Pain (VAS rating)		Physical Fatigue		Mental Fatigue		General Fatigue		Anxious Symptoms		Depressive Symptoms		Vitality	
	$\beta$	$\Delta R^2$	$\beta$	$\Delta R^2$	$\beta$	$\Delta R^2$	$\beta$	$\Delta R^2$	$\beta$	$\Delta R^2$	$\beta$	$\Delta R^2$	$\beta$	$\Delta R^2$	$\beta$	$\Delta R^2$
LPA	-.02	.000	-.06	.003	-.08	.006	-.11*	.011†	-.07	.005	-.04	.002	-.14*	.018†	.13*	.016†
Walking	.04	.002	.06	.003	-.11*	.010†	-.01	.000	-.06	.003	.02	.000	-.12*	.012†	.15*	.020†
Exercise	.04	.001	.01	.000	-.19*	.033†	-.04	.001	-.12*	.013†	-.04	.001	-.09*	.008	.07	.005
Sedentary Time	-.08	.006	-.02	.000	.19*	.032†	.04	.001	.07	.005	-.04	.001	.08	.006	-.09	.008

Model 2: expanded Model 1 (covariates age, gender, education, living situation, general COVID-19 concern and functional disability only) by adding LPA, walking, exercise and ST as individual predictors for each dependent variable.

$\beta$  represents the standardised beta coefficient and  $\Delta R^2$  represents the proportion of the variance that is explained by the addition of the predictor independent variable to the model relative to Model 1.  $\beta$  coefficients for covariates were broadly similar as those reported in Table 1. Therefore, to improve readability,  $\beta$ 's are only reported for the associations between the independent variables (LPA, walking, exercise, ST) and the dependent variables. Full information of all Model 2 regressions, including the 95% CI intervals to determine significance,  $F$ -value with associated  $p$ -values and the  $\beta$  for each covariate are reported in Supplementary Tables S1-S4

LPA light physical activity, Concern general COVID-19 concern, MPQ McGill Pain Questionnaire, VAS Visual Analogue Scale, CI confidence interval

\*significant  $\beta$  (standardised beta) coefficients derived using bootstrapped 95% CI (see Supplementary Tables S1-S4); † $p < .05$  for  $\Delta R^2$  values determining the significance of the overall model

## Discussion

This is the first study to show associations between activity behaviours with indicators of mental health and psychological wellbeing in people with RA during COVID-19. LPA and walking were associated with lower physical and mental fatigue and depressive symptoms and higher vitality.

Exercise was related to lower physical and general fatigue and fewer depressive symptoms and ST was related to higher physical fatigue. In addition, COVID-19 living situation moderated some associations between LPA and walking with physical and mental fatigue and vitality.

The finding that LPA and walking were associated with higher vitality is in line with epidemiological research demonstrating a relationship between LPA with wellbeing in

older adults [39]. Our results point to the importance of LPA and walking for wellbeing in RA during COVID-19. From a behaviour change perspective, encouraging non-exercise LPA (e.g. household chores) and walking may be perceived as more feasible and accessible for RA, a population experiencing significant disease-related barriers to PA [40].

PA was associated with fewer depressive symptoms, aligned with the previous arthritis research [2, 15]. Associations were of a similar, although opposite, magnitude to associations between COVID-19 concerns and depressive symptoms. Fear or concern about the virus has been related to depression [4], and our findings suggest that PA counteracts this negative impact on depressive symptoms in RA, in line with findings in college students and older adults [22, 41]. Importantly, associations between PA with depressive symptoms and vitality were independent of functional disability in this study and others [2]. Thus, the activity at any intensity should be promoted in all people with RA to improve mental health and wellbeing, regardless of functional disability.

PA was associated with lower and ST with higher physical and mental fatigue, in line with interventions promoting PA and reducing ST improving fatigue in RA [3, 42, 43]. Our findings emphasise the importance of the multidimensional aspects of fatigue in RA. Specifically, LPA was negatively associated with mental fatigue, whereas walking, exercise and ST were related to physical fatigue. Our results suggest that different PA types could be related to different aspects of fatigue in people living with RA. Aligned with present findings, exercise interventions have been reported to be particularly effective for physical fatigue in RA [30].

PA can lead to improved mental health and wellbeing during lockdown situations through several pathways. For example, PA can distract from negative thoughts and worries [44], being active can have an immediate positive effect on mood [44], outdoor environment can induce mental stimulation [23] and PA can provide structure when daily routine is disrupted due to lockdown, with the resulting sense of control improving wellbeing [45]. Thus, recommendations promoting PA may offer an avenue to support clinical populations to cope with the impact of COVID-19 on their mental health [41, 46]. Given that maintaining PA during COVID-19 is associated with better mental health in older adults [41], the present results imply that increasing PA may positively impact mental health and wellbeing in RA during COVID-19. Given the duration of the pandemic and recent return of restrictions, longitudinal studies during COVID-19 are needed to understanding how changes in PA contribute to better mental health and wellbeing in RA. This can inform guidance on management of wellbeing during these difficult times, not just in RA, but also other clinical populations.

PA and ST were not associated with anxious symptoms in this study. There are mixed findings related to anxiety and PA during COVID-19 [22, 23, 47]. The impact of PA on anxiety during COVID-19 may be influenced by prior activity levels. Indeed, research suggests inactive people who increased PA during COVID-19 reported lower anxiety compared to those who became less active, but these associations were not seen in people who were classed as active prior to COVID-19 [23]. Thus, perhaps instead of looking at absolute values, it might be more important to examine changes in PA during a pandemic in relation to anxiety.

People with RA in this study did not report associations between PA and ST with pain, which is in contrast to previous observational studies of RA [48]. This may be due to the possible bi-directional association between PA with mental health and wellbeing [49] and also the association between pain and functional disability [50], which may have affected levels of PA observed depending on COVID-19 living status. Specifically, correlational analysis revealed walking was negatively related to pain (data not reported), but this association was no longer significant when adjusting for functional disability. Given our findings suggested that people with RA who were self-isolating had higher levels of pain and functional disability and lower levels of LPA and walking, compared to those leaving the house, it could be assumed that individuals with the least pain and disability were more likely to leave the house for PA. This lower variability in pain among those who were accruing some form of PA (through leaving the house), could mean associations between PA and pain are less likely to be observed.

The lower levels of PA observed among people with RA who were self-isolating, is in agreement with previous studies [11]. Moderation analysis showed LPA was only related to mental fatigue and vitality in those not self-isolating, whereas walking was only associated with physical fatigue in those self-isolating. As those self-isolating did significantly less walking than those not self-isolating, this could suggest that walking specifically, should be encouraged among individuals self-isolating to reduce physical fatigue. However, as few significant moderation effects of COVID-19 living were observed, these findings should be interpreted with care.

Except for physical fatigue, ST was not associated with indicators of mental health or wellbeing, contrasting previous COVID-19 research in the general population [11]. Our measure of ST reflected total ST and did not differentiate between different sedentary behaviours; e.g. sitting while being intellectually stimulated, e.g. during work, is suggested to have a less negative impact health and wellbeing compared to sitting watching TV or using electronic devices [51]. Consequently, future studies are required to understand the specific role of different sedentary behaviours for mental



health and wellbeing in RA, both during and beyond the pandemic.

The current study included a large sample of people with RA during stringent lockdown conditions. Therefore, it was only possible to collect self-report data for all outcomes, which should be acknowledged as a limitation. In addition, the associations reported are cross-sectional, so causality cannot be inferred. Therefore, exploring these associations over time during COVID-19 will help to better understand the implications of this pandemic on the link between activity behaviours and mental health and wellbeing and how PA can support mental health and wellbeing throughout the pandemic.

In summary, PA, specifically LPA and walking, was positively associated with mental health and psychological wellbeing in RA during COVID-19. These findings support recommendations from different governments to encourage PA during lockdown restrictions, to attenuate the negative impact of a pandemic on mental health and wellbeing. Given the known barriers for PA in RA [40] and the reported additional barriers experienced during COVID-19 [10], these findings emphasise the importance of appropriate support and recommendations for PA in people with RA and potentially other clinical populations, particularly in those self-isolating, during a pandemic to maintain mental health and psychological wellbeing.

**Acknowledgements** The authors would like to thank the National Rheumatoid Arthritis Society (NRAS) for their support in the development of this study and the recruitment for this study. In addition, the authors would like to thank all participants for their time to complete to questionnaire pack.

**Author contributions** Conception and design of the study: SB, SF, GM, AB, GK, JVvZ. Data acquisition: SB, SF, AB, JVvZ. Data analysis: SB, SF, JVvZ. Data interpretations and drafting of manuscript: all authors. Final approval of manuscript: all authors.

**Funding** This work was completed as part of a PhD studentship supported by the Medical Research Council (MRC)-Versus Arthritis Centre for Musculoskeletal Ageing Research (CMAR) (grant number: MR/K00414X/1).

## Compliance with ethical standards

**Conflict of interest** The authors declare no conflicts of interest.

**Ethics approval** This study was performed in line with the principles of the Declaration of Helsinki. Ethical approval was obtained from the University of Birmingham ethics committee (ERN\_20-0475).

**Consent to participate and consent to publish** Informed consent was obtained from all individual participants included in the study. The authors affirm that human research participants provided informed consent for publication.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

1. WHO. Coronavirus disease (COVID-2019) situation reports. Coronavirus disease (COVID-2019) situation reports 2020 [cited 2020 July 2020]; Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>
2. Fenton SAM, Veldhuijzen van Zanten JJCS, Metsios GS, Rouse PC, Yu CA, Kitas GD et al (2018) Autonomy support, light physical activity and psychological well-being in rheumatoid arthritis: a cross-sectional study. *Mental Health Phys Act* 14:11–18
3. Rongen-van Dartel S, Repping-Wuts H, Flendrie M (2015) Effect of aerobic exercise training on fatigue in rheumatoid arthritis: a meta-analysis. *Arthritis Care Res* 67(8):1054–1062
4. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N et al (2020) The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet* 395(10227):912–920
5. Veldhuijzen van Zanten JJ, Fenton SAM, Brady S, Metsios GS, Duda JL, Kitas GD (2020) Mental health and psychological wellbeing in rheumatoid arthritis during COVID-19—can physical activity help? *Mediterr J Rheumatol* 31(Suppl 2):284–287
6. Huang Y, Zhao N (2020) Generalized anxiety disorder, depressive symptoms and sleep quality during COVID-19 outbreak in China: a web-based cross-sectional survey. *Psychiatry Res* 288:112954
7. Lai J, Ma S, Wang Y, Cai Z, Hu J, Wei N et al (2020) Factors associated with mental health outcomes among health care workers exposed to Coronavirus Disease 2019. *JAMA Netw Open* 3(3):e203976
8. Rodriguez-Rey R, Garrido-Hernansaiz H, Collado S (2020) Psychological impact and associated factors during the initial stage of the Coronavirus (COVID-19) pandemic among the general population in Spain. *Front Psychol* 11:1540
9. Harper C, Satchell L, Fido D, Latzman R (2020) Functional fear predicts public health compliance in the COVID-19 pandemic. *Int J Ment Health Addict*. <https://doi.org/10.1007/s11469-020-00281-5>
10. Michaud K, Wipfler K, Shaw Y, Simon T (2020) Experiences of patients with rheumatic diseases in the united states during early days of the COVID-19 pandemic. *ACR Open Rheumatol* 2(6):335–343
11. Meyer J, McDowell C, Lansing J, Brower C, Smith L, Tully M et al (2020) Changes in physical activity and sedentary behavior in response to COVID-19 and their associations with mental health in 3052 US adults. *Int J Environ Res Public Health* 17(18):E6469
12. Fiest K, Hitchon CA, Bernstein C, Peschken C (2017) Systematic review and meta-analysis of interventions for depression and

- anxiety in persons with rheumatoid arthritis. *J Clin Rheumatol* 23(8):425–434
13. Ziade N, El Kibbi L, Hmamouchi I, Abdulateef N, Halabi H, Hamdi W, et al (2020) Impact of the COVID-19 pandemic on patients with chronic rheumatic diseases: a study in 15 Arab countries. *Int J Rheum Dis*. <https://doi.org/10.1111/1756-185X.13960>
  14. Chekroud SR, Gueorguieva R, Zheutlin AB, Paulus M (2018) Association between physical exercise and mental health in 1.2 million individuals in the USA between 2011 and 2015: a cross-sectional study. *Lancet Psychiatry* 5(9):739–746
  15. Kelley GA, Kelley KS, Hootman JM (2015) Effects of exercise on depression in adults with arthritis: a systematic review with meta-analysis of randomized controlled trials. *Arthritis Res Ther* 17(1):21
  16. Stenstrom CH (1994) Home exercise in rheumatoid arthritis functional class II: goal setting versus pain attention. *J Rheumatol* 21(4):627–634
  17. Rouse PC, Veldhuijzen Van Zanten JJ, Ntoumanis N, Metsios GS, Yu CA, Kitas GD et al (2015) Measuring the positive psychological well-being of people with rheumatoid arthritis: a cross-sectional validation of the subjective vitality scale. *Arthritis Res Ther* 17:312
  18. Pépin JL, Bruno RM, Yang R-Y, Vercamer V, Jouhaud P, Escourrou P et al (2020) Wearable activity trackers for monitoring adherence to home confinement during the COVID-19 pandemic worldwide: data aggregation and analysis. *J Med Internet Res*. 22(6):e19787
  19. Fenton SA, Kitas GD (2016) Rheumatoid arthritis: sedentary behaviour in RA—a new research agenda. *Nat Rev Rheumatol* 12(12):698–700
  20. Pinto AJ, Rezende D, Siczekowska SM, Meireles K, Bonfiglioli K, Ribeiro ACM, et al (2020) Increased prolonged sitting in rheumatoid arthritis patients during the COVID-19 pandemic: a within-subjects, accelerometer-based study. medRxiv 2020.09.09.20191395
  21. Fenton SAM, Veldhuijzen van Zanten J, Duda JL, Metsios GS, Kitas GD (2018) Sedentary behaviour in rheumatoid arthritis: definition, measurement and implications for health. *Rheumatology (Oxford)* 57(2):213–226
  22. Zhang Y, Zhang H, Ma X, Di Q (2020) Mental Health Problems during the COVID-19 Pandemics and the Mitigation Effects of Exercise: a Longitudinal Study of College Students in China. *Int J Environ Res Public Health* 17(10):3722
  23. Lesser IA, Nienhuis CP (2020) The impact of COVID-19 on physical activity behavior and well-being of Canadians. *Int J Environ Res Public Health* 17(11):3899
  24. Gierach GL, Chang SC, Brinton LA, Lacey JV Jr, Hollenbeck AR, Schatzkin A et al. (2009) Physical activity, sedentary behavior, and endometrial cancer risk in the NIH-AARP Diet and Health Study. *Int J Cancer* 124(9):2139–2147
  25. Cleland C, Ferguson S, Ellis G, Hunter R (2018) Validity of the International Physical Activity Questionnaire (IPAQ) for assessing moderate-to-vigorous physical activity and sedentary behaviour of older adults in the United Kingdom. *BMC Med Res Methodol* 18(176):176
  26. Melzack R (1987) The short-form McGill pain questionnaire. *Pain* 30(2):191–197
  27. Scott J, Huskisson E (1976) Graphic representation of pain. *Pain* 2:175–184
  28. Van Lankveld W, Pad Bosch P, Van de Putte L, Van der Staak C, Naring G (1992) Pain in rheumatoid arthritis measured with the visual analogue scale and the Dutch version of the McGill Pain Questionnaire. *Ned Tijdschr Geneesk* 136(24):1166–1170
  29. Smets EM, Garssen B, Bonke B, Haes J (1995) The Multidimensional Fatigue Inventory (MFI) psychometric qualities of an instrument to assess fatigue. *J Psychosom Res* 39(3):315–325
  30. Rupp I, Boshuizen HC, Jacobi CE, Dinant HJ, van den Bos GA (2004) Impact of fatigue on health-related quality of life in rheumatoid arthritis. *Arthritis Rheum* 51(4):578–585
  31. Zigmond AS, Snaith RP (1983) The hospital anxiety and depression scale. *Acta Psychiatr Scand* 67(6):361–370
  32. Covic T, Pallant J, Tennant A, Cox S, Emery P, Conaghan PG (2009) Variability in depression prevalence in early rheumatoid arthritis: a comparison of the CES-D and HAD-D Scales. *Bmc Musculoskelet Disorders* 10:18
  33. Ryan RM, Frederick C (1997) On energy, personality, and health: Subjective vitality as a dynamic reflection of well-being. *J Pers* 65(3):529–565
  34. Kirwan JR, Reeback JS (1986) Stanford health assessment questionnaire modified to assess disability in british patients with rheumatoid arthritis. *Br J Rheumatol* 25(2):206–209
  35. Loppenthin K, Esbensen BA (2015) Sleep quality and correlates of poor sleep in patients with rheumatoid arthritis. *Clin Rheumatol* 34:2029–2039
  36. Hayes AF (2012) PROCESS: a versatile computational tool for observed variable mediation, moderation, and conditional process modeling [White Paper]. Retrieved from: <http://www.afhayes.com/public/process2012.pdf>
  37. Preacher KJ, Hayes AF (2008) Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behav Res Methods* 40(3):879–891
  38. du Prel J-B, Hommel G, Röhrig B, Blettner M (2009) Confidence interval or p-value?: part 4 of a series on evaluation of scientific publications. *Dtsch Arztebl Int* 106(19):335–339
  39. Buman MP, Hekler EB, Haskell WL, Pruitt L, Conway TL, Cain KL et al (2010) Objective light-intensity physical activity associations with rated health in older adults. *Am J Epidemiol* 172(10):1155–1165
  40. Veldhuijzen van Zanten JJ, Rouse PC, Hale ED, Ntoumanis N, Metsios GS, Duda JL et al (2015) Perceived barriers, facilitators and benefits for regular physical activity and exercise in patients with rheumatoid arthritis: a review of the literature. *Sports Med*. 45(10):1401–1412
  41. Carriedo A, Cecchini JA, Fernandez-Rio J, Mendez-Gimenez A (2020) COVID-19, psychological well-being and physical activity levels in older adults during the nationwide lockdown in Spain. *Am J Geriatr Psychiatry* 28(11):1146–1155
  42. Katz P, Margaretten M, Gregorich S, Trupin L (2018) Physical activity to reduce fatigue in rheumatoid arthritis: a randomized controlled trial. *Arthritis Care Res* 70(1):1–10
  43. Thomsen T, Aadahl M, Beyer N, Hetland ML, Loppenthin K, Midtgaard J et al (2017) The efficacy of motivational counselling and SMS reminders on daily sitting time in patients with rheumatoid arthritis: a randomised controlled trial. *Ann Rheum Dis* 76(9):1603–1606
  44. Mikkelsen K, Stojanovska L, Polenakovic M (2017) Exercise and mental health. *Maturitas* 106:48–56
  45. Ryan RM, Deci EL (2017) Self-determination theory: basic psychological needs in motivation, development, and well-ness. Guildford Press, New York
  46. Matias T, Dominski FH, Marks DF (2020) Human needs in COVID-19 isolation. *J Health Psychol* 25(7):871–882
  47. Antunes R, Frontini R, Amaro N, Salvador R, Matos R, Morouco P et al (2020) Exploring lifestyle habits, physical activity, anxiety and basic psychological needs in a sample of portuguese adults during COVID-19. *Int J Environ Res Public Health* 17(12):4360
  48. Hakkinen A, Sokka T, Kotaniemi A, Hannonen P (2001) A randomized two-year study of the effects of dynamic strength training on muscle strength, disease activity, functional capacity, and bone mineral density in early rheumatoid arthritis. *Arthritis Rheum* 44(3):515–522

49. Lwin MN, Serhal L, Holroyd C, Edwards CJ (2020) Rheumatoid arthritis: the impact of mental health on disease: a narrative review. *Rheumatol Ther* 7(3):457–471
50. Luyster FS, Chasens ER, Wasko MCM, Dunbar-Jacob J (2011) Sleep quality and functional disability in patients with rheumatoid arthritis. *J Clin Sleep Med* 7(1):49–55
51. Saidj M, Jorgensen T, Jacobsen RK, Linneberg A, Aadahl M (2014) Differential cross-sectional associations of work- and leisure-time sitting, with cardiorespiratory and muscular fitness among working adults. *Scand J Work Environ Health* 40(5):531–538

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

## Affiliations

Sophia M. Brady<sup>1,2,3</sup> · Sally A. M. Fenton<sup>1,2,3</sup> · George S. Metsios<sup>2,4</sup> · Ailsa Bosworth<sup>5</sup> · Joan L. Duda<sup>1,3</sup> · George D. Kitas<sup>1,2</sup> · Jet J. C. S. Veldhuijzen van Zanten<sup>1,2,3</sup> 

Sophia M. Brady  
sxb1075@student.bham.ac.uk

Sally A. M. Fenton  
s.a.m.fenton@bham.ac.uk

George S. Metsios  
g.metsios@wlv.ac.uk

Ailsa Bosworth  
ailsa@nras.org.uk

Joan L. Duda  
j.l.duda@bham.ac.uk

George D. Kitas  
George.kitas@nhs.net

<sup>1</sup> School of Sport, Exercise and Rehabilitation Sciences, University of Birmingham, Birmingham B15 2TT, UK

<sup>2</sup> Rheumatology Department, Dudley Group NHS Foundation Trust, Dudley, UK

<sup>3</sup> Medical Research Council-Versus Arthritis Centre for Musculoskeletal Ageing, University of Birmingham, Birmingham, UK

<sup>4</sup> Faculty of Education, Health and Wellbeing, University of Wolverhampton, Wolverhampton, UK

<sup>5</sup> National Rheumatoid Arthritis Society, Maidenhead, UK