

Editorial on plants as alternative hosts for human and animal pathogens

Holden, Nicola J.; Jackson, Robert W.; Schikora, Adam

DOI:

[10.3389/fmicb.2015.00397](https://doi.org/10.3389/fmicb.2015.00397)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Holden, NJ, Jackson, RW & Schikora, A 2015, 'Editorial on plants as alternative hosts for human and animal pathogens', *Frontiers in Microbiology*, vol. 6, 397. <https://doi.org/10.3389/fmicb.2015.00397>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Editorial on plants as alternative hosts for human and animal pathogens

Nicola J. Holden^{1*}, Robert W. Jackson² and Adam Schikora³

¹ Human and Animal Pathogen - Plant Interactions, The James Hutton Institute, Dundee, UK, ² School of Biological Sciences, University of Reading, Reading, UK, ³ Institute of Phytopathology and Applied Zoology, Justus Liebig University of Giessen, Giessen, Germany

Keywords: *Salmonella enterica*, *Escherichia coli*, *Arabidopsis thaliana*, fresh produce, PAMP-triggered immunity, effectors, mRNA extraction, microbiome

Fresh produce, in the form of raw or minimally processed fruits and vegetables, is recognized to be an important source of food-borne disease. Sporadic cases or outbreaks can arise from bacterial, viral, or parasitic contamination. Importantly, while the latter both require animal hosts for proliferation and cannot grow on or within crop plants, bacteria can do so and are able to use plants as secondary hosts. This type of interaction means that human pathogenic bacteria need to undergo adaptation to the plant host, which presents an environment quite distinct to animals or humans. Important differences lie in both physiochemical and biological properties, e.g., physiology, immunity, native microflora, physical barriers, mobility, and temperature. There is now good published evidence to describe different aspects of the interactions between plants and human pathogenic bacteria, and the subsequent effects on the outcome of colonization. However, in comparison to the interactions between the pathogens and their animal hosts, for which we know a great deal about both host and microbial factors, this area is relatively new with many important knowledge gaps. Research in this area is of most relevance to food safety, since the results can be and indeed, are already applied by public health agencies and food producers. It also reveals fascinating aspects of the bacterial life-cycle that were hitherto under appreciated.

One of the important features of plant host colonization is the adaptation of pathogens to the host defense response. Innate immunity is the first line of defense with both physical barriers and biological recognition of human pathogens, and the different layers of the response are reviewed in Melotto et al. (2014). An intriguing aspect is the finding that some human pathogens are able to subvert the plant defense, shown using *Salmonella enterica* and *Arabidopsis* as an example (Garcia and Hirt, 2014). Perception of human pathogens is based on recognition of PAMPs (e.g., flagellin) similar to those in phytopathogens and other plant-associated bacteria; perhaps unsurprising given that the mechanisms of recognition tend to be based on evolutionary conserved proteins. What was unexpected was the fact that some *Salmonella* serovars encode flagellin variants that are no longer recognized by the plant, presumably as a result of evolutionary selection (Garcia and Hirt, 2014).

There are good parallels in microbial suppression of the innate response of animal and plant hosts, for example via effectors secreted through the type 3 secretion (T3S) apparatus. Despite structural differences in the apparatus between human- and phyto-pathogens, the general function of host subversion is shared. Commonalities (and differences) in which pathways in animal and plant hosts are targeted by human pathogens are highlighted in Brunner and Fraiture (2014), showing where there is evolutionary conservation in the cellular hubs. Detailed examination of the function for one of the T3S effectors from *S. enterica*, SpvC, shows that it targets a signaling pathway that is shared in plant and animal hosts (Neumann et al., 2014). This effector is a phosphothreonine lyase able to de-phosphorylate activated MAP kinases, therefore attenuating the immune response in *Arabidopsis*.

OPEN ACCESS

Edited by:

Giulia De Lorenzo,
Sapienza Università di Roma, Italy

Reviewed by:

Simone Ferrari,
Sapienza Università di Roma, Italy

*Correspondence:

Nicola J. Holden,
nicola.holden@hutton.ac.uk

Specialty section:

This article was submitted to
Plant-Microbe Interaction,
a section of the journal
Frontiers in Microbiology

Received: 25 February 2015

Accepted: 17 April 2015

Published: 30 April 2015

Citation:

Holden NJ, Jackson RW and Schikora
A (2015) Editorial on plants as
alternative hosts for human and animal
pathogens. *Front. Microbiol.* 6:397.
doi: 10.3389/fmicb.2015.00397

Examination of mechanisms of colonization shows that although some aspects are broadly the same, there are important differences in the interaction between *S. enterica* and either plant or animal hosts (Wiedemann et al., 2015). To better understand the molecular basis of the interaction, it is necessary to determine the protein-protein interactions (PPIs) between the pathogen and the plant host. An important comparison of PPIs between *Salmonella* and *Arabidopsis* proteins to known interactions between bacterial and animal proteins was presented in Schleker et al. (2015). In order to characterize the interactions on a global scale, an approach was used that combines several new as well as previously described algorithms predicting new PPIs (Kshirsagar et al., 2015). However, while this approach yields candidates worthy of further investigation, it also serves to highlight the paucity of reported functional data. To further explore the molecular interactions, methods must be developed that are appropriate to the bacteria-plant system, for example for the analysis of bacterial gene expression. An improved method for the extraction of RNA from mixed samples was presented in Holmes et al. (2014), which allows the analysis of expression patterns in bacteria and the plant host at the same time.

Colonization of plant hosts by human pathogens rarely, if ever, occurs in isolation and successful colonization is dependent on the ability of the bacteria to compete with the native microflora. Microbiome approaches show that crops not obviously associated with human pathogens may harbor potential pathogens. As shown with an example of grapevines, crop plants can support

the endophytic growth of human pathogen, in this case *Propionibacterium acnes*, which probably arose from human contamination in the first place (Yousaf et al., 2014). Application of bulky organic fertilizers, such as manure, to crops may increase the chances of transmission of bacteria into the food chain. However, the potential for transmission clearly shows some specificity, not least from differences between species and sub-species of bacteria and the host plants (Hofmann et al., 2014). A combination of axenic system with susceptible plants (e.g., spinach) revealed that 400 bacteria in one ml are sufficient to successfully colonize the crop plant.

This collection of articles helps to highlight the underpinning mechanisms, but also shows the complexity of interactions between human pathogenic bacteria and plant hosts. It is becoming clear that it is not possible to apply a broad set of rules to these interactions: there is a great deal of specificity that depends on multiple factors. This is particularly important in the consideration of how to address issues related to food safety, for practices applied at the pre-harvest stage or for prevention of transmission and contamination during post-harvest processing.

Acknowledgments

NH is supported by Scottish Government's Rural and Environment Science and Analytical Services Division AS is supported by the Federal Office for Agriculture and Food (BLE), Germany.

References

- Brunner, F., and Fraiture, M. (2014). Killing two birds with one stone: trans-kingdom suppression of PAMP/MAMP-induced immunity by T3E from enteropathogenic bacteria. *Front. Microbiol.* 5:320. doi: 10.3389/fmicb.2014.00320
- Garcia, A. V., and Hirt, H. (2014). *Salmonella enterica* induces and subverts the plant immune system. *Front. Microbiol.* 5:141. doi: 10.3389/fmicb.2014.00141
- Hofmann, A., Fischer, D., Hartmann, A., and Schmid, M. (2014). Colonization of plants by human pathogenic bacteria in the course of organic vegetable production. *Front. Microbiol.* 5:191. doi: 10.3389/fmicb.2014.00191
- Holmes, A., Birse, L., Jackson, R. W., and Holden, N. (2014). An optimised method for the extraction of bacterial mRNA from plant roots infected with *Escherichia coli* O157:H7. *Front. Microbiol.* 5:286. doi: 10.3389/fmicb.2014.00286
- Kshirsagar, M., Schleker, S., Carbonell, J., and Klein-Seetharaman, J. (2015). Techniques for transferring host-pathogen protein interactions knowledge to new tasks. *Front. Microbiol.* 6:36. doi: 10.3389/fmicb.2015.00036
- Melotto, M., Panchal, S., and Roy, D. (2014). Plant innate immunity against human bacterial pathogens. *Front. Microbiol.* 5:411. doi: 10.3389/fmicb.2014.00411
- Neumann, C., Fraiture, M., Hernández-Reyes, C., Akum, F. N., Virlogeux-Payant, I., Chen, Y., et al. (2014). The *Salmonella* effector protein SpvC, a phosphothreonine lyase is functional in plant cells. *Front. Microbiol.* 5:548. doi: 10.3389/fmicb.2014.00548
- Schleker, S., Kshirsagar, M., and Klein-Seetharaman, J. (2015). Comparing human-*Salmonella* with plant-*Salmonella* protein-protein interaction predictions. *Front. Microbiol.* 6:45. doi: 10.3389/fmicb.2015.00045
- Wiedemann, A., Virlogeux-Payant, I., Chaussé, A.-M., Schikora, A., and Velge, P. (2015). Interactions of *Salmonella* with animals and plants. *Front. Microbiol.* 5:791. doi: 10.3389/fmicb.2014.00791
- Yousaf, S., Bulgari, D., Bergna, A., Pancher, M., Quaglino, F., Casati, P., et al. (2014). Pyrosequencing detects human and animal pathogenic taxa in the grapevine endosphere. *Front. Microbiol.* 5:327. doi: 10.3389/fmicb.2014.00327

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Holden, Jackson and Schikora. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.