

Collaboration and infrastructure is needed to develop an African perspective on micro(nano)plastic pollution

Nel, Holly A; Naidoo, Trishan; Akindele, Emmanuel O; Nhiwatiwa, Tamuka; Fadare, Oluniyi O; Krause, Stefan

DOI:

[10.1088/1748-9326/abdaeb](https://doi.org/10.1088/1748-9326/abdaeb)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Nel, HA, Naidoo, T, Akindele, EO, Nhiwatiwa, T, Fadare, OO & Krause, S 2021, 'Collaboration and infrastructure is needed to develop an African perspective on micro(nano)plastic pollution', *Environmental Research Letters*, vol. 16, no. 2, 021002. <https://doi.org/10.1088/1748-9326/abdaeb>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

PERSPECTIVE • OPEN ACCESS

Collaboration and infrastructure is needed to develop an African perspective on micro(nano)plastic pollution

To cite this article: Holly A Nel *et al* 2021 *Environ. Res. Lett.* **16** 021002

View the [article online](#) for updates and enhancements.

ENVIRONMENTAL RESEARCH
LETTERS

PERSPECTIVE

OPEN ACCESS

PUBLISHED
29 January 2021

Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Collaboration and infrastructure is needed to develop an African perspective on micro(nano)plastic pollution

Holly A Nel¹ , Trishan Naidoo² , Emmanuel O Akindele³ , Tamuka Nhiwatiwa⁴ , Oluniyi O Fadare⁵ and Stefan Krause^{1,6}

¹ School of Geography, Earth and Environmental Sciences, University of Birmingham, Edgbaston, Birmingham B15 2TT, United Kingdom

² School of Life Sciences, University of KwaZulu-Natal Westville Campus, Biology Building on University Road, Private Bag X54001, Durban 4000, South Africa

³ Department of Zoology, Obafemi Awolowo University, Ile-Ife, Nigeria

⁴ University of Zimbabwe, PO MP167 Mt. Pleasant, Harare, Zimbabwe

⁵ State Key Laboratory of Environmental Chemistry and Ecotoxicology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, 18 Shuangqing Road, Beijing 100085, People's Republic of China

⁶ LEHNA- Laboratoire d'ecologie des hydrosystemes naturels et anthropiques, University of Lyon, Lyon, France

E-mail: h.a.nel@bham.ac.uk**Keywords:** Pan-African, networks, research nodes, inclusivity, capacity buildingSupplementary material for this article is available [online](#)**Abstract**

Our current understanding of environmental micro(nano)plastic (MNP) pollution is driven by field and lab-based studies performed predominantly by and in wealthier countries. However, mismanaged waste and its consequences affect low- and middle-income countries over-proportionately. Evidence suggests that studies on MNP pollution in Africa are critically limited by the scientific infrastructure available, restricting research activities to opportunities for external collaboration with established research laboratories in the Global North. The development of Pan-African research networks and analytical nodes, is required to support intra-African research exchange and training, and drive evidence-based policy relevant to an African context. This can facilitate more inclusive conversations around the harmonization and standardization of methods currently mainly available to the Global North.

1. Introduction

While the global rise in plastic pollution represents one of the most pressing environmental challenges of our lifetime, some of the poorest countries in the world, many of them located on the African continent, are affected over-proportionally by the consequences of plastic waste mismanagement. Despite our current efforts, Borrelle *et al* (2020) suggests that by 2030, global plastic emissions may reach 53 million metric tons per year, an increase from 8 million metric tons in 2010. The state of plastic pollution in parts of Africa is dramatic, as a result of weak policies and indiscriminate dumping of solid waste (Bundhoo 2018). Five African countries (i.e. Algeria, Egypt, Morocco, Nigeria and South Africa) are listed among the top 20 countries estimated to contribute the most to marine plastic pollution (Jambeck *et al* 2015). Africa's contribution to plastic pollution

is expected to increase due to predicted population growth, rapid urbanization and the shift to a more plastic dominated consumer market (Jambeck *et al* 2018). The environmental impacts of plastic pollution in Africa will likely be accelerated, especially if plans to flood 'poorer' countries with plastic waste are to go ahead (Tabuchi *et al* 2020). This perspective will focus on the smaller size fractions, namely microplastic (≤ 5 mm) and nanoplastic (≤ 1 μ m) particles (MNP). Although these particles may contribute only a small fraction, by weight, to the total global plastic pollution crisis (Ryan *et al* 2020), their overall abundance and potential toxicity, to humans, animals and plants, increase as sizes decrease (see for example Choi *et al* (2020)). Research looking at the full size range of plastic pollution is needed (see for example Ryan *et al* (2020)), however, specific hurdles faced when investigating MNP are discussed here especially as the availability, accessibility and affordability

of technologies required for accurate quantification decreases as we shift from 'larger' to 'smaller' debris. In consequence, quantitative MNP research faces specific challenges and limitations that are different to investigations of macroplastics for instance.

While globally, MNP research has increased exponentially in the last few decades, with over 3800 articles published between 2005 and 2020. Many studies come from specific geographical regions, with the top contributors being the People's Republic of China (22%), USA (13%), Germany (10%), England (9%) and Italy (8%). In contrast, South Africa, Nigeria, Tunisia, Egypt, Kenya, and other African countries, cumulatively contributed a mere 3% (ISI Web of Knowledge, accessed in 2020). Given the sheer magnitude of the environmental plastic pollution problem in Africa, and the fact that microplastic research on the continent began in the 1980s (Alimi *et al* 2021), one should expect a higher contribution.

2. Challenges MNP research in Africa is currently facing

Most knowledge on MNPs in the environment have been based on studies performed outside of Africa, in areas that are likely to differ in their sources, concentrations, and types of MNP pollution. Associated surface contaminants may also differ, as well as degradation, fouling, and aggregation rates of MNPs, due to different environmental and anthropogenic factors (Shahul Hamid *et al* 2018). As these factors influence exposure mechanisms, one can also expect ecological toxicity to vary (Krause *et al* 2021). To advance MNP research in Africa we need to provide analytical capacity that enables investigation of the specific geographical context of MNP sources, pathways, and mechanisms. For example, indiscriminate dumping of solid waste often occurs along rivers in low- and middle-income countries (Bundhoo 2018). However, the role these sites play in the contribution of MNPs to the aquatic environment is under-studied. The burning of plastic items may also be a major MNP source rather than a termination point of plastic pollution (Yang *et al* 2021). The African blind spot of MNP research causes a critical information gap that urgently needs to be closed, as scientists call for the generation of accurate and comparable datasets that are regionally relevant and simultaneously contribute to our global understanding of MNP transport and fate through environmental compartments.


Compared to other environmental contaminants, the investigation of MNP pollution is still in its infancy, with scientific standards for analysis, reporting, and interpretation only now being developed. In many respects, the current formative phase of defining and agreeing on guidelines, standards, and protocols will shape the way MNP research is conducted in the future. It is hence paramount to ensure that widespread access to agreed protocols and standards will

be available to enable and even foster inclusivity of the most affected regions.

Our ability to study plastic pollution across wider size fractions, with higher precision, level of detail, and sample throughput, is driven by analytical advances over the last couple of years. It is, therefore, obvious that access to analytical facilities is instrumental for advancing MNP research, especially considering the range of scientific equipment and reagents needed to accurately quantify MNPs in various matrices. However, identifying particles <0.5 mm requires state-of-the-art technologies (Shim *et al* 2017) that at present do not appear to be available without external partnerships (figure 1 and table S1 (available online at stacks.iop.org/ERL/16/021002/mmedia)). There were only two publications without external partnerships that claimed to have local capacity to detect particles <0.5 mm, which could not be verified here as detection limits of the used analytical equipment were not reported (figure 1 and table S1).

As corroborated by Naidoo *et al* (2020a) MNP research in developing countries is often critically impeded by the lack of access to instrumentation. As a result of this, we see large fractions of scientific papers by African labs being published without polymer identification (Alimi *et al* 2021), with authors frequently citing the lack of access to relevant equipment as the main reason. Polymer identification has, however, become a reporting standard when investigating environmental MNP pollution and is increasingly seen as a requirement for publication. This is because polymer identification confirms visual accuracy and offers insight into possible pollution sources.

To overcome access restrictions, African researchers successfully collaborate with established research laboratories outside the continent. For instance, Akindele *et al* (2019) investigated microplastic contamination associated with gastropods in Nigeria, with polymer identification being supported through the German Academic Exchange Service (DAAD). Similarly, Naidoo *et al* (2020b) were supported by the Association of Commonwealth Universities, Blue Charter Fellowship to examine microplastics within juvenile fish associated with mangroves in South Africa. While these represent successful examples of initiatives to train the next generation of early career researchers, those African researchers are restricted in their ability to apply their newly gained knowledge when confronted with a lack of access to the required analytical facilities back home, locking them into a dependency to Northern hosts if they want to perform polymer identification. The inaccessibility of analytical techniques, frequently available to researchers from established Northern MNP laboratories, leaves African MNP scholars isolated, and reliant on gaining access to external facilities; a concept of scientific dependence that has relevance beyond the research of plastic pollution (Hountondji 1990).



3. Perspective on possible solutions

Advances in analytical techniques are key to progress, and scholars with the available resources should continue to evolve the field. However, given the complexity and geographical context of MNP pollution, the global research landscape would benefit from solutions that better include the perspectives and experiences of regions with limited infrastructural and resource capacity. Here we attempt to initiate this conversation, while outlining recommendations that may assist scholars outside of geographical MNP ‘hubs’ to accelerate internal research and method development. Our main goal is for governments and funding agencies to acknowledge the need to develop networks in Africa where Pan-African collaboration is fostered. While our secondary goal is to make the global MNP community aware that analytical advances and conversations surrounding harmonization need to include regions where

data, expertise, and infrastructure is lacking, and where climatic conditions may adversely affect equipment developed in more temperate regions. Through inclusive and innovative conversations surrounding research and data requirements, scholars in low to middle-income countries can contribute towards a more holistic understanding of the problem within different climatic and socio-economic conditions.

To date, collaboration between scholars, working on MNP pollution, across African countries is minimal (figure 1). This is largely due to a lack of opportunity, since the need for access to scientific equipment for polymer identification currently requires collaborations with research laboratories outside Africa. While the results of these mainly bi-lateral collaborations provide initial insights into site-specific MNP pollution, the lack of lateral Pan-African integration and coordination limits a systematic analysis and interpretation of the Africa specific context

(i.e. relating findings to respective pollution sources, transport and fate mechanisms, and associated impacts).

We advocate the development of a Pan-African network of research groups and facilities with dedicated plastic research programmes. Based on experiences from networks established in the UK (Microplastic Network, www.ukcnp.co.uk/uk-microplastics-network/) and Germany (Plastic Network—PLA-NET, www.plastic-network.org/), this Pan-African network should facilitate the sharing of facilities and expertise to foster research on MNP in the African context. Networking can facilitate interdisciplinary, cross-sectoral, and international collaboration, coordination of training and capacity building, and the development of multi-national, multi-institutional research strategies. A development of such magnitude will, however, require support by regional and global agents in the field (e.g. the African Union, African Development Bank, World Bank, and the United Nations Environment Programme) to initiate a development that can be sustained over the long-term.

A Pan-African network would not only require the establishment of adequate laboratories and research institutions, but also provide access to critical research infrastructure by supporting free movement of personnel between research groups and providing a long-term strategy ensuring facility maintenance (Oman *et al* 2006). Therefore, government and funding agencies should discuss innovative ways to encourage multi-national collaborations within Africa; while continuing partnerships with more established MNP laboratories globally. Lateral exchange programmes and training workshops are imperative, whereby visiting researchers are able to facilitate the development of African laboratories, while simultaneously obtaining a more inclusive understanding of site-specific drivers of plastic pollution. For example, in 2019 representatives from Centre for Environment, Fisheries and Aquaculture Science (Cefas), UK established a microplastic research facility, in South Africa, as part of the Commonwealth Litter Programme (CLiP) by providing infrastructure, training, and access to standard operating procedures (www.cefas.co.uk/clip/locations/africa/). This allowed scholars previously unable to visit established facilities the opportunity to develop the skills needed to set-up regional spaces for MNP research.

A first step towards establishing a Pan-African research network could be to collate an open-access database of expertise and equipment available nationally, regionally or across the continent, as well as a data sharing portal specific to Africa. Such information would provide critical evidence of core competences and interests, laying the foundation for where to establish collaborative research nodes. These collaborative research nodes will then

become inclusive facilities whereby researchers can visit to perform various steps along the MNP analysis pipeline, access training workshops and discuss innovative pathways for future research with like-minded scholars. This will include training on requirements for contamination control in laboratories and the development of ‘clean rooms’ for analyses which follow international standards. Learning from TREND Africa (<https://trendinafrica.org/about/>) an NGO committed to creating a scientifically independent Africa, we need to increase access to infrastructure and training while encouraging do-it-yourself (DIY) method development through the ideology of open scientific hardware. To date, none of the equipment used for polymer identification or their replaceable parts are manufactured on the continent, making routine and mandatory maintenance of such facilities difficult and cost-intensive. Research nodes should therefore disseminate vital knowledge on the development and use of low-cost extraction tools (Nel *et al* 2019), screening techniques (Maes *et al* 2017, Nel *et al* 2021) and identification keys while also providing access to state-of-the-art analytical techniques either housed on-site or through external partnerships.

4. Conclusion

We, therefore, call for global investment into analytical research infrastructure and the establishment of a collaborative Pan-African research network that will enable the step change in African MNP research that gives justice to the critical and complex challenges caused by MNP pollution. To support the bottom-up development of such Pan-African networks we are instigating a series of international webinars on MNP research in Africa that will be held online in 2021.

Data availability statements

All data that support the findings of this study are included within the article (and any supplementary files).

Acknowledgments

The authors thank Chantal Jackson for her assistance in creating figure 1, and the editors and anonymous reviewers for their valuable comments during the review process.

Author contribution

All co-authors contributed to the drafting of the manuscript, and all have approved the final version.

Conflict of interest

The authors declare that they have no known competing financial interests or personal relationships that

could have appeared to influence the work reported in this paper.

Funding

The authors would like to acknowledge funding provided by the Leverhulme Trust Leverhulme Trust research grants (RPG-2017-377) awarded to S K; and by the Institute for Global Innovation—University of Birmingham (ID: 3020) awarded to H A N.

ORCID iDs

Holly A Nel  <https://orcid.org/0000-0002-0571-2678>

Trishan Naidoo  <https://orcid.org/0000-0003-2004-6228>

Emmanuel O Akindele  <https://orcid.org/0000-0002-4911-8857>

Tamuka Nhiwatiwa  <https://orcid.org/0000-0002-5295-0724>

Oluniyi O Fadare  <https://orcid.org/0000-0002-8712-0448>

Stefan Krause  <https://orcid.org/0000-0003-2521-2248>

References

- Akindele E O, Ehlers S M and Koop J H E 2019 First empirical study of freshwater microplastics in West Africa using gastropods from Nigeria as bioindicators *Limnologica* **78** 9
- Alimi O S, Fadare O O and Okoffo E D 2021 Microplastics in African ecosystems: current knowledge, abundance, associated contaminants, techniques, and research needs *Sci. Total Environ.* **755** 142422
- Borrelle S B et al 2020 Predicted growth in plastic waste exceeds efforts to mitigate plastic pollution *Science* **369** 1515
- Bundhoo Z M A 2018 Solid waste management in least developed countries: current status and challenges faced *J. Mater. Cycles Waste Manage.* **20** 1867–77
- Choi J S, Hong S H and Park J-W 2020 Evaluation of microplastic toxicity in accordance with different sizes and exposure times in the marine copepod *Tigriopus Japonicus* *Mar. Environ. Res.* **153** 104838
- Hountondji P 1990 Scientific dependence in Africa today *Res. Afr. Lit.* **21** 5–15
- Jambeck J R, Geyer R, Wilcox C, Siegler T R, Perryman M, Andrady A, Narayan R and Law K L 2015 Plastic waste inputs from land into the ocean *Science* **347** 768
- Jambeck J et al 2018 Challenges and emerging solutions to the land-based plastic waste issue in Africa *Mar. Policy* **96** 256–63
- Krause S et al 2021 Gathering at the top? Environmental controls of microplastic uptake and biomagnification in freshwater food webs *Environ. Pollut.* **268** 115750
- Maes T, Jessop R, Wellner N, Haupt K and Mayes A G 2017 A rapid-screening approach to detect and quantify microplastics based on fluorescent tagging with Nile red *Sci. Rep.* **7** 10
- Naidoo T, Rajkaran A and Sershen 2020a Impacts of plastic debris on biota and implications for human health: a South African perspective *South Afr. J. Sci.* **116** 43–50
- Naidoo T, Sershen , Thompson R C and Rajkaran A 2020b Quantification and characterisation of microplastics ingested by selected juvenile fish species associated with mangroves in KwaZulu-Natal, South Africa *Environ. Pollut.* **257** 8
- Nel H A, Chetwynd A J, Kelleher L, Lynch I, Mansfield I, Margenat H, Onoja S, Goldberg Oppenheimer P, Sambrook Smith G H and Krause S 2021 Detection limits are central to improve reporting standards when using Nile red for microplastic quantification *Chemosphere* **263** 127953
- Nel H, Krause S, Sambrook Smith G H and Lynch I 2019 Simple yet effective modifications to the operation of the sediment microplastic isolation unit to avoid polyvinyl chloride (PVC) contamination *MethodsX* **6** 2656–61
- Oman C B, Gamaniel K S and Addy M E 2006 Properly functioning scientific equipment in developing countries *Anal. Chem.* **78** 5273–6
- Ryan P G, Weideman E A, Perold V and Moloney C L 2020 Toward balancing the budget: surface macro-plastics dominate the mass of particulate pollution stranded on beaches *Front. Mar. Sci.* **7** 929
- Shahul Hamid F, Bhatti M S, Anuar N, Mohan P and Periathamby A 2018 Worldwide distribution and abundance of microplastic: how dire is the situation? *Waste Manage. Res.* **36** 873–97
- Shim W J, Hong S H and Eo S E 2017 Identification methods in microplastic analysis: a review *Anal. Methods* **9** 1384–91
- Tabuchi H, Corkery M and Mureithi C 2020 Big oil is in trouble. its plan: flood Africa with plastic *The New York Times*
- Yang Z, Lü F, Zhang H, Wang W, Shao L, Ye J and He P 2021 Is incineration the terminator of plastics and microplastics? *J. Hazard Mater.* **401** 123429