

# Should we establish standards of care for management of elderly patients with inflammatory bowel disease?

Ghosh, Subrata; Iacucci, Marietta

DOI:  
[10.1155/2015/862458](https://doi.org/10.1155/2015/862458)

License:  
Creative Commons: Attribution-NonCommercial (CC BY-NC)

*Document Version*  
Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*  
Ghosh, S & Iacucci, M 2015, 'Should we establish standards of care for management of elderly patients with inflammatory bowel disease?', *Canadian Journal of Gastroenterology and Hepatology*, vol. 29, no. 5, 862458, pp. 236-236. <https://doi.org/10.1155/2015/862458>

[Link to publication on Research at Birmingham portal](#)

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

## Should we establish standards of care for management of elderly patients with inflammatory bowel disease?

Subrata Ghosh MBBS MD FRCPC FRCP FRCPE, Marietta Iacucci MD PhD

Elderly patients with inflammatory bowel disease (IBD) have poorer outcomes than their younger counterparts with increased inpatient mortality rates, infection rates, lymphoma rates (1), and intolerance to immunosuppressive agents or tumour necrosis factor inhibitors (TNFi) (2). These may lead to hesitancy in prescribing effective immunosuppressive drugs and TNFi to elderly patients, especially in individuals with comorbid conditions. Overuse of corticosteroids, however, has significant deleterious effects in elderly patients including osteoporosis, hypertension, diabetes, ocular complications, infections and postsurgical complications. Some elderly patients with IBD may also have very indolent disease and it is important not to overtreat them.

Several studies, including the report by Stepaniuk et al (3) in the current issue of the *Journal* (pages 273-277), have clearly shown that elderly patients receive less TNFi and less immunosuppressive drugs than their younger counterparts with IBD (4). While whether such undertreatment contributes to poorer outcome and complications in elderly patients still requires robust evidence, it is also of concern that elderly patients may be treated and monitored less rigorously and objective end points, such as mucosal healing, may never be reached. Introduction of newer therapies, such as the integrin inhibitor vedolizumab, may offer potentially certain safety advantages. TNFi may also be considered in this population as monotherapy rather than combination therapy in the presence of comorbid conditions such as chronic obstructive airways disease. Rapid control of more severe disease is essential to prevent increased morbidity and complications in elderly individuals.

Several areas of management of IBD in elderly patients require specific attention (Box 1). These include establishment of diagnosis by robust exclusion of several conditions such as colon cancer, infections, especially *Clostridium difficile*, ischemic colitis, diverticular colitis and angiodysplasia. Monitoring by regular use of C-reactive protein, fecal calprotectin, complete blood count and liver function tests is necessary to avoid the necessity of frequent invasive investigations and avoid hematological or other complications. Transabdominal ultrasonography for distal ileal Crohn disease may be a very useful noninvasive monitoring modality because it is well tolerated. Endoscopic surveillance to detect dysplasia may be especially challenging in elderly patients due to the high prevalence of diverticular disease and polypoid lesions leading to difficulty in deciding on local endoscopic resection strategies (5) versus panproctocolectomy with potentially increased morbidity. It is also important to vaccinate elderly patients against common infective preventable diseases and be aware of the high incidence of herpes zoster. The use of steroid-sparing agents, such as azathioprine, methotrexate and TNFi, is important in elderly patients to avoid excessive use of steroids in moderate to severe IBD and ensure mucosal healing. Finally, inpatients need careful prophylaxis against thromboembolism, vigilance against nosocomial infections and surgery if required in a timely manner before advanced disease, poor nutrition and infections lead to postoperative complications and poor outcome including mortality. Elderly patients are often troubled by fecal incontinence after pouch surgery and a permanent ileostomy is often a preferred approach. However, elderly

patients with adequate sphincter function may do well after pouch surgery. Multidisciplinary management of these elderly patients is especially important given the comorbid conditions that are often present.

While there are many guidelines and an increasing number of standards of care to optimize management of IBD, there are no specific guidelines for comprehensive management of elderly patients with IBD. This appears to be an omission that requires attention by national and international organizations. Many of the issues in the elderly population also require further generation of evidence because these patients form only a small fraction of patients in clinical trials.

### BOX 1

#### Specific issues influencing management of IBD in elderly patients

**Differential diagnosis:** Distinguish among ischemic colitis, colon cancer, diverticular colitis, *Clostridium difficile*, lymphocytic colitis, angiodysplasia.

**Management:** Variation in care compared with nonelderly patients, especially lower use of immunosuppressive drugs and TNFi. Risk for poor control of disease.

**Complications:** Increased risk for infections, lymphoma, herpes zoster, mortality after surgery. Fecal incontinence after ileal-pouch anal anastomosis.

**Dysplasia surveillance:** Increased prevalence of colonic polypoid lesions and challenge to determine local resection versus panproctocolectomy.

**Challenges to therapy:** Comorbid conditions such as heart failure, diabetes, hypertension, chronic obstructive lung disease, concurrent or previous malignancy, drug interactions, thromboembolism and osteoporosis.

### REFERENCES

1. Beaugerie L, Brousse N, Bouvier AM, et al. Lymphoproliferative disorders in patients receiving thiopurines for inflammatory bowel disease: a prospective observational cohort study. *Lancet* 2009;374:1617-25.
2. Desai A, Zator ZA, de Silva P, et al. Older age is associated with higher rate of discontinuation of anti-TNF therapy in patients with inflammatory bowel disease. *Inflamm Bowel Dis* 2013;19:309-15.
3. Stepaniuk P, Bernstein CN, Nugent Z, Singh H. Characterization of inflammatory bowel disease in hospitalized elderly patients in a large central Canadian health region. *Can J Gastroenterol Hepatol* 2015;29:273-7.
4. Juneja M, Baidoo L, Schwartz MB, Barrie A, Regueiro M, Dunn M, Binion DG. Geriatric inflammatory bowel disease: Phenotype presentation, treatment patterns, nutritional status, outcomes, and comorbidity. *Dig Dis Sci* 2012; 57:2408-2415.
5. Iacucci M, Uraoka T, Fort Gasia M, Yahagi N. Novel diagnostic and therapeutic techniques for surveillance of dysplasia in patients with inflammatory bowel disease. *Can J Gastroenterol Hepatol* 2014;28:361-70.

Department of Medicine, Division of Gastroenterology, University of Calgary, Calgary, Alberta

Correspondence: Dr Subrata Ghosh, Division of Gastroenterology, Department of Medicine, University of Calgary, 3280 Hospital Drive Northwest, Calgary, Alberta T2N 4N1. Telephone 403-944-8222, fax 403-944-1095, e-mail [subrata.ghosh@albertahealthservices.ca](mailto:subrata.ghosh@albertahealthservices.ca)

Received and accepted for publication April 2, 2015



**Hindawi**  
Submit your manuscripts at  
<http://www.hindawi.com>

