UNIVERSITY^{OF} BIRMINGHAM University of Birmingham Research at Birmingham

The role of the environment in transmission of Dichelobacter nodosus between ewes and their lambs

Muzafar, Mohd; Calvo-Bado, Leo A.; Green, Laura E.; Smith, Edward M.; Russell, Claire L.; Grogono-Thomas, Rosemary; Wellington, E. M. H.

DOI: 10.1016/j.vetmic.2015.04.010

License: Creative Commons: Attribution (CC BY)

Document Version Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Muzafar, M, Calvo-Bado, LA, Green, LÉ, Smith, EM, Russell, CL, Grogono-Thomas, R & Wellington, EMH 2015, 'The role of the environment in transmission of Dichelobacter nodosus between ewes and their lambs', *Veterinary Microbiology*, vol. 179, no. 1-2, pp. 53-59. https://doi.org/10.1016/j.vetmic.2015.04.010

Link to publication on Research at Birmingham portal

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Contents lists available at ScienceDirect

Veterinary Microbiology

journal homepage: www.elsevier.com/locate/vetmic



Mohd Muzafar^a, Leo A. Calvo-Bado^a, Laura E. Green^a, Edward M. Smith^a, Claire L. Russell^{b,1}, Rose Grogono-Thomas^b, Elizabeth M.H. Wellington^{a,*}

^a School of Life Sciences, University of Warwick, Gibbet Hill Road, Coventry CV4 7AL, UK
 ^b Department of Clinical Veterinary Sciences, University of Bristol, Langford House, Langford BS40 5DU, UK

ARTICLE INFO

Article history: Received 26 December 2014 Received in revised form 10 April 2015 Accepted 12 April 2015

Keywords: D. nodosus Footrot Pathogen transmission Disease reservoirs

ABSTRACT

Dichelobacter nodosus (D. nodosus) is the essential causative agent of footrot in sheep. The current study investigated when D. nodosus was detectable on newborn lambs and possible routes of transmission. Specific qPCR was used to detect and quantify the load of D. nodosus in foot swabs of lambs at birth and 5–13 h post-partum, and their mothers 5– 13 h post-partum; and in samples of bedding, pasture, soil and faeces. D. nodosus was not detected on the feet of newborn lambs swabbed at birth, but was detected 5-13 h after birth, once they had stood on bedding containing naturally occurring D. nodosus. Multiple genotypes identified by cloning and sequencing a marker gene, pgrA, and by multi locus variable number tandem repeat analysis (MLVA) of community DNA from swabs on individual feet indicated a mixed population of *D. nodosus* was present on the feet of both ewes and lambs. There was high variation in pgrA tandem repeat number (between 3 and 21 repeats), and multiple MLVA types. The overall similarity index between the populations on ewes and lambs was 0.45, indicating moderate overlap. Mother offspring pairs shared some alleles but not all, suggesting lambs were infected from sources(s) other than just their mother's feet. We hypothesise that D. nodosus is transferred to the feet of lambs via bedding containing naturally occurring populations of *D. nodosus*, probably as a result of transfer from the feet of the group of housed ewes. The results support the hypothesis that the environment plays a key role in the transmission of D. nodosus between ewes and lambs.

© 2015 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

1. Introduction

Footrot is an economically important disease of sheep. The aerotolerant anaerobe *Dichelobacter nodosus* (*D. nodosus*) is the essential causative agent (Beveridge, 1941) and

(Elizabeth M.H. Wellington).

Fusobacterium necrophorum has been suggested as a secondary bacterium after the development of disease (Beveridge, 1941; Witcomb et al., 2014). The disease is present worldwide and accounts for annual losses of between £24 and £84 million to the UK sheep industry alone (Nieuwhof and Bishop, 2005; Wassink et al., 2010). The severity of ovine footrot can vary from mild interdigital dermatitis (synonymous with benign footrot in Australian research) to virulent footrot causing severe under-running of the hoof horn with separation from the underlying tissue (Stewart, 1989). *D. nodosus* can be detected on the feet of sheep with no sign of disease (Calvo-Bado et al., 2011b; Witcomb et al., 2014) but the load is higher both before and

http://dx.doi.org/10.1016/j.vetmic.2015.04.010





CrossMark

^{*} Corresponding author at: School of Life Sciences, University of Warwick, Gibbet Hill Road, Coventry CV4 7AL, UK. Tel.: +44 02476 523184; fax: +44 02476 522052.

E-mail address: E.M.H.Wellington@warwick.ac.uk

¹ Present address: Imerys Ltd., Par Moor Centre, Par Moor Rd, Par, Cornwall, PL24, UK.

^{0378-1135/© 2015} The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/ 4.0/).

during episodes of interdigital dermatitis and virulent footrot than on healthy feet (Witcomb et al., 2014).

Temporal clustering of footrot between mothers and offspring was observed in a state transition study of factors associated with development of, and recovery from, footrot. Given that families cluster spatially this suggests spatiotemporal transmission of *D. nodosus* between family members (Kaler et al., 2010). D. nodosus has been isolated from pasture and barns where sheep are kept, indicating that contamination of the environment occurs (Witcomb, 2012). Contaminated holding areas have also been shown to cause disease in sheep put into such environments up to 2 weeks from initial seeding (Beveridge, 1941; Whittington, 1995). Recent work has indicated that D. nodosus can survive up to 14 days at 5 °C in soil, and at least 24 days when hoof material was present (Cederlof et al., 2013) and under certain conditions, D. nodosus has survived for at least 40 days in soil microcosms (unpublished data), however, further work is required to determine if survival is at a dose that could cause disease in sheep.

Multiple strains of *D. nodosus* detected by serogroup typing have been reported to co-exist in individual feet during subclinical and clinical infections (Claxton et al., 1983; Hindmarsh and Fraser, 1985; Jelinek et al., 2000; Moore et al., 2005). Molecular detection of strain differences is now possible using typing the *pgr* locus and by MLVA of *D. nodosus* (Calvo-Bado et al., 2011a; Russell et al., 2014).

The aims of this study were to investigate whether *D. nodosus* was present on the feet of newborn lambs at or after birth and the potential role played by the environment in pathogen transmission.

2. Materials and methods

2.1. Selection of animals

In April 2011 10 ewes with no clinical signs of disease and one lamb per ewe were convenience selected from a flock of 99 Mule and Suffolk crossbred ewes. Ewes were housed on the 28th March 2011, and samples collected on the 1st–6th April 2011 (Supplementary Table 1). Lambs were born in a large communal straw bedded pen, ewes and their lambs were moved to individual pens once the ewe had given birth to all her lambs. Sampled lambs were marked with tape so they could be identified for subsequent sampling.

2.2. Collection of environmental and foot swab samples

Environmental samples were taken in March prior to lambing and included swab samples of 30 fresh hoof prints in soil, four soil samples from the area around water containers, 10 samples of faecal material on the ground and compacted in the interdigital space and three straw samples collected from the storage area. In April, 10 straw bedding samples were collected from the communal pen where pregnant ewes were housed. All samples were stored at 4 °C for transportation and at -80 °C until analysed. All four feet of each lamb was swabbed using sterile cotton swabs (EUROTUBO collection swab; Delta lab, Rubi, Spain) directly after birth and before the lamb touched the ground. The lamb and its dam were sampled 5–13 h later once the lamb had stood and been transferred, with its mother, to an individual pen. Swabs were stored at 4 °C for transportation and at -80 °C on arrival at the laboratory.

2.3. Detection limit assay by direct PCR and nested PCR from swabs

The *D. nodosus* strain VCS1703A was used as a positive control for all PCR reactions. To determine the PCR detection limits, cells were harvested from a 5 d culture grown on 2% hoof agar, and 10-fold serial dilutions $(10^{-1} \text{ to } 10^{-10})$ were made in triplicate in sterile phosphate buffered saline (PBS). The numbers of cells in the initial concentration and all dilutions were counted using a haemocytometer. Sterile swabs were inoculated with 500 µl of each dilution, and frozen at $-20 \,^{\circ}$ C to produce swabs containing a known bacterial load. Microbial DNA was extracted from swabs as described below and the DNA used to determine assay detection limits.

2.4. DNA extraction from swabs

Total genomic DNA was extracted using the NucleoSpin Tissue Kit (Macherey-Nagel, GmbH and Co, Düren, Germany) with modifications. Swabs were thawed at 4 °C and 400 µl of buffer T1 was added followed by 40 µl of proteinase K. The samples were vortexed twice for 5 s and incubated for 10 min at 56 °C. The mixtures were transferred to microcentrifuge tubes and 400 µl of buffer B3 was added. The samples were vortexed twice for 5 s and incubated for 5 min at 70 °C then allowed to cool before adding 400 µl of 100% ethanol. The samples were again vortexed twice and the supernatant transferred to a NucleoSpin Tissue column and centrifuged at $11,000 \times g$ for 1 min. The flow-through was discarded, the membrane was washed with 500 µl of buffer B5 and centrifuged at $11,000 \times g$ for 1 min. The flow-through was again discarded, the column was washed with 600 µl of buffer B5 and centrifuged at $11,000 \times g$ for 1 min. The flow-through was again discarded and the membrane dried by centrifugation at $11,000 \times g$ for 1 min to remove residual ethanol. The DNA was eluted into 40 µl of elution buffer, warmed to 70 °C and centrifuged at $11,000 \times g$ for 1 min and the resultant DNA was stored at -20 °C.

2.5. DNA extraction from soil and faeces

DNA was extracted from soil and faecal samples (one gram each) using the Fast DNA Spin Kit for soil (QBiogene, Carlsbad, CA, USA) according to the manufacturer's instructions, and eluted in 70 μ l DES (DNase/Pyrogen Free Water). Sterile soil (autoclaved twice at 121 °C for 15 min) was used as negative controls for each set of extractions. The resultant DNA was stored at -20 °C.

2.6. DNA extraction from bedding

One gram of each of 10 bedding samples was thawed, and suspended in 40 ml of transport buffer (sterile PBS

containing 20 mM Na₂ EDTA; pH 8.0). The samples were shaken for 1 h at 37 °C followed by centrifugation for 15 min at $13,523 \times g$ at 4 °C. The supernatant was removed and the pellet resuspended in 2 ml sterile PBS. DNA was extracted from 200 µl of this solution using the NucleoSpin Blood kit (Macherey-Nagel, GmbH and Co, Düren, Germany) according to the manufacturer's recommendations, and DNA was stored at -20 °C.

2.7. End point and nested PCR

PCR amplifications were performed using an Eppendorf vapo.protect Mastercycler (Eppendorf, Hamburg, Germany). Each 50 µl reaction contained 25 µl Promega PCR master mix (Promega, Southampton, UK), 1.0 µl each of primers Cc and Ac [10 mM] (La Fontaine et al., 1993) (Table 1), 2.5 µl of dimethyl sulfoxide (Fisher Scientific, Leicestershire, UK), 2 µl bovine serum albumin [10 mg ml⁻¹] (Sigma–Aldrich Ltd., Poole, Dorset, UK), 16.5 µl of nuclease free water and 2 µl of template DNA. For direct detection of D. nodosus, PCR was performed using Cc and Ac primers (La Fontaine et al., 1993) (Table 1) under the following conditions: 1 cycle of 95 °C for 2 min, 40 cycles of 95 °C for 1 min. 60 °C for 45 s and 72 °C for 2 min and a final extension step of 72 °C for 5 min. Samples that were negative using this approach, were tested further using nested PCR. In the first round 16S rRNA universal primers 27F and 1525R (Table 1) (Baker et al., 2003; Lane, 1991) were used in the conditions described above but with an annealing temperature of 55 °C, 1 µl of this product was used in the second round of PCR as described above. The PCR products were visualised under UV light.

Table 1

All primers and probes used in the study.

2.8. Quantitative PCR of D. nodosus

The load of D. nodosus was determined using the Applied Biosystems 7500 Fast real-time detection system (Applied Biosystems, Warrington, UK). The qPCR targeted the rpoD gene (RNA polymerase sigma 70 factor, single copy number in *D. nodosus* genome) as described previously (Calvo-Bado et al., 2011b). All PCR reactions were performed in triplicate and each contained 12.5 µl TagMan Universal Master Mix (Applied Biosystems, Warrington, UK), 2.25 µl each of rpoDF and rpoDR $[10 \text{ pmol } \mu l^{-1}], 0.625 \text{ ml} \text{ rpoD} \text{ probe} [10 \text{ pmol } \mu l^{-1}]$ (Table 1), 1.25 μ l bovine serum albumin [10 mg ml⁻¹], 5.375 μ l nuclease free water and 1 μ l of template DNA. DNA dilutions of 1:10 were also used to investigate potential inhibitors of the reaction. In addition, known concentrations of target DNA were added to negative samples as internal controls. A non-template control (nuclease free water) was included in triplicate in all PCR reactions. The reaction was carried out under the following conditions: one cycle at 50 °C for 2 min, one cycle at 95 °C for 10 min, followed by 40 cycles at 95 °C for 15 s and 55 °C for 1 min. The *rpoD* copy number was estimated based on the standard curve obtained from analysis of 10fold serial dilutions of DNA extracted from *D. nodosus* strain VCS1703A.

2.9. Quantitative PCR assay for pgrA and pgrB

A fluorescent PCR was designed to increase the sensitivity of detection of *pgrA* and *B* (Table 1). The specificity of the *pgrA* and *B* primers was tested against genomic DNA isolated from 11 bacterial species and from a

| Primer (5'–3') | Sequence | Expected size in VCS1703A (BP) | Reference | | |
|-------------------|---------------------------------|-----------------------------------|---------------------------------------|--|--|
| (c | ΤΟΟΟΤΑΟΟΟΑΟΤΑΤΤΤΟΤΑΟΟΟΑΑΟΑΟΟΤ | 783 | La Fontaine et al. (1993) | | |
| Ac | CCCCTTATCTACCTTCC | 783 | La Fontaine et al. (1993) | | |
| 27F | | 1500 | La romanic et al. (1995) | | |
| 1525P | | 1500 | Lane (1991) ; Baker et al. (2003) | | |
| ngrAF1 | CCTCCACCATCCTTCTTAAA | 290 | Calvo-Bado et al. $(2011a)$ | | |
| ngrAR1 | CCTCTTCCTCCTTTCCCTAT | 290 | Calvo-Bado et al. $(2011a)$ | | |
| M13F | CTAAAACCACCCCCAC | N/A | Supplied in the cloning kit | | |
| M13R | CAGGAAACAGCTATGAC | N/A | Supplied in the cloning kit | | |
| DNTR02F | (FAM)-GATCCATCGTTTCATCGTCA | 549 | Russell et al (2014) | | |
| DNTR02R | CGCACTTTAGCCGTTATGTTT | 549 | Russell et al. (2014) | | |
| DNTR09F | (VIC)-GGCGTAAACGAAATGCCTAA | 987 | Russell et al. (2014) | | |
| DNTR09R | ATCGGCGGAAGATTGTCTC | 987 | Russell et al. (2014) | | |
| DNTR10F | (NED)-CCGTCTATCCACCCGATTTA | 626 | Russell et al. (2014) | | |
| DNTR10R | TTGAACCGCGTCACTATCAG | 626 | Russell et al. (2014) | | |
| DNTR19F | (PET)-CCCGTCGAATCACTCCAG | 854 | Russell et al. (2014) | | |
| DNTR19R | GGTAGCGCCGAAGAAAGA | 854 | Russell et al. (2014) | | |
| rpoDF | GCTCCCATTTCGCGCATAT | 61 | Calvo-Bado et al. (2011b) | | |
| rpoDR | CTGATGCAGAAGTCGGTAGAACA | 61 | Calvo-Bado et al. (2011b) | | |
| rpoD Taqman probe | (6FAM)-CATTCTTACCGGKCG-(BBQ) | 61 | Calvo-Bado et al. (2011b) | | |
| pgrAF | CATGAATGATAATATTTACCTTTTCGTT | 298 | | | |
| pgrAR | AAGATTGATGATGCTCCAGAAGAAG | 298 | | | |
| pgrA Taqman probe | (6FAM)-CCTGCACCATGCTTGTTAAACTCT | 298 | | | |
| | AATTTT-(BBQ) | | | | |
| pgrBF | AAAGGTGATCTCAACTGTATCGTCAT | N/A | | | |
| pgrBR | AATYARCARMGCCARAATTAGAGCTTAAT | N/A | | | |
| pgrB Taqman probe | (6FAM)-TTTACCCGCACCGTKCT-(BBQ) | N/A | | | |

FAM - Carboxyfluorescein, BBQ (Black Berry Quencher). BP is the size of fragment in base pairs.

range of diverse environmental samples. *D. nodosus* DNA from VCS1703A was used as a positive control and genomic DNA from *Escherichia coli* and sterile water as negative and no target controls respectively (Supplementary Table 2). The cycling conditions were modified slightly from those described above by reducing the number of cycles to 38 and 36 for *pgrA* and *pgrB* respectively.

2.10. Cloning and sequencing of pgrA amplicons

The *pgrA* gene was amplified using the primers *pgrA*F1 and pgrAR1 (Table 1; (Calvo-Bado et al., 2011a)) from the DNA extracted from the foot swabs of five ewes (n = 14)feet) and their lambs (n = 10 feet) and cloned using the TOPO TA Cloning Kit (Invitrogen Ltd., Paisley, UK). The cloning reactions were set up following the manufacturer's recommendations. Transformations were carried out using chemically competent (TOP10) E. coli cells (Invitrogen Ltd., Paisley, UK) with $50 \,\mu g \,ml^{-1}$ kanamycin. One-hundred microliters of the resulting solution was cultured on LB plates containing 50 µg ml⁻¹ kanamycin. Fifty colonies per sample were inoculated into individual wells of a 96-well plate, each containing 50 µl sterile water. The samples were heated to 75 °C for 10 min, and 1 µl of this solution was used as a template for PCR, resulting in analysis of 1200 transformants. The PCR products were run on a 1% high resolution agarose gel and visualised under UV light. The clones that showed variation in size within each foot were inoculated into LB media to provide sufficient biomass for plasmid DNA extraction using the Qiagen MiniPrep kit (Qiagen, West Sussex, UK). The plasmid DNA was digested using EcoR1 and sequenced by GATC Biotech (London, UK) using the supplied M13f and M13r primers (Invitrogen, Paisley, UK; Table 1). All sequences were deposited in GenBank database under accession numbers KR105403-KR105413.

2.11. Multi Locus VNTR analysis

The four D. nodosus tandem repeat (DNTR) loci were amplified individually as described previously (Russell et al., 2014) (Table 1) from the DNA isolated from the feet of all ewes and lambs. Primer specificity was tested against genomic DNA isolated from 11 bacterial species and from a range of diverse environmental samples (Supplementary Table 2). D. nodosus strain VCS1703A was included as a positive control for all PCR reactions. The amplified products from the community DNA were pooled in the ratio of 1:1:1:1 and submitted for fragment analysis to the University of Dundee (DNA Sequencing & Services, Dundee, Scotland). GeneScan 1200 LIZ dye (Applied Biosystems, Warrington, UK) was used as a size standard in fragment analysis. The data obtained were processed using Peak Scanner Software (Applied Biosystems, Warrington, UK) and analysed using T-REX (Culman et al., 2009) with a minimum fragment length cut off values of 300 bp and 500 bp for DNTR10 and DNTR19 respectively, peak height baseline threshold of 40 and bin range of 4 bp.

2.12. Statistical analysis

To determine the strain overlap between all ewes and lambs defined by MLVA, the coincidence index of overlap was calculated using the formula: C = 2B/(E+L), where C = coincidence index of overlap, B = occurrence of the same allele in ewes and lambs, E = occurrence of the allele in ewes, L = occurrence of the allele in lambs. Results can range from C = 0, no overlap between ewes and lambs to C = 1, identical strains occur in both ewes and lambs (Dice, 1945).

The *rpoD* copy number in ewes and lambs was not normally distributed. Therefore, a Mann–Whitney U test (Mann and Whitney, 1947) was used to test for differences in copy number between ewes and lambs.

3. Results

3.1. D. nodosus copy number detection limit and its persistence in the environment

End point PCR and qPCR data of the inoculated swabs suggested that the minimum detection level was 10⁴ cells and 10^2 rboD genome equivalents (i.e. 10^2 cells) per swab respectively assuming 100% DNA recovery. Below this concentration, detection was not reproducible. D. nodosus was detected by end point PCR in 5/10 faecal samples/balls from the interdigital space, 2/10 straw bedding samples collected after ewes were housed, 5/30 fresh hoof prints and 2/4 soil samples taken from the areas surrounding water containers. However qPCR analysis revealed that *D. nodosus* was present at loads of $10^2 - 10^4$ rpoD genome equivalents per gram in all the used straw bedding samples and $10^3 - 10^4$ rpoD genome equivalents per gram in all the faecal samples. Quantitative PCR also confirmed that D. nodosus was not detectable in the three stored straw samples (Fig. 1).

3.2. Quantification of D. nodosus in ewes and lambs

D. nodosus was not detected in lambs' feet at birth but was detected in all lambs and ewes 5–13 h later after their feet had touched the floor initially in a large communal pen, and subsequently in an individual pen. *D. nodosus* was detected on 39/40 (97.5%) of ewes' feet and 39/40 (97.5%) of lambs' feet (Fig. 2). Whilst overall the population loads were significantly higher in ewes than lambs (Mann–Whitney U test; *p*-value < 0.001); analysis of ewe/lamb pairs suggested only ewes 2, 3 and 5 had a higher load than their lambs. The *D. nodosus* load ranged from 10³ to 10⁵ *rpoD* genome equivalents per swab in lambs and 10^2 – 10^7 *rpoD* genome equivalents per swab in ewes (Supplementary Fig. 1).

3.3. Detection of pgr variants in the community DNA

pgrA was detected on 23/40 (57.5%) of ewes' feet and 15/40 (37.5%) of lambs' feet, whereas *pgrB* was detected on 27/40 (67.5%) of ewes' feet and 22/40 (55%) of lambs' feet (Fig. 2). Both variants *pgrA* and *pgrB* were detected on eight ewes and nine lambs.

Forty-two *pgrA* clones were sequenced from 14 foot swabs from five ewes and 10 foot swabs from five lambs. This resulted in the detection of 11 variants containing 3–21 tandem repeats in the R1 region, with 2–6 variants per animal (Table 2). Multiple *pgrA* variants with varying



Fig. 1. Presence of *D. nodosus* in the environmental samples. Absolute quantification of *rpoD* gene in bedding samples from the storage area, used bedding samples and faeces compacted within the interdigital space. Each bar is the average of triplicate analyses, error bars represent \pm standard deviation.

numbers of tandem repeats were observed in a single foot swab (data not shown).

3.4. Molecular typing MLVA from the community DNA

Alleles at *D. nodosus* tandem repeat DNTR19 were detected on the feet of six ewes and their lambs whereas for DNTR10 alleles were only detected on two ewe/lamb

pairs (Table 3) (Supplementary Fig. 2). The fluorescent data for DNTR09 was below the peak height threshold level (40 fluorescence units) and DNTR02 primers demonstrated some non-specific binding, so were excluded from the analysis. For both loci (DNTR10 and DNTR19), one or two alleles occurred in lambs, but there was greater diversity in ewes, with up to six alleles detected (Table 3). As detectable diversity in ewes increased, so did the likelihood of



Fig. 2. Detection of *pgr* variants in the community DNA. Presence of *pgrA*, *pgrB* and *rpoD* on the feet of 10 ewes and 10 lambs. *pgrA/B* was absent/below the detection limit in the samples where no data is shown. (E = Ewe; L = Lamb).

 Table 2

 Distribution of *pgrA* R1 tandem repeats in five pairs of ewes and lambs (14 ewe and 10 lamb feet).

| Ewe/Lamb ID | Number of clones sequenced | Number of <i>pgrA</i> tandem repeats in the R1 region |
|-------------|-------------------------------|---|
| E 1 | 6 | 3, 4, 5, 11, 13, 16 |
| L 1 | 4 | 4, 11, 12, 16 |
| E 2 | 3 | 4, 11, 15 |
| L 2 | 5 | 3, 4, 6, 11, 16 |
| E 3 | 4 | 6, 15, 16, 21 |
| L 3 | 2 | 16, 20 |
| E 4 | 4 | 4, 6, 11, 13 |
| L 4 | 5 | 4, 6, 11, 12, 16 |
| E 5 | 4 | 5, 12, 15, 16 |
| L 5 | 5 | 4, 11, 12, 15, 16 |

detecting the same strain on its offspring. The overall coincidence index of overlap between ewes and lambs was 0.45.

4. Discussion

In the current study we demonstrated, for the first time, that lambs are born free from D. nodosus (given the sensitivity of the tests used) but that we were able to detect high levels of D. nodosus on foot swabs within a few hours of birth. The most likely route of transmission is from standing on contaminated bedding in communal pens, however, recent work (Witcomb, 2012) indicated that D. nodosus can be detected in the mouths of ewes, so transmission might also have occurred as the lamb was being cleaned by the ewe after birth. Quantitative PCR with specific primers is widely used to determine bacterial load from swabs (Fredricks et al., 2009; Jaton et al., 2006; Lund et al., 2004) especially when studying disease development over time (Srinivasan et al., 2010). Here, as elsewhere (Witcomb et al., 2014) gPCR of rpoD was used to determine D. nodosus load on the feet of ewes and lambs. It was striking that such a large *D. nodosus* population was present on the feet of lambs within a few hours of birth. This suggests that D. nodosus might be an early coloniser of

 Table 3

 DNTR19 and DNTR10 allelic distribution between six and two pairs of ewes and lambs.

| ID | DNTR19 | | | | | | DNTR10 | | | | | | |
|-----|--------|---|---|---|---|---|--------|---|---|---|---|----|-------|
| | 3* | 4 | 5 | 6 | 7 | 8 | Total | 3 | 4 | 7 | 9 | 10 | Total |
| E 1 | 1 | 1 | 1 | 1 | 1 | 1 | 6 | 1 | 1 | 1 | 1 | 1 | 5 |
| L 1 | 0 | 0 | 0 | 1 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 1 |
| E 2 | 1 | 1 | 0 | 1 | 0 | 0 | 3 | 1 | 0 | 0 | 1 | 0 | 2 |
| L 2 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 1 |
| Е З | 0 | 1 | 0 | 0 | 0 | 0 | 1 | | | | | | |
| L 3 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | | | | | | |
| E 4 | 0 | 1 | 1 | 0 | 0 | 0 | 2 | | | | | | |
| L 4 | 0 | 0 | 0 | 1 | 0 | 0 | 1 | | | | | | |
| Е 5 | 1 | 1 | 1 | 1 | 1 | 0 | 5 | | | | | | |
| L 5 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | | | | | | |
| E 6 | 0 | 1 | 0 | 0 | 0 | 0 | 1 | | | | | | |
| L 6 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | | | | | | |

⁺ The numbers given in the table heading are the number of tandem repeats.

lambs' feet, although, it could be that naïve feet are simply colonized by the first bacterial species they encounter.

Several serogroups of *D. nodosus* on individual feet have been reported elsewhere (Claxton et al., 1983; Hindmarsh and Fraser, 1985; Jelinek et al., 2000; Moore et al., 2005) but not using the typing methods used in the current study where both *pgr* data and MLVA provided evidence for the occurrence of multiple strains in most animals studied. Not all the *D. nodosus* strains detected on the feet of lambs were on their mother's feet, indicating that the strains detected on lambs' feet originated from sources additional to their mothers' feet, most likely contaminated bedding in the communal pen. In addition, not all the strains detected in ewes were present in their offspring. It is possible that the specific variants on the ewes were not detected by chance, or were present below the minimum detection level or that only some of the strains were transferred via bedding.

Given the survival time of *D. nodosus* off host and the *D. nodosus*-negative stored straw samples, the most probable source of *D. nodosus* was the population of ewes in the communal pen. Methods to reduce the load of *D. nodosus* in ewes and the environment would probably have the biggest impact on transmission to newborn lambs. Management strategies that have been linked to a reduction in footrot prevalence and incidence include rapid appropriate treatment of diseased sheep (Kaler et al., 2010; Wassink et al., 2010), segregation of diseased sheep and footbathing healthy sheep (Wassink et al., 2004).

The identification of transmission routes and understanding the role of the environment is critical for the control of footrot. Previous studies have highlighted that diseased sheep are a reservoir of infection (Green et al., 2007; Kaler et al., 2010; Smith et al., 2014; Whittington, 1995), although transmission occurs indirectly via contaminated pasture or floors (Beveridge, 1941; Whittington, 1995). The current study has demonstrated that the environment potentially forms at least a temporary reservoir of infection for lambs because *D. nodosus* was detected in the straw bedding in the communal pens, and lambs have strains of *D. nodosus* on their feet not detected on their mother's feet.

The fact that lambs were *D. nodosus*-negative at birth suggests that it might be possible to produce *D. nodosus*-free lambs without the need for a caesarean birth. This finding is of interest to those performing challenge studies on pathogen-free individuals. At the practical, farm level this information might also be useful in countries that have eradicated footrot. In countries where footrot is endemic, even if lambs could be kept *D. nodosus*-free around lambing time, it is highly unlikely that this status could be maintained because of the high levels of infection in ewes. Reducing initial exposure (as suggested above) might be beneficial to subsequent disease severity, however, it might be detrimental if later age at first exposure increases disease severity.

5. Conclusions

We have provided evidence that lambs are born *D. nodosus*-negative, but within hours of birth several strains of *D. nodosus* are detectable on their feet. The

strains detected were a combination of those present on their mother's feet and on the feet of other ewes. Straw bedding in the communal pen was *D. nodosus*-positive and the most likely source of *D. nodosus* for newborn lambs.

Conflict of interest

None of the authors have any conflict of interest.

Acknowledgements

This work was supported in part by the Combating Endemic Diseases of Farmed Animals for Sustainability (CEDFAS) initiative, Grant No. BBE01870X1 from the Biotechnology and Biological Sciences Research Council (BBSRC). MM was in receipt of a Warwick Chancellor's International Scholarship Ref: 1150257/School of Life Sciences.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at http://dx.doi.org/10.1016/j. vetmic.2015.04.010.

References

- Baker, G.C., Smith, J.J., Cowan, D.A., 2003. Review and re-analysis of domain-specific 16S primers. J. Microbiol. Meth. 55, 541–555.
- Beveridge, 1941. A transmissible disease due to infection with Fusiformis nodusus (n.sp.): studies on its cause, epidemiology, and control, Bulletin of the Council of Scientific and Industrial Research Australia (Australia), pp. 1–58.
 Calvo-Bado, L.A., Green, L.E., Medley, G.F., Ul-Hassan, A., Grogono-Thomas,
- Calvo-Bado, L.A., Green, L.E., Medley, G.F., Ul-Hassan, A., Grogono-Thomas, R., Buller, N., Kaler, J., Russell, C.L., Kennan, R.M., Rood, J.I., Wellington, E.M., 2011a. Detection and diversity of a putative novel heterogeneous polymorphic proline-glycine repeat (Pgr) protein in the footrot pathogen Dichelobacter nodosus. Vet. Microbiol. 147, 358–366.
- Calvo-Bado, L.A., Oakley, B.B., Dowd, S.E., Green, L.E., Medley, G.F., Ul-Hassan, A., Bateman, V., Gaze, W., Witcomb, L., Grogono-Thomas, R., Kaler, J., Russell, C.L., Wellington, E.M., 2011b. Ovine pedomics: the first study of the ovine foot 16S rRNA-based microbiome. ISME J. 5, 1426–1437.
- Cederlof, S.E., Hansen, T., Klaas, I.C., Angen, O., 2013. An evaluation of the ability of *Dichelobacter nodosus* to survive in soil. Acta Vet. Scand. 55, 4.
- Claxton, P.D., Ribeiro, L.A., Egerton, J.R., 1983. Classification of Bacteroides nodosus by agglutination tests. Aus. Vet. J. 60, 331–334.
- Culman, S., Bukowski, R., Gauch, H., Cadillo-Quiroz, H., Buckley, D., 2009. T-REX: software for the processing and analysis of T-RFLP data. BMC Bioinformatics 10, 171.
- Dice, L.R., 1945. Measures of the amount of ecologic association between species. Ecology 26, 297–302.
- Fredricks, D.N., Fiedler, T.L., Thomas, K.K., Mitchell, C.M., Marrazzo, J.M., 2009. Changes in vaginal bacterial concentrations with intravaginal metronidazole therapy for bacterial vaginosis as assessed by quantitative PCR. J. Clin. Microbiol. 47, 721–726.
- Green, L.E., Wassink, G.J., Grogono-Thomas, R., Moore, L.J., Medley, G.F., 2007. Looking after the individual to reduce disease in the flock: a binomial mixed effects model investigating the impact of individual

sheep management of footrot and interdigital dermatitis in a prospective longitudinal study on one farm. Prev. Vet. Med. 78, 172–178.

- Hindmarsh, F., Fraser, J., 1985. Serogroups of Bacteroides nodosus isolated from ovine footrot in Britain. Vet. Rec. 116, 187–188.
- Jaton, K., Bille, J., Greub, G., 2006. A novel real-time PCR to detect Chlamydia trachomatis in first-void urine or genital swabs. J. Med. Microbiol. 55, 1667–1674.
- Jelinek, P.D., Depiazzi, L.J., Galvin, D.A., Spicer, I.T., Palmer, M.A., Pitman, D.R., 2000. Occurrence of different strains of *Dichelobacter nodosus* in new clinical lesions in sheep exposed to footrot associated with multi-strain infections. Aust. Vet. J. 78, 273–276.
- Kaler, J., Medley, G.F., Grogono-Thomas, R., Wellington, E.M.H., Calvo-Bado, L.A., Wassink, G.J., King, E.M., Moore, L.J., Russell, C., Green, L.E., 2010. Factors associated with changes of state of foot conformation and lameness in a flock of sheep. Prev. Vet. Med. 97, 237–244.
- La Fontaine, S., Egerton, J.R., Rood, J.I., 1993. Detection of *Dichelobacter* nodosus using species-specific oligonucleotides as PCR primers. Vet. Microbiol. 35, 101–117.
- Lane, D.J., 1991. 16S'23S rRNA sequencing. In: Stackebrandt, E., Goodfellow, M. (Eds.), Nucleic Acid Techniques in Bacterial Systematics. Wiley, Chichester, pp. 115–175.
- Lund, M., Nordentoft, S., Pedersen, K., Madsen, M., 2004. Detection of *Campylobacter spp.* in chicken fecal samples by real-time PCR. J. Clin. Microbiol. 42, 5125–5132.
- Mann, H.B., Whitney, D.R., 1947. On a Test of Whether one of Two Random Variables is Stochastically Larger than the Other, pp. 50–60.
- Moore, L.J., Wassink, G.J., Green, L.E., Grogono-Thomas, R., 2005. The detection and characterisation of *Dichelobacter nodosus* from cases of ovine footrot in England and Wales. Vet. Microbiol. 108, 57–67.
- Nieuwhof, G.J., Bishop, S.C., 2005. Costs of the major endemic diseases of sheep in Great Britain and the potential benefits of reduction in disease impact. Anim. Sci. 81, 23–29.
- Russell, C.L., Smith, E.M., Calvo-Bado, L.A., Green, L.E., Wellington, E.M., Medley, G.F., Moore, L.J., Grogono-Thomas, R., 2014. Multiple locus VNTR analysis highlights that geographical clustering and distribution of *Dichelobacter nodosus*, the causal agent of footrot in sheep, correlates with inter-country movements. Infect. Genet. Evol. 22, 273–279.
- Smith, E.M., Green, O.D., Calvo-Bado, L.A., Witcomb, L.A., Grogono-Thomas, R., Russell, C.L., Brown, J.C., Medley, G.F., KilBride, A.L., Wellington, E.M., Green, L.E., 2014. Dynamics and impact of footrot and climate on hoof horn length in 50 ewes from one farm over a period of 10 months. Vet. J. 201, 295–301.
- Srinivasan, S., Liu, C., Mitchell, C.M., Fiedler, T.L., Thomas, K.K., Agnew, K.J., Marrazzo, J.M., Fredricks, D.N., 2010. Temporal variability of human vaginal bacteria and relationship with bacterial vaginosis. PloS One 5, e10197.
- Stewart, D.J., 1989. Footrot of sheep. In: Egerton, J.R., Yong, W.K., Riffin, G.G. (Eds.), Footrot and Foot Abscess of Ruminants. CRC Press, Boca Raton, Florida, USA.
- Wassink, G.J., Grogono-Thomas, R., Moore, L.J., Green, L.E., 2004. Risk factors associated with the prevalence of interdigital dermatitis in sheep from 1999 to 2000. Vet. Rec. 154, 551–555.
- Wassink, G.J., King, E.M., Grogono-Thomas, R., Brown, J.C., Moore, L.J., Green, L.E., 2010. A within farm clinical trial to compare two treatments (parenteral antibacterials and hoof trimming) for sheep lame with footrot. Prev. Vet. Med. 96, 93–103.
- Whittington, R.J., 1995. Observations on the indirect transmission of virulent ovine footrot in sheep yards and its spread in sheep on unimproved pasture. Aust. Vet. J. 72, 132–134.
- Witcomb, L.A., 2012. The In Situ Analysis of the Microbial Community Associated with Footrot of Sheep. University of Warwick Ph.D. thesis.
- Witcomb, L.A., Green, L.E., Kaler, J., Ul-Hassan, A., Calvo-Bado, L.A., Medley, G.F., Grogono-Thomas, R., Wellington, E.M., 2014. A longitudinal study of the role of *Dichelobacter nodosus* and *Fusobacterium necrophorum* load in initiation and severity of footrot in sheep. Prev. Vet. Med. 115, 48–55.