

The journey towards safely restarting faecal microbiota transplantation services in the UK during the COVID-19 era

Quraishi, Mohammed Nabil; Shabir, Sahida; Manzoor, Susan E; Green, Christopher A; Sharma, Naveen; Beggs, Andrew D; Iqbal, Tariq H

DOI:

[10.1016/S2666-5247\(21\)00036-7](https://doi.org/10.1016/S2666-5247(21)00036-7)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Quraishi, MN, Shabir, S, Manzoor, SE, Green, CA, Sharma, N, Beggs, AD & Iqbal, TH 2021, 'The journey towards safely restarting faecal microbiota transplantation services in the UK during the COVID-19 era', *The Lancet. Microbe*, vol. 2, no. 4, pp. e133-e134. [https://doi.org/10.1016/S2666-5247\(21\)00036-7](https://doi.org/10.1016/S2666-5247(21)00036-7)

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

The journey towards safely restarting faecal microbiota transplantation services in the UK during the COVID-19 era



Faecal microbiota transplantation (FMT) is a potentially life-saving treatment for patients with refractory and recurrent *Clostridioides difficile* infection with cure rates of 90%.¹ Substantial implications with regard to the safe provision of FMT were made evident with emerging reports of detectable SARS-CoV-2 virus in stool even before the declaration of the COVID-19 pandemic.² Furthermore, data suggesting prolonged and potentially infectious faecal virus shedding in stool samples in asymptomatic carriers who tested negative on nasopharyngeal swabs meant FMT donor screening protocols could no longer rely on symptoms and a nasopharyngeal swab alone.^{3,4} The Medicine and Healthcare Regulatory Agency approved University of Birmingham Microbiome Treatment Centre (UoBMTC), was the first to mandate SARS-CoV-2 testing of stool as part of donor screening and called for a halt of FMT production globally until a validated stool assay was available⁵ with an evidence-based international consensus on guidance for donor screening being published.⁶

FMT production at UoBMTC was paused in March, 2020 and FMT that was already manufactured from February, 2020, was quarantined. Following consensus agreements with FMT providers globally and a review of updated evidence on safety and efficacy, the shelf life of FMT collected before December, 2019, was extended from 6 months to 12 months. This allowed our centre to treat a further 68 patients with recurrent *C difficile* infection until the stocks were exhausted by August, 2020.

In conjunction with FMT providers globally, we established new donor screening protocols that now include multiple levels of questionnaire-based and molecular screening for SARS-CoV-2 infection or enteric carriage at several timepoints of the donor and FMT production pathway (appendix). However, restarting FMT manufacture principally relied on a validated assay for detection of SARS-CoV-2 in stool, which is not commercially available. Our centre worked towards validating a sensitive and specific assay that uses multiple technologies in parallel for the detection of SARS-CoV-2 in stool.⁷ A combination

that includes real-time PCR, digital droplet PCR, and nanopore sequencing technology was validated with stool viral spiking experiments and testing of stool samples from more than 100 COVID-19 positive and COVID-19 negative hospitalised patients (on the basis of nasopharyngeal PCR). The assay was approved by the Medicine and Healthcare Regulatory Agency in January, 2021, and has enabled the resumption of donor screening and FMT services in the COVID-19 era. We have now released quarantined FMT aliquots from those manufactured in February, 2020, and have treated several patients successfully.

Although FMT is a safe and a highly effective treatment for recurrent *C difficile* infection, FDA safety alerts in the USA, following serious adverse events as a result of transmission of enteric pathogens, emphasise that there can be no room for complacency.⁸ Although our revised protocols are highly robust in screening asymptomatic carriers of SARS-CoV-2, both FMT providers and users need to remain vigilant for new threats during the changing viral pandemic and beyond.

We declare no competing interests.

Copyright © 2021 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.

*Mohammed Nabil Quraishi, Sahida Shabir, Susan E Manzoor, Christopher A Green, Naveen Sharma, Andrew D Beggs, Tariq H Iqbal
m.n.quraishi@bham.ac.uk

University of Birmingham Microbiome Treatment Centre, Birmingham, UK (MNQ, SS, SEM, CAG, NS, ADB, THI); University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK (MNQ, CAG, NS, ADB, THI); Institute of Cancer and Genomic Sciences, University of Birmingham, UK (MNQ, ADB); Institute of Microbiology and Infection, University of Birmingham, UK (CAG, THI)

- 1 Mullish BH, Quraishi MN, Segal JP, et al. The use of faecal microbiota transplant as treatment for recurrent or refractory *Clostridium difficile* infection and other potential indications: joint British Society of Gastroenterology (BSG) and Healthcare Infection Society (HIS) guidelines. *Gut* 2018; **67**: 1920–41.
- 2 Zhang JJ, Dong X, Cao YY, et al. Clinical characteristics of 140 patients infected with SARS-CoV-2 in Wuhan, China. *Allergy* 2020; **75**: 1730–41.
- 3 Hindson J. COVID-19: faecal-oral transmission? *Nat Rev Gastroenterol Hepatol* 2020; **17**: 259.
- 4 Wu Y, Guo C, Tang L, et al. Prolonged presence of SARS-CoV-2 viral RNA in faecal samples. *Lancet Gastroenterol Hepatol* 2020; **5**: 434–35.
- 5 Green CA, Quraishi MN, Shabir S, et al. Screening faecal microbiota transplant donors for SARS-CoV-2 by molecular testing of stool is the safest way forward. *Lancet Gastroenterol Hepatol* 2020; **5**: 531.

Published Online
February 19, 2021
[https://doi.org/10.1016/S2666-5247\(21\)00036-7](https://doi.org/10.1016/S2666-5247(21)00036-7)

See Online for appendix

- 6 Ianiro G, Mullah BH, Kelly CR, et al. Reorganisation of faecal microbiota transplant services during the COVID-19 pandemic. *Gut* 2020; **69**: 1555–63.
- 7 Mansoor S, Zaman S, Whalley C, et al. Multi-modality detection of SARS-CoV-2 in faecal donor samples for transplantation and in asymptomatic emergency surgical admissions. *medRxiv* 2021; published online Feb 3. <https://doi.org/10.1101/2021.02.02.21250934> (preprint).
- 8 US FDA. Important safety alert regarding use of fecal microbiota for transplantation and risk of serious adverse reactions due to transmission of multi-drug resistant organisms. June 13, 2019; <https://www.fda.gov/vaccines-blood-biologics/safety-availability-biologics/important-safety-alert-regarding-use-fecal-microbiota-transplantation-and-risk-serious-adverse> (accessed Feb 1, 2021).