

Comparison of respiratory health impacts associated with wood and charcoal biomass fuels

Woolley, Katherine; Bartington, Suzanne; Kabera, Telesphore ; Lao, Xiang Qian; Pope, Francis; Greenfield, Sheila; Price, Malcolm; Thomas, G Neil

DOI:
[10.3390/ijerph18179305](https://doi.org/10.3390/ijerph18179305)

License:
Creative Commons: Attribution (CC BY)

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (Harvard):
Woolley, K, Bartington, S, Kabera, T, Lao, XQ, Pope, F, Greenfield, S, Price, M & Thomas, GN 2021, 'Comparison of respiratory health impacts associated with wood and charcoal biomass fuels: a population-based analysis of 475,000 children from 30 low- and middle-income countries', *International Journal of Environmental Research and Public Health*, vol. 18, no. 17, 9305. <https://doi.org/10.3390/ijerph18179305>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



Article

Comparison of Respiratory Health Impacts Associated with Wood and Charcoal Biomass Fuels: A Population-Based Analysis of 475,000 Children from 30 Low- and Middle-Income Countries

Katherine E. Woolley ¹, Suzanne E. Bartington ^{1,*}, Telesphore Kabera ², Xiang-Qian Lao ³, Francis D. Pope ⁴, Sheila M. Greenfield ¹, Malcolm J. Price ^{1,5} and G. Neil Thomas ¹

- ¹ Institute of Applied Health Research, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK; KEW863@student.bham.ac.uk (K.E.W.); S.M.Greenfield@bham.ac.uk (S.M.G.); M.Price.2@bham.ac.uk (M.J.P.); g.n.thomas@bham.ac.uk (G.N.T.)
 - ² College of Science and Technology, University of Rwanda, Avenue de l'Armee, Kigali P.O. Box 3900, Rwanda; kaberafris@yahoo.fr
 - ³ The Jockey Club School of Public Health and Primary Care, The Chinese University of Hong Kong, Hong Kong; xqlao@cuhk.edu.hk
 - ⁴ School of Geography, Earth and Environmental Sciences, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK; F.Pope@bham.ac.uk
 - ⁵ NIHR Birmingham Biomedical Research Centre, University Hospitals Birmingham NHS Foundation Trust and University of Birmingham, Birmingham B15 2TT, UK
- * Correspondence: s.bartington@bham.ac.uk



Citation: Woolley, K.E.; Bartington, S.E.; Kabera, T.; Lao, X.-Q.; Pope, F.D.; Greenfield, S.M.; Price, M.J.; Thomas, G.N. Comparison of Respiratory Health Impacts Associated with Wood and Charcoal Biomass Fuels: A Population-Based Analysis of 475,000 Children from 30 Low- and Middle-Income Countries. *Int. J. Environ. Res. Public Health* **2021**, *18*, 9305. <https://doi.org/10.3390/ijerph18179305>

Received: 14 July 2021

Accepted: 30 August 2021

Published: 3 September 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Background: The World Health Organisation reported that 45% of global acute respiratory infection (ARI) deaths in children under five years are attributable to household air pollution, which has been recognised to be strongly associated with solid biomass fuel usage in domestic settings. The introduction of legislative restrictions for charcoal production or purchase can result in unintended consequences, such as reversion to more polluting biomass fuels such as wood; which may increase health and environmental harms. However, there remains a paucity of evidence concerning the relative health risks between wood and charcoal. This study compares the risk of respiratory symptoms, ARI, and severe ARI among children aged under five years living in wood and charcoal fuel households across 30 low- and middle-income countries. Methods: Data from children ($N = 475,089$) residing in wood or charcoal cooking households were extracted from multiple population-based Demographic and Health Survey databases (DHS) ($N = 30$ countries). Outcome measures were obtained from a maternal report of respiratory symptoms (cough, shortness of breath and fever) occurring in the two weeks prior to the survey date, generating a composite measure of ARI (cough and shortness of breath) and severe ARI (cough, shortness of breath and fever). Multivariable logistic regression analyses were implemented, with adjustment at individual, household, regional and country level for relevant demographic, social, and health-related confounding factors. Results: Increased odds ratios of fever (AOR: 1.07; 95% CI: 1.02–1.12) were observed among children living in wood cooking households compared to the use of charcoal. However, no association was observed with shortness of breath (AOR: 1.03; 95% CI: 0.96–1.10), cough (AOR: 0.99; 95% CI: 0.95–1.04), ARI (AOR: 1.03; 95% CI: 0.96–1.11) or severe ARI (AOR: 1.07; 95% CI: 0.99–1.17). Within rural areas, only shortness of breath was observed to be associated with wood cooking (AOR: 1.08; 95% CI: 1.01–1.15). However, an increased odds ratio of ARI was observed in Asian (AOR: 1.25; 95% CI: 1.04–1.51) and East African countries (AOR: 1.11; 95% CI: 1.01–1.22) only. Conclusion: Our population-based observational data indicates that in Asia and East Africa there is a greater risk of ARI among children aged under 5 years living in wood compared to charcoal cooking households. These findings have major implications for understanding the existing health impacts of wood-based biomass fuel usage and may be of relevance to settings where charcoal fuel restrictions are under consideration.

Keywords: acute respiratory infection; biomass fuel; household air pollution; respiratory symptoms; low-and middle-income countries

1. Introduction

Exposure to household air pollution (HAP) is associated with adverse child and maternal health outcomes, including morbidity and mortality in children under five years old [1–3], acute respiratory infection (ARI) [4], child growth failure [5], low birth weight, and stillbirths [6]. Vulnerability to ARI, the leading cause of mortality in children under five years worldwide [7], is high among children due to a greater level of pollutant inhalation from the same external concentration as their adult counterparts, and more susceptible pulmonary physiology [4]. HAP includes carbon monoxide (CO), particulate matter (PM), sulphur dioxide (SO₂) and nitrogen dioxide (NO₂) produced from burning biomass (wood, dung, charcoal and crop residue) for cooking, heating and lighting. Despite the known high HAP exposures in low- and middle-income countries (LMICs), there are few sustainable “cleaner fuel” interventions available to these populations, due to multiple barriers to sustained uptake, including low financial and infrastructural capabilities, lack of awareness, and appropriate policies [8]. Research has indicated harm reduction approaches such as outdoor cooking [9–12] and cooking with charcoal compared to wood [13,14], lowers exposure; with two small scale studies rural/peri-urban and urban settings, providing evidence for a respiratory health difference between wood and charcoal users [15,16]. However, the evidence in support of such approaches in LMIC settings remains limited.

Some governments have adopted legislative approaches to restrict the use of charcoal due to the recognised environmental and health impacts [17]. Evidence from domestic and commercial kitchens suggests that charcoal cooking is associated with high levels of PM [18] and CO [19] above the World Health Organisation’s Indoor Air Quality Guidelines (WHO-IAQ) [18,20]. Introduction of charcoal fuel-based legislative changes or fiscal disincentives are typically intended to improve population health, including shifting to cleaner fuels such as electricity and liquid petroleum gas (LPG) alternatives. However, such changes may also generate unintended consequences [21,22], such as substitution with more polluting biomass fuels (e.g., wood, dung, straw) [23] which are typically readily available and cheaper alternatives [24]. Wood is the most common fuel used globally and it is therefore preferred as it suits traditional cooking practices [25]. In addition, LPG adoption is not likely in the imminent future due to multiple barriers, including equipment and fuels access, cost, and safety concerns [26]. Societal and economic issues with uncertainty can also affect fuel choices [27,28], meaning fuel transition often does not occur in a linear fashion [29,30]. In the advent of policy measures to restrict charcoal use, it is possible that charcoal could be replaced by wood fuel by end-users, presenting overall health risks given that wood produces more PM than charcoal [18]. But there remains a paucity of evidence in the relative health effects between wood and charcoal cooking, on a global scale.

We report the association of under-five respiratory health (respiratory symptoms, ARI, severe ARI) with wood and charcoal fuel use for cooking, in over 475,000 children from 30 LMICs, using comprehensive population-based data obtained from the Demographic and Health Survey (DHS).

2. Materials and Methods

2.1. Data Sources

A cross-sectional study across 30 LMIC countries was conducted using data obtained from the most recently available national population-based Demographic and Health Survey (DHS) [31], with LMIC status defined using the Development Assistance Committee (DAC) list 2020 [32]. Criteria for country inclusion included: (i) DHS survey data available within the last 10 years, (ii) presence of wood and charcoal cooking fuel use (iii) presence of the outcome variables of interest (Appendix A: Figure A1). Each country

followed the same two-stage stratified DHS sampling methodology with proportionate random sampling and standardised questionnaires with fieldwork supported by United States Agency for International Development (USAID). Eligible participants were identified through the residential household survey and included ever-married (has been married at least once in their life) women aged 15–49 years and men aged 15–59 years, who resided in the household the night before the survey [33]. Non-response households at the time of data collection and those with institutional living arrangements (e.g., boarding schools, police camps, army barracks, and hospitals) were excluded.

All countries followed the standard core questionnaire from Phases VI, VII, and VIII of the DHS Program model, with country-specific modifications to non-core questions to reflect the population and health issues most relevant to that country. USAID standardises and provides training to government agencies and health authorities to complete surveys, with internal training and supervision of local data collectors and data entry. The questionnaire is translated into the main language(s) for each country and validated on approximately 100–200 households. Data for this current analysis were obtained from (i) household dataset containing situational and household characteristics; (ii) woman's dataset containing maternal characteristics; (iii) children's dataset containing health and individual characteristics. All primary data collection has ethical approval from the relevant government authority within each country, with all data being anonymised and aggregated for DHS online data archive [31]. The archive is publicly available and authorisation for data access has been gained for this study.

2.2. Modified Wealth Index

The wealth index provided by DHS is calculated through principal component analysis, including cooking fuel as an indicator variable [34], therefore to prevent effect underestimation due to circularity, a modified wealth index was calculated [35] following the DHS provided guide [36] using SPSS [37], to calculate a modified wealth index. The new wealth index included indicator variables for the source of drinking water, house construction material (wall, roof and floor), toilet facility and assets. The assets included vary by country [37] and have been documented in Appendix B: Table A1. The wealth index was then ranked by household to provide tertiles of wealth.

2.3. Outcome Variables-Measure of Respiratory Symptoms and Acute Respiratory Infection

Maternal respondents were asked to report the presence of respiratory symptoms (shortness of breath, cough and fever) during the two weeks prior to the survey among all children under the age of five years living in their household. Respiratory symptoms were modelled as binary outcomes (yes, no), included short rapid breaths or difficulty breathing, cough, and fever. These respiratory symptoms were used to form the composite measures for ARI (both shortness of breath and cough [38]), and severe ARI (each of shortness of breath, cough, and fever [39–41]). Composite measures for ARI and severe ARI were then modelled as binary (yes, no) outcomes.

2.4. Measure of Exposure to HAP

Cooking fuel use was recorded from self-report for each household that undertook cooking activities. Fuels were categorised as “Cleaner fuels” (electricity, LPG, natural gas, biogas) and “Solid biomass fuels and kerosene” (kerosene, coal/lignite, charcoal, wood, straw/shrubs/grass, agricultural crop, animal dung). Wood and charcoal cooking household fuels were extracted and modelled as a binary variable.

2.5. Explanatory Variables

Individual child characteristics included child's age (0–11, 12–23, 24–35, 36–48, 48–59 months), sex (male, female), mode of delivery (caesarean, vaginal), birth order (first, not first born), breastfeeding status (ever, never), vitamin A supplementation in the last 6 months (yes, no), iron supplementation (yes, no). Maternal characteristics in-

cluded age of mother (15–24, 25–35, 36–49 years), mother's highest attained educational level (none, primary, secondary/higher). Household characteristics included: number of household members (≤ 6 , >6), household smoking (yes, no), cooking location (indoor, outdoor), and modified wealth index (lowest, low, middle, high, highest) [34]. Situational variables included geographical region of residence and area of residence (rural, urban). All co-variables were modelled as categorical variables.

2.6. Missing Data

Data that were identified to be missing at random with less than 50% missing data [42,43] underwent multiple imputations of 50 iterations [44,45], at a country level, using the MICE package [46] in R studio [47].

2.7. Data Analysis

Using R studio [47], descriptive statistics were tabulated with the number of cases (n), and percentage (%) for categorical outcome variables within the combined dataset. The association between the health outcome variables and exposure to HAP was assessed using a multivariable logistic regression using the Survey package [48] in R to account for the sampling strategy. Adjusted odds ratios (AOR) and 95% confidence intervals (95% CI) for each country were obtained and presented on a forest plot, with a summary result for the combined dataset. Additional exploratory analyses of a subset of countries were undertaken, incorporating breastfeeding, birthweight, and household smoking, which were missing or incomplete in a number of countries. Stratified analyses were undertaken to investigate the association in rural and urban settings, indoor and outdoor cooking status, geographic location and before or after 2014 (mid-time point of included studies), separately.

3. Results

3.1. Cooking Fuel Usage and Number of Respiratory Outcomes

Out of the 30 included country datasets, there was substantial variation in the type of fuels used within the country (Figure 1), however, wood was the predominant cooking fuel (range: 2.5–94.9%). Indonesia, Afghanistan, Peru, Pakistan and India have a large proportion of “cleaner” fuel use (range: 48.9–56.6%), with low charcoal usage (range: 0.4–2.1%). Within the pooled dataset before imputation ($N = 475,089$), 88.7% used wood cooking fuel compared to 11.1% using charcoal cooking fuel (Table 1). Overall, there were 23,490 cases of severe ARI (5.3%), 36,657 of ARI (8.3%), with shortness of breath being reported in 38,703 children (8.8%), cough in 82,523 children (18.7%), and fever in 89,621 children (20.3%) (Table 1).

3.2. Risk of Respiratory Symptoms, ARI, and Severe ARI

After adjusting for individual and situational potential confounding factors, children who resided in wood cooking households were observed to have increased adjusted odds ratios (Figure 2) with fever only (AOR: 1.07; 95% CI: 1.02–1.12), in the pooled dataset. No association was observed with ARI (AOR: 1.03; 95% CI: 0.96–1.11) or severe ARI (AOR: 1.07; 95% CI: 0.99–1.17). However, at a country level Afghanistan (AOR: 4.24; 95% CI: 1.66–10.83), Pakistan (AOR: 2.44; 95% CI: 1.29–4.61), Burundi (AOR: 1.73; 95% CI: 1.21–2.46), Zambia (AOR: 1.62; 95% CI: 1.16–2.26), Philippines (AOR: 1.44; 95% CI: 1.04–2.00) and Uganda (AOR: 1.34; 95% CI: 1.02–1.76), were all observed to have increased adjusted odds ratios of severe ARI in children residing in wood cooking households compared to charcoal cooking. This observed increase in adjusted odds ratios was also present in Afghanistan (AOR: 3.38; 95% CI: 1.23–9.29), Pakistan (AOR: 2.71; 95% CI: 1.45–5.07), Zambia (AOR: 1.43; 95% CI: 1.12–1.83), Burundi (AOR: 1.40; 95% CI: 1.06–1.84) and Uganda (AOR: 1.26; 95% CI: 1.00–1.58) for ARI. Little change was observed in the effect estimate when controlling for birthweight, breastfeeding and household smoking in those countries with available data (Appendix C: Table A2).




Figure 1. Proportion of clean, kerosene, wood, charcoal, other biomass (dung, crop residue) fuel use within each country, ordered by geographical region.

Table 1. Descriptive statistics before imputation for respiratory health outcomes (N = 475,089).

	Cough (N = 442,450, Missing = 6.9%)			Shortness of Breath (N = 442,040, Missing = 7.0%)			Fever (N = 442,550, Missing = 6.9%)			ARI (N = 441,987, Missing = 7.0%)			Severe ARI (N = 441,627, Missing = 7.0%)		
	No (N = 359,927) n (%)	Yes (N = 82,523) n (%)	p Value	No (N = 403,337) n (%)	Yes (N = 38,703) n (%)	p Value	No (N = 352,929) n (%)	Yes (N = 89,621) n (%)	p Value	No (N = 405,330) n (%)	Yes (N = 36,657) n (%)	p Value	No (N = 418,137) n (%)	Yes (N = 23,490) n (%)	p Value
Household cooking fuel			<0.001			<0.001			<0.001			<0.001			0.790
Charcoal	36,279 (10.1%)	12,296 (14.9%)		43,749 (10.8%)	4799 (12.4%)		37,859 (10.7%)	10,774 (12.0%)		44,033 (10.9%)	4500 (12.3%)		45,933 (11.0%)	2546 (10.8%)	
Wood	323,648 (89.9%)	70,227 (85.1%)		359,588 (89.2%)	33,904 (87.6%)		315,070 (89.3%)	78,847 (88.0%)		361,297 (89.1%)	32,156 (87.7%)		372,204 (89.0%)	20,944 (89.2%)	
Child's sex			<0.001			0.064			<0.001			<0.001			<0.001
Male	183,227 (50.9%)	42,317 (51.3%)		205,018 (50.8%)	20,329 (52.5%)		179,258 (50.8%)	46,364 (51.7%)		206,066 (50.8%)	19,249 (52.5%)		212,678 (50.9%)	12,463 (53.1%)	
Child's Age (months)			<0.001			<0.001			<0.001			<0.001			<0.001
0–11	76,440 (21.2%)	18,754 (22.7%)		85,537 (21.2%)	9575 (24.7%)		75,566 (21.4%)	19,644 (21.9%)		86,053 (21.2%)	9055 (24.7%)		89,380 (21.4%)	5693 (24.2%)	
12–23	69,761 (19.4%)	20,002 (24.2%)		80,043 (19.8%)	9614 (24.8%)		66,964 (19.0%)	22,806 (25.4%)		80,529 (19.9%)	9124 (24.9%)		83,344 (19.9%)	6261 (26.7%)	
24–35	69,555 (19.3%)	16,566 (20.1%)		78,494 (19.5%)	7556 (19.5%)		67,785 (19.2%)	18,370 (20.5%)		78,899 (19.5%)	7140 (19.5%)		81,311 (19.4%)	4671 (19.9%)	
36–47	72,650 (20.2%)	14,756 (17.9%)		80,785 (20.0%)	6552 (16.9%)		71,814 (20.3%)	15,601 (17.4%)		81,120 (20.0%)	6206 (16.9%)		83,422 (20.0%)	3785 (16.1%)	
48–59	71,521 (19.9%)	12,445 (15.1%)		78,479 (19.5%)	5405 (14.0%)		70,799 (20.1%)	13,200 (14.7%)		78,729 (19.4%)	5133 (14.0%)		80,680 (19.3%)	3080 (13.1%)	
Birth order			<0.001			<0.001			<0.001			<0.001			<0.001
Not first born	256,340 (71.2%)	60,619 (73.5%)		288,083 (71.4%)	28,569 (73.8%)		250,638 (71.0%)	66,433 (74.1%)		289,685 (71.5%)	26,928 (73.5%)		299,038 (71.5%)	17,343 (73.8%)	
Mode of delivery *			0.005			<0.001			0.099			<0.001			<0.001
Caesarean	25,107 (7.0%)	6134 (7.5%)		28,378 (7.1%)	2846 (7.4%)		25,064 (7.2%)	6182 (7.0%)		28,487 (7.1%)	2735 (7.5%)		29,437 (7.1%)	1777 (7.6%)	

Table 1. Cont.

	Cough (N = 442,450, Missing = 6.9%)			Shortness of Breath (N = 442,040, Missing = 7.0%)			Fever (N = 442,550, Missing = 6.9%)			ARI (N = 441,987, Missing = 7.0%)			Severe ARI (N = 441,627, Missing = 7.0%)		
	No (N = 359,927) n (%)	Yes (N = 82,523) n (%)	p Value	No (N = 403,337) n (%)	Yes (N = 38,703) n (%)	p Value	No (N = 352,929) n (%)	Yes (N = 89,621) n (%)	p Value	No (N = 405,330) n (%)	Yes (N = 36,657) n (%)	p Value	No (N = 418,137) n (%)	Yes (N = 23,490) n (%)	p Value
Birthweight*			<0.001			<0.001			<0.001			<0.001			<0.001
Low	63,970 (28.8%)	11,814 (23.3%)		69,945 (28.0%)	5799 (25.3%)		62,841 (28.6%)	12,935 (24.4%)		70,163 (27.9%)	5578 (25.7%)		72,017 (27.8%)	3687 (26.5%)	
Breastfeeding status*			<0.001			<0.001			<0.001			<0.001			<0.001
Never Breast-fed	14,047 (5.1%)	2197 (3.2%)		15,209 (4.9%)	1021 (3.2%)		13,850 (5.1%)	2402 (3.1%)		15,244 (4.8%)	981 (3.2%)		15,537 (4.8%)	674 (3.4%)	
Vitamin A supplementation			<0.001			<0.001			<0.001			<0.001			<0.001
Yes	193,115 (54.6%)	49,265 (60.9%)		219,310 (55.3%)	22,867 (60.2%)		190,098 (54.8%)	52,338 (59.4%)		220,415 (55.3%)	21,741 (60.5%)		227,872 (55.4%)	14,113 (61.3%)	
Iron supplementation *			0.106			<0.001			0.018			<0.001			<0.001
Yes	63,042 (18.4%)	12,534 (16.6%)		69,285 (18.1%)	6245 (17.5%)		60,992 (18.2%)	14,587 (17.3%)		69,500 (18.1%)	6026 (17.9%)		71,308 (18.0%)	4156 (19.1%)	
Maternal age (years)			0.012			<0.001			<0.001			<0.001			<0.001
15–24	121,452 (33.7%)	27,738 (33.6%)		135,727 (33.7%)	13,329 (34.4%)		119,212 (33.8%)	29,974 (33.4%)		136,313 (33.6%)	12,723 (34.7%)		140,647 (33.6%)	8244 (35.1%)	
25–35	189,953 (52.8%)	42,587 (51.6%)		212,443 (52.7%)	19,888 (51.4%)		186,522 (52.8%)	46,110 (51.5%)		213,507 (52.7%)	18,795 (51.3%)		220,127 (52.6%)	11,987 (51.0%)	
36–49	48,523 (13.5%)	12,198 (14.8%)		55,167 (13.7%)	5486 (14.2%)		47,195 (13.4%)	13,537 (15.1%)		55,509 (13.7%)	5138 (14.0%)		57,363 (13.7%)	3259 (13.9%)	
Maternal education level			<0.001			<0.001			<0.001			<0.001			<0.001
No education	143,818 (40.0%)	25,418 (30.8%)		155,811 (38.6%)	13,185 (34.1%)		136,969 (38.8%)	32,317 (36.1%)		156,599 (38.6%)	12,384 (33.8%)		160,619 (38.4%)	8196 (34.9%)	
Primary	98,665 (27.4%)	30,379 (36.8%)		114,886 (28.5%)	14,076 (36.4%)		98,262 (27.8%)	30,839 (34.4%)		115,753 (28.6%)	13,178 (36.0%)		120,526 (28.8%)	8302 (35.3%)	
Secondary /Higher	117,428 (32.6%)	26,724 (32.4%)		132,623 (32.9%)	11,442 (29.6%)		117,685 (33.3%)	26,461 (29.5%)		132,961 (32.8%)	11,095 (30.3%)		136,974 (32.8%)	6991 (29.8%)	

Table 1. Cont.

	Cough (N = 442,450, Missing = 6.9%)			Shortness of Breath (N = 442,040, Missing = 7.0%)			Fever (N = 442,550, Missing = 6.9%)			ARI (N = 441,987, Missing = 7.0%)			Severe ARI (N = 441,627, Missing = 7.0%)		
	No (N = 359,927) n (%)	Yes (N = 82,523) n (%)	p Value	No (N = 403,337) n (%)	Yes (N = 38,703) n (%)	p Value	No (N = 352,929) n (%)	Yes (N = 89,621) n (%)	p Value	No (N = 405,330) n (%)	Yes (N = 36,657) n (%)	p Value	No (N = 418,137) n (%)	Yes (N = 23,490) n (%)	p Value
Household wealth index			<0.001			<0.001			<0.001			<0.001			<0.001
Lowest	100,447 (27.9%)	21,240 (25.7%)		110,928 (27.5%)	10,698 (27.6%)		97,093 (27.5%)	24,616 (27.5%)		111,451 (27.5%)	10,167 (27.7%)		114,771 (27.4%)	6773 (28.8%)	
Middle	80,643 (22.4%)	17,910 (21.7%)		90,194 (22.4%)	8240 (21.3%)		78,956 (22.4%)	19,595 (21.9%)		90,616 (22.4%)	7810 (21.3%)		93,332 (22.3%)	5000 (21.3%)	
Highest	27,563 (7.7%)	8875 (10.8%)		32963 (8.2%)	3434 (8.9%)		28,989 (8.2%)	7464 (8.3%)		33,149 (8.2%)	3237 (8.8%)		34,533 (8.3%)	1811 (7.7%)	
Household smoking *			<0.001			<0.001			<0.001			<0.001			<0.001
Yes	126,552 (36.2%)	24,092 (31.4%)		138,907 (35.6%)	11,612 (32.2%)		123,113 (36.1%)	27,526 (32.2%)		139,279 (35.5%)	11,231 (33.0%)		142,874 (35.4%)	7518 (34.3%)	
Household cooking location			0.118			0.395			<0.001			0.001			<0.001
Indoors	262,449 (73.2%)	60,621 (73.8%)		293,885 (73.1%)	28,871 (75.0%)		258,651 (73.5%)	64,479 (72.4%)		295,282 (73.1%)	27433 (75.3%)		304,800 (73.2%)	17,651 (75.6%)	
Number of household member *			<0.001			<0.001			<0.001			<0.001			<0.001
≤6	174,300 (48.5%)	44,496 (53.9%)		198,285 (49.2%)	20,349 (52.6%)		172,461 (48.9%)	46,326 (51.7%)		199,396 (49.2%)	19,201 (52.4%)		206,328 (49.4%)	12,071 (51.4%)	
Place of residence			0.476			<0.001			0.048			<0.001			0.578
Urban	66,652 (18.5%)	17,978 (21.8%)		77,206 (19.1%)	7349 (19.0%)		67,436 (19.1%)	17,232 (19.2%)		77,496 (19.1%)	7049 (19.2%)		80,271 (19.2%)	4201 (17.9%)	

N = observation number, n = category observation number, % = column percentage for category. p value = Chi-Squared. * Missing data = Mode of delivery = 0.7%, Breastfeeding status = 22.8%, Birthweight = 38.3%, Vitamin A supplementation = 1.7%, Iron Supplementation = 53.4%, Mother's education = 0.004%, Household smoking = 3.6%, Cooking location = 0.4%, Number of household members = 0.06%.




Figure 2. Forest plot illustrating the adjusted odds ratio (AOR) for respiratory symptoms, ARI and severe ARI with wood cooking fuel compared to charcoal for all countries. The summary measure is adjusted for: age, birth order, mode of delivery, vitamin A supplementation, mother’s age, mother’s education level, wealth status, number of household members, rural/urban residence and location of cooking.

3.3. Role of Rural and Urban Residence

In the rural and urban sub-analysis cough was observed to be associated with an increased odds ratio (AOR: 1.08; 95% CI: 1.01–1.15) among children residing in wood compared to charcoal fuel households in rural areas only.

3.4. Role of Outdoor Cooking

In the analyses of the pooled dataset, for indoor cooking children under five years residing in households using wood had increased adjusted odds ratios of fever (AOR: 1.07; 95% CI: 1.00–1.13). No other differences were observed (Table 2).

Table 2. Summary effects (AOR–95% CI) for respiratory symptoms, ARI and severe ARI with wood cooking of the whole, exploratory and sub-analysis.

Analysis (N)	Cough AOR (95%CI)	Shortness of Breath AOR (95%CI)	Fever AOR (95%CI)	ARI AOR (95%CI)	Severe ARI AOR (95%CI)
Whole (N = 482,644)	0.99(0.95–1.04)	1.03(0.96–1.10)	1.07(1.02–1.12) ^b	1.03(0.96–1.11)	1.07(0.99–1.17)
Sub-analysis					
Urban areas (N = 89,661)	0.93(0.87–1.00)	0.99(0.90–1.09)	1.03(0.96–1.10)	0.99(0.89–1.09)	1.02(0.90–1.14)
Rural Area (N = 392,983)	1.08(1.01–1.15) ^c	1.00(0.90–1.11)	1.05(0.98–1.13)	1.02(0.92–1.14)	1.05(0.92–1.20)
Indoor (N = 368,647)	1.03(0.97–1.09)	1.05(0.96–1.14)	1.07(1.00–2.13) ^c	1.06(0.97–1.15)	1.08(0.97–1.20)
Outdoor (N = 113,997)	0.96(0.89–1.04)	0.98(0.89–1.09)	1.07(0.99–1.16)	1.01(0.91–1.13)	1.10(0.96–1.25)
Africa (N = 245,363)	1.02(0.97–1.08)	1.00(0.93–1.08)	1.02(0.97–1.08)	1.01(0.93–1.09)	1.04(0.94–1.14)
Asia (N = 233,091)	1.05(0.92–1.20)	1.25(1.04–1.51) ^b	1.06(0.93–1.20)	1.25(1.04–1.51) ^c	1.24(0.99–1.54)
Central Africa (N = 47,710)	0.97(0.86–1.11)	0.99(0.79–1.24)	0.99(0.86–1.13)	0.99(0.79–1.25)	1.04(0.80–1.36)
East Africa (N = 105,543)	1.09(1.02–1.17) ^b	1.10(1.00–1.20) ^c	1.01(0.94–1.09)	1.11(1.01–1.22) ^c	1.10(0.98–1.24)
West Africa (N = 92,110)	0.96(0.87–1.06)	0.86(0.76–0.97) ^b	1.00(0.92–1.09)	0.88(0.78–1.00)	0.93(0.80–1.09)
Surveys undertaken during or before 2014 (N = 134,225)	0.97(0.91–1.04)	0.99(0.89–1.09)	1.01(0.94–1.08)	0.99(0.89–1.09)	1.00(0.89–1.14)
Surveys undertaken during or after 2015 * (N = 348,419)	1.04(0.97–1.12)	1.09(1.00–1.20)	1.18(1.10–1.26) ^a	1.11(1.01–1.22) ^c	1.17(1.04–1.31) ^b

AOR = adjusted odds ratio for wood cooking compared to charcoal, 95% CI = 95% confidence interval, a = $p \leq 0.001$, b = $p \leq 0.01$, c = $p \leq 0.05$, bold = $p \leq 0.05$, N = number of observations in the SARI analysis. * Surveys that were undertaken across 2014–2015 (n = 2) were included within during or after 2015 (Total countries = 15).

3.5. Role of Geographic Location

In the stratified sub-analysis by geographic location, an association with ARI in children under five years old living in wood compared to charcoal cooking households was observed in East Africa (AOR: 1.11; 95% CI: 1.01–1.22) and Asia (AOR: 1.25; 95% CI: 1.04–1.51) (Table 2). An increase in the adjusted odds ratio with wood cooking compared to charcoal was also observed with shortness of breath in Asia (AOR: 1.25; 95% CI: 1.04–1.51) and East Africa (AOR: 1.10; 95% CI: 1.00–1.20), whereas a decrease in the adjusted odds ratio was observed in West Africa (AOR: 0.86; 95% CI: 0.76–0.97). An association was observed with cough (AOR: 1.09; 95% CI: 1.02–1.17) in East Africa only.

3.6. Role of Time Period Survey Was Undertaken

In the stratified sub-analysis of those surveys undertaken during or after 2015, an association within an increase in the adjusted odds ratio of fever (AOR: 1.18; 95% CI: 1.10–1.26), ARI (AOR: 1.11; 95% CI: 1.01–1.22), severe ARI (AOR: 1.17; 95% CI: 1.04–1.31) in children under five years old living in wood compared to charcoal cooking households (Table 2). However, no associations were observed in surveys undertaken during or before 2014.

4. Discussion

In our large cross-sectional multi-country study (30 countries; 475,089 participants), increased odds ratios of ARI were observed in Asia (AOR: 1.09; 95% CI: 1.05–1.13) and East Africa only (AOR: 1.04; 95% CI: 1.00–1.08) among children living in wood cooking households compared to charcoal cooking households. The risk of ARI varies between countries, and this may reflect different wood fuel choices, cultural differences, access to healthcare [49], elevation [50] and seasonal or climatic differences [51], which could not be accounted for in our analyses. Moreover, the variation of the observed outcome results between countries indicates the need to take current country and regional level characteristics into consideration when developing HAP interventions for reducing ARI in children aged under five years. This is further highlighted as only the most recent surveys (post-2015) have an observed association with fever (AOR: 1.18; 95% CI: 1.10–1.26), ARI (AOR: 1.11; 95% CI: 1.01–1.22), severe ARI (AOR: 1.17; 95% CI: 1.04–1.31). Prevention

of ARI in children aged under five years would reduce child mortality and long-term morbidities, exert health and fiscal benefits; in addition to supporting progress towards the Sustainable Development Goals (SDGs) (namely SDG 3).

Given the wider environmental and health impacts of charcoal production and use, there is an argument for restricting the use of charcoal. However, the clear financial and structural difficulties of provision of clean fuels in many LMIC settings means charcoal restriction could potentially result in health harms by some users reverting to other biomass fuels. Although charcoal use presents significant health harms [52], it has previously been shown to produce lower pollutant levels than wood cooking in laboratory and field studies [15,18], and our results reflect these findings and their effect on child health, indicating that compared to charcoal, wood cooking is associated with increased risk of ARI in East Africa and Asia. It is evident that a package of measures is required for fuel transition policies which include charcoal restrictions, to limit increased uptake of wood alternatives. Adoption of charcoal restrictions should be carefully considered, in terms of the potential health harms, in LMIC settings, in the context of clean fuel access, availability and affordability. Given the volatility of fuel costs, for example in response to disruptive changes such as COVID-19, such policies must also consider the longer-term resilience of domestic fuel supplies, in each specific context.

Fuel choice, preparation, cooking characteristics, and cumulative exposure have been shown to vary between countries [53]. We also explored how cooking location (indoor vs outdoor) and exposure to second-hand smoke from household smoking could potentially contribute to the risk of ARI and severe ARI. In the main analysis outdoor cooking was associated with reduced risk of shortness of breath, cough, fever and ARI (AOR: 0.96; 95% CI: 0.94–0.98), which is another potential interim harm mitigation behavioural intervention promoted to reduce the adverse health effects of HAP exposure [10]. However, in the sub-analysis of outdoor cooking and indoor cooking separately, an association was only observed with fever (AOR: 1.07; 95% CI: 1.00–2.13) in indoor cooking households, which may have resulted from the small sample size or influence of seasonal factors [51]. However, a more detailed country-specific assessment by differing solid biomass fuels [40] would be required to understand the overall potential benefits of cooking outdoors; in addition, to the combined health effects and pollutants level benefits when changing cooking fuel to charcoal and moving cooking outside.

Household smoking could not be accounted for in the main summary analysis, due to missing data in Peru, the Philippines, and Kenya DHS surveys. In the exploratory analysis, limited effect of household smoking was observed upon outcome measures. Smoking is not only an alternative source of HAP exposure but is also a recognised risk factor for respiratory infections in infants [54]. However, an association was only observed in households with a smoker with cough (AOR: 1.05; 95% CI: 1.01–1.08) and fever (AOR: 1.04; 95% CI: 1.01–1.07), compared to non-smoking households, indicating the limited potential of smoking causing the occurrence of ARI.

Urban and rural areas also have additional differing situational contexts, including housing type, co-inhabiting with livestock, food security, WASH, household crowding, malnutrition, access to healthcare, wealth [55] and ambient pollution levels [56]. Differences in changes over time were investigated through the sub-analysis comparing surveys between 2010–2014 ($n = 15$) and 2015–2018 ($n = 15$), which indicated potential differing situational contexts. The role of alternative sources of HAP and differing situational characteristics both within and between countries highlights the complexities that need to be considered to understand the context-specific needs and acceptability of behavioural harm reduction HAP interventions.

Although there is an environmental and health need for reducing the reliance on charcoal cooking fuel, legislative approaches to restricting charcoal use should take into consideration the potential unintended or unanticipated health consequences of targeted fiscal policies. Wood fuels are readily available in most settings, as they are typically free to collect, thus may be reverted to as a fuel of choice [57], along with being strongly

linked with poverty [58]; as seen in this study with the wealth index. This combination of factors increases the vulnerability of households to the health-harms of solid biomass cooking. Other approaches to improving the sustainability of charcoal such as improving sustainable production and the use of ICS for improved burning efficiency [17,59,60], could be considered as alternative mitigation measures in the short to medium term. However, the longer-term solution is to support the sustained adoption of cleaner fuels, with maintained strong supply links [61] to prevent fuel switching, as seen in the COVID-19 lockdowns [23,27]; and also provide education for health-harms of using solid biomass cooking fuel. Any policy mitigation measures for HAP to reduce ARIs in children under five should also consider wider protective health behaviours against ARIs, e.g., encouraging breastfeeding, especially within the first 6 months of life [62,63], childhood vaccinations, undernutrition [4], reducing the incidence of HIV, TB [64], and reducing the risk of low birth weight [65].

Although the use of fuel type as a proxy for HAP exposure, self-reported respiratory symptoms, changes over time, weaknesses in the potential to control for all confounding factors and the observational nature of the data generate study limitations, this population-based approach provides a large sample size and global comparison, detailing the widespread impact of a potential harm reduction intervention for HAP exposure. In addition, many potential associations were investigated; therefore, some association would be expected to be down to chance. Further research implications include the need to characterise exposure levels and exposure-response functions for key health outcomes and increased clinical diagnostic confirmation to improve aetiological specificity. Further research is also necessary to understand the specific physiological mechanism between specific pollutant exposure and ARI risk in children aged under five years, including by specific wood and charcoal types and combustion techniques. Further, we recommend consideration of the implications of wood to charcoal transition for climate change, specifically for CO₂ emissions [66] and environmental degradation associated with charcoal production [67]. This study has global implications and provides the evidence to support a clear policy recommendation for safer domestic cooking practices.

5. Conclusions

Our population-based observational study indicates that in Asia and East Africa there is a greater risk of ARI among children aged under 5 years living in wood compared to charcoal cooking households. Users of domestic wood fuels are among the most vulnerable sub-populations worldwide and our findings support the need for ensuring long-term uptake of clean domestic energy alternatives in resource-poor settings worldwide. Policy-makers should adopt an evidence-based approach, to ensure long-term sustained uptake of clean domestic energy alternatives and to prevent unintended consequences of biomass fuel restriction policies.

Author Contributions: Conceptualization, K.E.W., S.E.B. and G.N.T.; methodology, K.E.W. and M.J.P.; software, K.E.W.; formal analysis, K.E.W.; data curation, K.E.W.; writing—original draft preparation, K.E.W.; writing—review and editing, S.E.B., T.K., X.-Q.L., F.D.P., S.M.G., M.J.P. and G.N.T.; visualization, K.E.W.; supervision, S.E.B., F.D.P., S.M.G., and G.N.T. All authors have read and agreed to the published version of the manuscript.

Funding: This study has been funded by a University of Birmingham Global Challenges Scholarship held by KW, the Institute for Global Innovation (IGI) 'Clean Air' and 'Resilient Cities' themes and UK Department for International Development (DFID) via the East Africa Research Fund (EARF) grant 'A Systems Approach to Air Pollution (ASAP) East Africa and EPSRC (EP/T030100/1). MJP is supported by the NIHR Birmingham Biomedical Research Centre at the University Hospitals Birmingham NHS Foundation Trust and the University of Birmingham. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR, or the Department of Health and Social Care. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The APC was funded by the University of Birmingham Library service.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study. Informed consent for the primary data collection can be found at <https://dhsprogram.com/> (accessed on 15 January 2020).

Data Availability Statement: Data is freely and publicly available at: <https://dhsprogram.com/> (accessed on 15 January 2020).

Acknowledgments: This study would not have been possible without the approval and access to the DHS data archive.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

Appendix A




Figure A1. Flow diagram for selection of final dataset for analysis from all household data sets available from the DHS program data archive [31] (a) Survey was excluded due to have low cell counts for wood or charcoal cooking ($n = 31$) included: Comoros 2012, Zimbabwe 2010–2011, Zimbabwe 2015, Senegal 2010–2017 (continuous dataset), Togo 2013–2014, Kyrgyz 2012, Tajikistan 2012, 2017, Papua New Guinea 2016–2017, Bangladesh 2011, 2014, Indonesia 2017, Maldives 2016–2017, Nepal 2011, 2016, Timor-Leste 2016, Albania 2017–2018, Armenia 2010, 2015–2016, Jordan 2012, Yemen 2012, Columbia 2010, 2015, Guatemala 2014–2015, Honduras 2011–2012, South Africa 2016, Namibia 2013, Lesotho 2014. (b) Surveys were excluded due to low cell counts with explanatory variable ($n = 3$) and low cell counts of missing entirely for wealth ($n = 3$) or three or more key variables (e.g., Breastfeeding, mode of delivery, household smoking) ($n = 3$). (c) Peru, Zambia and Kenya were excluded due to <50% missing data for breastfeeding. (d) Household smoking data was not collected for Kenya, Peru and the Philippines. (e) Chad, Ethiopia, Kenya, Guinea, Liberia, Afghanistan, Myanmar, Pakistan, Nigeria were excluded due to <50% missing data for birthweight.

Appendix C

Table A2. Summary effects (AOR–95% CI) for respiratory symptoms, ARI and severe ARI with wood cooking for the exploratory analysis.

Analysis (N)	Cough AOR (95%CI)	Shortness of Breath AOR (95%CI)	Fever AOR (95%CI)	ARI AOR (95%CI)	Server ARI AOR (95%CI)
Controlling for birthweight ¹ (N = 405,839)	0.97(0.93–1.03)	1.02(0.94–1.10)	1.08(1.02–1.14) ^a	1.02(0.94–1.10)	1.02(0.94–1.10)
Controlling for breastfeeding ² (N = 448,769)	0.97(0.92–1.02)	1.01(0.94–1.09)	1.07(1.01–1.12) ^b	1.02(0.94–1.10)	1.07(0.98–1.18)
Controlling for household smoking ³ (N = 455,289)	0.98(0.93–1.03)	1.01(0.94–1.09)	1.06(1.01–1.12) ^c	1.02(0.94–1.10)	1.07(0.97–1.17)

AOR = adjusted odds ratio for wood cooking compared to charcoal, 95% CI = 95% confidence interval, a = $p \leq 0.001$, b = $p \leq 0.01$, c = $p \leq 0.05$, bold = $N = p \leq 0.05$, number of observations in the SARI analysis. ¹ = Chad, Ethiopia, Kenya, Guinea, Liberia, Afghanistan, Myanmar, Pakistan, Nigeria were excluded due to < 50% missing data for birthweight. ² = Peru, Zambia and Kenya were excluded due to < 50% missing data for breastfeeding. ³ = Household smoking data not collected for Kenya, Peru and the Philippines.

References

- Cohen, A.J.; Brauer, M.; Burnett, R.; Anderson, H.R.; Frostad, J.; Estep, K.; Balakrishnan, K.; Brunekreef, B.; Dandona, L.; Dandona, R.; et al. Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: An analysis of data from the Global Burden of Diseases Study 2015. *Lancet* **2017**, *389*, 1907–1918. [[CrossRef](#)]
- Bruce, N.G.; Dherani, M.K.; Das, J.K.; Balakrishnan, K.; Adair-Rohani, H.; Bhutta, Z.A.; Pope, D. Control of household air pollution for child survival: Estimates for intervention impacts. *BMC Public Health* **2013**, *13* (Suppl. 3), S8. [[CrossRef](#)] [[PubMed](#)]
- Smith, K.; Bruce, N.; Balakrishnan, K.; Adair-Rohani, H.; Balmes, J.; Chafe, Z.; Dherani, M.; Hosgood, H.D.; Mehta, S.; Pope, D.; et al. Millions dead: How do we know and what does it mean? Methods used in the comparative risk assessment of household air pollution. *Annu. Rev. Public Health* **2014**, *35*, 185–206. [[CrossRef](#)] [[PubMed](#)]
- Gordon, S.B.; Bruce, N.G.; Grigg, J.; Hibberd, P.L.; Kurmi, O.P.; Lam, K.H.; Mortimer, K.; Asante, K.P.; Balakrishnan, K.; Balmes, J.; et al. Respiratory risks from household air pollution in low and middle income countries. *Lancet Respir. Med.* **2014**, *2*, 823–860. [[CrossRef](#)]
- Tielsch, J.M.; Katz, J.; Thulasiraj, R.D.; Coles, C.L.; Sheeladevi, S.; Yanik, E.L.; Rahmathullah, L. Exposure to indoor biomass fuel and tobacco smoke and risk of adverse reproductive outcomes, mortality, respiratory morbidity and growth among newborn infants in south India. *Int. J. Epidemiol.* **2009**, *38*, 1351–1363. [[CrossRef](#)] [[PubMed](#)]
- Amegah, A.K.; Quansah, R.; Jaakkola, J.J.K. Household air pollution from solid fuel use and risk of adverse pregnancy outcomes: A systematic review and meta-analysis of the empirical evidence. *PLoS ONE* **2014**, *9*, e0113920. [[CrossRef](#)] [[PubMed](#)]
- United Nations Inter-Agency Group for Child Mortality Estimation (UN IGME) Levels & Trends in Child Mortality: Report 2019-Estimates Developed by the UN Inter-Agency Group for Child Mortality Estimation. Available online: <https://www.unicef.org/media/60561/file/UN-IGME-child-mortality-report-2019.pdf> (accessed on 18 August 2020).
- Puzzolo, E.; Stanistreet, D.; Pope, D.; Bruce, N.; Rehfuess, E. *Factors Influencing the Large-Scale Uptake by Households of Cleaner and more Efficient Household Energy Technologies*; EPPI-Centre, Social Science Research Unit, Institute of Education, University of London: London, UK, 2013; ISBN 9781907345623.
- Van Vliet, E.D.S.; Asante, K.; Jack, D.W.; Kinney, P.L.; Whyatt, R.M.; Chillrud, S.N.; Abokyi, L.; Zandoh, C.; Owusu-Agyei, S. Personal exposures to fine particulate matter and black carbon in households cooking with biomass fuels in rural Ghana. *Environ. Res.* **2013**, *127*, 10–48. [[CrossRef](#)]
- Langbein, J.; Peters, J.; Vance, C. Outdoor cooking prevalence in developing countries and its implication for clean cooking policies. *Environ. Res. Lett.* **2017**, *12*, 115008. [[CrossRef](#)]
- Yamamoto, S.S.; Louis, V.R.; Sié, A.; Sauerborn, R. Biomass smoke in Burkina Faso: What is the relationship between particulate matter, carbon monoxide, and kitchen characteristics? *Environ. Sci. Pollut. Res.* **2014**, *21*, 2581–2591. [[CrossRef](#)] [[PubMed](#)]
- Sidhu, M.K.; Ravindra, K.; Mor, S.; John, S. Household air pollution from various types of rural kitchens and its exposure assessment. *Sci. Total Environ.* **2017**, *586*, 419–429. [[CrossRef](#)]
- Fullerton, D.G.; Semple, S.; Kalambo, F.; Suseno, A.; Malamba, R.; Henderson, G.; Ayres, J.G.; Gordon, S.B. Biomass fuel use and indoor air pollution in homes in Malawi. *Occup. Environ. Med.* **2009**, *66*, 777–783. [[CrossRef](#)] [[PubMed](#)]
- Ellegård, A. Cooking fuel smoke and respiratory symptoms among women in low-income areas in Maputo. *Environ. Health Perspect.* **1996**, *104*, 980–985. [[CrossRef](#)]

15. Taylor, E.T.; Nakai, S. Prevalence of acute respiratory infections in women and children in Western Sierra Leone due to smoke from wood and charcoal stoves. *Int. J. Environ. Res. Public Health* **2012**, *9*, 2252–2265. [CrossRef] [PubMed]
16. Sana, A.; Meda, N.; Badoum, G.; Kafando, B.; Bouland, C. Primary Cooking Fuel Choice and Respiratory Health Outcomes among Women in Charge of Household Cooking in Ouagadougou, Burkina Faso: Cross-Sectional Study. *Int. J. Environ. Res. Public Health* **2019**, *16*, 1040. [CrossRef]
17. Vos, J.; Vis, M. *Making Charcoal Production in Sub Sahara Africa Sustainable*; Report Prepared for the Ministry of Economic Affairs, Agriculture and Innovation; Government of the Netherlands: Amsterdam, The Netherlands, 2010.
18. Titcombe, M.E.; Simcik, M.; Titcombe, M.E.; Simcik, M. Personal and indoor exposure to PM 2.5 and polycyclic aromatic hydrocarbons in the southern highlands of Tanzania: A pilot-scale study. *Environ. Monit. Assess.* **2011**, *180*, 461–476. [CrossRef]
19. Woolley, K.; Bartington, S.E.; Pope, F.D.; Price, M.J.; Thomas, G.N.; Kabera, T. Biomass cooking carbon monoxide levels in commercial canteens in Kigali, Rwanda. *Arch. Environ. Occup. Health* **2020**, *76*, 75–85. [CrossRef] [PubMed]
20. Bartington, S.E.; Bakolis, I.; Devakumar, D.; Kurmi, O.P.; Gulliver, J.; Chaube, G.; Manandhar, D.S.; Saville, N.M.; Costello, A.; Osrin, D.; et al. Patterns of domestic exposure to carbon monoxide and particulate matter in households using biomass fuel in Janakpur, Nepal. *Environ. Pollut.* **2017**, *220*, 38–45. [CrossRef] [PubMed]
21. Goldemberg, J.; Martinez-Gomez, J.; Sagar, A.; Smith, K.R. Household air pollution, health, and climate change: Cleaning the air. *Environ. Res. Lett.* **2018**, *13*, 030201. [CrossRef]
22. Farabi-Asl, H.; Taghizadeh-Hesary, F.; Chapman, A.; Mohammadzadeh Bina, S. Energy Challenges for Clean Cooking in Asia, the Background and Possible Policy solutions. Available online: <https://www.adb.org/publications/energy-challenges-clean-cooking-asia> (accessed on 19 April 2021).
23. Zhang, Y.; Li, Z. COVID-19's Impact on the Transition to Clean Cooking Fuels: Initial Findings from a Case Study in Rural Kenya. Available online: <https://openknowledge.worldbank.org/handle/10986/35258> (accessed on 19 April 2021).
24. Zulu, L.C. The forbidden fuel: Charcoal, urban woodfuel demand and supply dynamics, community forest management and woodfuel policy in Malawi. *Energy Policy* **2010**, *38*, 3717–3730. [CrossRef]
25. Hooper, L.G.; Dieye, Y.; Ndiaye, A.; Diallo, A.; Sack, C.S.; Fan, V.S.; Neuzil, K.M.; Ortiz, J.R. Traditional cooking practices and preferences for stove features among women in rural Senegal: Informing improved cookstove design and interventions. *PLoS ONE* **2018**, *13*, e0206822. [CrossRef] [PubMed]
26. Stanistreet, D.; Hyseni, L.; Puzzolo, E.; Higgerson, J.; Ronzi, S.; Anderson de Cuevas, R.; Adekoje, O.; Bruce, N.; Mbatchou Ngahane, B.; Pope, D. Barriers and facilitators to the Adoption and Sustained use of cleaner fuels in Southwest Cameroon: Situating 'lay' knowledge within evidence-based policy and practice. *Int. J. Environ. Res. Public Health* **2019**, *16*, 4702. [CrossRef] [PubMed]
27. Shupler, M.; Mwitari, J.; Gohole, A.; Anderson de Cuevas, R.; Puzzolo, E.; Čukić, I.; Nix, E.; Pope, D. COVID-19 impacts on household energy & food security in a Kenyan informal settlement: The need for integrated approaches to the SDGs. *Renew. Sustain. Energy Rev.* **2021**, *144*, 111018. [CrossRef]
28. Ravindra, K.; Kaur-Sidhu, M.; Mor, S.; Chakma, J.; Pillarisetti, A. Impact of the COVID-19 pandemic on clean fuel programmes in India and ensuring sustainability for household energy needs. *Environ. Int.* **2021**, *147*, 106335. [CrossRef] [PubMed]
29. Masera, O.R.; Saatkamp, B.D.; Kammen, D.M. From linear fuel switching to multiple cooking strategies: A critique and alternative to the energy ladder model. *World Dev.* **2000**, *28*, 2083–2103. [CrossRef]
30. Hiemstra-van der Horst, G.; Hovorka, A.J. Reassessing the "energy ladder": Household energy use in Maun, Botswana. *Energy Policy* **2008**, *36*, 3333–3344. [CrossRef]
31. The Demographic and Health Survey Program (DHS) The DHS Program US-AID. Available online: <https://dhsprogram.com/> (accessed on 7 June 2019).
32. Development Assistance Committee (DAC) DAC List of ODA Recipients Effective for Reporting on 2020 Flows. Available online: <https://www.oecd.org/dac/financing-sustainable-development/development-finance-standards/DAC-List-of-ODA-Recipients-for-reporting-2020-flows.pdf> (accessed on 2 September 2020).
33. Croft, T.; Marshall, A.M.J.; Courtney, A.K.; Al, E. *Guide to DHS Statistics*; ICF: Rockville, MD, USA, 2018.
34. Rutstein, S.O.; Johnson, K. *The DHS Wealth Index*; DHS Comparative Reports No. 6; DHS: Calverton, MD, USA, 2004.
35. Tusting, L.S.; Bottomley, C.; Gibson, H.; Kleinschmidt, I.; Tatem, A.J.; Lindsay, S.W.; Gething, P.W. Housing improvements and malaria risk in Sub-Saharan Africa: A multi-country analysis of survey data. *PLoS Med.* **2017**, *14*, e1002234. [CrossRef] [PubMed]
36. Rutstein, S.O. *Steps to Constructing the New DHS Wealth Index*; ICF: Rockville, MD, USA, 2015.
37. IBM. *Corp IBM SPSS Statistics for Windows. Version 27.0*; IBM: Armonk, NY, USA, 2020.
38. Simoes, E.A.F.; Cherian, T.; Chow, J.; Shahid-Salles, S.A.; Laxminarayan, R.; John, T.J. Acute respiratory infections in children. In *Disease Control Priorities in Developing Countries*; Breman, J.D., Measham, A., Eds.; Oxford University Press: New York, NY, USA; The International Bank for Reconstruction and Development/The World Bank: Washington, DC, USA, 2006; Volume 2, pp. 1180–1184. ISBN 0821361791.
39. Madhi, S.A.; Klugman, K.P. Acute respiratory infections. In *Disease and Mortality in Sub-Saharan Africa*; Makgoba, M., Hofman, K.J., Feacham, R.G.A., Bainga, F., Rogo, K.O., Bos, E.R., Jamison, D.T., Eds.; The International Bank for Reconstruction and Development/The World Bank: Washington, DC, USA, 2006; p. 149. ISBN 9781617798887.
40. Sk, R.; Rasooly, M.H.; Barua, S. Do fuel type and place of cooking matter for acute respiratory infection among Afghan children? Evidence from the Afghanistan DHS 2015. *J. Biosoc. Sci.* **2020**, *52*, 140–153. [CrossRef]

41. World Health Organization (WHO). WHO Global Technical Consultation: Global Standards and Tools for Influenza Surveillance. Geneva. Available online: https://www.who.int/influenza/resources/documents/technical_consultation/en/ (accessed on 6 April 2020).
42. Mishra, S.; Khare, D. On comparative performance of multiple imputation methods for moderate to large proportions of missing data in clinical trials: A simulation study. *J. Med. Stat. Inform.* **2014**, *2*, 9. [[CrossRef](#)]
43. Madley-Dowd, P.; Hughes, R.; Tilling, K.; Heron, J. The proportion of missing data should not be used to guide decisions on multiple imputation. *J. Clin. Epidemiol.* **2019**, *110*, 63–73. [[CrossRef](#)]
44. White, I.R.; Royston, P.; Wood, A.M. Multiple imputation using chained equations: Issues and guidance for practice. *Stat. Med.* **2011**, *30*, 377–399. [[CrossRef](#)]
45. Bodner, T.E. What improves with increased missing data imputations? *Struct. Equ. Model. A Multidiscip. J.* **2008**, *15*, 651–675. [[CrossRef](#)]
46. Van Buuren, S.; Groothuis-Oudshoorn, K. Mice: Multivariate imputation by chained equations in R. *J. Stat. Softw.* **2011**, *45*, 1–67. [[CrossRef](#)]
47. R Core Team R. *A Language and Environment for Statistical Computing*, R version 3.6.0; R Foundation for Statistical Computing: Vienna, Austria, 2020.
48. Lumley, T. *Survey: Analysis of Complex Survey Samples*; R Package Version 4.0; 2020.
49. Yaya, S.; Bishwajit, G. Burden of acute respiratory infections among under-five children in relation to household wealth and socioeconomic status in bangladesh. *Trop. Med. Infect. Dis.* **2019**, *4*, 36. [[CrossRef](#)] [[PubMed](#)]
50. Rosati, J.A.; Yoneda, K.Y.; Yasmeen, S.; Wood, S.; Eldridge, M.W. Respiratory health and indoor air pollution at high elevation. *Arch. Environ. Occup. Health* **2005**, *60*, 96–105. [[CrossRef](#)] [[PubMed](#)]
51. Buchner, H.; Rehfues, E.A. Cooking and season as risk factors for acute lower respiratory infections in African children: A cross-sectional multi-country analysis. *PLoS ONE* **2015**, *10*, e0128933. [[CrossRef](#)] [[PubMed](#)]
52. World Health Organization (WHO). Burning Opportunity: Clean Household Energy for Health, Sustainable Development, and Wellbeing of Women and Children. Luxembourg. Available online: <https://www.who.int/airpollution/publications/burning-opportunities/en/> (accessed on 25 June 2020).
53. Muller, C.; Yan, H. Household fuel use in developing countries: Review of theory and evidence. *Energy Econ. J.* **2018**, *70*, 429–439. [[CrossRef](#)]
54. Jones, L.L.; Hashim, A.; McKeever, T.; Cook, D.G.; Britton, J.; Leonardi-Bee, J. Parental and household smoking and the increased risk of bronchitis, bronchiolitis and other lower respiratory infections in infancy: Systematic review and meta-analysis. *Respir. Res.* **2011**, *12*, 1–11. [[CrossRef](#)] [[PubMed](#)]
55. Boadi, K.O.; Kuitunen, M. Factors affecting the choice of cooking fuel, cooking place and respiratory health in the Accra metropolitan area, Ghana. *J. Biosoc. Sci.* **2006**, *38*, 403–412. [[CrossRef](#)] [[PubMed](#)]
56. Kashima, S.; Yorifuji, T.; Tsuda, T.; Ibrahim, J.; Doi, H. Effects of traffic-related outdoor air pollution on respiratory illness and mortality in children, taking into account indoor air pollution, in Indonesia. *J. Occup. Environ. Med.* **2010**, *52*, 340–345. [[CrossRef](#)]
57. Nasir, Z.A.; Murtaza, F.; Colbeck, I. Role of poverty in fuel choice and exposure to indoor air pollution in Pakistan. *J. Integr. Environ. Sci.* **2015**, *12*, 107–117. [[CrossRef](#)]
58. Barria, M.R.P. Indoor Air Pollution by Particulate Matter from Wood Fuel: An Unresolved Problem. *Environ. Pollut. Clim. E Chang.* **2016**, *1*. [[CrossRef](#)]
59. Adam, J.C. Improved and more environmentally friendly charcoal production system using a low-cost retort-kiln (Eco-charcoal). *Renew. Energy* **2009**, *34*, 1923–1925. [[CrossRef](#)]
60. Njenga, M.; Karanja, N.; Munster, C.; Iiyama, M.; Neufeldt, H.; Kithinji, J.; Jamnadass, R. Charcoal production and strategies to enhance its sustainability in Kenya. *Dev. Pract.* **2013**, *23*, 359–371. [[CrossRef](#)]
61. Puzzolo, E.; Zerriffi, H.; Carter, E.; Clemens, H.; Stokes, H.; Jagger, P.; Rosenthal, J.; Petach, H. Supply considerations for scaling up clean cooking fuels for household energy in low- and middle-income countries. *GeoHealth* **2019**, *3*, 370–390. [[CrossRef](#)] [[PubMed](#)]
62. Adane, M.M.; Alene, G.D.; Mereta, S.T.; Wanyonyi, K.L. Prevalence and risk factors of acute lower respiratory infection among children living in biomass fuel using households: A community-based cross-sectional study in Northwest Ethiopia. *BMC Public Health* **2020**, *20*, 1–3. [[CrossRef](#)] [[PubMed](#)]
63. Geleta, D.; Tessema, F.; Ewnetu, H. Determinants of community acquired pneumonia among children in Kersa District, Southwest Ethiopia: Facility based case control study. *J. Pediatr. Neonatal Care* **2016**, *5*, 00179. [[CrossRef](#)]
64. Tazinya, A.A.; Halle-Ekane, G.E.; Mbuagbaw, L.T.; Abanda, M.; Atashili, J.; Obama, M.T. Risk factors for acute respiratory infections in children under five years attending the Bamenda Regional Hospital in Cameroon. *BMC Pulm. Med.* **2018**, *18*, 7. [[CrossRef](#)] [[PubMed](#)]
65. Taksande, A.M.; Yeole, M. Risk factors of acute respiratory infection (ARI) in under-fives in a rural hospital of Central India. *J. Pediatr. Neonatal Individ. Med.* **2016**, *5*, 50105. [[CrossRef](#)]
66. Rosenthal, J.; Quinn, A.; Grieshop, A.P.; Pillarisetti, A.; Glass, R.I. Clean cooking and the SDGs: Integrated analytical approaches to guide energy interventions for health and environment goals. *Energy Sustain. Dev.* **2018**, *42*, 152–159. [[CrossRef](#)] [[PubMed](#)]
67. Chidumayo, E.N.; Gumbo, D.J. The environmental impacts of charcoal production in tropical ecosystems of the world: A synthesis. *Energy Sustain. Dev.* **2013**, *17*, 86–94. [[CrossRef](#)]