

Higher levels of physical activity are associated with reduced tethering and migration of pro-inflammatory monocytes in males with central obesity

Wadley, Alex; Roberts, Matthew J.; Creighton, Jade; Thackray, Alice E.; Stensel, David J.; Bishop, Nicolette C.

License:

None: All rights reserved

Document Version

Peer reviewed version

Citation for published version (Harvard):

Wadley, A, Roberts, MJ, Creighton, J, Thackray, AE, Stensel, DJ & Bishop, NC 2021, 'Higher levels of physical activity are associated with reduced tethering and migration of pro-inflammatory monocytes in males with central obesity', *Exercise Immunology Review*, vol. 27, pp. 54-66.

<https://repository.lboro.ac.uk/articles/journal_contribution/Higher_levels_of_physical_activity_are_associated_with_reduced_tethering_and_migration_of_pro-inflammatory_monocytes_in_males_with_central_obesity/14717157>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This document is the Author Accepted Manuscript version of a published work which appears in its final form in *Exercise Immunology Review*. The final Version of Record can be found at: <http://eir-isei.de/2021/eir-2021-054-article.pdf>

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Higher levels of physical activity are associated with reduced tethering and migration of pro-inflammatory monocytes in males with central obesity

Alex J. Wadley^{1} Matthew J. Roberts¹ Jade Creighton^{1*} Alice E. Thackray¹ David J. Stensel¹ & Nicolette C. Bishop¹*

¹ National Centre for Sport and Exercise Medicine, School of Sport, Exercise and Health Sciences, Loughborough University, Epinal Way, Loughborough LE11 3TU

See current author affiliations in acknowledgements *

Keywords: Migration; obesity; leukocytes; exercise; immunity

Running Title: Physical activity, adiposity and monocyte migration

Word Count: 5015

Corresponding author:

Professor Nicolette Bishop
National Centre for Sport and Exercise Medicine
School of Sport, Exercise and Health
Loughborough University
Loughborough
Leicestershire
LE11 3TU

Email: n.c.bishop@lboro.ac.uk

Phone: 01509 226385

Abstract

Despite evidence that monocyte migration is accentuated by central adiposity, the impact of physical activity (PA) and exercise, particularly in the post-prandial state, on limiting migration are not established. We hypothesised that PA and a single bout of walking exercise would be associated with reduced *ex vivo* monocyte tethering and migration in middle-aged males with central obesity (CO). Objective levels of PA were measured for 7 days in lean males (LE, N=12, mean (SD) age 39 (10) years, waist circumference 81.0 (6.3) cm) and males with CO (N=12, mean (SD) age 40 (9) years, waist circumference 115.3 (13.9) cm), followed by donation of a fasted blood sample. On the same day, CO undertook a bout of walking exercise, before donation of a second fasted blood sample. An *ex vivo* assay, coupled to flow cytometry, determined tethering and migration of classical, intermediate, and non-classical monocytes. C-C and CXC chemokine receptor (CCR2, CCR5 and CX3CR1) expression were also determined on total and classical monocytes. Monocyte subsets (total, classical, intermediate and CCR2+ monocytes), metabolic (glucose and lipids) and inflammatory (C-reactive protein) markers were greater in CO vs. LE (lower high-density lipoprotein); however, adjustments for PA mitigated group differences for glucose, lipids, and monocyte subsets. *Ex vivo* tethering and migration (absolute and relative) of most monocyte subsets was greater in CO vs LE. Relative monocyte tethering and migration was largely not influenced by PA; however, higher PA was associated with reduced absolute migration and tethering of CD16 expressing monocytes in CO. Prior walking had no impact on these variables. These results highlight that regular PA, not single exercise bouts may limit the migration of pro-inflammatory monocytes in CO. These changes may relate to physiological parameters in blood (i.e. number of cells and their adhesion), rather than differences in chemokine receptor expression.

1 **Introduction**

2 Cardiovascular, metabolic, and neurological diseases are threatening to reach epidemic
3 proportions, presenting an enormous health, economic and societal challenge. Common to all
4 these conditions is their chronic inflammatory aetiology (40). Increased central adiposity is
5 known to significantly elevate the risk of developing these diseases, with a heightened
6 migration of immune cells from blood into various tissues, namely monocytes into adipose
7 tissue (44). An increase in adipose tissue-resident macrophages can initiate a perpetual
8 inflammatory cycle whereby proteins that tether (e.g. cellular adhesion molecules) (34), attract
9 (e.g. chemokines) (32), and subsequently cause damage (e.g. pro-inflammatory cytokines) (35)
10 are chronically elevated, thus drawing more cells into tissues. Over time, excessive monocyte
11 migration in individuals with central obesity (CO) drives a dysfunctional interaction that
12 promotes metabolic dysfunction and an increased risk of chronic disease (44).

13 This risk of developing cardiovascular, metabolic, and neurological diseases is exaggerated
14 by multiple factors, including physical inactivity and diet (41). For the latter, prolonged
15 elevations of triglycerides in the bloodstream after high fat meals, termed postprandial lipaemia
16 (PPL), is a known independent predictor of cardiovascular disease (41). There is evidence that
17 PPL is also associated with higher systemic inflammation (8, 41), with monocyte activation
18 (29), adhesion and propensity for foam cell formation higher following a high fat meal (28).
19 Given the central role of monocytes in the aetiology of chronic inflammatory disease,
20 monocyte blood profiles have been used to predict current and future cardiovascular (22),
21 metabolic (37), and neurological health (38). Recent evidence has documented that individuals
22 with obesity have a more pro-inflammatory and pro-migratory monocyte profile than
23 individuals who are lean (15, 33). This is reflected by higher concentrations, and an altered
24 composition of the three monocyte sub-populations; e.g. decreased classical (CM:
25 CD14⁺⁺CD16⁻), higher intermediate (IM: CD14⁺⁺CD16⁺), and higher non-classical (NCM:

26 CD14+CD16++) monocytes (15, 20, 33). A reduced percentage of CM occurs as a result of
27 preferential tissue migration in response to cues from inflamed tissues, whereas IM and NCM
28 proportions increase due to their role in patrolling the circulation by adhering to the
29 endothelium (31). These functions are governed by higher expression of C-C and CXC
30 chemokine receptors (e.g. CCR2, CCR5 and CXCR3), which positively correlate with body
31 weight, body mass index, fat mass, and insulin insensitivity on total monocytes ⁶. Chemokine
32 receptors have high affinity for complementary chemokine ligands (e.g. CCL2, CCL5 and
33 CXCL1), which are released from cells or tissues under metabolic and inflammatory stress.
34 Obesity-associated inflammation also causes an increase in circulating cellular adhesion
35 molecule concentration (34), which enhances the potential for monocytes to tether to the
36 surface of these cells and tissues. A higher number of circulating monocytes with a higher
37 capacity to tether and migrate therefore generates a favourable gradient for the progressive
38 movement of monocytes towards metabolically active tissues or the endothelial layer in
39 individuals with CO, increasing the risk of cardiovascular diseases (14). Given the substantial
40 economic and social impact of managing and treating these diseases, cost-effective
41 interventions to prevent monocyte tethering and migration are urgently required.

42 Increased levels of physical activity are associated with reduced blood markers of chronic
43 inflammation (e.g. inflammatory cytokines and adhesion molecules), and a reduced risk of
44 chronic inflammatory disease, compared with sedentary individuals (16, 24). Studies in adults
45 with obesity indicate that regular structured exercise can reduce the pro-inflammatory
46 monocyte phenotype in blood (13, 42), as well as the expression of chemokine receptors on
47 total monocytes (2). It is well documented that single sessions of exercise result in acute
48 elevations in chemokine concentrations in blood plasma (12, 17, 39). It has been suggested that
49 this response may result in the internalisation of chemokine receptors on the surface of immune
50 cells that with regular physical activity, drive a reduction in expression level (2, 6, 24). Indeed,

51 recent evidence has highlighted that moderate intensity exercise lowered the expression of
52 CCR2 on intermediate monocytes within an hour of exercise cessation in individuals with CO
53 (6). When considering how this transient internalisation of CCR2 on monocytes may translate
54 into chronic changes in expression that have been reported separately (2), the impact of PPL-
55 induced inflammation must also be considered, particularly in individuals with CO as this is
56 associated with unfavourable metabolic and inflammatory profiles (14). A large body of
57 evidence supports a role for single sessions of exercise being an effective strategy to attenuate
58 PPL within 24 hours of a high-fat meal (23). Despite this, there is limited available data to
59 support a reduction in PPL-associated inflammation over this 24 hour period with prior
60 exercise. At present, our collective understanding of how regular physical activity and single
61 sessions of exercise, particularly in the PPL state, independently impact chemokine expression
62 on monocyte populations in individuals with CO is lacking. Furthermore, how these changes
63 then relate to the tethering and migration of monocytes is yet to be established.

64 We, and others, have previously quantified the *ex vivo* migratory capacity of peripheral
65 blood mononuclear cells (PBMCs) towards a fixed chemokine gradient over time (3, 33). Using
66 this method, there is evidence that monocyte migration is greater in individuals who are obese
67 than those who are lean (33). Evidence from our group indicates that a single session of exercise
68 can reduce the *ex vivo* migration of T-helper lymphocytes towards chemokine-rich supernatants
69 in healthy individuals (3). To our knowledge, there is no data indicating that regular physical
70 activity or single sessions of exercise in a state of PPL, can reduce monocyte migration in
71 individuals with CO. Importantly, previous investigations into monocyte migration are limited
72 by only quantifying the number of cells that migrate in response to a fixed chemokine gradient
73 (33), rather than their phenotype, as done in our laboratory previously with lymphocytes (3).
74 Adopting this approach would indicate more about the inflammatory characteristics of
75 monocytes and potential to cause damage within metabolically active tissues (i.e. CM) or the

76 vasculature (i.e. IM and NCM). Furthermore, it has been highlighted that mimicking conditions
77 of physiological blood flow better maintains monocyte phenotype when implementing *ex vivo*
78 models (45), and this is often overlooked (3, 33).

79 By validating a dynamic experimental platform, which coupled *ex vivo* monocyte tethering and
80 migration under conditions of physiological blood flow to digital flow cytometry, the aims of
81 this study were to: 1) cross-sectionally determine the association of central adiposity and
82 physical activity levels with monocyte tethering and migration in middle-aged males and 2)
83 determine whether a single session of exercise can impact the tethering and migration of
84 monocytes in middle-aged males with CO under conditions of PPL.

85

86 **Materials and Methods**

87 **Participants**

88 This project involved cross-sectional assessments in males who were lean (LE, N=12)
89 and males with central obesity (CO, N=12), followed by a walking intervention in CO only.
90 Using our preliminary data, we based our sample size calculation on mean resting (and SD)
91 differences in total monocytes between CO and LE. Using GPower 3.1.9.7, we calculated we
92 would need 12 participants in each group to detect similar differences in the present study, with
93 an effect size of 1.1, 80% power and α of 5%. Age and ethnicity-matched participants (White
94 European (WE), N=6 and South Asian (SA), N=6 in both LE and CO) gave their informed
95 written consent and the investigation was approved by the ethical review committee at
96 Loughborough University (ethics code: R18-P120) in accordance with the Declaration of
97 Helsinki. Central obesity was defined as a waist circumference ≥ 94 cm in White European men
98 and ≥ 90 cm in South Asian men according to the International Diabetes Federation cut-off
99 points (1). Males who were lean were classified based on a waist circumference under the

100 aforementioned boundaries. All participants had stable weight for the preceding 3 months, were
101 non-smokers and had not taken any anti-inflammatory drugs (e.g. NSAIDs) for 4 weeks prior
102 to taking part. Participants were screened for diabetes using an HbA1c test, and a health screen
103 questionnaire was used to screen for other underlying health conditions, plus lifestyle factors
104 that may influence the results such as smoking. In addition, participants were required to refrain
105 from any strenuous physical activity or consumption of alcoholic or caffeine-based beverages
106 in the 48 hours prior to or during the experimental session.

107

108 Experimental Procedures

109 All participants first visited the laboratory for screening of height and weight using a
110 fixed wall stadiometer with a digital weighing scale built in (Seca Ltd, Hamburg, Germany).
111 Hip and waist circumference were measured using a flexible, non-elastic tape (Hokanson,
112 Washington, USA) whilst adhering to established measurement guidelines (1). An assessment
113 of peak oxygen uptake ($\dot{V}O_2$ peak) was undertaken in CO only using the modified Bruce
114 treadmill test (43). Heart rate was continuously measured using short-range telemetry (*Polar*
115 *T31; Polar Electro, Kempele, Finland*) and subjective effort measured using the rating of
116 perceived exertion scale (7). At the end of the first visit, participants were fitted with an
117 accelerometer (*ActiGraph GT3X, ActiGraph corporation, Florida, USA*) to quantify levels of
118 habitual physical activity for a period of 7 days. Data were analysed over 15 second epochs
119 using specialised software (*Actilife, Actigraph corporation, Florida, USA*). Accelerometer data
120 were screened for wear time using standard methods (10). Time spent in a defined intensity of
121 activity was determined by summing together counts per minute and categorising this based on
122 widely used cut points (19).

123 Prior to the experimental period, all participants were asked to maintain their normal
124 habitual diet for a period of 7 days. The day before, both groups undertook an overnight fast
125 from 22:00 (except plain water). The next morning (08:00), both groups returned to the
126 laboratory and consumed 250mL of water prior to bioelectrical impedance analysis (*Seca*
127 *mBCA 515, Seca Ltd, Hamburg, Germany*) to measure body fat percentage. Participants then
128 donated a blood sample via venepuncture to an antecubital vein. After this, CO remained rested
129 in the laboratory throughout the day, with standardised meals (57% fat, 32% carbohydrate,
130 11% protein, 14.2 kcal per kg of body mass) provided, before undertaking a 60-minute session
131 of walking exercise at 60-65% of their $\dot{V}O_2$ peak (15:00-16:00). Walking intensity was
132 confirmed using a portable metabolic cart which analysed breath-by-breath gases (*Metalyzer*
133 *3B, Cortex, Leipzig, Germany*) and subjective measures of perceived exertion were obtained
134 using the Borg Scale (6-20) (7). Participants were then free to leave the laboratory and were
135 provided with a standardised evening meal to consume before 22:00. After this, participants
136 fasted overnight (except plain water) and returned to the laboratory the next morning (08:00)
137 to donate a second rested blood sample.

138

139 Blood collection and analysis

140 Whole blood (40.9mL) was collected via venepuncture into EDTA (4.9mL) and sodium
141 heparin-coated (36mL) monovettes. Heparinised blood was used for the isolation of peripheral
142 blood mononuclear cells (PBMCs) by density gradient centrifugation. Briefly, whole blood
143 was diluted 1:1 with D-PBS and layered on top of Histopaque 1077 (2:1). Blood was
144 centrifuged at 400g for 30 minutes at 21 degrees (brake off) and the PBMC layer aspirated and
145 washed in PBS and RPMI. EDTA monovettes were centrifuged at 3500g for 10 minutes at 4
146 degrees and plasma isolated for future analysis of triacylglycerol (TAG), total cholesterol (TC),

147 high density lipoprotein (HDL), low density lipoprotein (LDL), non-esterified fatty acids
148 (NEFA), glucose, and C-Reactive Protein (CRP).

149

150 *Ex vivo* Migration Assay

151 The *ex vivo* migration assay was adapted from previously published protocols published
152 in our laboratory (3) and others (33), as well as internal validation experiments. Whole blood
153 cell counts were initially performed using a haematology analyser (*Yumizen H500, Horiba,*
154 *Northampton, UK*) to determine the circulating monocyte count ($\approx 0.2 - 1.0 \times 10^6/\text{mL}$). PBMC
155 counts were then determined by using CountBright™ Absolute Counting Beads on a BD
156 C6 Accuri Flow Cytometer (*Becton Dickinson, Oxford, UK*). The seeding density of monocytes
157 for the migration assay matched the monocyte concentration in whole blood, thus mimicking
158 physiological conditions. Peripheral blood mononuclear cells (in 2mL RPMI) were added in
159 duplicate to fibronectin-coated polyester (PET) inserts (5 μm pore size) and placed into 6-well
160 non-tissue culture treated plates containing 3mL of heat-inactivated 10% fetal bovine serum
161 (FBS) or RPMI (background migration control). Monocytes were then allowed to migrate for
162 3 hours at 37 degrees (5% CO₂). To further generate conditions that closely mimic moderate
163 physiological flow within the circulation (3.2 dyn/cm³) (9) and to maintain monocyte
164 phenotype, the PBMC suspension was oscillated on a 2-dimensional orbital shaker at 77 rpm
165 during this 3 hour period (45). Fetal bovine serum concentration and incubation times were
166 based on in-house validation experiments from our laboratory. After 3 hours, non-adherent
167 cells were removed from the upper side of the PET insert, and this was then washed twice with
168 1mL of D-PBS. *Tethered* PBMCs were then removed from the upper side of the PET inserts
169 by adding 1mL of enzyme free, EDTA-based dissociation media (4 degrees, 30 minutes),
170 followed by five washes with D-PBS (1ml). The underside of the PET inserts and the wells of

171 the tissue culture plate beneath were treated identically to collect cells that had *migrated* across
172 the PET insert. *Tethered* and *migrated* cells were collected into separate tubes and washed in
173 D-PBS ready for counting and phenotyping using flow cytometry.

174

175 Flow Cytometry

176 Pre-migration (baseline) and collected tethered and migrated PBMCs were counted as
177 above. For each sample, 1.75×10^5 PBMCs were then stained using fluorescently conjugated
178 antibodies for identification of monocytes subsets and chemokine receptor expression (e.g.
179 CCR2, CCR5 and CX3CR1) using four-colour flow cytometry. Cells were incubated with
180 CD14-FITC, CD16-PE, CX3CR1-APC, CCR5-APC, CCR2-Alexa Fluor-647 antibodies, and
181 7-AAD for 30 minutes at 4°C in the dark. Cells were then twice washed with FACS buffer (D-
182 PBS supplemented with 0.5% bovine serum albumin and 2 mM EDTA; pH=7.2) for 5 minutes
183 at 300 x g. Compensation was adjusted weekly by using single stained controls and gates
184 established using fluorescence minus one controls. Confirmation of non-specific antibody
185 binding was determined by using isotype-matched controls.

186 Flow cytometry data were analysed using BD C6 Accuri software (*Becton Dickinson,*
187 *Oxford, UK*). Briefly, monocytes were gated on forward versus side scatter. Doublets were
188 discriminated using FSC-A vs FSC-H plots, and non-viable monocytes (i.e. 7-AAD+)
189 excluded. CD14+ and CD16+ positive populations were then used to determine classical
190 (CD14++CD16-), intermediate (CD14++CD16+), and non-classical (CD14+CD16++)
191 monocyte proportions (%) as previously described (18). Histogram plots of the cells in the total
192 monocyte, and CM regions that positively expressed CCR2, CCR5, and CX3CR1 were
193 subsequently used to calculate the percentage of chemokine receptor positive cells. Mean
194 fluorescent intensity (MFI), an indicator of the density of chemokine receptor expression on

195 each cell, was also determined for total monocyte and CM populations. A sub-population of
196 cells was defined as ≥ 1000 positive cells in a gate. For baseline, the percentage of monocyte
197 subsets and CCR2+, CCR5+, and CX3CR1+ monocytes were used with whole blood cell
198 counts to determine the circulating number of chemokine receptor positive monocytes for each
199 sub-population. Similarly, the percentages of these cells following tethering and migration
200 were used with the total number of tethered and migrated cells to determine absolute changes
201 at each assay stage.

202

203 Metabolic Parameters

204 Concentrations of TC, HDL, LDL, TAG, glucose, CRP (Horiba Medical, Montpellier,
205 France), and NEFA (Randox Laboratories Ltd., County Antrim, UK) were determined
206 spectrophotometrically using commercially available kits (high detection for CRP) and a
207 benchtop analyser (Pentra 400, Horiba Medical, Montpellier, France). Concentrations of
208 HbA1c were determined using a bench top analyser (Quo-lab HbA1c POC, EKF Diagnostics,
209 Penarth, UK).

210

211 Data and Statistical Analyses

212 Data were analysed using the statistical package for social sciences (SPSS version 24).
213 Residuals for the outcomes were explored using histograms. Normally distributed data are
214 presented as the arithmetic mean (95% CIs). Skewed data underwent natural log-
215 transformation and were back transformed which gave a similar variance to non-logged data
216 and a reasonable estimate which is presented as the geometric mean (95% CI) (4). Log-
217 transformed data are presented as the ratio of geometric means and 95% CIs for the ratio
218 difference between geometric means (5). An effect size of 0.2 was considered the minimal

219 value for a meaningful difference, 0.5 for moderate and 0.8 for large (11). Statistical
220 significance was accepted as $P < 0.05$.

221 Physical and behavioural characteristics were compared using linear mixed models
222 (LMM) with waist circumference category (CO vs LE) and ethnicity (WE vs. SA) as fixed
223 factors. Blood and *ex vivo* migration variables were compared using LMM with the same fixed
224 factors mentioned above. These variables were then compared in CO only using LMM with
225 day (day 1 vs day 2) and ethnicity as fixed factors. Chemokine receptor expression was
226 compared before (baseline) and after the *ex vivo* assay (tethered and migrated) using LMM
227 with stage (baseline vs migrated vs tethered) as a fixed factor. Adjusted models were performed
228 to account for differences in moderate-vigorous PA (MVPA) and daily steps between CO and
229 LE. From here on adjustment refers to the adjusted model for daily steps and MVPA, and no
230 difference refers to a statistical difference.

231

232 **Results**

233 **Cross-sectional comparison of CO vs. LE**

234 Participant characteristics are displayed in Table 1. CO participants demonstrated
235 higher BMI, waist circumference, hip circumference, waist-to-hip ratio, body fat percentage,
236 and lean mass than the LE group ($ES \geq 2.30$, $P \leq 0.001$). CO demonstrated lower daily steps and
237 MVPA ($ES \geq 1.47$, $P \leq 0.001$).

238 There were no differences in participant characteristics between WE and SA within or
239 between CO and LE groups. Due to low numbers of participants and inconclusive data on
240 ethnic differences between CO and LE participants for the remaining variables, ethnic data are
241 not presented throughout the rest of the manuscript.

242

243 *[Insert Table 1 here]*

244

245 Metabolic markers

246 Unadjusted fasting metabolic marker concentrations are presented in Table 1.
247 Concentrations of HDL were lower (ES=1.22, P=0.003), and concentrations of TC, LDL, TAG,
248 glucose, CRP, and NEFA were higher in CO vs LE (ES \geq 0.93, P \leq 0.025). There was no
249 difference for HbA1c (ES=-0.43, P=0.100). Adjustment eliminated the difference for TC,
250 TAG, glucose, and NEFA (P \geq 0.101). No difference for HbA1c remained (P=0.265). HDL was
251 still lower, and LDL and CRP remained higher in CO vs. LE (P \leq 0.048).

252

253 Monocyte phenotype in blood

254 A representative gating strategy for monocyte subsets and their respective chemokine
255 receptor expression are presented in Figure 1. Unadjusted fasting blood monocyte subset
256 concentrations are presented in Table 1. Higher concentrations of monocytes, CM, IM, CCR2+
257 monocytes, and CCR2+CM subsets were seen in CO vs. LE (ES \geq 0.90, P \leq 0.037). No
258 differences for NCM, CCR5+, and CX3CR1+ monocytes were seen (ES \leq 0.77, P \geq 0.066).
259 Adjustment eliminated all differences for monocyte subset concentrations (P \geq 0.155). There
260 were no differences in the unadjusted or adjusted models for the percentages of monocyte
261 subsets (P \geq 0.154).

262

263 *[Insert Figure 1 here]*

264

265 Absolute and relative *ex vivo* migration

266 Unadjusted and adjusted absolute migrated monocyte subsets are presented in Figure 2
267 panel A. Higher absolute migration of monocytes, CM, IM, NCM, CCR2+ monocytes,
268 CCR2+CM, and CX3CR1+monocytes were seen in CO vs. LE (ES \geq 0.90, P \leq 0.046). No
269 difference was seen for the absolute migration of CCR5+ monocytes (ES=0.02, P=0.953).
270 Adjustment maintained higher absolute migration of monocytes, CM, CCR2+ monocytes, and
271 CCR2+ CM subsets (P \leq 0.023), eliminated the differences for IM, NCM, and CX3CR1+
272 monocytes (P \geq 0.465), and maintained no difference for CCR5+ monocytes (P=0.763).

273 Unadjusted and adjusted percentages of migrated monocyte subsets are presented in
274 Figure 2 panel B. Higher relative migration of CX3CR1+ monocytes were seen in CO vs. LE
275 (ES=1.01, P=0.017). No difference was seen for the relative migration of the other subsets
276 (ES \leq 0.85, P \geq 0.058). Adjustment eliminated the difference for CX3CR1+ monocytes
277 (P=0.091), but revealed higher relative migration of monocytes, CM, CCR2+ monocytes, and
278 CCR2+CM in CO (P \leq 0.040).

279

280 *[Insert Figure 2 here]*

281

282 Absolute and relative *ex vivo* tethering

283 Unadjusted and adjusted absolute tethered monocyte subsets are presented in Figure 3
284 panel A. Higher absolute tethering of monocytes, CM, IM, NCM, CCR2+ monocytes, and
285 CX3CR1+ monocytes were seen in CO vs. LE (ES \geq 0.92, P \leq 0.029). No difference was seen
286 for CCR5+ monocytes (ES=0.65, P=0.125). Adjustment maintained higher absolute tethering

287 of monocytes, CM, NCM, CCR2+ monocytes, and CCR2+CM in CO vs. LE ($P \leq 0.039$). The
288 differences for IM and CX3CR1M were eliminated ($P \geq 0.070$), and there was still no statistical
289 difference for CCR5+ monocytes ($P = 0.238$).

290 Unadjusted and adjusted percentages of tethered monocyte subsets are presented in
291 Figure 3 panel B. Higher relative tethering of IM, NCM, and CX3CR1+ was seen in CO vs.
292 LE ($ES \geq 1.11$, $P \leq 0.008$). No difference was seen for the relative tethering of other subsets
293 ($ES \leq 0.78$, $P \geq 0.057$). Adjustment maintained a higher relative tethering of IM, NCM, and
294 CX3CR1+ monocytes ($P \leq 0.003$), and revealed a higher relative tethering for monocytes, CM,
295 CCR2+ monocytes, and CCR2+CM in CO vs. LE ($P \leq 0.019$). No difference was maintained
296 for CCR5+ monocytes ($P = 0.437$).

297

298 *[Insert Figure 3 here]*

299

300 Chemokine receptor expression

301 Unadjusted expression of CCR2, CCR5, and CX3CR1 on total monocytes between CO
302 and LE at baseline, and at stages of the *ex vivo* assay (baseline, tethered and migrated) are
303 presented in Table 2. No difference for CCR2, CCR5, or CX3CR1 at baseline was found in the
304 unadjusted ($ES \leq 0.82$, $P \geq 0.060$) or adjusted model ($P \geq 0.081$). When comparing CO vs. LE, no
305 central obesity x stage interactions for CCR2, CCR5, or CX3CR1 were detected for the
306 unadjusted ($ES \leq 0.34$, $P \geq 0.743$) or adjusted ($P \geq 0.612$) models.

307 To explore changes in chemokine receptor expression during different stages of the *ex*
308 *vivo* assay in isolation, we evaluated changes in both groups combined. Unadjusted expression
309 of CCR2, CCR5, and CX3CR1 on total monocytes between stages of the assay for CO and LE

310 combined (N=24) revealed lower CCR2 receptor expression on migrated (mean difference = -
311 610, 95% CI, -1023 to -196) and tethered (mean difference = -644, 95% CI, -1058 to -231)
312 monocytes compared to baseline. No difference was found between the other stages for the
313 monocyte subsets (ES≤0.41, P≥0.073). Adjustment maintained the differences for CCR2
314 (P≤0.011), CCR5 and CX3CR1 receptor expression (P≥0.083).

315

316 *[Insert Table 2 here]*

317

318 **Walking exercise intervention**

319 Physiological Responses

320 Participants walked at 59% (SD=3) $\dot{V}O_2$ peak and consumed 1.87 (SD=0.67) litres of O₂ per
321 minute. Average heart rate was 126 beats per minute (SD=9) and a subjective rating of
322 perceived exertion score of 12 (SD=2).

323

324 Metabolic and monocyte phenotype in blood

325 Unadjusted concentrations of TAG, NEFA, and CRP were similar between day 1 and day 2
326 (ES≤0.33, P≥0.481). Unadjusted concentrations of glucose were lower on day 2 (ES=1.43,
327 P=0.003). Adjustment maintained no difference for TAG, NEFA, and CRP between day 1 and
328 day 2 (P≥0.457), and lower concentrations of glucose on day 2 (P=0.006).

329 Fasting concentrations of monocyte subsets and monocytes positive for CCR2, CCR5, and
330 CX3CR1 were similar between day 1 and day 2 for the unadjusted (ES≤0.21, P≥0.653) and

331 adjusted models ($P \geq 0.615$). There were no differences in the percentages of monocyte subsets
332 for the unadjusted ($ES \leq 0.23$, $P \geq 0.600$) and adjusted models ($P \geq 0.533$).

333

334 Absolute and relative ex vivo migration and tethering

335 The number of migrated monocyte subsets were similar between day 1 and day 2 for the
336 unadjusted ($ES \leq 0.37$, $P \geq 0.446$) and adjusted models ($P \geq 0.363$). Percentage of migrated
337 monocyte subsets and monocytes positive for CCR2, CCR5M, and CX3CR1 were similar
338 between day 1 and day 2 for the unadjusted ($ES \leq 0.45$, $P \geq 0.314$) and adjusted models
339 ($P \geq 0.244$).

340 The number of tethered monocyte subsets were similar between day 1 and day 2 for the
341 unadjusted ($ES \leq 0.49$, $P \geq 0.293$) and adjusted models ($P \geq 0.319$). The percentage of tethered
342 monocyte subsets and monocytes positive for CCR2, CCR5M and CX3CR1 were similar
343 between day 1 and day 2 for the unadjusted ($ES \leq 0.58$, $P \geq 0.130$) and adjusted ($P \geq 0.147$)
344 models.

345

346 Chemokine receptor expression

347 Unadjusted expression of CCR2, CCR5, and CX3CR1 on day 1 and day 2, and at different
348 stages of the *ex vivo* assay (baseline, tethered and migrated) are presented in Table 3. No
349 difference between days was seen for CCR2, CCR5, or CX3CR1 expression ($ES \leq 0.17$,
350 $P \geq 0.718$). Adjustment maintained no difference ($P \geq 0.749$). No day x stage interactions were
351 detected for the unadjusted ($ES \leq 0.15$, $P \geq 0.927$) or adjusted ($P \geq 0.878$) models.

352

353 *[Insert Table 3 here]*

354 **Discussion**

355 The results of the current study provide novel evidence for the independent associations
356 of adiposity, physical activity levels, and single sessions of exercise on aspects of obesity-
357 driven monocyte tethering and migration. By validating a dynamic methodological approach
358 that quantified the movement of monocytes towards chemokine-rich serum under conditions
359 of physiological blood flow, for the first time we established that both *absolute* and *relative*
360 monocyte tethering and migration were greater in CO vs. LE for most monocyte subsets.
361 Higher levels of physical activity (i.e. daily MVPA and step count) were associated with
362 reduced absolute tethering and migration of CD16 expressing monocytes, but not classical
363 monocytes. Under controlled laboratory conditions, a single bout of walking exercise had no
364 impact on monocyte tethering and migration in males with CO 16 hours after the exercise bout.
365 Taken together, these data indicate that regular physical activity was associated with reduced
366 movement of pro-inflammatory monocytes towards chemokines in males with CO. These
367 findings have important implications for the potential anti-inflammatory effects of physical
368 activity independent of weight status.

369 The movement of immune cells (e.g. monocytes, lymphocytes, and dendritic cells)
370 from blood into vascular walls and metabolically active tissues is enhanced by increased
371 adiposity (44). In particular, monocytes are metabolically plastic cells that can migrate into
372 tissues (i.e. CM) and transendothelial sites (i.e. IM and NCM) to form macrophages in
373 individuals with CO, which increases the risk of several chronic diseases (25, 31). The
374 enhanced movement of monocytes stimulates haematopoiesis in the bone marrow, resulting in
375 higher numbers of these monocytes in the circulation (30). In the present study, most
376 circulating monocyte counts were higher in CO vs. LE (Table 1). To subsequently determine
377 physiologically relevant differences in *ex vivo* monocyte tethering and migration (*absolute*
378 change), our method examined how monocytes at this concentration tethered and migrated

379 towards a fixed chemokine stimulus under conditions mimicking physiological blood flow
380 (45). This approach controlled for important systemic variables that are known to differ
381 between individuals who are lean and obese, e.g. the number of monocyte subsets and receptors
382 that govern their adhesion and subsequent migration. Our data demonstrate that *absolute* total
383 monocyte tethering and migration were greater in CO vs LE, independent of levels of physical
384 activity (Figures 2A and 3A). Within the monocyte pool, CM and those expressing CCR2
385 appeared to be the main subsets driving these group differences. This supports previous work
386 indicating that monocytes expressing CCR2 have the greatest propensity to migrate via
387 chemokine gradients towards inflamed tissues (31), and specifically adipose tissue in
388 individuals with obesity (36). Importantly, adjustments for MVPA and step count removed
389 group differences for the *absolute* migration of IM, NCM and monocyte expressing CX3CR1.
390 Group differences for the *absolute* tethering of IM and CX3CR1+ monocytes were also lost.
391 CD16 expressing monocytes, such as IM primarily migrate towards transendothelial sites and
392 are a major source of pro-inflammatory cytokines, such as tumour necrosis factor- α (TNF- α),
393 interleukin (IL)-1 β , IL-6, and IL-8 (46). These results therefore highlight an association
394 between higher physical activity levels and reduced IM tethering and migration. Adjustments
395 for PA also mitigated group differences in the blood concentration of IM, suggesting that
396 reduced tethering and migration may be driven by a reduced number of cells within the
397 circulation. This provides some support for other work highlighting reduced percentages of IM
398 and NCM after structured exercise training in individuals who are lean (42) and obese (13). In
399 addition, an impact of PA on the number of CX3CR1+ monocytes that tethered and migrated
400 also indicates reduced monocyte adhesion. These results therefore highlight that although PA
401 does not impact the migration of CM with high tissue homing-potential, it is associated with
402 reduced transendothelial migration of IM and NCM populations, a key step in the development
403 of chronic systemic inflammation.

404 In addition to the notable *absolute* differences between groups, *relative* tethering and
405 migration of most monocyte subsets were also higher in CO vs. LE (Figures 2B and 3B). This
406 suggests that differences at the cellular level are also important in driving monocyte migration.
407 With regards to chemokine receptors however, we observed no difference in the protein
408 expression of CCR2, CCR5, and CX3CR1 between LE and CO. Previous studies conducted in
409 females with obesity have reported higher expression of CCR2, CCR5, and CX3CR1 on
410 monocytes at both the mRNA (15) and protein level (15, 33). Despite similar screening (i.e. no
411 metabolic disease) and anthropometric measures in these studies, serum CRP levels were
412 markedly higher (mean: 7.0 (33) and 9.8 (15) mg/L) when compared to the present study
413 (mean: 1.8 mg/L), indicating a much more heightened state of systemic inflammation.
414 Furthermore, these studies noted differences in the relative numbers of blood monocyte subsets
415 between obese vs. lean (i.e. higher percentage of IM), also indicative of systemic inflammation.
416 To further interrogate our data, we explored changes in chemokine receptor expression after *ex*
417 *vivo* tethering and migration between CO and LE (Table 2). After binding to their cognate
418 chemokines, chemokine receptors are typically desensitized and internalized via endocytosis
419 to limit the magnitude and duration of the stimulus (21). In support of this, CCR2 receptor
420 expression was lower after *ex vivo* tethering and migration. This indicates that receptor
421 internalisation may have played a role in driving monocyte migration in our *ex vivo* model;
422 however, no statistical differences were noted between CO and LE. Collectively, we can
423 therefore speculate that in individuals with CO that have relatively low levels of systemic
424 inflammation, chemokine receptor expression and internalisation does not explain the higher
425 *relative* rates of tethering and migration vs. lean individuals. Although not measured in our
426 study, this again suggests that properties of monocyte adhesion could explain these differences.
427 Importantly, there were no associations between physical activity and chemokine receptor
428 expression and internalisation.

429 On the same day as the cross-sectional analysis, CO undertook a single bout of walking
430 exercise under controlled lifestyle and dietary conditions, with *ex vivo* monocyte tethering and
431 migration measured the morning afterwards (i.e. 24 hours after the first sample). This
432 experimental model has been previously used in obese populations to demonstrate that prior
433 brisk walking can lower postprandial lipaemia (PPL) after intake of a high fat meal (23). Acute
434 elevations in soluble inflammatory markers (i.e. IL-6 and TNF- α) also accompany PPL;
435 however there is limited evidence to support the notion that prior walking reduces PPL-
436 associated inflammation (8, 41). By applying our *ex vivo* assay to this experimental model, we
437 explored the impact of prior walking on changes in functional immunity for the first time. We
438 report no differences in circulating monocyte counts, *ex vivo* monocyte tethering and migration,
439 or chemokine receptor expression (absolute or migration mediated loss, Table 3) after prior
440 brisk walking. Previous studies have reported elevated chemokine concentrations in the
441 circulation after exercise (12, 17, 39). It has been suggested that these changes may drive
442 internalisation of chemokine receptors that lower their surface expression over time (24), in
443 turn reducing migration. Despite a recent study reported lower expression of CCR2 on IM
444 immediately after (but not 1h and 2h after) a bout of cycling exercise (6), our data importantly
445 highlight that any acute loss of monocyte chemokine receptor expression was not maintained
446 the morning afterwards.

447 The current results support an association between higher levels of physical activity
448 and reduced migration of specific pro-inflammatory monocytes, as well as important markers
449 of metabolic health (TC, TAG, glucose and NEFA, Table 1). Epidemiological studies report
450 that being more physically active is associated with reduced blood markers of inflammation
451 (26, 27); however it is unclear whether this is directly related to changes in adiposity (16, 24).
452 Our data provide further support for the notion that the movement of specific populations of
453 monocytes (i.e. IM) may be reduced independently of adiposity (2, 13). Given that our data

454 reveals no short-term impact of walking exercise on monocyte migration, future studies should
455 examine the impact of regular PA on immune cell migration, in the context of energy balance
456 and weight loss.

457 We must acknowledge some limitations to the current study. Our *ex vivo* method when
458 coupled to flow cytometry permits much more detailed phenotypic analysis of monocyte sub-
459 populations; however, analysis of rarer cell populations was challenging. For example, we did
460 not present data on chemokine receptor positive cells for IM and NCM due to low event counts
461 during acquisition (< 500). This was a limitation of the number of monocytes that tethered and
462 migrated, which was lower in LE vs. CO. Repeated blood measures after the bout of walking
463 exercise collected before 16 hours post-exercise may have been beneficial, but given assay
464 logistics and ethical considerations of blood volume, this was not possible. Finally, we must
465 acknowledge that the walking intervention was only carried out in CO, and therefore a direct
466 comparison with LE was not made; however, this would not have influenced the interpretations
467 of the present study.

468

469 **Conclusion**

470 The current data adds to the growing body of evidence highlighting that central
471 adiposity is a major driver of monocyte migration in individuals with CO. By exploring both
472 absolute and relative changes, we established that the concentration of monocytes in the
473 circulation of middle-aged males with CO may drive their heightened tethering and migration
474 towards an *ex vivo* chemokine stimulus, compared to lean individuals. Cellular level changes
475 likely also contribute (e.g. adhesion); however, we highlight that this was not chemokine
476 receptor-mediated in individuals with a relatively low level of systemic inflammation. Regular
477 physical activity did not impact the capacity of classical monocytes to tether and migrate;
478 however, specific pro-inflammatory subsets were reduced by MVPA and step count. Further

479 research is needed to establish the significance of these changes by monitoring individuals who
480 are lean and centrally obese over time and under controlled lifestyle interventions.

481

482 **Funding**

483 This research was supported by the National Institute for Health Research (NIHR)
484 Leicester Biomedical Research Centre. The views expressed are those of the authors and not
485 necessarily those of the NHS, the NIHR or the Department of Health.

486

487 **Author Contribution Statement**

488 AJW, MR, AET, DJS, and NCB were involved in the conception and design of the
489 study. AW and NCB developed the laboratory methods. MR undertook recruitment and
490 participant testing. AJW, JC and MR carried out all data acquisition. MR, AJW, and NCB
491 carried out statistical analysis and data presentation. Drafting of the article for important
492 intellectual content was undertaken by AJW and MR and all authors undertook revision and
493 final approval of the manuscript.

494

495 **Conflict of Interest**

496 None of the authors declare a conflict of interest.

497

498 **Current Author Affiliations (*)**

499 AW: School of Sport, Exercise & Rehabilitation Sciences, College of Life & Environmental
500 Sciences, University of Birmingham, B15 2TT

501 JC: Department of Health, University of Bath, Bath, BA2 7AY

502

503

504 **Table Legends**

505 **Table 1.** Groups were ethnicity matched, with a 1:1 ratio of WE and SA males. Data were
506 analysed using linear mixed models with the ethnic specific waist circumference category (CO
507 vs LE) as a fixed factor. Values are mean and standard deviations. 95% CI, 95% confidence
508 interval of the difference between the groups. *Main effect of central obesity ($P \leq 0.032$). ¹Body
509 fat % determined by bioelectrical impedance. ^aFor HbA1c, n=10 for the LE group.

510 MVPA, moderate to vigorous PA; TC, total cholesterol; HDL, high-density lipoprotein
511 cholesterol; LDL, low-density lipoprotein cholesterol; TAG, triacylglycerol; HbA1c, glycated
512 haemoglobin; CRP, C-reactive protein; NEFA, non-esterified fatty acids; CM, classical
513 monocytes; IM, intermediate monocytes; NCM, non-classical monocytes; CCR, C-C
514 chemokine receptor; CX3CR1, CX3C chemokine receptor-1

515

516 **Table 2.** Values for CCR2 and CX3CR1 are arithmetic means (95% confidence intervals) and
517 between group differences are absolute differences (95% confidence intervals). Values for
518 CCR5 are geometric means (95% confidence intervals) and between group differences are 95%
519 confidence limits of the ratio difference between geometric means.

520

521 **Table 3:** Values for CCR2 and CX3CR1 are arithmetic means (95% confidence intervals) and
522 between group differences are absolute differences (95% confidence intervals). Values for
523 CCR5 are geometric means (95% confidence intervals) and between group differences are 95%
524 confidence limits of the ratio difference between geometric means.

525

526

527 **Figure Legends**

528 **Figure 1:** Representative gating strategy for monocyte subsets and their respective chemokine
529 receptor expression levels. The example presented includes data from a male with central
530 obesity on *Day 1* of the study. The example data indicate how monocyte subsets and then
531 CCR2+ classical monocytes were determined: **(A)** Monocytes were gated on forward light
532 scatter (FSC) vs. side light scatter (SSC); **(B)** doublets were discriminated using FSC-Area vs.
533 FSC-Height; **(C)** non-viable cells were excluded using a 7-AAD; **(D)** monocyte subsets were
534 identified using a CD14-Area vs. CD16-Area bivariate plot; **(E)** the positive gate for CCR2+
535 classical monocytes was determined using a fluorescence minus one control; **(F)** CCR2+
536 classical monocytes (%) were determined on CCR2-Area vs. SSC-Area.

537 7-AAD, 7-Aminoactinomycin D; CM, classical monocytes; IM, intermediate monocytes;
538 NCM, non-classical monocytes; CCR2+ CM, C-C chemokine receptor-2+ classical
539 monocytes.

540

541 **Figure 2.** Unadjusted and adjusted absolute (Panel A) and relative (Panel B) monocyte subset
542 migration between men with central obesity (CO, N=12) and men who were lean (LE, N=12).
543 Linear mixed models were used with the ethnic specific waist circumference category (CO vs
544 LE) as a fixed factor (unadjusted). The adjusted model accounted for group differences in
545 moderate-vigorous physical activity (MVPA) and daily steps. Data are presented as the
546 arithmetic / geometric mean (95% confidence intervals).

547 CM, classical monocytes; CCR2+M, C-C chemokine receptor-2+ monocytes; CCR2+CM, C-
548 C chemokine receptor-2+ classical monocytes; IM, intermediate monocytes; NCM, non-
549 classical monocytes; CCR5+M, C-C chemokine receptor-5+ monocytes; CX3CR1+M, CX3C
550 chemokine receptor-1+ monocytes.

551 *Main effect of central obesity ($P \leq 0.046$).

552

553 **Figure 3.** Unadjusted and adjusted absolute (Panel A) and relative (Panel B) monocyte subset
554 tethering between men with central obesity (CO, N=12) and men who were lean (LE, N=12).
555 Linear mixed models were used with the ethnic specific waist circumference category (CO vs
556 LE) as a fixed factor (unadjusted). The adjusted model accounted for group differences in
557 moderate-vigorous physical activity (MVPA) and daily steps. Data are presented as the
558 arithmetic / geometric mean (95% confidence intervals).

559 CM, classical monocytes; CCR2+M, C-C chemokine receptor-2+ monocytes; CCR2+CM, C-
560 C chemokine receptor-2+ classical monocytes; IM, intermediate monocytes; NCM, non-
561 classical monocytes; CCR5+M, C-C chemokine receptor-5+ monocytes; CX3CR1+M, CX3C
562 chemokine receptor-1+ monocytes.

563 *Main effect of central obesity ($P \leq 0.039$).

564

565

566

567

568

569

570

571

572

573 **References**

574

575 1. **Alberti KGMM, Zimmet P, Shaw J.** International Diabetes Federation: a consensus
576 on Type 2 diabetes prevention. *Diabet Med* 24: 451–63, 2007. doi: 10.1111/j.1464-
577 5491.2007.02157.x.

578 2. **Barry JC, Simtchouk S, Durrer C, Jung ME, Little JP.** Short-Term Exercise
579 Training Alters Leukocyte Chemokine Receptors in Obese Adults. *Med Sci Sports*
580 *Exerc* 49: 1631–1640, 2017. doi: 10.1249/MSS.0000000000001261.

581 3. **Bishop NC, Walker GJ, Gleeson M, Wallace F a, Hewitt CR a.** 7. Human T
582 lymphocyte migration towards the supernatants of human rhinovirus infected airway
583 epithelial cells: influence of exercise and carbohydrate intake. *Exerc Immunol Rev* 15:
584 127–44, 2009.

585 4. **Bland JM, Altman DG.** Transformations, means, and confidence intervals. *BMJ* 312:
586 1079, 1996. doi: 10.1136/bmj.312.7038.1079.

587 5. **Bland JM, Altman DG.** The use of transformation when comparing two means. *BMJ*
588 312: 1153, 1996. doi: 10.1136/bmj.312.7039.1153.

589 6. **Blanks AM, Wagamon TT, Lafratta L, Sisk MG, Senter MB, Pedersen LN,**
590 **Bohmke N, Shah A, Mihalick VL, Franco RL.** Impact of physical activity on
591 monocyte subset CCR2 expression and macrophage polarization following moderate
592 intensity exercise. *Brain, Behav Immun - Heal* 2: 100033, 2020. doi:
593 10.1016/j.bbih.2019.100033.

594 7. **Borg G.** The psychophysical bases of peceived exertion. *Med Sci Sport Exerc* 14: 377–
595 381, 1982. doi: 10.1249/00005768-198205000-00012.

596 8. **Brown M, McClean CM, Davison GW, Brown JCW, Murphy MH.** Preceding
597 exercise and postprandial hypertriglyceridemia: Effects on lymphocyte cell DNA
598 damage and vascular inflammation. *Lipids Health Dis* 18: 1–12, 2019. doi:
599 10.1186/s12944-019-1071-y.

600 9. **Buchanan CF, Verbridge SS, Vlachos PP, Rylander MN.** Flow shear stress
601 regulates endothelial barrier function and expression of angiogenic factors in a 3D
602 microfluidic tumor vascular model. *Cell Adh Migr* 8: 517–524, 2014.

603 10. **Chomistek AK, Yuan C, Matthews CE, Troiano RP, Bowles HR, Rood J, Barnett**
604 **JB, Willet WC, Rimm EB, D.R. BJ.** Physical Activity Assessment with the
605 ActiGraph GT3X and Doubly Labeled Water. *Med Sci Sport Exerc* 49: 1935–1944,
606 2017.

607 11. **Cohen J.** *Statistical Power Analysis for the Behavioural Sciences, 2nd Edition.* 1988.

608 12. **Crabb EB, Franco RL, Caslin HL, Blanks AM, Bowen MK, Acevedo EO.** The

- 609 effect of acute physical and mental stress on soluble cellular adhesion molecule
610 concentration. *Life Sci* 157: 91–96, 2016. doi: 10.1016/j.lfs.2016.05.042.
- 611 13. **de Matos MA, Garcia BCC, Vieira DV, de Oliveira MFA, Costa KB, Aguiar PF,**
612 **Magalhães F de C, Brito-Melo GA, Amorim FT, Rocha-Vieira E.** High-intensity
613 interval training reduces monocyte activation in obese adults. *Brain Behav Immun* 80:
614 818–824, 2019. doi: 10.1016/j.bbi.2019.05.030.
- 615 14. **Després J-P, Lemieux I.** Abdominal obesity and metabolic syndrome. *Nature* 444:
616 881–7, 2006. doi: 10.1038/nature05488.
- 617 15. **Devêvre EF, Renovato-Martins M, Clément K, Sautès-Fridman C, Cremer I,**
618 **Poitou C.** Profiling of the Three Circulating Monocyte Subpopulations in Human
619 Obesity. *J Immunol* 194: 3917–3923, 2015. doi: 10.4049/jimmunol.1402655.
- 620 16. **Dungey M, Young HML, Churchward DR, Burton JO, Smith AC, Bishop NC.**
621 Regular exercise during haemodialysis promotes an anti-inflammatory leucocyte
622 profile. *Clin Kidney J* 10: 813–821, 2017. doi: 10.1093/ckj/sfx015.
- 623 17. **Fico BG, Whitehurst M, Slusher AL, Mock JT, Maharaj A, Dodge KM, Huang**
624 **C-J.** The comparison of acute high-intensity interval exercise vs. continuous moderate-
625 intensity exercise on plasma calprotectin and associated inflammatory mediators.
626 *Physiol Behav* 183: 27–32, 2018. doi: 10.1016/j.physbeh.2017.10.015.
- 627 18. **Figuroa-Vega N, Marín-Aragón CI, López-Aguilar I, Ibarra-Reynoso L, Pérez-**
628 **Luque E, Manuel Malacara J.** Analysis of the percentages of monocyte subsets and
629 ILC2s, their relationships with metabolic variables and response to hypocaloric
630 restriction in obesity. *PLoS One* 15: 1–19, 2020. doi: 10.1371/journal.pone.0228637.
- 631 19. **Freedson PS, Melanson E, Sirard J.** Calibration of the Computer Science and
632 Applications, Inc. Accelerometer. *Med Sci Sport Exerc* 30: 777–781, 1998.
- 633 20. **Friedrich K, Sommer M, Strobel S, Thrum S, Blüher M, Wagner U, Rossol M.**
634 Perturbation of the monocyte compartment in human obesity. *Front Immunol* 10: 1–
635 10, 2019. doi: 10.3389/fimmu.2019.01874.
- 636 21. **García Lopez MA, Aguado Martínez A, Lamaze C, Martínez-A. C, Fischer T.**
637 Inhibition of dynamin prevents CCL2-mediated endocytosis of CCR2 and activation of
638 ERK1/2. *Cell Signal* 21: 1748–1757, 2009. doi: 10.1016/j.cellsig.2009.07.010.
- 639 22. **Garofallo SB, Portal VL, Markoski MM, Dias LD, Quadrosa AS De, Marcadenti**
640 **A.** Correlations between traditional and nontraditional indicators of adiposity,
641 inflammation, and monocyte subtypes in patients with stable coronary artery disease. *J*
642 *Obes* 2019, 2019. doi: 10.1155/2019/3139278.
- 643 23. **Gill JMR, Al-Mamari A, Ferrell WR, Cleland SJ, Packard CJ, Sattar N, Petrie**
644 **JR, Caslake MJ.** Effects of prior moderate exercise on postprandial metabolism and
645 vascular function in lean and centrally obese men. *J Am Coll Cardiol* 44: 2375–82,

- 646 2004. doi: 10.1016/j.jacc.2004.09.035.
- 647 24. **Gleeson M, Bishop NC, Stensel DJ, Lindley MR, Mastana SS, Nimmo MA.** The
648 anti-inflammatory effects of exercise: mechanisms and implications for the prevention
649 and treatment of disease. *Nat Rev Immunol* 11: 607–615, 2011.
- 650 25. **Gómez-Olarte S, Bolaños NI, Echeverry M, Rodríguez AN, Cuéllar A, Puerta CJ,**
651 **Mariño A, González JM.** Intermediate Monocytes and Cytokine Production
652 Associated With Severe Forms of Chagas Disease. *Front Immunol* 10: 1671, 2019.
653 doi: 10.3389/fimmu.2019.01671.
- 654 26. **Hamer M, Hackett RA, Bostock S, Lazzarino AI, Carvalho LA, Steptoe A.**
655 Objectively assessed physical activity, adiposity, and inflammatory markers in people
656 with type 2 diabetes. *BMJ Open Diabetes Res Care* 2: e000030, 2014. doi:
657 10.1136/bmjdr-2014-000030.
- 658 27. **Hamer M, Sabia S, Batty GD, Shipley MJ, Tabak AG, Singh-Manoux A,**
659 **Kivimaki M.** Physical activity and inflammatory markers over 10 years follow up in
660 men and women from the Whitehall II cohort study. *Circulation* 126: 928–933, 2012.
661 doi: 10.1038/jid.2014.371.
- 662 28. **Henning AL, Venable AS, Vingren JL, Hill DW, McFarlin BK.** Consumption of a
663 high-fat meal was associated with an increase in monocyte adhesion molecules,
664 scavenger receptors, and Propensity to Form Foam Cells. *Cytometry B Clin Cytom* 94:
665 606–612, 2018. doi: 10.1002/cyto.b.21478.
- 666 29. **Hyson DA, Paglieroni TG, Wun T, Rutledge JC.** Postprandial lipemia is associated
667 with platelet and monocyte activation and increased monocyte cytokine expression in
668 normolipemic men. *Clin Appl Thromb Hemost* 8: 147–55, 2002. doi:
669 10.1177/107602960200800211.
- 670 30. **Idzkowska E, Eljaszewicz A, Miklasz P, Musial WJ, Tycinska AM, Moniuszko M.**
671 The Role of Different Monocyte Subsets in the Pathogenesis of Atherosclerosis and
672 Acute Coronary Syndromes. *Scand J Immunol* 82: 163–173, 2015. doi:
673 10.1111/sji.12314.
- 674 31. **Kapellos TS, Bonaguro L, Gemünd I, Reusch N, Saglam A, Hinkley ER, Schultze**
675 **JL.** Human monocyte subsets and phenotypes in major chronic inflammatory diseases.
676 *Front Immunol* 10: 1–13, 2019. doi: 10.3389/fimmu.2019.02035.
- 677 32. **Kim CS, Park HS, Kawada T, Kim JH, Lim D, Hubbard NE, Kwon BS, Erickson**
678 **KL, Yu R.** Circulating levels of MCP-1 and IL-8 are elevated in human obese subjects
679 and associated with obesity-related parameters. *Int J Obes* 30: 1347–1355, 2006. doi:
680 10.1038/sj.ijo.0803259.
- 681 33. **Krinninger P, Ensenuer R, Ehlers K, Rauh K, Stoll J, Krauss-Etschmann S,**
682 **Hauner H, Laumen H.** Peripheral Monocytes of Obese Women Display Increased
683 Chemokine Receptor Expression and Migration Capacity. *J Clin Endocrinol Metab* 99:

- 684 2500–2509, 2014. doi: 10.1210/jc.2013-2611.
- 685 34. **Miller MA, Cappuccio FP.** Cellular adhesion molecules and their relationship with
686 measures of obesity and metabolic syndrome in a multiethnic population. *Int J Obes*
687 30: 1176–1182, 2006. doi: 10.1038/sj.ijo.0803264.
- 688 35. **Park HS, Park JY, Yu R.** Relationship of obesity and visceral adiposity with serum
689 concentrations of CRP, TNF-alpha and IL-6. *Diabetes Res Clin Pract* 69: 29–35, 2005.
690 doi: 10.1016/j.diabres.2004.11.007.
- 691 36. **Pecht T, Haim Y, Bashan N, Shapiro H, Harman-Boehm I, Kirshtein B, Clément
692 K, Shai I, Rudich A.** Circulating Blood Monocyte Subclasses and Lipid-Laden
693 Adipose Tissue Macrophages in Human Obesity. *PLoS One* 11: 1–14, 2016. doi:
694 10.1371/journal.pone.0159350.
- 695 37. **Ratter JM, Rooijackers HMM, Jacobs CWM, de Galan BE, Tack CJ, Stienstra R.**
696 Hypoglycaemia induces recruitment of non-classical monocytes and cytotoxic
697 lymphocyte subsets in type 1 diabetes. *Diabetologia* 61: 2069–2071, 2018. doi:
698 10.1007/s00125-018-4683-2.
- 699 38. **Saresella M, Marventano I, Calabrese E, Piancone F, Rainone V, Gatti A,
700 Alberoni M, Nemni R, Clerici M.** A Complex Proinflammatory Role for Peripheral
701 Monocytes in Alzheimer’s Disease. *J Alzheimer’s Dis* 38: 403–413, 2013. doi:
702 10.3233/JAD-131160.
- 703 39. **Serviente C, Troy LM, de Jonge M, Shill DD, Jenkins NT, Witkowski S.**
704 Endothelial and inflammatory responses to acute exercise in perimenopausal and late
705 postmenopausal women. *Am J Physiol Regul Integr Comp Physiol* 311: R841–R850,
706 2016. doi: 10.1152/ajpregu.00189.2016.
- 707 40. **Szczepanska-Sadowska E, Cudnoch-Jedrzejewska A, Ufnal M, Zera T.** Brain and
708 cardiovascular diseases: common neurogenic background of cardiovascular, metabolic
709 and inflammatory diseases. *J Physiol Pharmacol* 61: 509–21, 2010.
- 710 41. **Teeman CS, Kurti SP, Cull BJ, Emerson SR, Haub MD, Rosenkranz SK.**
711 Postprandial lipemic and inflammatory responses to high-fat meals: A review of the
712 roles of acute and chronic exercise. *Nutr Metab* 13: 1–14, 2016. doi: 10.1186/s12986-
713 016-0142-6.
- 714 42. **Timmerman KL, Flynn MG, Coen PM, Markofski MM, Pence BD.** Exercise
715 training-induced lowering of inflammatory (CD14+CD16+) monocytes: a role in the
716 anti-inflammatory influence of exercise? *J Leukoc Biol* 84: 1271–8, 2008. doi:
717 10.1189/jlb.0408244.
- 718 43. **Trabulo M, Mendes M, Mesquita A, Seabra-Gomes R.** Does the Modified Bruce
719 Protocol Induce Physiological Stress Equal to That of the Bruce Protocol? *Rev Port*
720 *Cardiol* 13: 753–60, 1994.

- 721 44. **Trim W, Turner JE, Thompson D.** Parallels in immunometabolic adipose tissue
722 dysfunction with ageing and obesity. *Front Immunol* 9, 2018. doi:
723 10.3389/fimmu.2018.00169.
- 724 45. **Tsubota Y, Frey JM, Raines EW.** Technical Advance: Novel ex vivo culture method
725 for human monocytes uses shear flow to prevent total loss of transendothelial
726 diapedesis function. *J Leukoc Biol* 95: 191–195, 2014. doi: 10.1189/jlb.0513272.
- 727 46. **Wong KL, Tai JJ-Y, Wong W-C, Han H, Sem X, Yeap W-H, Kourilsky P, Wong**
728 **S-C.** Gene expression profiling reveals the defining features of the classical,
729 intermediate, and nonclassical human monocyte subsets. *Blood* 118: e16–e31, 2011.
730 doi: 10.1182/blood-2010-12-326355.

731

732