

## Exploring women's decisions of where to give birth in the Peruvian Amazon; why do women continue to give birth at home? A qualitative study

Gardiner, Esme; Lai, Jo Freda; Khanna, Divya; Meza, Graciella; De Wildt, Gilles; Taylor, Beck; Kiguli, Juliet

DOI:

[10.1371/journal.pone.0257135](https://doi.org/10.1371/journal.pone.0257135)

License:

Creative Commons: Attribution (CC BY)

### Document Version

Publisher's PDF, also known as Version of record

### Citation for published version (Harvard):

Gardiner, E, Lai, JF, Khanna, D, Meza, G, De Wildt, G, Taylor, B & Kiguli, J (ed.) 2021, 'Exploring women's decisions of where to give birth in the Peruvian Amazon; why do women continue to give birth at home? A qualitative study', *PLoS One*, vol. 16, no. 9, e0257135. <https://doi.org/10.1371/journal.pone.0257135>

[Link to publication on Research at Birmingham portal](#)

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

## RESEARCH ARTICLE

# Exploring women's decisions of where to give birth in the Peruvian Amazon; why do women continue to give birth at home? A qualitative study

Esme Gardiner<sup>1\*</sup>, Jo Freda Lai<sup>1</sup>, Divya Khanna<sup>1</sup>, Graciella Meza<sup>2</sup>, Gilles de Wildt<sup>1</sup>, Beck Taylor<sup>3</sup>

**1** College of Medical and Dental Sciences, University of Birmingham, Birmingham, United Kingdom, **2** Facultad de Medicina Human, Universidad Nacional de la Amazonía Peruana, Iquitos, Peru, **3** Institute of Applied Health Research, University of Birmingham, Birmingham, United Kingdom

\* [esmegardiner@btinternet.com](mailto:esmegardiner@btinternet.com)



## OPEN ACCESS

**Citation:** Gardiner E, Lai JF, Khanna D, Meza G, de Wildt G, Taylor B (2021) Exploring women's decisions of where to give birth in the Peruvian Amazon; why do women continue to give birth at home? A qualitative study. PLoS ONE 16(9): e0257135. <https://doi.org/10.1371/journal.pone.0257135>

**Editor:** Juliet Kiguli, Makerere University, School of Public Health, UGANDA

**Received:** November 22, 2020

**Accepted:** August 24, 2021

**Published:** September 10, 2021

**Peer Review History:** PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0257135>

**Copyright:** © 2021 Gardiner et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** This study is based on a dataset of 25 qualitative interview transcripts. However, the authors did not seek ethical

## Abstract

### Background

Despite improvements in maternal mortality globally, hundreds of women continue to die daily. The World Health Organisation therefore advises all women in low-and-middle income countries to give birth in healthcare facilities. Barriers to seeking intrapartum care have been described in Thaddeus and Maine's Three Delays Model, however these decisions are complex and often unique to different settings. Loreto, a rural province in Peru has one of the highest homebirth rates in the country at 31.8%. The aim of this study was to explore facilitators and barriers to facility births and explore women's experiences of intrapartum care in Amazonian Peru.

### Methods

Through purposive sampling, postnatal women were recruited for semi-structured interviews (n = 25). Interviews were transcribed verbatim and thematically analysed. A combination of deductive and inductive coding was used. Analytical triangulation was undertaken, and data saturation was used to determine when no further interviews were necessary.

### Results

Five themes were generated from the data: 1) Financial barriers; 2) Accessing care; 3) Fear of healthcare facilities; 4) Importance of seeking care and 5) Comfort and traditions of home. Generally, participants realised the importance of seeking skilled care however barriers persisted, across all areas of the Three Delays Model. Barriers identified included fear of healthcare facilities and interventions, direct and indirect costs, continuation of daily activities, distance and availability of transport. Women who delivered in healthcare facilities had mixed experiences, many reporting good attention, however a selection experienced poor treatment including abusive behaviour.

permission from the participants, nor the ethics committee, for the data to be used for anything other than this particular research study. The authors therefore do not have explicit permission for data sharing, re-analysis nor future studies and so would be inappropriate and unethical to make them available in the public domain. Furthermore, the data contains potentially identifying patient information. However, qualified individuals can direct queries by contacting Dr Ruth Riley ([r.riley@bham.ac.uk](mailto:r.riley@bham.ac.uk)) - chair of the University of Birmingham BMedSci Intercalation Internal Ethics Review Committee.

**Funding:** This work was supported by the University of Birmingham and an intercalated bursary award from the Topham Bursary Fund (EG). Support was also provided by the National Institute for Health Research (NIHR) Applied Research Centre (ARC) West Midlands (BT). The views expressed are those of the author and not necessarily those of the NIHR or the Department of Health and Social Care. The funders had no role in the study design, data collection and analysis, decision to publish or preparation of the manuscript.

**Competing interests:** The authors have declared that no competing interests exist.

**Abbreviations:** COREQ, consolidated criteria for reporting qualitative research; EsSalud, Social Health Insurance; IPC, Intrapartum Care; LMIC, Lower-middle Income Country; MDGs, Millennium Development Goals; MMR, Maternal Mortality Rate; SBA, Skilled Birth Attendant; SDGs, Sustainable Development Goals; SIS, Seguro Integral de Salud; TDM, Three Delays Model; WHO, World Health Organisation.

## Conclusion

Despite free care, women continue to face barriers seeking obstetric care in Amazonian Peru, including fear of hospitals, cost and availability of transport. However, women accessing care do not always receive positive care experiences highlighting implications for changes in accessibility and provision of care. Minimising these barriers is critical to improve maternal and neonatal outcomes in rural Peru.

## Introduction


Despite advances in recent years, maternal mortality remains a major global issue. The maternal mortality rate (MMR) is defined as the number of maternal deaths per 100,000 live births, caused by conditions related to or aggravated by pregnancy [1]. Despite aspirations within the Millennium Development Goals (MDGs) to reduce MMR by 75% by 2015, only a 38% reduction was achieved, and hundreds of women continue to die daily [2–4]. Many of these deaths are preventable, highlighting the importance of high quality antenatal and intrapartum care (IPC) [5]. The World Health Organisation (WHO) state that all women have a right to access high quality care during pregnancy and parturition, however less than half of births in low and middle-income countries (LMIC) are attended by healthcare professionals [6, 7]. Preventing these deaths is a key priority for the WHO, specifically addressing inequalities in access to and quality of maternity services [6, 8].

The Sustainable Development Goals (SDGs) aim to reduce MMR to less than 70 in every country and achieve universal access to skilled birth attendants (SBAs) [9]. SBAs are defined as “accredited health professionals who has been educated and trained to proficiency in the skills needed to manage an uncomplicated pregnancy and childbirth and identify, manage and refer complications” [10]. Clear evidence links the presence of SBAs with improved maternal outcomes [11–13]. Therefore, the WHO advises women to give birth in healthcare facilities, allowing access to SBAs and timely referrals if required [10].

The Three Delays Model (TDM), described by Thaddeus and Maine, outlines barriers to accessing maternity care in LMICs (Fig 1) [14]. These range from deciding to seek care to receiving adequate care. Understanding factors which prevent or facilitate women attending obstetric care is pivotal to achieve universal access to SBAs and prevent maternal deaths.

With an MMR of 68, Peru has achieved the SDGs, however when compared to other upper-middle income countries, Peru's statistic is 50% higher [15, 16]. The WHO and Pan American Health Organisation recently announced an aim to further reduce MMR to 65 by 2021. 93.1% of all births in Peru are attended by an SBA [9]. However, deliveries outside of healthcare settings, and therefore often without the presence of SBAs, are up to 9.3 times more common in rural areas [17]. Loreto is the largest region of Peru, located within the Amazon basin and has one of the highest rates of MMRs [119.5] and homebirths in the country at 31.8% [18, 19].

As far as the authors are aware, there are no publications in English, or with an English language abstract, about why women continue to give birth at home in Peru or Amazonia. Most of the global literature is based in Sub-Saharan Africa where MMRs are highest [4]. Barriers documented include lack of education about childbirth, cultural disparities with healthcare, distance, quick onset of labour and poor treatment from HCPs [20–32]. Cost is also often cited as a barrier for accessing IPC in many LMICs. Since 2002, Peru has had a Decentralised National Healthcare System to help achieve comprehensive healthcare for all the population



**Fig 1. The Three Delays Model: Thaddeus and Maine (1994).**

<https://doi.org/10.1371/journal.pone.0257135.g001>

[33]. The Ministry of Health provides free antenatal and IPC through the government funded *Seguro Integral de Salud* (SIS) programme for the poorest of the population and *Social Health Insurance* (EsSalud) for employees [34].

Marsland et al., conducted a study about women's perceptions of antenatal and IPC in Loreto; women reported free appointments facilitated antenatal care attendance and IPC was often poor, including discontinuity of care [35]. Westgard et al., also reported that women in rural Peru understood the importance of seeking obstetric care, particularly with complicated pregnancies [36]. However, no research has been conducted in Peru or Amazonia as to why women give birth at home. Literature from other countries has shown that these decisions are complex and multi-factorial, so in-depth qualitative research is required to help address barriers and improve uptake of obstetric care [37].

The aim of this study was to research why women in Loreto choose to give birth at home or in healthcare facilities. This involved exploring facilitators and barriers to health-facility births, to understand the multi-factorial reasons for the low uptake of IPC. Additionally, this study explored women's experiences of IPC in healthcare facilities to highlight possible improvements. Recent publication by the Regional Government of Loreto indicated key areas for health research, including exploring sociocultural factors that influence location of child-birth and reasons for delays in accessing obstetric care [38].

## Methods

### Ethical approval

Written ethical approval was provided locally by the Regional Directorate of Health, Loreto (385-2019-GRL-DRSL/30.09.01) and The University of Birmingham–BMedSc Population Sciences and Humanities Internal Research Ethics Committee (IREC2019/1548737). All participants provided written, informed consent prior to conducting the interviews.

### Methodological approach

This paper details an interpretive, descriptive qualitative study aligned with the *consolidated criteria for reporting qualitative research* (COREQ) checklist (S1 File) [39]. Qualitative methods were chosen to allow participants to express beliefs, feelings and motivations that underpin

their behaviour [40]. Semi-structured interviews were chosen over focus groups, to allow for potentially sensitive content [41, 42].

## Setting

Loreto, a large rural province in north-eastern Peru has one of the highest rates of homebirths in the country at 31.8% [17]. Despite Peru's economic development, Loreto continues to experience poverty and poor healthcare provision [43]. Iquitos, the capital of Loreto, and neighbouring villages are surrounded by the Amazon, Nanay and Itaya rivers; making them inaccessible by road [44, 45]. The majority of the population describe themselves as indigenous or descendants of indigenous people and the predominant language is Spanish (92.6%) [46]. The literacy rate of the population is 92.6% and over 75% of the population is Roman Catholic [47].

## Participant characteristics

Women who had given birth at home and in healthcare facilities were included in the study to allow exploration of both facilitators and barriers to facility births. The eligibility criteria are detailed in Table 1. Participant demographics were collected through a questionnaire to contextualise findings and allow reporting of characteristics (Table 2). Birth location was reported from participants' last pregnancy. However, an additional two women had previous experience of homebirths.

## Recruitment and sampling

Participants were recruited from January to February 2020, through a purposive method in primary healthcare centres within Iquitos and a door-to-door approach. The healthcare centres, Centro-de-Salud San Juan and Centro-de-Salud Moronacocha, serve the San Juan Bausitica and Iquitos regions respectively. These healthcare centres or '*postas*', provide an array of services including general medicine, paediatrics, dentistry and maternity care. All care is provided free of charge through the SIS. Participants were also recruited in six villages and communities within three hours of Iquitos, accessible by either road or boat (Fig 2). Due to time and financial restraints of the project, a pragmatic approach was taken, and other villages could not be visited. A visualisation of the recruitment process is detailed in S2 File.

## Data collection

Data was collected through face-to-face semi-structured interviews (n = 25, average length 34 minutes), including one pilot interview. Interviews were structured with a topic guide (S3 File), developed for this study from the TDM, other studies and discussion between authors [14, 25, 35, 49]. The topic guide was refined iteratively and used to ensure consistency between interviews whilst allowing the researcher flexibility to explore topics [50, 51]. Interviews were transcribed concurrently with data collection to allow a constant comparative approach.

**Table 1. Eligibility criteria.**

Inclusion Criteria	Exclusion Criteria
Women who had given birth within the previous 18 months	Women who did not have capacity to consent Serious illness or death of the new-born
Speak English or Spanish as their first language	
Permanent resident in Loreto	
Over the age of 18	

<https://doi.org/10.1371/journal.pone.0257135.t001>

Table 2. Socioeconomic characteristics of participants.

Characteristic	Number of participants (%)
<b>Location of most recent birth</b>	
Hospital	12 (48)
Posta	4 (16)
EsSalud	3 (12)
Home	6 (24)
<b>Age</b>	
18–24	8 (32)
25–29	8 (32)
30–34	6 (24)
≥35	3 (12)
<b>Parity</b>	
1	4 (16)
2	8 (32)
3	8 (32)
4	3 (12)
≥5	2 (8)
<b>Ethnicity</b>	
Mestizo	19 (76)
Other	6 (24)
<b>Education completed</b>	
No schooling complete	2 (8)
Primary	9 (36)
Secondary	10 (40)
Further education	4 (16)
<b>Occupation</b>	
Housewife	20 (80)
Student	2 (8)
Other	3 (12)

<https://doi.org/10.1371/journal.pone.0257135.t002>

All interviews were conducted in Spanish by EG via an experienced interpreter, independently recruited from the Universidad Nacional de la Amazonia Peruana. Neither the researcher nor interpreter were involved in the care of participants and the interpreter was asked to sign a confidentiality agreement. All interviews were digitally recorded and supplemented with reflective field notes.

## Data analysis

All interviews were transcribed verbatim into the English and Spanish segments by the principal research and interpreter respectively. Hybrid verbatim was chosen to ensure narrative flow whilst including fillers and interjections [52]. The Spanish was translated and compared against the English to assess the quality of translation and to ensure all interview data was included. Following the interviews, reflections and a list of interview topics were made. After 23 interviews, it was recognised that no additional issues were being discussed. A further two interviews were conducted, and it was decided by the research team that analytical saturation had been achieved [53, 54]. Recruitment and further data collection was then ceased.

Data was managed using NVIVO12 and thematically analysed following the 6-step approach described by Braun and Clarke [41]. A combination of deductive and inductive



**Fig 2. Map of recruitment sites including primary healthcare facilities and neighbourhoods.** Adaptation of image from Google Maps [48].

<https://doi.org/10.1371/journal.pone.0257135.g002>

coding was used; with concepts from the TDM labelled deductively whilst identifying other elements inductively [55]. A set of four transcripts were coded by two research assistants and compared to the principal researchers to ensure rigour [56]. Following the deductive and inductive rounds of coding, codes were arranged into themes and reviewed using a constant comparative approach [57]. The final themes were refined and named through discussion between authors.

## Results

Five key themes were developed from the data (Table 3). Extracts of data are presented alongside participant details; some quotations have been adapted to ensure participant anonymity. A further breakdown of the themes is available (S4 File).

### Theme 1—Financial barriers

Healthcare in Peru is supported by two health insurance schemes which provide free antenatal and IPC. Since the introduction of both schemes, rates of births with SBAs have increased

**Table 3. Themes and subthemes.**

Themes	Subthemes
Financial Barriers	Insurance
	Additional costs
	Poverty
Accessing care	Transport
	Distance
	Nature of labour–speed and pain of labour
Fear of Healthcare Facilities	Fear of hospital
	Fear of intervention
	Discomfort with hospital care
Importance of seeking care	Realisation of the importance of care
	Poor education
	Healthcare practitioner's advice
Comfort and Traditions of Home	Comfort of home
	Care available at home
	Continuation of daily activities
	Avoiding hospital experiences
	Autonomy

<https://doi.org/10.1371/journal.pone.0257135.t003>

[58]. Throughout the interviews, women suggested the availability of these systems encouraged facility births. However, for a handful of women, despite having access to either the SIS or EsSalud, other barriers arose, preventing women accessing this care; including indirect costs of facility-based deliveries.

*“I didn’t go to the hospital, I didn’t have any money. I mean, although in the hospital they don’t ask for money. . . I didn’t have enough for the ticket to get out of here.”*– 18yrs, Home

Furthermore, many of the women discussed insufficiencies in the system whereby additional costs were incurred, notably for medication.

*“Because in the hospital not. . . not everything is covered by SIS. . . SIS only covers the simple, most necessary thing you need. The doctors asked us for the medicine. . . we had to cover it from our pocket.”*– 21yrs, Hospital

These additional costs can be even greater when complications arise during labour:

*“Suddenly you get a. . . a few stitches or you get another disease and. . . that isn’t covered by the SIS”*– 39yrs, Home

The high rates of poverty in Loreto were also evident [19]. This combined with the unpredictable costs of IPC influenced women’s decision to choose to give birth at home without the assistance of SBAs.

*“I thought. . . that it costs a lot in the hospital, and at home I am not going to spend. . . I thought about it, the hospital is going to take a lot of money from me”*– 36yrs, Home

Despite the free IPC available to many women in Loreto, participants experienced and feared additional costs. This has resulted in financial barriers to accessing IPC, ultimately resulting in a few women deciding to give birth at home.



## Theme 2—Accessing care

Problems accessing care were discussed by many, including distance and cost of transport, as previously mentioned. In addition, the availability of transport influenced women's decision of where to give birth. Generally, women living within Iquitos reported good availability of 'motokars'. However, connectivity fell within a couple of kilometres of the city. In rural villages, women relied on the river and public boats for transport. This caused challenges to accessing care, made worse during the dry seasons.

*"Going in the month of October when the river was dry to get out of here is a little difficult, to go to the city. That's why I stopped going to the city, because the river is far away."*— 44yrs, Posta

Additionally, the river transport system made accessing IPC particularly difficult with the onset of labour at night when public boats do not run.

*"Only the boats work from. . . in the morning. From 4 am in the morning until 5 or 6 pm in the afternoon, from then on there are no boats. . . At night, there is no longer any available"*— 44yrs, Posta

Distance and travel time to healthcare facilities also influenced many women's decisions; one woman discussed the fear of delivering en route.

*"I told them the [labour] pains were coming quickly. . . and the ambulance won't come soon. It's more dangerous to deliver my baby on the road and so I have my baby here"*— 18yrs, Home

To enable a facility-based delivery, two women who lived in rural villages chose to travel to Iquitos prior to their delivery dates to ensure reaching healthcare facilities.

*"That's why I stayed there in Iquitos so I could get to the post faster."*— 44yrs, Posta

The speed and pain of labour also influenced many women's ability to access care.

*"I got a pain. . . I couldn't walk anymore so I had my son here in my house. I couldn't go to the posta anymore."*— 25yrs, Home

Accessing care presented as a major barrier to accessing IPC for many women. Within Iquitos, transport is easily accessible however, within a couple of miles of the city, transport services become less reliable and regular.

## Theme 3—Fear of healthcare facilities

Throughout many of the interviews, women expressed fear and unfamiliarity of healthcare facilities.

*"I have been told [to give birth in hospital], but I was afraid because I've never been to the hospital"*— 28yrs, Posta

One woman feared attending hospital due to the death of a child in hospital.

*"I feel like having [the baby] in my house, umm because going to the hospital, I feel shy since I lost my baby"*— 39yrs, Home

In addition, participants feared a hospital birth in which interventions were required and autonomy was lost, including a fear of 'cutting' and a desire for a 'normal' vaginal birth.

*"I wanted to have my baby boy this way without cuts, but no, I couldn't. . . they had already done an evaluation for me, and I was not going to be able to have a natural, normal baby. They told me. . . they were going to do a C-section"* – 29yrs, EsSalud

One woman also discussed how women's partners wanted them to have a natural delivery with the fear of deserting the woman if she required an intervention. This stemmed from a desire for women to have the 'courage' for a natural birth.

*[Why didn't you want the C-section?] "Because, no one is safe in life, suddenly my husband leaves me"* – 39yrs, Home

As well as this, participants often discussed feeling uncomfortable with other aspects of care associated with healthcare facilities, including exposure to HCPs and a fear of 'touching'.

*"They stick their finger in and they say. . . if the baby is coming or is about to come. . . in the house, you just feel the baby coming, and you tell the person helping that it's already born. . . but in the hospital, after a while they touch you again."* – 18yrs, Home

This woman directly compared the care provided to women at hospital to home, with less intervention occurring at home. Another woman also reported feeling uncomfortable with repeated examinations and implied a discomfort towards male HCPs.

*"They started to check me inside . . . it turns out that the baby was in a different position. Lots of doctors came in, first one comes in, touches me, inserts his fingers in the vagina, touches. . . then another doctor comes in. . . another doctor does the same thing, a male comes in, also the same thing, and two. . . two more doctors come in, just the same."* – 29yrs, Hospital

As well as direct inferences to fear of hospitals, many women discussed their experiences of hospital care, which can have implications for others decisions. A handful of women had negative experiences in hospital, including poor attention from HCPs.

*"I don't know if it's because the hospital is big, they don't give you adequate attention, they don't listen to you"* – 44yrs, Posta

Women also reported poor treatment from HCPs.

*"Some obstetricians don't treat you properly. . . I had this experience that they yelled at me, this one [obstetrician], went crazy because it hurt a lot"* – 20yrs, Hospital

Experiences like this could feed into other women's fears of hospital care. Finally, a selection of women feared separation from their babies during hospital stays.

*"While she was in the hospital, I was uncomfortable because she was out of my sight and I was in the hospital as a sick person, and my little girl was in the incubator. I didn't see her."* – 39yrs, Home

Many of the participants experienced a fear of healthcare facilities or interventions and practices deterring them from accessing care. However, a handful of women also experienced poor attention and treatment in facilities which could prevent them and others accessing care in the future.

#### **Theme 4—Importance of seeking care**

Throughout the interviews, women's awareness and knowledge of diseases in pregnancy and childbirth varied. Many women realised the importance of seeking IPC, discussing their fears of homebirths.

*"I have to be in a hospital because if I gave birth in a house or in some other place outside the hospital, my mother thought that suddenly something would happen to me, that is, she thought that suddenly I would die or my babies would"*— 21yrs, Hospital

Fear sometimes stemmed from stories they had heard about homebirths:

*"They arrived with their dead babies already. . . that's what I saw, so that's why I decided not to have it in my house. . . because it's going to happen to me. . . so I decided to go to hospital, for my safety and for my baby."*— 31yrs, Posta

Additionally, women talked about the availability of medication, equipment and HCPs in supporting their decision to give birth in healthcare facilities.

*"In an emergency, anything that you don't expect to happen, can happen in labour. . . the positive side is that you have specialists by your side, and they can take care of you. If you give birth at home, I think there is a higher death rate of pregnant women"*— 34yrs, Hospital

However, this awareness of risk was not universal, and a couple of women did not believe they were at risk when delivering at home.

*"My mum took care of all my sisters-in-law to have their babies at home, and that gave me the courage to have my baby here in my house because I knew nothing would go wrong."*— 18yrs, Home

Furthermore, other women who had homebirths believed that 'normal' antenatal appointments and a healthy pregnancy meant they did not need to attend hospital. Additionally, women believed that if the labour became complicated, they could then attend hospital.

*"If the baby is well, it's fine, if the baby is bad, we can go [to the hospital]."*— 39yrs, Home

However, the majority of women followed advice given at ANC appointments, particularly those with previously successful hospital births, and attended healthcare facilities for their delivery.

*"When I was going through my pregnancy appointments, the gynaecologist and obstetrician told me, in an emergency we can go to any health centre, the closest thing for the wellbeing of me and my baby."*— 20yrs, EsSalud

### Theme 5—Comfort and traditions of home

The final theme generated from the data was the comfort of homebirths and traditions that prevented women seeking help. This included having loved ones around and the familiarity of home.

*“I’m better off at home, I’m not worried that I’m in another bed, I know. . . I’m being treated, they’re giving me my warmth, my children, my husband.”*— 39yrs, Home

As well as this, many women discussed the traditional remedies that were available at home, with some women mentioning these could not be taken into hospitals or postas.

*“My sister-in-law was making me [the hot drink], so that my baby comes out quickly.”*  
*[What does it do?] “make the pain go away faster and to make the baby come out”*— 18yrs, Home

Additionally, all the woman who delivered at home discussed the normality of homebirths and due to previous successful home deliveries, they were more inclined to have another. Participants also discussed how after a homebirth it was possible for them to seek professional care by attending a posta within the following days for reassurance.

*“On the second day [after the birth] I go to the posta, they give him his vaccination and after 8 days I go for his check-ups.”*— 39yrs, Home

Giving birth at home also meant for many women that they could continue with day-to-day activities.

*“I’m having my babies normally, I have to take medicine, do my laundry, that’s all when you have them at home.”*— 36yrs, Home

For many, a key part of this was the ability to care for their other children.

*“In the home sometimes. . . they don’t have relatives who look after their children, and. . . that’s why they decide to have them in their home, because so many things are being known in these times, there is rape. . . that’s why I’m going to be thinking in the hospital, how is my daughter, how is my son?”*— 30yrs, Home

Finally, a couple of the women preferred giving birth at home for the autonomy it gave. As previously mentioned, women preferred a natural birth with the avoidance of interventions. Delivering at home also meant women had the ability to choose their birthing position.

*“It’s possible for you to have your baby lying down or sitting down, or squatting. . . I had my baby sitting down. That’s the difference. I mean, in the hospital, they make you lie down”*— 18yrs, Home

For many of the participants who gave birth at home, the comfort and familiarity played a large role in their decision making. This combined with other factors such as the cost and availability of transport and the fear of healthcare centres resulted in women choosing to have homebirths.

## Discussion

### Principal findings

Many barriers persist for women accessing IPC in Amazonian Peru, highlighting the complex and multifactorial nature of accessing SBAs in LMICs [14]. Despite free IPC provided by the government, many financial barriers were still cited. This included indirect costs of facility births, such as transport, and the additional costs of medication. Furthermore, the tradition of homebirths and fear of hospitals and interventions prevented women accessing care. Several other barriers were also identified including distance to healthcare facilities and fast onset of labour. Women who had delivered in healthcare facilities had mixed experiences, some discussing poor care, including poor attention and verbal abuse. Women's awareness of risks associated with childbirth were mixed; many seeking medical care for the safety of themselves and the new-born and others unaware of the importance of SBAs.

### Comparison with literature

Barriers within all aspects of Thaddeus and Maine's model were identified (Table 4). Many of the themes identified from the data confirms work from other settings, whilst adding new insight into barriers in rural Peru.

Financial barriers have been reported by women worldwide [21, 26, 32, 37, 59–61]. Due to the existence of the government funded SIS and EsSalud and recent literature from Loreto, the authors initially thought finances wouldn't be a key barrier [35]. However, costs, and the fear of hidden costs, were experienced by women including for transport and medication [62]. Several studies from other settings with free IPC also highlighted that women experienced similar issues [25, 27, 37, 63], including in Laos, where women were required to pay for medical equipment [31].

**Table 4. Summary of findings summarised within the Three Delays Model.**

Aspect of the Three Delays Model	Finding
Phase 1 delays—Decision to seek care	Direct and indirect costs of healthcare facility birth
	Fear and unfamiliarity of healthcare facilities
	Previous healthcare experiences
	Fear of interventions in hospital
	Poor awareness of risks and diseases associated with pregnancy
	Familiarity of home
	Normality of homebirth
	Availability of IPC at home
Phase 2 delays—Identifying and reaching medical facilities	Interruption to daily activities and childcare
	Cost of transport
	Distance/time to healthcare facility
	Fear of delivering en route to healthcare facilities
	Unavailability of transport
	Onset of labour at night—lack of transport
Phase 3 delays—Receipt of adequate and appropriate treatment	Reliance on public transport
	Speed and pain of labour
	Poor attention from HCPs
	Verbal abuse from HCPs
	Unfamiliarity with hospital care

<https://doi.org/10.1371/journal.pone.0257135.t004>

Many women's birth location decisions were driven by fear; women delivering at home feared hospitals and interventions, whilst women choosing to deliver in healthcare facilities feared the risks for mother and new-born with homebirths. The fear of hospital births has been found in other studies, including beliefs that every woman attending hospital for child-birth receives 'cutting' [37, 64]. Women favoured a natural birth and it was noted that women's partners also preferred women to deliver naturally. To the authors' knowledge this has not been reported before and contrasts other publications such as Pazandeh et al., who reported that women in Iran feared their future sexual appeal and satisfaction of their husbands, following pelvic floor injury with a natural delivery [65].

A selection of women did not seek care due to poor knowledge of risks. This has previously been documented, including the belief that hospital care is only necessary when experiencing obstetric complications [20, 23, 27, 29, 37, 63, 64, 66, 67]. The final phase 1 barrier, the comforts of home, also agreed with many findings amongst the global literature. One aspect included the ability to be cared by family members, in contrast to hospitals where they may not be permitted in the room [29, 30, 32, 64]. Furthermore, a study from Burkina Faso found women were unable to take traditional drinks into hospitals [21]. Likewise, in the Peruvian Amazon, Westgard et al., found women feared prenatal vitamins and preferred traditional remedies [36]. Other studies have also found women prefer homebirths to enable continuation of daily activities, including care for other children [37, 61]. However, the fear for the safety of their children whilst at hospital has not previously been identified as a barrier. Finally, women preferred giving birth at home to allow choice in birthing position. This confirms other studies where women said in hospital, they have to be in a supine position [20, 21, 30, 32]. Women in Rural Northern Ghana also reported this; More flexibility in birthing positions were possible with traditional birth attendants who allowed any position, given it would not harm the mother or baby [32].

Similarly, to Iquitos, transport is often reported by women as a primary reason for not being able to seek IPC, including unavailability of transport at night and lack of suitable transport [26, 31, 32, 37, 59, 60, 64, 68, 69]. A selection of studies also found that women had transport issues dependent on seasonal rainfall. In contrast to Loreto, where the dry season caused difficulties, the rainy season, including flooding and landslides, caused accessibility problems for other women elsewhere [21, 28, 32, 70].

A selection of women experienced negative treatment in healthcare facilities including poor attention and verbal abuse. These abusive behaviours are not confined to women in this setting with a recent paper published reporting that 41.6% of observed women in 4 LMICs experienced some form of abuse, stigma or discrimination [71]. Other qualitative papers note that poor attention or treatment in healthcare facilities influences women's future birth location decisions, both in high- and low-income settings [21, 26, 32, 37, 72].

Whilst similar research is being conducted in other LMICs and women are encouraged to deliver in healthcare facilities, many high-income countries are now supporting women to deliver at home [73, 74]. For low-risk women, this is being shown to be a safe choice with reduced rates of interventions and complications [75, 76]. However, for this to be safe, women need access to trained midwives, a good referral system and reliable transport.

## Strengths and limitations

As far as the authors are aware, this is the first study to explore barriers to facility births in Amazonian Peru. All women recruited had given birth in the prior 18 months, improving participant's recall. Furthermore, through a diverse recruitment strategy, women who had delivered at healthcare facilities and home were included to ensure facilitators and barriers were

explored. The presence of a local interpreter ensured comfort for participants and accurate translations of local dialects. To reduce the likelihood of misinterpretation, both the English and Spanish from the recordings were transcribed and compared [77, 78].

Due to time restraints of the project, it was not possible to do respondent validation. To improve the analysis of data and increase credibility, analyst triangulation was performed, and the final results were discussed between the authors [41, 79]. Cultural differences are likely to have impacted the data however the researcher made attempts to remain unbiased and reflexive throughout the process, including de-briefing with the local supervisor and a reflexive diary. Furthermore, interviews were conducted in settings where participants were comfortable, and a local interpreter was always present.

## Implications

Despite the provision of free IPC in Peru, women continue to face financial barriers. Until these cost barriers are removed it is likely that difficulties will persist. Further research needs to be conducted into the additional costs incurred by those in the Peruvian Amazon across different fields of healthcare and methods to eliminate or reduce them. Increasing the provision of care in primary healthcare centres could help reduce transport costs and distance, helping to achieve UHC for childbirth. However, providing UHC, requires both the utilisation of care provided and good quality care [80, 81]; Research has shown that delivering in healthcare facilities does not always improve maternal outcomes [82–84]. A recent publication modelled a service delivery redesign to ensure women's outcomes were maximised, by encouraging women to deliver in larger, better equipped hospitals [85]. To enable access to better resourced facilities, maternity waiting homes (MWH) could be introduced. MWHs are residences located near hospitals, enabling access to obstetric care and removing the unpredictability of onset of labour, similarly to participants in this study who stayed with relatives prior to parturition [86]. Prior to their establishment, a "needs assessment" would be required to assess the health services available and geographical inaccessibility and acceptability to the local population [86]. It would also be important to consider potential barriers, including care for children and cost of travelling to MWHs [87]. MWHs have been established in other parts of Peru, for example Cuzco, where the Ministry of Health provided training to ensure culturally acceptable care and rates of homebirths have subsequently fallen [88]. Although MWHs may not remove all barriers to facility births, primarily financial, they would help to reduce problems associated with distance and the unpredictable nature of labour. Additionally, other methods to tackle travel expenses should be considered to address financial and geographical inequalities, for example travel cards for those who do not have access to personal transport or additional funds from the SIS [89]. These could also be used in conjunction with MWHs.

In addition to this, as a result of the negative care experiences, staff need to ensure women are treated with dignity and respect. This includes restricting the frequency of vaginal examinations [90]. Furthermore, changes to practice need to be made to alleviate fears about hospitals and interventions, including educating and reassuring women about the reasons for interventions. Additionally, as recommended by the WHO, women need to be given autonomy during childbirth, including birthing position [90]. To help reduce the cultural differences with care in hospitals and reduce fear, women should be allowed a birthing partner of choice [90, 91]. Ideally infrastructure would also be amended to ensure mothers and newborns remain together postpartum or allowing access to neonatal care units.

Further research should be targeted in other parts of the Peruvian Amazon, particularly further away from Iquitos where fewer facilities are available and accessing SBAs is more challenging. Furthermore, research should be conducted to establish how best to educate women

about the risks of childbirth and the importance of SBAs. This study can also aid the development of safe motherhood initiative and public health policies [92].

## Conclusion

Despite the WHO encouraging women to deliver in healthcare facilities with SBAs, women in the Peruvian Amazon continue to face barriers accessing IPC [10]. Several barriers found in this setting concur with the global literature including fear of hospitals, lack of transport and financial hurdles. However, barriers unique to this setting were also found; fear for children's safety whilst in hospital and a fear of caesareans, partly due partners leaving women if interventions are required. Women in Loreto have also experienced abusive behaviour in health facilities. Changes to practice and facilities are required, including changes to the attitude of staff and modifications to ensure mothers and babies remain together postpartum. Further research should be conducted to assess the suitability of MWHs in the region and explore women's barriers to IPC in other parts of Loreto.

## Supporting information

**S1 File. Consolidated criteria for reporting qualitative studies (COREQ): 32-item checklist.**  
(PDF)

**S2 File. Visualisation of the recruitment process.**  
(PDF)

**S3 File. Summary of topic guide.**  
(PDF)

**S4 File. Themes, subthemes & codes breakdown.**  
(DOCX)

## Acknowledgments

We thank the healthcare staff in primary healthcare centres in Iquitos for helping in the recruitment of participants and the interpreter for helping with the conduction of interviews and transcription. Secondly, we would like to thank the women who gave their time to be involved in the research and share their experiences and beliefs.

## Author Contributions

**Conceptualization:** Esme Gardiner, Graciella Meza, Gilles de Wildt, Beck Taylor.

**Data curation:** Esme Gardiner, Beck Taylor.

**Formal analysis:** Esme Gardiner, Beck Taylor.

**Funding acquisition:** Esme Gardiner.

**Investigation:** Esme Gardiner.

**Methodology:** Esme Gardiner, Gilles de Wildt, Beck Taylor.

**Project administration:** Esme Gardiner, Graciella Meza, Gilles de Wildt.

**Resources:** Esme Gardiner, Graciella Meza.

**Software:** Esme Gardiner.

**Supervision:** Graciella Meza, Gilles de Wildt, Beck Taylor.



**Validation:** Jo Freda Lai, Divya Khanna, Beck Taylor.

**Visualization:** Esme Gardiner.

**Writing – original draft:** Esme Gardiner.

**Writing – review & editing:** Esme Gardiner, Jo Freda Lai, Divya Khanna, Graciella Meza, Gilles de Wildt, Beck Taylor.

## References

1. World Health Organisation. Maternal mortality ratio (per 100 000 live births) [Internet]. [cited 2019 Nov 15]. Available from: <https://www.who.int/healthinfo/statistics/indmaternalmortality/en/>
2. World Health Organisation. MDG 5: improve maternal health [Internet]. 2015 [cited 2020 Mar 17]. Available from: [https://www.who.int/topics/millennium\\_development\\_goals/maternal\\_health/en/](https://www.who.int/topics/millennium_development_goals/maternal_health/en/)
3. Sustainable Development Goals Knowledge Platform. Transforming our world: the 2030 Agenda for Sustainable Development [Internet]. [cited 2020 Mar 17]. Available from: <https://sustainabledevelopment.un.org/post2015/transformingourworld>
4. UNICEF. Maternal mortality [Internet]. 2019 [cited 2019 Oct 10]. Available from: <https://data.unicef.org/topic/maternal-health/maternal-mortality/>
5. Maternal health [Internet]. [cited 2020 Mar 17]. Available from: <https://www.who.int/westernpacific/health-topics/maternal-health>
6. World Health Organisation. Maternal mortality [Internet]. 2019 [cited 2019 Nov 15]. Available from: <https://www.who.int/news-room/fact-sheets/detail/maternal-mortality>
7. United Nations. SDG Indicators [Internet]. [cited 2020 Mar 17]. Available from: <https://unstats.un.org/sdgs/indicators/database/>
8. World Health Organisation. Strategies toward ending preventable maternal mortality (EPMM) [Internet]. 2015 [cited 2020 Mar 17]. Available from: <https://data.unicef.org/resources/strategies-toward-ending-preventable-maternal-mortality/>
9. Sustainable Development Goals Knowledge Platform. Sustainable Development Goals [Internet]. [cited 2020 Mar 17]. Available from: <https://sustainabledevelopment.un.org/?menu=1300>
10. World Health Organisation. Skilled birth attendants [Internet]. [cited 2019 Oct 11]. Available from: [https://www.who.int/reproductivehealth/topics/mdgs/skilled\\_birth\\_attendant/en/](https://www.who.int/reproductivehealth/topics/mdgs/skilled_birth_attendant/en/)
11. Graham WJ, Bell JS, Bullough CHW. Can skilled attendance at delivery reduce maternal mortality in developing countries? In: Safe Motherhood Strategies: A Review of the Evidence (eds. De Brouwere, V.; Van Lerberghe, W.), Studies in Health Services Organisation and Policy. In: Safe Motherhood Strategies: A Review of the Evidence (eds. De Brouwere, V.; Van Lerberghe, W.), Studies in Health Services Organisation and Policy. 2001
12. Scott S, Ronsmans C. The relationship between birth with a health professional and maternal mortality in observational studies: a review of the literature. *Trop Med Int Health* TM IH. 2009 Dec; 14(12):1523–33. <https://doi.org/10.1111/j.1365-3156.2009.02402.x> PMID: 19793070
13. Utz B, Siddiqui G, Adegoke A, Broek NVD. Definitions and roles of a skilled birth attendant: a mapping exercise from four South-Asian countries. *Acta Obstet Gynecol Scand*. 2013 Sep; 92(9):1063–9. <https://doi.org/10.1111/aogs.12166> PMID: 23656549
14. Thaddeus S, Maine D. Too far to walk: Maternal mortality in context. *Soc Sci Med*. 1994 Apr 1; 38(8):1091–110. [https://doi.org/10.1016/0277-9536\(94\)90226-7](https://doi.org/10.1016/0277-9536(94)90226-7) PMID: 8042057
15. World Health Organisation. Maternal mortality in 2000–2017 (Peru) [Internet]. [cited 2020 Mar 17]. Available from: [https://www.who.int/gho/maternal\\_health/countries/per.pdf?ua=1](https://www.who.int/gho/maternal_health/countries/per.pdf?ua=1)
16. The World Bank. Maternal mortality ratio (modeled estimate, per 100,000 live births) [Internet]. [cited 2020 Mar 17]. Available from: <https://data.worldbank.org/indicator/SH.STA.MMRT>
17. Instituto Nacional de Estadística e Informatic. Peru Encuesta Demografica y de Salud Familiar. 2017 [Internet]. [Cited 2019 Nov 19]. Available from: [https://www.inei.gob.pe/media/MenuRecursivo/publicaciones\\_digitales/Est/Lib1525/index.html](https://www.inei.gob.pe/media/MenuRecursivo/publicaciones_digitales/Est/Lib1525/index.html)
18. dl Carpio Ancaya L. Situation of maternal mortality in Peru, 2000–2012. *Rev Peru Med Exp Salud Publica*. 2013 Jul; 30(3):461–4. PMID: 24100823
19. Knoema. Peru—Loreto—Data and Statistics [Internet]. [cited 2020 Apr 17]. Available from: <https://knoema.com/atlas/Peru/Loreto>

20. Bedford J, Gandhi M, Admassu M, Girma A. 'A normal delivery takes place at home': a qualitative study of the location of childbirth in rural Ethiopia. *Matern Child Health J.* 2013 Feb; 17(2):230–9. <https://doi.org/10.1007/s10995-012-0965-3> PMID: 22359241
21. Some TD, Sombie I, Meda N. Women's perceptions of homebirths in two rural medical districts in Burkina Faso: a qualitative study. *Reprod Health.* 2011 Jan 28; 8:3. <https://doi.org/10.1186/1742-4755-8-3> PMID: 21276252
22. Kifle MM, Kesete HF, Gaim HT, Angosom GS, Araya MB. Health facility or home delivery? Factors influencing the choice of delivery place among mothers living in rural communities of Eritrea. *J Health Popul Nutr.* 2018 22; 37(1):22. <https://doi.org/10.1186/s41043-018-0153-1> PMID: 30348219
23. Kouanda S, Bado A, Meda IB, Yameogo GS, Coulibaly A, Haddad S. Home births in the context of free health care: The case of Kaya health district in Burkina Faso. *Int J Gynaecol Obstet Off Organ Int Fed Gynaecol Obstet.* 2016 Nov; 135 Suppl 1:S39–44. <https://doi.org/10.1016/j.ijgo.2016.08.009> PMID: 27836083
24. N'Gbichi C, Ziraba AK, Wambui DW, Bakibinga P, Kisiangani I, Njoroge P, et al. 'If there are no female nurses to attend to me, I will just go and deliver at home': a qualitative study in Garissa, Kenya. *BMC Pregnancy Childbirth.* 2019 Sep 10; 19(1):332. <https://doi.org/10.1186/s12884-019-2477-2> PMID: 31500582
25. Ochieng CA, Odhiambo AS. Barriers to formal health care seeking during pregnancy, childbirth and postnatal period: a qualitative study in Siaya County in rural Kenya. *BMC Pregnancy Childbirth.* 2019 Sep 18; 19(1):339. <https://doi.org/10.1186/s12884-019-2485-2> PMID: 31533640
26. Tancred T, Marchant T, Hanson C, Schellenberg J, Manzi F. Birth preparedness and place of birth in Tandahimba district, Tanzania: what women prepare for birth, where they go to deliver, and why. *BMC Pregnancy Childbirth.* 2016 16; 16(1):165. <https://doi.org/10.1186/s12884-016-0945-5> PMID: 27422526
27. Caulfield T, Onyo P, Byrne A, Nduba J, Nyagero J, Morgan A, et al. Factors influencing place of delivery for pastoralist women in Kenya: a qualitative study. *BMC Womens Health.* 2016 Aug 9; 16:1–11. <https://doi.org/10.1186/s12905-015-0282-2> PMID: 26729344
28. Kumbani L, Bjune G, Chirwa E, Malata A, Äyvind Odland J. Why some women fail to give birth at health facilities: a qualitative study of women's perceptions of perinatal care from rural Southern Malawi. *Reprod Health.* 2013 Jan; 10(1):9–20. <https://doi.org/10.1186/1742-4755-10-9> PMID: 23394229
29. Shah R, Rehfuess EA, Paudel D, Maskey MK, Delius M. Barriers and facilitators to institutional delivery in rural areas of Chitwan district, Nepal: a qualitative study. *Reprod Health.* 2018 Jun 20; 15(1):110. <https://doi.org/10.1186/s12978-018-0553-0> PMID: 29925398
30. Adatara P, Strumpher J, Ricks E, Mwini-Nyaledzigbor PP. Cultural beliefs and practices of women influencing home births in rural Northern Ghana. *Int J Womens Health.* 2019; 11:353–61. <https://doi.org/10.2147/IJWH.S190402> PMID: 31239788
31. Sato C, Phongluxa K, Toyama N, Gregorio ER, Miyoshi C, Nishimoto F, et al. Factors influencing the choice of facility-based delivery in the ethnic minority villages of Lao PDR: a qualitative case study. *Trop Med Health.* 2019; 47:50. <https://doi.org/10.1186/s41182-019-0177-2> PMID: 31516363
32. Adatara P, Strumpher J, Ricks E. Exploring the reasons why women prefer to give birth at home in rural northern Ghana: a qualitative study. *BMC Pregnancy Childbirth.* 2020 Aug 28; 20(1):500. <https://doi.org/10.1186/s12884-020-03198-y> PMID: 32859165
33. Pan American Health Organisation. Peru [Internet]. [cited 2020 Mar 17]. Available from: <https://www.paho.org/salud-en-las-americas-2017/?p=3232>
34. WHO | Peru [Internet]. [cited 2020 Mar 17]. Available from: <https://www.who.int/workforcealliance/countries/per/en/>
35. Marsland H, Meza G, de Wildt G, Jones L. A qualitative exploration of women's experiences of antenatal and intrapartum care: The need for a woman-centred approach in the Peruvian Amazon. East CE, editor. *PLOS ONE.* 2019 Jan 7; 14(1):e0209736. <https://doi.org/10.1371/journal.pone.0209736> PMID: 30615634
36. Westgard CM, Rogers A, Bello G, Rivadeneyra N. Health service utilization, perspectives, and health-seeking behavior for maternal and child health services in the Amazon of Peru, a mixed- methods study. *Int J Equity Health.* 2019 Oct 15; 18(1):155. <https://doi.org/10.1186/s12939-019-1056-5> PMID: 31615516
37. Bohren MA, Hunter EC, Munthe-Kaas HM, Souza JP, Vogel JP, Gülmezoglu AM. Facilitators and barriers to facility-based delivery in low- and middle-income countries: a qualitative evidence synthesis. *Reprod Health.* 2014 Sep 19; 11(1):71. <https://doi.org/10.1186/1742-4755-11-71> PMID: 25238684
38. Mera KC, Palomino YA, Alvarez CA, D'Orazi DG, Del Castillo LR, Vasquez RG, et al. Identificación de Prioridades Regionales de Investigación Para la Salud 2015–2021. Loreto: Dirección Regional de Salud de Loreto; 2014.

39. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *Int J Qual Health Care*. 2007 Dec 1; 19(6):349–57. <https://doi.org/10.1093/intqhc/mzm042> PMID: 17872937
40. Berkwits M, Inui TS. Making Use of Qualitative Research Techniques. *J Gen Intern Med*. 1998 Mar; 13(3):195–9. <https://doi.org/10.1046/j.1525-1497.1998.00054.x> PMID: 9541377
41. Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol*. 2006 Jan; 3(2):77–101.
42. Gill P, Stewart K, Treasure E, Chadwick B. Methods of data collection in qualitative research: interviews and focus groups. *Br Dent J*. 2008 Mar; 204(6):291–5. <https://doi.org/10.1038/bdj.2008.192> PMID: 18356873
43. Newman DE, Shapiro MC. Obstacles faced by general practitioners in Loreto Department, Peru in pursuing residency training. *Rural Remote Health*. 2010 Jun; 10(2):1256. PMID: 20707591
44. Lonely Planet. Iquitos travel [Internet]. [cited 2019 Oct 11]. Available from: <https://www.lonelyplanet.com/peru/amazon-basin/iquitos>
45. Instituto Nacional de Estadística e Informatic. Población. [Internet]. [cited 2020 Mar 18]. Available from: <https://www.inei.gob.pe/estadisticas/indice-tematico/poblacion-y-vivienda/>
46. Rullier HB, Tanco ED, Carrión DD, Portal JC, Samanez CA, Guillermo EM, et al. RESPONSABLES DEL ESTUDIO:308.
47. City Population. Iquitos (District, Peru) Population Statistics, Charts, Map and Location [Internet]. [cited 2020 Mar 18]. Available from: <https://www.citypopulation.de/php/peru-distr.php?adm2id=160101>
48. Iquitos [Internet]. Iquitos, Peru: Google Maps; 2020 [cited 2020 Sep 19]. Available from: <https://www.google.co.uk/maps/>
49. Naylor Smith J, Taylor B, Shaw K, Hewison A, Kenyon S. 'I didn't think you were allowed that, they didn't mention that.' A qualitative study exploring women's perceptions of home birth. *BMC Pregnancy Childbirth*. 2018 Apr 18; 18(1):105. <https://doi.org/10.1186/s12884-018-1733-1> PMID: 29669527
50. Hancock B, Ockleford E, Windridge K. An Introduction to Qualitative Research. *Qual Res*. 2009; 39.
51. Holloway I. and Galvin K., 2016. *Qualitative Research In Nursing And Healthcare*. 4th ed. Wiley-Blackwell.
52. Kim Y. The Pilot Study in Qualitative Inquiry: Identifying Issues and Learning Lessons for Culturally Competent Research. *Qual Soc Work*. 2011 Jun 1; 10(2):190–206.
53. Saunders B, Sim J, Kingstone T, Baker S, Waterfield J, Bartlam B, et al. Saturation in qualitative research: exploring its conceptualization and operationalization. *Qual Quant*. 2018 Jul; 52(4):1893–907. <https://doi.org/10.1007/s11135-017-0574-8> PMID: 29937585
54. Hennink M, Kierser B, Marconi V. Code Saturation Versus Meaning Saturation: How Many Interviews Are Enough?. *SAGE*. 2017; 27(4):591–608.
55. Punch KF, Oancea A, 2014. *Introduction to Research Methods in Education*. 2nd ed. SAGE Publications
56. Barbour RS. Checklists for improving rigour in qualitative research: a case of the tail wagging the dog? *BMJ*. 2001 May 5; 322(7294):1115–7. <https://doi.org/10.1136/bmj.322.7294.1115> PMID: 11337448
57. Fugard A, Potts HWW. Thematic Analysis. *SAGE Research Methods* [Internet]. 2019 [cited 2020 Apr 20]. Available from: <https://methods.sagepub.com/foundations/thematic-analysis>
58. World Health Organisation. Peru Country Profile [Internet]. [cited 2020 Apr 17]. Available from: [https://www.who.int/maternal\\_child\\_adolescent/events/2008/mdg5/countries/final\\_cp\\_peru\\_19\\_09\\_08.pdf](https://www.who.int/maternal_child_adolescent/events/2008/mdg5/countries/final_cp_peru_19_09_08.pdf)
59. Shiferaw S, Spigt M, Godefrooij M, Melkamu Y, Tekie M. Why do women prefer home births in Ethiopia? *BMC Pregnancy Childbirth*. 2013 Jan 16; 13(1):5. <https://doi.org/10.1186/1471-2393-13-5> PMID: 23324550
60. Mason L, Dellicour S, Ter Kuile F, Ouma P, Phillips-Howard P, Were F, et al. Barriers and facilitators to antenatal and delivery care in western Kenya: a qualitative study. *BMC Pregnancy Childbirth*. 2015 Feb 13; 15(1):26. <https://doi.org/10.1186/s12884-015-0453-z> PMID: 25886593
61. Milne L, van Teijlingen E, Hundley V, Simkhada P, Ireland J. Staff perspectives of barriers to women accessing birthing services in Nepal: a qualitative study. *BMC Pregnancy Childbirth*. 2015 Jul 2; 15(1):142. <https://doi.org/10.1186/s12884-015-0564-6> PMID: 26133977
62. Amnesty International. Fatal Flaws—Barriers to Maternal health in Peru. 2008 [cited 2020 Apr 20]. Available from: <https://www.amnesty.org/download/Documents/44000/amr460082009eng.pdf>
63. Moshi F, Nyamhanga T. Understanding the preference for homebirth; an exploration of key barriers to facility delivery in rural Tanzania. *Reprod Health*. 2017 Oct 17; 14(1):132. <https://doi.org/10.1186/s12978-017-0397-z> PMID: 29041972

64. Roro MA, Hassen EM, Lemma AM, Gebreyesus SH, Afework MF. Why do women not deliver in health facilities: a qualitative study of the community perspectives in south central Ethiopia? *BMC Res Notes*. 2014; 7: 556. <https://doi.org/10.1186/1756-0500-7-556> PMID: 25143017
65. Pazandeh F, Potrata B, Huss R, Hirst J, House A. Women's experiences of routine care during labour and childbirth and the influence of medicalisation: A qualitative study from Iran. *Midwifery*. 2017 Oct; 53:63–70. <https://doi.org/10.1016/j.midw.2017.07.001> PMID: 28763721
66. Titaley CR, Hunter CL, Dibley MJ, Heywood P. Why do some women still prefer traditional birth attendants and home delivery?: a qualitative study on delivery care services in West Java Province, Indonesia. *BMC Pregnancy Childbirth*. 2010 Aug 11; 10(1):43. <https://doi.org/10.1186/1471-2393-10-43> PMID: 20701762
67. Kawakatsu Y, Sugishita T, Oruenjo K, Wakhule S, Kibosia K, Were E, et al. Determinants of health facility utilization for childbirth in rural western Kenya: cross-sectional study. *BMC Pregnancy Childbirth*. 2014 Aug 9; 14(1):265. <https://doi.org/10.1186/1471-2393-14-265> PMID: 25106432
68. Lerberg PM, Sundby J, Jammeh A, Fretheim A. Barriers to skilled birth attendance: a survey among mothers in rural Gambia. *Afr J Reprod Health*. 2014 Mar; 18(1):35–43. PMID: 24796167
69. Kisiangani I, Elmi M, Bakibinga P, Mohamed SF, Kisia L, Kibe PM, et al. Persistent barriers to the use of maternal, newborn and child health services in Garissa sub-county, Kenya: a qualitative study. *BMC Pregnancy Childbirth*. 2020 May 7; 20(1):277. <https://doi.org/10.1186/s12884-020-02955-3> PMID: 32380975
70. Belton S, Myers B, Ngana FR. Maternal deaths in eastern Indonesia: 20 years and still walking: an ethnographic study. *BMC Pregnancy Childbirth*. 2014 Jan 22; 14(1):39.
71. Bohren MA, Mehrtash H, Fawole B, Maung TM, Balde MD, Maya E, et al. How women are treated during facility-based childbirth in four countries: a cross-sectional study with labour observations and community-based surveys. *The Lancet*. 2019 Nov; 394(10210):1750–63. [https://doi.org/10.1016/S0140-6736\(19\)31992-0](https://doi.org/10.1016/S0140-6736(19)31992-0) PMID: 31604660
72. Bohren MA, Vogel JP, Hunter EC, Lutsiv O, Makh SK, Souza JP, et al. The Mistreatment of Women during Childbirth in Health Facilities Globally: A Mixed-Methods Systematic Review. *PLoS Med*. 2015 Jun; 12(6):e1001847; discussion e1001847. <https://doi.org/10.1371/journal.pmed.1001847> PMID: 26126110
73. Nygaard SS, Kesmodel US. Home births—Where are we heading? *Acta Obstet Gynecol Scand*. 2018 Oct 1; 97(10):1155–6. <https://doi.org/10.1111/aogs.13441> PMID: 30198112
74. The National Institute for Health and Care Excellence. Choosing where to have your baby—Information for the public [Internet]. 2014 [cited 2020 Apr 29]. Available from: <https://www.nice.org.uk/guidance/cg190/ifp/chapter/Choosing-where-to-have-your-baby>
75. de Jonge A, Mesman J, Mannien J, Zwart J, van Dillen J, van Roosmalen J. Severe adverse maternal outcomes among low risk women with planned home versus hospital births in the Netherlands: nationwide cohort study. *BMJ*. 2013; 346(jun13 2):f3263–f3263. <https://doi.org/10.1136/bmj.f3263> PMID: 23766482
76. Hollowell J, Rowe R, Townend J, Knight M, Li Y, Linsell L, et al. The Birthplace in England national prospective cohort study: further analyses to enhance policy and service delivery decision-making for planned place of birth. *Health Serv Deliv Res*. 2015 Aug; 3(36):1–264. <https://doi.org/10.3310/hsdr03360> PMID: 26334076
77. Squires A, Sadarangani T, Jones S. Strategies for overcoming language barriers in research. *J Adv Nurs*. 2020 Feb; 76(2):706–14. <https://doi.org/10.1111/jan.14007> PMID: 30950104
78. Squires A. Methodological challenges in cross-language qualitative research: A research review. *Int J Nurs Stud*. 2009 Feb 1; 46(2):277–87. <https://doi.org/10.1016/j.ijnurstu.2008.08.006> PMID: 18789799
79. Nowell LS, Norris JM, White DE, Moules NJ. Thematic Analysis: Striving to Meet the Trustworthiness Criteria. *Int J Qual Methods*. 2017 Dec 1; 16(1):1609406917733847.
80. Kruk ME, Gage AD, Joseph NT, Danaei G, García-Saisó S, Salomon JA. Mortality due to low-quality health systems in the universal health coverage era: a systematic analysis of amenable deaths in 137 countries. *The Lancet*. 2018 Nov 17; 392(10160):2203–12. [https://doi.org/10.1016/S0140-6736\(18\)31668-4](https://doi.org/10.1016/S0140-6736(18)31668-4) PMID: 30195398
81. Das J, Woskie L, Rajbhandari R, Abbasi K, Jha A. Rethinking assumptions about delivery of healthcare: implications for universal health coverage. *BMJ* [Internet]. 2018 May 21 [cited 2020 Apr 29];361. Available from: <https://www.bmj.com/content/361/bmj.k1716> <https://doi.org/10.1136/bmj.k1716> PMID: 29784870
82. Godlonton S, Okeke EN. Does a ban on informal health providers save lives? Evidence from Malawi. *J Dev Econ*. 2016 Jan 1; 118:112–32. <https://doi.org/10.1016/j.jdeveco.2015.09.001> PMID: 26681821

83. Powell-Jackson T, Mazumdar S, Mills A. Financial incentives in health: New evidence from India's Janani Suraksha Yojana. *J Health Econ*. 2015 Sep; 43:154–69. <https://doi.org/10.1016/j.jhealeco.2015.07.001> PMID: 26302940
84. Mohanan M, Bauhoff S, La Forgia G, Babiarz KS, Singh K, Miller G. Effect of Chiranjeevi Yojana on institutional deliveries and neonatal and maternal outcomes in Gujarat, India: a difference-in-differences analysis. *Bull World Health Organ*. 2014 Mar 1; 92(3):187–94. <https://doi.org/10.2471/BLT.13.124644> PMID: 24700978
85. Gage AD, Carnes F, Blossom J, Aluvaala J, Amatya A, Mahat K, et al. In Low- And Middle-Income Countries, Is Delivery In High-Quality Obstetric Facilities Geographically Feasible? *Health Aff (Millwood)*. 2019 Sep 1; 38(9):1576–84. <https://doi.org/10.1377/hlthaff.2018.05397> PMID: 31479351
86. World Health Organisation. Maternity Waiting Homes: A review of experiences [Internet]. 1996 [cited 2020 Apr 29]. Available from: [https://apps.who.int/iris/bitstream/handle/10665/63432/WHO\\_RHT\\_MSM\\_96.21.pdf?sequence=1](https://apps.who.int/iris/bitstream/handle/10665/63432/WHO_RHT_MSM_96.21.pdf?sequence=1)
87. Hodin S. Maternal Health Task Force. Maternity Waiting Homes: A Viable Solution for Rural Women? [Internet]. 2017 [cited 2020 Apr 29]. Available from: <https://www.mhtf.org/2017/11/08/maternity-waiting-homes-a-viable-solution-for-rural-women/>
88. UNICEF. Adapting maternity services to the cultures of rural Peru [Internet]. [cited 2020 Apr 29]. Available from: <https://www.unicef.org/sowc09/docs/SOWC09-Panel-2.5-EN.pdf>
89. Montagu D, Sudhinaraset M, Diamond-Smith N, Campbell O, Gabrysch S, Freedman L, et al. Where women go to deliver: understanding the changing landscape of childbirth in Africa and Asia. *Health Policy Plan*. 2017 Oct 1; 32(8):1146–52. <https://doi.org/10.1093/heapol/czx060> PMID: 28541422
90. World Health Organisation. WHO recommendations: intrapartum care for a positive childbirth experience [Internet]. [cited 2020 Apr 28]. Available from: <http://www.who.int/reproductivehealth/publications/intrapartum-care-guidelines/en/>
91. Bohren MA, Berger BO, Munthe-Kaas H, Tunçalp Ö. Perceptions and experiences of labour companionship: a qualitative evidence synthesis. *Cochrane Database Syst Rev* [Internet]. 2019 Mar 18 [cited 2020 May 18]; 2019(3). Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6422112/> <https://doi.org/10.1002/14651858.CD012449.pub2> PMID: 30883666
92. World Health Organisation. The Safe Motherhood Initiative and beyond [Internet]. [cited 2020 May 20]. Available from: <https://www.who.int/bulletin/volumes/85/10/07-045963/en/>