

## Attention-deficit hyperactivity disorder (ADHD) and glial integrity

Oades, Robert D.; Myint, Aye Mu; Dauvermann, Maria R.; Schimmelmann, Benno G.; Schwarz, Markus J.

DOI:

[10.1186/1744-9081-6-32](https://doi.org/10.1186/1744-9081-6-32)

License:

Creative Commons: Attribution (CC BY)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Oades, RD, Myint, AM, Dauvermann, MR, Schimmelmann, BG & Schwarz, MJ 2010, 'Attention-deficit hyperactivity disorder (ADHD) and glial integrity: an exploration of associations of cytokines and kynurenine metabolites with symptoms and attention', *Behavioral and Brain Functions*, vol. 6, 32.  
<https://doi.org/10.1186/1744-9081-6-32>

[Link to publication on Research at Birmingham portal](#)

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

RESEARCH

Open Access

# Attention-deficit hyperactivity disorder (ADHD) and glial integrity: an exploration of associations of cytokines and kynurenine metabolites with symptoms and attention

Robert D Oades\*<sup>1</sup>, Aye-Mu Myint<sup>2</sup>, Maria R Dauvermann<sup>1</sup>, Benno G Schimmelmann<sup>3</sup> and Markus J Schwarz<sup>2</sup>

## Abstract

**Background:** In contrast to studies of depression and psychosis, the first part of this study showed no major differences in serum levels of cytokines and tryptophan metabolites between healthy children and those with attention-deficit/hyperactivity disorder of the combined type (ADHD). Yet, small decreases of potentially toxic kynurenine metabolites and increases of cytokines were evident in subgroups. Therefore we examined predictions of biochemical associations with the major symptom clusters, measures of attention and response variability.

**Methods:** We explored systematically associations of 8 cytokines (indicators of pro/anti-inflammatory function) and 5 tryptophan metabolites with symptom ratings (e.g. anxiety, opposition, inattention) and continuous performance test (CPT) measures (e.g. movement, response time (RT), variability) in 35 ADHD (14 on medication) and 21 control children. Predictions from linear regressions (controlled by the false discovery rate) confirmed or disconfirmed partial correlations accounting for age, body mass and socio-economic status.

**Results:** (1) Total symptom ratings were associated with increases of the interleukins IL-16 and IL-13, where relations of IL-16 (along with decreased S100B) with hyperactivity, and IL-13 with inattention were notable. Opposition ratings were predicted by increased IL-2 in ADHD and IL-6 in control children. (2) In the CPT, IL-16 related to motor measures and errors of commission, while IL-13 was associated with errors of omission. Increased RT variability related to lower TNF- $\alpha$ , but to higher IFN- $\gamma$  levels. (3) Tryptophan metabolites were not significantly related to symptoms. But increased tryptophan predicted errors of omission, its breakdown predicted errors of commission and kynurenine levels related to faster RTs.

**Conclusions:** Many associations were found across diagnostic groups even though they were more marked in one group. This confirms the quantitative trait nature of these features. Conceptually the relationships of the pro- and anti-inflammatory cytokines distinguished between behaviours associated more with cognitive or more with motor control respectively. Further study should extend the number of immunological and metabolic markers to confirm or refute the trends reported here and examine their stability from childhood to adolescence in a longitudinal design.

## Background

Essential components of the diagnosis of attention-deficit/hyperactivity disorder of the combined type (ADHD) are clinical impairments of attention, impulsivity and hyperactivity.

Further, the variability of the expression of these symptoms and of neuropsychological measures in the laboratory is viewed as central to the syndrome and has been proposed as an endophenotype of the disorder [1-4]. Neuronal firing must be maintained to sustain a behavioural response. A recent hypothesis [5] proposed that an inefficient supply of energy from glial cells (the lactate shuttle: [6]) might underlie the intra-individual variability

\* Correspondence: robert.oades@uni-due.de

<sup>1</sup> Clinic for Child and Adolescent Psychiatry and Psychotherapy, University of Duisburg-Essen, 45147 Essen, Germany

Full list of author information is available at the end of the article

in behaviour and the underlying neuronal activity of children with ADHD.

The cytokine and growth factor S100B is a potential marker of glial function. Serum levels arise largely but not exclusively from astrocytes [7]. Increases have been associated with brain damage, major depression, psychosis and dementia [8-10]. However, our pilot study of children with ADHD found that levels of S100B were not unusual, although a tendency for levels to decrease in those with internalizing symptoms was noted [11].

Tryptophan metabolism along the kynurenine pathway takes place primarily in glia [12]. Our study [11] also recorded levels of these metabolites and some cytokines that influence the metabolism [13] as further potential indicators of glial integrity. Relevant to our hypothesis on glial function is that metabolites of kynurenine can be neuroprotective (e.g. kynurenate) or potentially toxic (3-hydroxy-kynurenine, 3-HK) and that the balance between these metabolic routes is modulated by the relative activity of the pro- and anti-inflammatory cytokines [13].

Our initial analysis [11] reported a modest imbalance of cytokines in children with ADHD (figure 1). This improved in those treated with methylphenidate, and thus arguably reflected an increased allostatic load [14]. In the kynurenine pathway, we noted lower levels of the potentially toxic 3HK in children with ADHD than controls that could reflect a maturational delay [15] of the neuronal pruning processes seen in typically developing children [16,17]. [However, an alternative metabolic route for kynurenine over nicotinic adenine dinucleotide (NAD(+)) rather than the toxic quinolinic acid cannot be excluded [18].]

As we had evidence for differences of the cytokine levels and their associations specific to sub-groups (see internalizing symptoms and S100B above, and interleukin IL-10 and IL-16 in those with experience of allergy), we proposed that there would be associations with the major symptom clusters and underlying processes. In particular, we focused on ratings of inattention, hyperactivity, opposition and anxiety, and laboratory measures of attention where the coefficient of variation as a measure of variability could be calculated. This aim has aetiological significance firstly as the inattentive symptoms are those most likely to persist into adulthood [19], and secondly the measure of variability relates directly to the glial hypothesis for the generation of features of ADHD [5]. On the basis of allostasis we predicted that poorer measures of attention would relate to increased levels of some cytokines and reflect an imbalance between pro-and anti-inflammatory activity.

The absence of previous work in this field makes precise predictions impractical. We know of only one study of cytokines in ADHD [20] and one relating to kynurenine [21]. The former found no extreme values for type-1

or type-2 interleukins among patients with ADHD, unlike those with schizophrenia or obsessive-compulsive disorder. The latter described a potential increase of kynurenine metabolism (over that for serotonin, 5-HT) in ADHD following stimulant medication. Indeed, there is substantial indirect evidence for considering a role for 5-HT in ADHD, and by implication the availability of its precursor tryptophan [22,23]. Thus, in the domain of impulsivity, there is evidence for the genetic control in ADHD of the expression of 5-HT receptors, its synthesis and availability [24]. In the domain of sustained attention, task performance related negatively to 5-HT metabolism, and increased 5-HT metabolism (vs. dopamine metabolism) was inversely related with the signal detection measure, d-prime [25]. In support of a continued consideration of 5-HT metabolism here, we reported moderately increased tryptophan availability in untreated ADHD children [11].

Thus, this study represents an exploration of the hypothesis that there are potentially functional associations of symptom ratings (e.g. inattention) and continuous performance task measures (CPT: sustained attention, impulsivity and variability) with 8 cytokines, tryptophan and 4 tryptophan metabolites. The cytokines include the interleukins IL-2, IL-6, interferon-gamma (IFN- $\gamma$ ), tumour necrosis factor-alpha (TNF- $\alpha$ ), IL-16, IL-10, IL-13 and S100B. The metabolites include 5-hydroxy-indoleacetic acid (5-HIAA), kynurenine, kynurenate, and 3HK.


## Methods

### Subjects

Twenty one medication-naïve, consecutively referred children were diagnosed with ADHD combined type (DSM-IV) and recruited to the ADHD-group. Fourteen older children had been given the same diagnosis 4 years earlier and were receiving medication (ADHDmed-group). Their 7 healthy siblings [11] are not further considered here. Controls for the ADHD-group were 21 normally developing children recruited by advertisement. For all groups exclusion criteria included a history of encephalitis, autism, epilepsy, Tourette syndrome, bipolar disorder, IQ  $\leq$  80, brain damage and any genetic or medical condition associated with externalizing behaviours that might mimic ADHD. The protocol, respecting the Declaration of Helsinki, was approved by the ethics committee of the medical faculty of the University of Duisburg-Essen. Verbal and written information on the study was given to the children and the parents or guardians who gave written consent to the procedures.

Diagnosis was based on a semi-structured interview of the parents (Parental Account of Children's Symptoms, PACS: [26,27]) on mood disorders, ADHD/hyperkinetic syndrome, disruptive behaviour and additional problems

### Z-values ( $\times 10^3$ ) for serum levels of 9 cytokines in 4 subject groups



**Figure 1** Shows serum levels (z-values) in 4 groups of subjects for 8 cytokines (the interleukins -1 $\beta$ , -2, -6, -16, TNF- $\alpha$ , IFN- $\gamma$ , -10, -13). Both control groups tended to have lower IFN- $\gamma$  levels than the ADHD groups, and the ADHDmed group more IL-1 $\beta$  than the group with ADHD alone. \*  $p = 0.03$ , #  $p = 0.1$ , ##  $p = 0.07$ .

of the children in structured and unstructured situations. Following training with written definitions of the behaviours replies were rated for the previous week and year (in a period without medication) on a 4-point scale. A standardized diagnostic algorithm based on the DSM-IV criteria for the symptoms, age of onset, situational pervasiveness and clinical impairment was applied to this information. This procedure has achieved good inter-rater reliability: ( $r = .76$  to  $.96$ ; [27]). Symptoms were rated with the Conners parent and teacher ratings scales (CPRS-R:L, CTRS-R:L: [28]). Internalizing and externalizing symptoms were recognized on the anxiety and oppositional subscales with  $T > 65$  on either the CTRS or CPRS (9/14 [ADHD], 6/2 [controls], and 6/9 [ADHDmed]).

Information on gender, age, body-mass index (BMI), IQ, socio-economic index (SES) and reported allergic sensitivities is given in table 1. The IQ was assessed with the CFT-20-R [29] or for children less than 9 years with the Kaufman assessment battery [30]. The current and at-

birth reported SES of the father and mother for profession/qualification was scored on a scale of 1-7 [31]. Allergies in the past ( $> 12$  months) and currently experienced were rated for severity (0-3).

#### Continuous performance test (CPT/Qb)

The CPT/Qb test provides data on sustained attention: accuracy, response times and their variability and restless movements [32]. It lasts 4-15 minutes depending on the speed of response. A grey circle (target), and another with a cross (nontarget) are presented in a random sequence for 100 ms every 2 sec on a computer monitor. Half of the 450 stimuli require a button-press within 2 sec. Accuracy is measured by the percentage errors of omission (sustained attention) and commission (impulsivity). Multiple and anticipatory responses ( $< 150$  ms) were excluded from the analysis. Response latency measures include the mean reaction time (RT) the variation (standard deviation, SD) and coefficient of variance (intra-individual variability,  $CV = 100 \times RT-SD/RT$ ). During the task the

**Table 1: Characteristics of the ADHD, control and ADHDmed groups of children (means and standard deviations)**

	<i>ADHD</i>		<i>Control</i>		<i>ADHDmed</i>	
	mean	SD	mean	SD	mean	SD
<b>N</b>	21		21		14	
<b>Gender (m/f)</b>	14m, 7f		20 m, 1 f		12 m, 2 f	
<b>Age (y)</b>	8.9**	1.4	11.0**	1.5	12.6	2.1
<b>IQ</b>	95.8*	10.7	114.1	14.4	106.7	12.4
<b>Height (cm)</b>	132.1	9.3	147.3	11.6	161.2	17.8
<b>Weight (kg)</b>	29.2	8.4	39.7	11.7	54.7	19.1
<b>BMI</b>	16.5***	2.4	17.9***	3.1	20.5	4.6
<b>Allergy (N)</b>	8		11		2	
<b>Allergy (grade)</b>	0.95	1.2	0.76	0.9	0.29	0.7
<b>SES-birth (M)</b>	4.2	1.6	3.6	0.8	3.7	1.4
<b>(F)</b>	4.3	1.6	3.3	2	3.7	1.6
<b>SES-current (M)</b>	4.3	2.3	3.3	1.5	4.4	2
<b>(F)</b>	4.3	2.2	3.5	2.1	4.1	1.9

BMI = Body mass index,

SES = socio-economic scale for the mother (M) and father (F) at child birth and currently (scale 1-7)

Allergy = the number of subjects reporting allergic sensitivity and the rated severity (scale 0-3)

\* p < 0.05 \*\* p < 0.02 \*\*\* p < 0.005

subject's movements are recorded by an infrared camera following a reflective marker on a headband. The camera reads the X/Y coordinates of the marker, sampling 50x/sec and scales (1/27) the measures to mm/sec. The duration of restlessness (time spent active [ $> 1$  cm/sec] as a percentage of task duration) and the number of microevents (movements of  $>1$  mm) are recorded [33].

### Biochemistry

Details are given in the accompanying report [11]. Briefly, fasting venous blood (20 ml) was drawn (08:00-09:00 a.m.), the serum separated and stored at  $-80^{\circ}\text{C}$  for analysis blind to the origin of the sample [13]. S100B was measured by an immuno-luminometric assay (detection limit  $0.02 \mu\text{g/L}$ : ELECSYS S100™, Roche Diagnostics, Switzerland). The cytokines IL-1 $\beta$ , IL-2, IL-6, IL-10, IL-13, IL-16, IFN- $\gamma$  and TNF- $\alpha$  were analysed with ELISA test kits from R&D systems. For IL-1 $\beta$ , IL-6, IL-10 and TNF- $\alpha$  high-sensitive assays were used. IL-1 $\beta$  was not detected in 40/63 samples (detection limit:  $0.063 \text{ pg/ml}$ ) and excluded from the correlational analysis.

For amino acid determination the AccQ Tag HPLC method was used (Waters, Milford, MA, USA) and concerned the 5 tryptophan-competing amino acids and 12 others [11]. The fluorescence wave lengths were  $\lambda_{\text{ex}} = 250 \text{ nm}$  and  $\lambda_{\text{em}} = 395 \text{ nm}$ . Analyses for tryptophan and the metabolites were run on a Waters 2695 chromatograph with a 2487 dual-l UV detector and a 2475 fluorescence detector. Tryptophan ( $\lambda_{\text{ex}}: 300 \text{ nm}$ ;  $\lambda_{\text{em}}: 350 \text{ nm}$ ) and 5-HIAA ( $\lambda_{\text{ex}}: 300 \text{ nm}$ ;  $\lambda_{\text{em}}: 340 \text{ nm}$ ) were measured with fluorescence, kynurenine (365 nm), kynurenate (330 nm) and 3-HK (365 nm) with UV detection. Three indices of metabolic activity were calculated: (1) *Tryptophan availability* [ $100 \times \text{tryptophan}/\text{sum of the competing amino acids}$ ], (2) *Tryptophan breakdown* [kynurenine/tryptophan] reflecting the summed metabolic enzymatic activity, and (3a) the *neuroprotective index* [kynurenate/3HK]. Neuroprotection would be increased with larger values. The third index differs from Myint et al. [13] who used the ratio of  $1000 \times \text{kynurenate}/\text{kynurenine}$  (index 3b) representing turnover within the protective pathway rather than the production of toxic metabolites.

### Data analysis

First, group comparisons (MANOVA) were performed for group characteristics (table 1), symptom ratings and CPT measures, and confirmed with Tukey tests. Symptom subscales for opposition, anxiety, and the DSM scales for inattention, hyperactive-impulsivity and sum scores were analysed on the parent and the teacher scales (table 2). Six CPT responses were selected as representative of motor activity (duration, microevents), attention (errors of omission and commission) and variability (coefficient of variance and RT: table 3). Covariates for gender, age,

BMI, SES and IQ were considered and are reported where relevant. Relevance was determined from predictions of group characteristics by the biochemical measures in a backwards step-wise regression. The first results' section summarises these analyses.

The second results section reports the associations sought between the group features and the functional domains of attention and symptoms with the 3 sets of biochemical measures (cytokines, tryptophan metabolism and metabolic indices). Initially in this exploration partial correlations were calculated accounting for the relevant covariates (see above) for each group. Second these results were confirmed/disconfirmed with linear regression analysis to test the hypotheses that the biochemical measures predict the symptoms and attentional features. This assists in the elimination of false positives in the correlational analysis. Applying the false discovery rate to the results of the multiple regressions identifies uncorrected probabilities reported as  $\leq 0.01$  as satisfying  $\alpha = 5\%$ : values above this threshold are described as trends or tendencies ( $p < 0.05/\alpha < 10\%$ ). Following each analytical section there is a short summary of the confirmed effects. For tryptophan metabolism measures of tryptophan, kynurenine, kynurenate, 3HK and 5-HIAA and separately the metabolic indices (above) were considered. For the cytokine associations analyses of S100B, IL-2, IL-6, IL-16, TNF- $\alpha$ , IFN- $\gamma$ , IL-10 and IL-13 measures were considered. A few missing values were replaced by the mean.

## Results

### (1) Group comparisons

#### *Group features and comparisons of their symptoms and attentive behaviour*

The three groups did not differ on gender distribution, socio-economic scale or incidence of allergy (table 1). ADHD children were younger than the controls and both were younger than the ADHDmed group ( $p < 0.02$ ): they also had a lower BMI than the ADHDmed group ( $p < 0.005$ ) and their IQ was lower than in either the ADHDmed or control groups ( $p < 0.05$ ).

Covarying for age and IQ, the CPRS symptom ratings were clearly higher for both ADHD groups than the controls on each scale except anxiety. The anxiety ratings were only a little higher in the ADHD groups than in controls. Similar but less marked differences were shown with the CTRS. There were no differences between the ADHD groups (table 2). Using the same covariates attenuated the group differences on CPT measures, but only that for the errors of commission lost significance. Compared to the ADHD groups controls showed less motor activity, a shorter RT with less variability and more errors of omission (table 3). While post-hoc comparisons of the two ADHD groups showed that medication did not

**Table 2: Conners Parent and Teacher Rating Scale Assessments (CPRS/CTRS) of ADHD Symptoms (means and standard errors)**

	ADHD	(N = 20) sem	CON	(N = 21) sem	ADHDmed	(N = 14) sem
<b>CPRS</b>						
Opposition	71.4	2.1	51.7**	2.1	71.5	2.6
Anxiety	56.3	2.3	51.1 <sup>#</sup>	2.3	58.2	2.9
DSM-IA	72.1	2.2	47.8**	2.2	69.3	2.8
DSM-HI	78.4	1.9	48.1**	1.8	72.8	2.3
DSM-sum	76.4	1.8	48.0**	1.7	72.6	2.2
<b>CTRS</b>						
opposition	63.3	2.5	50.8	2.4	56.0	3.1
Anxiety	60.3	2.8	54.8	2.6	62.3	3.4
DSM-IA	65.8	2.5	48.6*	2.3	59.9	3.0
DSM-HI	67.7	2.5	48.4*	2.3	51.5	3.0
DSM-sum	68.1	2.3	48.5*	2.2	57.0	2.9

For all subscales (except anxiety) the ADHD groups scored higher than children without a diagnosis (independent of 4 covariates): IA, inattention; HI, hyperactivity-impulsivity: <sup>#</sup>  $F(2,49) = 2.7$   $p = 0.077$ ,  $\eta^2 = 0.099$ ; \*  $F(2,49) = 7.11$ ,  $p < 0.002$   $\eta^2 = 0.23-0.32$ ; \*\*  $F(2,49) = 34.74$ ,  $p < 0.000$ ,  $\eta^2 = 0.49-0.69$ ;

significantly reduce RT or errors of omission, the ADHDmed group did show fewer microevents, a shorter duration of activity, less RT variability and fewer errors of commission.

**Group comparisons: tryptophan metabolism and cytokines**

Tryptophan levels and availability were modestly higher and breakdown indices lower in those with a diagnosis of ADHD after controlling for age and gender (figure 2, [11]). But this was not reflected in altered levels of 5-HIAA or kynurenine. Taking account of age and BMI, the ratio of kynurenate to 3HK, was lower in children with ADHD than the comparison groups which showed slightly increased levels of 3HK.

There were no group differences in S100B levels, but reductions in ADHD children reflected those with internalizing rather than conduct problems [11]. Except for IL-1 $\beta$  and TNF- $\alpha$  the normalised cytokine values (figure 1) tended to be higher in the ADHD than in the control group. All values in the ADHDmed group returned to more normal levels.

**(2) Relations of biochemical measures to group features, symptoms and attention**

**Group features (relations to tryptophan metabolism and the cytokines)**


Across all children tryptophan levels increased with BMI and age ( $R^2$  0.14,  $F(1,54) = 8.69$ :  $\beta +0.37$ ,  $p = 0.005$ ): this was evident in both the ADHD and ADHDmed groups ( $R^2$  0.22/0.60,  $F(1,19) = 5.33/8.16$ :  $\beta + 0.47/+0.96$ ,  $p = 0.03/0.002$ , respectively). 3HK levels also tended to increase in controls ( $R^2$  0.20,  $F(1,19) = 4.69$ ,  $\beta +0.45$ ,  $p = 0.04$ ). IQ was unrelated, but low SES was predicted by lower tryptophan and increasing kynurenate levels ( $R^2$  0.15,  $F(2,53) = 4.70$ :  $\beta + 0.33/-0.34$ ,  $p = 0.018/0.015$ , respectively). This was supported by modest associations with increased tryptophan breakdown and lower neuroprotection ratio-B ( $\beta -0.3/-0.28$ ,  $p = 0.02/0.03$ ). Lastly, across all children (and within each group), those with an experience of allergy tended to be predicted by low levels of 5-HIAA ( $R^2$  0.07,  $F(1,54) = 3.93$ :  $\beta -0.26$ ,  $p = 0.05$ ).

**Table 3: Continuous performance task measures (CPT/Qb) of motor activity, response speed/variability and accuracy/errors**

CPT/Qb measure	Group	N	Mean	SD	df 2,50
<b>Duration active</b> (% > 1 cm/sec)	ADHD	21	43.4	23.7	
	CON	21	15.6	11	F 5.6, p = 0.006, $\eta^2 = 0.18$
	ADHDmed	13	23.5 <sup>2</sup>	21.9	
<b>Microevents</b> (N > 1 mm)	ADHD	21	10167	5624	
	CON	21	4390	1962	F 5.0, p = 0.01, $\eta^2 = 0.17$
	ADHDmed	13	5531 <sup>1</sup>	3626	
<b>RT (ms)</b>	ADHD	21	581	85	
	CON	21	444	92	F 4.1, p = 0.02, $\eta^2 = 0.14$
	ADHDmed	13	533	141	
<b>CV</b> 100 × RT-SD/RT	ADHD	21	44.1	15.4	
	CON	21	29.2	5.1	F 5.2, p = 0.009, $\eta^2 = 0.17$
	ADHDmed	13	33 <sup>3</sup>	7.7	
<b>Errors</b>					
<b>Omissions</b>	ADHD	21	20.4	17	
	CON	21	2	3	F 5.6, p = .006, $\eta^2 = 0.18$
	ADHDmed	13	19.4	24.2	
<b>Commissions</b>	ADHD	21	17.3	9.1	
	CON	21	9.7	7.4	NS
	ADHDmed	13	6.4 <sup>4</sup>	9.7	

ADHD vs. ADHDmed: 1. p = 0.006, 2. p = 0.014, 3. p = 0.013, 4. p = 0.002





**Figure 2** shows a comparison for the 4 groups of subjects of the levels of tryptophan, its metabolites and the indices for tryptophan availability, breakdown and neuroprotection (A and B). Tryptophan levels and availability were higher in both ADHD groups ( $p < 0.02$ ). Descriptively higher levels of 3HK in the controls ( $p < 0.08$ ) related closely to Kynurenine levels ( $r = +0.85$ ,  $p < 0.000$ ), but a correlation was absent in the ADHD group ( $p = 0.35$ ).

Group features were only marginally predicted by the levels of a few proinflammatory cytokines. Across all subjects increasing TNF- $\alpha$  and decreasing IFN- $\gamma$  levels tended to predict increasing age ( $R^2 0.084$ ,  $F(2,53) = 0.09$ :  $\beta +0.28/-0.31$ ,  $p = 0.077/0.049$ , respectively). BMI also tended to be predicted by increasing TNF- $\alpha$  and IL-6 levels ( $\beta +0.32/+0.45$ ,  $p = 0.03/0.001$ , respectively).

Across subjects allergic experience tended to relate to decreasing proinflammatory TNF- $\alpha$  but increasing anti-inflammatory IL-10 levels ( $R^2 0.16$ ,  $F(2,53) = 5.0$ ,  $p = 0.01$ :  $\beta -0.27/ +0.31$ ,  $p = 0.039/0.017$ , respectively). In the ADHD group, in addition to the TNF- $\alpha$  decrease, ( $\beta -0.63$ ,  $p = 0.03$ ) increasing anti-inflammatory IL-13 levels predicted response ( $\beta +0.56$ ,  $p = 0.01$ ). In contrast, for the controls alone the increase in IL-10 was masked by decreases of IL-13 in the prediction of allergy severity ( $R^2 0.42$ ,  $F(2,18) = 6.6$ ,  $p = 0.007$ :  $\beta +0.33/-0.13$ ,  $p = 0.08/0.01$ , respectively).

#### Summary

Tryptophan levels increased with age and BMI, but lower SES status facilitated metabolism in the kynurenine pathway. Some proinflammatory cytokine levels decreased

(IFN- $\gamma$ ) or increased with BMI (TNF- $\alpha$ , IL-6). Independent of diagnosis, children with allergies tended to show lower levels of 5-HT metabolism. Most children with experience of allergy showed decreases of the proinflammatory TNF- $\alpha$ . However, in contrast to the controls who showed decreases of the anti-inflammatory IL-13, the ADHD group showed increased IL-13 levels.

#### Symptom ratings: relations of tryptophan metabolism

In general, symptoms were not associated clearly with tryptophan metabolism. Partial correlations accounting for age, BMI and SES suggested that inattentive symptoms were related to tryptophan levels across all children ( $r = +0.32$ ,  $p = 0.026$ ). Alone for controls anxiety related to 5-HIAA levels ( $r = 0.52$ ,  $p = 0.038$ ) and inattentive ratings with 3HK ( $r = +0.49$ ,  $p = 0.037$ ). Only the latter result (3HK) approached the threshold for significance in the regression ( $R^2 0.19$ ,  $F(1,19) = 4.5$ :  $\beta +0.44$ ,  $p = 0.048$ ).

#### Symptom ratings: relations of cytokine measures

Overall, considering total symptom ratings in the ADHD group the best predictors were increasing levels of IL-13 and IL-16 along with decreasing levels of S100B ( $R^2 0.57$ ,  $F(4,16) = 5.21$ ,  $p = 0.007$ :  $\beta +0.59, +0.51, -0.54$ ,  $p = 0.024$ ,

0.019, 0.013, respectively) with a minor contribution from TNF- $\alpha$ .


Partial correlations accounting for age, BMI and SES pointed to associations for pro-inflammatory cytokines with two symptom groupings, namely the oppositional and hyperactive-impulsive ratings. In the ADHD group decreasing IL-2 ( $r = -0.64$ ,  $p = 0.026$ ) and TNF- $\alpha$  ( $r = -0.62$ ,  $p = 0.018$ : figure 3) correlated with increased ratings of opposition. Regression confirmed the IL-2 result ( $R^2 0.41$ ,  $F(2,18) = 6.2$ ,  $p = 0.009$ :  $\beta -0.55$ ,  $p = 0.007$ : figure 4) and implied a modest predictive role for increasing levels of S100B ( $\beta +0.37$   $p = 0.057$ ). Of interest is that in the controls another proinflammatory cytokine (IL-6) correlated with oppositional ratings ( $r +0.48$ ,  $p = 0.05$ ): the predictive role was confirmed in the regression ( $R^2 0.30$ ,  $F(1,19) = 7.99$ ,  $p = 0.01$ :  $\beta +0.54$ ,  $p = 0.01$ ).

The association of hyperactive-impulsive ratings in the ADHD group with IL-16 ( $r +0.66$ ,  $p = 0.015$ ) was confirmed by the regression ( $R^2 0.52$ ,  $F(2,18) = 9.82$ ,  $p =$


$0.001$ :  $\beta +0.80$ ,  $p = 0.000$ ) with modest support from decreasing levels of S100B ( $\beta -0.42$ ,  $p = 0.035$ ). Anxiety ratings were without relationships to the cytokines. But, intriguingly ratings of inattention in the ADHD group tended to relate positively to levels of the antiinflammatory IL-13, yet negatively to IL-13 in the controls ( $R^2 0.33/0.45$ ,  $F(2/3, 18/17) = 4.5/4.6$ ,  $p = 0.026/0.015$ :  $\beta +0.44/-0.67$ ,  $p = 0.03/0.007$ , respectively: figure 5). The ADHD result was modestly supported by increasing IL-6 levels ( $\beta +0.38$ ) and the control result by increasing S100B ( $\beta +0.64$ ).

#### Summary

There were modest associations of tryptophan levels and its metabolism to the potentially toxic 3HK with inattention, and 5-HT metabolism to anxiety. Proinflammatory cytokines were associated with oppositional symptoms: decreases of IL-2 in the ADHD group contrasted with increases of IL-6 in the controls. In general, ADHD symptoms were associated with increases of the antiin-



**Figure 3** For children with ADHD, ratings of oppositional behaviour vs. levels of TNF- $\alpha$  (linear regression).



**Figure 4** For children with ADHD, ratings of oppositional behaviour vs. levels of IL-2 (linear regression).

flammatory IL-13 (inattention), increases of IL-16 (hyperactive-impulsive symptoms) and with decreases of S100B (total symptoms).


#### **CPT performance: relations of tryptophan metabolism**

Across all subjects motor activity (duration, microevents) showed a very modest tendency to increase with lower levels of 3HK ( $R^2 0.07$ ,  $F(1,54) = 4.0/3.4$ ;  $\beta -0.26/-0.24$ ,  $p = 0.05/0.07$ , respectively), but this was not evident in the partial correlations and single group analyses.

In control children faster RTs were associated with increasing kynurenine levels ( $R^2 0.32$ ,  $F(1,19) = 9.13$ ;  $\beta -0.57$ ,  $p = 0.007$ ; figure 6). This confirmed the partial correlation ( $r = -0.52$ ,  $p = 0.026$ ). Unexpectedly, the opposite relationship appeared in the ADHDmed group ( $r = +0.93$ ,  $p = 0.000$ ;  $R^2 0.48$ ,  $F(1,12) = 10.93$ ;  $\beta +.69$ ,  $p = 0.006$ ) with no association evident in the non-medicated cases. Across all subjects, there was a partial correlation for the coefficient of RT variance with tryptophan levels ( $r +0.30$ ,  $p = 0.035$ ), yet a predictive role for tryptophan was not

confirmed in the regression. Interestingly, the decrease of tryptophan (and variability) in controls (vs. the ADHD group, figure 2) tended to be reflected by increases of 5-HIAA ( $r = -0.45$ ,  $R^2 0.17$ ,  $F(1,19) = 3.95$ ;  $\beta -0.42$ ,  $p = 0.06$ ). But, no associations were evident in the ADHD groups. The interpretation of the RT associations in terms of 5-HT metabolism is supported by the partial correlations of the turnover indices. While tryptophan availability was positively related to RT ( $r +0.36$ ,  $p = 0.012$ ), the breakdown to kynurenine was negatively associated ( $r -0.31$ ,  $p = 0.03$ ).

The error analysis suggested that omissions were positively associated with tryptophan levels across subjects ( $r = +0.38$ ,  $p = 0.006$ ;  $R^2 0.14$ ,  $F(1,54) = 9.03$ ;  $\beta +0.38$ ,  $p = 0.004$ ). This positive association between tryptophan levels and omission errors was evident as a trend in the ADHDmed group where the relationship to commission errors was negative ( $R^2 0.35/0.54$ ,  $F(1,12)/(2,11) = 6.37/6.42$ ;  $\beta +0.59/-0.92$ ,  $p = 0.027/0.004$ , respectively). This



negative trend with commission errors is upheld by their association with the tryptophan breakdown index ( $R^2$  0.49,  $F(1,12) = 11.8$ ;  $\beta +0.70$ ,  $p = 0.005$ ).


**CPT performance: relations of cytokine measures**

Considering motor activity across all subjects, the partial correlations pointed to a positive association of the duration measure with IL-16 levels ( $r+0.30$ ,  $p = 0.038$ ). This was confirmed by the regression ( $R^2$  0.08,  $F(1,54) = 4.56$ ;  $\beta +0.28$ ,  $p = 0.037$ ). The motor associations appeared to be driven by IL-16 in the ADHD group for both duration and microevent measures ( $R^2$  0.55/0.47,  $F(3,17) = 6.9/5.0$ ,  $p = 0.003/0.01$ ;  $\beta +0.55/+0.47$ ,  $p = 0.007/0.028$ , respectively). [The other degrees of freedom here refer to minor contributions from decreases of the pro-inflammatory IL-6 and TNF- $\alpha$ .] In contrast, the effect of medication introduced a potential modest antiinflammatory influ-

ence (e.g. IL-10 with microevents,  $R^2$  0.35,  $F(1,12) = 6.3$ ;  $\beta +0.59$ ,  $p = 0.027$ ).

Several analyses indicated minor negative associations of RT with cytokines (e.g. IL-2, S100B) but only the effect of medication showed that decreasing RTs related to increasing levels of IL-2 (ADHDmed:  $\beta -0.73$ ,  $p = 0.003$ ; figure 7). This contrasts with RT variability, where across subjects the coefficient of variance was influenced in opposite ways by TNF- $\alpha$  and IFN- $\gamma$  ( $R^2$  0.18,  $F(2,53) = 6.1$ ,  $p = 0.004$ ;  $\beta -0.46/+0.04$ ,  $p = 0.002/0.007$ , respectively; figure 8). While these two relationships were not driven by processes related to diagnosis, the association with TNF- $\alpha$  was most marked in the medicated ADHD group ( $\beta -0.70$ ,  $p = 0.007$ ).

In the error analyses errors of commission were predicted by levels of IL-16 ( $r+0.42$ ,  $p = 0.003$ ;  $R^2$  0.17,  $F(1,54) = 10.75$ ;  $\beta +0.41$ ,  $p = 0.002$ ; figure 9). But, as with



variability (above) this was most apparent in the ADHDmed group ( $\beta +0.75$ ,  $p = 0.000$ ). In this group alone, again as with variability, decreases of proinflammatory  $\text{TNF-}\alpha$  ( $\beta -0.36$ ,  $p = 0.016$ ) and  $\text{IL-6}$  ( $\beta -0.39$ ,  $p = 0.032$ ) also tended to predict errors of commission. For omission errors only the negative associations of  $\text{IL-13}$  (ADHD:  $\beta -0.47$ ,  $p = 0.03$ ) and  $\text{IL-16}$  (ADHDmed:  $\beta -0.63$ ,  $p = 0.004$ ) were of note.


#### Summary

Measures of motor activity were not related to tryptophan metabolism but were positively predicted by  $\text{IL-16}$  levels. In contrast, shorter RTs were associated with increasing kynurenine levels in controls, but by decreasing levels in the ADHDmed group. Here, the cytokines were without any major influence. Interestingly, decreased RT variability in controls was reflected by modest increases of catabolism to 5-HIAA. With regard to the cytokines, across subjects, levels of variability reflected the opposing influences of  $\text{TNF-}\alpha$  and  $\text{IFN-}\gamma$ . In

the error analysis, while the build up of tryptophan predicted omission errors, its breakdown predicted errors of commission. But only the errors of commission were predicted by  $\text{IL-16}$  levels. Medication may moderate interactions of pro- and anti-inflammatory cytokines in motor activity ( $\text{IL-2}/\text{RT}$ ,  $\text{IL-10}/\text{microevents}$ ) and the influences of proinflammatory  $\text{TNF-}\alpha$ ,  $\text{IL-6}$  (variable RT) and  $\text{IL-16}$  (impulsivity).

#### Discussion

Over 20 years ago Kim and Choi [34] demonstrated that the breakdown of tryptophan and kynurenine can produce toxic metabolites (3HK and quinolinate) in neocortical tissue, but an alternative pathway produces kynurenate that can be protective. Kynurenine metabolism in the brain occurs largely in the glia that envelop glutamatergic synapses [12]. Among the numerous responses of the glia to excitotoxicity are alterations in the proinflammatory cytokines, such as  $\text{TNF-}\alpha$  and  $\text{IFN-}\gamma$ .



$\gamma$  [35], where the balance with antiinflammatory cytokines influences tryptophan metabolism [13]. Recently there have been numerous reports of associations of kynurenine metabolites and cytokines with brain damage and psychiatric disorders [8,36-40]. While we only found modest changes of selected markers of these cytokines and metabolites in our pilot study of ADHD [11], we hypothesized here that key symptoms and poor attentional function would reflect alterations of the activity of these markers. Features of the kynurenine metabolic pathway and neuroinflammatory activity of cytokines have not been explored before in the context of ADHD.


We report a preliminary exploration of whether products of tryptophan metabolism and the cytokines that modulate this metabolism predict some of these symptoms, and their laboratory measures in children with a strict diagnosis of ADHD (combined type) contrasted with those developing normally. The study is predicated on increasing evidence that alterations in these two bio-

chemical domains are reflected in both the features of ADHD [21,41] and animal models of these functions [42,43].

Those with the diagnosis of ADHD had clearly more symptoms of inattention, hyperactivity-impulsivity and opposition than those without the diagnosis. On a test of sustained attention the children with ADHD showed more motor activity, errors of omission, a slower RT and more RT variability. Most cytokine levels were modestly increased in children with ADHD (not S100B and IL-1 $\beta$ ) and all tended to normalize on psychostimulant medication. While tryptophan availability was slightly increased in the ADHD group, 3HK levels tended to be lower than in the controls [11].

#### The cytokines and associations with symptoms

The DSM ratings of total symptom severity in the ADHD group were predicted by increasing levels of IL-16, and the anti-inflammatory IL-13, but by decreasing levels of




**Figure 8** For all subjects, the coefficient of variance (CV) vs. levels of TNF- $\alpha$  (linear regression).

S100B. The anti-inflammatory relationship was modestly evident for inattentive symptoms in the ADHD group, and arguably effective in countering dysfunction in that higher levels of IL-13 in the controls were related to fewer symptoms of inattention. The role of IL-16 can be pro- or anti-inflammatory in different situations [44,45]. Here, in the ADHD group, IL-16 levels additionally predicted hyperactive-impulsive symptoms that were also modestly associated with decreased S100B. The IL-16 relationship with hyperactive symptoms might not be expected in so far as 2 genetic polymorphisms have been associated with the inattentive type of ADHD [46]. However, the difference may reflect other features of the two populations influencing the neuroinflammatory role of IL-16, such as allergy (see below). The association of symptom severity with decreasing S100B levels is consistent with our prediction of inefficient glial function in ADHD [11] rather than an overproduction resulting from apoptosis and

brain damage reported for other major psychiatric disorders [10,47,48].

Only proinflammatory cytokine activity related to oppositional symptom ratings. Decreasing levels of IL-2, and to a minor extent TNF- $\alpha$ , were associated with increasing signs of opposition. We suggest the negative direction of association mirrors consistently the frequent reports of increased IL-2 activity in disorders such as schizophrenia and Parkinsonism with a very different clinical profile [40,49,50]. On the other hand we note a relationship between increasing IL-6 and oppositional behaviour in the control group. We tentatively suggest that, as IL-6 levels increased with BMI and the controls were marginally older than the patients, such an increase reflects the normal parallel development of boisterous behaviour often associated with adolescence onset. This interpretation is also consistent with the well-studied associations of IL-6 levels with physical and psychological stress [51].



**Figure 9** For all subjects, commission errors on the continuous performance test vs. levels of IL-16 (linear regression).

Sensitivity to allergy reflected opposing trends for pro- and antiinflammatory cytokines. Children sensitive to allergy showed 15% more IL-16, 32% more IL-13 and similar levels of IL-10 [11]. But for the ADHD children who experienced allergic responses, these responses tended to be predicted by decreasing levels of the proinflammatory  $\text{TNF-}\alpha$  and increases of the antiinflammatory IL-13. For controls the trend went in the opposite direction for IL-13, but this was balanced by modest increases of another antiinflammatory cytokine, IL-10. The difference between groups in the proinflammatory response is quantitative, but that for the antiinflammatory response is qualitative. This physiological response may underlie a subgroup of ADHD children with high allergic responsiveness and for whom ADHD symptoms are alleviated by removal of the allergens from the diet [52-54]. Further, IL-10 secretion is facilitated by physical or psychological stress [55] and could be responsible for the decreases of tryptophan metabolism [56] discussed below.

#### The cytokines and associations with CPT performance

During the CPT both the duration of major movements and the quantity of small fidgety activity (microevents) were directly measured. Increasing IL-16 levels not only predicted hyperactive symptoms (above) but clearly predicted quantitative measures of motor activity, especially (but not only) in ADHD children. The associations extended to error omission and commission in the medicated children that may reflect motor as well as cognitive control. The associations of IL-16 with motor activity and allergy in this study clearly add to the concept of this being a pleiotropic cytokine, for which not all the roles have yet been elaborated.

Accuracy in terms of the errors made is more usually thought of as a cognitive than a motoric feature. Our data suggest that decreasing levels of the anti-inflammatory IL-13 may not only relate to inattentive symptoms (above), but specifically to increases of errors of omission. But, both this association and the tendency for errors of



commission in medicated children to be reduced as pro-inflammatory TNF- $\alpha$  and IL-6 levels increased were modest trends. This should be examined closely in future studies. Nonetheless the increase of IL-2 with decreased RT in the ADHDmed group supports the proinflammatory effect on attention-related function.

Associations of the coefficient of variance of the RT is of especial interest, first as variability is the core feature of the hypothesis of glial impairment in ADHD [5], and secondly as it has been proposed as an endophenotype for ADHD (introduction). Here, across all subjects, increasing RT variability was related to decreasing levels of TNF- $\alpha$  and with increasing levels of IFN- $\gamma$ .

Numerous studies confirm the involvement of pro-inflammatory cytokines in cognitive performance both in animal models and in the treatment of cancer patients. For example IL-2 treatment can interfere with long term potentiation in the hippocampal slice [57] and the formation of working memory in patients [58]. Treating rodents with IL-6 can impair measures of memory [59,60], yet in knockout mice recall can be facilitated [61,62]: the relationship of IL-6 to cognitive function may be linear or U-shaped [63]. However, IL-6 expression depends on the nature of the allele carried (CC/GC genotype) and this variable has too rarely been considered [64,65]: the C allele may dominate in ADHD while the protective G allele may predominate in typically developing children [66]. The characteristics of TNF- $\alpha$  and IFN- $\gamma$  have much in common with other proinflammatory interleukins. Too much [67] or too little TNF- $\alpha$  [68] can impair cognitive function, but moderate levels are essential [69]. Variation of these levels can influence the turnover of each of the monoamines [70]. Thus it is no surprise that TNF- $\alpha$  levels relate to accuracy and the variability of response in ADHD. However, the relationship of the effects with those of IFN- $\gamma$  require closer study, as the physiological response to each varies with the measure and over time [70].

#### **Tryptophan metabolism and associations with symptoms and CPT performance**

Even after controlling for age, BMI and SES there were tendencies across subjects for a build-up of tryptophan (and its increased availability) to be associated with symptoms of inattention, RT and its variability as well as increases of errors of omission. In contrast decreased tryptophan levels (more breakdown) related to errors of commission. However, tryptophan metabolism was not related to motor activity. This was unexpected as the motor regions of the brain receive a dense 5-HT innervation [71].

For controls, slower RTs but for the ADHDmed group faster RTs were clearly associated with lower kynurenine levels. There was an indication that low levels of kynure-

nine for controls was reflected by the trend for reduced variability and an associated increase of metabolism in the 5-HT pathway. For medicated children the implication is of further glial metabolism in the kynurenine pathway that is facilitated by psychostimulant treatment [72]. Other trend relationships (e.g. 3HK with inattention, 5-HIAA with anxiety) were equivocal. The lack of anticipated changes of 5-HT metabolism related to antisocial/oppositional behaviour [73,74] may simply reflect the extent of 5-HT metabolism from peripheral sources.

The absence of clear associations for tryptophan metabolism with symptoms or attention related performance in ADHD is itself of interest. An overexpression of indoleamine 2, 3 dioxygenase facilitates degradation to kynurenine (and its potentially toxic metabolites) and reduced 5-HT metabolism [75]. Such activity is promoted by proinflammatory cytokines [76]. This scenario fits evidence for the development of psychotic symptoms [38] and major depression [13]. But, our results tend to support the opposite effect in ADHD with increases of tryptophan availability relating to inattention and variable responses. Indeed, here there was even modest support for previous evidence for an involvement of 5-HT metabolism [77].

We had anticipated that the cytokine balance would favour kynurenine breakdown to either kynurenate or 3HK. On the basis of their glutamatergic antagonist and agonist activity these metabolites (and the 3HK metabolite quinolinate) have been ascribed a neuroprotective and toxic role, respectively [78]. But in fact our typically developing children rather than the ADHD group showed descriptively higher levels of kynurenate and 3HK. Yet, these measures did not predict symptoms or CPT performance. We speculate that the ADHD children in fact exhibit a relative lack of some of the other characteristics of these metabolites. For example, kynurenate also has  $\alpha$ 7 cholinergic antagonist properties [79] and higher levels antagonize rather than facilitate AMPA receptor transmission [80]. 3HK has antioxidant capability ([81]) as well as the toxic potential to assist in developmental pruning [16]. Indeed, it is unclear as yet whether energy metabolism in developing children would be impaired by the NMDA agonism of quinolinate [82] or facilitated through further metabolism to NAD(+) [83].

#### **Limitations**

Our conclusions are severely curtailed by not having measured other metabolites (picolinic and, quinolinic acid, NAD(+)), enzyme activity or the expression of genetic variants for these. Also in a pilot study the robustness of the data is naturally compromised by the small number of children that could be recruited and persuaded to offer a blood sample. However, this exploration with 56 children indicates a number of lines of inquiry

that are not worth pursuing and others that might merit further study. Indeed, although the number of variables examined raises questions about the prevalence of false positive results, the statistical procedure included confirmatory tests and false discovery assessment to limit this. Cytokine measures, renowned for their variability, were relatively consistent with remarkably few outliers. Future work should seek replication with a longitudinal design to demonstrate consistency of the data in a larger sample. Such a design should account for developmental age, match for gender, and control for medication yet recruit cases with a range of comorbid attributes to clarify their contribution to the diagnostic condition.

## Conclusions

We hypothesised that even if there are no major dichotomies for this selection of metabolites and cytokines between the groups studied, there would be some associations between symptom dimensions or attentional features and the activity of individual markers. In this exploration we have been able to point out some such associations that bear further study. Some regressions predicted 10 to 40% of the variance of the variable in question.

While in most cases the elements of the kynurenine pathway analysed did not show strong associations with behaviour, we are aware that those metabolites determining toxicity or energy supply were missing (e.g. quinolinic acid, NAD(+)). Only such a determination can link unequivocally putative maturation-related pruning (3HK levels) or glial energy supply [5] to metabolism. However, independent of kynurenine metabolism, among the cytokines there appears to be a conceptual division between the activity of anti-inflammatory members with motor-control related features (symptoms, microevents) and pro-inflammatory members with cognitive-control features (opposition, variability). In several instances the associations extended across all subjects implying their quantitative trait nature. We recommend replications to include the pro-inflammatory IL-1 $\beta$  and anti-inflammatory IL-4 and comparisons to examine stability of these associations before and after adolescence.

## Abbreviations

ADHD: Attention-deficit/hyperactivity disorder; BMI: Body mass index; DSM-IV: Diagnostic and Statistical Manual of the American Psychiatric Association, 4th edition; 5-HIAA: 5-hydroxy-indole-acetic acid; 5-HT: serotonin; HPLC: High Performance Liquid Chromatography; 5-HT: Serotonin; IL: Interleukin; IFN- $\gamma$ , Interferon gamma; 3HK: 3-hydroxy-kynurenine; PACS: Parental Account of Children's Symptoms; RT: Response time; SES: Socio-economic scale; SD: Standard Deviation; TNF- $\alpha$ : Tumour necrosis factor alpha

## Competing interests

RDO, MD and BGS declare that they have no competing interests beyond the source of finance for the study (see below). AMM & MJS have patented the use of tryptophan pathway metabolites as neurodegenerative markers for depression and related psychiatric diseases

## Authors' contributions

RDO was involved in conceiving and organizing the study as well as analysing the data and writing the report. The organization was refined by BGS who with MD was involved in recruitment, diagnosis and logistics. MJS and AMM advised on the biochemical items for study, organized and ran the biochemical analyses and contributed to the strategy for the study and report. The ideas and hypotheses derived from discussion of the work of AMM and RDO. All authors contributed to and approved the final manuscript.

## Authors' information

The study design was intentionally exploratory and the interpretation accordingly conditional. The presentation is aimed at informing both the disciplines of ADHD (psychiatric) and psychoimmunology (basic research) for whom certain elements may be more or less familiar and include negative and positive findings that may both guide future investigations.

Some of these data were communicated at the 10<sup>th</sup> Psychoimmunology Expert meeting at Ulm/Günzburg, 12-14<sup>th</sup> Nov. 2009 [84].

## Acknowledgements

We are very grateful to UCB Pharma GmbH for financial support. We would like to thank Victoria Kirchhoff, Adriana Banozic and Ellen Uslar for their assistance in running this study in Essen, and Johana Zach for her help in the biochemical analyses in Munich.

## Author Details

<sup>1</sup>Clinic for Child and Adolescent Psychiatry and Psychotherapy, University of Duisburg-Essen, 45147 Essen, Germany, <sup>2</sup>Laboratory for Psychoneuroimmunology, Ludwig Maximilian's University Psychiatric Hospital, 8036 Munich 2, Germany and <sup>3</sup>Child and Adolescent Psychiatry, University of Bern, Effingerstr. 12, 3011 Bern, Switzerland

Received: 9 December 2009 Accepted: 9 June 2010

Published: 9 June 2010

## References

1. Castellanos FX, Tannock R: **Neuroscience of attention-deficit/hyperactivity disorder: the search for endophenotypes.** *Nat Rev Neurosci* 2002, **3**:617-628.
2. Kuntsi J, Asherson P: **An interdisciplinary approach to ADHD.** In *Attention Deficit Hyperactivity Disorder Research Developments* Edited by: Larimer MP. Hauppauge, NY: Nova Science Publishers, Inc; 2005:1-30.
3. Klein C, Wendling K, Huettner P, Ruder H, Peper M: **Intra-subject variability in attention-deficit hyperactivity disorder (ADHD).** *Biol Psychiatry* 2006, **60**:1088-1097.
4. Uebel H, Albrecht B, Asherson P, Boerger N, Butler L, Chen W, Christiansen H, Heise A, Kuntsi J, Schäfer U, Andreou P, Manor I, Marco R, Miranda A, Mulligan A, Oades RD, van der Meere JJ, Faraone SV, Rothenberger A, Banaschewski T: **Performance variability, impulsivity errors and the impact of incentives as gender-independent endophenotypes for ADHD.** *J Child Psychol Psychiatry* 2010, **51**:210-218.
5. Russell VA, Oades RD, Tannock R, Auerbach J, Killeen PR, Johansen EB, Sagvolden T: **Response variability in attention-deficit/hyperactivity disorder: a neuronal and glial energetics hypothesis.** *Behav Brain Funct* 2006, **2**:30.
6. Gallagher CN, Carpenter KLH, Grice P, Howe DJ, Mason A, Timofeev I, Menon DK, Kirkpatrick PJ, Pickard JD, Sutherland GR, Hutchinson PJ: **The human brain utilizes lactate via the tricarboxylic acid cycle: a 13C-labelled microdialysis and high-resolution nuclear magnetic resonance study.** *Brain* 2009, **132**:2839-2849.
7. Steiner J, Bernstein H-G, Bielau H, Berndt A, Brisch R, Mawrin C, Keilhof G, Bogerts B: **Evidence for a wide extra-astrocytic distribution of S100B in human brain.** *BMC Neurosci* 2007, **8**:2.
8. Arolt V, Peters M, Erfurth A, Wiesmann M, Missler U, Rudolf S, Kirchner H, Rothermundt M: **S100B and response to treatment in major depression: a pilot study.** *Eur Neuropsychopharmacol* 2003, **13**:235-239.
9. Park ES, Park C, Choi KS, Choi I-H, Shin J-S: **Over-expression of S100B protein in children with cerebral palsy or delayed development.** *Brain Dev* 2004, **26**:190-196.
10. Steiner J, Bernstein H-G, Bielau H, Farkas N, Winter J, Dobrowolny H, Brisch R, Gos T, Mawrin C, Myint AM, Bogerts B: **S100B-immunopositive glia is**

- elevated in paranoid as compared to residual schizophrenia: A morphometric study. *J Psychiatr Res* 2008, **42**:868-876.
11. Oades RD, Dauvermann MR, Schimmelmann BG, Schwarz MJ, Myint AM: **Attention-deficit hyperactivity disorder (ADHD) and glial integrity: S100B, cytokines and kynurenine metabolism - effects of medication.** *Behav Brain Funct* 2010.
  12. Kiss C, Ceresoli-Borroni G, Guidetti P, Zielke CL, Zielke HR, Schwarz R: **Kynurenate production by cultured human astrocytes.** *J Neural Transm* 2003, **110**:1-14.
  13. Myint AM, Kim YK, Verkerk R, Scharpe S, Steinbusch HWM, Leonard BE: **Kynurenine pathway in major depression: Evidence of impaired neuroprotection.** *J Affect Disord* 2007, **98**:143-151.
  14. Chapman BP, Khan A, Harper M, Stockman D, Fiscella K, Walton J, Duberstein P, Talbot N, Lyness JM, Moynihan J: **Gender, race/ethnicity, personality, and interleukin-6 in urban primary care patients.** *Brain Behav Imm* 2009, **23**:636-642.
  15. Shaw P, Eckstrand K, Sharp W, Blumenthal J, Lerch JP, Greenstein D, Clasen L, Evans A, Giedd JN, Rapoport JL: **Attention-deficit/hyperactivity disorder is characterized by a delay in cortical maturation.** *Proc Natl Acad Sci (USA)* 2007, **104**:19649-19654.
  16. Huttenlocher PR: **Synapse elimination and plasticity in developing human cerebral cortex.** *Am J Ment Defic Res* 1984, **88**:488-496.
  17. Bourgeois JP, Rakic P: **Changes of synaptic density in the primary visual cortex of the macaque monkey from fetal to adult stage.** *J Neurosci* 1993, **13**:2801-2820.
  18. di Luccio E, Wilson DK: **Comprehensive X-ray structural studies of the quinolinate phosphoribosyl transferase (BNA6) from *Saccharomyces cerevisiae*.** *Biochemistry* 2008, **47**:4039-4050.
  19. Nigg JT, Stavro G, Ettenhofer M, Hambrick DZ, Miller T, Henderson JM: **Executive Functions and ADHD in Adults: Evidence for Selective Effects on ADHD Symptom Domains.** *J Abnorm Psychol* 2005, **114**:706-717.
  20. Mittelman BB, Castellanos FX, Jacobsen LK, Rapoport JL, Swedo SE, Shearer GM: **Cerebrospinal fluid cytokines in pediatric neuropsychiatric disease.** *J Immunol* 1997, **159**:2994-2999.
  21. Uzbekov MG: **Hyperkinetic syndrome as a manifestation of a disturbance of metabolism and mental development.** In *Attention-Deficit/Hyperactivity Disorder and the Hyperkinetic Syndrome: Current Ideas and Ways Forward* Edited by: Oades RD. Hauppauge, New York: Nova Science Publishers, Inc; 2006:133-154.
  22. Oades RD: **The role of the serotonin system in ADHD: treatment implications.** *Expert Rev Neurotherapeutics* 2007, **7**:1357-1374.
  23. Oades RD: **Dopamine-serotonin interactions in attention-deficit/hyperactivity disorder (ADHD).** *Prog Brain Res* 2008, **172**:543-565.
  24. Oades RD, Lasky-Su J, Christiansen H, Faraone SV, Sonuga-Barke EJS, Banaschewski T, Chen W, Anney R, Buitelaar JK, Ebstein RP, Franke B, Gill M, Miranda A, Rothenberger A, Sergeant JA, Steinhausen H-C, Taylor EA, Thompson M, Asherson P: **The influence of serotonin and other genes on impulsive behavioral aggression and cognitive impulsivity in children with attention-deficit/hyperactivity disorder (ADHD): Findings from a family-based association test (FBAT) analysis.** *Behav Brain Funct* 2008, **4**:48.
  25. Oades RD: **Differential measures of sustained attention in children with attention-deficit/hyperactivity or tic disorders: relationship to monoamine metabolism.** *Psychiatry Res* 2000, **93**:165-178.
  26. Taylor EA, Sandberg S, Thorley G, Giles S: *The epidemiology of childhood hyperactivity* Oxford, UK: Oxford University Press; 1991.
  27. Chen W, Taylor EA: **Parental Account of Children's Symptoms (PACS), ADHD phenotypes and its application to molecular genetic studies.** In *Attention-Deficit/Hyperactivity Disorder and the Hyperkinetic Syndrome: Current Ideas and Ways Forward* Edited by: Oades RD. Hauppauge, NY 11788: Nova Science Publishing Inc; 2006:3-20.
  28. Conners CK: *Manual for Conners' rating scales, Revised edn* Edited by: Tonoawanda N. NY: Multi-Health Systems Inc; 2002.
  29. Weiss RH: *CFT-20-R: Grundintelligenztest* 2nd edition. Göttingen: Hogrefe Verlag GmbH & Co KG; 1996.
  30. Kaufman AS, Kaufman NL: *Kaufman-Assessment Battery for Children* 5th edition. Edited by: Melchers P, Preuß U. Frankfurt am Main, Germany: Swets & Zeitlinger; 2001.
  31. Brauns H, Haun D, Steinmann S: **The construction of an internationally comparable classification by class.** In *Erwerbsstatistische Besonderheiten am Beispiel von Labour Force Surveys der Bundesrepublik Deutschland, Frankreichs, Großbritanniens und Ungarns* Mannheim, Germany; 1997.
  32. Bergfalk H: *QbTest Technical Manual* Gothenburg, Sweden: Qbtech; 2006.
  33. Brocki KC, Tillman CM, Bohlin G: **CPT performance, motor activity, and continuous relations to ADHD symptom domains: A developmental study.** *Eur J Dev Psychol* 2010, **7**:178-197.
  34. Kim JP, Choi DW: **Quinolate neurotoxicity in cortical cell culture.** *Neurosci* 1987, **23**:423-432.
  35. Kumar A, Babu GN: **In Vivo Neuroprotective Effects of Peripheral Kynurenine on Acute Neurotoxicity Induced by Glutamate in Rat Cerebral Cortex.** *Neurochem Res* 2010, **35**:636-644.
  36. Tanaka Y, Marumo T, Omura T, Yoshida S: **Early increases in serum S100B are associated with cerebral hemorrhage in a rat model of focal cerebral ischemia.** *Brain Res* 2008, **1227**:248-254.
  37. Andreatza AC, Cassini C, Rosa AR, Leite MC, de Almeida LMV, Nardin P, Cunha ABN, Cereser KM, Santin A, Gottfried C, Salvador M, Kapzinski F, Goncalves C-A: **Serum S100B and antioxidant enzymes in bipolar patients.** *J Psychiatr Res* 2007, **41**:523-529.
  38. Kim Y-K, Myint AM, Verkerk R, Scharpe S, Steinbusch HWM, Leonard BE: **Cytokine Changes and Tryptophan Metabolites in Medication-Naïve and Medication-Free Schizophrenic Patients.** *Neuropsychobiology* 2009, **59**:123-129.
  39. Barry S, Clarke G, Scully P, Dinan TG: **Kynurenine pathway in psychosis: evidence of increased tryptophan degradation.** *J Psychopharmacology* 2009, **23**:287-294.
  40. Stöber G, Ben-Shachar D, Cardon M, Falkai P, Fonteh AN, Gawlik M, Glenthøj BY, Grünblatt E, Jablensky AV, Kim Y-K, Kornhuber J, Müller N, Oranje B, Saito T, Saoud M, Schmitt A, Schwartz M, Thome J, Uzbekov MG, Durany N, Riederer P: **Schizophrenia: From the brain to peripheral markers. A consensus paper of the WFSBP task force on biological markers.** *World J Biol Psychiatry* 2009, **10**:127-155.
  41. Oades RD: **The Role of Serotonin in Attention-Deficit Hyperactivity Disorder (ADHD).** In *The Handbook of Behavioral Neurobiology of Serotonin* Edited by: Müller C, Jacobs B. Amsterdam: Elsevier; 2010:565-584.
  42. Wu H-Q, Rassoulpour A, Schwarz R: **Kynurenine acid leads, dopamine follows: A new case of volume transmission in the brain?** *J Neural Transm* 2007, **114**:33-41.
  43. O'Sullivan JB, Ryan KM, Curtin NM, Harkin A, Connor TJ: **Noradrenaline reuptake inhibitors limit neuroinflammation in rat cortex following a systemic inflammatory challenge: implications for depression and neurodegeneration.** *Int J Neuropsychopharmacology* 2009, **12**:687-699.
  44. Deng J-M, Shi H-Z: **Interleukin-16 in asthma.** *Chin Med J* 2006, **119**:1017-1025.
  45. Little FF, Lynch E, Fine G, Center DM, Cruikshank WW: **Tumor necrosis factor-alpha-induced synthesis of interleukin-16 in airway epithelial cells: priming for serotonin stimulation.** *Am J Respir Cell Mol Biol* 2003, **28**:354-362.
  46. Lasky-Su J, Neale BM, Franke B, Anney R, Zhou K, Maller JB, Vasquez AA, Chen W, Asherson P, Buitelaar JK, Banaschewski T, Ebstein RP, Gill M, Miranda A, Mulas F, Oades RD, Roeyers H, Rothenberger A, Sergeant JA, Sonuga-Barke EJS, Steinhausen H-C, Taylor EA, Daly M, Laird N, Lange C, Faraone SV: **Genome-wide association scan of quantitative traits for Attention Deficit Hyperactivity Disorder identifies novel associations and confirms candidate gene associations.** *Am J Med Genet B* 2008, **147B**:1345-1354.
  47. Van Eldik LJ, Wainwright MS: **The Janus face of glial-derived S100B: beneficial and detrimental functions in the brain.** *Restor Neurol Neurosci* 2003, **21**:97-108.
  48. Castellani C, Bimbashi P, Rutenstock E, Sacherer P, Stojakovic T, Weinberg A-M: **Neuroprotein S100B - a useful parameter in paediatric patients with mild traumatic brain injury?** *Acta Paediatr* 2009, **98**:1607-1612.
  49. Breece C, Rapoport MH: **Persistently increased serum soluble interleukin-2 receptors in continuously ill patients with schizophrenia.** *Int J Neuropsychopharmacology* 2009, **12**:861-865.
  50. Brodacki B, Staszewski J, Toczyłowska B, Kozłowska E, Drela N, Chalimoniuk C, Stepien A: **Serum interleukin (IL-2, IL-10, IL-6, IL-4), TNF, and INFγ concentrations are elevated in patients with atypical and idiopathic parkinsonism.** *Neurosci Lett* 2008, **441**:158-162.
  51. Shizuya K, Komori T, Fujiwara R, Miyahara S, Ohmori M, Nomura J: **The expressions of mRNAs for interleukin-6 (IL-6) and the IL-6 receptor (IL-6R) in the rat hypothalamus and midbrain during restraint stress.** *Life Sci* 1998, **62**:2315-2320.

52. Rapp DJ: Does diet affect hypersensitivity? *J Learn Disabil* 1978, **11**:56-62.
53. Tryphonas H, Trites RL: Food allergy in children with hyperactivity, learning disabilities and/or minimal brain dysfunction. *Ann Allergy* 1979, **42**:22-27.
54. Egger J, Carter CM, Graham PJ, Gumley D, Soothill JF: Controlled trial of oligoantigenic treatment in the hyperkinetic syndrome. *Lancet* 1985, **14**:540-545.
55. Curtin NM, Boyle NT, Mills KHG, Connor TJ: Psychological stress suppresses innate IFN-g production via glucocorticoid receptor activation: Reversal by the anxiolytic chlordiazepoxide. *Brain Behav Imm* 2009, **23**:535-542.
56. Chiarugi A, Calvani M, Meli E, Traggiai E, Moroni F: Synthesis and release of neurotoxic kynurenine metabolites by human monocyte-derived macrophages. *J Neuroimmunol* 2001, **120**:190-198.
57. Di Filippo M, Sarchielli P, Picconi B, Calabresi P: Neuroinflammation and synaptic plasticity: theoretical basis for a novel, immune-centred, therapeutic approach to neurological disorders. *Trends Pharmacol Sci* 2008, **29**:402-412.
58. Capuron L, Ravaut A, Dantzer R: Timing and specificity of the cognitive changes induced by interleukin-2 and interferon-alpha treatments in cancer patients. *Psychosom Med* 2001, **63**:376-386.
59. Monje ML, Toda H, Palmer TD: Inflammatory blockade restores adult hippocampal neurogenesis. *Science* 2003, **302**:1760-1765.
60. Marsland AL, Gianaros PJ, Abramowitch SM, Manuck SB, Hariri AR: Interleukin-6 Covaries Inversely with Hippocampal Grey Matter Volume in Middle-Aged Adults. *Biol Psychiatry* 2008, **64**:484-490.
61. Braidia D, Sacerdote P, Panerai AE, Bianchi M, Aloisi AM, Iosue S, Sala M: Cognitive function in young and adult IL(interleukin)-6 deficient mice. *Behav Brain Res* 2004, **153**:423-429.
62. Hryniewicz A, Bialuk I, Kaminski KA, Winnicka MM: Impairment of recognition memory in interleukin-6 knock-out mice. *Eur J Pharmacol* 2007, **577**:219-220.
63. Baier PC, May U, Scheller J, Rose-John S, Schiffelholz T: Impaired hippocampus-dependent and -independent learning in IL-6 deficient mice. *Behav Brain Res* 2009, **200**:192-196.
64. Baune BT, Ponath G, Rothermundt M, Riess O, Funke H, Berger K: Association between genetic variants of IL-1 $\beta$ , IL-6 and TNF- $\alpha$  cytokines and cognitive performance in the elderly general population of the MEMO-study. *Psychoneuroendocrinology* 2008, **33**:68-76.
65. Harding D, Brull D, Humphries SE, Whitelaw A, Montgomery H, Marlow N: Variation in the interleukin-6 gene is associated with impaired cognitive development in children born prematurely: a preliminary study. *Pediatr Res* 2005, **58**:117-120.
66. Drtilkova I, Sery O, Theiner P, Uhrova A, Zackova M, Balastikova B, Znojil V: Clinical and molecular genetic markers of ADHD in children. *Neuroendocrinol Lett* 2008, **29**:320-327.
67. Medeiros R, Figueiredo CP, Pandolfo P, Duarte FS, Prediger RDS, Passos GF, Calixto JB: The role of TNF- $\alpha$  signaling pathway on COX-2 upregulation and cognitive decline induced by b-amyloid peptide. *Behav Brain Res* 2010, **209**:165-173.
68. McAfoose J, Koerner H, Baune BT: The effects of TNF deficiency on age-related cognitive performance. *Psychoneuroendocrinology* 2009, **34**:615-619.
69. Baune BT, Wiede F, Braun A, Golledge J, Arolt V, Koerner H: Cognitive dysfunction in mice deficient for TNF $\alpha$  and its receptors. *Am J Med Genet B* 2008, **147B**:1056-1064.
70. Hayley S, Brebner K, Lacosta S, Merali Z, Anisman H: Sensitization to the effects of tumor necrosis factor- $\alpha$ : neuroendocrine, central monoamine, and behavioral variations. *J Neurosci* 1999, **19**:5654-5665.
71. Jacobs BL, Fornal CA: Serotonin and Behavior: a General Hypothesis. In *Psychopharmacology: The Fourth Generation of Progress* Edited by: Bloom FE, Kupfer DJ. New York: Raven Press; 1995:461-469.
72. Todd RD, Botteron KN: Is attention-deficit/hyperactivity disorder an energy deficiency syndrome? *Biol Psychiatry* 2001, **50**:151-158.
73. Oades RD, Slusarek M, Velling S, Bondy B: Serotonin platelet-transporter measures in childhood attention-deficit/hyperactivity disorder (ADHD): clinical versus experimental measures of impulsivity. *World J Biol Psychiatry* 2002, **3**:96-100.
74. Flory JD, Newcorn JH, Miller C, Harty S, Halperin JM: Serotonergic function in children with attention - deficit hyperactivity disorder Relationship to later antisocial personality disorder. *Br J Psychiatry* 2007, **190**:410-414.
75. Myint AM, Schwarz MJ, Steinbusch HWM, Leonard BE: Neuropsychiatric disorders related to interferon and interleukins treatment. *Metabol Brain Dis* 2009, **24**:55-68.
76. Byrne GI, Lehmann LK, Kirschbaum JG, Borden EC, Lee CM, Brown RR: Induction of tryptophan degradation in vitro and in vivo: a gamma-interferon-stimulated activity. *J Interferon Res* 1986, **6**:389-396.
77. Oades RD: Dopamine may be 'hyper' with respect to noradrenaline metabolism, but 'hypo' with respect to serotonin metabolism in children with ADHD. *Behav Brain Res* 2002, **130**:97-101.
78. Stone TW, Darlington LG: Endogenous kynurenines as targets for drug discovery and development. *Nat Rev Drug Discovery* 2002, **1**:609-620.
79. Alkondon M, Pereira EF, Yu P, Arruda EZ, Almeida LE, Guidetti P, Fawcett WP, Sapko MT, Randall WR, Schwarcz R, Tagle DA, Albuquerque EX: Targeted deletion of the kynurenine aminotransferase ii gene reveals a critical role of endogenous kynurenine acid in the regulation of synaptic transmission via alpha7 nicotinic receptors in the hippocampus. *J Neurosci* 2004, **24**:4635-4648.
80. Prescott C, Weeks AM, Staley KJ, Partin KM: Kynurenine acid has a dual action on AMPA receptor responses. *Neurosci Lett* 2006, **402**:108-112.
81. Leipnitz G, Schumacher C, Dalcin KB, Scussiato K, Solano A, Funchal C, Dutra-Filho CS, Wyse ATS, Wannmacher CMD, Latini A, Wajner M: In vitro evidence for an antioxidant role of 3-hydroxykynurenine and 3-hydroxyanthranilic acid in the brain. *Neurochem Int* 2006, **50**:83-94.
82. Ribeiro CAJ, Grando V, Filho CSD, Wannmacher CMD, Wajner M: Evidence that quinolinic acid severely impairs energy metabolism through activation of NMDA receptors in striatum from developing rats. *J Neurochem* 2006, **99**:1531-1542.
83. Costantino G: New promises for manipulation of kynurenine pathway in cancer and neurological diseases. *Expert Opin Ther Targets* 2009, **13**:247-258.
84. Oades RD, Dauvermann MR, Schwarz MJ, Myint AM: Does glial function underlie ADHD variability? Evidence from measures of S100B, interleukins, tryptophan metabolism and the kynurenine metabolic pathway. *Neurol Psychiatry Brain Res* 2009:44.

doi: 10.1186/1744-9081-6-32

Cite this article as: Oades et al., Attention-deficit hyperactivity disorder (ADHD) and glial integrity: an exploration of associations of cytokines and kynurenine metabolites with symptoms and attention *Behavioral and Brain Functions* 2010, **6**:32

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

