

Papain-like cysteine proteases are required for the regulation of photosynthetic gene expression and acclimation to high light stress

Alomrani, Sarah; Kunert, Karl J; Foyer, Christine H

DOI:

[10.1093/jxb/erab101](https://doi.org/10.1093/jxb/erab101)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Alomrani, S, Kunert, KJ & Foyer, CH 2021, 'Papain-like cysteine proteases are required for the regulation of photosynthetic gene expression and acclimation to high light stress', *Journal of Experimental Botany*, vol. 72, no. 9, pp. 3441-3454. <https://doi.org/10.1093/jxb/erab101>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

RESEARCH PAPER

Papain-like cysteine proteases are required for the regulation of photosynthetic gene expression and acclimation to high light stress

Sarah Alomrani^{1,2}, Karl J. Kunert³ and Christine H. Foyer^{1,*} 

¹ School of Biosciences, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK

² Centre for Plant Sciences, School of Biology, Faculty of Biological Sciences, University of Leeds, Leeds LS2 9JT, UK

³ Department of Plant and Soil Sciences, Forestry and Agricultural Biotechnology Institute, University of Pretoria, Pretoria, 0002, South Africa

* Correspondence: C.H.Foyer@bham.ac.uk

Received 29 September 2020; Editorial decision 26 February 2021; Accepted 1 March 2021

Editor: Robert Sharwood, Western Sydney University, Australia

Abstract

Chloroplasts are considered to be devoid of cysteine proteases. Using transgenic *Arabidopsis* lines expressing the rice cystatin, oryzacystatin I (OC-I), in the chloroplasts (PC lines) or cytosol (CYS lines), we explored the hypothesis that cysteine proteases regulate photosynthesis. The CYS and PC lines flowered later than the wild type (WT) and accumulated more biomass after flowering. In contrast to the PC rosettes, which accumulated more leaf chlorophyll and carotenoid pigments than the WT, the CYS lines had lower amounts of leaf pigments. High-light-dependent decreases in photosynthetic carbon assimilation and the abundance of the Rubisco large subunit protein, the D1 protein, and the phosphorylated form of D1 proteins were attenuated in the CYS lines and reversed in the PC lines relative to the WT. However, the transgenic lines had higher amounts of *LHC*, *rbcs*, *pasbA*, and *pasbD* transcripts than the WT, and also showed modified chloroplast to nucleus signalling. We conclude that cysteine proteases accelerate the re-configuration of the chloroplast proteome after flowering and in response to high-light stress. Inhibition of cysteine proteases, such as *AtCEP1*, slows chloroplast protein degradation and stimulates photosynthetic gene expression and chloroplast to nucleus signalling, enhancing stress tolerance traits.

Keywords: Acclimation, carbon assimilation, chloroplast to nucleus signalling, D1, GUN pathway, high light, photosynthetic gene expression, protein turnover, Rubisco.

Introduction

While plant cysteine proteases fulfil many important functions in plants, they have never been shown to localize to chloroplasts, except in barley (Frank *et al.*, 2019). The major chloroplast protease families (Clp, FtsH, and DegP) are constitutively

expressed and fulfil roles in protein quality control and the maintenance of photosynthesis rather than large-scale remodelling of the chloroplast proteome. Plastid functions are regulated by the ubiquitin–proteasome system (UPS) through the action of the SP1 ubiquitin E3 ligase and the SP2 channel protein that are localized on the outer chloroplast envelope

membrane (Ling *et al.*, 2012, 2019). SP1 and SP2 target the plastid protein import machinery [TOC (translocon of the outer membrane) proteins] for degradation in a novel proteolytic pathway, which is called chloroplast-associated protein degradation (CHLORAD). This pathway regulates the TOC machinery to facilitate reconfiguration of the plastid proteome, which is essential during stages of plant development that involve the interconversion of different plastid types, for example reconfiguration of plastid proteins when chloroplasts are converted into gerontoplasts during leaf senescence (Ling *et al.*, 2012, 2019). Moreover, CHLORAD has an additional role in abiotic stress responses (Ling *et al.*, 2019).

Cysteine proteases can be divided into 14 superfamilies, each with a catalytic triad or dyad in a different structural fold (Rawlings *et al.*, 2018). Small ubiquitin-like modifier (SUMO) cysteine proteases regulate the SUMO cycle by cleaving SUMO from SUMOylated proteins, providing specificity to which proteins become SUMOylated and generating free SUMO (Botha *et al.*, 2017; Morrell and Sadanandom, 2019). The papain-like cysteine proteases (PLCPs) are classified into clan CA because of the structural similarity of their conserved catalytic residues to papain (Rawlings *et al.*, 2018). Thirty-one genes encoding PLCPs have been identified in the Arabidopsis genome. These enzymes play a key role in the regulation of protein turnover, particularly in leaf senescence and the recycling of nutrients from the senescing leaves to growing leaves and reproductive organs (Díaz-Mendoza *et al.*, 2014; Liu *et al.*, 2018). The activities of PLCPs are tightly controlled via auto-catalytic post-translational modifications, as well as by cystatins and serpins (van der Linde *et al.*, 2012; Lampl *et al.*, 2013). PLCPs contain the conserved catalytic triad (Cys, His, Asn). They perform a nucleophilic attack of the thiol group at the substrate C-terminus where His acts as a proton acceptor (base) for the catalytic Cys, and Asn plays an important role for the orientation of His (Rawlings *et al.*, 2018).

Cystatins are proteins that contain a Gln–Xaa–Val–Xaa–Gly motif in the centre of the polypeptide chain (where Xaa is any amino acid), a Pro–Trp (or Leu–Trp) dipeptide motif in the C-terminal region, and a conserved Gly residue in the N-terminus (Benchabane *et al.*, 2010). Cystatins bind to the active site of their target cysteine proteases and inhibit enzyme activity in an irreversible manner. Studies of *in vivo* interactions between a cystatin and cathepsin L-like cysteine protease showed co-localization to the endoplasmic reticulum and the Golgi complex (Martinez *et al.*, 2009). The functions of PLCPs and phytocystatins have been intensively studied, particularly in relation to plant development, stress responses, and pest control (Van der Hoorn, 2008; Kunert *et al.*, 2015). In particular, oryzacystatin I (OC-I), which is a small protein of 120 amino acids that is a competitive inhibitor of PLCPs (Benchabane *et al.*, 2010), has been extensively studied in a range of transgenic plants including tobacco (Van der Vyver *et al.*, 2003; Prins *et al.*, 2008), soybean, and Arabidopsis (Quain

et al., 2014). These studies showed that cysteine proteases play a role in chloroplast protein turnover (Prins *et al.*, 2008).

Despite intensive efforts, molecular-genetic and biochemical studies have failed to identify chloroplast-localized cysteine proteases in Arabidopsis and plant species. However, a barley C1A family cysteine protease, called HvPAP14, was recently found in the thylakoid lumen of the chloroplasts, as well as the endoplasmic reticulum and vesicular bodies (Frank *et al.*, 2019). HvPAP14 is involved in the degradation of the light-harvesting Chl *a/b*-binding proteins and the large subunit of Rubisco (Frank *et al.*, 2019). There is only one other report which has provided evidence of cysteine protease activity in the thylakoid lumen (Sokolenko *et al.*, 1997).

While a small number of proteins are encoded by chloroplast genes, most of the photosynthetic proteins are encoded by the nuclear genome. This requires the coordinated regulation of the transcriptional activity of both genomes, a process that involves bi-directional communication from the nucleus to the chloroplasts (anterograde signalling), and from chloroplasts to the nucleus (retrograde signalling). Retrograde signals provide information concerning the metabolic and energy state of the chloroplasts that facilitates appropriate responses to environmental cues (Chan *et al.*, 2016). Mature chloroplasts produce a wide range of signalling molecules that act as retrograde signals, such as tetrapyrroles, that affect the expression of plastid redox-associated nuclear genes (PHANGs) in the nucleus in response to environmental cues (Chan *et al.*, 2016). Retrograde signalling between the chloroplasts and nuclei has been intensively studied using the carotenoid synthesis inhibitor norflurazon (NF) and/or the plastid translation inhibitor, lincomycin (LINCO). These inhibitors cause oxidation of the cytosol and nuclei, leading to the repression of nuclear genes encoding components of the photosynthetic electron transfer chain such as the light-harvesting Chl *a/b*-binding complex proteins and plastocyanin (Grubler *et al.*, 2017; Karpinska *et al.*, 2017). Screens using these inhibitors have revealed the plastid-localized genomes uncoupled (GUN) retrograde signalling pathway (Shimizu *et al.*, 2019).

This study was conducted to test the hypothesis that cysteine proteases play a role in reconfiguration of the chloroplast proteome and photosynthesis in response to changes in environmental stimuli such as light intensity. In order to test whether cysteine protease activities localized within chloroplasts play a role in this regulation, we prepared transgenic Arabidopsis lines expressing the rice cystatin OC-I targeted to the chloroplast stroma. We compared shoot traits and photosynthetic properties in transgenic lines with OC-I-dependent inhibition of cysteine proteases targeted to the chloroplasts with those where OC-I was expressed with no intracellular targeting and hence the product of the transgene is presumed to be localized in the cytosol. We show that the growth and shoot phenotypes of transgenic lines expressing OC-I in the chloroplasts (PC lines) or in the cytosol (CYS lines) were modified in the

OC-I-expressing lines compared with wild-type (WT) plants. In addition, evidence is presented showing that exposure to high light (HL), which is widely accepted as a stress that triggers local and systemic signalling to regulate stomatal closure and activate the expression of defence-related genes (Karpinski *et al.*, 1999; Devireddy *et al.*, 2018), leads to a substantial increase in leaf protease activities, particularly cysteine protease activities in the WT but not in the PC and CYS lines. The absence of HL-induced increases in leaf cysteine protease activities protects photosynthesis against HL-induced inhibition by limiting the decreases in the abundance of the Rubisco large subunit protein, the D1 protein, and the phosphorylated form of D1 that were observed in the WT plants. Moreover, photosynthetic gene expression was altered in the PC and CYS lines, in a manner that suggests that cysteine proteases are involved in chloroplast to nucleus signalling.

Materials and methods

Production of transgenic Arabidopsis lines expressing OC-I targeted to the chloroplasts

The OC-I gene was cloned as a *SacI*–*XbaI* fragment into the plasmid pLBR19 to create plasmid pLBR Cys-1. The pLBRPRKCys-1 plasmid contained the OC-I gene with expression under the control of the double 35S promoter of the *Cauliflower mosaic virus* (CaMV), an ampicillin resistance gene, and a CaMV terminator. The pLBRPRKCys-1 plasmid also contained the sequence encoding the signal peptide of the stomatal enzyme phosphoribulokinase (PRK), leading to targeting of the OC-I protein in the chloroplast stroma. Transgenic plants expressing the plasmid were prepared by the floral dip method from the *Agrobacterium tumefaciens* strain GV3101. Successive generations were screened for herbicide resistance and expression of the transgene. T₄ generation plants that were homozygous for the transgene were used in the following experiments.

Plant material and growth conditions

Seeds of WT *Arabidopsis thaliana* (L.) and six independent transgenic lines overexpressing OC-I either with the chloroplast PRK targeting sequence (PC2, PC7, and PC9) or without a targeting sequence for localization in the cytosol (CYS1, CYS3, and CYS4; Makgopa, 2014; Quain *et al.*, 2014) were used in the following experiments. Unless otherwise stated below or in the figure legends, each experiment involved six T₄ generation plants from each line, together with six WT plants. Plants were grown in 10 cm pots containing potting compost for up to 12 weeks in controlled-environment chambers with a 16 h/8 h day/night photoperiod (light intensity of 250 $\mu\text{mol m}^{-2} \text{s}^{-1}$) at 20 °C, 60% humidity. All genotypes were grown at the same time in the same controlled-environment chambers.

High-light treatments

The transgenic and WT *Arabidopsis* plants were grown in compost in controlled-environment chambers under a moderate light (LL) intensity (250 $\mu\text{mol m}^{-2} \text{s}^{-1}$) for 5 weeks. Thereafter, before the start of the next photoperiod, half of the plants were transferred to HL (800 $\mu\text{mol m}^{-2} \text{s}^{-1}$) supplied by light-emitting diodes (LEDs) in the controlled-environment chambers and subjected to HL for the whole 8 h photoperiod, while the remaining half were maintained for 8 h at 250 $\mu\text{mol m}^{-2} \text{s}^{-1}$. Leaf samples were collected between 7 h and 8 h into the 8 h photoperiod. The metabolism of harvested material was arrested immediately by immersion

in liquid nitrogen. Thereafter, frozen samples were stored at –80 °C for further analysis. Six plants per line were used in each experiment. All lines were grown together in each experiment and subjected to the different light treatments simultaneously. Each experiment was repeated at least three times.

Shoot growth analysis

At the time points indicated in the figure legends, the WT, PC, and CYS lines were harvested and separated into shoots and roots. The shoots were immediately weighed and then placed in an oven at 80 °C for 2 d to determine shoot dry biomass.

The time of flowering—the point at which the plants produce flowering stems—was also recorded.

Leaf pigment analysis

Leaf chlorophyll and protein contents were determined in leaf samples that had been ground in liquid nitrogen. Pigments extracted in 96% (v/v) ethanol were determined according to Lichtenthaler and Wellburn (1983). The soluble protein content was determined according to the method of Bradford (1976).

Photosynthetic pigments were extracted from frozen leaf samples that had been ground in liquid nitrogen by the addition of 80% (v/v) acetone and were measured as described by Lichtenthaler (1987).

Leaf protein analysis

Leaf proteins were extracted using protein extraction buffer (AS08 300, Agrisera). Protein concentrations (mg g^{-1}) were determined using the Pierce Microplate BCA Protein Assay Kit (Thermo Scientific) employing a Fluostar Omega plate reader (BMG Labtech GmbH).

Western blot analysis

Protein samples (10 μg) were separated on 4–20% Mini-PROTEAN[®] TGX™ pre-cast gels (Bio-Rad). A typical example of one Coomassie Brilliant Blue-stained gel is shown in Supplementary Fig. S1. After electrophoresis, proteins were transferred to (0.45 μm) nitrocellulose membranes (Amersham 10600003). Individual membranes were then incubated with the following primary antibodies at a 1:10 000 dilution: Rubisco large subunit (RbcL) form I and form II (AS03 037, Agrisera, Sweden), the PSII D1 protein (AS05 084, Agrisera, Sweden), and PsbA (D1) and the phosphorylated form of D1 (AS13 2669, Agrisera, Sweden). A typical example of an individual Western blot is shown in Supplementary Fig. S1, together with a typical loading control gel.

Total protease activity and cysteine protease activity

Protease activities were determined using the Abcam Assay Kit (ab111750), which incorporates fluorescein isothiocyanate (FITC)-labelled casein as a general protease substrate. The FITC-casein substrate is broken down into small peptides by the leaf proteases, resulting in a decrease in fluorescence quenching. The fluorescence of peptide fragments was estimated at an excitation/emission (Ex/Em) wavelength of 485/530 nm. Cysteine protease activities were determined by pre-incubation of the samples with the specific cysteine protease inhibitor E-64 (5 mM; Sigma).

Photosynthetic gas exchange measurements

All genotypes were grown in pots together in the same controlled-environment chambers under LL (250 $\mu\text{mol m}^{-2} \text{s}^{-1}$) for 5 weeks. Thereafter, half of the plants were transferred to HL (800 $\mu\text{mol m}^{-2} \text{s}^{-1}$) for 8 h, while the remaining half were maintained for 8 h at 250 μmol

$\text{m}^{-2} \text{s}^{-1}$. Photosynthetic gas exchange measurements were performed on plants in the controlled-environment chambers between 7 h and 8 h into the photoperiod under each irradiance.

Measurements were performed on fully expanded leaves essentially as described by Soares *et al.* (2008) using a LICOR portable photosynthetic gas exchange system (Model LI-6400XT).

Photosynthetic CO_2 assimilation rates were measured on whole leaves at 420 ppm CO_2 , 20 °C, at an irradiance of 400 $\mu\text{mol m}^{-2} \text{s}^{-1}$. Leaves were allowed to reach steady-state gas exchange conditions by maintaining the leaves for 10 min in the light and CO_2 conditions of the chamber prior to measurements. All experiments were conducted at 60% relative humidity. Vapour water deficits were kept constant throughout the assays.

Chl *a* fluorescence quenching measurements

Chl *a* fluorescence quenching measurement were performed on intact leaves using the LICOR portable photosynthetic gas exchange system (Model LI-6400XT) equipped with the leaf chamber fluorometer between 7 h and 8 h into the photoperiod under each irradiance. Plants were dark adapted for 45 min before each procedure (Plumb *et al.*, 2018). F_m and F_o , the maximum and minimum yields of fluorescence, respectively, were measured in dark-adapted leaves. Dark-adapted F_v/F_m , Φ_{PSII} , and non-photochemical quenching (NPQ) were calculated as $F_v/F_m = (F_m - F_o)/F_m$, $\Phi_{\text{PSII}} = (F_m' - F_o)/F_m'$, and $\text{NPQ} = (F_m - F_m')/F_m'$, respectively.

Norflurazon and lincomycin treatments

Seeds of the transgenic *A. thaliana* and WT lines were grown in the absence or the presence of inhibitors using a standard lab protocol (Karpinska *et al.*, 2017). CYS, PC, and WT seeds were grown on half-strength Murashige and Skoog medium containing 0.1 g l^{-1} myoinositol, 10 g l^{-1} sucrose, and 0.5 g l^{-1} MES buffer (pH 5.7) supplemented with either NF (5 μM), LINCO (500 μM), or ethanol as a control. Seedlings were grown for 7 d in a controlled-environment chamber with a 16 h/8 h day/night photoperiod and a daytime irradiance of 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$. Seedlings were then harvested for RNA extraction and further analysis. Each experiment, which involved 20 seedlings per line, was repeated at least three times.

Quantitative real-time reverse transcription-PCR (qRT-PCR)

For these analyses, plants were grown on compost in controlled-environment chambers at 250 $\mu\text{mol m}^{-2} \text{s}^{-1}$ for 5 weeks. Thereafter, half of the plants were transferred to 800 $\mu\text{mol m}^{-2} \text{s}^{-1}$ for 8 h. Six plants per line were used in each experiment, and each experiment was repeated three times. Leaf samples were collected between 6 h and 8 h into the photoperiod. Metabolism was arrested immediately in liquid nitrogen, and samples were stored at -80 °C for analysis. RNAs were extracted from leaves using the Spectrum™ plant total RNA kit protocol. Following synthesis of cDNA using the QuantiTect Reverse Transcription Kit (Qiagen), qRT-PCR was performed using a Quantifast SYBR Green RT-PCR Kit (Qiagen). The expression of the genes of interest was performed using the primer sequences given in Supplementary Table S1 relative to two (actin and SAND) control transcripts.

Statistical analysis

All datasets were analyzed by a two-ANOVA. If an independent variable (e.g. light) had a significant effect on the response ($P \leq 0.05$), Tukey's multiple comparison test was used to compare means between genotypes, while Sidak's multiple comparison test was applied to compare light effects within each genotype. For datasets which were normally distributed, Tukey's HSD (honestly significant difference) was used as a post-hoc test

at a stringency level of $P < 0.05$. Asterisks indicate statistical significance as follows: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, and **** $P < 0.0001$. Each measurement in the phenotypic analysis involved 24 plants per line, and three replicates were performed for each experiment. All statistical analysis was performed using SPSS v.13 for windows (Statistical Package for Social Sciences, Chicago, IL, USA).

Results

Shoot phenotypes

The plants expressing OC-I without a targeting sequence (CYS1, CYS3, and CYS4) had a similar shoot phenotype to the WT at 6 weeks (Fig. 1A). However, the transformed lines expressing OC-I in the chloroplast (PC2, PC7, and PC9) were visibly smaller than the WT at 6 weeks. Moreover, the WT plants had flowered at 6 weeks but the PC lines were without flowers (Fig. 1A). All of the transgenic lines flowered later than the WT (Fig. 1B), the PC lines showing the greatest delay in flowering (Fig. 1C).

The CYS (Fig. 2A) and PC (Fig. 2B) lines had a lower shoot biomass than the WT during vegetative development but, after flowering, shoot biomass was greater than that of the WT. The CYS leaves had similar amounts of chlorophyll (Fig. 3A) and carotenoid (Fig. 3B) pigments and protein (Fig. 3C) to the WT during vegetative development. However, after flowering, the CYS leaves had significantly lower amounts of chlorophyll (Fig. 3A) and carotenoid (Fig. 3B) pigments than the WT, but higher amounts of leaf protein (Fig. 3C). In contrast to the CYS lines, the PC lines had higher amounts of chlorophyll (Fig. 3A) and carotenoid (Fig. 3B) pigments than the WT throughout development. After flowering, the PC leaves accumulated greater amounts of leaf protein (Fig. 3C) than the WT.

Effects of high light on photosynthesis

Photosynthetic CO_2 assimilation rates were slightly lower in the CYS and PC lines than in the WT under the growth light intensity (250 $\mu\text{mol m}^{-2} \text{s}^{-1}$) conditions (Fig. 4A, B). However, after transfer to HL (800 $\mu\text{mol m}^{-2} \text{s}^{-1}$), photosynthesis rates were significantly decreased in the WT plants. In contrast, photosynthetic CO_2 assimilation rates were increased in the CYS (Fig. 4A) and PC lines (Fig. 4B) under HL compared with LL, and were significantly higher in the transgenic lines than in the WT under HL conditions. Photosynthesis was 60% higher in the PC lines than in the WT under HL conditions (Fig. 4B).

The calculated values for the flexible NPQ mechanisms of the photosynthetic electron transport system that allow the dissipation of excess light energy as heat were lower in the CYS and PC lines than in the WT under LL (Fig. 4C, D). While exposure to HL slight decreased NPQ values in the WT, NPQ levels were significantly increased in the CYS and PC lines under HL (Fig. 4C, D).




Fig. 1. The phenotype of wild-type *A. thaliana* (WT) and six independent transgenic lines expressing OC-I either in the cytosol (CYS1, CYS3, and CYS4) or in the chloroplast (PC2, PC7, and PC9). Representative images of the shoot phenotypes of 6-week-old plants (A), the number of flowering plants per week (B), and flowering time trends for the different lines (C). $n=24$ plants per line. Scale bar=3 cm.

Effects of high light on leaf protease activities

The total protease activities of the leaves were significantly lower in the CYS lines than in the WT under LL conditions (Fig. 5A, B). Exposure to HL for 8 h significantly increased

WT leaf total protease activities (Fig. 5A, B). However, the HL-dependent increases in the leaf protease activities were much lower in the CYS lines than in the WT (Fig. 5A). In addition, no HL-dependent increases in leaf protease activities




Fig. 2. A comparison of shoot biomass (dry mass per plant) in the *A. thaliana* plants expressing OC-I either in the cytosol (A: CYS1, CYS3, and CYS4) or in the chloroplast (B; PC2, PC7, and PC9) compared with the WT plants at 4, 6, 8, 10, and 12 weeks after sowing. Bars show the means \pm SD ($n=24$ plants per line). Statistical significance is indicated by asterisks: P -value ≤ 0.05 , $**P$ -value ≤ 0.01 .

were observed in the PC lines, which had even lower total protease activities under HL than under LL (Fig. 5B). HL significantly increased the leaf cysteine protease activities of the WT (Fig. 5C, D). However, the HL-dependent increases in leaf cysteine protease activities were, however, significantly lower in the CYS and PC lines than in the WT. While the CYS and PC lines had similar cysteine protease activities under LL (Fig. 5D), the PC lines had higher cysteine protease activities than the CYS lines under HL conditions (Fig. 5C).

The abundance of the Rubisco large subunit protein, the D1 protein, and the phosphorylated form of the D1 protein was lower in the leaves of the CYS lines than in those of the WT under LL conditions (Fig. 6). In contrast, the amounts of these photosynthetic proteins were similar in the PC and WT lines (Fig. 6). The amounts of the Rubisco large subunit protein, the D1 protein, and the phosphorylated form of D1 were lower under HL than under LL in the WT plants (Fig. 6). In contrast, the amounts of the photosynthetic proteins were further higher in the Cys and PC lines under HL than under LL (Fig.

6). The HL-dependent increases in the amounts of the chloroplast proteins were most marked in the PC lines.

Photosynthetic gene expression

We next compared the effects of OC-I expression on the abundance of transcripts encoding chloroplast proteins in the leaves of soil-grown plants. The amounts of transcripts arising from the nuclear-localized genes encoding the light-harvesting Chl *a/b*-binding proteins (LHCA, LHCB1, and LHCB2) and chloroplast-localized genes (*rbcS*, *psbA*, and *psbD*) encoding the small subunit of Rubisco, the PSII D1 protein, and the PSII D2 protein, respectively, were increased in the CYS and PC lines compared with the WT under LL conditions (Fig. 7). In general, the amounts of these transcripts were generally higher in the PC lines than in the CYS lines under LL (Fig. 7A). While the amounts of all of these transcripts was even further increased in the CYS lines relative to the WT under HL conditions, the effect of HL was less pronounced in the PC lines than in the CYS lines (Fig. 7). Moreover, in contrast to all other mRNAs measured in this study, the amounts of transcripts encoding the LHCB2 protein were lower in the PC lines than in the WT under HL (Fig. 7). Hence, there are clear differences in the effects of targeting OCI to chloroplasts or the cytosol in the responses of the transcript profile to HL (Fig. 7).

The data shown in Fig. 7 indicate that the presence of OC-I alters the expression of genes encoding photosynthetic proteins in both the nucleus and chloroplasts. Moreover, differential effects of the OC-I-dependent inhibition of cysteine proteases in chloroplasts on the transcript profile were observed compared with OC-I-dependent inhibition of cysteine proteases in the cytosol. These results suggest that cysteine proteases in both the chloroplasts and the cytosol play roles (and possibly different roles) in chloroplast to nucleus signalling. To explore this possibility further, seedlings were grown for 7 d on agar plates at an irradiance of $100 \mu\text{mol m}^{-2} \text{s}^{-1}$, with or without the chloroplast inhibitors LINCO or NF. The inhibitor treatments led to bleaching of the seedlings (Fig. 8A). Under these LL conditions in the absence of inhibitors, the amounts of *LHCA* (Fig. 8B), *LHCB1* (Fig. 8C), and *LHCB2* (Fig. 8D) transcripts were significantly lower in the CYS and PC lines than in the WT in the absence of inhibitors, but the abundance of *LHCB1* (Fig. 8C) and *LHCB2* (Fig. 8D) mRNAs tended to be higher in the PC lines than in the CYS lines under these conditions.

The LINCO treatment decreased the amounts of the *LHCA* (Fig. 8B), *LHCB1* (Fig. 8C), and *LHCB2* (Fig. 8D) transcripts in the WT but greatly increased the amounts of these mRNAs in the CYS and PC lines, the stimulatory effect being greater in the former than in the latter. In comparison with the effects of LINCO on *LHCA* transcripts in the CYS and PC lines (Fig. 8B), the presence of this inhibitor led to smaller increases in the amounts of *LHCB1* (Fig. 8C) and *LHCB2* (Fig. 8D) transcripts.




Fig. 3. A comparison of the (A) chlorophyll and (B) carotenoid pigment, and (C) soluble protein amounts in the leaves of *A. thaliana* plants expressing OC-I either in the cytosol (CYS1, CYS3, and CYS4) or in the chloroplast (PC2, PC7, and PC9) compared with the wild-type (WT) plants at 4, 6, 8, 10, and 12 weeks after sowing. Bars represent means \pm SD ($n=24$ plants per line). Statistical significance as determined by *t*-testing is indicated by asterisks: *P*-value ≤ 0.05 , ***P*-value ≤ 0.01 .

The NF treatment decreased the amounts of the *LHCA* (Fig. 8B) *LHCB1* (Fig. 8C), and *LHCB2* (Fig. 8D) transcripts in the WT but greatly increased the amounts of these mRNAs in the CYS lines. In contrast, this inhibitor increased the amounts of *LHCA* (Fig. 8B) and *LHCB2* (Fig. 8D) transcripts in the PC lines, but decreased the as of *LHCB1* (Fig. 8C) transcripts to values lower than those observed in the WT under these conditions.

Discussion

Chloroplasts belong to the plastid family of plant organelles. Unlike etioplasts that are found in leaves that emerge in the dark and gerontoplasts, which are dominant in senescent leaves,

chloroplasts are photosynthetically competent. The different plastid types present in leaves interconvert in response to environmental and developmental triggers as well as in response to biotic and abiotic stresses. These transitions require a complete remodelling of the plastid proteome that transforms the functions and developmental fate of the organelles. Degradation of chloroplast proteins is initiated by chloroplast proteases within the organelle, but the final steps of protein degradation occur outside the plastids through the action of other proteases. Serine, cysteine, aspartic, and metalloprotease expression and activities are detected in senescing leaves, but studies have not revealed the identities of the proteases responsible for chloroplast protein breakdown (Buet *et al.*, 2019). Cysteine proteases,




Fig. 4. The effects of high light (HL) on photosynthetic CO₂ assimilation rates and on the non-photochemical quenching of Chl *a* fluorescence (NPQ) in the leaves of CYS (A and C) and PC (B and D) lines compared with the WT. The CYS, PC, and WT plants were grown under moderate light (LL: 250 μmol m⁻² s⁻¹) for 5 weeks. Thereafter, half of the plants were transferred to HL (800 μmol m⁻² s⁻¹) for 8 h, while the remaining half were maintained for 8 h at 250 μmol m⁻² s⁻¹). Photosynthetic CO₂ assimilation and NPQ were measured at the end of the photoperiod in all lines. Bars represent means ± SD (*n*=24 plants per line). Statistical significance is indicated by asterisks: **P*-value ≤ 0.05.

which are important in leaves undergoing rapid protein turnover, are largely located in the central vacuole and other lytic compartments (Buet *et al.*, 2019). In addition, the chloroplast protein complement is reconfigured by regulation of the cytosolic ubiquitin–proteasome system by the CHLORAD-dependent regulation of the TOC machinery at different stages of plant development (Ling *et al.*, 2012, 2019). The data presented here show that cysteine proteases play an important role in determining the chloroplast transcript and proteomic profiles, particularly under different HL conditions.

Cystatins are thiol proteinase inhibitors that are ubiquitously present in plants, animals, and other organisms. The reaction of plant cystatins with cysteine proteases from the C1A papain-like family is highly specific. Cystatins bind to the active site of cysteine proteases as pseudo-substrates, thus preventing cleavage of peptide bonds (Barrett, 1987). While approaches, such as sites, have directed mutagenesis at positively selected amino acid sites have been used to generate cystatin variants with improved inhibitory potency and specificity against specific targets, such as the gut cysteine proteases of herbivorous insects (Goulet *et al.*, 2008), there is no record in the literature of cystatin-dependent inhibition of other types of proteases.

We have not identified which cysteine proteases are involved in the control of chloroplast gene expression and photosynthetic functions, but the results presented here clearly highlight their functional significance, particularly in relation to the chloroplast signalling and responses to HL stress. The study of transgenic plants, in which a cysteine protease inhibitor is expressed in the chloroplasts, allows precise characterization of the functions of putative chloroplast-localized cysteine proteases. In comparison, the consequences of OC-I expression in the cytosol allow characterization of the roles of chloroplast-associated cysteine proteases. We have already characterized the shoot phenotypes of a range of transgenic plants with untargeted expression of OC-I, including tobacco (Prins *et al.*, 2008) and soybean (Quain *et al.*, 2014). The results presented here are the first to characterize transgenic plants in which OC-I expression is targeted to the chloroplasts.

During vegetative growth, the CYS and PC lines had less biomass than the WT line, but the transgenic lines accumulated significantly more biomass during reproductive growth (Fig. 2). These data show that cysteine proteases accelerate shoot biomass production during vegetative growth but they have a negative impact on biomass production after the initiation of flowering. We conclude that the function of cysteine




Fig. 5. A comparison of the maximal extractable total protease activities and cysteine protease activities from the leaves of the CYS (A and C) and PC (B and D) lines compared with wild-type (WT) plants grown under moderate light (LL) and high light (HL) conditions. The CYS, PC, and WT plants were grown under moderate light ($250 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 5 weeks. Thereafter, half of the plants were transferred to HL ($800 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 8 h, while the remaining half were maintained for 8 h at $250 \mu\text{mol m}^{-2} \text{s}^{-1}$. The total protease activities and the cysteine protease activities of the leaves were measured between 7 h and 8 h into the photoperiod in all lines. Bars represent means \pm SD ($n=24$ individual plants per line). Statistical significance is indicated by asterisks: * P -value ≤ 0.05 .




Fig. 6. Western blot analysis of the Rubisco large subunit protein (RbcL), the D1 protein, and the phosphorylated form of the D1 protein in the leaves of the CYS and PC lines compared with the wild type (WT). The CYS, PC, and WT plants were grown under moderate light ($250 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 5 weeks. Thereafter, half of the plants were transferred to HL ($800 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 8 h, while the remaining half were maintained for 8 h at $250 \mu\text{mol m}^{-2} \text{s}^{-1}$. Soluble proteins were extracted from leaves that were harvested 8 h into the photoperiod. Proteins were loaded onto polyacrylamide gels ($10 \mu\text{g}$ of protein per well). After electrophoresis, proteins were transferred to nitrocellulose membranes and analysed with specific antibodies for each protein form.

proteases is to accelerate the turnover of chloroplast proteins in leaves after flowering and that OC-I-dependent inhibition of these enzymes has a positive effect on shoot biomass. The

slower vegetative growth of the CYS lines agrees with previously published results for OC-I-expressing transgenic tobacco plants (van der Vyver *et al.*, 2003; Prins *et al.*, 2008). The time to flowering was delayed in the CYS and PC lines compared with the WT (Fig. 1A, B). This delay was greatest in the PC lines, suggesting that chloroplast-localized proteins are important in the regulation of flowering and that chloroplast proteases that are blocked by OC-I are important determinants of the timing of flowering (Bryant *et al.*, 2011; Feng *et al.*, 2016).

The targeting of the OC-I transgene to the chloroplasts has a marked effect on leaf pigment contents, an effect that was absent from the CYS lines (Fig. 3A, B). The CYS lines had less chlorophyll (Fig. 3A) and carotenoid pigments (Fig. 3B) than the WT, particularly in the later stages of development. In contrast, the PC rosettes accumulated more leaf pigments than the WT at the later stages of leaf development, suggesting that senescence was delayed in the PC lines. These findings suggest that OC-I-inhibited protease targets in the chloroplasts are involved in pigment biosynthesis and/or degradation. Possible candidates are the degradation of the pigment-protein complexes (Buet *et al.*, 2019; Frank *et al.*, 2019) and the tetrapyrrole biosynthesis pathway (Eckhardt *et al.*, 2004; Tanaka and Tanaka, 2007). Both of these pathways can generate signals, such as singlet oxygen that are involved in chloroplast to nucleus signalling (Rintamäki *et al.*, 2009; Leister, 2019).




Fig. 7. A comparison of the relative abundance of nuclear-encoded chloroplast-targeted photosynthesis transcripts and chloroplast-encoded photosynthesis transcripts in the leaves of CYS, PC, and WT plants grown under moderate light (LL) and high light (HL) conditions. The CYS, PC, and WT plants were grown under moderate light ($250 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 5 weeks. Thereafter, half of the plants were transferred to HL ($800 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 8 h, while the remaining half were maintained for 8 h at $250 \mu\text{mol m}^{-2} \text{s}^{-1}$. The amounts of transcripts were measured in the leaves of the CYS, PC, and WT plants harvested between 7 h and 8 h into the photoperiod. *LHCA*, light-harvesting Chl *a*-binding protein A; *LHCB1*, light-harvesting Chl *a/b*-binding protein B1; *LHCB2*, light-harvesting Chl *a/b*-binding protein B2; *rbcS*, the small subunit of Rubisco; *psbA*, PSII D1 protein; *psbD*, PSII D2 protein. Bars represent means \pm SD ($n=24$ individual plants per line). Statistical significance is indicated by asterisks: * P -value ≤ 0.05 , ** P -value ≤ 0.01 .

Photosynthetic carbon assimilation was better protected against HL-induced inhibition in the CYS and PC lines than in the WT (Fig. 4). This finding may be related to the higher turnover of chloroplast proteins in the WT that is associated with a general increase in leaf protease activities, particularly the cysteine proteases. The amounts of chloroplast proteins (Fig. 6) and the abundance of transcripts (Fig. 7) encoding these proteins were differentially changed in the PC and CYS lines relative to the WT. For example, the Rubisco large subunit and the D1 protein were more abundant in the transgenic lines than in the WT following exposure to HL stress (Fig. 6). Rubisco degradation is a complex process that involves a network of different pathways. Oxidative modification and partial cleavage of the Rubisco protein occurs inside the chloroplasts and thereafter the protein is transported to the vacuole (Ono *et al.*, 2013). The transport of Rubisco and other stromal proteins to vacuolar compartments involves

several pathways. Stromal proteins can be degraded by a type of autophagosome known as the Rubisco-containing body (RCB) that allows efficient amino acid recycling (Ishida *et al.*, 2014). Autophagy was found to remove HL-damaged and collapsed chloroplasts in *A. thaliana* (Izumi *et al.*, 2017). Other pathways are independent of autophagy and involve small proteolytic vacuolar compartments (senescence-associated vacuoles) that accumulate during senescence (Martinez *et al.*, 2008). A further pathway is dependent on the chloroplast vesiculation (CV) protein, a process that again involves the formation of vesicles that contain stromal proteins, envelope membrane proteins, and thylakoid membrane proteins (Wang and Blumwald, 2014). In addition, singlet oxygen induces a pathway of chloroplast degradation that is induced by ubiquitination, a process that requires a cytoplasmic ubiquitin E3 ligase called PLANT U-BOX 4 (Woodson *et al.*, 2015). The data presented here implicate chloroplast and cytosolic cysteine proteases in the




Fig. 8. The effects of OC-I expression on chloroplast to nucleus signalling. Wild-type *A. thaliana* (WT) and six independent transgenic lines expressing OC-I either in the cytosol (CYS1, CYS3, and CYS4) or in the chloroplast (PC2, PC7, and PC9) were grown on agar plates under low light ($100 \mu\text{mol m}^{-2} \text{s}^{-1}$) with or without lincomycin (LINCO) or norflurazon (NF). (A) Comparisons of representative CYS1, CYS3, CYS4, PC2, PC7, PC9, and WT seedlings grown on agar plates containing either no added inhibitors, LINCO ($500 \mu\text{M}$), or NF ($5 \mu\text{M}$) for 7 d. The effect of LINC and NF on the amounts of (B) *LHCA*, (C) *LHCB1*, and (D) *LHCB2* transcripts in the WT, CYS, and PC lines. Bars represent means \pm SD ($n=24$ plants per line). Statistical significance as determined by *t*-testing is indicated by asterisks: **P*-value ≤ 0.05 .

Rubisco degradation process. These data agree with a recent report concerning the barley HvPAP14 protease, which is involved in the degradation of the large subunit of Rubisco and thylakoid membrane proteins (Frank *et al.*, 2019).

We undertook a phylogenetic analysis to identify Arabidopsis cysteine proteases with homology to HvPAP14. Of the 280 sequences found in the Arabidopsis genome, CEP1 was identified as the closest orthologue to the barley HvPAP14 protein, which was detected in the thylakoid lumen, endoplasmic reticulum, and cytosol (Frank *et al.*, 2019). Like HvPAP14, AtCEP1 carries a C-terminal KDEL motif, which leads to localization in the endoplasmic reticulum, but it has been suggested that AtCEP1 is also active in the cytosol (Howing *et al.*, 2017). Given the observed effects of OC-I expression on the turnover of the D1 protein and the light-harvesting Chl *a/b*-binding proteins, the chloroplast-localized cysteine protease should localize to the thylakoid lumen of the chloroplasts. However, further work is required to investigate the intracellular localization and possible roles of AtCEP1 in chloroplast protein turnover.

The amounts of transcripts encoding LHCA, LHCB1, LHCB2, and the D1 protein were increased in the CYS and PC lines relative to the WT (Fig. 7), suggesting that cysteine proteases in the chloroplasts and cytosol may be important for chloroplast to nucleus signalling. We therefore investigated effects on chloroplast to nucleus signalling through the GUN pathway using the chloroplast inhibitors NF and LINCO (Fig. 8) While the amounts of LHCA, LHCB1, and LHCB2 transcripts were higher in the CYS and PC lines than in WT when plants were grown in soil, the opposite was found when the plants were grown on plates in the absence of these inhibitors. The discrepancy in the effects of OC-I on photosynthetic gene expression reported in Figs 7 and 8 is likely to lead to differences in the growth conditions. For example, the plants on plates are grown at a much lower irradiance ($100 \mu\text{mol m}^{-2} \text{s}^{-1}$) than those grown on soil ($250 \mu\text{mol m}^{-2} \text{s}^{-1}$). Similarly, plants grown on plates are enclosed, with lids limiting the exchange of gases with the atmosphere. However, the observed differences in the amounts of transcripts encoding photosynthetic proteins in the CYS and PC lines under HL suggest that different cysteine proteases are involved in this regulation in the chloroplasts and cytosol.

The GUN pathway of chloroplast to nucleus signalling is impaired by OC-I expression in the chloroplasts and cytosol. As with the amounts of transcripts encoding photosynthetic proteins in the different genotypes grown on soil discussed above, variations in the responses of transcripts that are markers for chloroplast to nucleus signalling were observed in the CYS and PC lines. This finding again suggests that different cysteine proteases and processes may underpin the observed effects. Several components of the GUN pathway have been characterized. For example, GUN1 interacts with the chloroplast chaperone HSC70-1 to promote import of nuclear-encoded chloroplast proteins such as the tetrapyrrole synthesis proteins

(Shimizu, *et al.*, 2019). As well as binding to the tetrapyrrole synthesis proteins, GUN1 binds to haem and other porphyrins that affect the flux through the tetrapyrrole synthesis pathway (Shimizu *et al.*, 2019). Further work is required to understand how cysteine proteases are involved in these mechanisms.

In summary, the data presented here show that OC-I expression targeted to the chloroplasts has effects on the HL-induced changes in leaf total protease activities and cysteine protease activities. The level of inhibition of cysteine protease activities achieved with the chloroplast-localized OCI (PC lines) is similar to, but not the same, as that observed in the cytosol-targeted OCI (CYS lines). This suggests that OCI has different targets in both compartments, particularly under HL conditions. Under LL conditions, OC-I expression in the chloroplasts had no detectable effect on the maximum extractable cysteine protease activities from leaves, suggesting that few OC-I-inhibited cysteine proteases are present in chloroplasts under these conditions. However, exposure to HL significantly increased the total protease activities and cysteine protease activities of the WT leaves. Intriguingly, OC-I expression targeted to the chloroplasts resulted in a decrease in both leaf total protease activities and cysteine protease activities, suggesting that cysteine proteases are localized in the chloroplasts, as well as other cellular compartments under HL. Since no cysteine proteases have been identified that have targeting sequences directing the enzyme proteins to the chloroplasts, it is likely that post-translational modifications cause changes in compartmentation, as described for other enzymes (Foyer *et al.*, 2020). The implications of the study are significant in terms of future research on proteases, the regulation of photosynthesis, and the overall reconfiguration of plastid proteins that occurs in response to environmental and developmental triggers. The data presented here demonstrate that cysteine proteases are recruited not only when chloroplasts are converted into gerontoplasts during leaf senescence but also during acclimation to environmental stresses such as exposure to HL. Manipulation of chloroplast-localized cysteine proteases simply using OC-I or similar inhibitors, perhaps under the control of different promoters, may be a useful strategy for the improvement of the efficiency of photosynthesis, particularly in response to changing environmental conditions. Moreover, these data demonstrate that decreasing the stress-induced increases in the turnover of key proteins such as Rubisco has advantages in terms of protection of photosynthesis. Taken together, these data demonstrate that increasing the life span of key proteins by targeting cysteine proteases has advantages in terms of stress tolerance and ultimate productivity.

Supplementary data

The following supplementary data are available at [JXB online](#).

Fig. S1. Examples of the original gels used for Western blot analysis of the Rubisco, D1, and phosphorylated D1 proteins in the leaves of the CYS and PC lines compared with WT

Arabidopsis plants grown under moderate light (LL) and high light (HL) conditions.

Table S1. List of forward and reverse primers used in the (qRT) PCR analysis

Acknowledgements

This manuscript is dedicated to the memory of Dr Eugene Makgopa, who produced all the transgenic Arabidopsis lines used in these experiments. SA thanks the Ministry of Education and University of Tabuk, Saudi Arabia for a PhD Scholarship through the government PhD programme.

Author contributions

CHF planned the studies and wrote the manuscript. SA performed all the experimental work and produced the figures. KK provided expert advice during the course of the studies and undertook critical reading of the manuscript and evaluation of the data.

Data availability

All of the original datasets are available upon request from the authors.

References

- Barrett AJ.** 1987. The cystatins: a new class of peptidase inhibitors. *Trends in Biochemical Sciences* **12**, 193–196.
- Benchabane M, Schlüter U, Vorster J, Goulet MC, Michaud D.** 2010. Plant cystatins. *Biochimie* **92**, 1657–1666.
- Botha AM, Kunert KJ, Cullis CA.** 2017. Cysteine proteases and wheat (*Triticum aestivum* L) under drought: a still greatly unexplored association. *Plant, Cell & Environment* **40**, 1679–1690.
- Bradford MM.** 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein–dye binding. *Analytical Biochemistry* **72**, 248–254.
- Bryant N, Lloyd J, Sweeney C, Myounga F, Meinke D.** 2011. Identification of nuclear genes encoding chloroplast-localized proteins required for embryo development in Arabidopsis. *Plant Physiology* **155**, 1678–1689.
- Buet A, Costa ML, Martínez DE, Guamet JJ.** 2019. Chloroplast protein degradation in senescing leaves: proteases and lytic compartments. *Frontiers in Plant Science* **10**, 747.
- Chan KX, Phua SY, Crisp P, McQuinn R, Pogson BJ.** 2016. Learning the languages of the chloroplast: retrograde signaling and beyond. *Annual Review of Plant Biology* **67**, 25–53.
- Díaz-Mendoza M, Velasco-Arroyo B, González-Melendi P, Martínez M, Díaz I.** 2014. C1A cysteine protease–cystatin interactions in leaf senescence. *Journal of Experimental Botany* **65**, 3825–3833.
- Devireddy AR, Zandalinas SI, Gomez-Cadenas A, Blumwald E, Mittler R.** 2018. Coordinating the overall stomatal response of plants: rapid leaf-to-leaf communication during light stress. *Science Signaling* **11**, 9514.
- Eckhardt U, Grimm B, Hörtensteiner S.** 2004. Recent advances in chlorophyll biosynthesis and breakdown in higher plants. *Plant Molecular Biology* **56**, 1–14.
- Feng P, Guo H, Chi W, et al.** 2016. Flowering regulation by chloroplasts. *Proceedings of the National Academy of Sciences, USA* **113**, 10708–10713.
- Foyer CH, Baker A, Wright M, Sparkes IA, Mhamdi A, Schippers JHM, Van Breusegem F.** 2020. On the move: redox-dependent protein relocation in plants. *Journal of Experimental Botany* **71**, 620–631.
- Frank S, Hollmann J, Mulisch M, Matros A, Carrión CC, Mock HP, Hensel G, Krupinska K.** 2019. Barley cysteine protease PAP14 plays a role in degradation of chloroplast proteins. *Journal of Experimental Botany* **70**, 6057–6069.
- Goulet MC, Dallaire C, Vaillancourt LP, Khalf M, Badri AM, Preradov A, Duceppe MO, Goulet C, Cloutier C, Michaud D.** 2008. Tailoring the specificity of a plant cystatin toward herbivorous insect digestive cysteine proteases by single mutations at positively selected amino acid sites. *Plant Physiology* **146**, 1010–1019.
- Grubler B, Merendino L, Twardziok SO, et al.** 2017. Light and plastid signals regulate different sets of genes in the albino mutant Pap7-1. *Plant Physiology* **175**, 1203–1219.
- Howing T, Dann M, Hoefle C, Hüchelhoven R, Gietl C.** 2017. Involvement of *Arabidopsis thaliana* endoplasmic reticulum KDEL-tailed cysteine endopeptidase 1 (AtCEP1) in powdery mildew-induced and AtCPR5-controlled cell death. *PLoS One* **12**, e0183870.
- Ishida H, Izumi M, Wada S, Makino A.** 2014. Roles of autophagy in chloroplast recycling. *Biochimica et Biophysica Acta* **1837**, 512–521.
- Izumi M, Ishida H, Nakamura S, Hidema J.** 2017. Entire photodamaged chloroplasts are transported to the central vacuole by autophagy. *The Plant Cell* **29**, 377–394.
- Karpinska B, Alomrani SO, Foyer CH.** 2017. Inhibitor-induced oxidation of the nucleus and cytosol in *Arabidopsis thaliana*: implications for organelle to nucleus retrograde signalling. *Philosophical Transactions of the Royal Society B: Biological Sciences* **372**, 1730.
- Karpinski S, Reynolds H, Karpinska B, Wingsle G, Creissen G, Mullineaux P.** 1999. Systemic signaling and acclimation in response to excess excitation energy in Arabidopsis. *Science* **284**, 654–657.
- Kunert KJ, van Wyk SG, Cullis CA, Vorster BJ, Foyer CH.** 2015. Potential use of phytocystatins in crop improvement, with a particular focus on legumes. *Journal of Experimental Botany* **66**, 3559–3570.
- LampI N, Alkan N, Davydov O, Fluhr R.** 2013. Set-point control of RD21 protease activity by AtSerpin1 controls cell death in Arabidopsis. *The Plant Journal* **74**, 498–510.
- Leister D.** 2019. Piecing the puzzle together: the central role of reactive oxygen species and redox hubs in chloroplast retrograde signaling. *Antioxidants & Redox Signaling* **30**, 1206–1219.
- Lichtenthaler HK.** 1987. Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. *Methods in Enzymology* **148**, 350–382.
- Lichtenthaler, H, Wellburn A.** 1983. Determinations of total carotenoids and chlorophylls a and b of leaf extracts in different solvents. *Biochemical Society Transactions* **603**, 591–592.
- Ling Q, Broad W, Trösch R, Töpel M, Sert TD, Lymeropoulos P, Baldwin A, Jarvis P.** 2019. Ubiquitin-dependent chloroplast-associated protein degradation in plants. *Science* **363**, eaav4467.
- Ling Q, Huang W, Baldwin A, Jarvis P.** 2012. Chloroplast biogenesis is regulated by direct action of the ubiquitin–proteasome system. *Science* **338**, 655–659.
- Liu H, Hu M, Wang Q, Cheng L, Zhang Z.** 2018. Role of papain-like cysteine proteases in plant development. *Frontiers in Plant Science* **9**, 1717.
- Makgopa ME.** 2014. Production and characterization of transgenic Arabidopsis and soybean (*Glycine max* (L.) Merrill.) plants over-expressing oryzacystatin I (OC-I). PhD thesis, University of Pretoria.
- Martínez DE, Costa ML, Gomez FM, Otegui MS, Guamet JJ.** 2008. ‘Senescence-associated vacuoles’ are involved in the degradation of chloroplast proteins in tobacco leaves. *The Plant Journal* **56**, 196–206.
- Martínez M, Cambra I, Carrillo L, Díaz-Mendoza M, Díaz I.** 2009. Characterization of the entire cystatin gene family in barley and their target cathepsin L-like cysteine-proteases, partners in the hordein mobilization during seed germination. *Plant Physiology* **151**, 1531–1545.
- Morrell R, Sadanandom A.** 2019. Dealing with stress: a review of plant SUMO proteases. *Frontiers in Plant Science* **10**, 1122.
- Ono Y, Wada S, Izumi M, Makino A, Ishida H.** 2013. Evidence for contribution of autophagy to Rubisco degradation during leaf senescence in *Arabidopsis thaliana*. *Plant, Cell & Environment* **36**, 1147–1159.

- Plumb W, Townsend AJ, Rasool B, Alomrani S, Razak N, Karpinska B, Ruban AV, Foyer CH.** 2018. Ascorbate-mediated regulation of growth, photoprotection, and photoinhibition in *Arabidopsis thaliana*. *Journal of Experimental Botany* **69**, 2823–2835.
- Prins A, van Heerden PD, Olmos E, Kunert KJ, Foyer CH.** 2008. Cysteine proteinases regulate chloroplast protein content and composition in tobacco leaves: a model for dynamic interactions with ribulose-1,5-bisphosphate carboxylase/oxygenase (Rubisco) vesicular bodies. *Journal of Experimental Botany* **59**, 1935–1950.
- Quain MD, Makgopa ME, Magama F, Márquez García B, Montrose A, Schnaubelt D, Fernandez-Garcia N, Olmos E, Kunert KJ, Foyer CH.** 2014. Ectopic phytoalexin expression leads to enhanced drought stress tolerance in soybean (*Glycine max*) and *Arabidopsis thaliana* through effects on strigolactone pathways, and can also result in improved seed traits. *Plant Biotechnology Journal* **12**, 903–913.
- Rawlings ND, Barrett AJ, Thomas PD, Huang X, Bateman A, Finn RD.** 2018. The MEROPS database of proteolytic enzymes, their substrates and inhibitors in 2017 and a comparison with peptidases in the PANTHER database. *Nucleic Acids Research* **46**, D624–D632.
- Rintamäki E, Lepistö A, Kangasjärvi S.** 2009. Implication of chlorophyll biosynthesis on chloroplast-to-nucleus retrograde signaling. *Plant Signaling & Behavior* **4**, 545–547.
- Shimizu T, Kacprzak SM, Mochizuki N, et al.** 2019. The retrograde signaling protein GUN1 regulates tetrapyrrole biosynthesis. *Proceedings of the National Academy of Sciences, USA* **116**, 24900–24906.
- Soares AS, Driscoll SP, Olmos E, Harbinson J, Arrabaça MC, Foyer CH.** 2008. Adaxial/abaxial specification in the regulation of photosynthesis and stomatal opening with respect to light orientation and growth with CO₂ enrichment in the C₄ species *Paspalum dilatatum*. *New Phytologist* **177**, 186–198.
- Sokolenko A, Altschmied L, Herrmann RG.** 1997. Sodium dodecyl sulfate-stable proteases in chloroplasts. *Plant Physiology* **115**, 827–832.
- Tanaka R, Tanaka A.** 2007. Tetrapyrrole biosynthesis in higher plants. *Annual Review of Plant Biology* **58**, 321–346.
- van der Hoorn RA.** 2008. Plant proteases: from phenotypes to molecular mechanisms. *Annual Review of Plant Biology* **59**, 191–223.
- van der Linde K, Hemetsberger C, Kastner C, Kaschani F, van der Hoorn RA, Kumlehn J, Doeblemann G.** 2012. A maize cystatin suppresses host immunity by inhibiting apoplastic cysteine proteases. *The Plant Cell* **24**, 1285–1300.
- Van der Vyver C, Schneidereit J, Driscoll S, Turner J, Kunert K, Foyer CH.** 2003. Oryzacystatin I expression in transformed tobacco produces a conditional growth phenotype and enhances chilling tolerance. *Plant Biotechnology Journal* **1**, 101–112.
- Wang S, Blumwald E.** 2014. Stress-induced chloroplast degradation in *Arabidopsis* is regulated via a process independent of autophagy and senescence-associated vacuoles. *The Plant Cell* **26**, 4875–4888.
- Woodson JD, Joens MS, Sinson AB, Gilkerson J, Salomé PA, Weigel D, Fitzpatrick JA, Chory J.** 2015. Ubiquitin facilitates a quality-control pathway that removes damaged chloroplasts. *Science* **350**, 450–454.