## **UNIVERSITY** OF BIRMINGHAM University of Birmingham Research at Birmingham

### Genomic classification and clinical outcome in rhabdomyosarcoma

Shern, Jack F; Selfe, Joana; Izquierdo, Elisa; Patidar, Rajesh; Chou, Hsien-Chao; Song, Young K.; Yohe, Marielle E; Sindiri, Sivasish; Wei, Jun; Wen, Xinyu; Rudzinski, Erin R.; Barkauskas, Donald A; Lo, Tammy; Hall, David; Linardic, Corinne M; Hughes, Debbie; Jamal, Sabri; Jenney, Meriel; Chisholm, Julia; Brown, Rebecca

DOI: 10.1200/JCO.20.03060

License: Creative Commons: Attribution (CC BY)

Document Version Publisher's PDF, also known as Version of record

### Citation for published version (Harvard):

Shern, JF, Selfe, J, Izquierdo, E, Patidar, R, Chou, H-C, Song, YK, Yohe, ME, Sindiri, S, Wei, J, Wen, X, Rudzinski, ER, Barkauskas, DA, Lo, T, Hall, D, Linardic, CM, Hughes, D, Jamal, S, Jenney, M, Chisholm, J, Brown, R, Jones, K, Hicks, B, Angelini, P, George, SL, Chesler, L, Hubank, M, Kelsey, A, Gatz, SA, Skapek, SX, Hawkins, DS, Shipley, JM & Khan, J 2021, 'Genomic classification and clinical outcome in rhabdomyosarcoma: a report from an international consortium', *Journal of Clinical Oncology*, vol. 39, no. 26, pp. 2859-2871. https://doi.org/10.1200/JCO.20.03060

Link to publication on Research at Birmingham portal

### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Sq

tra

### **Genomic Classification and Clinical Outcome Rhabdomyosarcoma: A Report From an** International Consortium

Jack F. Shern, MD<sup>1,2</sup>; Joanna Selfe, PhD<sup>3</sup>; Elisa Izquierdo, MD<sup>4</sup>; Rajesh Patidar, MS<sup>1</sup>; Hsien-Chao Chou, PhD<sup>1</sup>; Young K. Song, PhD<sup>1</sup>; Marielle E. Yohe, MD, PhD<sup>2</sup>; Sivasish Sindiri, MS<sup>1</sup>; Jun Wei, PhD<sup>1</sup>; Xinyu Wen, MS<sup>1</sup>; Erin R. Rudzinski, MD<sup>5</sup>;

Donald A. Barkauskas, PhD<sup>6,7</sup>; Tammy Lo, MPH<sup>7</sup>; David Hall, MS<sup>7</sup>; Corinne M. Linardic, MD, PhD<sup>8</sup>; Debbie Hughes, PhD<sup>9</sup>;

Sabri Jamal, MSc<sup>4</sup>; Meriel Jenney, MD<sup>10</sup>; Julia Chisholm, MD<sup>11</sup>; Rebecca Brown, MD<sup>3,12</sup>; Kristine Jones, PhD<sup>13</sup>; Belynda Hicks, PhD<sup>13</sup>; Paola Angelini, MD<sup>11</sup>; Sally George, MD<sup>9,11</sup>; Louis Chesler, MD<sup>9</sup>; Michael Hubank, MD<sup>4</sup>; Anna Kelsey, MD<sup>14</sup>; Susanne A. Gatz, MD<sup>3,15</sup>; Stephen X. Skapek, MD<sup>16</sup>; Douglas S. Hawkins, MD<sup>17</sup>; Janet M. Shipley, PhD<sup>3</sup>; and Javed Khan, MD<sup>1</sup>

PURPOSE Rhabdomyosarcoma is the most common soft tissue sarcoma of childhood. Despite aggressive therapy, the 5-year survival rate for patients with metastatic or recurrent disease remains poor, and beyond PAX-FOX01 fusion status, no genomic markers are available for risk stratification. We present an international consortium study designed to determine the incidence of driver mutations and their association with clinical Ct outcome.

**PATIENTS AND METHODS** Tumor samples collected from patients enrolled on Children's Oncology Group trials (1998-2017) and UK patients enrolled on malignant mesenchymal tumor and RMS2005 (1995-2016) trials were subjected to custom-capture sequencing. Mutations, indels, gene deletions, and amplifications were identified, and survival analysis was performed.

RESULTS DNA from 641 patients was suitable for analyses. A median of one mutation was found per tumor. In FOXO1 fusion-negative cases, mutation of any RAS pathway member was found in > 50% of cases, and 21% had no putative driver mutation identified. BCOR (15%), NF1 (15%), and TP53 (13%) mutations were found at a higher incidence than previously reported and TP53 mutations were associated with worse outcomes in both fusion-negative and FOXO1 fusion-positive cases. Interestingly, mutations in RAS isoforms predominated in infants < 1 year (64% of cases). Mutation of MYOD1 was associated with histologic patterns beyond those previously described, older age, head and neck primary site, and a dismal survival. Finally, we provide a searchable companion database (ClinOmics), containing all genomic variants, and clinical annotation including survival data.

**CONCLUSION** This is the largest genomic characterization of clinically annotated rhabdomyosarcoma tumors to date and provides prognostic genetic features that refine risk stratification and will be incorporated into prospective trials.

### J Clin Oncol 39:2859-2871. © 2021 by American Society of Clinical Oncology

Licensed under the Creative Commons Attribution 4.0 License (C)

ASSOCIATED CONTENT See accompanying editorial on page 2851 Appendix

### **Data Supplement**

Author affiliations and support information (if applicable) appear at the end of this article.

Accented on May 7. 2021 and published at ascopubs.org/journal/ jco on June 24, 2021: DOI https://doi.org/10. 1200/JC0.20.03060

### **INTRODUCTION**

Rhabdomyosarcoma (RMS) is the most common soft tissue sarcoma of childhood.<sup>1</sup> With the development of multimodal chemotherapy regimens, relapse-free survival rates have improved to 70%-80% in patients with localized disease, albeit with significant toxicity.<sup>2</sup> Unfortunately, despite aggressive therapy, the 5-year survival rate for patients with metastatic disease remains poor, but variable.<sup>3</sup> Therapy assignment in North American and European trials is currently based on clinicopathologic features and not molecular or genetic markers, with the exception of the recent incorporation of *FOXO1* fusion status by the Children's Oncology Group (COG) and European paediatric Soft

tissue sarcoma Study Group (EpSSG).<sup>4-6</sup> Although clinical features reasonably stratify patients into broad treatment cohorts, prognostic imprecision hampers efforts to successfully escalate or de-escalate therapy. Particularly problematic is the COG intermediate risk category, defined as localized FOXO1 fusion-positive (FP) RMS and localized, incompletely resected (clinical group III) FOXO1 fusion-negative (FN) RMS arising from an unfavorable anatomic site; this category comprises approximately 50% of cases and has a heterogeneous clinical outcome.<sup>4,7,8</sup> This suggests that some of these children could be treated with less aggressive therapy or alternatively should be considered to have more aggressive disease.



### **Journal** of Clinical Oncology<sup>®</sup>

Volume 39. Issue 26 2859

Check for updates

Downloaded from ascopubs.org by 2.29.8.68 on September 17, 2021 from 002.029.008.068 Copyright © 2021 American Society of Clinical Oncology. All rights reserved.

### CONTEXT

### **Key Objective**

Rhabdomyosarcoma (RMS) is a sarcoma of childhood with a poor 5-year survival rate for patients with metastatic or recurrent disease. No genomic markers are currently available for risk stratification except for *PAX-FOXO1* fusion gene status. This study performed sequencing to determine the mutational status of genes implicated in RMS oncogenesis and correlated these results with clinical outcomes.

### **Knowledge Generated**

The genetic and clinical characteristics associated with primary tumors from two international cohorts are presented. Survival analysis demonstrated that *TP53* and *MYOD1* mutations were associated with worse event-free survival.

### Relevance

This study nominates mutant genes *MYOD1* and *TP53* as indicators of poor prognosis in fusion-negative RMS, and *TP53* alterations as a biomarker of more aggressive disease in fusion-positive RMS. Mutation of *MYOD1* was not restricted to spindle histology, and the association with adverse outcome highlights the need to accurately diagnose *MYOD1* mutations and develop novel treatment strategies for these patients.

Previous comprehensive genomic sequencing studies of RMS have been completed, but outcome analysis was limited by sample size or incomplete clinical annotation of the included samples.<sup>9,10</sup> To genetically classify RMS and refine risk stratification, we formed an international collaborative group and performed standardized sequencing of a large cohort of clinically annotated cases. Herein, we report the summary findings and detail the importance of incorporation of genomic data into prospectively enrolling RMS clinical trials. Adopting molecular features to RMS risk stratification should improve clinical outcomes for patients with RMS by allowing further tailoring of therapies to match an individual patient's risk and mutational profile. To ensure that this critical data set is available to the broader research community, the generated sequencing data are available within dbGAP (accession phs000720.v4.p1) and the clinical and mutational data are publicly accessible.<sup>11</sup>

### PATIENTS AND METHODS

### **Study Population and Clinical Annotation**

Samples from the two cohorts included in this work were collected on institutional review board–approved clinical trials or tissue banking studies. COG samples included samples collected on ARST0331, ARST0431, D9602, D9803, and D9902. UK samples from patients treated on protocols through the malignant mesenchymal tumor and RMS2005 protocols<sup>12-14</sup> were collected and approved for study through local and national ethical approvals (CCR2015 and 06/MRE04/71, respectively) (Appendix Table A1, online only). Because there are subtle differences in risk stratification between the COG and EpSSG (reviewed in Chen et al<sup>15</sup>), an overarching simplified risk stratification definition was used to enable merging data. Additional population and annotation details are provided in the Data Supplement (online only).

### **Gene Panel Sequencing**

A custom-capture sequencing assay targeting 39 genes previously implicated in RMS (Appendix Table A2, online only) was performed, and variants were called using previously published sequencing algorithms.<sup>9,16-19</sup> Detailed sequencing and variant calling methods are provided in the Data Supplement.

### Statistical Methods

Patient characteristics were summarized using medians and ranges or frequencies and percentages. Associations between pairs of gene markers were tested using the exact conditional test of proportions (Fisher's exact test). Eventfree survival (EFS) was defined as time from diagnosis (United Kingdom) or enrollment on a study (COG) until event (relapse, second malignant neoplasm, or death) or last contact. Each gene predictor variable was tested for univariate association with EFS and overall survival using the log-rank test and for association within a multivariate Cox proportional hazard regression. The survival analysis was done separately for each cohort of patients: COG and United Kingdom and within each cohort, separately for FP and FN patients. Detailed statistical methods are provided in the Data Supplement.

### RESULTS

### Patient Population

Clinically annotated cases from patient samples were assembled on COG biology study ARST14B1Q and a parallel cohort assembled from UK malignant mesenchymal tumor and RMS2005 studies, to generate two large cohorts of samples. The clinical details are summarized in Table 1. In total, 641 cases had adequate DNA to generate sequencing libraries of minimum quality to be included in the study. The median age of the combined cohort was 5.9

Characteristic	All (N = 641)	COG (n = 344)	UK (n = 297)
Sex, No. (%)			
Male	421 (66)	232 (66)	189 (66)
Female	220 (34)	112 (34)	108 (34)
Age at presentation, years			
Median	5.9	6.4	5.3
Range	0.02-37.8	0.02-37.8	0.1-23.1
Tumor histology, No. (%)			
Alveolar	151 (24)	68 (20)	83 (28)
Embryonal	447 (70)	254 (74)	187 (63)
Mixed alveolar and embryonal	3 (< 1)	2 (1)	1 (< 1)
Spindle cell RMS	18 (3)	18 (5)	7 (2)
NOS	20 (3)	2 (1)	18 (6)
Pleomorphic	2 (< 1)	0 (< 1)	1 (< 1)
Anatomic location, No. (%)			
Bladder/prostate	50 (8)	25 (7)	25 (7)
Extremity	92 (14)	52 (15)	40 (13)
Female GU	18 (3)	7 (2)	11 (4)
Head and neck	57 (9)	31 (9)	26 (9)
Orbital	45 (7)	25 (7)	20 (7)
Others	21 (3)	3 (1)	18 (6)
Parameningeal	127 (20)	63 (18)	64 (22)
Paratesticular	125 (20)	66 (19)	59 (20)
Peritoneum/trunk	101 (16)	72 (21)	29 (10)
Unknown	5 (< 1)	0 (0)	5 (2)
Risk group, No. (%)			
Low	220 (34)	124 (36)	96 (32)
Intermediate	299 (47)	147 (43)	152 (51)
High	115 (18)	73 (21)	42 (14)
Unknown	7 (1)	0 (0)	7 (2)
Variant calls, No. (%)			
Median	1	1	1
Range	0-5	0-5	0-4

 TABLE 1. Clinical Characteristics of Included Patients

Abbreviations: COG, Children's Oncology Group; GU, genitourinary; NOS, not otherwise specified; RMS, rhabdomyosarcoma.

years (range 0.02-37.8) (Appendix Fig A1, online only). The male:female ratio was 1.9:1, slightly higher than the generally accepted ratio of 1.5:1,<sup>20</sup> reflecting an enrichment of paratesticular tumors within this cohort. The most common anatomic locations were the parameningeal (20%) and paratesticular tumors (20%) followed by tumors of the retroperitoneum, peritoneum, or trunk (16%) and the extremity (14%). A simplified risk stratification algorithm was used to harmonize COG and UK cohorts (Data Supplement) and using this method, the population had representation

of cases of the low-risk (34%), intermediate-risk (47%), and high-risk populations (18%).

### **Mutation Frequency Observations**

Overall, the sequenced tumors had a median of one mutation call per tumor (range 0-5). The most frequently observed gene mutations are presented in Table 2. Consistent with prior reports, the genomic profiles of the FP and FN populations were distinct. The most frequently observed lesions in FP tumors were the focal amplification of CDK4 (13%) or MYCN (10%). The genes BCOR (6%), NF1 (4%), TP53 (4%), and PIK3CA (2%) were found in a small number of FP RMS cases, verifying previous observations.<sup>9</sup> In contrast, the most frequently observed genetic alteration in FN tumors were RAS isoform mutations NRAS (17%), KRAS (9%), and HRAS (8%), with any RAS isoform mutation noted in 32% (n = 167 of 515) of FN tumors. Mutation of an RAS pathway gene (defined as NRAS, KRAS, HRAS, FGFR4, NF1, and PIK3CA) could be found in 56% (n = 288 of 515) of all FN samples. Recurrence of mutations in tumor suppressor genes in FN RMS, TP53 (13%), NF1 (15%), and BCOR (15%), was higher than reports in previous studies.<sup>9</sup> Hotspot mutations in FGFR4 (13%), CTNNB1 (6%), PIK3CA (5%), and MYOD1 (3%) were observed at similar frequencies as previously reported and seen at similar percentages within the two independent international patient cohorts.9,10 No mutations were found in 14 genes previously associated with RMS (MTOR, PKN1, ALK, SOS1, SOS2, ROBO1, PDGFRA, GAB1, BRAF, CCND1, CCND2, ATM, AKT, and SMARCA4), although variants of unknown significance were observed in each of these genes. The median age of presentation of the patients correlated with alteration of individual genes, with a notable increase in MYOD1 mutations, CDK4 amplification, and MYCN amplification in patients older than 10 years and *HRAS* mutations in infants < 1 year (Fig 1A).

### Mutations Summarized by Anatomic Distribution

RMS tumors arise in diverse anatomic locations throughout the body, and the site of disease is known to correlate with clinical outcome.<sup>21</sup> Current clinical risk stratification assigns tumors arising in the orbit, nonparameningeal head and neck, and the male or female genital tracts as favorable. The distribution of mutations by anatomic location is presented in Figure 1A. As previously described,<sup>22</sup> DICER1 mutations had a predilection for tumors arising within the female genitourinary tract. Within the sequenced tumors, 33% (6 of 18) of all female genitourinary cases harbored a mutation in DICER1. Interestingly, FN cases of the extremity had an enrichment for TP53 mutations or MDM2 amplifications, whereby 42% (15 of 35) of FN cases with a primary tumor of the extremity had an alteration of one of these two genes (Appendix Table A3, online only). MYOD1mutant tumors also showed a distinct anatomic enrichment with 88% (15 of 17) of MYOD1-mutant tumors observed in either the head and neck or parameningeal region.

TABLE 2.	Mutation	Frequency ( Total Fi (N =	Dbservations I N Cohort 515)	by Cohort COG FN (n =	Cohort 275)	UK FN (n =	Cohort 240)		Total (N					
Gene	Total Cases	Total Mutant FN Cohort	Total FN (% of all FN cases)	COG Mutant FN Cohort	COG (% of FN cases)	UK Mutant FN Cohort	UK (% of FN cases)	Mutant <i>ALV</i> or <i>FOX01</i> +	Mutant <i>ALV</i> Fusion Unknown	Mutant F0X01+	<i>ALV</i> or <i>FOXO1</i> + (% of FP cases)			
NRAS	88	87	17	44	16	43	18	1	1	0	1			
BCOR	85	78	15	48	17	30	13	7	5	2	6			
NF1	80	75	15	40	15	35	15	5	1	4	4			
TP53	74	69	13	34	12	35	15	5	3	2	4			
FGFR4	65	65	13	28	10	37	15	0	0	0	0			
KRAS	45	44	9	32	12	12	5	1	1	0	1			
HRAS	44	41	8	21	8	20	8	3	2	1	2			
CTNNB1	32	32	6	17	6	15	6	0	0	0	0			
<i>РІКЗСА</i>	28	26	5	19	7	7	3	2	1	1	2			
MDM2	27	26	5	17	6	9	4	1	1	0	1			
CDKN2A	23	23	4	17	6	6	3	0	0	0	0			
FBXW7	18	18	3	13	5	5	2	0	0	0	0			
MYOD1	17	17	3	11	4	6	3	0	0	0	0			
CDK4	17	1	0	1	0	0	0	16	11	5	13			
MYCN	13	0	0	0	0	0	0	13	7	6	10			
DICER1	12	12	2	4	1	8	3	0	0	0	0			
ARID1A	11	11	2	7	3	4	2	0	0	0	0			
IGF1R	9	8	2	4	1	4	2	1	1	0	1			
PTEN	5	5	1	5	2	0	0	0	0	0	0			
MET	5	4	1	4	1	0	0	1	1	0	1			
FGFR1	4	4	1	3	1	1	0	0	0	0	0			
PTPN11	3	3	1	1	0	2	1	0	0	0	0			
PTCH1	3	3	1	2	1	1	0	0	0	0	0			
ERBB2	2	2	0	2	1	0	0	0	0	0	0			

Abbreviations: COG, Children's Oncology Group; FN, fusion-negative; FP, fusion-positive.

### FN Tumors Frequently Harbor More Than One Genetic Driver Alteration

Mutational heterogeneity has previously been described in FN RMS.<sup>23</sup> In this study, the presence of multiple driver mutations within individual tumors was evident within FN samples with 41% (213 of 515) of tumors having one mutation, 37% (193 of 515) of tumors having two or more mutations, and 21% (109 of 515) containing no alteration of a candidate gene (Appendix Fig A2A, online only). Greater than two mutations within a tumor was a significant marker in terms of worse EFS (P = .01, hazard ratio [HR] 2.014 [1.010-4.015]) within the COG cohort; however, this observation was not replicated in the UK cohort for EFS (P = .39, HR 1.098 [0.633-1.904]) (Appendix Fig A2B). To establish the most common pairings of gene interactions that drive RMS, we analyzed mutational data across the cohort and observed that mutations of tumor suppressor genes such as NF1, TP53, and BCOR frequently co-occurred

with other mutations. Significant interactions included *BCOR* with *NRAS* (P = .01) or *NF1* (P = .03), and *MYOD1* with *PIK3CA* (P < .0001) or *CDKN2A* (P = .0049) (Fig 2B, Appendix Fig A2C). *MYOD1* mutations were exceptional in that they always occurred with an additional gene being mutated. Although *NRAS*, *KRAS*, *HRAS*, and *FGFR4* were mutually exclusive in most cases, surprisingly, there were individual tumors with co-occurrence of multiple hotspot mutations in these genes.

### CDK4, MYCN, and TP53 in FP Tumors

Although *PAX3-FOXO1* fusion is itself of prognostic value, no molecular markers are currently available for risk substratification of FP tumors. In total, 126 FP tumors were evaluated (69 COG and 57 UK) with a dedicated fusion assay performed on 80 of 126 (64%) and centrally reviewed alveolar histology used as a proxy in the remaining cases. Small numbers of *CDK4-* and *MYCN-*mutant cases

A	Gene	No. of Cases		Age (median), years	Low (n = 220)	Intermediate (n = 299)	High (n = 115)	Bladder/prostate (n = 50)	Extremity (n = 92)	Female GU (n = 18)	Head and Neck (n = 57)	Orbital (n = 45)	Others $(n = 21)$	Parameningeal (n = 127)	Paratesticular (n = 125)	Peritoneum/trunk (n = 101)		
	NRAS	88		6.4	25	9	5	16	2	22	26	20	5	9	22	9		
	BCOR	85		6.7	20	11	7	12	3	11	16	27	10	12	18	14		
	NF1	80		5.1	11	14	10	22	3	11	7	13	19	14	13	16		
	TP53	74		4.2	11	12	11	8	16	17	19	18	10	9	2	16		
F	GFR4	65		4.7	11	11	6	16		11	9	18	5	17	6	12		
	KRAS	45		4.6	9	6	5	4	1		7	2	4.0	8	11	13		
	HRAS	44		2.8 4 0	8	7	4	14	3	11	2	2	10	2	12	9		
СТ	NNB1	32		4.3 5 1	0	5	3	10	2	6	4	4		2	/	6		
	IK3CA	20		5.1 6.6	5	4	4	2	2	6	2	2		4	6	3		
	NDNZ	23		0.0 7.6	3	4	4	-	3	U	5	7		6	Ū	6		
FI	RXW7	18		6.7	6	1	3	2			2	2		1	8	4		
M	YOD1	17	1	10.8	2	4	2		1		7			9				
	CDK4	17	1	1.3		2	10		12		2			2		3	i - 1	
л	мүсм	13	1	10.5		2	5		4		2		10	2		4		
D	ICER1	12		6.0	2	2	2			33			10			4		
A	RID1A	11		8.2	2	2		2			2	4		3	1	2		
				1%	5%	10%	Perc	renta	0 0 0	frasi	es wi	ith a	muta	tion	in tł	nat d	roun	
				170	070	1070	i cit	Jointa	ge o	1 003	0.5 00	in a		т	₽	iut g	oup	
B	NRAS	HRAS	KRAS	FGFR4	NF1	PIK3CA	FBXW7	MYOD1	TP53	MDM2	DICER1			CDNNZY	CTNNB	IGF1R	CDK4	MYCN
NR/	4 <i>S</i> 35	2	3	2	9	5	7					מ מ						
HR,	4 <i>S</i> 2	24	1	1	3		/	4	11	3	1	2	1	1	2	1		
KR/	45 3	1			-	1	/	4 1	11 2	3 1	1	2 6	1 ·	1	2 4	1		
FGF	-		24	2	4	1 5	1	4	11 2 2	3	1	2 2 6	1 : ; :	1 2 2	2 4 2	1		
Δ. A	R4 2	1	24 2	2 26	4 9	1 5 4	7 1 4	4	11 2 2 6	3 1 1	1		1 : ; : ; :	1 2 2 4	2 4 2 5	1		
/	R4 2 IF1 9	1 3	24 2 4	2 26 9	4 9 24	1 5 4 4	7 1 4 3	4 1 1	11 2 2 6 11	3 1 1 1	1		1 · · · · · · · · · · · · · · · · · · ·	1 2 2 4 6	2 4 2 5 6	1 1 1		
PIK30	R4 2 IF1 9 CA 5	1 3 1	24 2 4 5	2 26 9 4	4 9 24 4	1 5 4 4 5	7 1 4 3 2	4 1 1 1 9	11 2 2 6 11 1	3 1 1 1 1	1	2 6 6 7 1 1 3	1 : ;	1 2 2 4 6 6	2 4 2 5 6	1 1 1		
PIK30 FBXV	R4 2 IF1 9 CA 5 V7 7	1 3 1	24 2 4 5 1	2 26 9 4 4	4 9 24 4 3	1 5 4 5 2	7 1 4 3 2 2	4 1 1 1 9	11 2 2 6 11 1 1	3 1 1 1 1	1			1 2 2 2 4 4 5 6 6 7 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	2 4 2 5 6 2	1 1 1		
PIK30 FBXV MYO	R4         2           IF1         9           CA         5           V7         7           D1         4	1 3 1 1	24 2 4 5 1	2 26 9 4 4	4 9 24 4 3 1	1 5 4 5 2 9	7 1 4 3 2 2	4 1 1 9 9	11 2 2 6 11 1 1	3 1 1 1 1 1 3				1 2 2 2 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4	2 4 2 5 6 2 2	1 1 1	1	
PIK30 FBXV MYO TP	R4     2       IF1     9       CA     5       V7     7       D1     4       253     11	1 3 1 1 2	24 2 4 5 1	2 26 9 4 4 6	4 9 24 4 3 1 11	1 5 4 5 2 9 1	7 1 4 3 2 2 1	4 1 1 9 0	11 2 2 6 11 1 1 30	3 1 1 1 1 1 3 4				1 2 2 4 4 6 6 6 1 1 4	2 4 2 5 6 2 2 1	1 1 1 1 1 1 1 1	1	1
PIK30 FBXV MYO. TP MDN	<ul> <li>R4 2</li> <li>IF1 9</li> <li>CA 5</li> <li>VV7 7</li> <li>D1 4</li> <li>V53 11</li> <li>M2 3</li> <li>R1 6</li> </ul>	1 3 1 1 2 1	24 2 4 5 1 2	2 26 9 4 4 6 1	4 9 24 4 3 1 11 11	1 5 4 5 2 9 1	7 1 4 3 2 2 1 1	4 1 1 9 0 3	111 2 2 6 111 1 1 1 300 4	3 1 1 1 1 3 4 11			1	1 2 4 6 6 1 1 4	2 4 2 5 6 2 1	1 1 1 1 1 1 1 1	1 1 2	1
PIK30 FBXV MYO TP MDM DICE	<ul> <li>R4 2</li> <li>IF1 9</li> <li>CA 5</li> <li>V7 7</li> <li>D1 4</li> <li>I1</li> <li>M2 3</li> <li>R1 1</li> <li>DR 5</li> </ul>	1 3 1 1 2 1	24 2 4 5 1 2	2 9 4 4 6 1	4 9 24 4 3 1 11 1 1	1 5 4 5 2 9 1	7 1 4 3 2 2 2 1 1 1	4 1 9 0 3	111 2 2 6 111 1 1 1 300 4 4	3 3 1 1 1 1 1 1 3 4 4 11 2 2			1	1 2 2 4 4 5 6 1 1 4 4 2 4	2 4 5 6 2 1 1	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1 1 2	
PIK30 FBXV MYO. TP MDM DICE BCC	R4         2           IF1         9           CA         5           D1         4           M2         3           R1         1           DR         21	1 3 1 1 2 1 1 6	24 2 4 5 1 2 2 6	2 9 4 4 6 1 7	4 9 24 4 3 1 11 1 1 1 8 6	1 5 4 5 2 9 1 1 3	1 1 4 3 2 2 2 1 1 1 4 4	4 1 9 0 3 3	111 2 2 6 111 1 1 1 300 4 4 4 6	3 3 1 1 1 1 3 4 4 11 2 2		2 6 6 7 1 1 3 3 4 4 1 1 5 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2		1 2 2 4 4 6 6 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7	2 4 2 5 6 2 2 1 1 1 6	1 1 1 1 1 1 1 1 1 1	1 1 2 2	1
PIK30 FBXV MYO TP MDM DICE BCC CDKN.	R4     2       IF1     9       CA     5       V7     7       D1     4       753     11       M2     3       R1     1       DR     21       2A     1       B1     2	1 3 1 1 2 1 1 6 2	24 2 4 5 1 2 2 6 2 2	2 9 4 4 6 1 7 4	4 9 24 4 3 1 11 1 1 1 8 6	1 5 4 5 2 9 1 1 3 6	1 1 4 3 2 2 2 1 1 1 1 4 1 2	4 1 9 9 0 3 3 1 4	111 2 2 6 111 1 1 1 300 4 4 6	3 1 1 1 1 3 4 11 2 2		2 6 6 7 1 3 3 4 4 1 1 6 8 2 2 2 2 2 2 2 2 4 4		1 2 2 4 4 5 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7	2 4 2 5 6 2 1 1 6	1 1 1 1 1 1 1 1 1	1 1 2 2	
PIK30 FBXV MYO. TP MDM DICE BCC CDKN. CTNN	R4     2       VF1     9       CA     5       V7     7       D1     4       V53     11       M2     3       R1     1       DR     21       B1     2       1B1     2	1 3 1 1 2 1 1 6 2 4	24 2 4 5 1 2 2 6 6 2 2 2	2 26 9 4 4 4 6 1 1 7 7 4 5	4 9 24 4 3 1 11 1 1 1 1 8 6 6	1 5 4 5 2 9 1 1 3 6	1 1 4 3 2 2 2 1 1 1 1 4 1 2	4 1 9 0 3 3 1 4	111 2 6 111 1 1 1 300 4 4 6 1 1	3 1 1 1 1 1 1 1 3 4 111 2 2 1 1 1 1 1 1 1 1 1 1 1 1 1		2 6 6 7 7 1 3 3 4 4 1 1 5 2 2 2 2 2 2 2 2 4 6 6 6 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7		1 2 2 4 6 6 6 6 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7	2 4 2 5 6 2 1 1 1 6 1 1 6	1 1 1 1 1 1 1 1 1 1		1
PIK30 FBXV MYO. TP DICE BCC CDKN. CTNN IGF	R4     2       IF1     9       CA     5       V7     7       D1     4       IT     3       R1     1       OR     21       2A     1       B1     2       1R     1	1 3 1 1 2 1 6 2 4	24 2 4 5 1 2 2 6 6 2 2 2 1	2 26 9 4 4 4 6 1 7 7 4 5	4 9 24 4 3 1 11 1 1 1 1 8 6 6 1	1 5 4 2 9 1 1 3 3 6	1 1 4 3 2 2 2 1 1 1 1 4 1 2	4 1 9 0 3 3 1 4 4	111 2 6 111 1 1 1 30 4 4 6 1 1 1	3 1 1 1 1 1 3 4 11 2 2 2 1 1		2 6 6 7 1 1 3 4 4 1 2 2 2 2 4 4 6 1 1 1 1 1 1 1 1 1 1 1 1 1		1 2 4 4 5 4 5	2 4 2 5 6 2 1 1 6 1 1 6	1 1 1 1 1 1 1 1 1 3		
PIK30 FBXV MYO. TP MDM DICE BCC CDKN. CTNN IGF CD	R4         2           IF1         9           CA         5           V7         7           D1         4           IS53         11           M2         3           R1         1           DOR         21           2A         1           B1         2           1R         1           CN         CN	1 3 1 2 1 6 2 4	24 2 4 5 1 2 2 2 6 6 2 2 2 1	2 26 9 4 4 4 6 1 1 7 7 4 5 5	4 9 24 4 3 1 11 1 1 1 1 1 8 6 6 1	1 5 4 9 9 1 1 3 3 6	1 1 4 3 2 2 2 1 1 1 1 1 4 1 2	4 1 9 0 3 3 1 4 1 4	111 2 2 6 111 1 1 1 300 4 4 6 1 1 1 1 1	3 1 1 1 1 1 1 1 1 1 1 1 1 1		2 6 6 7 1 3 3 4 4 1 1 5 2 2 6 2 2 2 6 6 6 1 1 2 2 1 1 2 2 1 1 1 1	1	1 2 2 4 4 5 1 4 4 5 5	2 4 2 5 6 2 1 1 6 1 1 6	1 1 1 1 1 1 1 1 1 1 3 1	1 1 2 2 1 1 10	
PIK30 FBXV MYO. TP MDN DICE BCC CDKN. CTNN IGF CD MYC	R4     2       IF1     9       CA     5       V7     7       D1     4       M2     3       R1     1       DDR     21       2A     1       B1     2       1R     1       K4     1       CN     1	1 3 1 2 1 2 1 4 6 2 4	24 2 4 5 1 2 2 2 6 6 2 2 2 1	2 26 9 4 4 6 1 7 7 4 5 5	4 9 24 4 3 1 11 1 1 1 1 1 1 8 6 6 1 1	1 5 4 2 9 1 1 3 3 6	1 4 3 2 2 2 1 1 1 1 4 1 2 2 1 1 1 1 2	4 1 9 0 3 3 1 4 1 4 1	111         2         6         11         1         1         300         4         6         11         1         1         1         1         1         1         1         1         1         1         1         1         1	3 3 1 1 1 1 3 4 11 2 2 2 1 1 2 2 1 1 2 2 1 1 2 2 1 1 1 1 1 1 1 1 1 1 1 1 1		2 6 7 11 3 3 4 1 1 5 2 2 6 6 6 1 1 2 2 1 1 1 1 1 1 1 1 1 1 1		1 2 2 4 4 5 6 6 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7 7	2 4 2 5 6 2 2 1 1 6 1 0	1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1 1 2 2 1 1 10	

**FIG 1.** Mutations summarized by anatomic location and co-occurrence. (A) Summary of mutations by occurrence within defined risk groups (low, intermediate, or high) or by anatomic location. The reported value is a percentage of the number of cases with a mutation in that gene within each group. (B) Co-occurrence of mutated genes reported as absolute number of cases. GU, genitourinary.

Downloaded from ascopubs.org by 2.29.8.68 on September 17, 2021 from 002.029.008.068 Copyright © 2021 American Society of Clinical Oncology. All rights reserved.



**FIG 2.** *RAS* isoform mutations. (A) Oncoprint of mutations observed in infants < 1 year old (n = 25) showed an enrichment for RAS mutations. (B) Distribution of RAS isoform mutations by age with a distinct peak of *HRAS* mutant cases discovered in the infant population. (C) *HRAS* mutations were frequently found to have a higher VAF indicating that the mutation occurred before a loss-of-heterozygosity event on chromosome 11p. VAF, variant allele frequency.

(COG: n = 14 and n = 10; UK: n = 2 and n = 3, respectively) were observed between the two cohorts, limiting conclusions about the prognostic significance of these genes (Appendix Fig A3A, online only). Interestingly, a small number of FP cases (COG: n = 3; UK: n = 2) had mutations in *TP53* and were universally fatal (Appendix Fig A3B).

### Survival Analysis of RAS Isoforms and Enrichment of RAS Mutations in Infants

A driving hypothesis of this study was that the presence of a mutation in a RAS isoform or RAS pathway gene would

correlate with poor outcomes in FN RMS, because of the observation in previous smaller cohorts of enrichment of RAS isoform mutations in high-risk cases.<sup>10</sup> Therefore, we examined each RAS isoform and individual RAS pathway members for correlation with survival. Neither mutation of any RAS isoform nor a RAS pathway gene was associated with a worse EFS or overall survival across the two cohorts (Appendix Fig A4, online only). One striking observation was the correlation of RAS isoform mutations longitudinally with age. Most notable was the enrichment of RAS isoform mutations in FN cases that occurred under the age of 1 year

(64% of cases, P = .0068) with a clear peak incidence of HRAS mutations (40% of all FN infants, P < .0001) within this age group and no enrichment of secondary mutations (Fig 2A). Although not significant, KRAS-mutant tumors were frequent within the toddler period (15% of cases at 3 years, P = .2368) and a peak of NRAS-mutant tumors was observed in adolescence (30% of all cases at 13 years, P = .4407) (Fig 2B). NRAS, HRAS, and KRAS isoforms had distinct codon and amino acid profiles, consistent with previous studies (Appendix Fig A5, online only). Interrogation of the observed allele frequency of RAS isoform mutations showed that the majority of HRAS mutations occur at variant allele frequency > 0.5, likely reflecting the occurrence of this mutation within the frequently observed uniparental disomy event that occurs on chromosome 11p (Fig 2C).<sup>24</sup>

### Association of *TP53* Alterations With Survival in FN Tumors

TP53 was found to be altered in 13% (n = 69 of 515) of the FN cohort, and the observed lesions included deep deletions, truncating mutations, and point mutations (Appendix Fig A6A, online only). The mutations occurred throughout the gene body with some enrichment seen within the DNAbinding domain of the protein (Appendix Fig A6B). The most recurrent mutations were found at the codons G245S (six cases), R248Q or W (six cases), R175H (four cases), and P72A (four cases). Given the lack of a matched normal sample, no determination was made if these lesions represent somatic or germline events. Univariate (EFS P = .0083; HR 2.067 [1.192-3.585]) and risk-stratified analysis (EFS P = .0146; HR 1.973 [1.132-3.438]) of survival data within the COG cohort demonstrated that the presence of a TP53 mutation imparted a worse EFS (Figs 3A and 3B). Evaluation of the UK cohort verified the significance of this observation in both non-risk-stratified (EFS P = .0079; HR 2.006 [1.187-3.390]) and risk-stratified (EFS P = .0055; HR 2.105 [1.230-3.604]) analysis (Figs 3C and 3D).

### Association of *MYOD1* Mutations With Survival in FN Tumors

Mutations in the transcription factor *MYOD1* were found in 3% (n = 17 of 515) of all FN cases and no FP cases (Fig 4A). The observed mutations were confined to the previously reported hotspot codon change L122R.<sup>25</sup> As noted, *MYOD1*-mutant tumors within this pediatric cohort occurred at an older mean age of 10.8 years (2.1-21.1 years) when compared with the rest of the cohort. Centrally reviewed pathology from COG and review of UK samples frequently noted spindle or sclerosing features of the tumor (Figs 4B1 and 4B2); however, interestingly, cases with densely packed cells that mimicked the dense pattern of embryonal rhabdomyosarcoma (ERMS) or RMS not otherwise specified were also found (Figs 4B3 and 4B4). EFS of those patients with *MYOD1* mutations within the COG cohort was dismal and associated with rapid progression in non-risk-stratified (EFS P < .0001; HR 6.839 [3.463-13.507]) and risk-stratified (EFS P < .0001; HR 5.579 [2.791-11.151]) analysis (Figs 4C and 4D). Parallel survival analysis of the UK cohort verified this observation in nonrisk-stratified (EFS P = .0133; HR 3.320 [1.212-9.099]) and risk-stratified (EFS P = .0111; HR 3.455 [1.247-9.571]) analysis (Figs 4E and 4F). Importantly, within the MYOD1-mutant group, 23% of patients were identified before treatment as low-risk and 65% were identified as intermediate-risk. Consistent with previous reports.<sup>26</sup> 53% of cases had a corresponding alteration of PIK3CA and MYOD1 mutations were not found to be mutually exclusive with RAS mutations, with coexisting lesions seen in NRAS (n = 4), HRAS (n = 1), and NF1 (n = 1). Interestingly, MYOD1-mutant tumors also frequently harbored deep deletions in CDKN2A (n = 4 of 17, 24%). Deep deletions or deleterious mutations of CDKN2A were present in 4% of all FN tumors and associated with a worse EFS (COG: *P* = .0031, HR 2.737 [1.363 to 5.494]; UK: *P* = .0031, HR 4.7 [1.896 to 11.648]) (Appendix Fig A7A, online only). This observation was independent of the co-occurrence of MYOD1 mutations (Appendix Fig A7B), however was specific for cases within the intermediate-risk group (Appendix Fig A7C).

### DISCUSSION

Integration of molecular features into risk stratification and therapeutic decision making remains a major challenge to improving the care of any patient with a rare tumor. Decades of clinical trials led to the development of a complicated system for risk stratification of patients with RMS, on the basis of information from both the pretreatment tumor staging and surgical grouping.<sup>4</sup> The imprecision of these assignments is known, which has important implications for how current therapy is delivered and how clinical trials incorporating novel agents are designed. Our international collaboration generated the largest cohort of clinically annotated and genomically characterized RMS tumors analyzed to date. The effort discovered critical genetic insights into the underpinnings of the disease and significant molecular markers that provide refinement to the current risk stratification of patients with RMS. On the basis of our results, we propose a new framework for the classification and treatment of RMS, using TP53 and MYOD1 mutations in addition to the FOXO1 fusion status, 5-7 which could be tested in prospective clinical trials (Appendix Table A4, online only).

The overall gene mutation frequency that was observed is consistent with previous sequencing studies<sup>9,27</sup> with the notable exception of an increased frequency of tumor suppressor genes *TP53*, *NF1*, and *BCOR*. The observed increase in frequency of mutation of these genes likely results from improved depth of sequencing using a targeted assay approach. Of interest are the approximately 20% of



**FIG 3.** *TP53* mutations and association with survival. (A) KM analysis of EFS within the COG FN cohort (n = 275) by the presence of a *TP53* Mut or absence of a *TP53* lesion (*TP53* WT). (B) KM analysis of EFS within the COG FN cohort by *TP53* status and RMS risk category. Total case numbers: low, n = 124; intermediate, n = 126; high, n = 35. (C) KM analysis of EFS within the UK FN cohort (n = 240) by the presence of a *TP53* Mut or absence of a *TP53* lesion (*TP53* WT). (D) KM analysis of EFS within the UK FN cohort by *TP53* status and RMS risk category. Total case numbers: low, n = 96; intermediate, n = 113; high, n = 25. Presented *P* values are log-rank and BH adj. HR with 95% Cl. BH adj, Benjamini-Hochberg-adjusted; COG, Children's Oncology Group; EFS, event-free survival; FN, fusion-negative; HR, hazard ratio; KM, Kaplan-Meier; RMS, rhabdo-myosarcoma; *TP53* Mut, *TP53* mutation; *TP53* WT, *TP53* wild type.

Genomic Classification and Outcome in Rhabdomyosarcoma



FIG 4. MYOD1 mutations and survival. (A) All tumors with a mutation in MYOD1 within the entire sequenced cohort (N = 641) are shown to visualize co-occurrence of MYDO1 mutations with other genes. All MYOD1 cases were observed (continued on following page)

**FIG 4.** (Continued). within FN cases. (B) Histology of *MYOD1* mutant cases demonstrated the frequent presence of sclerosing or spindle morphology (case 1 and case 2). Cases 3 and 4 show that some cases demonstrated areas more typical of dense embryonal histology. (C) KM analysis of EFS within the COG FN cohort (n = 275) by the presence of a *MYOD1* Mut or absence of a *MYOD1* lesion (*MYOD1* WT). (D) KM analysis of EFS within the COG FN cohort by *MYOD1* status and RMS risk category. Total case numbers: low, n = 124; intermediate, n = 126; high, n = 35. (E) KM analysis of EFS within the UK FN cohort (n = 240) by the presence of a *MYOD1* Mut or absence of a *MYOD1* Mut or absence of a *MYDO1* lesion (*MYOD1* WT). (F) KM analysis of EFS within the UK FN cohort by *MYOD1* status and RMS risk category. Total case numbers: low, n = 96; intermediate, n = 113; high, n = 25. Presented *P* values are log-rank and BH adj. HR with 95% CI. BH adj, Benjamini-Hochberg-adjusted; COG, Children's Oncology Group; EFS, event-free survival; FN, fusion-negative; HR, hazard ratio; KM, Kaplan-Meier; *MYOD1* Mut, *MYOD1* mutation; *MYOD1* WT, *MYOD1* wild type; RMS, rhabdomyosarcoma.

FN tumors that have no driver mutation of a candidate gene. Beyond genome-wide aneuploidy and focal loss of heterozygosity of 11p15, the previous comprehensive analyses,<sup>9,10,23,28</sup> and this focused analysis, have failed to discover recurrent genetic driver genes in this group of tumors. This suggests the need for continued comprehensive genomic and epigenetic evaluations that would allow identification of genes mutated at a low recurrence frequency, as well as alternative mechanisms of oncogenesis. The discovery that FN tumors are frequently driven by alteration of multiple coexisting mutations mirrors previous work that showed FN tumors are composed of multiple subclones that follow an evolutionary selection.<sup>23</sup> This finding suggests that clonal evolution of these tumors may be significant, and elegant genomic work has highlighted how these processes might drive relapsed or refractory disease.<sup>10</sup> Although the trend toward a worse outcome in patients with multiple mutations in the COG cohort is intriguing, this observation was not replicated in the UK cohort and may be confounded by the different therapeutic regimens that the patients received. Comprehensive, prospective assessment is required to address the validity of the survival correlations. In addition, studies designed to assay sequential tumor biopsies will be required to fully interrogate the mechanisms of metastasis and relapse in RMS.

Mutations in RAS isoforms have long been described as a driver of FN RMS.<sup>29</sup> Our study clearly determines that the presence at diagnosis of a mutation in an RAS isoform or RAS pathway gene does not portend a poor prognosis, in contrast to previous smaller cohorts finding enrichment for RAS isoform mutations in high-risk cases.<sup>10</sup> Although RAS was not found to be a prognostic predictor, we highlight an interesting observation that RAS isoform mutations appear to have some age-specific correlations, with HRAS occurring in the infants, KRAS occurring in the toddlers, and NRAS mutations with a peak in adolescence. Infants have previously been shown to have an inferior 5-year failure-free survival as compared with older patients (67% v 81%).<sup>30</sup> This difference is attributed to the general reluctance to use more aggressive local control, including radiation, in these patients.<sup>31</sup> Our results indicate that incorporation of targeted therapeutic agents such as tipifarnib (NCT04284774) or AMG510 (NCT03600883) may be particularly beneficial to this vulnerable and high-risk population. In addition,

further mechanistic surveys of the developmental biology underlying these observations might have important implications for the generation of accurate preclinical models of RMS.

No molecular markers are currently used for risk stratification of FP RMS tumors. Amplification of the chromosomal regions 2p24 and 12q13-q15 and the implicated genes MYCN and CDK4, respectively, are the most recurrent lesions associated with a FOXO1 fusion. Previous work dedicated to assigning the prognostic value of these lesions identified amplification of 12q13-q14, but not 2p24, as a marker of an aggressive subset of FP tumors.<sup>32</sup> Other efforts discovered that in ARMS, overexpression or gain of genomic copies of MYCN was significantly associated with adverse outcome.<sup>33</sup> The current study found inconsistent results for MYCN and CDK4 amplification, with nonreproducible correlations noted between the two cohorts. There is evidence of a small subset of FP tumors that harbor a mutation of *TP53* at diagnosis and appear to be particularly aggressive. Ultimately, prospective consortium-level trials should include profiling of each of these genes to define their prognostic value and the biologic role they play in FP RMS.

From the seminal report of Li-Fraumeni syndrome,<sup>34</sup> the role of TP53 in ERMS oncogenesis has long been established; however, the association of TP53 mutations with clinical outcome has previously been unknown. Given the lack of a corresponding germline sample, our study could not determine whether the discovered TP53 mutation was germline or somatic. Despite this, we demonstrated that the presence of a TP53 mutation was predictive of a worse outcome. This finding is consistent with reports from several cancer types that found mutation of TP53 is associated with poor response and survival.<sup>35</sup> This is also consistent with higher levels of TP53 protein in metastatic versus localized ERMS<sup>36</sup> and also observations in zebra fish models of ERMS, where tp53 mutations are linked to more aggressive and metastatic disease.<sup>37</sup> Determination of TP53 status in all cases of RMS therefore is critical, both for prognostic value and the implications that germline mutations have for genetic counseling.

*MYOD1* mutation of the L122R codon was reported by independent groups in 2014.<sup>25,38</sup> *MYOD1*-mutant tumors make up only 3% of FN RMS, and this study highlights the importance of *MYOD1* mutations within the RMS

population. These tumors have unique demographic, anatomic, and histologic characteristics, but none of these appear to definitively capture all *MYOD1*-mutant tumors. This suggests the need to incorporate sequencing of this gene into the diagnostic workup of FN RMS. Our observation that *MYOD1* mutations invariably co-occur with mutation in a second gene, most notably *PIK3CA* and *CDKN2A*, is consistent with a recent report.<sup>26</sup> The co-occurrence with *CDKN2A* is of interest given that a recent large survey of soft tissue sarcomas of multiple histologies, including a small number of RMS

### **AFFILIATIONS**

<sup>1</sup>Genetics Branch, Oncogenomics Section, Center for Cancer Research, National Institutes of Health, Bethesda, MD

<sup>2</sup>Pediatric Oncology Branch, Center for Cancer Research, National Institutes of Health, Bethesda, MD

<sup>3</sup>Sarcoma Molecular Pathology Team, Divisions of Molecular Pathology and Cancer Therapeutics, The Institute of Cancer Research, London, United Kingdom

<sup>4</sup>Molecular Diagnostics Department, The Institute of Cancer Research and Clinical Genomics, The Royal Marsden NHS Foundation, London, United Kingdom

<sup>5</sup>Department of Laboratories, Seattle Children's Hospital, University of Washington, Seattle, WA

<sup>6</sup>Department of Preventive Medicine, Keck School of Medicine of the University of Southern California, Los Angeles, CA

<sup>7</sup>Children's Oncology Group, Monrovia, CA

<sup>8</sup>Duke University School of Medicine, Durham, NC

<sup>9</sup>Paediatric Tumour Biology, Division of Clinical Studies, The Institute of Cancer Research, London, United Kingdom

<sup>10</sup>Cardiff and Vale UHB, Paeds Oncology, Cardiff, United Kingdom

<sup>11</sup>Children and Young People's Unit, Royal Marsden NHS Foundation Trust, London, United Kingdom

<sup>12</sup>Department of Pathology, Aberdeen Royal Infirmary, Aberdeen, United Kingdom

<sup>13</sup>Cancer Genomics Research Laboratory, Leidos Biomedical Research, Frederick National Laboratory for Cancer Research, Frederick, MD

<sup>14</sup>Department of Paediatric Histopathology, Manchester University NHS Foundation Trust Royal Manchester Childrens Hospital, Manchester, United Kingdom

<sup>15</sup>Cancer Research UK Clinical Trials Unit, Institute of Cancer and Genomic Sciences, University of Birmingham, Birmingham, United Kingdom

<sup>16</sup>Division of Hematology/Oncology, Department of Pediatrics, University of Texas Southwestern Medical Center, Dallas, TX

<sup>17</sup>Department of Pediatrics, Seattle Children's Hospital, Fred Hutchinson Cancer Research Center, University of Washington, Seattle, WA

### **CORRESPONDING AUTHOR**

Jack F. Shern, MD, Pediatric Oncology Branch, National Cancer Institute, Building 10, Room 1W-3750, 9000 Rockville Pike, Bethesda, MD 20892; e-mail: john.shern@nih.gov.

### **STUDY GROUPS**

International Consortium: Children's Oncology Group; Children's Cancer and Leukaemia Group, Young Onset Sarcoma Subgroup; and the National Cancer Institute. tumors, implicated *CDKN2A* as a biomarker of poor prognosis.<sup>39</sup> Although our study indicates that *CDKN2A* alterations may have prognostic value independent of *MYOD1*, this conclusion is based on low overall numbers. Therefore, we recommend that *CDKN2A* alterations be evaluated in prospective studies. Regardless, *MYOD1*mutant tumors have an aggressive nature and have limited responses to current therapeutic regimens, which highlights the impetus to identify these cases and develop novel therapeutic trials for this rare subset of patients with RMS.

#### DISCLAIMER

The content of this publication does not necessarily reflect the views or policies of the Department of Health and Human Services, nor does mention of trade names, commercial products, or organizations imply endorsement of the US government.

### **EQUAL CONTRIBUTION**

 $\sf J.F.S.$  and  $\sf J.S.$  contributed equally to this work.  $\sf J.M.S.$  and  $\sf J.K.$  are equal senior authors.

### **PRIOR PRESENTATION**

Presented at the Pediatric Oncology Session of the ASCO annual meeting, June 3, 2018; the plenary session of the Children's Oncology Group annual meetings on October 3, 2018 and March 19, 2020; and the plenary session of the EpSSG Winter meeting, December 6, 2018.

### **SUPPORT**

Supported by the St Baldrick's Hero Fund in memory of Peyton Arens and by Grants No. U10CA098543, U10CA098413, U10CA180899, and U10CA180886 from the National Cancer Institute, Bethesda, MD, to the Children's Oncology Group. United Kingdom funding was provided by the Chris Lucas Trust, the Talan's Trust, Alice's Arc charity, and Cancer Research UK (Grant No. C5066/A10399).

### AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Disclosures provided by the authors are available with this article at DOI https://doi.org/10.1200/JC0.20.03060.

### AUTHOR CONTRIBUTIONS

**Conception and design:** Jack F. Shern, Susanne A. Gatz, Douglas S. Hawkins, Janet M. Shipley, Javed Khan

Financial support: Jack F. Shern, Janet M. Shipley, Javed Khan Administrative support: Jack F. Shern, Javed Khan

Provision of study materials or patients: Xinyu Wen, Paola Angelini, Louis Chesler, Anna Kelsev

**Collection and assembly of data:** Jack F. Shern, Joanna Selfe, Young K. Song, Sivasish Sindiri, Jun Wei, Xinyu Wen, Erin R. Rudzinski, Tammy Lo, David Hall, Corinne M. Linardic, Debbie Hughes, Julia Chisholm, Rebecca Brown, Kristine Jones, Belynda Hicks, Paola Angelini, Sally George, Louis Chesler, Michael Hubank, Anna Kelsey, Douglas S. Hawkins, Janet M. Shipley, Javed Khan

**Data analysis and interpretation:** Jack F. Shern, Joanna Selfe, Elisa Izquierdo, Rajesh Patidar, Hsien-Chao Chou, Marielle E. Yohe, Sivasish Sindiri, Jun Wei, Xinyu Wen, Erin R. Rudzinski, Donald A. Barkauskas, David Hall, Corinne M. Linardic, Sabri Jamal, Meriel Jenney, Stephen X. Skapek, Douglas S. Hawkins, Janet M. Shipley, Javed Khan Manuscript writing: All authors

Final approval of manuscript: All authors

Accountable for all aspects of the work: All authors

### ACKNOWLEDGMENT

The authors thank the core sequencing facility of the NCI Division of Cancer Epidemiology and Genetics. Processing of the data was made possible by the NIH Helix computer cluster. The authors also thank the Children's Cancer and Leukaemia Group for facilitating sample collection, the National Cancer Research Institute Young Onset Sarcoma Subgroup and EpSSG for endorsing this study, and the Clinical Trials Unit in Birmingham, UK, for providing clinical trial data.

### REFERENCES

- 1. Skapek SX, Ferrari A, Gupta AA, et al: Rhabdomyosarcoma. Nat Rev Dis Primers 5:1, 2019
- Malempati S, Hawkins DS: Rhabdomyosarcoma: Review of the Children's Oncology Group (COG) Soft-Tissue Sarcoma Committee experience and rationale for current COG studies. Pediatr Blood Cancer 59:5-10, 2012
- 3. Oberlin O, Rey A, Lyden E, et al: Prognostic factors in metastatic rhabdomyosarcomas: Results of a pooled analysis from United States and European cooperative groups. J Clin Oncol 26:2384-2389, 2008
- Hibbitts E, Chi YY, Hawkins DS, et al: Refinement of risk stratification for childhood rhabdomyosarcoma using FOXO1 fusion status in addition to established clinical outcome predictors: A report from the Children's Oncology Group. Cancer Med 8:6437-6448, 2019
- Missiaglia E, Williamson D, Chisholm J, et al: PAX3/FOXO1 fusion gene status is the key prognostic molecular marker in rhabdomyosarcoma and significantly improves current risk stratification. J Clin Oncol 30:1670-1677, 2012
- Selfe J, Olmos D, Al-Saadi R, et al: Impact of fusion gene status versus histology on risk-stratification for rhabdomyosarcoma: Retrospective analyses of patients on UK trials. Pediatr Blood Cancer 64, 2017
- Skapek SX, Anderson J, Barr FG, et al: PAX-FOXO1 fusion status drives unfavorable outcome for children with rhabdomyosarcoma: A Children's Oncology Group report. Pediatr Blood Cancer 60:1411-1417, 2013
- Meza JL, Anderson J, Pappo AS, et al: Analysis of prognostic factors in patients with nonmetastatic rhabdomyosarcoma treated on intergroup rhabdomyosarcoma studies III and IV: The Children's Oncology Group. J Clin Oncol 24:3844-3851, 2006
- Shern JF, Chen L, Chmielecki J, et al: Comprehensive genomic analysis of rhabdomyosarcoma reveals a landscape of alterations affecting a common genetic axis in fusion-positive and fusion-negative tumors. Cancer Discov 4:216-231, 2014
- 10. Chen X, Stewart E, Shelat AA, et al: Targeting oxidative stress in embryonal rhabdomyosarcoma. Cancer Cell 24:710-724, 2013
- 11. ClinOmics Genomics Portal. https://clinomics.ccr.cancer.gov/clinomics/public/
- 12. Stevens MC, Rey A, Bouvet N, et al: Treatment of nonmetastatic rhabdomyosarcoma in childhood and adolescence: Third study of the International Society of Paediatric Oncology—SIOP Malignant Mesenchymal Tumor 89. J Clin Oncol 23:2618-2628, 2005
- Oberlin O, Rey A, Sanchez de Toledo J, et al: Randomized comparison of intensified six-drug versus standard three-drug chemotherapy for high-risk nonmetastatic rhabdomyosarcoma and other chemotherapy-sensitive childhood soft tissue sarcomas: Long-term results from the International Society of Pediatric Oncology MMT95 study. J Clin Oncol 30:2457-2465, 2012
- 14. Bisogno G, De Salvo GL, Bergeron C, et al: Vinorelbine and continuous low-dose cyclophosphamide as maintenance chemotherapy in patients with high-risk rhabdomyosarcoma (RMS 2005): A multicentre, open-label, randomised, phase 3 trial. Lancet Oncol 20:1566-1575, 2019
- 15. Chen C, Dorado Garcia H, Scheer M, et al: Current and future treatment strategies for rhabdomyosarcoma. Front Oncol 9:1458, 2019
- 16. Izquierdo E, Yuan L, George S, et al: Development of a targeted sequencing approach to identify prognostic, predictive and diagnostic markers in paediatric solid tumours. Oncotarget 8:112036-112050, 2017
- 17. George SL, Izquierdo E, Campbell J, et al: A tailored molecular profiling programme for children with cancer to identify clinically actionable genetic alterations. Eur J Cancer 121:224-235, 2019
- Chang W, Brohl AS, Patidar R, et al: MultiDimensional ClinOmics for precision therapy of children and adolescent young adults with relapsed and refractory cancer: A report from the center for cancer research. Clin Cancer Res 22:3810-3820, 2016
- 19. Talevich E, Shain AH, Botton T, et al: CNVkit: Genome-wide copy number detection and visualization from targeted DNA sequencing. PLoS Comput Biol 12: e1004873, 2016
- 20. Ognjanovic S, Linabery AM, Charbonneau B, et al: Trends in childhood rhabdomyosarcoma incidence and survival in the United States, 1975-2005. Cancer 115:4218-4226, 2009
- 21. Crist W, Gehan EA, Ragab AH, et al: The third intergroup rhabdomyosarcoma study. J Clin Oncol 13:610-630, 1995
- Dehner LP, Jarzembowski JA, Hill DA: Embryonal rhabdomyosarcoma of the uterine cervix: A report of 14 cases and a discussion of its unusual clinicopathological associations. Mod Pathol 25:602-614, 2012
- 23. Chen L, Shern JF, Wei JS, et al: Clonality and evolutionary history of rhabdomyosarcoma. PLoS Genet 11:e1005075, 2015
- 24. Robbins KM, Stabley DL, Holbrook J, et al: Paternal uniparental disomy with segmental loss of heterozygosity of chromosome 11 are hallmark characteristics of syndromic and sporadic embryonal rhabdomyosarcoma. Am J Med Genet A 170:3197-3206, 2016
- Kohsaka S, Shukla N, Ameur N, et al: A recurrent neomorphic mutation in MYOD1 defines a clinically aggressive subset of embryonal rhabdomyosarcoma associated with PI3K-AKT pathway mutations. Nat Genet 46:595-600, 2014
- Agaram NP, LaQuaglia MP, Alaggio R, et al: MYOD1-mutant spindle cell and sclerosing rhabdomyosarcoma: An aggressive subtype irrespective of age. A reappraisal for molecular classification and risk stratification. Mod Pathol 32:27-36, 2019
- 27. Seki M, Nishimura R, Yoshida K, et al: Integrated genetic and epigenetic analysis defines novel molecular subgroups in rhabdomyosarcoma. Nat Commun 6: 7557, 2015
- 28. Casey DL, Wexler LH, Pitter KL, et al: Genomic determinants of clinical outcomes in rhabdomyosarcoma. Clin Cancer Res 26:1135-1140, 2020
- Stratton MR, Fisher C, Gusterson BA, et al: Detection of point mutations in N-ras and K-ras genes of human embryonal rhabdomyosarcomas using oligonucleotide probes and the polymerase chain reaction. Cancer Res 49:6324-6327, 1989
- Malempati S, Rodeberg DA, Donaldson SS, et al: Rhabdomyosarcoma in infants younger than 1 year: A report from the Children's Oncology Group. Cancer 117: 3493-3501, 2011

- Joshi D, Anderson JR, Paidas C, et al: Age is an independent prognostic factor in rhabdomyosarcoma: A report from the soft tissue sarcoma Committee of the Children's Oncology Group. Pediatr Blood Cancer 42:64-73, 2004
- Barr FG, Duan F, Smith LM, et al: Genomic and clinical analyses of 2p24 and 12q13-q14 amplification in alveolar rhabdomyosarcoma: A report from the Children's Oncology Group. Genes Chromosomes Cancer 48:661-672, 2009
- Williamson D, Lu YJ, Gordon T, et al: Relationship between MYCN copy number and expression in rhabdomyosarcomas and correlation with adverse prognosis in the alveolar subtype. J Clin Oncol 23:880-888, 2005
- 34. Li FP, Fraumeni JF Jr: Rhabdomyosarcoma in children: Epidemiologic study and identification of a familial cancer syndrome. J Natl Cancer Inst 43:1365-1373, 1969
- Petitjean A, Achatz MI, Borresen-Dale AL, et al: TP53 mutations in human cancers: Functional selection and impact on cancer prognosis and outcomes. Oncogene 26:2157-2165, 2007
- 36. Leuschner I, Langhans I, Schmitz R, et al: p53 and mdm-2 expression in rhabdomyosarcoma of childhood and adolescence: Clinicopathologic study by the Kiel Pediatric Tumor Registry and the German Cooperative Soft Tissue Sarcoma Study. Pediatr Dev Pathol 6:128-136, 2003
- Ignatius MS, Hayes MN, Moore FE, et al: tp53 deficiency causes a wide tumor spectrum and increases embryonal rhabdomyosarcoma metastasis in zebrafish. Elife 7:e37202, 2018
- Szuhai K, de Jong D, Leung WY, et al: Transactivating mutation of the MYOD1 gene is a frequent event in adult spindle cell rhabdomyosarcoma. J Pathol 232: 300-307, 2014
- Bui NQ, Przybyl J, Trabucco SE, et al: A clinico-genomic analysis of soft tissue sarcoma patients reveals CDKN2A deletion as a biomarker for poor prognosis. Clin Sarcoma Res 9:12, 2019

### **ASCO**<sup><sup>•</sup></sup> Journals

### Gain Recognition as an ASCO Journal Reviewer

Participate as a journal reviewer and play an integral role in maintaining the quality and integrity of ASCO journals. Benefits and recognition for reviewing ASCO journals content include:

- Staying up to date on the latest oncology research
- Increased exposure to key leaders in oncology
- Career advancement opportunities
- ASCO members who become reviewers can earn FASCO points

Get started today at ascopubs.org/reviewers



#### **AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST**

#### Genomic Classification and Clinical Outcome in Rhabdomyosarcoma: A Report From an International Consortium

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated unless otherwise noted. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to www.asco.org/rwc or ascopubs.org/jco/authors/author-center.

Open Payments is a public database containing information reported by companies about payments made to US-licensed physicians (Open Payments).

Donald A. Barkauskas Employment: Genentech Stock and Other Ownership Interests: Genentech Patents, Royalties, Other Intellectual Property: US patent on the basis of PhD research in glioblastoma

Corinne M. Linardic

Leadership: Grid Therapeutics

Stock and Other Ownership Interests: Pfizer, LabCorp, Thermo Fisher Scientific Honoraria: AstraZeneca

Patents, Royalties, Other Intellectual Property: My spouse has a patent for his therapeutic antibody developed in his laboratory at Duke and licensed by Grid

Julia Chisholm

Consulting or Advisory Role: Roche, Bayer, Roche/Genentech

#### Belynda Hicks

Employment: United Therapeutics Stock and Other Ownership Interests: United Therapeutics Travel, Accommodations, Expenses: United Therapeutics

#### Michael Hubank

Honoraria: Boehringer Ingelheim, Incyte, Qiagen Consulting or Advisory Role: Guardant Health, Illumina, AstraZeneca, Bristol Myers Squibb Foundation, Bayer, Roche, Lilly, Amgen Anna Kelsey Honoraria: Bayer Travel, Accommodations, Expenses: Bayer

Susanne A. Gatz Consulting or Advisory Role: Tesaro, Bayer Travel, Accommodations, Expenses: AstraZeneca

### **Douglas S. Hawkins**

Research Funding: Loxo, Bristol Myers Squibb, Merck Sharp & Dohme, Bayer, Lilly, Eisai, Amgen, Seattle Genetics, Incyte, Jazz Pharmaceuticals Travel, Accommodations, Expenses: Loxo, Bayer, AstraZeneca

#### Javed Khan

Research Funding: Lentigen, Taiho Oncology

Patents, Royalties, Other Intellectual Property: Monoclonal antibody-based therapeutics targeting fibroblast growth factor receptor 4 (FGFR4) for potential treatment of human cancers expressing FGFR4

No other potential conflicts of interest were reported.

# APPENDIX



FIG A1. Histogram presentation of the age distribution of cases within the cohort (N = 641).

Shern et al



**FIG A2.** Survival analysis on the basis of number of called mutations. (A) All FN samples (n = 515) were distributed into groups having no observed mutations (black), one mutation (red), or two or more genes mutated (blue). (B) Kaplan-Meier analysis of EFS from the COG and UK FN cohorts on the basis of the number of observed mutations within a tumor. The COG cohort (n = 275) had 49 patients with no observed mutation, 113 patients with one observed mutation, and 113 patients with two or more observed mutations. The UK cohort (n = 240) had 60 patients with no observed mutation, 100 patients with one observed mutation, and 80 patients with two or more observed mutations. (C) Pairwise testing of the significance of gene-gene interactions was performed to identify interactions that occurred at higher (red and purple) or lower (green) frequencies than expected. COG, Children's Oncology Group; EFS, event-free survival; FN, fusion-negative; HR, hazard ratio.



**FIG A3.** *CDK4, MYCN*, and *TP53* mutations in FP RMS. (A) Survival analysis for all three genes within the COG and UK cohorts with and without risk stratification. The log-rank and BH adj *P* value are presented. (B) Kaplan-Meier analysis of EFS and OS in FP cases within the COG cohort (n = 69) and UK cohort (n = 57) for *TP53* Mut or WT. COG cohort has three *TP53* mutant cases and UK cohort has two *TP53* mutant cases. BH adj, Benjamini-Hochberg–adjusted; COG, Children's Oncology Group; EFS, event-free survival; FP, fusion-positive; HR, hazard ratio; Mut, mutated; OS, overall survival; RMS, rhabdomyosarcoma; WT, wild type.

Journal of Clinical Oncology

		COG Coho	rt EFS		UK Cohort EFS								
	P Values No	BH Adj No	P Values Risk	BH Adj Risk	P Values No	BH Adj No	<b>P</b> Values Risk	BH Adj Risk					
Biomarker	Stratification	Stratification	-Stratified	-Stratified	Stratification	Stratification	-Stratified	-Stratified					
FGFR4	.4032	0.5700	.6002	0.7804	.9249	0.9432	.9239	0.9239					
HRAS	.2072	0.4463	.1562	0.4860	.8993	0.9432	.8337	0.9239					
KRAS	.3743	0.5700	.6132	0.7804	.9432	0.9432	.9126	0.9239					
NF1	.6759	0.7570	.3211	0.5994	.0275	0.1719	.0432	0.1800					
NRAS	.0494	0.1537	.2987	0.5994	.1669	0.5216	.2659	0.6648					
PIK3CA	.0056	0.0392	.0243	0.1075	.8650	0.9432	.8973	0.9239					
RAS isoform mutation	.1565	0.3652	.3117	0.5994	.4842	0.7261	.6329	0.8328					
RAS pathway mutation	.1004	0.2736	.3244	0.5994	.3937	0.7030	.5965	0.8285					
		COG Coh	ort OS		UK Cohort OS								
	P Values No	BH Adj No	P Values Risk	BH Adj Risk	P Values No	BH Adj No	P Values Risk	BH Adj Risk					
Biomarker	Stratification	Stratification	-Stratified	-Stratified	Stratification	Stratification	-Stratified	-Stratified					
FGFR4	.9455	0.9455	.7423	0.8231	.6701	0.7615	.5605	0.8132					
HRAS	.1918	0.4882	.1287	0.4004	.6593	0.7615	.5263	0.8132					
KRAS	.6453	0.7827	.8291	0.8598	.8699	0.9220	.8700	0.9168					
NF1	.1789	0.4882	.0588	0.2058	.1819	0.7579	.2279	0.8132					
NRAS	.2767	0.6456	.6996	0.8231	.1802	0.7579	.2855	0.8132					
PIK3CA	.0834	0.2919	.2487	0.5803	.8851	0.9220	.5855	0.8132					
RAS isoform mutation	.7303	0.7982	.7612	0.8231	.5831	0.7615	.7725	0.9168					
<b>RAS</b> pathway mutation	.7412	0.7982	.8630	0.8630	.6441	0.7615	.8475	0.9168					

FIG A4. RAS isoform and RAS pathway analysis. (A) RAS isoforms and pathway member *P* values for both EFS and OS. The log-rank *P* value and the BH adj value are presented. BH adj, Benjamini-Hochberg–adjusted; COG, Children's Oncology Group; EFS, event-free survival; OS, overall survival.



FIG A5. RAS codon and amino acid usage summary. Absolute number of cases with a mutation in the specified codon by amino acid change and isoform (NRAS, blue; HRAS, red; KRAS, green).

© 2021 by American Society of Clinical Oncology



FIG A6. TP53 mutations. (A) Association of TP53 mutations with other gene mutations across the entire rhabdomyosarcoms cohort. (B) Location and number of mutations across TP53 protein.



FIG A7. CDKN2A and association with survival. (A) EFS and OS by CDKN2A deletion in the COG cohort (blue) and the UK cohort (orange). (B) EFS and OS by CDKN2A status after removal of any cases with a co-occurring mutation of MYOD1: (continued on following page)

© 2021 by American Society of Clinical Oncology

Downloaded from ascopubs.org by 2.29.8.68 on September 17, 2021 from 002.029.008.068 Copyright © 2021 American Society of Clinical Oncology. All rights reserved. **FIG A7.** (Continued). COG cohort (blue) and UK cohort (orange). (C) EFS within the COG cohort demonstrated that the observed effect of *CDKN2A* alteration was largely within the intermediate-risk group. BH adj, Benjamini-Hochberg–adjusted; COG, Children's Oncology Group; EFS, event-free survival; HR, hazard ratio; Mut, mutated; OS, overall survival; WT, wild type.

Study	Cohort	No. of Cases	Percentage of Cohort
ARST0331	COG	50	14.5
ARST0431	COG	12	3.5
ARST0531	COG	68	19.8
ARST08P1	COG	9	2.6
D9602	COG	10	2.9
D9802	COG	3	0.9
D9803	COG	21	6.1
D9902	COG	171	49.7
EpSSG RMS2005	UK	48	16
MMT	UK	249	84

TABLE A1. Number of Patients Profiled From Each Trial

Abbreviations: COG, Children's Oncology Group; EpSSG, European paediatric Soft tissue sarcoma Study Group; MMT, malignant mesenchymal tumor; RMS, rhabdomyosarcoma.

### Journal of Clinical Oncology

TABLE A2. Includ	led Genes in the Capture Assay			
Gene	Genomic Alteration	Affected Exon	Published Frequency (%) <sup>a</sup>	References
AKT1				
ALK				10
ARID1A				26
ATM	Deletion and mutation			
BCOR	Multiple	All exons	7	9
BRAF	V600E	15	1	9
CCND1	Amplification			9,26
CCND2				
CDK4	Amplification			
CDKN2A	Deletion			
CTNNB1	Mutation			9,26
DICER1	Mutation			
ERBB2	Amplification			
FBXW7	R387P, R441G, R367	9	7	9
FGFR1	Amplification			
FGFR4	G528C, N535D/K, V550E/L/M	9, 10	9	9,10,26
GAB1				
HRAS	G12C, G13R, Q61K	2, 3	4	9,10,26
IGF1R	Amplification			10
IGF2	LOH			
KRAS	G12A/C/D	2	6	9,10,26
MDM2	Amplification			
MET	Amplification			9
MTOR				
MYCN	Amplification			
MYOD1	Point mutation			
NF1	Mutation, deletion, and LOH	All exons	5	9,10,26
NRAS	Q61H/K/R	3	12	9,10,26
PDGFRA				
РІКЗСА	E542K, E545K, Q546	9, 20	7	9
PKN1	Mutation			22
PTCH1				
PTEN	Deletion			22
PTPN11	E69K, A72T, E76A	3	2.5	9
ROBO1				26
SMARCA4				
SOS1	Mutation, duplication			
SOS2				
TP53	Multiple	5, 6, 7, 8	5	9,10,26

NOTE. Total estimated capture area 349,636 base pairs; estimated percentage of target bases covered = 98%. Abbreviation: LOH, loss of heterozygosity.

<sup>a</sup>Published frequency if available.

 ~		~

TABLE A3.	Fusion-Negative	Cases	Summarized	by	Mutation	and	Anatomic	Location
-----------	-----------------	-------	------------	----	----------	-----	----------	----------

Gene	Total No. of Gene Mutations in Cohort	Bladdes/ Prostate Mutant Cases	% of Gene Mutations Are Bladder/ Prostate	Tetal Cases Bladden/ Prostate	Mutant % in Bladder/ prostate	Extremity Mutant Cases	% of Gene Mutations Are Extremity	Total Cases Extremity	Mutant % in Extremity	Female GU Mutant Cases	% of Dene Mutations Are Female GU	Total Cases Female GU	Mutant % in Female GU	Head and Neck Mutant Cases	% of Dene Mutations Are Head and Neck	Total Cases Head and Neck	Mutant % in Head and Neck	Orbital Mutant Cases	% of Dene Mutations Are Orbital	Total Cases Orbital	Mutant % in Orbital	Other Mutant Cases	% of Gene Mutations Are Other	Total Cases Other	Mutant % in Other	Parameningeal Mutant Cases	% of Dene Mutations Are Parameningeal	Total Cases Parameningeal	Mutant % in Parameningeal	Paratesticular Motant Cases	% of Gene Mutations Are Paratesticular	Total Cases Paratesticular	Mutant % in Paratesticular	Retroperitonesse, Peritonesse, or Trusk Mutant Cases	% of Gene Mutations Are Retroperitoneum, Peritoneum, or Trunk	Tatal Cases Retroperitoneum, Peritoneum, er Trunk	Mutant % in Retroperitoneum, Peritoneum, or Trank
NRAS	87	8	8/87 (9)	49	849 (16)	2	2/87 (2)	35	2/35 (6)	4	4/87 (5)	18	4/18 (22)	15	15/87 (17)	47	15/47 (32)	9	9/87 (10)	43	9/43 (21)	1	1/87 (1)	7	1/7 (14)	11	11/87 (13)	105	11/105 (10)	28	28/87 (32)	120	28/120 (23)	9	9/87 (10)	87	9/87 (10)
BCOR	78	6	6/78 (8)	49	6/49 (12)	0	0/78 (0)	35	035 (0)	2	2/78 (3)	18	2/18 (11)	9	9/78 (12)	47	9/47 (19)	12	12/78 (15)	43	12/43 (28)	2	2/78 (3)	7	2/7 (29)	13	13/78 (17)	105	13/105 (12)	22	22/78 (28)	120	22/120 (18)	12	12/78 (15)	87	12/87 (14)
NF1	75	11	11/75 (15)	49	11/49 (22)	3	3/75 (4)	35	3/35 (9)	2	2/75 (3)	18	2/18 (11)	3	3/75 (4)	47	3/47 (6)	5	5/75 (7)	43	5/43 (12)	2	2/75 (3)	7	2/7 (29)	18	18/75 (24)	105	18/105 (17)	15	15/75 (20)	120	15/120 (13)	16	16/75 (21)	87	16/87 (18)
TP53	69	4	4/69 (6)	49	4/49 (8)	12	12/69 (17)	35	12/35 (34)	3	3/69 (4)	18	3/18 (17)	11	11/69 (16)	47	11/47 (23)	8	8/69 (12)	43	8/43 (19)	2	2/69 (3)	7	2/7 (29)	12	12/69 (17)	105	12/105 (11)	3	3/69 (4)	120	3/120 (3)	14	14/69 (20)	87	14/87 (16)
FGFR4	65	8	8/65 (12)	49	849 (16)	0	0/65 (0)	35	035 (0)	2	2/65 (3)	18	2/18 (11)	5	5/65 (8)	47	5/47 (11)	8	8/65 (12)	43	8/43 (19)	1	1/65 (2)	7	1/7 (14)	21	21/65 (32)	105	21/105 (20)	8	8/65 (12)	120	8/120 (7)	12	12/65 (18)	87	12/87 (14)
KRAS	44	2	2/44 (5)	49	249 (4)	1	1/44 (2)	35	1/35 (3)	0	044 (0)	18	0/18 (0)	4	4/44 (9)	47	4/47 (9)	1	1/44 (2)	43	1/43 (2)	0	0/44 (0)	7	0/7 (0)	9	9/44 (20)	105	9/105 (9)	14	14/44 (32)	120	14/120 (12)	13	13/44 (30)	87	13/87 (15)
HRAS	41	7	7/41 (17)	49	7/49 (14)	2	2/41 (5)	35	2/35 (6)	2	241 (5)	18	2/18 (11)	1	1/41 (2)	47	1/47 (2)	1	1/41 (2)	43	1/43 (2)	1	1/41 (2)	7	1/7 (14)	2	2/41 (5)	105	2/105 (2)	15	15/41 (37)	120	15/120 (13)	9	941 (22)	87	9/87 (10)
CTNNB1	32	5	5/32 (16)	49	5/49 (10)	0	0/32 (0)	35	035 (0)	1	1/32 (3)	18	1/18 (6)	2	2/32 (6)	47	2/47 (4)	2	2/32 (6)	43	2/43 (5)	0	0/32 (0)	7	0/7 (0)	2	2/32 (6)	105	2/105 (2)	9	9/32 (28)	120	9/120 (8)	11	11/32 (34)	87	11/87 (13)
PIK3CA	26	0	0/26 (0)	49	049 (0)	1	1/26 (4)	35	1/35 (3)	1	1/26 (4)	18	1/18 (6)	3	3/26 (12)	47	3/47 (6)	1	1/26 (4)	43	1/43 (2)	0	0/26 (0)	7	0/7 (0)	12	12/26 (46)	105	12/105 (11)	2	2/26 (8)	120	2/120 (2)	6	6/26 (23)	87	6/87 (7)
MDM2	26	2	6/26 (8)	49	249 (4)	6	6/26 (8)	35	6/35 (17)	1	1/26 (4)	18	1/18 (6)	1	1/26 (4)	47	1/47 (2)	1	1/26 (4)	43	1/43 (2)	0	0/26 (0)	7	0/7 (0)	5	5/26 (19)	105	5/105 (5)	7	7/26 (27)	120	7/120 (6)	3	3/26 (12)	87	3/87 (3)
CDKN/2A	23	0	0/23 (0)	49	049 (0)	3	3/23 (13)	35	3/35 (9)	0	0/23 (0)	18	0/18 (0)	3	3/23 (13)	47	3/47 (6)	3	3/23 (13)	43	3/43 (7)	0	0/23 (0)	7	0/7 (0)	8	8/23 (35)	105	8/105 (8)	0	0/23 (0)	120	0/120 (0)	6	6/23 (26)	87	6/87 (7)
FBXW7	18	1	1/18 (6)	49	149 (2)	0	0/18 (0)	35	035 (0)	0	0/18 (0)	18	0/18 (0)	1	1/18 (6)	47	1/47 (2)	1	1/18 (6)	43	1/43 (2)	0	0/18 (0)	7	0/7 (0)	1	1/18 (6)	105	1/105 (1)	10	10/18 (56)	120	10/120 (8)	4	4/18 (22)	87	4/87 (5)
MYOD1	17	0	0/17 (0)	49	049 (0)	1	1/17 (6)	35	1/35 (3)	0	0/17 (0)	18	0/18 (0)	4	4/17 (24)	47	4/47 (9)	0	0/17 (0)	43	0/43 (0)	0	0/17 (0)	7	0/7 (0)	11	11/17 (65)	105	11/105 (10)	0	11/17 (65)	120	0/120 (0)	0	0/17 (0)	87	0/87 (0)
CDK4	1	0	0/1 (0)	49	049 (0)	0	0/1 (0)	35	035 (0)	0	0/1 (0)	18	0/18 (0)	1	1/1 (100)	47	1/47 (2)	0	1/1 (100)	43	0/43 (0)	0	0/1 (0)	7	0/7 (0)	0	0/1 (0)	105	0/105 (0)	0	0/1 (0)	120	0/120 (0)	0	0/1 (0)	87	0/87 (0)
MYCN	0	0	00 (0)	49	049 (0)	0	Q/Q (0)	35	035 (0)	0	Q/D (0)	18	0/18 (0)	0	0/0 (0)	47	0/47 (0)	0	0/0 (0)	43	0/43 (0)	0	0/0 (0)	7	0/7 (0)	0	Q/D (0)	105	0/105 (0)	0	0/0 (0)	120	0/120 (0)	0	0/0 (0)	87	0/87 (0)
DICER1	12	0	0/12 (0)	49	049 (0)	0	0/12 (0)	35	035 (0)	6	6/12 (50)	18	6/18 (33)	0	0/12 (0)	47	Q/47 (0)	0	0/12 (0)	43	0/43 (0)	2	2/12 (17)	7	2/7 (29)	0	0/12 (0)	105	0/105 (0)	0	0/12 (0)	120	0/120 (0)	4	4/12 (33)	87	4/87 (5)
ARID1A	11	1	1/11 (9)	49	149 (2)	0	0/11 (0)	35	035 (0)	0	0/11 (0)	18	0/18 (0)	1	1/11 (9)	47	1/47 (2)	2	2/11 (18)	43	2/43 (5)	0	0/11 (0)	7	2/7 (5)	4	4/11 (36)	105	4/105 (4)	1	1/11 (9)	120	1/120 (1)	2	2/11 (18)	87	2/87 (2)

Abbreviation: GU, genitourinary.

<b>Risk Stratification</b>	FFS, %	Fusion Status	Stage	Group	Anatomy	Metastatic Sites	Genetic Marker
Low	> 85						
		Negative	l or ll	l or ll			
		Negative	I	III	Orbit only		
Intermediate	60-75						
		Negative	Any	III	Nonorbit		
		Negative	111	l or ll			
		Negative	IV	IV		1	
		Negative	Any low risk				TP53 mutant
		Positive	I, II, or III	I, II, or III			<i>TP53</i> WT
High	< 40						
		Negative	IV	IV		> 1	
		Negative	Any intermedi	ate risk			TP53 mutant
		Positive	IV	IV			<i>TP53</i> WT
Ultrahigh	< 20						
		Negative	Any	Any		Any	MYOD1 mutant
		Positive	Any	Any		Any	TP53 mutant

TABLE A4. Proposed Risk Stratification With the Incorporation of Genetic Markers

Abbreviations: FFS, failure-free survival; WT, wild type.