UNIVERSITY^{OF} BIRMINGHAM University of Birmingham Research at Birmingham

Pain catastrophizing mediates rapid benefits of accessing in-person chiropractic care during the COVID-19 lockdown

Gevers-Montoro, Carlos; Deldar, Zoha; Conesa-Buendía, Francisco Miguel; Lazar, Eric Arthur; Mahillo-Fernandez, Ignacio; Khatibi, Ali; Ortega de Mues, Arantxa

DOI: 10.1002/ejp.1872

License: Other (please specify with Rights Statement)

Document Version Peer reviewed version

Citation for published version (Harvard):

Gevers-Montoro, C, Deldar, Z, Conesa-Buendía, FM, Lazar, EA, Mahillo-Fernandez, I, Khatibi, A & Ortega de Mues, A 2021, 'Pain catastrophizing mediates rapid benefits of accessing in-person chiropractic care during the COVID-19 lockdown', *European Journal of Pain (United Kingdom)*, vol. 26, no. 2, pp. 463-479. https://doi.org/10.1002/ejp.1872

Link to publication on Research at Birmingham portal

Publisher Rights Statement:

This is the peer reviewed version of the article, which has been published in final form at https://doi.org/10.1002/ejp.1872. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions. This article may not be enhanced, enriched or otherwise transformed into a derivative work, without express permission from Wiley or by statutory rights under applicable legislation. Copyright notices must not be removed, obscured or modified. The article must be linked to Wiley's version of record on Wiley Online Library and any embedding, framing or otherwise making available the article or pages thereof by third parties from platforms, services and websites other than Wiley Online Library must be prohibited.

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Pain catastrophizing mediates rapid benefits of accessing in-person chiropractic care during the COVID-19 lockdown

Running Title: Benefits of in-person care during COVID-19 lockdown

C. Gevers-Montoro^{a,b}, Z. Deldar^{b,c}, F.M. Conesa-Buendía^d, E.A. Lazar^a, I. Mahillo-Fernandez^e, A. Khatibi^{f,g}, A. Ortega de Mues^{a,*}

^a Madrid College of Chiropractic – RCU María Cristina, Paseo de los Alamillos 2, 28200 San Lorenzo de El Escorial, Madrid, Spain.

^b Department of Anatomy, Université du Québec à Trois-Rivières, 3351 Boul. Des Forges, C.P. 500, Trois- Rivières, QC, Canada, G9A 5H7.

^c Department of Psychology, McGill University, 2001 McGill College Ave, Montreal, Quebec, H3A 1G1

^d Bone and Joint Research Unit, Institute of Health Research (IIS–Fundación Jiménez Díaz), Madrid, Spain.

^e Epidemiology and Biostatistics Unit (IIS-Fundación Jiménez Díaz), Madrid, Spain.

^f Centre of Precision Rehabilitation for Spinal Pain, University of Birmingham, Birmingham, UK

^gCentre for Human Brain Health, University of Birmingham, Birmingham, UK

*Corresponding author:

Arantxa Ortega de Mues Madrid College of Chiropractic Real Centro Universitario Escorial-María Cristina Paseo de los Alamillos, 2 28200 San Lorenzo de El Escorial, Madrid, Spain Phone: 34-91890 45 45 (ext. 211); Fax: 34-918906609 E-mail: aortega@rcumariacristina.com

Number of pages: 34 Number of figures: 5 (plus 2 supplemental) Number of tables: 3 (plus 2 appendices)

Funding sources: None declared

Conflicts of interest: None declared

What's already known about this topic?

- COVID-19 has directly and indirectly affected the course of pain conditions
- Limitations on care provision, psychological stress and physical inactivity all lead to increased pain intensity and interference
- Little is known about how to mitigate the negative impact of the pandemic on these conditions

What does this study add?

- Providing in-person chiropractic care, under strict sanitary measures, may provide quick pain relief and decrease interference
- Pain catastrophizing partially mediates these effects, while kinesiophobia and baseline pain intensity act as moderators
- In-person care during social distancing may be prioritized to a subset of patients by assessing pain intensity and cognitions

ABSTRACT

Background: In March 2020, the COVID–19 pandemic forced the Spanish government to declare a state of emergency. A stringent lockdown was enforced, restricting access to healthcare services, including chiropractic. The combination of reduced access to care provision with psychological stress, social isolation and physical inactivity during lockdown was shown to negatively influence pain conditions. However, data on strategies to mitigate the impact of the pandemic on these conditions is lacking.

Methods: Upon easing of restrictions in May 2020, fifty-one chiropractic clinics throughout Spain pseudo-randomly invited patients, recruiting a total of 385 participants. During a 14-day period, participants were exposed to in-person chiropractic care in either one (n=177) or multiple encounters (n=109), or to no care (n=99). The effects of access to chiropractic care on patients' pain-related and psychological outcomes were assessed online through validated self-reported questionnaires before and after the period of care. Coprimary outcomes included pain intensity, pain interference and pain cognitions.

Results: When comparing to participants without access to care, pain intensity and interference were significantly decreased at follow-up, irrespective of the number of encounters. Kinesiophobia was also significantly reduced at follow-up, though only after multiple encounters. The relationship between fear of movement, changes in pain intensity and interference was mediated by catastrophizing.

Conclusion: Access to in-person chiropractic care may provide pain relief, associated with reductions in interference and pain cognitions. Prioritizing in-person care for patients

with maladaptive pain cognitions may help dampen the detrimental consequences of the pandemic on physical and psychological well-being.

Keywords: chiropractic; COVID–19; lockdown; pain interference; kinesiophobia; pain cognitions; mediation.

1. Introduction

Shortly after the Coronavirus disease 2019 (COVID–19) was declared a pandemic by the World Health Organization (Mahase, 2020), the Spanish government imposed a state of national emergency and the country went into full lockdown (Spain Government, 2020b). Citizens remained confined to their homes except for essential business. Nonessential services were interrupted, limiting access to healthcare (Henriquez et al., 2020; Tanne et al., 2020), with consequences for patient management (Clauw et al., 2020; Lugli et al., 2020). Home confinement led to substantial decreases in physical activity and increases in sitting time (Ammar et al., 2020; Tison et al., 2020). Depression, anxiety and stress increased in over 25% of the population (Garcia-Alvarez et al., 2020; Ozamiz-Etxebarria et al., 2020; Rodriguez-Rey et al., 2020). Considering previous evidence on quarantine and sedentarism, the pandemic and policy changes to mitigate their impact can both be presumed to result in deleterious effects for mental and physical well-being (Brooks et al., 2020; Hall et al., 2020).

Direct and indirect effects of the pandemic are expected to result in a higher morbidity of pain conditions (Thacker, 2020), disproportionately affecting chronic pain (Clauw et al., 2020). Pandemic-induced psychological stress and restricted access to healthcare services during lockdown were linked to increased pain severity and interference, partially mediated by pain catastrophizing (Fallon et al., 2020; Hruschak et al., 2021; Nieto et al., 2020; Page et al., 2021). Previous data showed that pain cognitions, particularly catastrophizing, strongly influenced pain intensity, pain interference, and their relationship (Hirsh et al., 2011). Catastrophizing itself is not only dependent on previous painful experiences but can also be influenced by environmental factors (Petrini and Arendt-Nielsen, 2020). During COVID–19, psychological distress, catastrophizing, sedentarism and social isolation may all contribute to swell the burden of pain (Alzahrani et al., 2019; Hammig, 2019; Joseph et al., 2020; Zhaoyang et al., 2020).

Nearly twelve million people in Spain suffer from spine pain (Cruz-Sánchez et al., 2012), the most frequent reason to pursue chiropractic care (Beliveau et al., 2017). Access to chiropractic services has been linked to positive clinical outcomes for spine-related conditions (Blanchette et al., 2016; Garner et al., 2007; Goertz et al., 2018; Prater et al., 2020). During the state of emergency in Spain, chiropractic services were not deemed essential and clinics were to remain shut (Asociacion Espanola de Quiropractica, 2020). Whilst the pandemic may further increase the burden of pain on already overwhelmed healthcare systems, data on how to effectively tackle this crisis are missing (Carrillo-de-la-Pena et al., 2021).

We aimed to investigate the effects of accessing in-person chiropractic care on pain and psychological outcomes upon easing of lockdown measures. Specifically, we were interested in measuring pain intensity, interference and cognitions in patients without access to care during lockdown, and the effect of accessing in-person services on these variables and their interaction. We hypothesized that accessing care would reduce pain intensity and interference by influencing pain cognitions and psychological factors. Further, we intended to identify which patients would benefit most from accessing care in this context.

2. Methods

2.1 Ethical approval

A pragmatic controlled clinical trial was performed in the settings of 51 private chiropractic clinics from 16 of the 17 autonomous regions of Spain. Ethical approval was obtained from the Madrid College of Chiropractic research ethics committee (reference 300420). The study followed the Consolidated Standards of Reporting Trials (CONSORT) guidelines (except for the allocation process since randomization was not possible) and was registered in ClinicalTrials.gov with the trial ID: NTC04573361.

2.2 Study setting

On May 3, 2020, the Spanish government announced the end of the most stringent phase of lockdown (Spain Government, 2020a), which would go on to last until June 21. Some services were allowed to reopen under strict hygiene and safety measures, including chiropractic clinics. In particular, the Spanish Chiropractic Association published a set of guidelines for their members to reopen, enforcing social distance, universal maskwearing, temperature control, ventilation and sanitization of surfaces, tables, instruments, and practitioners' hands. The trial was conducted between May 4 and May 26, 2020, during the initial phase of lockdown *de-escalation*, as referred to by the Spanish healthcare authorities. Not all chiropractic clinics were open during this early phase, depending on practitioner preference and clinic capacity. Personal protective equipment was scarce and adapting the clinic to new sanitary regulations required time and monetary investments, making reopening difficult or infeasible in some cases. This set the stage for two possible scenarios: one in which patients had access to care and one in which they did not.

2.3 Patient recruitment

All chiropractors registered in the Spanish Chiropractic Association were invited via e-mail to participate in the recruitment process, and 51 private chiropractic clinics throughout Spain accepted. Participants were chiropractic patients from these clinics over 16 years of age, presenting with pain at the time of recruitment. Only active patients at the time of the closure of chiropractic clinics in March were to be included. We intentionally excluded new patients or patients who had not been under care for long periods of time before the pandemic. One of the reasons was to minimize the number of patients who perceived no need to resume care. As lockdown measures were being eased, clinics contacted regular patients to provide information on the status of the clinic reopening, and when possible for both sides, to schedule an appointment. Participant clinics were requested to generate a contact list and to pseudo-randomly invite one of every three patients contacted to participate in the study, independently of whether they would be scheduling an appointment or not. This was done to reduce a recruitment bias on behalf of the person contacting potential participants. Contacts were made via phone calls, e-mails or text messages by the chiropractor or the administrative staff.

Patients accepting to participate were provided a link to access the initial online questionnaire that was to be completed during the first seven days of the trial (May 4 to May 11), always before any treatment was initiated (see Fig. 1). Access to the initial questionnaire was closed on May 11. Participants in the study were instructed to fill out a follow-up questionnaire 14 days after the date of completion of their initial response. Access was secured until the end of the study (May 18 to May 26) and a total of 500 responses were received. After removal of duplicates, non-paired and non-valid responses, participants receiving chiropractic teleconsultations or answering incorrectly all three screening questions, 385 patients were included (see supplemental Figure S1 for a patient inclusion and participation flowchart). A description of all demographic data is available in Table 1.

An a priori power analysis was conducted using G^{*}Power for between-within interactions in ANOVA (Faul et al., 2007). The parameters for this analysis were: an alpha of 0.05, a power of 0.95, a conservative estimate of the correlation among repeated measures of r = 0, for three groups, two measurements and an effect size of f = 0.268, based on pain intensity data reported in a pragmatic trial of chiropractic spinal manipulation for low back pain (Goertz et al., 2013). The analysis yielded a required sample of at least 111 participants, 37 participants in each group to reach statistical significance. Thus, our sample size of 385 largely exceeded this threshold.

2.4 Study protocol

The present study is a pragmatic controlled clinical trial in which chiropractic patients were contacted by their clinic and were invited to participate in a pseudo-random order (one of every three patients contacted was invited, independently of future treatment). Therapists remained blinded to which patients eventually took part or did not, and investigators were blinded to participant recruitment and treatment received. During the 14 days following completion of the initial self-administered online questionnaire, patients participating in the study either resumed care with one (Group 1) or more than one visit (Group 2) to their chiropractor, or else did not have access to care (Group 0). This assigned patients to one of three groups, depending on opening times and availability of each clinic and the individual patient. The follow-up questionnaire was completed 14 days later, determining the end of the trial period (see Fig. 1).

2.5 Hands-on chiropractic care

Registered chiropractors, who were not aware of the study hypotheses and group allocation, provided routine care based on individual patient needs. Chiropractic is a type of conservative spine care, which consists mainly of the application of manual therapy (Nelson et al., 2005). Nearly four out of five chiropractic patients receive spinal manipulation, while soft tissue therapy, patient education and exercise therapy are less frequently part of care (Beliveau et al., 2017). The most common reasons for seeking chiropractic care are low back pain, neck pain and extremity complaints (Beliveau et al., 2017). In accordance with previous reports, clinics participating in the current trial provided in-person care primarily based on spinal manipulation (100% of patients), followed by patient education and exercise prescription. For details on the frequency of use of different interventions, see Table 2.

2.6 Patient-reported outcome measures

Both the initial and follow-up questionnaires were designed and completed online using Google Forms (Google Inc., Mountain View, CA, US). The link to access the initial questionnaire was provided by the chiropractic clinics, while the link for the follow-up questionnaires was provided either by the clinic staff or one of two investigators not involved in data analysis (AOdM and FMC-B) for patients who had provided contact information. When planning the study, two representatives of the national chiropractic patient association (Asociación Española de Usuarios de **Ouiropráctica**: http://www.aeuq.es/) provided feedback on the format and length of the questionnaires, as well as on the adequacy of the outcome measures used. The questionnaires comprised three sections with the following data: demographics (first section), pain-related variables (second section) and variables on the psychosocial impact of COVID–19 (third section). At the beginning of the initial questionnaire, participants provided informed consent to

participate in the study and to use their responses. All questions and answers were provided in Spanish, using the validated versions in Spanish of structured questionnaires when available. Sections and individual items are detailed in Appendices I and II.

2.6.1 Coprimary outcomes

Following the demographic questions, the second section of both the initial and follow-up questionnaires contained pain variables, including the coprimary outcomes: current pain intensity (numerical rating scale from 0 to 10, where 0 = no pain at all, and 10 = maximum pain imaginable), pain interference from the brief pain inventory (BPI) (de Andres Ares et al., 2015; Keller et al., 2004), the short version of the pain catastrophizing scale (PCS-4) (Bot et al., 2014; Olmedilla Zafra et al., 2013) and the short version of the Tampa Scale of Kinesiophobia (TSK-11) (Gómez-Pérez et al., 2011). The validated version in Spanish of the BPI is widely used for noncancer pain and its interference scale showed good reliability with a Cronbach's a of 0.93 (de Andres Ares et al., 2015). Considering that the confinement measures explicitly forbid walking outdoors, the patient representatives suggested excluding the item related to interference with "walking", hence it was not used. The PCS-4 has shown good internal consistency (a = 0.86) and almost perfect correlation with the long version (r = 0.96) (Bot et al., 2014), which has been validated in Spanish (Olmedilla Zafra et al., 2013). The Spanish version of the TSK-11 has been validated and has shown good reliability and validity with a 2factor solution, namely activity avoidance and harm (Gómez-Pérez et al., 2011).

A 30% reduction from baseline has been proposed as a minimal clinically important difference for both pain intensity and disability, particularly for low back pain (Ostelo et al., 2008; Rowbotham, 2001). Alternatively, a 10-20% decrease in pain intensity and a one-point reduction in the BPI pain interference were recommended as

cut-offs for minimally important differences in chronic pain patients (Dworkin et al., 2008). Considering the limited timeframe and the passive character of the intervention, clinically meaningful differences were only expected for pain intensity and interference, but not for pain cognitions.

2.6.2 Secondary outcomes

Other pain variables from the second section of the questionnaire such as pain duration, frequency and perceived evolution were secondary outcomes. Perceived evolution referred to patients' self-reported perceived changes in their pain during lockdown (before the trial) and during the trial.

The third section comprised questions on symptoms, diagnosis, treatment and exposure to COVID–19, as well as on the impact, emotional responses, fears and stresses evoked by the pandemic and associated restrictions (details available in Appendices I and II), followed by the validated version in Spanish of the Generalized Anxiety Disorder scale (GAD-7) (García-Campayo et al., 2010).

2.6.3 Access to healthcare services variables

The third section also evaluated access to healthcare services during lockdown and the trial period. One specific question concerning access to chiropractic care was included only in the follow-up questionnaire. The answer to this question determined the group in which patients were placed: no access (Group 0), one visit (Group 1) or more than one visit (Group 2) to their chiropractor. Patients receiving care were asked to rate four dimensions of their satisfaction (hygiene, safety, effectiveness, global satisfaction) using a 1–4 Likert scale. Patients were also asked whether they had visited a healthcare professional other than their chiropractor for their pain or other reasons (at baseline and follow-up).

Included in the different sections were three instructional manipulation checks, screening questions that are known to increase the reliability of the dataset (Oppenheimer et al., 2009). These are questions similar in format, length and answers to the experimental questions. However, instead of providing experimental data, they are used as a mechanism to confirm that the participant is correctly reading the questions (Oppenheimer et al., 2009). This resulted in a total of 73 and 71 items initially and at follow-up, respectively.

2.7 Statistical analysis

Statistical analyses were performed with Statistica (v13.3; Dell Inc., Tulsa, OK, USA), SPSS (v25 statistical package IBM SPSS Statistics, Armonk, NY, USA) and the PROCESS module for SPSS (v3.3) (Hayes, 2017). A value of p < 0.05 was used as a threshold to determine statistical significance.

For all demographic variables, means and standard deviations or percentages from the total samples were calculated. Separate Chi-Square analyses were used to compare categorical variables between groups. To measure the influence of access to chiropractic care, changes in all pain-related variables (Section 2), as well as those of fear/stress and anxiety linked to the pandemic (Section 3) over time, were compared between experimental groups using mixed model ANOVAs. Equality of variance was assessed by using the Brown-Forsythe test (Brown and Forsythe, 1974). To control for potential interactions of pain duration and of access to another healthcare professional, these were both introduced as covariates in separate covariance analyses. Significant interactions were decomposed by using Bonferroni-corrected planned comparisons (3 comparisons) to test *a priori* hypotheses. We hypothesized that patients with access to care, independent of the number of encounters, experience reductions in pain intensity and interference compared to patients without. Further, we hypothesized that pain cognitions, pandemicrelated stress and emotions would improve for patients visiting their chiropractor. Effect sizes are expressed as partial eta squared values (η^2_p) . Small effect size is considered when η^2_p values ≥ 0.01 , medium when $\eta^2_p \ge 0.06$ and large $\eta^2_p \ge 0.14$. Correlation coefficients (Pearson's *r*) were calculated for all numerical coprimary and secondary outcomes, at baseline and follow-up. These analyses were conducted to generate hypotheses regarding which coprimary or secondary variables may act as mediators of the changes in pain intensity and interference. The results of the correlation analyses are available as supplemental material in Figure S2.

In order to examine the mediatory role of pain catastrophizing in the relationship between the change in pain intensity ratings and the change in pain interference, a simple moderation analysis (model 4 of Hayes) was used (Hayes, 2017). In this model, the influence of the predictor variable (change in pain intensity) over the outcome (change in pain interference) is mediated by the mediator variable (pain catastrophizing). Further, Hayes' model 7 (moderated mediation) was used to test the moderating role of baseline pain intensity in the mediatory relationship between change in pain ratings and pain catastrophizing (PCS). In this model, the moderating variable (baseline pain intensity) moderates the relationship between the predictor and the mediator (Hayes, 2017). Another mediation analysis was performed to test the mediatory effect of PCS on the relationship between fear of movement at baseline and change in pain intensity ratings.

3. Results

3.1 Demographics

All demographic data for the whole sample and for each individual group are summarized in Table 1. The final sample of 385 participants was comprised of 99 patients who reported receiving no care (Group 0), 177 patients reportedly accessing chiropractic care on one occasion (Group 1) and 109 patients visiting their chiropractor more than once (Group 2). Baseline scores in pain-related variables for all participants and groups are available in Table 3.

3.2 Coprimary outcomes

3.2.1 Pain intensity

Mixed ANOVAs for pain intensity ratings with TIME as the within-subjects factor and GROUP (access to care) as the between-subjects factor revealed a significant main effect of TIME ($F_{(l,382)}$ =90.97, p<0.001, η^2_p =0.19). Homogeneity of variances was respected for baseline and follow-up data (p >0.95 for both). For the whole sample, mean pain intensity significantly decreased 1.23 points over time [95% confidence interval (C.I.) -1.46 to -1.0]. Pain intensity decreased by 0.53 (-0.97 to -0.08), 1.53 (-1.88 to -1.17) and 1.38 (-1.77 to 0.98) points in Groups 0, 1 and 2, respectively. There was a significant TIME x GROUP interaction ($F_{(2.382)}$ =6.46, p=0.002, η^2_p =0.03). Planned comparisons used to describe this interaction revealed a significant difference in the evolution over time when comparing Group 0 to both Groups 1 and 2 ($F_{(l,382)}$ =12.26, p=0.002 and $F_{(l,382)}$ =7.25, p=0.02, respectively). However, no differences were found over time between Groups 1 and 2 ($F_{(l,382)}$ =0.29, p=1.0). Fig. 2 provides a graphic representation of these results.

3.2.2 Pain interference with activity

Pain interference showed a negative main effect of TIME ($F_{(l,382)}=14.17, p<0.001$, $\eta^2_p=0.04$). Variances were not significantly different between groups (p=0.08 and p=0.17at baseline and follow-up respectively). There was a significant TIME x GROUP interaction ($F_{(2,382)}=7.30$, p<0.001, $\eta^2_p=0.04$). Planned comparisons showed that interference was significantly different over time when comparing Group 0 to Group 1 ($F_{(l,382)}=12.79$, p=0.001) and to Group 2 ($F_{(l,382)}=10.04$, p=0.005), but not between Groups 1 and 2 ($F_{(l,382)}=0.01, p=1.0$). Interference scores showed a mean increase in 1.34 points for Group 0 (95% C.I. -1.06 to 3.75), and a mean reduction in 4.8 points for Group 1 (-6.98 to -2.62) and 4.68 points for Group 2 (-7.13 to -2.23). See Fig. 3 for a depiction of these results.

3.2.3 Pain Catastrophizing and Fear of Movement (Kinesiophobia)

Mixed ANOVAs showed a significant main effect of TIME for PCS and TSK $(F_{(1,382)}=58.24, p<0.001, \eta^2_p=0.13; F_{(1,382)}=4.10, p=0.04, \eta^2_p=0.01, respectively)$. For both variables, the variances were homogenous at baseline and follow-up (all *p*'s >0.3). Since gender differences in catastrophizing have been extensively reported in the literature (Sullivan et al., 2000), the PCS scores were tested for gender differences with the use of one-way ANOVA. The scores were higher for women than for men, though not significantly (6.7 vs. 6.4 at baseline, *p*=0.32; 5.8 vs. 5.4 at follow-up, *p*=0.22).

Only TSK was influenced by exposure to interventions (TIME x GROUP: $F_{(2,382)}$ =4.29, p=0.01, η_p^2 =0.02). Planned comparisons revealed that TSK scores did not have a significantly different evolution when comparing no exposure to one ($F_{(1,382)}$ =5.44, p=0.06), however, it was the case for multiple exposures ($F_{(1,382)}$ =7.83, p=0.02). No differences were found between both groups exposed (p=1.0). Group 0 showed an increase in 0.47 points (95% C.I. -0.27 to 1.22), Group 1 a decrease in 0.68 points (-1.22)

to -0.13) and Group 2 in 1.06 points (-1.89 to -0.22). Results are shown in Fig. 4. and correlations between these coprimary outcomes are represented as supplemental material in Figure S2.

3.3 Influence of pain duration and access to other healthcare services

Patients were classified as acute or chronic according to their self-reported symptom duration (< 3 and \geq 3 months, respectively). The sample classified as "chronic" showed significantly higher levels of pain catastrophizing ($F_{(1,379)}=12.39$; p<0.001) and kinesiophobia ($F_{(1,379)}=9.25$; p=0.003). In order to determine whether pain duration was a factor influencing the evolution of participants allocated to different groups, it was introduced as a covariate in a set of separate ANCOVAs. Introducing pain duration did not result in significant effects on the interactions TIME x GROUP x PAIN DURATION (pain intensity: $F_{(2,379)}=2.86$, p=0.06, $\eta^2_p=0.06$; pain interference: $F_{(2,379)}=0.21$, p=0.81, $\eta^2_p=0.001$; kinesiophobia: $F_{(2,379)}=1.35$, p=0.26, $\eta^2_p=0.01$).

During the trial, some participants reported visiting another healthcare professional (HCP) for their pain (12%, 9% and 8% of patients in Groups 0, 1 and 2, respectively). No significant interactions TIME x GROUP x HCP were observed for pain interference ($F_{(2,379)}$ =0.46, p=0.6, η^2_p =0.002) or kinesiophobia ($F_{(2,379)}$ =0.73, p=0.5, η^2_p =0.004). Accessing another HCP only had a significant influence on pain intensity (interaction: $F_{(2,379)}$ =4.6, p=0.01, η^2_p =0.02). Interestingly, patients visiting another professional did not have a significant pain reduction (-1.75 to 0.75, 95% C.I.), whereas patients only visiting their chiropractor did (-2.16 to -1.32, 95% C.I.).

3.4 Mediation effects by pain catastrophizing

The mediatory effect of baseline pain catastrophizing (PCS) on the relationship between change in pain intensity (predictor) and change in pain interference between pre and post-test (outcome variable) was assessed (Fig. 5-A). The indirect effect of change in pain intensity on change in pain interference was found to be statistically significant (b=0.144, 0.032<C.I.<0.291). Further, testing moderated mediation using Hayes model number 7 (Hayes, 2017) showed that the relationship between the difference in pain intensity and PCS was moderated by the intensity of pain the individual had at baseline ($F_{(l,38l)}$ =5.70, p=0.02). The moderation effect suggests that when an individual has a lower pain rating at baseline, a reduction in pain is associated with lower pain catastrophizing. Correspondingly, an increase in pain among those with a high intensity at baseline is associated with high pain catastrophizing (Fig. 5-B).

Next, the mediatory role of baseline pain catastrophizing in the relationship between baseline fear of movement and change in pain ratings was investigated. The overall model with change in pain intensity as the dependent variable was significant ($R^2=0.039$, $F_{(3,381)}=5.17$, p=0.002). Only the indirect effect of fear of movement on pain was significant (0.022 < C.I. < 0.085). When gender was included in the above-mentioned model, there was a trend towards significance when PCS was the outcome variable (p=0.057). Therefore, the analysis was repeated for each gender separately. Among women, the indirect effect of PCS in the relationship between TSK and changes in pain intensity was significant (Fig. 5-C) (0.024 < C.I. < 0.097), but not among men (Fig. 5-D) (-0.022 < C.I. < 0.096).

A subsequent moderation analysis was performed with the change in pain intensity as the outcome variable for analysis, and fear of harm ("harm" subscale of the TSK) as a moderator variable. The model in the full sample showed that the interaction between harm and initial pain intensity was not significant. When the model was tested among those visiting their practitioner more than once, the interaction between harm and initial pain intensity showed a trend towards significance ($F_{(1,105)}$ =3.75, p=0.055). Contrarily, the observed effect was not significant or close to significant in the other two groups. For Group 2 exclusively, the conditional effect of all three levels of the moderator was significant. Individuals with higher pain intensity at baseline and higher harm scores had less reduction in pain at follow-up, while those with similar pain ratings at baseline but lower levels of harm had more pain reduction (more than 1.7 times compared to the high harm group). There was not such a difference between individuals with low baseline pain intensity.

3.5 Secondary outcomes

3.5.1 Perceived evolution

Patients rated their perceived evolution during the lockdown period and at the completion of the study. During the lockdown period, two out of three patients (65%) reported that their pain had worsened. After completion of the trial, pain had improved or was gone in 61% and 77% of patients accessing care once and more than once, respectively. The proportion of patients improving was significantly lower (χ^2 =39.8, df=2, p<0.001) in patients without access to care (34%).

3.5.2 Pandemic-related emotional profile

Mixed ANOVAs showed a significant main effect of TIME for general anxiety $(F_{(1,382)}=19.27, p<0.001, \eta^2_p=0.05)$, which decreased in the trial period (95% C.I. -0.98 to -0.38). However, there was no significant interaction by GROUP $(F_{(2,382)}=1.65, p=0.19)$.

A set of questions assessed the degree (from 0 to 10) to which emotions were evoked by the pandemic (sadness, worry, loneliness, anger/rage, helplessness, anxiety, surprise, relief and hope). Of the emotions reported, only *Hope* showed a significant interaction of TIME with GROUP ($F_{(2,382)}=3.79$, p=0.02, $\eta^2_p=0.02$). The assumption of variance equality was met (p > 0.3 at baseline and follow-up). Changes in the degree that this emotion was reported significantly differed when comparing Group 0 to Group 1 only: $F_{(1,382)}=7.27$, p=0.02 (Group 0 vs. Group 2: p=0.8; Group 1 vs. Group 2: p=0.4). *Hope* only increased for participants in groups receiving care once (0.29 to 1.32, 95% C.I.); patients in the other two groups did not experience significant changes (-0.85 to 0.27 for Group 0 and -0.41 to 0.83 for Group 2, 95% C.I.).

Concerning the questions on pandemic stress and/or fear, only stress in the face of the COVID–19 pandemic showed a significant evolution over time ($F_{(1,382)}$ =9.90, p=0.002, η^2_p =0.03) and a trend for significance by group ($F_{(2,382)}$ =2.95, p=0.054, η^2_p =0.015). The variances were homogenous between groups at baseline and follow-up (p=0.06 and p=0.14 respectively). Despite not finding significant differences between groups, a significant pre-post decrease (-1.16 to -0.31, 95% C.I.) in stress from COVID– 19 was seen for Group 2 but not for the rest (-0.39 to 0.35 for Group 0 and -0.64 to 0.02 for Group 1, 95% C.I.). (Correlations between the numerical secondary outcomes are illustrated in supplemental Figure S2).

Patients with access to chiropractic care rated their satisfaction with different aspects of care in a Likert scale (1–4) as follows: hygiene 3.53 to 3.74; safety 3.51 to 3.72; effectiveness 3.48 to 3.70; and overall satisfaction 3.46 to 3.68 (all 95% C.I.).

4. Discussion

This pragmatic non-randomized trial evaluated access to in-person chiropractic care at the end of the most stringent phase of lockdown taking place in Spain during the spring of 2020. When lockdown measures were eased, accessing care rapidly decreased pain intensity, interference with activity, and fear of movement. Pain catastrophizing was identified as the mediator of the relationships between these variables. Patients visiting their chiropractor in this period experienced a significant increase in feelings of hope with regards to the COVID–19 pandemic. Satisfaction with care was high (~ 3.6/4 overall) and none of these results seemed to be influenced by exposure to other healthcare professionals or pain management services.

To redistribute and prioritize healthcare resources during COVID–19, access to and provision of services were limited or delayed (Lazzerini et al., 2020; Minhas et al., 2020). Restricting access to care forced patients to change their treatment, leading to increased pain intensity and interference (Fallon et al., 2020; Hruschak et al., 2021; Lacasse et al., 2021; Nieto et al., 2020; Page et al., 2021). Correspondingly, two-thirds of our patients reported aggravation of their pain during lockdown. In the context of COVID–19 social distancing, self-perceived increases in pain intensity and interference were independently predicted by catastrophizing (Fallon et al., 2020; Hruschak et al., 2021). In the present study, pain catastrophizing also emerged as an important factor explaining the relationship between pain intensity and interference. This is aligned with previous data on the mediating role of catastrophizing on disability/interference (Hirsh et al., 2011; Marshall et al., 2017). Individuals' cognitions about pain affect their perception on how it interferes with daily activities (Sullivan et al., 2001). However, the relationship between pain and catastrophizing is not a direct one. Our findings supported that severity at baseline moderates this relationship: higher baseline pain intensity and catastrophizing were associated with pain aggravation after exposure. Indeed, state pain catastrophizing may be influenced by a person's experience of pain (Sturgeon and Zautra, 2013).

These findings add support to the premise that pain intensity and catastrophizing may influence levels of physical activity (Zhaoyang et al., 2020). It was proposed that catastrophizing also mediates the relationship between pain and fear (Marshall et al., 2017). This is compatible with the findings of our second mediation analysis. Fear of movement influenced changes in pain intensity indirectly through pain catastrophizing. Kinesiophobia has been associated with higher pain intensity and disability, predicting the evolution of the latter (Luque-Suarez et al., 2019). High scores, particularly in the harm subscale, were linked to low levels of physical activity (Elfving et al., 2007), while reducing kinesiophobia could stimulate engagement in physical activity for patients with back pain (Luning Bergsten et al., 2012). Currently, the promotion of physical activity is a public health priority (Damiot et al., 2020; Pinto et al., 2020; J. F. Sallis et al., 2020; Stamatakis and Bull, 2020), particularly as inactivity is associated with a higher risk for severe COVID-19 (R. Sallis et al., 2021). In the present study, fear of movement was significantly reduced after longer exposure to hands-on care. Though this may have influenced the patients' degree of engagement in physical activity, this variable was not assessed. Activity levels may explain augmented pain during lockdown (Fallon et al., 2020; Yoshimoto et al., 2021), and may change in response to care, or vice versa. This variable should be monitored and controlled for in future studies.

Our results are in line with literature supporting that pain catastrophizing and fear of movement modulate treatment efficacy (Flink et al., 2010; Wertli et al., 2014a; Wertli et al., 2014b). Participants were mainly exposed to manual therapy, for which the mediators of treatment success are poorly understood. Pain relief from spinal manipulation was found to partly rely on influencing psychological variables, including fear of movement (Ellingsen et al., 2018; Williams et al., 2007). Catastrophizing may play an important role, but the direction of the effect remains unclear (Bishop et al., 2015). It was found to moderate changes in pain sensitivity (Alonso-Perez et al., 2017), but also to increase the chances of response to manual therapy (Verhagen et al., 2010). The current study provides evidence for pain catastrophizing as a mediator of the relationships between kinesiophobia, changes in pain intensity and in pain interference.

Kinesiophobia, particularly the harm subscale of the TSK-11, showed a tendency to decrease the likelihood of benefitting from access to care in individuals with more intense pain at baseline. This is consistent with high levels of fear avoidance beliefs decreasing the likelihood of responding favourably to spinal manipulation (Childs et al., 2004; Cleland et al., 2007) and greater pain intensity moderating the response to it (Schellingerhout et al., 2008). However, a recent systematic review did not find these effects to be clinically meaningful (de Zoete et al., 2021).

There is currently little data on the effects of re-introducing specific services and novel initiatives to counteract the consequences of delaying the provision of care for non–COVID conditions (Carrillo-de-la-Pena et al., 2021). Previous evidence supports that the introduction of chiropractic services significantly reduces pain and disability (Garner et al., 2007; Goertz et al., 2018; Prater et al., 2020). In the present investigation, hands-on chiropractic care resulted in minimal clinically important differences for both outcomes (approaching 30% from baseline), particularly after the first visit. Exposure to additional visits or other healthcare services did not yield better outcomes. This may be explained

by a strong influence of in-person care during lockdown on psychological factors. Modulation of distressing factors such as fear, anxiety and beliefs may contribute to the short-term clinical benefits of spinal manipulation (Williams et al., 2007). Similarly, feelings of hope were leveraged after the first encounter and stress diminished after multiple interactions. These changes likely reflect the influence of contextual factors linked to hands-on care in times of high uncertainty (Kaptchuk et al., 2020) and stress associated to the pandemic (Taylor et al., 2020).

In the context of COVID–19, the effect of touch, context and therapeutic alliance associated to chiropractic care may be particularly relevant for individuals with a specific psychological profile (Bishop et al., 2015). These individual differences should be considered when planning patient access to care during restrictions. A careful selection process may be necessary to prioritize care provision for patients who will benefit more from in-person access, as opposed to those who may wait or find alternatives, such as remote interventions. Telehealth services offer a safer alternative to in-person care during times when social distancing is pivotal (Eccleston et al., 2020). It remains to be determined whether they provide equivalent benefits and levels of satisfaction (Carrillo-de-la-Pena et al., 2021).

The interpretation of the study results is limited by the lack of treatment randomization. We did not consider it ethical to assign a specific treatment after two months of strict lockdown, as it could further delay care for participants with more urgency (higher pain intensity, interference or anxiety) or increase the exposure of those at higher risk of infection. It is possible that some patients declined to receive treatment if they perceived that the risks outweighed the potential benefits amidst the pandemic. A cluster randomization was not deemed feasible due to different barriers or accelerators influencing the reopening scenario for each clinic, such as economic factors, personnel, infrastructure and personal protective equipment availability. Instead, the recruitment process was pseudo-randomized, attempting to reduce sampling bias. However, this does not mean that the experimental design is devoid of other biases. It is possible that participants in the groups receiving care were more likely to be lost to follow-up if the results did not meet their expectations. This arises as a high risk for attrition bias (Nunan et al., 2018), which was moderated by using a mixed model analysis (Bell et al., 2013). Other limitations come from the conditions during the pandemic. A pragmatic trial with self-reported online data was deemed the most appropriate design to provide "real-world" data (Christian et al., 2020; Gartlehner et al., 2006), while ensuring social distancing. This offered a short window of opportunity to measure rapid effects of access to chiropractic care, during the two weeks that made up the initial phase of lockdown easing.

To our knowledge, this is the first study to measure the impact on pain conditions of access to a healthcare service that was limited due to restrictions and made available anew in the midst of the COVID–19 pandemic. This provided a unique setting to evaluate the effects of accessing chiropractic services –of limited access during the pandemic– on pain and psychological variables. The sample (covering the majority of the Spanish territory) and the mediation analyses suggest potential benefits of hands-on care mediated by pain catastrophizing and moderated by baseline pain intensity and kinesiophobia. The provision of in-person chiropractic care under strict regulations concerning safety and hygiene, may be considered to mitigate some of the negative consequences of the pandemic on patients with pain. However, individual differences in pain perception and cognitions should be examined in order to effectively prioritize access to care during the COVID–19 pandemic.

Acknowledgments

We thank the trial participants for their time and interest in the study; the Spanish chiropractic patient association (Asociación Española de Usuarios de Quiropráctica) for their advice and help in designing the study; the Spanish Chiropractic Association (Asociación Española de Quiropráctica) for their help in enrolling chiropractic clinics; the professionals for recruiting patients; the University of Birmingham for its support; Camino García-Balboa for advice on statistical analyses; and Erfan Ghalibaf for his help with data visualization. Figure 1 was created with <u>BioRender.com</u>.

Author contributions: CG-M, ZD and AOdM contributed to conceive and plan the overall trial design. The questionnaires were developed by CG-M, ZD, FMC-B, AK and AOdM. CG-M, FMC-B and AOdM contributed to the administration of questionnaire trial data collection, CG-M and AK performed the statistical analyses, AOdM is the principal investigator, IMF the guarantor of statistical quality. CG-M wrote the first version of the manuscript. EL contributed to manuscript writing and editing. All the authors reviewed the manuscript. The team met regularly during the trial period and contributed as a whole to discussions of the data.

Conflicts of interest: This study was fully self-funded, and the authors received no specific funding for this work. All authors declare no financial relationships with any organizations that might have an interest in the submitted work and no other relationships or activities that could appear to have influenced the submitted work.

References

- Alonso-Perez, J. L., Lopez-Lopez, A., La Touche, R., Lerma-Lara, S., Suarez, E., Rojas, J., . . . Fernandez-Carnero, J. (2017). Hypoalgesic effects of three different manual therapy techniques on cervical spine and psychological interaction: A randomized clinical trial. *J Bodyw Mov Ther*, 21(4), 798-803. doi:10.1016/j.jbmt.2016.12.005
- Alzahrani, H., Mackey, M., Stamatakis, E., Zadro, J. R., & Shirley, D. (2019). The association between physical activity and low back pain: a systematic review and meta-analysis of observational studies. *Sci Rep*, 9(1), 8244. doi:10.1038/s41598-019-44664-8
- Ammar, A., Brach, M., Trabelsi, K., Chtourou, H., Boukhris, O., Masmoudi, L., . . . Hoekelmann, A. (2020). Effects of COVID-19 Home Confinement on Eating Behaviour and Physical Activity: Results of the ECLB-COVID19 International Online Survey. *Nutrients*, 12(6). doi:10.3390/nu12061583
- Asociacion Española de Quiropractica (2020). COVID-19: Cierre De Centros Quiroprácticos. Retrieved from <u>https://quiropractica-aeq.net/covid-19-cierre-de-centros-quiropracticos/</u>
- Beliveau, P. J. H., Wong, J. J., Sutton, D. A., Simon, N. B., Bussières, A. E., Mior, S. A., & French, S. D. (2017). The chiropractic profession: a scoping review of utilization rates, reasons for seeking care, patient profiles, and care provided. *Chiropractic & Manual Therapies*, 25(1). doi:10.1186/s12998-017-0165-8
- Bell, M. L., Kenward, M. G., Fairclough, D. L., & Horton, N. J. (2013). Differential dropout and bias in randomised controlled trials: when it matters and when it may not. *BMJ*, 346, e8668. doi:10.1136/bmj.e8668
- Bishop, M. D., Torres-Cueco, R., Gay, C. W., Lluch-Girbes, E., Beneciuk, J. M., & Bialosky, J. E. (2015). What effect can manual therapy have on a patient's pain experience? *Pain Manag*, 5(6), 455-464. doi:10.2217/pmt.15.39
- Blanchette, M. A., Stochkendahl, M. J., Borges Da Silva, R., Boruff, J., Harrison, P., & Bussieres, A. (2016). Effectiveness and Economic Evaluation of Chiropractic Care for the Treatment of Low Back Pain: A Systematic Review of Pragmatic Studies. *PLoS One*, 11(8), e0160037. doi:10.1371/journal.pone.0160037
- Bot, A. G., Becker, S. J., Bruijnzeel, H., Mulders, M. A., Ring, D., & Vranceanu, A.-M. (2014). Creation of the abbreviated measures of the Pain Catastrophizing Scale and the short health Anxiety inventory: the PCS-4 and SHAI-5. *Journal of Musculoskeletal Pain*, 22(2), 145-151.
- Brooks, S. K., Webster, R. K., Smith, L. E., Woodland, L., Wessely, S., Greenberg, N., & Rubin, G. J. (2020). The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet*, 395(10227), 912-920. doi:10.1016/S0140-6736(20)30460-8
- Brown, M. B., & Forsythe, A. B. (1974). Robust tests for the equality of variances. Journal of the American Statistical Association, 69(346), 364-367.
- Carrillo-de-la-Pena, M. T., Gonzalez-Villar, A., & Trinanes, Y. (2021). Effects of the COVID-19 pandemic on chronic pain in Spain: a scoping review. *Pain Rep, 6*(1), e899. doi:10.1097/PR9.00000000000899
- Childs, J. D., Fritz, J. M., Flynn, T. W., Irrgang, J. J., Johnson, K. K., Majkowski, G. R., & Delitto, A. (2004). A clinical prediction rule to identify patients with low back pain most likely to benefit from spinal manipulation: a validation study. *Ann Intern Med, 141*(12), 920-928. doi:10.7326/0003-4819-141-12-200412210-00008

- Christian, J. B., Brouwer, E. S., Girman, C. J., Bennett, D., Davis, K. J., & Dreyer, N. A. (2020). Masking in Pragmatic Trials: Who, What, and When to Blind. *Ther Innov Regul Sci*, 54(2), 431-436. doi:10.1007/s43441-019-00073-7
- Clauw, D. J., Hauser, W., Cohen, S. P., & Fitzcharles, M. A. (2020). Considering the potential for an increase in chronic pain after the COVID-19 pandemic. *Pain*, *161*(8), 1694-1697. doi:10.1097/j.pain.00000000001950
- Cleland, J. A., Childs, J. D., Fritz, J. M., Whitman, J. M., & Eberhart, S. L. (2007). Development of a clinical prediction rule for guiding treatment of a subgroup of patients with neck pain: use of thoracic spine manipulation, exercise, and patient education. *Phys Ther*, 87(1), 9-23. doi:10.2522/ptj.20060155
- Cruz-Sánchez, E., Torres-Bonete, M., García-Pallarés, J., Gascón-Cánovas, J., Valero-Valenzuela, A., & Pereñíguez-Barranco, J. (2012). Dolor de espalda y limitación de la actividad física cotidiana en la población adulta española. Paper presented at the Anales del Sistema Sanitario de Navarra.
- Damiot, A., Pinto, A. J., Turner, J. E., & Gualano, B. (2020). Immunological Implications of Physical Inactivity among Older Adults during the COVID-19 Pandemic. *Gerontology*, 66(5), 431-438. doi:10.1159/000509216
- de Andres Ares, J., Cruces Prado, L. M., Canos Verdecho, M. A., Penide Villanueva, L., Del Valle Hoyos, M., Herdman, M., . . . Velazquez Rivera, I. (2015). Validation of the Short Form of the Brief Pain Inventory (BPI-SF) in Spanish Patients with Non-Cancer-Related Pain. *Pain Pract*, 15(7), 643-653. doi:10.1111/papr.12219
- de Zoete, A., de Boer, M. R., Rubinstein, S. M., van Tulder, M. W., Underwood, M., Hayden, J. A., . . . International, I. P. D. S. M. T. g. (2021). Moderators of the Effect of Spinal Manipulative Therapy on Pain Relief and Function in Patients with Chronic Low Back Pain: An Individual Participant Data Meta-analysis. *Spine (Phila Pa 1976), 46*(8), E505-E517. doi:10.1097/BRS.00000000003814
- Dworkin, R. H., Turk, D. C., Wyrwich, K. W., Beaton, D., Cleeland, C. S., Farrar, J. T., ... Zavisic, S. (2008). Interpreting the clinical importance of treatment outcomes in chronic pain clinical trials: IMMPACT recommendations. *J Pain*, 9(2), 105-121. doi:10.1016/j.jpain.2007.09.005
- Eccleston, C., Blyth, F. M., Dear, B. F., Fisher, E. A., Keefe, F. J., Lynch, M. E., ...
 Williams, A. C. C. (2020). Managing patients with chronic pain during the COVID-19 outbreak: considerations for the rapid introduction of remotely supported (eHealth) pain management services. *Pain*, 161(5), 889-893. doi:10.1097/j.pain.00000000001885
- Elfving, B., Andersson, T., & Grooten, W. J. (2007). Low levels of physical activity in back pain patients are associated with high levels of fear-avoidance beliefs and pain catastrophizing. *Physiother Res Int*, *12*(1), 14-24. doi:10.1002/pri.355
- Ellingsen, D. M., Napadow, V., Protsenko, E., Mawla, I., Kowalski, M. H., Swensen, D., ... Loggia, M. L. (2018). Brain Mechanisms of Anticipated Painful Movements and Their Modulation by Manual Therapy in Chronic Low Back Pain. J Pain, 19(11), 1352-1365. doi:10.1016/j.jpain.2018.05.012
- Fallon, N., Brown, C., Twiddy, H., Brian, E., Frank, B., Nurmikko, T., & Stancak, A. (2020). Adverse effects of COVID-19-related lockdown on pain, physical activity and psychological well-being in people with chronic pain. *Brit J Pain*, 1-12. doi:<u>https://doi.org/10.1177/2049463720973703</u>
- Faul, F., Erdfelder, E., Lang, A. G., & Buchner, A. (2007). G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods*, 39(2), 175-191. doi:10.3758/bf03193146

- Flink, I. K., Boersma, K., & Linton, S. J. (2010). Catastrophizing moderates the effect of exposure in vivo for back pain patients with pain-related fear. *Eur J Pain*, 14(8), 887-892. doi:10.1016/j.ejpain.2010.02.003
- Garcia-Alvarez, L., de la Fuente-Tomas, L., Garcia-Portilla, M. P., Saiz, P. A., Lacasa, C. M., Dal Santo, F., ... Bobes, J. (2020). Early psychological impact of the 2019 coronavirus disease (COVID-19) pandemic and lockdown in a large Spanish sample. J Glob Health, 10(2), 020505. doi:10.7189/jogh.10.020505
- García-Campayo, J., Zamorano, E., Ruiz, M. A., Pardo, A., Pérez-Páramo, M., López-Gómez, V., . . . Rejas, J. (2010). Cultural adaptation into Spanish of the generalized anxiety disorder-7 (GAD-7) scale as a screening tool. *Health and Quality of Life Outcomes*, 8(1), 8.
- Garner, M. J., Aker, P., Balon, J., Birmingham, M., Moher, D., Keenan, D., & Manga, P. (2007). Chiropractic care of musculoskeletal disorders in a unique population within Canadian community health centers. *J Manipulative Physiol Ther*, 30(3), 165-170. doi:10.1016/j.jmpt.2007.01.009
- Gartlehner, G., Hansen, R. A., Nissman, D., Lohr, K. N., & Carey, T. S. (2006). A simple and valid tool distinguished efficacy from effectiveness studies. *J Clin Epidemiol*, 59(10), 1040-1048. doi:10.1016/j.jclinepi.2006.01.011
- Goertz, C. M., Long, C. R., Hondras, M. A., Petri, R., Delgado, R., Lawrence, D. J., ... Meeker, W. C. (2013). Adding chiropractic manipulative therapy to standard medical care for patients with acute low back pain: results of a pragmatic randomized comparative effectiveness study. *Spine (Phila Pa 1976), 38*(8), 627-634. doi:10.1097/BRS.0b013e31827733e7
- Goertz, C. M., Long, C. R., Vining, R. D., Pohlman, K. A., Walter, J., & Coulter, I. (2018). Effect of Usual Medical Care Plus Chiropractic Care vs Usual Medical Care Alone on Pain and Disability Among US Service Members With Low Back Pain: A Comparative Effectiveness Clinical Trial. JAMA Netw Open, 1(1), e180105. doi:10.1001/jamanetworkopen.2018.0105
- Gómez-Pérez, L., López-Martínez, A. E., & Ruiz-Párraga, G. T. (2011). Psychometric properties of the Spanish version of the Tampa Scale for Kinesiophobia (TSK). *The Journal of Pain*, 12(4), 425-435.
- Hall, G., Laddu, D. R., Phillips, S. A., Lavie, C. J., & Arena, R. (2020). A tale of two pandemics: How will COVID-19 and global trends in physical inactivity and sedentary behavior affect one another? *Prog Cardiovasc Dis.* doi:10.1016/j.pcad.2020.04.005
- Hammig, O. (2019). Health risks associated with social isolation in general and in young, middle and old age. *PLoS One*, 14(7), e0219663. doi:10.1371/journal.pone.0219663
- Hayes, A. F. (2017). Introduction to mediation, moderation, and conditional process analysis: A regression-based approach: Guilford publications.
- Henriquez, J., Gonzalo-Almorox, E., Garcia-Goni, M., & Paolucci, F. (2020). The first months of the COVID-19 pandemic in Spain. *Health Policy Technol*, 9(4), 560-574. doi:10.1016/j.hlpt.2020.08.013
- Hirsh, A. T., Bockow, T. B., & Jensen, M. P. (2011). Catastrophizing, pain, and pain interference in individuals with disabilities. *Am J Phys Med Rehabil*, 90(9), 713-722. doi:10.1097/PHM.0b013e31822409b2
- Hruschak, V., Flowers, K. M., Azizoddin, D. R., Jamison, R. N., Edwards, R. R., & Schreiber, K. L. (2021). Cross-sectional study of psychosocial and pain-related variables among patients with chronic pain during a time of social distancing

imposed by the coronavirus disease 2019 pandemic. *Pain, 162*(2), 619-629. doi:10.1097/j.pain.00000000002128

- Joseph, S. J., Shoib, S., Sg, T., & Bhandari, S. S. (2020). Psychological concerns and musculoskeletal pain amidst the COVID-19 lockdown. *Open J Psychiatry Allied Sci, 11*(2), 137-139. Retrieved from <u>https://www.ncbi.nlm.nih.gov/pubmed/32775694</u>
- Kaptchuk, T. J., Hemond, C. C., & Miller, F. G. (2020). Placebos in chronic pain: evidence, theory, ethics, and use in clinical practice. *BMJ*, 370, m1668. doi:10.1136/bmj.m1668
- Keller, S., Bann, C. M., Dodd, S. L., Schein, J., Mendoza, T. R., & Cleeland, C. S. (2004). Validity of the brief pain inventory for use in documenting the outcomes of patients with noncancer pain. *Clin J Pain*, 20(5), 309-318. doi:10.1097/00002508-200409000-00005
- Lacasse, A., Pagé, M. G., Dassieu, L., Sourial, N., Janelle-Montcalm, A., Dorais, M., . . . Moor, G. (2021). Impact of the COVID-19 pandemic on the pharmacological, physical, and psychological treatments of pain: findings from the Chronic Pain & COVID-19 Pan-Canadian Study. *PAIN Reports*, 6(1), e891.
- Lazzerini, M., Barbi, E., Apicella, A., Marchetti, F., Cardinale, F., & Trobia, G. (2020).
 Delayed access or provision of care in Italy resulting from fear of COVID-19.
 Lancet Child Adolesc Health, 4(5), e10-e11. doi:10.1016/S2352-4642(20)30108-5
- Lugli, G., Ottaviani, M. M., Botta, A., Ascione, G., Bruschi, A., Cagnazzo, F., . . . Portaluri, T. (2020). The impact of the SARS-CoV-2 pandemic on healthcare provision in Italy to non-COVID patients: a systematic review. *medRxiv*.
- Luning Bergsten, C., Lundberg, M., Lindberg, P., & Elfving, B. (2012). Change in kinesiophobia and its relation to activity limitation after multidisciplinary rehabilitation in patients with chronic back pain. *Disabil Rehabil*, 34(10), 852-858. doi:10.3109/09638288.2011.624247
- Luque-Suarez, A., Martinez-Calderon, J., & Falla, D. (2019). Role of kinesiophobia on pain, disability and quality of life in people suffering from chronic musculoskeletal pain: a systematic review. Br J Sports Med, 53(9), 554-559. doi:10.1136/bjsports-2017-098673
- Mahase, E. (2020). Covid-19: WHO declares pandemic because of "alarming levels" of spread, severity, and inaction. *BMJ*, *368*, m1036. doi:10.1136/bmj.m1036
- Marshall, P. W. M., Schabrun, S., & Knox, M. F. (2017). Physical activity and the mediating effect of fear, depression, anxiety, and catastrophizing on pain related disability in people with chronic low back pain. *PLoS One*, 12(7), e0180788. doi:10.1371/journal.pone.0180788
- Minhas, Z., Ganau, M., Thakar, C., Reynolds, J., Rothenfluh, D., Bojanic, S., . . . Nnadi, C. (2020). COVID-19: new challenges, risks, and the future provision of care in spinal services. *Bone Joint J*, 102-B(6), 655-657. doi:10.1302/0301-620X.102B6.BJJ-2020-0686
- Nelson, C. F., Lawrence, D. J., Triano, J. J., Bronfort, G., Perle, S. M., Metz, R. D., . . . LaBrot, T. (2005). Chiropractic as spine care: a model for the profession. *Chiropr Osteopat*, 13, 9. doi:10.1186/1746-1340-13-9
- Nieto, R., Pardo, R., Sora, B., Feliu-Soler, A., & Luciano, J. V. (2020). Impact of COVID-19 Lockdown Measures on Spanish People with Chronic Pain: An Online Study Survey. J Clin Med, 9(11). doi:10.3390/jcm9113558
- Nunan, D., Aronson, J., & Bankhead, C. (2018). Catalogue of bias: attrition bias. *BMJ Evid Based Med*, 23(1), 21-22. doi:10.1136/ebmed-2017-110883

- Olmedilla Zafra, A., Ortega Toro, E., & Cano, L. (2013). Validation of the Pain Catastrophizing Scale in Spanish athletes. *Cuadernos de Psicología del Deporte,* 13(1), 83-93.
- Oppenheimer, D. M., Meyvis, T., & Davidenko, N. (2009). Instructional manipulation checks: Detecting satisficing to increase statistical power. *Journal of experimental social psychology*, *45*(4), 867-872.
- Ostelo, R. W., Deyo, R. A., Stratford, P., Waddell, G., Croft, P., Von Korff, M., . . . de Vet, H. C. (2008). Interpreting change scores for pain and functional status in low back pain: towards international consensus regarding minimal important change. *Spine (Phila Pa 1976), 33*(1), 90-94. doi:10.1097/BRS.0b013e31815e3a10
- Ozamiz-Etxebarria, N., Idoiaga Mondragon, N., Dosil Santamaria, M., & Picaza Gorrotxategi, M. (2020). Psychological Symptoms During the Two Stages of Lockdown in Response to the COVID-19 Outbreak: An Investigation in a Sample of Citizens in Northern Spain. *Front Psychol*, 11, 1491. doi:10.3389/fpsyg.2020.01491
- Page, M. G., Lacasse, A., Dassieu, L., Hudspith, M., Moor, G., Sutton, K., . . . Choiniere, M. (2021). A cross-sectional study of pain status and psychological distress among individuals living with chronic pain: the Chronic Pain & COVID-19 Pan-Canadian Study. *Health Promot Chronic Dis Prev Can*, 41(5). doi:10.24095/hpcdp.41.5.01
- Petrini, L., & Arendt-Nielsen, L. (2020). Understanding Pain Catastrophizing: Putting Pieces Together. *Front Psychol*, 11, 603420. doi:10.3389/fpsyg.2020.603420
- Pinto, A. J., Dunstan, D. W., Owen, N., Bonfa, E., & Gualano, B. (2020). Combating physical inactivity during the COVID-19 pandemic. *Nat Rev Rheumatol*, 16(7), 347-348. doi:10.1038/s41584-020-0427-z
- Prater, C., Tepe, M., & Battaglia, P. (2020). Integrating a Multidisciplinary Pain Team and Chiropractic Care in a Community Health Center: An Observational Study of Managing Chronic Spinal Pain. J Prim Care Community Health, 11, 2150132720953680. doi:10.1177/2150132720953680
- Rodriguez-Rey, R., Garrido-Hernansaiz, H., & Collado, S. (2020). Psychological Impact and Associated Factors During the Initial Stage of the Coronavirus (COVID-19) Pandemic Among the General Population in Spain. *Front Psychol*, 11, 1540. doi:10.3389/fpsyg.2020.01540
- Rowbotham, M. C. (2001). What is a "clinically meaningful" reduction in pain? *Pain*, 94(2), 131-132. doi:10.1016/S0304-3959(01)00371-2
- Sallis, J. F., Adlakha, D., Oyeyemi, A., & Salvo, D. (2020). An international physical activity and public health research agenda to inform coronavirus disease-2019 policies and practices. J Sport Health Sci, 9(4), 328-334. doi:10.1016/j.jshs.2020.05.005
- Sallis, R., Young, D. R., Tartof, S. Y., Sallis, J. F., Sall, J., Li, Q., ... Cohen, D. A. (2021). Physical inactivity is associated with a higher risk for severe COVID-19 outcomes: a study in 48 440 adult patients. *Br J Sports Med.* doi:10.1136/bjsports-2021-104080
- Schellingerhout, J. M., Verhagen, A. P., Heymans, M. W., Pool, J. J. M., Vonk, F., Koes, B. W., & de Vet, H. C. W. (2008). Which subgroups of patients with non-specific neck pain are more likely to benefit from spinal manipulation therapy, physiotherapy, or usual care? *Pain*, 139(3), 670-680. doi:10.1016/j.pain.2008.07.015
- Spain, Government of. (2020a). Orden SND/387/2020, de 3 de mayo, por la que se regula el proceso de cogobernanza con las comunidades autónomas y ciudades de Ceuta

y Melilla para la transición a una nueva normalidad. Boletin Oficial del Estado -State's Official Bulletin: Government of Spain Retrieved from https://www.boe.es/eli/es/o/2020/05/03/snd387

- Spain, Government of. (2020b). Real Decreto 463/2020, de 14 de marzo, por el que se declara el estado de alarma para la gestión de la situación de crisis sanitaria ocasionada por el COVID-19. Boletin Oficial del Estado State's Official Bulletin: Government of Spain Retrieved from https://www.boe.es/buscar/pdf/2020/BOE-A-2020-3692-consolidado.pdf
- Stamatakis, E., & Bull, F. C. (2020). Putting physical activity in the 'must-do' list of the global agenda. Br J Sports Med, 54(24), 1445-1446. doi:10.1136/bjsports-2020-103509
- Sturgeon, J. A., & Zautra, A. J. (2013). Psychological resilience, pain catastrophizing, and positive emotions: perspectives on comprehensive modeling of individual pain adaptation. *Curr Pain Headache Rep, 17*(3), 317. doi:10.1007/s11916-012-0317-4
- Sullivan, M. J., Thorn, B., Haythornthwaite, J. A., Keefe, F., Martin, M., Bradley, L. A., & Lefebvre, J. C. (2001). Theoretical perspectives on the relation between catastrophizing and pain. *Clin J Pain*, 17(1), 52-64. doi:10.1097/00002508-200103000-00008
- Sullivan, M. J., Tripp, D. A., & Santor, D. (2000). Gender differences in pain and pain behavior: the role of catastrophizing. *Cognitive Therapy and Research*, 24(1), 121-134.
- Tanne, J. H., Hayasaki, E., Zastrow, M., Pulla, P., Smith, P., & Rada, A. G. (2020). Covid-19: how doctors and healthcare systems are tackling coronavirus worldwide. *BMJ*, 368, m1090. doi:10.1136/bmj.m1090
- Taylor, S., Landry, C. A., Paluszek, M. M., Fergus, T. A., McKay, D., & Asmundson, G. J. G. (2020). COVID stress syndrome: Concept, structure, and correlates. *Depress Anxiety*, 37(8), 706-714. doi:10.1002/da.23071
- Thacker, M. M., M. (2020). Staring Down the Barrel of a Gun: A Potential Pain Pandemic Following COVID-19 Infection. Pain and Rehabilitation-the Journal of Physiotherapy Pain Association, 2020(49), 1-8.
- Tison, G. H., Avram, R., Kuhar, P., Abreau, S., Marcus, G. M., Pletcher, M. J., & Olgin, J. E. (2020). Worldwide Effect of COVID-19 on Physical Activity: A Descriptive Study. Ann Intern Med, 173(9), 767-770. doi:10.7326/M20-2665
- Verhagen, A. P., Karels, C. H., Schellingerhout, J. M., Willemsen, S. P., Koes, B. W., & Bierma-Zeinstra, S. M. (2010). Pain severity and catastrophising modify treatment success in neck pain patients in primary care. *Man Ther*, 15(3), 267-272. doi:10.1016/j.math.2010.01.005
- Wertli, M. M., Burgstaller, J. M., Weiser, S., Steurer, J., Kofmehl, R., & Held, U. (2014a). Influence of catastrophizing on treatment outcome in patients with nonspecific low back pain: a systematic review. *Spine (Phila Pa 1976), 39*(3), 263-273. doi:10.1097/BRS.00000000000110
- Wertli, M. M., Rasmussen-Barr, E., Held, U., Weiser, S., Bachmann, L. M., & Brunner, F. (2014b). Fear-avoidance beliefs-a moderator of treatment efficacy in patients with low back pain: a systematic review. *Spine J, 14*(11), 2658-2678. doi:10.1016/j.spinee.2014.02.033
- Williams, N. H., Hendry, M., Lewis, R., Russell, I., Westmoreland, A., & Wilkinson, C. (2007). Psychological response in spinal manipulation (PRISM): a systematic review of psychological outcomes in randomised controlled trials. *Complement Ther Med*, 15(4), 271-283. doi:10.1016/j.ctim.2007.01.008

- Yoshimoto, T., Fujii, T., Oka, H., Kasahara, S., Kawamata, K., & Matsudaira, K. (2021). Pain Status and Its Association with Physical Activity, Psychological Stress, and Telework among Japanese Workers with Pain during the COVID-19 Pandemic. Int J Environ Res Public Health, 18(11). doi:10.3390/ijerph18115595
- Zhaoyang, R., Martire, L. M., & Darnall, B. D. (2020). Daily pain catastrophizing predicts less physical activity and more sedentary behavior in older adults with osteoarthritis. *Pain, 161*(11), 2603-2610. doi:10.1097/j.pain.00000000001959

Figure legends

Figure 1. Trial Protocol

Schema of the study protocol. Participants received the link to participate in the initial and follow-up questionnaires on their phone or via e-mail. Subsequently, according to the care received, they would be placed in one of three groups. A flow-chart based on the CONSORT recommendations is available as supplemental material.

Figure 2. Pain intensity ratings at baseline and follow-up by group

Pain intensity reported at baseline and follow-up on a numerical rating scale from 0 to 10 in Groups 0, 1 and 2. Between-group and within-group statistical significance obtained via planned comparisons of mixed ANOVAs.

p = p < 0.05, p = p < 0.01, p = p < 0.001.

Figure 3. Pain interference at baseline and follow-up by group

Pain interference reported at baseline and follow-up from 0 to 60 according to the brief pain inventory in Groups 0, 1 and 2. Between-group and within-group statistical significance obtained via planned comparisons of mixed ANOVAs.

p = p < 0.05, p = p < 0.01, p = p < 0.001.

Figure 4. Fear of movement at baseline and follow-up by group

Scores for the short version of the Tampa Scale of Kinesiophobia reported at baseline and follow-up from 11 to 44 in Groups 0, 1 and 2. Between-group and within-group statistical significance obtained via planned comparisons of mixed ANOVAs.

$$* = p < 0.05, ** = p < 0.01, *** = p < 0.001.$$

Figure 5. Mediation effects by pain catastrophizing

Mediatory effect of pain catastrophizing (PCS) on the relationship between changes in pain intensity ratings and in pain interference (A), moderating role of pain ratings at baseline in the relationship between pain catastrophizing and change in pain ratings (B). For patients with lower pain intensity at baseline (3/10, red line), low levels of PCS were associated with larger reductions in pain intensity. For patients with higher baseline pain intensity (~7/10, blue line), higher PCS scores were associated with pain increases.

Mediatory effects of pain catastrophizing in the relationship between fear of movement (TSK) and changes in pain intensity ratings in women (C) and men (D).

* = p < 0.05; ** = p < 0.01*.

Supplemental figure legends

Figure S1. Flowchart of participant inclusion and participations

Figure S2. Heatmap of correlations

Heatmap of correlations between all numerical coprimary (Pain intensity, Pain interference, TSK = Tampa scale of Kinesiophobia, PCS = Pain Catastrophizing Scale) and secondary outcomes (GAD = General Anxiety Disorder scale, emotions evoked by the pandemic: Sadness, Worry, Loneliness, Anger, Helplessness, Anxiety, Surprise, Relief, Hope, Stress_COVID = Stress caused by the pandemic, Stress_Restrictions = Stress caused by the restrictions, Stress_Health = Stress about one's health) at baseline (Pre) and follow-up (Post). The right *y*-axis provides a key for the correlation coefficients for each two variables (from -1.0 to 1.0). The statistical significance of these correlations was not considered, instead, variables with an $r \ge 0.5$ were considered to confirm hypotheses during the mediation analyses.