UNIVERSITY^{OF} BIRMINGHAM University of Birmingham Research at Birmingham

Best response dynamics on random graphs

Chellig, Jordan; Durbac, Calina; Fountoulakis, Nikolaos

DOI: 10.1016/j.geb.2021.11.003

License: Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version Peer reviewed version

Citation for published version (Harvard):

Chellig, J, Durbac, C & Fountoulakis, N 2022, 'Best response dynamics on random graphs', Games and Economic Behaviour, vol. 131, pp. 141-170. https://doi.org/10.1016/j.geb.2021.11.003

Link to publication on Research at Birmingham portal

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)

•Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

BEST RESPONSE DYNAMICS ON RANDOM GRAPHS

JORDAN CHELLIG, CALINA DURBAC, AND NIKOLAOS FOUNTOULAKIS

ABSTRACT. We consider evolutionary games on a population whose underlying topology of interactions is determined by a binomial random graph G(n, p). Our focus is on 2player symmetric games with 2 strategies played between the incident members of such a population. Players update their strategies synchronously: each player selects the strategy that is the best response to the current set of strategies its neighbours play. We show that such a system reduces to generalised majority and minority dynamics. We show rapid convergence to unanimity for p in a range that depends on a certain characteristic of the payoff matrix. In the presence of a bias among the pure Nash equilibria, we determine a sharp threshold on p above which the largest connected component reaches unanimity with high probability. For p below this critical value, we identify those substructures inside the largest component that block unanimity.

1. INTRODUCTION

The need for a dynamic game theory was pointed out by von Neumann and Morgenstern in their seminal book [37] which set the foundations of modern game theory. Research on dynamic games on populations was stimulated by settings in evolutionary biology. In 1973 John Maynard Smith and George Price [28] set the foundations of *evolutionary game theory*, trying to explain phenomena that arise in animal fighting, which contradict the traditional Darwinian theory.

Maynard Smith and Price [28] considered a variation of the well-known hawk-dove game (see below for a precise description of its payoff matrix) with more than two strategies, which range from purely altruistic to hawkish behaviour. They argued that altruistic behaviour in such a population may have higher payoff compared to the payoff in a pure

E-mail addresses: jac555@student.bham.ac.uk, cmd639@bham.ac.uk, n.fountoulakis@bham.ac.uk. 2010 *Mathematics Subject Classification*. Primary 05C80, 91A22; Secondary 91A05.

Key words and phrases. random graphs, evolutionary games, unanimity.

School of Mathematics, University of Birmingham B15 2TT, Birmingham, United King-Dom.

School of Mathematics, University of Birmingham B15 2TT, Birmingham, United Kingdom.

School of Mathematics, University of Birmingham B15 2TT, Birmingham, United Kingdom.

This research started during the Undergraduate Research Bursary 17-18 91 (for Calina Durbac under the supervision of Nikolaos Fountoulakis) funded by the London Mathematical Society and the School of Mathematics in Birmingham. Jordan Chellig is supported by an EPSRC scholarship. Nikolaos Fountoulakis has also been partially supported by an Alan Turing Fellowship EP/N5510129/1 and the EPRSC grant EP/P026729/1.

population consisting only of hawks. Introducing the notion of an *evolutionary stable strategy*, they argued that natural selection would favour the co-existence of behaviours in a given population. In this setting, every member of the population may interact with any other member. That is, in graph-theoretic terms, this is an evolutionary system on a complete graph.

Later on, Nowak and May [33] considered settings with non-trivial underlying topology. They considered a population of agents that are located on the vertices of a 2-dimensional lattice, in which every agent interacts only with its four neighbours. The interaction is that any two adjacent agents play the prisoner's dilemma (which we describe below) in which they may *cooperate* or *defect*. The dynamics that was considered there depends on the total payoff of each agent that is accumulated by the four games it plays with its neighbours (or three or two, if it is located on the boundary of the lattice). The agents update their strategies synchronously, with each agent adopting the strategy of a neighbour who had the largest total payoff in the previous round. The main observation in [33] is the *co-existence* of the two strategies in the long term, despite the fact that the defect strategy is a *Nash equilibrium* for this game: that is, in a game between two players who both defect none has interest in cooperating, given that the other does not. Similar dynamics were studied by Santos and Pacheko [36] in the preferential attachment model, which typically yields graphs that have some properties of complex networks.

In this paper, we consider a class of dynamics of games on networks in which agents seek to choose a *local dominating strategy*. In this process, every agent switches their strategy to the one which maximises their own payoff, given the strategies of its opponents. In other words, in each round each agent tries to give their *best response* to the current choice of strategies of their opponents. This dynamics was first considered by Gilboa and Matsui [18] in continuous-time setting and for a population with no underlying topology, where everyone interacts with everyone else. Gilboa and Matsui showed the existence of *cyclically stable sets*. Roughly speaking, these are sets of configurations of the population in which the best response dynamics is trapped.

We consider a discrete setting and study the evolution of best response dynamics on a random graph. Adjacent agents are able to interact with each other via a symmetric game with 2 strategies, on a round-by-round basis. In a given round, each agent will choose one of the two (pure) strategies, which they will use against all of their neighbours. For each game played, the agent will receive a payoff determined by the corresponding entry in the payoff matrix; subsequently, each agent will also receive a total payoff, which is the sum of all of the individual payoffs that the agent received in that round. In the next round, agents will choose to execute the strategy which would have given them the largest payoff in the previous round. In essence, each agent is focused on maximising their own total payoff by utilising information gained from the previous round. Our central result concerns the rapid rate at which agents will settle into a global unanimous strategy on a typical sample of the underlying random graph.

In our analysis, we consider arbitrary 2-player symmetric¹ games and we show that best response dynamics reduce to generalised majority or minority dynamics. These terms refer to a general class of discrete processes on graphs where vertices have two states and at each round a vertex adopts the state of the majority or the minority of its neighbourhood, respectively. Majority games reward the individuals who follow strategies which are in line with the popular opinion; such games are clearly co-operative, and thus agents will tend to form large unanimous coalitions; see [20]. On the other hand, minority games capture the idea that agents will benefit from making choices that oppose the popular consensus. The idea of a minority game was introduced to characterise the behaviour of agents within the *El Farol Bar problem*. The problem describes a fixed population who will repeatedly attempt to synchronously choose their favourite evening location; however, only the individuals who choose less crowded locations will be rewarded [11, 13]. This problem captures the underlying tension of minority games. Choosing the lucrative option will clearly reward agents with the greatest payoff, however this payoff is easily spoiled if too many of the participants think alike. If agents adopt this line of thinking, then we deduce that the entire population will reject the optimal payoff, regardless of the fact that the most lucrative option will tend to be uncontested. In the case of the El Farol Bar problem, any deterministic strategy will ultimately fail to satisfy anyone.

While the study of evolutionary games on populations with arbitrary mutual interactions is well established, a new focus has now been given to systems of agents which possess an underlying topology. Within these systems, agents are only able to interact with their topological neighbours and not necessarily with everyone. The topology of these interactions is commonly represented by means of an underlying network. Consideration of an underlying topology allows us to analyse how local decisions in the system can cascade out to form a global consensus. In the context of best response dynamics Lelarge [26] shows that a small set of agents which oppose the current consensus can cause a large contagion of opposing opinions to spread throughout the network. Therein, a coordination mechanism described by Morris [30] is considered on a population whose topology is that of a random graph with a given degree sequence. As we shall see shortly, this coordination mechanism falls under the framework of generalised majority dynamics. Furthermore, all agents update their strategies synchronously. Lelarge [26] gives a condition for the parameter of the coordination game, which determines whether contagion occurs with high probability. Here, the term "contagion" does not necessarily mean complete domination of the invading strategy, but adoption of this strategy by a positive fraction of the population.

An asynchronous setting has been considered by Kreindler and Young [25] as well as by Arieli et al. [3]. Kreindler and Young [25] consider the setting where agents update their strategies at random times (following a Poisson clock independently of any other agent) according to a stochastic best response rule. The update procedure takes place with some probability that may be bounded away from 0 and 1. To this end, they use the logit function. The underlying topology is that of a regular graph which they go on to generalise to a graph with a given degree sequence. They show that the expected time

¹Here, the term "symmetric" means that the roles of the two players are interchangable.

until at least half of the agents adopt the invading strategy is bounded by a constant. This has been generalised by Arieli et al. [3] where arbitrary response functions are considered and furthermore the underlying network may be evolving.

Other classes of mechanisms which model or represent the spread of an invading strategy (or an innovation) within a network of agents have also been studied. However, these have non-game-theoretic flavour. Such is the class of *bootstrap percolation processes* where the agents synchronously adopt a certain strategy, if this has been adopted by a certain number of their neighbouring agents. Also, there are variations of these processes where a strategy is adopted if a certain fraction of the neighbouring agents have adopted it. Such processes have their origins in statistical physics [12], but have been studied in a variety of networks (random and non-random) - see for example [2, 5-7, 15, 22, 24, 29]. A quantity that is of interest in this context is the final size of the set of agents who have adopted this strategy. Note that these mechanisms are monotone in the sense that once the particular strategy has been adopted it cannot change. Another mechanism of non-game-theoretic flavour which is non-monotone is the *voter model*. Here, the update of each agent's strategy takes place asynchronously. At these random times each agent adopts the strategy of one of its neighbours that has been randomly selected according to some distribution. This model was introduced by Holley and Liggett in the context of interacting particle systems on an infinite setting [21, 27]. Bounds on the expected time to consensus on some finite (random and non-random) graphs results were derived by Yildiz et al. [38] in the case where the choice is made uniformly among the neighbouring agents.

The underlying topology which is the main focus of our work is that of a binomial random graph G(n, p). This classic model has been central in the study of random graphs and was first considered by Gilbert [17]. In this model, on a set of n vertices each edge appears independently with probability p = p(n). We shall consider the evolution of best response dynamics on G(n, p) for an arbitrary symmetric game with two players played on the vertex set of a G(n, p) random graph, starting with a random assignment of strategies. As we shall see in our analysis, best response dynamics will reduce to the analysis of generalised majority or minority dynamics on G(n, p).

Our results show that in a wide range of densities best response dynamics stabilises rapidly and one of the strategies dominate within the population entirely or almost entirely. We prove a general result which shows that this is achieved in at most 4 rounds when $np = \Omega(n^{1/2})$. However, if the game exhibits a certain form of bias in terms of its pure Nash equilibria, then we get very precise results on how this dynamics evolves for p such that $np \gg 1$. Let us recall that in this range G(n, p) may not be connected. However, typically its largest connected component contains almost all vertices, whereas each one of the other components is a tree and has order at most log n. (The books [10,23] provide a detailed description of the typical structure and the evolution of G(n, p).) For p such that G(n, p) is not connected with high probability, it is the largest connected component we focus our study on. In the presence of such a bias, we determine a sharp threshold on pabove which the largest connected component reaches consensus among its vertices. For p below this critical value, we identify those substructures inside the largest component that are in disagreement with the majority of the vertices therein. Hence, we are able to characterise the co-existence of strategies very precisely.

We layout our paper as follows: In Section 2 we provide key definitions for our analysis and define the *interacting node system* on a graph. In Section 3, we consider our first approaches to analysing the model and state our two main results. The first concerns the formation of a consensus when the underlying networks is both random and suitably dense. The second result shows that a consensus can be achieved in a sparser regime of G(n, p), given that there exists a certain form of bias among the pure Nash equilibria of the game. This is analysed in Section 4. In this case, we identify a sharp threshold for p above which consensus is reached in the largest component of G(n, p) in at most $\beta \log n$ rounds, for a positive constant β , where $\log \cdot$ is the natural logarithm. This critical value is determined by the payoff matrix and is below the well-known connectivity threshold of G(n, p) which is $\log n/n$. Furthermore, we show that if $np > c \log n$, for some c > 1 which depends on the parameters of the system, then in fact the system reaches consensus only after one round.

In Section 5, we consider games where this bias is no longer present. In this scenario, we observe that the interacting node system reduces to the so-called *majority or minority dynamics* on G(n, p). We sample a selection of results from [16], which concern the rapid stabilisation of agent strategies when playing a majority game on a dense random graph. Using these results as a basis, we proceed to prove our result concerning the formation of consensus strategies in the minority game analogue of the random graph majority game. By combining the above results, we may readily deduce that for any 2×2 real-valued payoff matrix and a suitably dense random graph, the agents of the interacting node system will reach a consensus after at most four rounds, with high probability.

2. Iterative Games on Graphs

Let $Q = (q_{i,j})$ be a 2 × 2 matrix with entries in \mathbb{R} ; we refer to Q as the *payoff matrix*. (The rows and columns of Q are indexed by $\{0, 1\}$, which are assumed to represent the strategies.) Each player will now choose a strategy from $\{0, 1\}$. Player 1 will then receive a payoff given by $q_{i,j}$, where i is the strategy chosen by Player 1, and j is the strategy chosen by Player 2. Analogously, Player 2 will receive a payoff $q_{i,i}$.

The interactions of the agents/players are represented by a fixed graph G = (V, E). The nodes of G represent the agents, and if two nodes are adjacent, then the corresponding agents interact with each other by playing the game with payoff matrix Q. We refer to this process as an *interacting node system* (G, Q, S), which we detail formally as follows. We fix a graph G, a payoff matrix Q, and for every $v \in V(G)$ an initial vertex strategy S(v)where $S : V(G) \to \{0, 1\}$. We consider a discrete time process. For each $t \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}$, we denote by $S_t(v)$ the strategy played by vertex v at step t. Thus, $S_t : V(G) \to \{0, 1\}$. Let $N_G(v)$ denote the set of neighbours of a vertex $v \in V(G)$ in G. For a vertex $v \in V(G)$, a step $t \in \mathbb{N}_0$ and $j \in \{0, 1\}$, we set $n_t(v; j) = |N_G(v) \cap \{u : S_t(u) = j\}|$. To each vertex $v \in V(G)$, we assign the initial state $S_0(v) = S(v)$. To progress from round t to round t + 1, the following evolution rule is applied: Each vertex v will play the game against each one of its neighbours, playing strategy $S_t(v)$. For each game played, the vertex will receive a payoff given by the corresponding entry in the payoff matrix Q. We denote the total payoff for a vertex v at time step t to be the sum of all payoffs that v received in that round. Therefore, for a vertex v with $S_t(v) = i$, we define the total payoff of v at time t as $T_t(v) := n_t(v; 0)q_{i,0} + n_t(v; 1)q_{i,1}$. We define the alternative payoff of v at time t, as the total payoff the vertex would have received, had they played the other available strategy. Hence, the alternative payoff for a vertex v with $S_t(v) = i$ is given as: $T'_t(v) := n_t(v; 0)q_{1-i,0} + n_t(v; 1)q_{1-i,1}$. The value of $S_{t+1}(v)$ is determined by comparing the values of the current payoff and the alternative payoff. If the alternative payoff is strictly greater than the total payoff, then the vertex will switch to strategy 1 - i in round t + 1; otherwise it will continue with its current strategy. For a vertex v at time t, we can succinctly express the evolution rule as follows:

$$S_{t+1}(v) = \begin{cases} S_t(v) & \text{if } T_t(v) \ge T'_t(v); \\ 1 - S_t(v) & \text{if } T_t(v) < T'_t(v). \end{cases}$$

We wish to analyse the global evolution of the strategies as time elapses. We say that a vertex's strategy is *periodic* if there exist some $T, p \in \mathbb{N}$, such that $S_T(v) = S_{T+kp}$ for all $k \in \mathbb{N}$. We call the least possible p which satisfies this definition the *period*. We say that the evolution of a system is *unanimous*, if there exists some $T \in \mathbb{N}$ such that for all $t \geq T$ and all distinct pairs of vertices $u, v \in V(G)$, we have that $S_t(u) = S_t(v)$. We say that the system is *stable* if there exists $T \in \mathbb{N}$ such that for all $v \in V(G)$ and $t \geq T$, we have that $S_t(v) = S_T(v)$.

3. Main results

The evolution of the node system is largely governed by the entries present in the payoff matrix Q. As a consequence of choosing specific entries for Q, it can be the case that one strategy is strictly more beneficial than the other. In this case the evolution of the system is trivial, as all agents will unanimously maximise their payoff; we capture this behaviour with the notion of degeneracy. We say that a payoff matrix $Q = (q_{i,j})$ with $i, j \in \{0, 1\}$ is non-degenerate, if one of the following holds:

- (i) We have that $q_{0,0} > q_{1,0}$ and $q_{0,1} < q_{1,1}$.
- (ii) Or, $q_{0,0} < q_{1,0}$ and $q_{0,1} > q_{1,1}$.

Otherwise, we say that Q is *degenerate*. The behaviour of interacting node systems with degenerate payoff matrices will be discussed in Section 3.2.

We consider an interacting node system where the underlying graph G is both random and suitably dense. We let G(n, p) be the *binomial random graph* on vertex set $V_n := [n] :=$ $\{1, 2, ..., n\}$, where each edge appears independently with probability p. We introduce a random binomial initial state, which we refer to as $S_{1/2} \in \{0, 1\}^{[n]}$. For every vertex $v \in V_n$, we have that $\mathbb{P}(S_{1/2}(v) = 1) = \mathbb{P}(S_{1/2}(v) = 0) = 1/2$, independently of any other vertex. We say that a sequence of events E_n defined on a sequence of probability spaces with probability measure \mathbb{P}_n occurs asymptotically almost surely (or a.a.s.) if $\mathbb{P}_n(E_n) \to 1$ as $n \to \infty$. We now state our first result which describes the vertex strategies of an interacting node system on G(n, p), with initial state $S_{1/2}$. We denote this system by $(G(n, p), Q, S_{1/2})$.

Theorem 3.1. Let Q be a 2 × 2 non-degenerate payoff matrix. For any $\varepsilon \in (0,1)$ there exist positive constants Λ, n_0 such that for all $n \ge n_0$, if $p > \Lambda n^{-\frac{1}{2}}$, then with probability at least $1 - \varepsilon$, across the product space of G(n, p) and $S_{1/2}$, the interacting node system $(G(n, p), Q, S_{1/2})$ will be unanimous after at most four rounds.

Proof sketch. Suppose we have a node system \mathcal{I} , with payoff matrix Q. We introduce the following quantity, which we refer to as the payoff skew, $\lambda = \lambda(Q) := (q_{1,1} - q_{0,1}) / (q_{0,0} - q_{1,0}).$ We do a case analysis, depending on whether or not λ is equal to one. In the case $\lambda \neq 1$ we can deduce that a.a.s all vertices will choose the same strategy after the first round. Indeed, consider a vertex and expose its neighbourhood. By the Chernoff bound (3) we can show that the ratio $n_0(v;0)/n_0(v;1)$ can be made arbitrarily close to one, for n large enough. The value of λ then imposes a bias which causes the vertex to switch to a specific strategy. It can then be shown that this applies to all vertices in the network. This argument is captured by Lemma 4.8, the proof of which can be found in Section 4.3. The case of $\lambda = 1$ is considered in Section 5. We show that the node system reduces to majority or minority dynamics, depending on the form of the matrix. In the case of majority dynamics, the theorem follows from Theorem 5.1, which states that with high probability majority dynamics achieves unanimity within four rounds. This was proved in [16]. Thereafter, we show the analogous result (Theorem 5.2) for minority dynamics. The idea here is that for a given minority game, we construct a majority game which approximates the behaviour of the vertices in the minority game. We show that there is a discordant set of vertices which resist this approximation. Nevertheless, this set is small enough so as to not affect the evolution of the system. So, essentially the system evolves as in majority dynamics and reaches unanimity within four rounds.

3.1. The skew of the payoff matrix Q. Suppose a vertex v at time t has picked a strategy $S_t(v) = i$. From the above discussion, we observe that the *incentive* for v to switch strategy is the condition that $T_t(v) < T'_t(v)$. Expanding both terms and re-arranging, we observe the condition for a vertex playing strategy i to switch to strategy 1-i is as follows:

$$n_t(v; 0) (q_{i,0} - q_{1-i,0}) < n_t(v; 1) (q_{1-i,1} - q_{i,1}).$$

If Q is non-degenerate, there are two possible cases: Either $q_{0,0} > q_{1,0}$ and $q_{1,1} > q_{0,1}$; or we have that $q_{0,0} < q_{1,0}$ and $q_{0,1} > q_{1,1}$.

We lead with the former case. By substituting values of $i \in \{0, 1\}$, we can rephrase the evolution conditions for each agent in terms of $n_t(v; 0)$ and $n_t(v; 1)$. We recall the definition of the *payoff skew* of the payoff matrix Q:

$$\lambda = \lambda(Q) := (q_{1,1} - q_{0,1}) / (q_{0,0} - q_{1,0}).$$

We remark that $\lambda(Q)$ is positive and well-defined, if and only if, Q is non-degenerate. If $S_t(v) = i$, then we can write:

$$S_{t+1}(v) = \begin{cases} 1-i & \text{if } n_t(v;i) < \lambda^{1-2i} n_t\left(v;1-i\right) \\ i & \text{otherwise} \end{cases}$$
(1)

Suppose now that the latter case holds, that is, $q_{0,0} < q_{1,0}$ and $q_{0,1} > q_{1,1}$. Then if $S_t(v) = i$,

$$S_{t+1}(v) = \begin{cases} 1-i & \text{if } n_t(v;i) > \lambda^{1-2i} n_t\left(v;1-i\right) \\ i & \text{otherwise} \end{cases}$$
(2)

We refer to the system governed by the evolution rules from (1) as the majority regime; while we refer to the system described in (2) as the minority regime. We will tackle each of these systems separately. The intuition for these names arises from how each individual agent tends to think of its neighbours. In the majority regime, agents will tend to follow the strategies which are shared by the majority of their neighbours; however, in the minority regime agents will generally choose the least popular strategy seen across their neighbours. The value of λ determines the strength of this tendency.

In game-theoretic terms, it is easy to see that payoff matrices in the majority regime give rise to two pure Nash equilibria: namely with both players simultaneously playing strategy one or strategy zero. On the other hand, payoff matrices in the minority regime have two pure Nash equilibria: those in which the two players play opposite strategies. (In both regimes, there is also a mixed Nash equilibrium.) Hence, games of the former type are sometimes called *coordination games*, whereas those of the latter type are called *anticoordination games*. From this point of view, the parameter λ can be seen as some form of bias between the two pure Nash equilibria.

Examples. We give two examples of games that are well studied in game theory and reflect the above classification.

1. Stag-Hare game In this game, two hunters need to decide whether to co-operate on hunting or not. There two possible species to hunt: a stag or a hare. However, hunting a stag is difficult for each one on their own to catch, so they can only do this if they work together. Nevertheless, each one of them can hunt on their own, but then it will only be possible to hunt for a hare, which offers a smaller reward. This setting can be described in the following payoff matrix $Q_{\rm S-H}$ in which the first column and row correspond to the stag strategy whereas the second column and row correspond to the hare strategy:

$$Q_{\mathrm{S-H}} = \left[\begin{array}{cc} S & 0\\ H & H/2 \end{array} \right],$$

where H < S. Here, S is the reward they both earn if they cooperate and go for a stag. If one goes for a stag but the other goes for a hare, the former will get no reward, as it is impossible to hunt a stag on their own, but the other will get some reward H < S which comes from catching a hare. Now, if they both go for a hare, they will both receive a smaller reward, H/2, as they will split what they will catch. This is an example of a coordination game with two pure Nash equilibria, namely when both players have a common strategy. This belongs to the majority regime.

2. The Hawk-Dove game. Here, a population of animals consists of two types of individuals which are differentiated by the amount of aggression they display during their interactions. There is the most aggressive type (hawk) and the least aggressive, or most cooperative, type (dove). When two of them interact over some fixed resource, the outcome depends on their types. If two hawks compete over the resource they get injured, because of the fighting, and one of them (at random) manages to grab the resource. Hence, if R is the gain from the resource, each is expected to gain R/2 out of the fight, but they pay a price P > R for their injuries, whereby their overall gain is (R - P)/2. If a hawk interacts with a dove, then the hawk grabs the resource gaining R, whereas the dove walks away with nothing. Finally, if two doves interact, then they share the resource each gaining R/2. Thus, if $Q_{\rm H-D}$ denotes the payoff matrix of the Hawk-and-Dove game, this is

$$Q_{\rm H-D} = \begin{bmatrix} \frac{R-P}{2} & R\\ 0 & \frac{R}{2} \end{bmatrix},$$

where first row and column correspond to the hawk strategy and the second row and column correspond to the dove strategy. Hence, this is a non-degenerate payoff matrix in the minority regime with

$$\lambda(Q_{\rm H-D}) = \frac{R}{P-R}.$$

3. The Prisoners dilemma. In this game, two individuals are arrested while committing a crime together, but they are put in different cells. The police do not have enough evidence to convict them, but they make an offer to each one separately. If one of them confesses (defects) but the other remains silent (cooperates), then the former is released but the other is sentenced to imprisonment of P > 0 years. If they both confess, they are sentenced to R > 0 years of imprisonment, for R < P. Finally, if both remain silent, they are both sentenced to S years imprisonment for some minor offence, as the police do not have enough evidence. In this case, S < R.

Thus, 0 < S < R < P. The payoff matrix of the Prisoners dilemma game $Q_{\rm PD}$ is

$$Q_{\rm PD} = \left[\begin{array}{cc} -R & 0\\ -P & -S \end{array} \right],$$

where the first row and column correspond to the Defect strategy and the second row and column correspond to the Cooperate strategy. (Here, we put the - sign in front of these quantities, as imprisonment is thought of as a loss.) Hence, this is a case of a degenerate payoff matrix.

Our second result concerns non-degenerate interacting node systems in a sparser regime compared to that covered in Theorem 3.1. In this setting, we assume that $\lambda \neq 1$ and we are able to consider smaller values of p, below the *connectivity threshold* of G(n, p). Among the first results on the theory of random graphs due to Gilbert [17] is that the connectivity of G(n, p) undergoes a sharp transition. More specifically, let $\omega : \mathbb{N} \to \mathbb{R}_+$ be a function such that $\omega(n) \to \infty$ as $n \to \infty$; if $np = \log n + \omega(n)$, then a.a.s. G(n, p) is connected, whereas if $np = \log n - \omega(n)$, then a.a.s. G(n, p) has at least one isolated vertex. Analogous results were proved by Erdős and Rényi [14] on the uniform model of random graphs with a given number of edges.

In the subcritical regime, we cannot hope for unanimity. For $1 \ll np < \log n$, all connected components of G(n, p) besides the largest one are trees of order at most $\log n$ and, in fact, many of them are isolated vertices (see for example [10] for a complete and precise description of the structure of G(n, p)). However, a.a.s. the largest connected component contains most vertices of G(n, p), when $np \gg 1$. So, if we want to make a statement about almost all vertices of G(n, p) it suffices to focus on its largest component. This is what lies behind the next theorem, which states that almost all vertices of G(n, p) reach uninanimity after a logarithmic (in n) number of rounds. Furthermore, for certain densities not much above the aforementioned connectivity threshold, we are able to show that the process reaches complete unanimity after a single round.

Theorem 3.2. Let $p = d/n \leq 1$, where $d \gg 1$, and let Q be a 2×2 non-degenerate payoff matrix. Suppose that $(G(n, p), Q, S_{1/2})$ is an interacting node system with payoff skew $\lambda \neq 1$. For any $\varepsilon > 0$ there exists $\beta = \beta(\lambda, \varepsilon) > 0$ such that a.a.s. at least $n(1 - \varepsilon)$ vertices in G(n, d/n) will be unanimous after at most $\beta \log n$ rounds. Moreover, there exists a constant $\alpha(\lambda) > 1$ such that if $d > \alpha(\lambda) \log n$, then a.a.s. G(n, d/n) will be unanimous after one round.

Proof sketch. The part of the above theorem for $np > c \log n$ with $c > \alpha(\lambda)$ implies and, in fact, it is much stronger than Theorem 3.1. That is, for $\lambda \neq 1$ the system goes into unanimity in one round for much lower densities than $n^{-1/2}$. The idea is to apply Lemma 4.8 from Section 4.3 in the first round. For even smaller densities, this lemma implies that the number of vertices following the minority strategy after one round is at most sublinear. We then consider the set of vertices in the giant component that do not have small degree; these vertices comprise of almost all of the giant component. We show that amongst these vertices, the number which utilise the most popular strategy increases by a constant factor of its current size during each round. Thus amongst this set of vertices, we achieve unanimity in a logarithmic number of rounds, thus providing the $1-\varepsilon$ proportion, if G is not connected. This procedure is outlined in Lemma 4.9 and proven in Appendix D. Furthermore, if $d > \alpha(\lambda) \log n$, the graph is connected with high probability and, therefore, the only vertices which are not unanimous are those of low degree. We prove a sequence of structural lemmas concerning how these vertices are connected, which in turn imply that structures which block unanimity do not appear. Thus the low degree vertices synchronise with the high degree vertices. The proofs of the structural lemmas can be found in Appendix C, while the full proof of this theorem can be found in Section 4.4.

We refine the above theorem focusing on the largest connected component of G(n,p)which we denote by $L_1(G(n,p))$ (formally, if there are at least two, we take the lexicographically smallest one). Let $u_n^{(1)}$ be the probability that $L_1(G(n,d/n))$ will eventually become unanimous. The next two theorems give the precise location of the threshold on p above which $u_n^{(1)}$ approaches 1. In fact, there are two different thresholds for the two regimes. We start with the majority regime. For $\lambda > 0$, let

$$\ell_{\lambda} := \lceil \max\{\lambda, \lambda^{-1}\} \rceil$$
 and $c_{\lambda} := \frac{1}{\ell_{\lambda} + 1}$.

(For a real number x > 0, we let $\lceil x \rceil = x$, if $x \in \mathbb{N}$, and $\lceil x \rceil = \lfloor x \rfloor + 1$, otherwise.) We write $\operatorname{Po}(\gamma)$ for the Poisson distribution with parameter $\gamma > 0$.

Theorem 3.3. Suppose that $\lambda \neq 1$ and $d = c_{\lambda} \log n + \log \log n + \omega(n)$. Then the following hold in the majority regime.

(1) If $\omega(n) \to +\infty$ as $n \to +\infty$, then

$$\lim_{n \to +\infty} u_n^{(1)} = 1.$$

(2) If $\omega(n) \to c \in \mathbb{R}$ as $n \to +\infty$, then

$$\limsup_{n \to +\infty} u_n^{(1)} \le \sum_{k=0}^{\infty} \left(1 - \frac{1}{2^{\ell_{\lambda}+1}} \right)^k \mathbb{P} \left(\operatorname{Po}(e^{c(\ell_{\lambda}+1)}/\ell_{\lambda}!) = k \right).$$

and

$$\liminf_{n \to +\infty} u_n^{(1)} \ge \sum_{k=0}^{\infty} \left(\frac{1}{2^{\ell_{\lambda}+1}}\right)^k \mathbb{P}\left(\operatorname{Po}(e^{c(\ell_{\lambda}+1)}/\ell_{\lambda}!) = k\right)$$

(3) If $\omega(n) \to -\infty$ as $n \to +\infty$, then

$$\lim_{n \to +\infty} u_n^{(1)} = 0$$

The analogous result for the minority regime is as follows. For $\lambda > 0$, we let

$$\ell_{\lambda}' = \lfloor \max\{\lambda, \lambda^{-1}\} \rfloor.$$

Theorem 3.4. Suppose that $\lambda \neq 1$ and $d = \frac{1}{2}\log n + \frac{1+\ell_{\lambda}'}{2}\log\log n + \omega(n)$. Then the following hold in the minority regime.

(1) If $\omega(n) \to +\infty$ as $n \to +\infty$, then

$$\lim_{n \to +\infty} u_n^{(1)} = 1.$$

(2) If $\omega(n) \to c \in \mathbb{R}$ as $n \to +\infty$, then

$$\limsup_{n \to +\infty} u_n^{(1)} \le \sum_{k=0}^{\infty} \left(\frac{3}{4}\right)^k \mathbb{P}\left(\operatorname{Po}(e^{2c}/\ell_{\lambda}'!) = k\right).$$

and

$$\liminf_{n \to +\infty} u_n^{(1)} \ge \mathbb{P}\left(\operatorname{Po}(e^{2c}/\ell_{\lambda}'!) = 0 \right).$$

(3) If $\omega(n) \to -\infty$ as $n \to +\infty$, then

$$\lim_{n \to +\infty} u_n^{(1)} = 0.$$

Proof sketch. Essentially both of these theorems capture the threshold needed for the existence of structures which block unanimity. The exact structure which these theorems describe is that of a blocking star; this is introduced in Section 4.1. Unanimity is achieved if and only if there is no induced copy of a suitable blocking star in the network. In each regime, when p is below the threshold a suitable blocking star appears, while no induced copies are found when p is above the threshold. It is worth noting that thresholds differ between regimes. This follows due to the fact that differing evolution rules, require different types of structures to block unanimity. The proofs of these results are supported by a sequence of lemmas concerning the distribution of blocking stars in G(n, p). These are stated in Section 4.1, and proved in Appendix B. The proofs of these two theorems can be found in Section 4.4.

The reason for the existence of two different thresholds is that the structures that block unanimity are different in the two regimes. Effectively, these are the thresholds for their disappearance as subgraphs of G(n, p) and, as it turns out, as subgraphs of $L_1(G(n, p))$. Nevertheless, as our arguments show, in both regimes the unanimity of $L_1(G(n, p))$ is achieved in $O(\log n)$ steps.

During our analysis, we will see that there are two different kinds of unanimity depending on whether the payoff matrix is in the majority or in the minority regime. In the majority regime, all vertices of $L_1(G(n, p))$ stabilise to one of the two strategies. However, in the minority regime, the vertices of $L_1(G(n, p))$ arrive at unanimity but they fluctuate incessantly between the two strategies. In other words, in the majority regime the subsystem of $L_1(G(n, p))$ becomes periodic with period 1, whereas in the minority regime the period is equal to 2. Furthermore, we identify the strategy that is played at each step once this subsystem has entered the periodic cycle.

We close this introductory section with a brief discussion on the case of degenerate payoff matrices.

3.2. Degenerate payoff matrices. If Q is degenerate, then the behaviour of the interacting node system is readily deduced. The system will either reach stability from the outset, or it will reach stable unanimity after one round. We summarise this behaviour in the following lemma.

Lemma 3.5. Let Q be a degenerate payoff matrix, G a connected graph, and S an initial configuration of vertex strategies. Then the interacting node system (G, Q, S), evolves as follows: Either the system is stable from T = 0, or the system is unanimous and stable from T = 1.

Proof. We divide our proof into a number of cases. Firstly, we note that if $q_{0,0} = q_{1,0}$ and $q_{0,1} = q_{1,1}$, then we must achieve stability from the initial state, since $T_0(v) = T'_0(v)$. Hence, for all $v \in V(G)$ we have that $S_0(v) = S_t(v)$ for $t \ge 0$.

Suppose that the above case does not occur, and we have $q_{0,0} > q_{1,0}$. For Q to be degenerate, we are forced to have that $q_{0,1} \ge q_{1,1}$. Consequently, we now have top row domination in Q. For any game played by vertex v, it is always optimal to play strategy

zero. As G is connected, every vertex will play at least one game, and, in particular, it is always optimal to play strategy 0, which corresponds to the top row of the payoff matrix. Therefore, for all v we have that $S_1(v) = 0$, and a stable unanimity is achieved. A similar row domination argument follows for the remaining possibilities of degenerate matrices.

As Lemma 3.5 holds when G is any connected graph, it then follows that this argument suffices as a proof of both Theorem 3.1 and Theorem 3.2 for the case that Q is a degenerate payoff matrix.

3.3. Notation and probabilistic tools. We finish the introductory section giving some notation. For two distinct vertices $v, u \in V$ of a graph G(V, E), we write $v \sim u$ to indicate that they are adjacent, that is, $uv \in E$. We also use the symbol \sim in the context of random variables. In particular, for a random variable X we will write $X \sim$ $\operatorname{Bin}(s,q)$ to indicate that the distribution of X is the binomial with parameters $s \in \mathbb{N}$ and $q \in [0,1]$. Furthermore, we write $f(n) \sim g(n)$ for two real-valued functions on \mathbb{N} , if $\lim_{n\to\infty} |f(n)|/|g(n)| = 1$. Also, we write $f \gg g$, if $\lim_{n\to\infty} |f(n)|/|g(n)| = \infty$.

For a vertex v in a given graph, we will set $d(v) := |N_G(v)|$ to be the degree of v within the graph G, and recall that $N_G(v)$ denotes the neighbourhood of v in G. The particular graph we refer to will be made specific by the context. If $S \subset V$, we will be writing e(S)for the number of edges that G spans inside S. Also, if $v \in V$ we denote by $d_S(v)$ the degree of v inside S.

When we consider the process on G(n, p) we shall use the symbols P_t and N_t to denote the subsets of vertices whose agents play strategy 1 and 0, respectively, after the *t*th step. More formally, for $t \ge 0$, we define $P_t = \{v \in V_n : S_t(v) = 1\}$ and $N_t = \{v \in V_n : S_t(v) = 0\}$.

The Chernoff bound. Throughout our arguments we will use the standard Chernoff bounds for concentration of binomially distributed random variables. The inequality we will use follows from Theorem 2.1 in [23]. If X is a random variable such that $X \sim Bin(N, q)$, then for any $\delta \in (0, 1)$ we have

$$\mathbb{P}(|X - Nq| \ge \delta Nq) \le 2e^{-\delta^2 Nq/3}.$$
(3)

4. Interacting node systems with skew $\lambda \neq 1$

4.1. Blocking structures and their properties. We will start with the identification of those structures/induced subgraphs of G(n, p) which, roughly speaking, will stay immune to what the rest of the graph is doing. Thus, these substructures act as obstructions to unanimity.

In particular, the structure we consider is an (ℓ, k) -blocking star. This is a star whose central vertex has degree $\ell + k$ in G(n, d/n) and, furthermore, ℓ leaves of the star have degree 1 inside G(n, d/n), whereas we impose no restriction on the degrees of the remaining k leaves. We call the latter leaves the *connectors* of the blocking star, whereas the ℓ leaves of degree 1 are called the *blocking leaves*. Such a structure is illustrated in Figure 1. An (ℓ, k) -blocking star has the property that it can block or absorb the influence of the external



FIGURE 1. A blocking star with central vertex v, blocking leaves v_1, \ldots, v_ℓ and connectors u_1, \ldots, u_k .

vertices depending on the choice of its parameters ℓ and k, respectively. Let us consider first the minority regime. It will turn out that it will be sufficient to consider (1, k)blocking stars for a suitable defined k. Let $i^* \in \{0, 1\}$ be such that $\lambda^{2i^*-1} = \max\{\lambda, \lambda^{-1}\}$. For $i, j \in \{0, 1\}$, we say that a star has the (i, j)-configuration if the blocking leaves play strategy i whereas the centre plays strategy j.

Claim 4.1. Consider a (1,k)-blocking star with $k \leq \lfloor \lambda^{2i^*-1} \rfloor$. In the minority regime, if the star ever gets into the $(i^*, 1 - i^*)$ -configuration, it will stay so forever.

Proof. Let v_1 denote the blocking leaf and v denote the centre of the star. Suppose that $S_t(v_1) = i^*$ but $S_t(v) = 1 - i^*$. By (2), the vertex v will change strategy, if $n_t(v; 1 - i^*) > \lambda^{2i^*-1}n_t(v; i^*)$. Since $S_t(v_1) = i^*$, then $n_t(v; 1 - i^*) \leq k \leq \lfloor \lambda^{2i^*-1} \rfloor$. But also $n_t(v; i^*) \geq 1$. So we would have $\lfloor \lambda^{2i^*-1} \rfloor > \lambda^{2i^*-1}$, which is impossible. Thus, $S_{t+1}(v) = 1 - i^*$. Furthermore, $S_{t+1}(v_1) = i^*$, since v_1 has no neighbours playing i^* .

Claim 4.2. Consider a (1, k)-blocking star with $k > \lfloor \lambda^{2i^*-1} \rfloor$. In the minority regime, if the k connectors simultaneously alternate between $1 - i^*$ and i^* , then the blocking leaf and centre will eventually synchronise with them.

Proof. Let v_1 be the blocking leaf, v be the centre and u_1, \ldots, u_k be k connectors of the star. Suppose that $S_t(u_j) = 1 - i^*$, for all $j = 1, \ldots, k$.

If $S_t(v) = 1 - i^*$, then v will change strategy if $n_t(v; 1 - i^*) > \lambda^{2i^*-1}n_t(v; i^*)$ (cf. (2)). But $n_t(v; 1 - i^*) \ge k$ and $n_t(v; i^*) \le 1$. Since $k > \lfloor \lambda^{2i^*-1} \rfloor$, the above inequality is indeed satisfied. Hence, $S_{t+1}(v) = i^*$.

On the other hand, if $S_t(v) = i^*$, then v will not change strategy if $n_t(v; i^*) \leq \lambda^{1-2i^*} n_t(v; 1-i^*)$. But $n_t(v; 1-i^*) \geq k > \lfloor \lambda^{2i^*-1} \rfloor$. Therefore, $n_t(v; 1-i^*) > \lambda^{2i^*-1}$. So $\lambda^{1-2i^*} n_t(v; 1-i^*) > 1$. But $n_t(v; i^*) \leq 1$ and the inequality is satisfied. Therefore, $S_{t+1}(v) = i^*$. However, now $S_{t+1}(v_1) = 1 - i^*$.

Suppose now that $S_t(u_j) = i^*$, for all j = 1, ..., k. If $S_t(v) = 1 - i^*$, then by (2) v will change strategy if $n_t(v; 1 - i^*) > \lambda^{2i^*-1}n_t(v; i^*)$. But now $n_t(v; 1 - i^*) \leq 1$ and $n_t(v; i^*) \geq k > \lambda^{2i^*-1}$. Thus, the above inequality is not satisfied and $S_{t+1}(v) = 1 - i^*$. Also, note that $S_{t+1}(v_1) = i^*$.

If $S_t(v) = i^*$, then v will not change strategy if $n_t(v; i^*) \leq \lambda^{1-2i^*} n_t(v; 1-i^*)$. Now, $n_t(v; i^*) \geq k > \lambda^{2i^*-1} > 1$ but $\lambda^{1-2i^*} n_t(v; 1-i^*) \leq \lambda^{1-2i^*} < 1$. So the above inequality is not satisfied and $S_{t+1}(v) = 1 - i^*$. Furthermore, $S_{t+1}(v_1) = 1 - i^*$.

We thus conclude that in any case the centre v will synchronise with the k connectors. Note that from the above four cases, we see that the blocking leaf v_1 will synchronise with v at the steps where it changes state. Hence, it will also synchronise with the k connectors too.

In the majority regime, it will turn out that we will need to consider $(\ell, 1)$ -blocking stars. We will show that if ℓ is sufficiently large, then the $(\ell, 1)$ -blocking star can block the influence of the external vertices and retain the strategy of its vertices.

Claim 4.3. Consider an $(\ell, 1)$ -blocking star with $\ell \geq \lceil \lambda^{2i^*-1} \rceil$. In the majority regime, if it is set to the $(1 - i^*, 1 - i^*)$ -configuration initially, then it will stay in this configuration forever.

Proof. Suppose that v_1, \ldots, v_{ℓ} are the ℓ -blocking leaves and v is the centre of the star. Assume that all $S_t(v) = S_t(v_1) = \cdots = S_t(v_{\ell}) = 1 - i^*$. By (1), the centre will change strategy at step t + 1, if $n_t(v; 1 - i^*) < \lambda^{2i^*-1}n_t(v; i^*)$. But $n_t(v; 1 - i^*) = \ell \ge \lceil \lambda^{2i^*-1} \rceil$ and $n_t(v; i^*) \le 1$. Hence, we should have $\lceil \lambda^{2i^*-1} \rceil < \lambda^{2i^*-1}$, which is impossible. So $S_{t+1}(v) = 1 - i^*$. Further, note that the ℓ -blocking leaves will adopt the strategy of the centre at step t + 1, since they have no other neighbours. Thus, $S_{t+1}(v_j) = 1 - i^*$, for any $j = 1, \ldots, \ell$, as well.

Claim 4.4. Consider an $(\ell, 1)$ -blocking star with $\ell \ge 1$. In the majority regime, if it is set to the (i^*, i^*) -configuration initially, then it will stay in this configuration forever.

Proof. Suppose that v_1, \ldots, v_{ℓ} are the ℓ -blocking leaves and v is the centre of the star. Assume that all $S_t(v) = S_t(v_1) = \cdots = S_t(v_{\ell}) = i^*$. By (1), the centre will change strategy at step t + 1, if $n_t(v; i^*) < \lambda^{1-2i^*} n_t(v; 1-i^*)$. But $n_t(v; i^*) \ge 1$ and $n_t(v; 1-i^*) \le 1$. So the above inequality is not satisfied, since $\lambda^{1-2i^*} < 1$, and, therefore, $S_{t+1}(v) = i^*$. Finally, the ℓ -blocking leaves will retain the strategy of the centre at step t + 1, since they have no other neighbours. Thus, $S_{t+1}(v_j) = i^*$, for any $j = 1, \ldots, \ell$, as well.

4.2. The distribution of blocking stars inside G(n, d/n). Now, we shall give a general condition that determines the distribution of the (ℓ, k) -blocking stars in G(n, d/n). The following results will be useful both for the subcritical and the supercritical regime that we analyse in the next subsection.

Let $X_{\ell,k,n}$ be the random variable which is the number of (ℓ, k) -blocking stars in G(n, d/n)and let $X_{\ell,k,n}^{(1)}$ be the number of those (ℓ, k) -blocking stars which are subgraphs of the largest component, $L_1(G(n, d/n))$. Clearly, $X_{\ell,k,n}^{(1)} \leq X_{\ell,k,n}$. However, the following result states that a.a.s. these two random variables are approximately equal. In particular, we consider the following two lemmas, the proofs of which can be found in Appendix B.

Lemma 4.5. Let $\ell \in \mathbb{N}$ and p = d/n, where $1 \ll d = d(n) = O(\log n)$. Then $\mathbb{E}X_{\ell,n} \sim n \frac{d^{\ell+k}}{\ell |k|} e^{-d(\ell+1)}$.

Furthermore, if $d = \frac{1}{\ell+1} \log n + \frac{\ell+k}{\ell+1} \log \log n + \omega(n)$, then the following hold.

- i. If $\omega(n) \to -\infty$ as $n \to +\infty$, then $\mathbb{E}X_{\ell,k,n} \to +\infty$ as $n \to +\infty$ and moreover, a.a.s. $X_{\ell,k,n} \ge \mathbb{E}X_{\ell,k,n}/2.$
- ii. If $\omega(n) \to c \in \mathbb{R}$ as $n \to +\infty$, then $\mathbb{E}X_{\ell,k,n} \to \frac{e^{(\ell+1)c}}{\ell!k!}$ as $n \to \infty$ and $X_{\ell,k,n} \xrightarrow{d} \operatorname{Po}\left(\frac{e^{(\ell+1)c}}{\ell!k!}\right)$.
- iii. If $\omega(n) \to +\infty$ as $n \to +\infty$, then $\mathbb{P}(X_{\ell,k,n} > 0) < 2e^{-(\ell+1)\omega(n)}$, for any n sufficiently large. Thus, a.a.s. $X_{\ell,k,n} = 0$.

Let $X_{\ell,k,n}^{(2)} = X_{\ell,k,n} - X_{\ell,k,n}^{(1)}$.

Lemma 4.6. We have $\mathbb{E}X_{\ell,k,n}^{(2)} = o(\mathbb{E}X_{\ell,k,n}).$

So Markov's inequality implies that a.a.s. $X_{\ell,k,n}^{(2)} \leq \mathbb{E}X_{\ell,k,n}/4$. By Lemma 4.5, if $d = \frac{1}{\ell+1}\log n + \frac{\ell+k}{\ell+1}\log\log n + \omega(n)$, with $\omega(n) \to -\infty$, then a.a.s.

$$X_{\ell,k,n}^{(1)} \ge \frac{1}{4} \mathbb{E} X_{\ell,k,n}.$$
 (4)

Furthermore, note that any two (ℓ, k) -blocking stars can share only their connector vertices. So, if we consider the initial assignment of strategies, each (ℓ, k) -blocking star inside $L_1(G(n, d/n))$ will be set into (i, j)-configuration with probability $1/2^{\ell+1}$, independently of each other. Thus, the weak law of large numbers together with (4) imply the following.

Lemma 4.7. Let p = d/n, where $d = \frac{1}{\ell+1} \log n + \frac{\ell+k}{\ell+1} \log \log n + \omega(n)$, for $k, \ell \in \mathbb{N}$. Let $i, j \in \{0, 1\}$.

- i. If $\omega(n) \to -\infty$ as $n \to +\infty$, then a.a.s. at least $\frac{1}{8} \cdot \frac{1}{2^{\ell+1}} \mathbb{E}X_{\ell,k,n}$ of the (ℓ, k) -blocking stars inside $L_1(G(n, d/n))$ will be set into (i, j)-configuration at the beginning of the process.
- ii. If $\omega(n) \to c \in \mathbb{R}$ as $n \to +\infty$, then the number of (ℓ, k) -blocking stars inside $L_1(G(n, d/n))$ will be set into (i, j)-configuration at the beginning of the process converges in distribution as $n \to \infty$ to a random variable distributed as $\operatorname{Bin}(\operatorname{Po}(e^{(\ell+1)c}/(\ell!k!)), \frac{1}{2^{\ell+1}}).$

4.3. Analysis of first round, and a procedure for declaring unanimity in vertices of high degree. In this subsection, we analyse the first round of the process. In Lemma 4.8 we show that after one round, a sublinear majority is formed. It then follows that if G(n, p) is dense enough, then this majority is in fact unanimity. In sparser cases we show that vertices of degree at least some (large) constant C, reach unanimity over a logarithmic number of rounds. This argument is presented in Lemma 4.9. Suppose that $\mathcal{I} = (G, Q, \mathcal{S})$ is an interacting node system with $\lambda = \lambda(Q) \neq 1$ and G = G(n, p). We denote by $m_t = \operatorname{argmin}\{|N_t|, |P_t|\}$ and $\mu_t = \min\{|N_t|, |P_t|\}$. The following lemma describes the first round of the evolution; notably it describes the formation of a large majority after a single round. Recall that $i^* \in \{0, 1\}$ is the strategy which satisfies $\lambda^{1-2i^*} < 1$. Note that since $\lambda \neq 1$, exactly one of 0 or 1 satisfies this. Note further that $\lambda^{1-2i^*} = \min\{\lambda, \lambda^{-1}\}$. Hence,

$$\max\{\lambda, \lambda^{-1}\} \cdot \lambda^{1-2i^*} = \max\{\lambda, \lambda^{-1}\} \cdot \min\{\lambda, \lambda^{-1}\} = \lambda \cdot \lambda^{-1} = 1.$$
(5)

We will use this identity later on.

Now, we will show that, in the majority regime, after one round strategy i^* will become the dominant strategy among the vertices of G(n, d/n). However, in the minority regime it will be strategy $1 - i^*$ that will dominate.

Lemma 4.8. Let p = d/n where $d \gg 1$. For any $0 < \lambda \neq 1$, there exists $\gamma > 0$ for which the following holds. A.a.s. across the product space of G(n, p) and $S_{1/2}$: For the interacting node system $\mathcal{I} = (G(n, p), Q, S_{1/2})$ with $\lambda(Q) = \lambda$, we have $\mu_1 \leq ne^{-\gamma d}$. In particular, after the first round, the majority of the vertices will be playing either strategy i^* (majority regime) or strategy $1 - i^*$ (minority regime). Furthermore, there exists $\alpha(\lambda) > 1$ such that if $d > \alpha(\lambda) \log n$, then a.a.s. $\mu_1 = 0$.

Proof. Suppose first that our system is in the majority regime. By (1), if a vertex plays strategy $1 - i^*$, then it will change strategy if $n_0(v; 1 - i^*) < \lambda^{1-2(1-i^*)}n_0(v; i^*) = \lambda^{2i^*-1}n_0(v; i^*)$. Also, if a vertex plays strategy i^* , then it will stay there, if $n_0(v; i^*) \geq \lambda^{1-2i^*}n_0(v; 1 - i^*)$.

Suppose now that our system is in the minority regime. By (2), if a vertex plays strategy $1 - i^*$, then it will remain there if $n_0(v; 1 - i^*) \leq \lambda^{1-2(1-i^*)}n_0(v; i^*) = \lambda^{2i^*-1}n_0(v; i^*)$. Also, if a vertex plays strategy i^* , then it will switch to strategy $1 - i^*$, if $n_0(v; i^*) > \lambda^{1-2i^*}n_0(v; 1 - i^*)$.

For each $v \in V_n$ and for $\delta \in (0, 1)$ we say that v is δ -balanced if for all $i \in \{0, 1\}$, we have that $|n_0(v; i) - \mathbb{E}[n_0(v; i)]| \leq \delta \mathbb{E}[n_0(v; i)]$. If v is not δ -balanced then we say that v is δ -unbalanced, and we denote the set of δ -unbalanced vertices as \mathcal{U}_{δ} . Now, note that if v is δ -balanced, then provided that $\delta = \delta(\lambda) \in (0, 1)$ is sufficiently small

$$\lambda^{1-2i^*} < \frac{1-\delta}{1+\delta} \le \frac{n_0(v; 1-i^*)}{n_0(v; i^*)} \le \frac{1+\delta}{1-\delta} < \lambda^{2i^*-1}.$$

In other words, if v is δ -balanced, then all the above four inequalities will be satisfied. We thus arrive at the following conclusions:

- 1. if v is δ -balanced and Q is in the majority regime, then $S_1(v) = i^*$;
- 2. if v is δ -balanced but Q is in the minority regime, then $S_1(v) = 1 i^*$.

Furthermore, if the majority of the vertices in V_n are δ -balanced, then $\mu_1 \leq |\mathcal{U}_{\delta}|$. We will show that a.a.s. the majority of the vertices in V_n are δ -balanced, whereby they adopt strategy i^* or $1 - i^*$ after one step, as described above.

We will show that an arbitrary vertex $v \in V_n$ is δ -balanced with probability 1-o(1). For any $v \in V_n$ the random variables $n_0(v; 0)$ and $n_0(v; 1)$ have identical distributions, namely the binomial distribution $\operatorname{Bin}(n-1, p/2)$. Therefore

$$\mathbb{E}[n_0(v;1)] = \mathbb{E}[n_0(v;0)] = (1-o(1))\frac{d}{2}.$$

Set $\bar{\gamma} = \delta^2/7$. We bound the probability that $v \in \mathcal{U}_{\delta}$. By Chernoff's inequality (3) we have:

$$\mathbb{P}\Big[\Big|n_0(v;0) - \mathbb{E}[n_0(v;0)]\Big| \ge \delta \mathbb{E}[n_0(v;0)]\Big] \le 2e^{-\frac{\delta^2}{3}\frac{(1-o(1))d}{2}} \le e^{-\bar{\gamma}d}$$

where the last inequality holds for n sufficiently large. The same holds for $n_0(v; 1)$ as it is identically distributed to $n_0(v; 0)$. Hence, by the union bound, for any $v \in V_n$

$$\mathbb{P}\left[v \in \mathcal{U}_{\delta}\right] \le 2e^{-\bar{\gamma}d}$$

Therefore, $\mathbb{E}\left[|\mathcal{U}_{\delta}|\right] \leq 2ne^{-\bar{\gamma}d}$.

By Markov's inequality we have:

$$\mathbb{P}[|\mathcal{U}_{\delta}| \ge n e^{-\bar{\gamma}d/2}] \le \frac{2n e^{-\bar{\gamma}d}}{n e^{-\bar{\gamma}d/2}} = 2e^{-\bar{\gamma}d/2} = o(1).$$

Therefore, a.a.s. we have that $|\mathcal{U}_{\delta}| < ne^{-\bar{\gamma}d/2}$. In turn, a.a.s. $\mu_1 \leq |\mathcal{U}_{\delta}| < ne^{-\bar{\gamma}d/2}$.

Finally, note that the last inequality above implies that if $d > \alpha(\lambda) \log n$, for some $\alpha(\lambda) > 1$ sufficiently large then the above is o(1), which shows the last part of the lemma.

It suffices to focus on the case where $d \leq \alpha(\lambda) \log n$, as otherwise a.a.s. the process reaches unanimity after one step by the above lemma. For the analysis of the subsequent rounds we will split the vertices of G(n, d/n) into two classes and we will consider their evolution separately. More specifically, for $C \in \mathbb{N}$ we set

$$H_n(C) := H(C, G(n, d/n)) := \{ v \in V_n : d(v) \ge C \}$$

and

$$L_n(C) := L(C, G(n, d/n)) := V_n \setminus H(C, G(n, d/n)) = \{ v \in V_n : d(v) < C \}.$$

The following lemma concerns the evolution of the vertices in $H_n(C)$. It shows that if one of the strategies occupies only a sublinear number of vertices in $H_n(C)$ (and we have almost unanimity), then after one more round the size of the minority strategy will contain only a fraction of these vertices. The proof of this lemma can be found in Appendix D.

Lemma 4.9. Let d = d(n) be such that $1 \ll d \leq \alpha(\lambda) \log n$, and let Q be a 2×2 nondegenerate payoff matrix. For any $\varepsilon > 0$ there exists $C_{\varepsilon,\lambda} \in \mathbb{N}$ such that for any $\gamma > 0$ and for any $C \geq C_{\varepsilon,\lambda}$ a.a.s. G(n, d/n) satisfies the following: for any initial configuration Swith $|m_0 \cap H_n(C)| < ne^{-\gamma d}$, the interacting node system $\mathcal{I} = (G(n, d/n), Q, S)$ will have $|m_1 \cap H_n(C)| \leq \varepsilon |m_0 \cap H_n(C)|$.

Remark 4.10. It is not very hard to show that in fact a.a.s. $H_n(C)$ consists of the majority of the vertices of G(n, d/n) if $d \gg 1$. (For example, an application of the Chernoff bound can show that most vertices have degree around d.) More specifically, if $d \gg 1$, then by (3) for any vertex $v \in V_n$

$$\mathbb{P}(d(v) \le C) = o(1).$$

Hence, $\mathbb{E}(|L_n(C)|) = o(n)$. By Markov's inequality for any $\varepsilon > 0$ a.a.s.

 $|L_n(C)| < \varepsilon n,$

as long as $d \gg 1$.

18

4.4. Small degree vertices, their structure and their role in unanimity: proof of Theorems 3.2, 3.3 and 3.4. Lemma 4.9 above implies that $H_n(C)$ will become unanimous after a number of rounds. So now, it remains to characterise when $L_n(C)$ will synchronise with $H_n(C)$. In order to do this we need to consider which structures can occur within $L_n(C)$ and observe what this implies for the evolution rules. We will now state a selection of lemmas which describe the structure of the subgraph induced by the vertices in $L_n(C)$, for any fixed integer $C \ge 2$. Their proofs can be found in Appendix C. Let us recall that $\ell_{\lambda} = \lceil \max\{\lambda, \lambda^{-1}\} \rceil$ and $c_{\lambda} = \frac{1}{\ell_{\lambda+1}}$.

Lemma 4.11. Suppose that p = d/n with $c_{\lambda} \log n \le d \le \alpha(\lambda) \log n$. A.a.s. there are no $\ell_{\lambda} + 2$ vertices in $L_n(C)$ that have a common neighbour.

The above lemma implies in particular that a.a.s. at most $\ell_{\lambda} + 1$ vertices in $L_n(C)$ are adjacent to each vertex in $H_n(C)$. Thus, we see that if all vertices in $H_n(C)$ play a certain strategy simultaneously and C is large compared to ℓ_{λ} , then they may stay unaffected by what the vertices in $L_n(C)$ do. The following two lemmas state that connected components in $L_n(C)$ are both small, and can not be too complex. We will apply these assertions to show unanimity occurring within $L_n(C)$.

Lemma 4.12. Let $\ell \in \mathbb{N}$ and let p = d/n where $\frac{1}{\ell+1} \log n \leq d = d(n) \leq \alpha(\lambda) \log n$. A.a.s. all connected sets of vertices in $L_n(C)$ have size at most $\ell + 1$.

Lemma 4.13. Let p = d/n where $c_{\lambda} \log n \le d = d(n) \le \alpha(\lambda) \log n$. A.a.s. all connected sets of vertices in $L_n(C)$ induce trees.

We now proceed with the proofs of Theorems 3.2, 3.3 and 3.4.

Proof of Theorem 3.2. Let us first point out that by Lemma 4.8, if $d \ge c \log n$ with $c > \alpha(\lambda)$, where $\alpha(\lambda)$ is as in the statement of that lemma, then a.a.s. $\mu_1 = 0$; so the last part of Theorem 3.2 follows. Hence, we now assume that $d \le \alpha(\lambda) \log n$. We say that G(n, d/n) has the minority decline property for some $\gamma > 0$, whenever for any configuration \mathcal{S} on V_n the node system $\mathcal{I} = (G(n, d/n), Q, \mathcal{S})$ is such that $\mu_0 \le ne^{-\gamma d}$, then $|m_1 \cap H_n(C)| \le |m_0 \cap H_n(C)|/10$. Observe now that if G(n, d/n) has the minority decline property, and $\mathcal{I} = (G(n, p, Q, \mathcal{S}))$ is a node system with $\mu_0 < ne^{-\gamma d}$, for some $\gamma > 0$, then the vertices in H_n will reach unanimity in a finite number of rounds, by repeated applications of this definition.

By Lemma 4.8, a.a.s. $\mu_1 < ne^{-\gamma d}$ for some $\gamma > 0$. But by Lemma 4.9, if $C > C_{1/10,\lambda}$, then a.a.s. G(n,p) has the minority decline property for $\gamma > 0$ as above. Thus, for every $t \ge 1$ we have $|m_t \cap H_n(C)| \le |m_0 \cap H_n(C)| 10^{-t}$. So for $R = \lceil (1/\log 10) \log (|m_1 \cap H_n(C)|) \rceil + 1 = O(\log n)$ we have

 $|m_R \cap H_n(C)| \le |m_1 \cap H_n(C)| 10^{-R} < 1.$

So $|m_R \cap H_n(C)| = 0$. We now show that if C is sufficiently large and every vertex in $H_n(C)$ does not have too many neighbours inside $L_n(C)$, then once $H_n(C)$ has reached unanimity, it will stay there. In particular, Lemma 4.11 states that a.a.s. no $\ell_{\lambda} + 2$ vertices in $L_n(C)$ have a common neighbour. Let us denote this event by \mathcal{D}_n . Thus, on \mathcal{D}_n all vertices in $H_n(C)$ have at most $\ell_{\lambda} + 1$ neighbours inside $L_n(C)$.

Claim 4.14. If $C > \max\{\ell_{\lambda} + 4, (\ell_{\lambda} + 1)^2\}$ and the vertices in $H_n(C)$ are unanimous just after step t, playing i^{*} when in the majority regime, then on the event \mathcal{D}_n , the vertices of $H_n(C)$ will stay unanimous after step t + 1.

Proof of Claim 4.14. Indeed, suppose that at step t all vertices of $H_n(C)$, for C to be determined, are unanimous at playing strategy $i \in \{0, 1\}$. Consider a vertex $v \in H_n(C)$. If the event \mathcal{D}_n is realised, then all but at most $\ell_{\lambda}+1 = \lceil \max\{\lambda, \lambda^{-1}\} \rceil + 1$ neighbours of v play strategy i. So $n_t(v; i) \ge C - (\ell_{\lambda}+1)$ and $n_t(v; 1-i) \le \lceil \max\{\lambda, \lambda^{-1}\} \rceil + 1 < \max\{\lambda, \lambda^{-1}\} + 2$.

Suppose first that we are in the majority regime. Then in this case $i = i^*$, by assumption. Vertex v will change strategy if $n_t(v; i^*) < \lambda^{1-2i^*} n_t(v; 1-i^*)$ (cf. (1)). But

$$\lambda^{1-2i^*} n_t(v; 1-i^*) < \lambda^{1-2i^*} (\ell_{\lambda}+1) \le \lambda^{1-2i^*} (\max\{\lambda, \lambda^{-1}\}+2) \stackrel{(5)}{<} 1+2=3$$

Therefore must have that $C - (\ell_{\lambda} + 1) < 3$. However, choosing $C \ge \ell_{\lambda} + 4$ leads to a contradiction.

If we are in the minority regime, then we want to show that v will switch strategy. By (2) if the entire set $H_n(C)$ plays strategy i^* , then $v \in H_n(C)$ will switch strategy if $n_t(v;i^*) > \lambda^{1-2i^*}n_t(v;1-i^*)$. But $n_t(v;i^*) \ge C - (\ell_{\lambda}+1)$ and $n_t(v;1-i^*) \le \ell_{\lambda}+1$. As seen above, if $C \ge \ell_{\lambda}+4$ then $n_t(v;i^*) \ge C - (\ell_{\lambda}+1) \ge 3 > \lambda^{1-2i^*}(\ell_{\lambda}+1) \ge \lambda^{1-2i^*}n_t(v;1-i^*)$. Now, if the entire set $H_n(C)$ plays strategy $1 - i^*$, then $v \in H_n(C)$ will switch strategy if $n_t(v;1-i^*) > \lambda^{1-2(1-i^*)}n_t(v;i^*)$. But $n_t(v;i^*) \le \ell_{\lambda}+1$ whereas $n_t(v;1-i^*) \ge C - (\ell_{\lambda}+1)$. Thus v will switch strategy if $C - (\ell_{\lambda}+1) \ge \lambda^{2i^*-1}(\ell_{\lambda}+1)$. Therefore, choosing $C \ge (\ell_{\lambda}+1)^2$ verifies it and completes the proof of the claim.

Remark 4.15. For the minority regime, the above claim and Lemma 4.8 imply that, when unanimity occurs within $H_n(C)$, its vertices will be playing strategy $1 - i^*$ at odd steps and strategy i^* at even steps. In the majority regime, they stabilise to strategy i^* .

Note that by Remark 4.10, if $d \gg 1$, then for any fixed $C \in \mathbb{N}$ we have a.a.s. $|H_n(C)| \ge n(1-o(1))$. Therefore, the above analysis implies that for any $\varepsilon > 0$ there exists $\beta = \beta(\varepsilon, \lambda) > 0$ such that if $d \gg 1$, then a.a.s. at least $n(1-\varepsilon)$ vertices in G(n, d/n) will be unanimous after at most $\beta \log n$ rounds.

Proof of Theorem 3.3. Suppose first that $d = c_{\lambda} \log n + \log \log n + \omega(n)$, where $\omega(n) \to -\infty$ as $n \to +\infty$. By Lemma 4.7 *i*., a.a.s. there are $(\ell_{\lambda}, 1)$ -blocking stars in $L_1(G(n, d/n))$ that are set to the $(1-i^*, 1-i^*)$ -configuration. By Claim 4.3, those will retain this configuration forever and, therefore, they will be in disagreement with the vertices in $H_n(C)$. Hence, $u_n^{(1)}$, the probability that $L_1(G(n, d/n))$ becomes eventually unanimous, tends to 0 as $n \to +\infty$.

We will now consider the cases where $d = c_{\lambda} \log n + \log \log n + \omega(n)$, where either $\omega(n) \to +\infty$ or $\omega(n) \to c \in \mathbb{R}$, as $n \to +\infty$. We will need the following claim.

Claim 4.16. Let Q be in the majority regime. If a vertex v has at most $\ell_{\lambda} - 1$ neighbours playing strategy $1 - i^*$ but at least one playing strategy i^* , it will play strategy i^* in the next round.

Proof of Claim 4.16. If v already plays strategy i^* , then it will change strategy if $n_t(v; i^*) < \lambda^{1-2i^*} n_t(v; 1-i^*)$. But $n_t(v; i^*) \ge 1$ and $n_t(v; 1-i^*) \le \ell_{\lambda} - 1$.

$$\lambda^{1-2i^*} n_t(v; 1-i^*) \le \lambda^{1-2i^*} (\ell_{\lambda} - 1) < \lambda^{1-2i^*} \cdot \max\{\lambda, \lambda^{-1}\} \stackrel{(5)}{=} 1.$$

So v will not change strategy. Now, if v already plays strategy $1-i^*$, then it will not change its strategy if $n_t(v; 1-i^*) \geq \lambda^{1-2(1-i^*)}n_t(v; i^*) = \lambda^{2i^*-1}n_t(v; i^*)$. But $n_t(v; 1-i^*) \leq \ell_{\lambda} - 1$ whereas $\lambda^{2i^*-1}n_t(v; i^*) \geq \lambda^{2i^*-1} > \ell_{\lambda} - 1$. Therefore, v will change its strategy into i^* . \Box

Suppose that $d = c_{\lambda} \log n + \log \log n + \omega(n)$ with $\omega(n) \to +\infty$ as $n \to +\infty$. Let $G^{(L)}(n, d/n)$ denote the subgraph of G(n, d/n) induced by the vertices in $L_n(C)$. By Lemma 4.12 (with $\ell = \ell_{\lambda}$) and Lemma 4.13, every connected component of $G^{(L)}(n, d/n)$ is a tree of order at most $\ell_{\lambda} + 1$. Let T be one of these connected components that is a subgraph of $L_1(G(n, d/n))$.

If $|T| \leq \ell_{\lambda}$, then all its vertices have degree at most $\ell_{\lambda} - 1$ in T. For $i \geq 1$, let $T^{(i)}$ denote the set of vertices in T that are at distance i from $H_n(C)$. Once the vertices in $H_n(C)$ have been unanimous on strategy i^* , they will stay there forever. By Claim 4.16 the vertices in $T^{(1)}$ will adopt strategy i^* and remain there forever. Assuming that the vertices $T^{(i)}$ have adopted strategy i^* for ever, then the vertices of $T^{(i+1)}$ will adopt strategy i^* too (provided it is non-empty) by Claim 4.16 and remain there forever. Hence, the entire vertex set of Twill adopt strategy i^* .

Suppose now that $|T| = \ell_{\lambda} + 1$. If all its vertices have degree at most $\ell_{\lambda} - 1$, then eventually the vertices of T adopt strategy i^* , by the above argument. If there is a vertex in T of degree $\ell_{\lambda} + 1$ within T, then T must be a star. However, by Lemma 4.7 *ii.* a.a.s. this is not an $(\ell_{\lambda}, 1)$ -blocking star. Thus, one of its leaves must have a neighbour in $H_n(C)$. Since it has degree $1 (\leq \ell_{\lambda} - 1)$ inside T, then by Claim 4.16 it adopts strategy i^* after $H_n(C)$ becomes unanimous and stays there forever. Subsequently, the centre of T will do so (it also has at most $\ell_{\lambda} - 1$ neighbours that are not playing strategy i^*) and finally the remaining leaves adopt it as well.

Moreover, the above argument shows that the only connected components of $G^{(L)}(n, d/n)$ which may not adopt strategy i^* are the $(\ell, 1)$ -blocking stars, for $\ell \leq \ell_{\lambda}$. If there are no $(\ell_{\lambda}, 1)$ -blocking stars, then $L_1(G(n, d/n))$ will then become unanimous.

Firstly, let us observe that if $\omega(n) \to c \in \mathbb{R}$ as $n \to +\infty$, then by Lemma 4.5 *iii*. a.a.s. there are no $(\ell_{\lambda} + 1, 1)$ -blocking stars as

$$\mathbb{P}(X_{\ell_{\lambda}+1,1,n} > 0) \le 2e^{-\Omega(\log n)} = o(1).$$

Furthermore, by Lemma 4.11 a.a.s. there are no $\ell_{\lambda} + 2$ vertices of degree 1 that have a common neighbour. Therefore, a.a.s. there are no $(\ell, 1)$ -blocking stars for any $\ell \ge \ell_{\lambda} + 1$.

Now, by Lemma 4.7 *ii*., the random variable $X_{\ell_{\lambda},1,n}^{(1)}$ converges in distribution as $n \to +\infty$ to a random variable distributed as $\operatorname{Po}(e^{c(\ell_{\lambda}+1)}/\ell_{\lambda}!)$. Thus, for any integer $k \ge 0$, we have

$$\mathbb{P}(X_{\ell_{\lambda},1,n}^{(1)}=k) \to \mathbb{P}\left(\operatorname{Po}(e^{c(\ell_{\lambda}+1)}/\ell_{\lambda}!)=k\right)$$

as $n \to +\infty$.

Suppose now that $X_{\ell_{\lambda},1,n}^{(1)} = k$, for some $k \in \mathbb{N}_0$. The case k = 0 was treated above and unanimity is attained a.a.s. (on the conditional space where $X_{\ell_{\lambda},1,n}^{(1)} = 0$). Let us consider the case $k \geq 1$. If an $(\ell_{\lambda}, 1)$ -blocking star is initially set to $(1 - i^*, 1 - i^*)$ -configuration, then by Claim 4.3 it will stay in this configuration forever. We thus conclude that if unanimity is achieved then no $(\ell_{\lambda}, 1)$ -blocking star is initially set to $(1 - i^*, 1 - i^*)$ -configuration. The probability of this is $(1 - 1/2^{\ell_{\lambda}+1})^k$. Also, if all $(\ell_{\lambda}, 1)$ -blocking stars attached to $L_1(G(n, d/n))$ are initially set to (i^*, i^*) -configuration, they will remain so forever (cf. Claim 4.4) and will be synchronised with the vertices of $H_n(C)$. Thus, $L_1(G(n, d/n))$ will be unanimous. The probability of this is $1/2^{k(\ell_{\lambda}+1)}$.

Consequently,

$$\limsup_{n \to +\infty} u_n^{(1)} \le \sum_{k=0}^{\infty} \left(1 - \frac{1}{2^{\ell_{\lambda}+1}} \right)^k \mathbb{P} \left(\operatorname{Po}(e^{c(\ell_{\lambda}+1)}/\ell_{\lambda}!) = k \right).$$

and

$$\liminf_{n \to +\infty} u_n^{(1)} \ge \sum_{k=0}^{\infty} \left(\frac{1}{2^{\ell_{\lambda}+1}}\right)^k \mathbb{P}\left(\operatorname{Po}(e^{c(\ell_{\lambda}+1)}/\ell_{\lambda}!) = k\right).$$

Since $H_n(C)$ will reach unanimity in at most $\beta \log n$ steps, the above case analysis implies that a.a.s. if $L_1(G(n, d/n))$ reaches unanimity, then this will happen within $\beta \log n + O(1)$ steps.

Proof of Theorem 3.4. Let us recall that $\ell'_{\lambda} = \lfloor \max\{\lambda, \lambda^{-1}\} \rfloor = \lfloor \lambda^{2i^*-1} \rfloor$. Suppose first that $d = \frac{1}{2} \log n + \frac{1+\ell'_{\lambda}}{2} \log \log n + \omega(n)$, where $\omega(n) \to -\infty$ as $n \to +\infty$. By Lemma 4.7 *i*. (setting $\ell = 1$ and $k = \ell'_{\lambda}$ therein), a.a.s. there are $(1, \ell'_{\lambda})$ -blocking stars in $L_1(G(n, d/n))$ that are initially set to the $(i^*, 1 - i^*)$ -configuration. By Claim 4.1, those will retain this configuration forever and, therefore, they will be in disagreement with the vertices in $H_n(C)$. Hence, $u_n^{(1)}$, the probability that $L_1(G(n, d/n))$ becomes eventually unanimous, tends to 0 as $n \to +\infty$.

Now, suppose that $d = \frac{1}{2} \log n + \frac{1+\ell_{\lambda}'}{2} \log \log n + \omega(n)$, where $\omega(n) \to +\infty$. As before, we let $G^{(L)}(n, d/n)$ denote the subgraph of G(n, d/n) induced by the vertices in $L_n(C)$. By Lemma 4.12 (with $\ell = 1$) a.a.s. every connected component of $G^{(L)}(n, d/n)$ is of order at most 2. That is, a.a.s. every component of $G^{(L)}(n, d/n)$ is either a vertex or an edge.

Let T be one of these connected components that is also a subgraph of $L_1(G(n, d/n))$. If T is a vertex, then it will synchronise with the vertices $H_n(C)$ after the R^{th} step, where the vertices of $H_n(C)$ arrive at unanimity. Thus, all its neighbours (which lie in $H_n(C)$) will play the same strategy, say i, by (2) this vertex will adopt strategy 1 - i in the next round and be in agreement with the vertices of $H_n(C)$ (cf. Claim 4.14).

Suppose now that T is an edge with one of its endpoints being adjacent to vertices in $H_n(C)$. Hence, T is a (1, k)-blocking star for some $k \in \mathbb{N}$. But in fact, $k > \ell'_{\lambda}$ as $\omega(n) \to +\infty$ and by Lemma 4.7 *iii*. a.a.s. there are no (1, k)-blocking stars with $k \leq \ell'_{\lambda}$. Such a (1, k)-blocking star with $k > \ell'_{\lambda}$, will have its k connectors inside $H_n(C)$. But recall that these will arrive at unanimity after step R and will start alternating simultaneously between states i^* and $1 - i^*$. So by Claim 4.2, the (1, k)-blocking star will synchronise with them.

Finally, suppose that $T = v_1 v_2$ is an edge where both its endpoints v_1 and v_2 have at least one neighbour in $H_n(C)$. Let $t \ge R$ be a step at which $S_t(v) = i^*$, for all $v \in H_n(C)$. Assume that v_1 and v_2 are not unanimous with $H_n(C)$. In particular, suppose that $S_t(v_1) =$ $S_t(v_2) = 1 - i^*$. Vertex v_1 will not switch strategy, if $n_t(v_1; 1 - i^*) \le \lambda^{2i^*-1} n_t(v_1; i^*)$. But $n_t(v_1; 1 - i^*) = 1$, $n_t(v_1; i^*) \ge 1$ and $\lambda^{2i^*-1} > 1$. So the inequality is satisfied. The same holds for v_2 . Thereby, $S_{t+1}(v_1) = S_{t+1}(v_2) = 1 - i^*$ and as $S_{t+1}(v) = 1 - i^*$ for all $v \in H_n(C)$, thereafter v_1, v_2 will be synchronised with $H_n(C)$.

Suppose now that $S_t(v_1) = 1 - i^*$ but $S_t(v_2) = i^*$. Then $S_{t+1}(v_1) = 1 - i^*$ as v_1 has no neighbours who play strategy $1 - i^*$. Also, $S_{t+1}(v_2) = 1 - i^*$, since $n_t(v_2; i^*) > \lambda^{1-2i^*}n_t(v_2; 1-i^*)$. The latter holds since $n_t(v_2; i^*) \ge 1$, $n_t(v_2; 1-i^*) = 1$ and $\lambda^{1-2i^*} < 1$. As $S_{t+1}(v) = 1 - i^*$ for all $v \in H_n(C)$, thereafter v_1, v_2 will stay synchronised with $H_n(C)$. By symmetry, analogous argument can be used for the case $S_t(v_1) = i^*$ but $S_t(v_2) = 1 - i^*$. We thus conclude that if $\omega(n) \to +\infty$, then $u_n^{(1)} \to 1$ as $n \to +\infty$.

Finally, suppose that $\omega(n) \to c \in \mathbb{R}$ as $n \to +\infty$. By Lemma 4.7 *ii.*, the random variable $X_{1,\ell'_{\lambda},n}^{(1)}$ converges in distribution as $n \to +\infty$ to a random variable distributed as $\operatorname{Po}(e^{2c}/\ell'_{\lambda}!)$. Thus, for any integer $k \geq 0$, we have

$$\mathbb{P}(X_{1,\ell'_{\lambda},n}^{(1)}=k) \to \mathbb{P}\left(\operatorname{Po}(e^{2c}/\ell'_{\lambda}!)=k\right)$$

as $n \to +\infty$.

Suppose now that $X_{1,\ell_{\lambda},n}^{(1)} = k$, for some $k \in \mathbb{N}_0$. The case k = 0 was treated above and unanimity is attained a.a.s. (on the conditional space where $X_{1,\ell_{\lambda},n}^{(1)} = 0$). Let us consider the case $k \ge 1$. If an $(1,\ell_{\lambda}')$ -blocking star is initially set to $(i^*, 1-i^*)$ -configuration, then by Claim 4.1 it will stay in this configuration forever. In other words, if unanimity is achieved, then no $(1,\ell_{\lambda}')$ -blocking star is initially set to $(i^*, 1-i^*)$ -configuration. The probability of this is $(1-1/4)^k = (3/4)^k$. Consequently,

$$\limsup_{n \to +\infty} u_n^{(1)} \le \sum_{k=0}^{\infty} \left(\frac{3}{4}\right)^k \mathbb{P}\left(\operatorname{Po}(e^{2c}/\ell_{\lambda}'!) = k\right).$$

Since $H_n(C)$ will reach unanimity in at most $\beta \log n$ steps, the above argument implies that a.a.s. if $L_1(G(n, d/n))$ reaches unanimity, then this will happen in at most $\beta \log n + O(1)$ steps.

5. Unbiased node systems in dense regimes: the $\lambda = 1$ case

5.1. Majority and minority dynamics. We recall that $S_t(v)$ is the state of a vertex v at a discrete time-step $t \ge 0$. We consider a process running on a suitably dense realisation of G(n, p), and also utilise the initial configuration $S_{1/2}$. If our interacting node system

 $(G(n, p), Q, \mathcal{S}_{1/2})$ is in the majority regime with $\lambda(Q) = 1$, then its evolution coincides with the majority dynamics process. The latter is defined by the following evolution rule:

$$S_{t+1}(v) = \begin{cases} 1 & \text{if } n_t(v;1) > n_t(v;0); \\ 0 & \text{if } n_t(v;1) < n_t(v;0); \\ S_t(v) & \text{if } n_t(v;1) = n_t(v;0). \end{cases}$$
(6)

In other words, in the majority dynamics process, a node will always choose to adopt the state shared by the majority of its neighbours. If there is a tie, then its state will remain unchanged. Goles and Olivos [19] showed that majority dynamics on a finite graph becomes eventually periodic with period at most 2. More specifically, there is a t_0 depending on the graph such that for any $t > t_0$ and for any vertex v we have $S_t(v) = S_{t+2}(v)$. Majority dynamics is also a special case of voting with at least two alternatives; see [31]. Results on the evolution of majority dynamics on the random graph G(n, p) were obtained recently by Benjamini et al. [8]. The last author in collaboration with Kang, and Makai [16] proved the following theorem confirming the rapid stabilisation of the majority dynamics process on a suitably dense G(n, p), confirming a conjecture stated in [8]. This result was also proved later by Berkowitz and Devlin [9] through a different approach. Let M_0 be the most popular vertex state seen across the initial configuration.

Theorem 5.1 [16]. For all $\varepsilon \in [0, 1)$ there exist Λ , n_0 such that for all $n > n_0$, if $p \ge \Lambda n^{-\frac{1}{2}}$, then G(n, p) is such that with probability at least $1 - \varepsilon$, across the product space of G(n, p) and $S_{1/2}$, the vertices in V_n following the majority dynamics rule, unanimously have state M_0 after four rounds.

We note that Theorem 5.1 allows us to conclude the stabilisation of the interacting node system with $\lambda = 1$ in the majority regime.

On the other hand, if we consider an interacting node system in the minority regime with $\lambda = 1$, then the process coincides with the *minority dynamics process*, described by the following:

$$S_{t+1}(v) = \begin{cases} 1 & \text{if } n_t(v;1) < n_t(v;0); \\ 0 & \text{if } n_t(v;1) > n_t(v;0); \\ S_t(v) & \text{if } n_t(v;1) = n_t(v;0). \end{cases}$$
(7)

Under these rules, nodes will update to the state shared by the minority of their neighbours. It can be readily checked from (1) and (2), respectively, that the evolution of the above systems are identical to an interacting node system with $\lambda = 1$.

We show that in the minority regime unanimity is also achieved within at most four rounds too. However, (7) implies that vertex strategies will alternate synchronously with period two. Our theorem concerning the evolution of minority dynamics is analogous to Theorem 5.1.

Theorem 5.2. For all $\varepsilon \in (0,1]$ there exist Λ, n_0 such that for all $n > n_0$, if $p \ge \Lambda n^{-\frac{1}{2}}$, then G(n,p) is such that with probability at least $1 - \varepsilon$, across the product space of G(n,p) and $S_{1/2}$, the vertices in V_n following the minority dynamics rule will unanimously have the same state after four rounds.

Throughout this section we consider p = d/n where $d \ge \Lambda\sqrt{n}$, for a suitably large constant Λ . Thus, in comparison to the previous section, we will now work in a denser regime. We will comment on sparser regimes in the discussion section of our paper. We consider an initial configuration of $S_{1/2}$, and apply the evolution rules from (6) in the majority regime, or the evolution rules from (7) in the minority regime. We refer to the node system using evolution rule (6) as the majority game which we denote as $(G(n, p), S_{1/2})^>$; while we refer to the system given by evolution rule (7) as the minority game, denoted $(G(n, p), S_{1/2})^<$. We show that in the minority game, unanimity is achieved after at most four rounds with high probability. As noted above, the majority game will give rise to stability, while the minority game produces a periodic system with period two.

The quantity $\eta_t := ||P_t| - |N_t||$ represents the size of the majority of the dominant strategy at time t. Due to the distribution of the $S_{1/2}$, we have with probability $1 - \varepsilon$ that the quantity $\eta_0 = ||P_0| - |N_0||$ will be sufficiently bounded away from zero.

Lemma 5.3 [16]. Given $\varepsilon > 0$, set $c = c(\varepsilon) = \sqrt{2\pi}\varepsilon/20$. Then across the probability space $S_{1/2}$,

$$\mathbb{P}\left[\eta_0 \ge 2c\sqrt{n}\right] \ge 1 - \varepsilon/4,$$

for any n sufficiently large.

The proof of this lemma is a direct consequence of the Local Limit Theorem (see Theorem 5.8 below). We define \mathcal{E}_c^+ to be the event that $|P_0| - |N_0| \ge 2c\sqrt{n}$, and \mathcal{E}_c^- to be the event that $|N_0| - |P_0| \ge 2c\sqrt{n}$. The events \mathcal{E}_c^+ and \mathcal{E}_c^- occur with equal probability; therefore by symmetry we may condition on either \mathcal{E}_c^+ or \mathcal{E}_c^- , without loss of generality.

5.2. Some results on the majority regime. We will consider a selection of results from [16] (Lemmas 3.5, 3.6 therein), which will be useful in the minority regime analysis. The first result concerns the expectation and variance of $n_1(v; 1)$, given that \mathcal{E}_c^+ has occurred.

For a vertex $v \in V_n$ let $\mathcal{N}(v) = \{ |d(v) - d| < d^{2/3} \}.$

Lemma 5.4. Consider the majority game $M = (G(n, p), S_{1/2})^>$ where p = d/n. Let $c = c(\varepsilon)$ be the constant given in Lemma 5.3. Then there exists a constant ζ , (independent of ε) such that for any $v \in V_n$ and any n sufficiently large the following holds: for any configuration $s_0 \in \mathcal{E}_c^+$ and any $k \in \mathbb{N}$ such that $|k - d| < d^{2/3}$:

$$\mathbb{E}\left[n_1(v;1) \mid \mathcal{S}_{1/2} = s_0, d(v) = k\right] \ge \frac{k}{2} + \frac{\zeta c}{7} \left(\frac{d^3}{n}\right)^{1/2}$$

Moreover, there exists a positive constant α , such that for any $k \in \mathbb{N}$ with $|k - d| < d^{2/3}$ we have

$$\operatorname{Var}\left[n_1(v;1) \mid \mathcal{S}_{1/2} = s_0, d(v) = k\right] \le \alpha d.$$

By applying Lemma 5.4, we now proceed to prove an adjustment to a result from [16]. In the modified result, we show that with high probability a vertex v will have a $n_1(v; 1)$ sufficiently bounded away from d(v)/2. In [16], the authors show that this quantity is at least d(v)/2. However, in order for us to utilise this result for the minority game, we will instead require $n_1(v; 1) \ge d(v)/2 + 2\gamma\sqrt{d}$, with high probability, for some positive constant γ . We elaborate on the reasoning behind this assertion in Section 5.3.

Lemma 5.5. Let $\varepsilon > 0$, and c a positive constant as given in Lemma 5.3. Consider the majority game $M = (G(n, p), \mathcal{S}_{1/2})^{>}$ where p = d/n. For any positive constant γ there exists a positive constant $\Lambda = \Lambda(\gamma, \varepsilon)$, such that for all n sufficiently large if $d \ge \Lambda n^{1/2}$ and $v \in V_n$, the followings holds:

$$\mathbb{P}\left[n_1(v;1) < \frac{d(v)}{2} + 2\gamma\sqrt{d} \mid \mathcal{E}_c^+\right] < \varepsilon.$$

Proof. This argument is a direct application of Chebyshev's inequality. Fix $s_0 \in \mathcal{E}_c^+$ and $k \in \mathbb{N}$ such that $|k - d| < d^{2/3}$. By applying Lemma 5.4 and subtracting $2\gamma\sqrt{d}$, from both sides we have that:

$$\left| \mathbb{E}\left[n_1(v;1) \mid \mathcal{S}_{1/2} = s_0, d(v) = k \right] - \left(\frac{d(v)}{2} + 2\gamma\sqrt{d} \right) \right| \ge \left(\frac{d^3}{n} \right)^{1/2} \frac{\zeta c}{7} - 2\gamma\sqrt{d}.$$

By Lemma 5.4 we have Var $[n_1(v; 1) | S_{1/2} = s_0, d(v) = k] \leq \alpha d$. We now apply Chebyshev's inequality to bound the probability that $n_1(v; 1) < d(v)/2 + 2\gamma\sqrt{d}$. This gives

$$\mathbb{P}\left[n_1(v;1) < \frac{d(v)}{2} + 2\gamma\sqrt{d} \mid \mathcal{S}_{1/2} = s_0, d(v) = k\right] \leq \frac{49\alpha d}{\left(\zeta c \left(\frac{d^3}{n}\right)^{1/2} - 14\gamma\sqrt{d}\right)^2} \leq \frac{49\alpha}{\zeta c \left(\frac{d}{\sqrt{n}}\right) \left[\zeta c \left(\frac{d}{\sqrt{n}}\right) - 28\gamma\right]}$$

We now recall that $d > \Lambda \sqrt{n}$. If we take $\Lambda > (1 + 28\gamma)/(\zeta c)$, then $\zeta c (d/\sqrt{n}) - 28\gamma \ge 1$. Applying this inequality to the denominator, and choosing $\Lambda > 2 \cdot \max\{(1 + 28\gamma)/\zeta c, 49\alpha/\varepsilon c\zeta\}$ we have:

$$\mathbb{P}\left[n_1(v;1) < \frac{d(v)}{2} + 2\gamma\sqrt{d} \mid \mathcal{S}_{1/2} = s_0, d(v) = k\right] \le \frac{49\alpha}{\Lambda c\zeta} \le \varepsilon/2.$$

Integrating over all possible choices of $s_0 \in \mathcal{E}_c^+$ and k such that $\mathcal{N}(v)$ is realised, we obtain

$$\mathbb{P}\left[n_1(v;1) < \frac{d(v)}{2} + 2\gamma\sqrt{d} \mid \mathcal{E}_c^+, \mathcal{N}(v)\right] \le \varepsilon/2.$$

But $\mathbb{P}[\mathcal{N}(v)] = 1 - o(1)$, (which follows by a standard application of the Chernoff bound (3)), we deduce that for *n* sufficiently large we have

$$\mathbb{P}\left[n_1(v;1) < \frac{d(v)}{2} + 2\gamma\sqrt{d} \mid \mathcal{E}_c^+\right] \le \varepsilon.$$

5.3. **Minority Regime.** We now work within the minority regime, proving Theorem 5.2. Recall that by (7), a vertex will update to the state shared across the minority of its neighbours. In the event of a tie, the vertex will remain in its current state.

We observe similarities with Theorem 5.1, namely the fact that unanimity occurs, and is achieved within at most four rounds. However the system is no longer stable, but will become periodic with period two after unanimity is reached.

We wish to relate the proof of Theorem 5.2 to Theorem 5.1. We first start by showing that the first round of the minority game $m = (G(n, p), S_{1/2})^{<}$ can be approximated by a specific majority game which starts on the *complementary configuration*. For a configuration Son V_n we define the *complementary configuration* \bar{S} as follows: for every $v \in V_n$ we set $\bar{S}(v) = 1 - S(v)$. Suppose we have an interacting node system $\mathcal{I} = (G, S_{1/2})^*$, where $* \in \{<,>\}$. For $v \in V_n$, we denote $S_t^{\mathcal{I}}(v)$ to be the strategy of a vertex v in the game \mathcal{I} at time t; similarly we define $n_t^{\mathcal{I}}(v; i) = |\{u : S_t^{\mathcal{I}}(u) = i\}|$. While these definitions are similar to their counterparts, we would like to emphasise the role of \mathcal{I} , as we will generally work with two different systems: a minority game $\mathcal{I} = m$ (when * is <) and a majority game $\mathcal{I} = M$ (when * is >). We now state the following lemma concerning the approximation of a minority game m to a suitably designed majority game M.

Lemma 5.6. Let G = (V, E) be a graph and $S : V \to \{0, 1\}$ be a configuration. Let $m = (G, S)^{<}$ be a minority game, and $M = (G, \overline{S})^{>}$ be a majority game with initial configuration \overline{S} . If for $v \in V(G)$ we have that $n_0^m(v; 0) \neq n_0^m(v; 1)$, then $S_1^m(v) = S_1^M(v)$. If $n_0^m(v; 0) = n_0^m(v; 1)$, then $S_1^m(v) = 1 - S_1^M(v)$.

Lemma 5.6 allows us to deduce the behaviour of a significant number of vertices in the first round of the minority process. The idea is to take a minority game, complement each of the vertex strategies, and then allow one round of the evolution to occur using the majority rules. As long as a vertex satisfies the condition $n_0^m(v; 0) \neq n_0^m(v; 1)$, it will have the same state as if it had just evolved using the minority rules on the original configuration. We refer to the additional condition $n_0^m(v; 0) = n_0^m(v; 1)$, as the equal neighbourhoods condition (ENC).

Proof of Lemma 5.6. We first assume that $n_0^m(v; 0) \neq n_0^m(v; 1)$. We split our analysis into cases which depend on both the current state of the vertex, along with which of $n_0^m(v; 0)$ and $n_0^m(v; 1)$ is larger. In all four cases the argument is identical; we simply must show that $S_1^m(v) = S_1^M(v)$.

Suppose $S_0^m(v) = 0$, and $n_0^m(v; 0) > n_0^m(v; 1)$. By applying the minority rules, we see that $S_1^m(v) = 1$. We now consider the complementary state, $\bar{S}_0^m(v) = S_0^M(v) = 1$. As we have that $n_0^m(v; 0) > n_0^m(v; 1)$, then by the definition of complementary initial configuration, it must be the case that $n_0^M(v; 0) < n_0^M(v; 1)$. By applying the majority rules to the vertex v we have that $S_1^M(v) = 1$; therefore, $S_1^m(v) = S_1^M(v)$.

For the case where $n_0^m(v; 0) = n_0^m(v; 1)$, we observe that in both games v has an equal of number of the vertices playing strategy one and zero in its neighbourhood. Therefore, it follows that $S_0^m(v) = S_1^m(v)$ and $S_0^M(v) = S_1^M(v)$. However, by the definition of complementary states we have that $S_0^m(v) = 1 - S_0^M(v)$, and thus $S_1^m(v) = 1 - S_1^M(v)$. \Box

The main application of the above lemma is to connect the games m and M on G(n, d/n) with initial configuration $S_{1/2}$. It is at that point where we apply Lemma 5.5 to the game m. However, we must consider which vertices satisfy the equal neighbourhoods condition. For each vertex $v \in V_n$, we say that $v \in \text{ENC}$ if $n_0^m(v; 0) = n_0^m(v; 1)$, and we define $\text{EQ}(v) = |\{w : w \in \text{ENC}\} \cap N_{G(n,d/n)}(v)|$. Let γ be a positive constant. We say that a vertex $v \in V_n$ has a γ -decisive neighbourhood in G(n, d/n) if $\text{EQ}(v) < \gamma \sqrt{d}$, and a vertex has a γ -abundant neighbourhood in G(n, d/n) if $n_1^M(v; 1) \ge 2\gamma\sqrt{d} + d(v)/2$. We say that a vertex $v \in V_n$ is γ -good if v has a γ -abundant and a γ -decisive neighbourhood in G(n, d/n). The following corollary illustrates the role of γ -good vertices.

Corollary 5.7. Let $\gamma > 0$. If a vertex $v \in V_n$ is γ -good in G(n, d/n) for a given initial configuration, then $S_2^m(v) = 0$.

Proof. Let us abbreviate G(n, d/n) by G. Given an initial configuration on V_n , set up the systems m and M as in Lemma 5.6, and assume that $v \in V_n$ is γ -good vertex. We show that $n_1^m(v;1) > d(v)/2$; we proceed by applying Lemma 5.6 to bound $n_1^m(v;1)$ from below. This lemma implies that if $u \notin \text{ENC}$, then $S_1^m(u) = S_1^M(u)$. Hence, for all $u \in N_G(v) \setminus \text{ENC}$, if $S_1^M(u) = 1$, then $S_1^m(u) = 1$. However for $w \in \text{ENC} \cap N_G(v)$, we have that $S_1^m(w) = 1 - S_1^M(w)$. If we assume that for all $w \in \text{ENC} \cap N_G(v)$ we have that $S_1^M(w) = 1$, then we would minimise the size of $n_1^m(v;1)$ as in that case $S_1^m(w) = 0$. As a direct consequence, we have that $n_1^m(v;1) \ge n_1^M(v;1) - \text{EQ}(v)$. Therefore, by combining this bound with the definitions of γ -abundance and γ -decisiveness, we have:

$$n_1^m(v;1) \ge n_1^M(v;1) - \operatorname{EQ}(v) > n_1^M(v;1) - \gamma\sqrt{d} \ge \frac{d(v)}{2} + 2\gamma\sqrt{d} - \gamma\sqrt{d}$$
$$= \frac{d(v)}{2} + \gamma\sqrt{d}.$$

5.4. Bounding the size of EQ(v). In light of Corollary 5.7, our aim is to show that there are a significant number of good vertices. We first show that with high probability, there are a sufficient number of vertices with a γ -decisive neighbourhood. The proof of this bound will require us to invoke the Local Limit Theorem for sums of Bernoulli-distributed random variables, which follows from Theorem VII.6 p.197 in [35]. We state this result as follows.

Theorem 5.8 [35]. Let X_1, \ldots, X_n be independent identically distributed Bernoulli-distributed random variables with $\mathbb{E}(X_1) = \mu$ and $\operatorname{Var}(X_1) = \mu - \mu^2 =: \sigma^2 > 0$. Let also $X = \sum_{i=1}^n X_i$. There exists ρ depending on μ for which:

$$\sup_{i \in \mathbb{N}_0} \left| \sqrt{\operatorname{Var}[X]} \cdot \mathbb{P}[X=i] - \frac{1}{2\pi} \exp\left(-\frac{(i-\mathbb{E}[X])^2}{2\operatorname{Var}[X]}\right) \right| < \frac{\rho}{\sqrt{\operatorname{Var}[X]}}.$$

The following lemma provides an upper bound on the number of vertices in the neighbourhood of $v \in V_n$ in G(n, p) which satisfy the equal neighbourhoods condition.

Lemma 5.9. Consider G(n,p) with $np =: d = d(n) \to \infty$ as $n \to \infty$ and let $v \in V_n$. For every $\varepsilon > 0$, there exist positive constants γ and n_0 such that for all $n > n_0$ we have

$$\mathbb{P}[\mathrm{EQ}(v) \ge \gamma \sqrt{d}] < \varepsilon.$$

Proof. For a vertex $v \in V_n$ we bound the value of $\mathbb{E}[\mathrm{EQ}(v)]$. Without loss of generality we may also assume that $S_0(v) = 1$. For a vertex $w \in V_n$ to belong to EQ(v), we must have that $w \in N_{G(n,p)}(v)$ and $w \in ENC$. Therefore, we write

$$\mathbb{E}[\mathrm{EQ}(v) \mid S_0(v) = 1] = \mathbb{E}\left[\sum_{w:w\neq v} \mathbb{1}_{\{w \sim v\}} \mathbb{1}_{\{w \in \mathrm{ENC}\}} \mid S_0(v) = 1\right]$$
$$= \sum_{w:w\neq v} \mathbb{P}\left[\{w \sim v\} \cap \{w \in \mathrm{ENC}\} \mid S_0(v) = 1\right].$$

By conditioning on the event $\{w \sim v\}$, we have the following:

$$\mathbb{E}[\mathrm{EQ}(v) \mid S_0(v) = 1] = \sum_{w:w \neq v} \mathbb{P}[w \sim v] \mathbb{P}[w \in \mathrm{ENC} \mid w \sim v, S_0(v) = 1]$$
$$= \frac{d}{n} \sum_{w:w \neq v} \mathbb{P}[w \in \mathrm{ENC} \mid w \sim v, S_0(v) = 1].$$

We now condition on the size of $N_{G(n,p)}(w) \setminus \{v\}$. The size of $N_{G(n,p)}(w) \setminus \{v\}$ can vary from zero to n-2. We set $\mu' = \mathbb{E}[\mathrm{EQ}(v)]$ and let $\mathcal{N}_k(w)$ be the event that $|N_{G(n,p)}(w) \setminus \{v\}| = k$. Note that for $w \in ENC$, we necessarily have k is odd. Applying the law of total probability and bounding the upper and lower extremes of k, we have that:

$$\mathbb{P}[w \in \text{ENC} \mid w \sim v, S_0(v) = 1] \leq \mathbb{P}\left[\left|\left|N_{G(n,p)}(w) \setminus \{v\}\right| - \mu'\right| \geq d^{3/4}\right] + \sum_{k : |k-\mu'| \leq d^{3/4}, k \text{ odd}} \mathbb{P}\left[\left|N_{G(n,p)}(w) \setminus \{v\}\right| = k\right] \mathbb{P}\left[w \in \text{ENC} \mid \{w \sim v\} \cap \mathcal{N}_k(w) \cap \{S_0(v) = 1\}\right].$$
(8)

As we have conditioned on $\{v \sim w\}$, we have that $|N_{G(n,p)}(w) \setminus \{v\}| \sim Bin(n-2, d/n)$. So, $\operatorname{Var}[|N_{G(n,p)}(w) \setminus \{v\}|] = (n-2)(d/n)(1-d/n)$ and therefore $\operatorname{Var}[|N_{G(n,p)}(w) \setminus \{v\}|] < d$. By Chebyshev's inequality,

$$\mathbb{P}\left[\left|\left|N_{G(n,p)}(w)\backslash\{v\}\right| - \mu'\right| \ge d^{3/4}\right] \le \frac{d}{d^{6/4}} = \frac{1}{\sqrt{d}}.$$
(9)

Recall that $\mathcal{N}_k(w)$ is the event that $|N_{G(n,p)}(w) \setminus \{v\}| = k$. It follows from (9) that: $\mathbb{E}[\mathrm{EQ}(v) \mid S_0(v) = 1] \le$ $\frac{d}{n} \sum_{w:w \neq v} \left(\frac{1}{\sqrt{d}} + \sum_{k:|k-\mu'| \leq d^{3/4}, k \text{ odd}} \mathbb{P}[\mathcal{N}_k(w)] \mathbb{P}\left[w \in \text{ENC} \middle| \{w \sim v\} \cap \mathcal{N}_k(w) \cap \{S_0(v) = 1\} \right] \right).$ We now turn our attention to the range where $\mu - d^{3/4} \leq k \leq \mu + d^{3/4}$. We wish to apply Theorem 5.8 to bound $\mathbb{P}[w \in \text{ENC} \mid \{w \sim v\} \cap \mathcal{N}_k(w) \cap \{S_0(v) = 1\}]$. For a vertex $w \in N_{G(n,p)}(v)$, we define $X_w^+ = |\{u \in N_{G(n,p)}(w) \setminus \{v\} : S_0(u) = 1\}|$. As we have conditioned on $v \sim w$ and $S_0(v) = 1$, for odd k the following holds:

$$\mathbb{P}\left[w \in \text{ENC} \mid \{w \sim v\} \cap \mathcal{N}_{k}(w) \cap \{S_{0}(v) = 1\}\right] = \mathbb{P}\left[X_{w}^{+} = (k-1)/2 \mid \{w \sim v\} \cap \mathcal{N}_{k}(w) \cap \{S_{0}(v) = 1\}\right].$$
(10)

Now for a given k, suppose we condition on the event $\mathcal{N}_k(w)$. By conditioning on $\mathcal{N}_k(w)$, we observe that $X_w^+ \sim \operatorname{Bin}(k, 1/2)$; so $\operatorname{Var}[X_w^+] = k/4$. We now apply Theorem 5.8, to bound the probability of the event given by (10): there exits a constant ρ such that for all k we have:

$$\mathbb{P}\left[X_{w}^{+}=\frac{k-1}{2}\middle| \{w\sim v\}\cap\mathcal{N}_{k}(w)\cap\{S_{0}(v)=1\}\right] \leq \frac{\rho}{\operatorname{Var}\left[X_{k}^{+}\right]}+\frac{1}{\sqrt{2\pi\operatorname{Var}\left[X_{k}^{+}\right]}}e^{-1/\left(8\operatorname{Var}\left[X_{k}^{+}\right]\right)} \leq \frac{2\rho}{\sqrt{\operatorname{Var}\left[X_{k}^{+}\right]}}=\frac{4\rho}{\sqrt{k}}.$$
(11)

Using (10) and (11) in (8), we finally deduce that $\mathbb{E}[\mathrm{EQ}(v) \mid S_0(v) = 1] <$

$$\frac{d(n-1)}{n} \left(\frac{1}{\sqrt{d}} + \max_{k : |k-\mu'| \le d^{3/4}, k \text{ odd}} \left\{ \mathbb{P} \left[X_k^+ = (k-1)/2 \mid \{w \sim v\} \cap \mathcal{N}_k(w) \cap \{S_0(v) = 1\} \right] \right\} \right)$$

Using the bound (11), the above expression becomes

$$\mathbb{E}[\mathrm{EQ}(v) \mid S_0(v) = 1] \le \frac{d(n-1)}{n} \left(\frac{1}{\sqrt{d}} + \frac{4\rho}{\sqrt{\mu - d^{3/4}}} \right).$$

A similar argument gives the same upper bound on $\mathbb{E}[\mathrm{EQ}(v) \mid S_0(v) = 0]$.

Since $\mu' = \Theta(d)$, therefore there exists a constant κ , such that for sufficiently large d we have that $\mu' - d^{3/4} \ge \kappa d$. Letting $\Gamma = 2(\sqrt{\kappa} + 4\rho)/\sqrt{\kappa}$, for sufficiently large d we have that:

$$\mathbb{E}[\mathrm{EQ}(v)] \le \frac{\Gamma d}{\sqrt{d}} = \Gamma \sqrt{d}$$

Fix $\varepsilon > 0$ and define $\gamma = \Gamma/\varepsilon$. Then by Markov's inequality

$$\mathbb{P}\left[\mathrm{EQ}(v) \ge \gamma \sqrt{d}\right] < \frac{\mathbb{E}[\mathrm{EQ}(v)]}{\gamma} \le \varepsilon.$$

5.5. A bound on the number of good vertices. We now have a suitable bound on EQ(v), which holds with high probability. Furthermore, we would like to show that there are a significant number of good vertices. We denote the set of γ -good vertices in G(n, p) as GD_{γ} . We recall that a vertex $v \in V_n$ of G(n, p) is γ -good if v has a γ -abundant and a γ -decisive neighbourhood. The following corollary asserts that our node system will have a significant number of good vertices. We also recall the definition of the definition of the

event \mathcal{E}_c^- : we say that the event \mathcal{E}_c^- occurs if $N_0 - P_0 \ge 2c\sqrt{n}$, where $P_0 = \sum_{v \in V_n} S_0(v)$ and $N_0 = n - P_0$.

Corollary 5.10. Let $\varepsilon > 0$ and $m = (G(n, d/n), S_{1/2})^{<}$, where $d > \Lambda n^{1/2}$. Suppose that \mathcal{E}_{c}^{-} has occurred for some constant c > 0. Then there exist positive constants Λ and γ , such that for all n large enough we have with probability at least $1 - \varepsilon$ that $|\text{GD}_{\gamma}| \ge n(1 - \varepsilon)$.

Proof. We fix $\varepsilon > 0$ and again define the majority game $M = (G(n, d/n), \overline{S}_{1/2})^{>}$ as above. As we have conditioned on \mathcal{E}_{c}^{-} in m, then M starts with an initial configuration which satisfies \mathcal{E}_{c}^{+} . By Lemma 5.9, there exists a constant γ such that for all n sufficiently large and all $v \in V_n$, we have that $\mathbb{P}\left[\mathrm{EQ}_{\gamma}(v) \geq \gamma \sqrt{d}\right] < \varepsilon^2/2$. Also, by Lemma 5.5, we may select Λ large enough, such that for all n sufficiently large and $v \in V_n$, we have that $\mathbb{P}\left[n_1^M(v;1) < d(v)/2 + 2\gamma\sqrt{d} \mid \mathcal{E}_c^+\right] < \varepsilon^2/2$.

Now, we denote the events that v has a γ -decisive neighbourhood by $D_{\gamma}(v)$ and that v has an γ -abundant neighbourhood by $A_{\gamma}(v)$. Note that $D_{\gamma}^{c}(v)$ and $A_{\gamma}^{c}(v)$ are the events of the previous paragraph for this particular γ . By the union bound, we have that:

$$\mathbb{P}[v \notin \mathrm{GD}_{\gamma}] = \mathbb{P}[A_{\gamma}^{c}(v) \cup D_{\gamma}^{c}(v)] \le \mathbb{P}[A_{\gamma}^{c}(v)] + \mathbb{P}[D_{\gamma}^{c}(v)] < \varepsilon^{2}.$$

By taking a sum across all vertices, and applying the above inequality we have that $\mathbb{E}[|V_n \setminus \mathrm{GD}_{\gamma}|] \leq \varepsilon^2 n$, where $|V_n \setminus \mathrm{GD}_{\gamma}|$ is the number of vertices which are not good. By Markov's inequality, we have that $\mathbb{P}[|V_n \setminus \mathrm{GD}_{\gamma}| > \varepsilon n] \leq \varepsilon$. Therefore, with probability at least $1 - \varepsilon$, we have that $|\mathrm{GD}_{\gamma}| \geq n(1 - \varepsilon)$.

Now, if there are at least $(1 - \varepsilon)n$ good vertices (which occurs with high probability), then by Corollary 5.7 we have that $|N_2| \ge (1 - \varepsilon)n$.

5.6. The final two rounds. In the remaining two rounds we claim that unanimity will occur. We note that as a consequence of Corollary 5.10, at time t = 2 our node system satisfies the hypothesis of Lemma 4.9. However, as we are working within a denser regime, it turns out that unanimity can be reached in the entire G(n, p) much faster than $\beta \log n$ rounds. We state a result from [16], which concerns the rapid formation of unanimity for majority dynamics, wherein one of the initial strategies has a linear majority. We recall the definitions $P_t = \{v \in V_n : S_t(v) = 1\}$, and $N_t = \{v \in V_n : S_t(v) = 0\}$. Suppose that S is an initial configuration of vertex states on V_n , and $\delta \in (0, 1)$ is fixed. We say that $S \in \hat{S}_{\delta}$, if we have that $|N_0| \ge (1 - \delta)n$ in S. Thus \hat{S}_{δ} is the collection of all initial vertex configurations where there are at least $(1 - \delta)n$ vertices in the zero state. We now consider the following result considering the two round evolution of games utilising initial states belonging to \hat{S}_{δ} for sufficiently small δ .

Lemma 5.11. Let $p \ge \Lambda n^{-\frac{1}{2}}$ and $\delta < 1/10$. On the probability space of G(n,p) we have that a.a.s. for any $S \in S_{\delta}$ the majority game $(G(n,p),S)^{>}$ will reach unanimity after at most two rounds.

Proof of Theorem 5.2. While the above lemma directly concerns the majority dynamics game, with only some minimal adjustments to the argument, we may apply it to the

minority game as well. We fix a minority game $(G, \mathcal{S})^{<}$, where $\mathcal{S} \in \hat{\mathcal{S}}_{\delta}$. By following the proof given in [16], after one round of applying the minority game rules we have with probability 1 - o(1) that $|N_1| \leq d/10$. However a standard application of the Chernoff bound can show that a.a.s. the minimum degree of G(n, p) is at least d/2. Therefore for all $v \in V_n$ we have that $n_1(v; 1) > n_1(v; 0)$. In turn, we deduce that $S_2(v) = 0$ for all $v \in V_n$. As unanimity is achieved, it is clear that the system will also display periodic behaviour of period two from the evolution rules of the minority dynamics (7).

6. DISCUSSION

In this paper we study the evolution of games on G(n, p) under the best response rule. Our first result concerns the rapid formation of unanimous strategies on the node system $(G(n, p), Q, S_{1/2})$ for $p \ge \Lambda n^{-1/2}$. As a byproduct of this analysis, we also prove an analogous result of Fountoulakis et al. [16] regarding the rapid formation of unanimity in the random graph minority game. Our second main result concerns the formation of unanimous strategies in sparser regimes, namely beyond the connectivity threshold, and on the presence of a strategic bias given by $\lambda \neq 1$. A natural question of the sparse regime is to ask whether we can remove the condition that $\lambda \neq 1$. As previously seen, the case $\lambda = 1$ reduces to the problem of majority and minority dynamics. However the study of majority dynamics for $p = o(n^{-1/2})$ imposes immense complications. Little is currently known about how these systems evolve apart from becoming eventually periodic. However the following was conjectured by Benjamini et al. [8].

Conjecture 6.1 [8]. With high probability over choice of random graph and choice of initial state, if $p \ge d/n$ then the following holds:

(1) If $d \gg 1$ then, for any $\varepsilon > 0$ and n sufficiently large we have,

$$\lim_{t \to \infty} \left| |P_{2t}| - |N_{2t}| \right| \in \left[(1 - \varepsilon)n, n \right].$$

(2) If d is bounded then for any $\varepsilon > 0$ and for n sufficiently large,

$$\lim_{t \to \infty} \left| |P_{2t}| - |N_{2t}| \right| \in \left[(1 - \varepsilon/2)n, (1 + \varepsilon/2)n \right].$$

These results suggest the idea of long term almost unanimity, when $d \gg 1$. Theorem 3.2 can be thought of as an approximate approach to studying these kinds of systems. By introducing the strategic bias $\lambda \neq 1$, we are able to show a stronger form of (1) in the above conjecture. In fact, we have been able to identify precisely the critical density of the random graph around which its largest connected component (which contains the overwhelming majority of its vertices) achieves unanimity. Furthermore, when non-unanimity occurs, we identified those substructures which play different strategies to the majority of the vertices in this component. However, it is still unclear how to approach these sparser interacting node systems when $\lambda = 1$.

Another direction for consideration is to remove the synchronicity of the decision updating. For the asynchronous setting, in each round we randomly and uniformly select a node from the network and update its strategy. Such dynamics have been considered in [4, 32]. A natural question is whether unanimity can be achieved in G(n, p) under asynchronous majority dynamics, or even best response, and determine bounds on how many steps this requires. If unanimity is indeed achieved, then we can observe this must happen in at least $\Omega(n \log n)$ steps. In any initial configuration there are at least a linear number of vertices in the "wrong state", and they must be sampled at some time-step during the process. Due to the way we sample the network, checking each one of these contrarian vertices follows a coupon-collector-like scheme.

Another direction for further study is the setting where players possess more than 2 strategies. For example, this setting is considered by Oyama and Takashaki [34]. The authors consider a modification of a game with two strategies A and B played on a network; further to this, players can also play a *bilingual strategy* AB, whose payoff is a combination of the payoffs of A and B respectively. In [34], conditions on contagion and impenetrability are considered for underlying graphs which are infinite. For more general settings, it is not clear whether one of the strategies prevails eventually or we have the co-existence of two or more strategies. Furthermore, it is not clear how our conditions can be generalised for higher-order pay-off matrices.

APPENDIX A. DERIVATION OF EVOLUTION RULES

In this section we briefly detail the derivation of equations (1) and (2). We recall that if an agent chooses strategy i in round t, then the incentive for them to switch to strategy 1 - i in round t + 1, is given by the following condition:

$$n_t(v;0) (q_{i,0} - q_{1-i,0}) < n_t(v;1) (q_{1-i,1} - q_{i,1}).$$

As we have assumed that Q is non-degenerate, we have two cases: Either $q_{0,0} > q_{1,0}$ and $q_{1,1} > q_{0,1}$, or $q_{0,0} < q_{1,0}$ and $q_{1,1} < q_{0,1}$. Suppose that the former case occurs, then we must check the incentive for i = 0 an i = 1. Suppose $S_t(v) = 0$, then the incentive for changing strategy is given as follows:

$$n_t(v; 0) (q_{0,0} - q_{1,0}) < n_t(v; 1) (q_{1,1} - q_{0,1}).$$

We now re-arrange the above, and recall that $\lambda = (q_{1,1} - q_{0,1})/(q_{0,0} - q_{1,0})$, to form the first evolution rule:

$$n_t(v;0) < n_t(v;1) \frac{q_{1,1} - q_{0,1}}{q_{0,0} - q_{1,0}} = n_t(v;1)\lambda.$$
 (A1)

Similarly if we instead have that $S_t(v) = 1$, then the incentive is expressed as:

$$n_t(v;0) \left(q_{1,0} - q_{0,0} \right) < n_t(v;1) \left(q_{0,1} - q_{1,1} \right).$$

We recall that in this case we have $q_{0,1} < q_{1,1}$, and therefore we re-arrange as follows:

$$n_t(v;1) < n_t(v;0) \frac{q_{1,0} - q_{0,0}}{q_{0,1} - q_{1,1}} = n_t(v;0) \frac{q_{0,0} - q_{1,0}}{q_{1,1} - q_{0,1}} = n_t(v;0) \frac{1}{\lambda}.$$
 (A2)

Combing equations (A1) and (A2) provides the required evolution rules as described by (1). For the second case, an identical argument will produce the rules given by (2).

Appendix B. The distribution of blocking stars

Lemma 4.5. Let $\ell \in \mathbb{N}$ and p = d/n, where $1 \ll d = d(n) = O(\log n)$. Then $\mathbb{E}X_{\ell,n} \sim n \frac{d^{\ell+k}}{\ell k!} e^{-d(\ell+1)}$.

Furthermore, if $d = \frac{1}{\ell+1} \log n + \frac{\ell+k}{\ell+1} \log \log n + \omega(n)$, then the following hold.

- i. If $\omega(n) \to -\infty$ as $n \to +\infty$, then $\mathbb{E}X_{\ell,k,n} \to +\infty$ as $n \to +\infty$ and moreover, a.a.s. $X_{\ell,k,n} \ge \mathbb{E}X_{\ell,k,n}/2.$
- ii. If $\omega(n) \to c \in \mathbb{R}$ as $n \to +\infty$, then $\mathbb{E}X_{\ell,k,n} \to \frac{e^{(\ell+1)c}}{\ell!k!}$ as $n \to \infty$ and $X_{\ell,k,n} \xrightarrow{d} \operatorname{Po}\left(\frac{e^{(\ell+1)c}}{\ell!k!}\right)$.
- iii. If $\omega(n) \to +\infty$ as $n \to +\infty$, then $\mathbb{P}(X_{\ell,k,n} > 0) < 2e^{-(\ell+1)\omega(n)}$, for any n sufficiently large. Thus, a.a.s. $X_{\ell,k,n} = 0$.

Proof of Lemma 4.5. We start with the expected value of $X_{\ell,k,n}$. Suppose S denotes a set of size $\ell + k + 1$ on which an (ℓ, k) -blocking star will be formed. There are $\binom{n}{\ell+k+1}$ ways to select these vertices and $(\ell + k + 1)\binom{\ell+k}{k}$ ways to select the centre v and the connector vertices u_1, \ldots, u_k . Suppose that the remaining vertices are v_1, \ldots, v_ℓ . We have $\mathbb{P}(d(v_1) = 1 | v_1 \sim v) = (1-p)^{n-1}$. For $j = 2, \ldots, \ell$,

$$\mathbb{P}(d(v_j) = 1 | v_i \sim v, d(v_i) = 1, \text{ for } i = 1, \dots, j-1) = (1-p)^{n-(j-1)} < e^{-d+d\ell/n}.$$

But since $j \leq \ell$, we have for n sufficiently large

$$e^{-d - (d/n)^2} \le (1 - p)^{n - (j-1)} < e^{-d + d\ell/n}.$$

Hence, using the assumption that $d = O(\log n)$

$$\mathbb{P}(d(v_1) = \dots = d(v_\ell) = 1 | v_i \sim v, \text{ for } i = 1, \dots, \ell) \sim e^{-d\ell}$$

Also, $\mathbb{P}(d_{V_n \setminus S}(v) = 0) = (1 - p)^{n - (\ell + k + 1)} \sim e^{-d}$ Thus, we obtain

$$\mathbb{E}X_{\ell,k,n} \sim \frac{n^{\ell+k+1}}{(\ell+k+1)!} \cdot (\ell+k+1) \binom{\ell+k}{k} \cdot \left(\frac{d}{n}\right)^{\ell+k} \cdot e^{-d(\ell+1)} \\ = \frac{n^{\ell+k+1}}{(\ell+k+1)!} \cdot (\ell+k+1) \frac{(\ell+k)!}{\ell!k!} \cdot \left(\frac{d}{n}\right)^{\ell+k} \cdot e^{-d(\ell+1)} \\ = n \frac{d^{\ell+k}}{\ell!k!} \cdot e^{-d(\ell+1)}.$$

This concludes the first part of the lemma.

Now, the value of $\lim_{n\to\infty} \mathbb{E}X_{\ell,k,n}$ is deduced as in parts *i., ii.* and *iii.* of the lemma follows by taking $d(\ell+1) = \log n + (\ell+k) \log \log n + (\ell+1)\omega(n)$, where either $\omega(n) \to +\infty$ or $\to c$ or $\to -\infty$, as $n \to \infty$, respectively.

For Part *iii.*, Markov's inequality implies that

$$\mathbb{P}(X_{\ell,k,n} > 0) \le \mathbb{E}X_{\ell,k,n} \stackrel{n \ large}{<} 2 \cdot e^{-(\ell+1)\omega(n)} \to 0, \text{ as } n \to +\infty$$

For Parts *i*. and *ii*. we will show that for any fixed integer $r \ge 2$, we have $\mathbb{E}(X_{\ell,k,n}(X_{\ell,k,n}-1)\cdots(X_{\ell,k,n}-(r-1))) \sim \mathbb{E}^r X_{\ell,k,n}$.

So the second statement in Part *i*. will follow from Chebyshev's inequality as for r = 2, the above implies that $\operatorname{Var}(X_{\ell,k,n}) = o(\mathbb{E}X_{\ell,k,n})$. The second statement in Part *ii*. will follow from Theorem 1.22 in [10] (Poisson convergence).

Consider r subsets $S_1, \ldots, S_r \subset V_n$ of size $\ell + k + 1$. They may all induce (ℓ, k) -blocking stars only if any two of them share at most k vertices. For $S \subset V_n$ let I_S be the indicator random variable that is equal to 1 if and only if S is an (ℓ, k) -blocking star. If $S_i \cap S_j = \emptyset$ for all $i \neq j$, then

$$\mathbb{P}(I_{S_1} = \dots = I_{S_r} = 1) \sim \left((\ell + k + 1) \binom{\ell + k}{k} p^{(\ell + k)} e^{-d(\ell + 1)} \right)^r.$$
(12)

There are $\prod_{i=1}^{r} \binom{n-(i-1)(\ell+k+1)}{\ell+k+1}$ to select the ordered *r*-tuple of pairwise disjoint sets (S_1, \ldots, S_r) . But

$$\prod_{i=1}^{r} \binom{n - (i-1)(\ell+k+1)}{\ell+k+1} \sim \left(\frac{n^{\ell+k+1}}{(\ell+k+1)!}\right)^{r}.$$
(13)

Now, let us assume that $S_i \cap S_j \neq \emptyset$ for some $i \neq j$. Note first that if $|S_i \cap S_j| > k$, then the two sets cannot induce (ℓ, k) -blocking stars simultaneously. If $S_i \cap S_j = \{u_1, \ldots, u_s\}$, with $s \leq k$ and S_i, S_j induce (ℓ, k) -blocking stars, then u_1, \ldots, u_s must be connector vertices in both of them. There are O(1) ways to select the vertices in $\bigcup_{i=1}^r S_i$ that will be the connectors of the (ℓ, k) -blocking stars. Let $S' \subset \bigcup_{i=1}^r S_i$ be such a choice; by the above observation, this contains any vertex which belongs to at least two members of the r-tuple. Set $S^{1..r} = \bigcup_{i=1}^r S_i$. Hence,

$$\mathbb{P}(I_{S_1} = \dots = I_{S_r} = 1) = O(1) \cdot \left[(1-p)^{\binom{\ell+k+1}{2} - (\ell+k)} p^{(\ell+k)} \right]^r \cdot \mathbb{P}(\forall v \in S^{1..r} \setminus S', d_{V_n \setminus S^{1..r}}(v) = 0).$$

Note that $|S^{1.r} \setminus S'| = r(\ell + 1)$. So, the latter probability is

$$\mathbb{P}(\forall v \in S^{1..r} \setminus S', d_{V_n \setminus S^{1..r}}(v) = 0) = (1-p)^{r(\ell+1)(n-|S^{1..r}|)} \sim e^{-dr(\ell+1)}.$$

Therefore,

$$\mathbb{P}(I_{S_1} = \dots = I_{S_r} = 1) = O(1) \cdot p^{r(\ell+k)} e^{-dr(\ell+1)}.$$
(14)

Now, such an ordered r-tuple can be selected into at most

$$\binom{n}{r(\ell+k+1)-1} = o(n^{r(\ell+k+1)})$$
(15)

ways. Thus,

$$\mathbb{E}(X_{\ell,k,n}(X_{\ell,k,n}-1)\cdots(X_{\ell,k,n}-(r-1))) = \sum_{(S_1,\dots,S_r):S_i \subset V_n, |S_i| = \ell+k+1} \mathbb{P}(I_{S_1} = \dots = I_{S_r} = 1)$$

$$= (1+o(1)) \left(\frac{n^{\ell+k+1}}{(\ell+k+1)!} \cdot \left((\ell+k+1)\binom{\ell+k}{k}\right) p^{\ell+k}e^{-d(\ell+1)}\right)^r \text{ by } (12), (13)$$

$$+o(n^{r(\ell+k+1)})p^{r(\ell+k)}e^{-dr(\ell+1)} \text{ by } (14), (15)$$

$$\sim \mathbb{E}^r X_{\ell,k,n}.$$

Lemma 4.6. We have $\mathbb{E}X_{\ell,k,n}^{(2)} = o(\mathbb{E}X_{\ell,k,n}).$

Proof of Lemma 4.6. In this lemma, we will bound the expected number of (ℓ, k) -blocking stars which do not belong to $L_1(G(n, d/n))$. Recall that the random variable which counts these is $X_{\ell,k,n}^{(2)}$.

We will give an upper bound on the expected number of connected components of order at most log n which contain an (ℓ, k) -blocking star. This suffices due to the following result about the structure of G(n, d/n).

Theorem B.1. Let p = d/n with $d \gg 1$. Then a.a.s. all connected components of G(n, d/n) apart from $L_1(G(n, d/n))$ have order at most log n.

This is a direct consequence of the proof of Theorem 6.10 (pp.143–146) in [10] and we refer the interested reader directly to this.

For $r \geq \ell + k + 1$, let $C_{\ell,k,r}$ denote the number of connected components which contain an (ℓ, k) -blocking star and have order r. We will give an upper bound on the expected value of $C_{\ell,k,r}$. For two sets $S \subset S' \subset V_n$ having $|S| = \ell + k + 1$ and |S'| = r, and $v, u_1, \ldots, u_k \in S$, we set $I(S, S', v, u_1, \ldots, u_k)$ to be the indicator random variable which is equal to 1 if and only if S' is a connected component in G(n, d/n), where in particular, Sinduces an (ℓ, k) -blocking star with centre v and connectors u_1, \ldots, u_k . Let $\mathcal{S}_{\ell,k,r}$ denote the set of pairs (S, S') such that $S \subset S' \subset V_n$ with $|S| = \ell + k + 1$ and |S'| = r. Moreover, for a set $S \subset V_n$ of size at least k + 1, let $S_{\neq}^{(k+1)}$ denote the set of all ordered k + 1-tuples of distinct vertices in S. With this notation, we can write

$$\mathbb{E}(C_{\ell,k}) \le \sum_{(S,S')\in\mathcal{S}_{\ell,k,r}} \sum_{\substack{(v,u_1,\dots,u_k)\in S^{(2)}_{\neq}}} \mathbb{E}(I(S',S,v,u_1,\dots,u_k)).$$
(16)

For $S \subset V_n$ having $|S| = \ell + k + 1$ and $(v, u) \in S_{\neq}^{(k+1)}$, we let $I_1(S, v, u_1, \ldots, u_k)$ be the indicator random variable that is equal to 1 if and only if S forms an ℓ -blocking star with centre v and connectors u_1, \ldots, u_k . Also, we take $I_2(S')$ to be the indicator random variable that is equal to 1 if and only if S' is a connected component of G(n, d/n).

For a given $(S, S') \in \mathcal{S}_{\ell,k}$ and $(v, u_1, \ldots, u_k) \in S_{\neq}^{(k+1)}$, we write

$$\mathbb{P}(I(S', S, v, u_1, \dots, u_k) = 1) = \mathbb{P}(I_1(S, v, u_1, \dots, u_k) = 1) \cdot \mathbb{P}(I_2(S') = 1 \mid I_1(S, v, u_1, \dots, u_k) = 1).$$
(17)

We will provide an upper bound on $\mathbb{P}(I_2(S') = 1 \mid I_1(S, v, u_1, \ldots, u_k) = 1)$. If S induces an (ℓ, k) -blocking star with centre v and connectors u_1, \ldots, u_k , then any spanning tree $T_{S'}$ on S' contains the star on $S \setminus \{u_1, \ldots, u_k\}$ centred at v as an induced subgraph and, moreover, one of the edges $u_i v$ is a cutting edge between $S' \setminus (S \setminus \{u_1, \ldots, u_k\})$ and $S \setminus \{u_1, \ldots, u_k\}$. Thus, $T_{S'} \setminus (S \setminus \{u_1, \ldots, u_k\})$ is a spanning tree of the subgraph induced by the set $S' \setminus (S \setminus \{u_1, \ldots, u_k\})$. Furthermore, if S' is a connected component in G(n, d/n), there are no edges between $S' \setminus (S \setminus \{u_1, \ldots, u_k\})$ and $V_n \setminus S'$.

With the above observations and p = d/n we can give the following bound:

$$\mathbb{P}(I_2(S') = 1 \mid I_1(S, v, u_1, \dots, u_k) = 1) \le (r - (\ell + 1))^{r - (\ell + 1) - 2} p^{r - (\ell + 1) - 1} (1 - p)^{(n - r)(r - (\ell + 1))}$$
(18)

We have that

$$(r - (\ell + 1))^{r - (\ell + 1) - 2} p^{r - (\ell + 1) - 1} \le (rp)^{r - (\ell + 1) - 1} = (rp)^{r - \ell - 2},$$

and for $r \leq \log n$,

$$(1-p)^{(n-r)(r-(\ell+1))} \sim e^{-dr} \leq 2e^{-d(r-\ell-1)} = 2e^{-d(r-\ell-1)}e^{-d}.$$

Using these in (18) we get

$$\mathbb{P}(I_2(S') = 1 \mid I_1(S, v, u_1, \dots, u_k) = 1) \le 2(rpe^{-d})^{r-\ell-2}e^{-d}.$$

Hence, the left-hand side in (17) is bounded as

$$\mathbb{P}(I(S', S, v, u_1, \dots, u_k) = 1) \le \mathbb{P}(I_1(S, v, u_1, \dots, u_k) = 1) \cdot (2(rpe^{-d})^{r-\ell-2}e^{-d}).$$

Thus, for n sufficiently large, (16) yields:

$$\mathbb{E}(C_{\ell,k,r}) \leq \sum_{(S,S')\in\mathcal{S}_{\ell,k,r}} \sum_{(v,u_1,\dots,u_k)\in S_{\neq}^{(k+1)}} \mathbb{P}(I_1(S,v,u_1,\dots,u_k) = 1) \cdot (2(rpe^{-d})^{r-\ell-2}e^{-d}) \\
= \sum_{S\subset V_n:|S|=\ell+k+1} \sum_{(v,u_1,\dots,u_k)\in S_{\neq}^{(k+1)}} \mathbb{P}(I_1(S,v,u_1,\dots,u_k) = 1) \times \\
\sum_{S':|S'|=r,S\subset S'} 2(rpe^{-d})^{r-\ell-2}e^{-d}.$$
(19)

For a fixed choice of S, we will provide an upper bound on the inner sum. This is

$$\begin{split} \sum_{S':|S'|=r,S\subset S'} (rpe^{-d})^{r-\ell-2}e^{-d} &= 2\sum_{r=\ell+k+1}^{\log n} \binom{n-\ell-k-1}{r-\ell-k-1} (rpe^{-d})^{r-\ell-k-1}e^{-d} \\ &\leq e^{-d}\sum_{r=\ell+k+2}^{\log n} \left(\frac{ne}{r-\ell-k-1}\right)^{r-\ell-k-1} \left(\frac{rde^{-d}}{n}\right)^{r-\ell-k-1} + e^{-d}((\ell+k+1)pe^{-d}) \\ & d \geq 1 \qquad e^{-d}\sum_{r=\ell+k+2}^{\log n} \left(\frac{ne}{r-\ell-k-1}\right)^{r-\ell-k-1} \left(\frac{rde^{-d}}{n}\right)^{r-\ell-k-1} + o(1) \\ & \overline{r}^{r}_{r-\ell-k-1} \leq e^{-d}\sum_{r=\ell+k+2}^{\log n} \left((\ell+k+2)ede^{-d}\right)^{r-\ell-k-1} + o(1) \\ &= e^{-d}\sum_{j=1}^{\log n+\ell+k+1} \left((\ell+k+2)ede^{-d}\right)^{j} + o(1) \\ & (\ell+k+2)ede^{-d} \leq e^{-d/2}, \text{ for n large} \quad e^{-d}\sum_{j=1}^{\log n+\ell+2} \left(e^{-d/2}\right)^{j} + o(1) \\ &\leq \frac{e^{-d}}{1-e^{-d/2}} + o(1) = o(1). \end{split}$$

Note that this upper bound holds for n sufficiently large uniformly over all choices of S. Thus,

$$\mathbb{E}(C_{\ell,k,r}) = o(1) \cdot \sum_{S \subset V_n : |S| = \ell + k + 1} \sum_{\substack{(v,u_1,\dots,u_k) \in S_{\neq}^{(k+1)}}} \mathbb{P}(I_1(S,v,u_1,\dots,u_k) = 1) = o(1) \cdot \mathbb{E}X_{\ell,k,n},$$

which concludes the proof of the lemma.

APPENDIX C. STRUCTURAL RESULTS IN G(n, p)

Claim C.1. Let $S \subset V_n$ be such that |S| < k, for some fixed $k \in \mathbb{N}$. Then

$$\mathbb{P}(\forall v \in S, d(v) \le C) \le \left(2C \left(de\right)^C \cdot e^{-d}\right)^{|S|}$$

Proof. If a vertex in S has degree at most C, then it has also degree at most C in $V_n \setminus S$. So we can write:

$$\mathbb{P}(\forall v \in S, d(v) \le C) \le \mathbb{P}(\forall v \in S, d_{V_n \setminus S}(v) \le C).$$

Observe that these degrees form an independent family as they are determined by mutually disjoint sets of edges. Thereby,

$$\mathbb{P}(\forall v \in S, d_{V_n \setminus S}(v) \le C) = \prod_{v \in S} \mathbb{P}(d_{V_n \setminus S}(v) \le C).$$

But $d_{V_n \setminus S}(v)$ is distributed as $\operatorname{Bin}(n-|S|, d/n)$ and, therefore, its expected value is d-o(1). Hence, $\mathbb{P}(d_{V_n \setminus S}(v) = k)$ is increasing as a function of k, if $d/k \to \infty$, as $n \to \infty$. Using $\binom{n}{k} \leq \left(\frac{ne}{k}\right)^k$ we can write the following bound:

$$\mathbb{P}(d_{V_n \setminus S}(v) \leq C) \leq C \cdot {\binom{n}{C}} \left(\frac{d}{n}\right)^C \left(1 - \frac{d}{n}\right)^{n-|S|-C} \\
\stackrel{|S| < k}{\leq} C \left(\frac{ne}{C}\right)^C \left(\frac{d}{n}\right)^C e^{-d+o(1)} \\
\leq 2C \left(\frac{de}{C}\right)^C \cdot e^{-d},$$
(20)

for n sufficiently large. Therefore,

$$\mathbb{P}(\forall v \in S, d_{V_n \setminus S}(v) \le C) \le \left(2C \left(de\right)^C \cdot e^{-d}\right)^{|S|},$$

and the claim now follows.

Lemma 4.11. Suppose that p = d/n with $c_{\lambda} \log n \leq d \leq \alpha(\lambda) \log n$. A.a.s. there are no $\ell_{\lambda} + 2$ vertices in $L_n(C)$ that have a common neighbour.

Proof. We will use a first moment argument to bound the expected number of collections of vertices in L_n of size $\ell_{\lambda} + 2$ that have a common neighbour. Let $S \subset V_n$ be a subset of vertices. We denote the degree of v outside the set S by $d_{V_n \setminus S}(v)$. Let $S \subset V_n$ be such that $S = \{u_1, \ldots, u_{\ell_{\lambda}+2}\}$ and $z \in V_n \setminus S$. The expected number of collections of $\ell_{\lambda} + 2$ vertices in $L_n(C)$ which have a common neighbour is at most

$$\binom{n}{\ell_{\lambda}+2} \cdot (n - (\ell_{\lambda}+2)) \cdot \left(\prod_{i=1}^{\ell_{\lambda}+2} \mathbb{P}(u_{i} \sim z)\right) \cdot \mathbb{P}\left(\forall i = 1, \dots, \ell_{\lambda+2} \ d_{V_{n}}(u_{i}) \leq C\right)$$

$$\stackrel{Claim \ C.1}{\leq} n^{\ell_{\lambda}+3} \cdot \left(\frac{d}{n}\right)^{\ell_{\lambda}+2} \left(2C \ (de)^{C} \cdot e^{-d}\right)^{\ell_{\lambda}+2}$$

$$= nd^{\ell_{\lambda}+2} \cdot \left(2C \ (de)^{C} \cdot e^{-d}\right)^{\ell_{\lambda}+2} = n \cdot e^{-d(\ell_{\lambda}+2)+O(\log d)}. \tag{21}$$

But $d \ge c_{\lambda} \log n = (\ell_{\lambda} + 1)^{-1} \log n$. So $d(\ell_{\lambda} + 2) - \log n = \Omega(\log n)$, whereby the above expected value is o(1).

Lemma 4.12. Let $\ell \in \mathbb{N}$ and let p = d/n where $\frac{1}{\ell+1} \log n \leq d = d(n) \leq \alpha(\lambda) \log n$. A.a.s. all connected sets of vertices in $L_n(C)$ have size at most $\ell + 1$.

The FKG inequality. Before we lead with the proof of Lemma 4.12 we also require a correlation inequality that is known as the FKG inequality. We state it in the context of the model G(n, p). We say that a graph property \mathcal{P} is non-decreasing if, for graphs G, H on V_n , whenever $G \in \mathcal{P}$ and $G \subseteq H$, then $H \in \mathcal{P}$. Similarly we say that \mathcal{P} is non-increasing, if whenever $H \notin \mathcal{P}$ and $H \subseteq G$, then $G \notin \mathcal{P}$ as well. We state the FKG inequality as follows.

Theorem C.2 The FKG inequality [1]. Let \mathcal{P}_1 be a non-decreasing graph property and \mathcal{P}_2 be a non-increasing graph property. Then for the binomial random graph, G(n, p), we have the following:

$$\mathbb{P}[G(n,p) \in \mathcal{P}_1 \cap G(n,p) \in \mathcal{P}_2] \le \mathbb{P}[G(n,p) \in \mathcal{P}_1] \cdot \mathbb{P}[G(n,p) \in \mathcal{P}_2].$$

Proof of Lemma 4.12. We will show that a.a.s. there are no connected sets of vertices in $L_n(C)$ of size $\ell + 2$ or more. If there is such a set, then in fact there must also be such a set of size exactly $\ell + 2$. So it suffices to show that a.a.s. no such subsets exist.

Let $S \subset V_n$ have $|S| = \ell + 2$. Then

$$\mathbb{P}(S \text{ is connected and } \forall v \in S, d(v) \leq C) \leq \mathbb{P}(S \text{ is connected}) \cdot \mathbb{P}(\forall v \in S, d(v) \leq C),$$

by the FKG inequality (Theorem C.2), since the graph property that $\{S \text{ is connected}\}\$ is non-decreasing whereas the property that $\{\forall v \in S, d(v) \leq C\}\$ is non-increasing. Now,

$$\mathbb{P}(S \text{ is connected}) \le |S|^{|S|-2} \cdot \left(\frac{d}{n}\right)^{|S|-1},$$

since if S induces a connected subgraph, then this has to have a spanning tree (selected in $|S|^{|S|-2}$ ways). By Claim C.1, we have

$$\mathbb{P}(\forall v \in S, d(v) \le C) \le \left(2C \left(de\right)^C \cdot e^{-d}\right)^{|S|}$$

Therefore,

$$\mathbb{P}(S \text{ is connected and } \forall v \in S, d(v) \le C) \le (\ell+2)^{\ell} \cdot \left(\frac{d}{n}\right)^{\ell+1} \cdot \left(2C \left(de\right)^C \cdot e^{-d}\right)^{\ell+2}$$

Hence, the expected number of such subsets is at most

$$O(1) \cdot {\binom{n}{\ell+2}} \cdot {\binom{d}{n}}^{\ell+1} \cdot {\binom{d^C \cdot e^{-d}}{\ell+2}} = n e^{-(\ell+2)d + O(\log \log n)}.$$

But $d \ge \frac{1}{\ell+1} \log n$. Thereby, $(\ell+2)d - \log n = \Omega(\log n)$, and the right-hand side is o(1). \Box

Lemma 4.13. Let p = d/n where $c_{\lambda} \log n \leq d = d(n) \leq \alpha(\lambda) \log n$. A.a.s. all connected sets of vertices in $L_n(C)$ induce trees.

Proof. By the previous lemma it suffices only to consider sets of size at most $\ell_{\lambda} + 1$. Let $S \subset V_n$ with $|S| \leq \ell_{\lambda} + 1$. Then

$$\mathbb{P}(e(S) \ge |S| \text{ and } \forall v \in S, d(v) \le C) \le \mathbb{P}(e(S) \ge |S|) \cdot \mathbb{P}(\forall v \in S, d(v) \le C),$$

by the FKG inequality (Theorem C.2), since the graph property that $\{e(S) \ge |S|\}$ is non-decreasing and the property that $\{\forall v \in S, d(v) \le C\}$ is non-increasing. But

$$\mathbb{P}(e(S) \ge |S|) = O(1) \cdot \left(\frac{d}{n}\right)^{|S|}$$

Combining this with Claim C.1 we get

$$\mathbb{P}(e(S) \ge |S| \text{ and } \forall v \in S, d(v) \le C) = O(1) \cdot \left(\frac{d}{n}\right)^{|S|} \left(d^C e^{-d}\right)^{|S|} = O(1) \cdot n^{-|S|} e^{-d|S| + O(\log \log n)}.$$

As there are $\binom{n}{|S|} = O(1)n^{|S|}$ choices for S, the expected number of such sets is

$$O(1)e^{-d|S|+O(\log\log n)} = o(1)$$

The lemma follows from the union bound, taking the union over all possible values of $|S| \leq \ell_{\lambda} + 1$.

Appendix D. Proof that G(n, d/n) has the minority decline property

Lemma 4.9. Let d = d(n) be such that $1 \ll d \leq \alpha(\lambda) \log n$, and let Q be a 2×2 nondegenerate payoff matrix. For any $\varepsilon > 0$ there exists $C_{\varepsilon,\lambda} \in \mathbb{N}$ such that for any $\gamma > 0$ and for any $C \geq C_{\varepsilon,\lambda}$ a.a.s. G(n, d/n) satisfies the following: for any initial configuration Swith $|m_0 \cap H_n(C)| < ne^{-\gamma d}$, the interacting node system $\mathcal{I} = (G(n, d/n), Q, S)$ will have $|m_1 \cap H_n(C)| \leq \varepsilon |m_0 \cap H_n(C)|$.

Proof. Suppose that initially the majority strategy is *i*. Assume first that \mathcal{I} is in the majority regime. Then in the random graph G(n, d/n) typically a vertex is expected to have many more neighbours among those playing strategy *i* than those playing strategy 1 - i. So one would expect that most vertices will adopt strategy *i* in the next round. If we revisit (1), we will see that if this does not happen, then $n_0(v; i) \leq \lambda^{1-2i}n_0(v; 1-i)$. Indeed, if *v* initially was playing strategy *i*, it switches to 1-i, if $n_0(v; i) < \lambda^{1-2i}n_0(v; 1-i)$. If *v* was initially playing 1 - i, then it does not switch if $n_0(v; 1 - i) \geq \lambda^{1-2(1-i)}n_0(v; i)$. Rearranging the latter, we also get that $n_0(v; i) \leq \lambda^{1-2i}n_0(v; 1-i)$.

Suppose now that \mathcal{I} is in the minority regime. In this case one would expect that most vertices will adopt strategy 1-i. Suppose that a vertex v initially plays i. By (2), it keeps on playing i after one round, if $n_0(v;i) \leq \lambda^{1-2i}n_0(v;i-1)$. Similarly, if v initially plays 1-i, then it switches to i, if $n_0(v;1-i) > \lambda^{1-2(1-i)}n_0(v;i)$. If we rearrange the latter, we get $n_0(v;i) < \lambda^{1-2i}n_0(v;1-i)$. Furthermore, to reduce notation we set $H_n := H_n(C)$, where C is to be determined later. Also, we set $L_n := L_n(C)$.

Assuming that initially the most popular strategy in H_n is *i*, we say that a vertex $v \in V_n \cap H_n$ is *i*-atypical, if $n_0(v;i) \leq \lambda^{1-2i}n_0(v;1-i)$. Let $A_n^{(i)}$ denote this set. We will show that a.a.s. provided that $|m_0| < ne^{-\gamma d}$, we have $|A_n^{(i)}| \leq \varepsilon |m_0 \cap H_n|$, for all partitions of H_n into two parts one of which has size less than $ne^{-\gamma d}$. If this happens, then all but at most $\varepsilon |m_0 \cap H_n|$ vertices in H_n will behave as expected and, therefore, $|m_1 \cap H_n| \leq \varepsilon |m_0 \cap H_n|$.

We proceed with showing the above (assuming that the majority strategy initially is *i*). So a vertex is *i*-atypical if $n_0(v; i) \leq \lambda^{1-2i}n_0(v; 1-i)$. We set $S_{1-i} = \{v \in V_n : S_0(v) = 1-i\}$. By assumption, we have that $|S_{1-i}| < ne^{-\gamma d}$ for some $\gamma \in [0, 1)$.

We will condition on the event of Lemma 4.11, which we refer to as \mathcal{D}_n . That is, \mathcal{D}_n denotes the event that no more than $\ell_{\lambda} + 1$ vertices in $L_n(C)$ have a common neighbour. According to Lemma 4.11, we have $\mathbb{P}(\mathcal{D}_n) = 1 - o(1)$. For a partition (U_i, U_{1-i}) of $H_n(C)$, we assume that the vertices in U_j are assigned strategy j, for $j \in \{0, 1\}$. As we pointed out previously, the event \mathcal{D}_n will allow us to ignore the influence of the vertices in $L_n(C)$ on the evolution of those in $H_n(C)$, provided that C is sufficiently large. To this end, we will say that a vertex is *i*-atypical with respect to (U_i, U_{1-i}) if $n_0(v; i) \leq \lambda^{1-2i} n_0(v; 1-i)$ for any initial assignment of strategies to the vertices of $L_n(C)$. We will show that a.a.s. for all configurations (U_i, U_{1-i}) of H_n with $|U_{1-i}| < ne^{-\gamma d}$ there is no collection of $\varepsilon |U_{1-i}|$ vertices in $H_n(C)$ which are *i*-atypical with respect to (U_i, U_{1-i}) . This will imply that G(n, p) is such that for any initial configuration (U_i, U_{1-i}) on $H_n(C)$, with $|U_{1-i}| < ne^{-\gamma d}$, and an arbitrary configuration for the vertices in $L_n(C)$, there will be at most $\varepsilon |U_{1-i}|$ vertices in $H_n(C)$ that will adopt strategy 1-i in the subsequent round.

We wish to bound the number of vertices in $H_n(C)$ which are *i*-atypical with respect to a given configuration (U_i, U_{1-i}) ; thus we define $\hat{S}_i = \{v \in U_i : n_0(v; 1-i) \ge \lambda^{2i-1}n_0(v; i)\}$ and $\hat{S}_{1-i} = \{v \in U_{1-i} : n_0(v; 1-i) \ge \lambda^{2i-1}n_0(v; i)\}$. If there are at least $\varepsilon |S_{1-i}|$ vertices which are *i*-atypical with respect to (U_i, U_{1-i}) , then either $|\hat{S}_i| \ge \varepsilon |U_{1-i}|/2$ or $|\hat{S}_{1-i}| \ge \varepsilon |U_{1-i}|/2$. We will show that

$$\mathbb{P}\left(\bigcup_{1 \le k < ne^{-\gamma d}} \bigcup_{(U_i, U_{1-i}): |U_{1-i}| = k} \left\{ |\hat{S}_i| \ge \varepsilon |U_{1-i}|/2 \right\} \bigcap \mathcal{D}_n \right) = o(1), \quad (22)$$

$$\mathbb{P}\left(\bigcup_{1\leq k< ne^{-\gamma d}} \bigcup_{(U_i, U_{1-i}): |U_{1-i}|=k} \left\{ |\hat{S}_{1-i}| \geq \varepsilon |U_{1-i}|/2 \right\} \bigcap \mathcal{D}_n \right) = o(1).$$
(23)

We will show that the union bound indeed suffices to show these. So, firstly, we will consider a fixed partition (U_i, U_{1-i}) as above. To bound $\mathbb{P}(|\hat{S}_i| \geq \varepsilon |U_i|/2)$ and $\mathbb{P}(|\hat{S}_{1-i}| \geq \varepsilon |U_i|/2)$, we translate the defining conditions of \hat{S}_i and \hat{S}_{1-i} into a condition on the degree of these vertices in U_{1-i} . On the event \mathcal{D}_n , there are at most $\ell_{\lambda} + 1$ neighbours of v in L_n . Consider the degree of v inside U_{1-i} , which we denote by $d_{U_{1-i}}(v)$. Similarly, we denote by $d_{S_{1-i}\cap L_n}(v)$ its degree inside $S_{1-i} \cap L_n$. Thus, $d_{U_{1-i}}(v) + d_{S_{1-i}\cap L_n}(v) = n_0(v; 1-i)$. But since \mathcal{D}_n is realised, we have $d_{S_{1-i}\cap L_n}(v) \leq \ell_{\lambda} + 1$, whereby

$$d_{U_{1-i}}(v) + \ell_{\lambda} + 1 \ge n_0(v; 1-i).$$
(24)

Furthermore, $n_0(v;i) + n_0(v;1-i) = d(v) \ge C$. If $\lambda^{2i-1}n_0(v;i) \le n_0(v;1-i)$, then

$$\frac{\lambda^{2i-1}+1}{\lambda^{2i-1}}n_0(v;1-i) \ge C.$$

We further bound $n_0(v; 1-i)$ using (24) and get

$$\frac{\lambda^{2i-1}+1}{\lambda^{2i-1}} \left(d_{U_{1-i}}(v) + \ell_{\lambda} + 1 \right) \ge C.$$

Rearranging this we deduce that

$$d_{U_{1-i}}(v) \ge \frac{\lambda^{2i-1}}{\lambda^{2i-1}+1}C - (\ell_{\lambda}+1) \ge \frac{\lambda^{2i-1}}{2(\lambda^{2i-1}+1)}C =: \psi_{\lambda}C,$$

provided that C is large enough. To summarise, we have proved that if $\lambda^{2i-1}n_0(v;i) \leq n_0(v;1-i)$ and \mathcal{D}_n is realised, then

$$d_{U_{1-i}}(v) \ge \psi_{\lambda} C. \tag{25}$$

We will start with (22). Using (25), we see that the event in (22) is included in the following event: there are disjoint set sets S, U with $1 \leq |U| < ne^{-\gamma d}$ and $|S| = \varepsilon |U|/2$ such that for any $v \in S$ we have $d_U(s) \geq \psi_{\lambda} C$.

Let us consider a set U with $1 \leq |U| \leq ne^{-\gamma d}$ and let $S \subset V_n \setminus U$ be such that $|S| = \varepsilon |U|/2$.

$$\mathbb{P}(\forall v \in S, \ d_U(v) \ge \psi_{\lambda}C) = \prod_{v \in S} \mathbb{P}(d_U(v) \ge \psi_{\lambda}C),$$

since these are events depending on pairwise disjoint sets of edges.

We now observe that for any $v \in S$ the random variable $d_U(v)$ is stochastically dominated by a random variable with distribution $\operatorname{Bin}(|U|, d/n)$. Using that $\binom{n}{k} \leq (en/k)^k$, we can bound the above probability in the following way:

$$\mathbb{P}(d_U(v) \ge \psi_{\lambda}C) \le \binom{|U|}{\psi_{\lambda}C} \left(\frac{d}{n}\right)^{\psi_{\lambda}C} \le \left(\frac{ed|U|}{\psi_{\lambda}n}\right)^{\psi_{\lambda}C}.$$

Substituting this bound into the above inequality, we finally get

$$\mathbb{P}(\forall v \in S, \ d_U(v) \ge \psi_{\lambda}C) \stackrel{|S|=\varepsilon|U|}{\le} \left(\frac{ed|U|}{\psi_{\lambda}n}\right)^{\psi_{\lambda}\varepsilon|U|C} \\ \stackrel{d\le\alpha(\lambda)\log n}{=} \exp\left(-\psi_{\lambda}\varepsilon|U|C\log(n/|U|)(1+o(1))\right).$$

Now we can bound

$$\mathbb{P}\left(\exists S : |S| = \varepsilon |U|/2, \ \forall v \in S, \ d_{U_{1-i}}(v) \ge \psi_{\lambda}C\right) \le \begin{pmatrix} n \\ \varepsilon |U|/2 \end{pmatrix} \cdot \exp\left(-\psi_{\lambda}\frac{\varepsilon}{2}|U|C\log\left(\frac{n}{|U|}\right)(1+o(1))\right) \\ \le \left(\frac{ne}{\varepsilon |U|/2}\right)^{\varepsilon |U|/2} \cdot \exp\left(-\psi_{\lambda}\frac{\varepsilon}{2}|U|C\log\left(\frac{n}{|U|}\right)(1+o(1))\right) \\ \le \exp\left(-(\psi_{\lambda}C-1)\psi_{\lambda}\frac{\varepsilon}{2}|U|\log\left(\frac{n}{|U|}\right)(1+o(1))\right).$$

We are now ready to show (22). We write

$$\mathbb{P}\left(\exists U, S: S \cap U = \emptyset, \ 1 \leq |U| < ne^{-\gamma d}, \ |S| = \varepsilon |U|/2, \ \forall v \in S, \ d_{U_{1-i}}(v) \geq \psi_{\lambda}C\right) \leq \sum_{1 \leq k < ne^{-\gamma d}} \binom{n}{k} \exp\left(-(\psi_{\lambda}C - 1)\frac{\varepsilon}{2}k\log\left(n/k\right)\left(1 + o(1)\right)\right) \\ \leq \sum_{1 \leq k < ne^{-\gamma d}} \left(\frac{ne}{k}\right)^{k} \exp\left(-(\psi_{\lambda}C - 1)\frac{\varepsilon}{2}k\log\left(n/k\right)\left(1 + o(1)\right)\right) \\ = \sum_{1 \leq k < ne^{-\gamma d}} \exp\left(k\log(n/k)\left(1 - (\psi_{\lambda}C - 1)\varepsilon/2\right)\left(1 + o(1)\right)\right) \\ \log(n/k) > \gamma d \leq \sum_{1 \leq k < ne^{-\gamma d}} \exp\left(\gamma dk(1 - (\psi_{\lambda}C - 1)\varepsilon/2)(1 + o(1))\right) = o(1),$$
(26)

provided that C is large enough, depending on ε, λ .

Now, we turn to (23). Consider a partition (U_i, U_{1-i}) of $H_n(C)$ with $|U_{1-i}|$ as specified above. If $|\hat{S}_{1-i}| > \varepsilon |U_{1-i}|/2$ and \mathcal{D}_n is realised, then there exists a set of $\varepsilon |U_{1-i}|/2$ vertices in U_{1-i} , whose degree inside U_{1-i} is at least $\psi_{\lambda}C$. Hence, the total degree of this set inside U_{1-i} must be at least $\frac{\varepsilon \psi_{\lambda}}{2} |U_{1-i}|C$. In turn, the number of edges in U_{1-i} is at least $\frac{\varepsilon \psi_{\lambda}}{4} |U_{1-i}|C$. Thus, if $e(U_{1-i})$ denotes the number of edges inside S_{1-i} we have

$$\mathbb{P}(|\hat{S}_{1-i}| \ge \varepsilon |U_{1-i}|/2|, \mathcal{D}_n) \le \mathbb{P}\left(e(U_{1-i}) \ge \frac{\varepsilon \psi_{\lambda}}{4} |U_{1-i}|C\right).$$

We will show that a.a.s. any subset $U \subset V_n$ with $1 \leq |U| < ne^{-\gamma d}$ has $e(U) \leq \frac{\varepsilon \psi_{\lambda}}{4} |U|C$. Now, e(U) is stochastically dominated from above by a binomially distributed random variable $Y \sim \text{Bin}(|U|^2, d/n)$. So

$$\mathbb{P}\left(e(U) \ge \frac{\varepsilon \psi_{\lambda}}{4} |U|C\right) \le \mathbb{P}\left(Y \ge \frac{\varepsilon \psi_{\lambda}}{4} |U|C\right).$$

We now bound the last probability as follows:

$$\mathbb{P}\left(Y \ge \frac{\varepsilon\psi_{\lambda}}{4}|U|C\right) \le \binom{|U|^2}{\frac{\varepsilon\psi_{\lambda}}{4}|U|C} \cdot \left(\frac{d}{n}\right)^{\frac{\varepsilon\psi_{\lambda}}{4}|U|C} \\ \le \left(\frac{|U|^2e}{\frac{\varepsilon\psi_{\lambda}}{4}|U|C} \cdot \frac{d}{n}\right)^{\frac{\varepsilon\psi_{\lambda}}{4}|U|C} \\ \le \left(\frac{4e}{\varepsilon\psi_{\lambda}} \cdot \frac{d}{C} \cdot \frac{|U|}{n}\right)^{\frac{\varepsilon\psi_{\lambda}}{4}|U|C}.$$

Since $d = O(\log n)$, we conclude that

$$\mathbb{P}\left(e(U) \ge \frac{\varepsilon \psi_{\lambda}}{4} |U|C\right) \le \exp\left(-(1+o(1))\frac{\varepsilon \psi_{\lambda}}{4} |U|C \log(n/|U|)\right).$$

Arguing as in the case of (22), we take the union bound over all choices of the subset U which satisfy the assumed conditions and a similar calculation as in (26) (the only difference being that $\varepsilon/2$ is replaced by $\varepsilon/4$) yields:

$$\mathbb{P}\left(\bigcup_{1\leq k< ne^{-\gamma d}} \bigcup_{(U_i, U_{1-i}): |U_{1-i}|=k} \left\{ |\hat{S}_{1-i}| \geq \varepsilon |U_{1-i}|/2 \right\} \bigcap \mathcal{D}_n \right) \leq \mathbb{P}\left(\exists U: 1\leq |U|< ne^{-\gamma d}, \ e(U) \geq \frac{\varepsilon \psi_{\lambda}}{4} |U|C \right) = o(1).$$

References

- [1] N. Alon and J. Spencer. *The Probabilistic Method.* Wiley, 2008.
- [2] H. Amini. Bootstrap percolation in living neural networks. Journal of Statistical Physics, 141:459–475, 2010.
- [3] I. Arieli, Y. Babichenko, R. Peretz, and H.P. Young. The speed of innovation diffusion in social networks. *Econometrica*, 88(2):569–594, 2020.
- [4] M. Bahrani, N. Immorlica, D. Mohan, and S.M. Weinberg. Asynchronous majority dynamics in preferential attachment trees. In A. Czumaj, A. Dawar, and Merelli. E., editors, 47th International Colloquium on Automata, Languages, and Programming, ICALP 2020, July 8-11, 2020, Saarbrücken, Germany (Virtual Conference), volume 168 of LIPIcs, pages 8:1-14, 2020.
- [5] J. Balogh and B. Bollobás. Bootstrap percolation on the hypercube. Probability Theory and Related Fields, 134(4):624–648, 2006.
- [6] J. Balogh, B. Bollobás, H. Duminil-Copin, and R. Morris. The sharp threshold for bootstrap percolation in all dimensions. *Trans. Amer. Math. Soc.*, 36:2667 – 2701, 2012.
- J. Balogh and B. G. Pittel. Bootstrap percolation on the random regular graph. Random Structures & Algorithms, 30:257–286, 2007.
- [8] I. Benjamini, S-O. Chan, R. O'Donnell, O. Tamuz, and L-Y. Tan. Convergence, unanimity and disagreement in majority dynamics on unimodular graphs and random graphs. *Stochastic Processes* and their Applications, 126:2719–2733, 2016.
- [9] R. Berkowitz and P. Devlin. Central limit theorem for majority dynamics: Bribing three voters suffices. 16 pp., arXiv:2010:08172v1, 2020.
- [10] B. Bollobás. Random graphs. Cambridge University Press, 2001.
- [11] A. S. Chakrabarti, B. K. Chakrabarti, A. Chatterjee, and M. Mitra. The Kolkata paise restaurant problem and resource utilization. *Physica A: Statistical Mechanics and its Applications*, 388:2420– 2426, 2009.
- [12] J. Chalupa, P. L. Leath, and G. R. Reich. Bootstrap percolation on a Bethe lattice. Journal of Physics C: Solid State Physics, 12:L31–L35, 1979.
- [13] M.A.R. De Cara, O. Pla, and F. Guinea. Competition, efficiency and collective behavior in the "El Farol" bar model. *The European Physical Journal B-Condensed Matter and Complex Systems*, 10:187– 191, 1999.
- [14] P. Erdős and A. Rényi. On random graphs I. Publ. Math. Debrecen, 6:290–297, 1959.
- [15] L. Fontes and R. Schonmann. Bootstrap percolation on homogeneous trees has 2 phase transitions. Journal of Statistical Physics, 132:839–861, 2008.
- [16] N. Fountoulakis, M. Kang, and T. Makai. Resolution of a conjecture on majority dynamics: rapid stabilisation in dense random graphs. *Random Structures and Algorithms*, 57:1134–1156, 2020.
- [17] E.N. Gilbert. Random graphs. Annals of Mathematical Statistics, 30:1141–1144, 1959.

- [18] I. Gilboa and A. Matsui. Social stability and equilibrium. *Econometrica*, 59(3):859–867, 1991.
- [19] E. Goles and J. Olivos. Periodic behaviour of generalized threshold functions. Discrete Mathematics, 30:187–189, 1980.
- [20] J. Hofbauer and K. Sigmund. Evolutionary game dynamics. Bulletin of the American mathematical society, 40:479–519, 2003.
- [21] R.A. Holley and T.M. Liggett. Ergodic theorems for weakly interacting infinite systems and the voter model. The Annals of Probability, 3(4):643 – 663, 1975.
- [22] A. E. Holroyd. Sharp metastability threshold for two-dimensional bootstrap percolation. Probability Theory and Related Fields, 125(2):195–224, 2003.
- [23] S. Janson, T. Luczak, and A. Ruciński. Random Graphs. Wiley & Sons, 2000.
- [24] S. Janson, T. Luczak, T. Turova, and T. Vallier. Bootstrap percolation on the random graph $G_{n,p}$. The Annals of Applied Probability, 22(5):1989–2047, 2012.
- [25] G.E. Kreindler and H.P. Young. Rapid innovation diffusion in social networks. Proceedings of the National Academy of Sciences, 111(3):10881–10888, 2014.
- [26] M. Lelarge. Diffusion and cascading behavior in random networks. Games and Economic Behavior, 75:752–775, 2012.
- [27] T.M. Liggett. Interacting Particle Systems. Springer, Berlin, 1985.
- [28] J. Maynard Smith and G.R. Price. The logic of animal conflict. Nature, 246:15–18, 1973.
- [29] R. Morris. Zero-temperature Glauber dynamics on Z^d. Probability Theory and Related Fields, 149:417– 434, 2009.
- [30] S. Morris. Contagion. The Review of Economic Studies, 67:57–78, 2000.
- [31] E. Mossel, J. Neeman, and O. Tamuz. Majority dynamics and aggregation of information in social networks. Autonomous Agents and Multi-Agent Systems, 28(3):408–429, 2014.
- [32] A. Mukhopadhyay, R. Mazumdar, and R. Roy. Voter and majority dynamics with biased and stubborn agents. *Journal of Statistical Physics*, 181(4):1239–1265, 2020.
- [33] M. Nowak and R. M. May. Evolutionary games and spatial chaos. *Nature*, pages 826–829, 1992.
- [34] D. Oyama and S. Takahashi. Contagion and uninvadability in local interaction games: the bilingual game and general supermodular games. *Journal of Economic Theory*, 157:100–127, 2015.
- [35] V. V. Petrov. Sums of independent random variables, volume 82. Springer Science & Business Media, 2012.
- [36] F.C. Santos and J. M. Pacheco. Scale-free networks provide a unifying framework for the emergence of cooperation. *Phys. Rev. Lett.*, 95, 098104, 2005.
- [37] J. von Neumann and O. Morgenstern. Theory of games and economic behavior. Princeton University Press, 1944.
- [38] M. Yildiz, R. Pagliari, A. Ozdaglar, and A. Scaglione. Voting models in random networks. In Information Theory and Applications Workshop (ITA), 2010, pages 1 – 7, 03 2010.