## UNIVERSITY<sup>OF</sup> BIRMINGHAM University of Birmingham Research at Birmingham

# Abdominal subcutaneous adipose tissue insulin resistance and lipolysis in patients with nonalcoholic steatohepatitis

Armstrong, M. J.; Hazlehurst, J. M.; Hull, D.; Guo, K.; Borrows, S.; Yu, J.; Gough, S. C.; Newsome, P. N.; Tomlinson, J. W.

DOI: 10.1111/dom.12272

License: Creative Commons: Attribution (CC BY)

Document Version Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Armstrong, MJ, Hazlehurst, JM, Hull, D, Guo, K, Borrows, S, Yu, J, Gough, SC, Newsome, PN & Tomlinson, JW 2014, 'Abdominal subcutaneous adipose tissue insulin resistance and lipolysis in patients with non-alcoholic steatohepatitis', *Diabetes, Obesity and Metabolism*, vol. 16, no. 7, pp. 651–660. https://doi.org/10.1111/dom.12272

Link to publication on Research at Birmingham portal

Publisher Rights Statement: Eligibility for repository : checked 07/04/2014

#### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

#### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

## Abdominal subcutaneous adipose tissue insulin resistance and lipolysis in patients with non-alcoholic steatohepatitis

M. J. Armstrong<sup>1</sup>, J. M. Hazlehurst<sup>2</sup>, D. Hull<sup>1</sup>, K. Guo<sup>1</sup>, S. Borrows<sup>3</sup>, J. Yu<sup>4</sup>, S. C. Gough<sup>5</sup>, P. N. Newsome<sup>1,†</sup> & J. W. Tomlinson<sup>2,†</sup>

<sup>1</sup>Centre for Liver Research and NIHR Liver Biomedical Research Unit, University of Birmingham, Birmingham, UK

<sup>2</sup> Centre for Endocrinology, Diabetes and Metabolism, Institute of Biomedical Research, School of Clinical and Experimental Medicine, University of Birmingham, Birmingham, UK

<sup>3</sup>NIHR/Wellcome Trust Clinical Research Facility, Queen Elizabeth Hospital, Birmingham, UK

<sup>4</sup> School of Sport, Exercise & Rehabilitation Sciences, University of Birmingham, Birmingham, UK

<sup>5</sup> Oxford Centre for Diabetes, Endocrinology and Metabolism, and NIHR Oxford Biomedical Research Centre, Oxford, UK

**Background:** Systemic insulin resistance (IR) is a primary feature in non-alcoholic steatohepatitis (NASH), however, there remain limited data on tissue-specific insulin sensitivity *in vivo*.

**Methods:** We examined tissue-specific (adipose, muscle and liver) insulin sensitivity and inflammation in 16 European Caucasian patients with biopsy-confirmed NASH and in 15 healthy controls. All underwent a two-step hyperinsulinaemic euglycaemic clamp incorporating stable isotope measurements of carbohydrate and lipid metabolism with concomitant subcutaneous adipose tissue (SAT) microdialysis.

**Results:** Hepatic and muscle insulin sensitivity were decreased in patients with NASH compared with controls, as demonstrated by reduced suppression of hepatic glucose production and glucose disposal (Gd) rates following insulin infusion. In addition, rates of lipolysis were higher in NASH patients with impaired insulin-mediated suppression of free fatty acid levels. At a tissue specific level, abdominal SAT in patients with NASH was severely insulin resistant, requiring >sixfold more insulin to cause 1/2-maximal suppression of glycerol release when compared with healthy controls. Furthermore, patients with NASH had significantly higher circulating levels of pro-inflammatory adipocytokines than controls.

**Conclusion:** NASH patients have profound IR in the liver, muscle and in particular adipose tissues. This study represents the first *in vivo* description of dysfunctional SAT in patients with NASH.

Keywords: adipose tissue, fatty liver, insulin sensitivity, lipolysis, steatohepatitis

Date submitted 12 December 2013; date of first decision 14 January 2014; date of final acceptance 6 February 2014

### Introduction

Non-alcoholic fatty liver disease (NAFLD) is reaching epidemic proportions, affecting up to 30% of the general population and 70–90% of individuals with type 2 diabetes and/or obesity [1], and is now expected to become the leading indication for liver transplantation by 2020 [2]. Furthermore, NAFLD and in particular the inflammatory (with or without fibrosis) component of non-alcoholic steatohepatitis (NASH), is associated with a significant risk of developing type 2 diabetes, chronic kidney disease, and cardiovascular morbidity and death [1,3]. A better understanding of the key components of the pathogenesis of NASH is therefore needed to provide new therapeutic approaches and thus prevent progressive liver disease and extra-hepatic complications.

E-mail: mattyarm2010@googlemail.com

<sup>†</sup>Joint senior authors.

Systemic insulin resistance (IR) is recognized as one of the main pathogenic factors in NASH [4,5]. Using hyperinsulinaemic euglycaemic clamp techniques (coupled with stable isotopes), several studies have identified the liver (with increased glucose production) and muscle [with decreased glucose disposal (Gd)] as the key sites of increased IR in patients with NASH [5-9]. Recent studies have recognized the importance of adipose tissue, as the principal source of fatty acids ( $\approx 60\%$ ) for the liver, in driving lipid synthesis in both healthy individuals [10] and NASH patients [11]. Adipose tissue is a highly insulin responsive tissue. In an insulin-sensitive state, insulin promotes lipid storage [through fatty acid uptake, re-esterification and de novo lipogenesis (DNL)] and inhibits triglyceride lipolysis, the process whereby triglycerides are hydrolysed to release non-esterified fatty acids (NEFAs) from their glycerol backbone. Studies in patients with NASH have inferred changes in adipose tissue insulin sensitivity through systemic measures of circulating NEFA which are elevated in both the fasting state and under hyperinsulinaemic conditions [6-8,12]. Importantly, this appears to be independent of the degree of obesity [13].

Adipose tissue dysfunction is considered to be a major contributory factor of NASH, by means of the resultant

*Correspondence to:* Dr Matthew J. Armstrong, Wellcome Trust Research Fellow & Hepatology Registrar, NIHR Centre for Liver Research, University of Birmingham, 5th Floor IBR, Birmingham B15 2TH, UK.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

'lipotoxicity' inducing both hepatic IR and skeletal muscle IR [14]. Studies that have been published to date have, however, solely focused on quantifying whole-body lipolysis using either circulating NEFA [i.e. quantified by adipose insulin resistance (ADIPO-IR) index = fasting NEFA  $\times$  insulin] [8,9,12] or the rate of systemic appearance of labelled glycerol/palmitate isotopes [6,13,15]. In particular, no studies have assessed the response of local adipose tissue to the action of insulin, which provides greater insights into the functional relevance of adipose tissue. A greater understanding of which adipose depots are dysfunctional in NASH patients would greatly enhance our knowledge in developing new targeted therapies. Traditionally, visceral adipose tissue (VAT) has been recognized as a major contributor to IR and metabolic conditions including NASH, due to its close proximity to the portal vein and abundance of pro-inflammatory mediators [16,17]. However, as VAT only contributes to 15-20% of circulating NEFA pool [18,19], researchers have questioned whether overspill from abdominal subcutaneous adipose tissue (SAT) plays a more significant role. Indeed, while several studies have linked abdominal SAT with indices of IR in subjects with and without metabolic syndrome [20-23], none have examined it in relation to NASH.

Adopting an integrative physiological approach with functional measures of lipid and carbohydrate flux, we have performed a clinical study to determine the relative contribution of tissue-specific insulin sensitivity, notably in SAT, in patients with biopsy-proven NASH in comparison with a healthy control cohort.

## **Research Design and Materials**

The clinical protocols received full ethical approval from Leicestershire, Northamptonshire and Rutland (ref. 10/H0402/32) and South Birmingham (ref. 10/H1207/15) Local Research Ethics Committees. All adult subjects gave informed written consent prior to participation.

### **Study Subjects**

NASH Patients. Sixteen patients with a definitive diagnosis of NASH on liver biopsy within 6 months of the study were recruited. The histological diagnosis was made using well-established criteria [24] by two independent liver histopathologists. The subjects were of adult age (18-70 years) and had a body mass index  $(BMI) \ge 25 \text{ kg/m}^2$ . Patients with co-existing type 2 diabetes were diet-controlled or were on a stable dose of metformin  $\pm$  gliclazide for a minimum of 3 months prior to the study and had a glycated haemoglobin 1c (HbA1c) < 9.0%. Participants were excluded if they had a history of excess alcohol consumption (females >14 units/week and males >21 units/week), liver disease of other aetiology, decompensated cirrhosis (Child's Pugh B or C), recent or concomitant drug use of inducers of hepatic steatosis/weightinducing therapy, and significant active medical illnesses (with the exception of type 2 diabetes). All patients with no previous diagnosis of type 2 diabetes underwent a 75 g oral glucose tolerance test (OGTT).

Healthy Volunteers. Fifteen healthy volunteers (9 males:6 females; mean age  $33 \pm 2$  years) were recruited by use of a local advertisement. All controls were asymptomatic, non-diabetic, were taking no regular medication and had no significant medical history of note. Female controls had pregnancy excluded and were not taking any form of hormonal contraception. In the healthy control cohort, all consumed alcohol within recommended limits, had normal liver function tests (LFTs) and had normal levels of noninvasive markers of hepatic injury (serum cytokeratin-18) and fibrosis [serum enhanced liver test, enhanced liver fibrosis (ELF)]. Furthermore, all controls had a negative NAFLD liver fat score (<-0.640) and estimated liver fat <3.0% based on the Kotronen et al. equations, which were originally validated with <sup>1</sup>H-magnetic resonance spectroscopy (MRS) [25]; 5 of 15 subjects underwent a hepatic MRS, as part of a separate study, and in keeping with the Kotronen equations, had hepatic steatosis excluded (<2.5%).

### **Study Design**

All participants underwent a two-step hyperinsulinaemic euglycaemic clamp incorporating stable isotopes with concomitant SAT microdialysis at the NIHR/Wellcome Trust Clinical Research Facility (WTCRF, Birmingham, UK) (Figure S1, Supporting information).

*Hepatic DNL*. At 17:00 hours, participants were admitted to the WTCRF and total body water was estimated by bioimpedance (Tanita BC418MA, Amsterdam, the Netherlands). A standardized meal (carbohydrate 45 g, protein 23 g and fat 20 g) was provided at 17:00 hours, after which participants remained fasted until the end of the clamp at 14:00 hours the next day. To determine rates of DNL, participants were given oral deuterated water,  ${}^{2}H_{2}O$  (3 g/kg total body water in two divided doses), at 18:00 and 22:00 hours followed by *ad libitum* drinking water enriched with 0.4%  ${}^{2}H_{2}O$ .

Two-Step Hyperinsulinaemic Euglycaemic Clamp. At 08:00 hours the next morning fasting blood samples were taken by standard needle venepuncture prior to starting the twostep hyperinsulinaemic euglycaemic clamp. Arterialized blood was sampled to determine the blood glucose concentration at which to maintain ('clamp') the participant throughout the study using an YSI 2700 machine (YSI life sciences, Fleet, Hampshire, UK). An intravenous bolus of U-[<sup>13</sup>C]glucose (2 mg/kg body weight; CK gas limited, Hook, UK) was administered over 1 min followed by a constant infusion rate (0.02 mg/kg/min) for 6 h until the end of the clamp. Steady state blood samples were taken at three time points during the final 30 min of the 2-h basal phase. At 10:00 hours, lowdose insulin (Actrapid; Novo Nordisk, Copenhagen, Denmark) was infused at 20 mU/m<sup>2</sup>/min. At 10:04 hours a concomitant infusion of 20% glucose enriched with U-[<sup>13</sup>C]-glucose to 4% was commenced. Arterialized blood samples were taken at 5 min intervals and the 20% glucose infusion rate was changed to maintain fasting glycaemic levels. Steady state blood samples were taken at three time points in the final 30 min of the 2-h low-dose insulin infusion. The insulin infusion rate was then increased to  $100 \text{ mU/m}^2/\text{min}$  (high-dose) for 2 h with sampling as described above. Rates of hepatic endogenous glucose production (EGP) and Gd were calculated by using modified versions of the Steele Equations [26,27].

Adipose Microdialysis. A microdialysis catheter (CMA microdialysis AB, Solna, Sweden) was inserted after local anaesthetic (5 ml 1% lignocaine) was injected into the abdominal SAT (minimum depth 1 cm), 10 cm lateral to the umbilicus, prior to commencing the clamp. Thereafter, micro-dialysate samples were collected into micro-vials ( $0.3 \mu$ l/min) every 30 min until the end of the clamp.

### **Data Collection and Analysis**

*Clinical and Biochemical Parameters.* Participant demographics and clinical/biochemical measures were recorded at the study visit. These included 75 g OGTT, anthropometry (including bioimpedance), fasting haematological/biochemical bloods and non-invasive serum markers of liver injury [cytokeratin-18 (CK-18), ELF test, Kotronen score)]. Serum insulin, NEFA and adipocytokines [adiponectin, leptin, resistin, tumour necrosis factor alpha (TNF- $\alpha$ ), high sensitivity C-reactive protein (hs-CRP), interleukin-6 (IL-6), IL-17, chemokine ligand-2 (CCL-2), CCL-3, CCL-4 and CCL-5] were measured using commercially available kits. Detailed descriptions of the above are available in Appendix S1.

*Abdominal SAT Microdialysis.* Microdialysate samples were analysed using a mobile photometric, enzyme-kinetic analyser (CMA Iscus Flex) for glycerol concentration. The rate of interstitial glycerol release represented the magnitude of SAT lipolysis in the fasted state and in response to insulin.

Stable Isotope Mass Spectrometry Analysis. The enrichment of U-[<sup>13</sup>C]-glucose in plasma was determined by gas chromatography-mass spectrometry (model 5973; Agilent technologies, Cheshire, UK). Deuterium enrichment of the body water pool was measured using the Gasbench II (www.thermo.com), an automated  $H_2/H_2O$  equilibration device, coupled on-line to a ThermoFinnigan Deltaplus XP Isotope Ratio Mass Spectrometer (IRMS; ThermoFinnigan MAT GmbH, Bremen, Germany). The full methods have been previously described in detail [28]. Deuterium enrichment in the palmitate fraction of total plasma triglycerides was measured on an automated GC/TC/IRMS system (ThermoFinnigan Delta Pus XP; www.thermo.com).

Contribution of Hepatic DNL to Total Palmitate Synthesis. The percentage contribution of hepatic DNL to endogenous palmitate synthesis was determined by the incorporation of  ${}^{2}$ H<sub>2</sub>O in the palmitate present in the plasma total triglyceride pool, as previously described [28]. This percentage was calculated from the increase in the  ${}^{2}$ H/ ${}^{1}$ H ratio in the palmitate methylester of the total triglyceride fraction and in the water of plasma samples taken before (17:00 hours, at admission) and 14 h after the initial ingestion of the  ${}^{2}$ H<sub>2</sub>O tracer (08:00 hours, before the start of the hyperinsulinaemic euglycaemic clamp). The following formula was used: % hepatic DNL contributes to endogenous palmitate synthesis = (delta

## original article

 $^{2}$ H/<sup>1</sup>H ratio in palmitate methylester)/(delta  $^{2}$ H/<sup>1</sup>H ratio in waterpool) × (34/22) × 100%. In the equation, 34 is the total number of H-atoms in palmitate methylester and 22 is the number of water molecules incorporated into palmitate via DNL as observed in previous rodent studies [29] and currently used in human studies [30].

### **Statistical Analysis**

Descriptive statistics was applied to characterize the NASH and healthy volunteer cohorts. Continuous clinical and laboratory variables are reported as means and standard error (s.e.) as all variables had parametric distribution on D'Agostino and Pearson Omnibus Normality testing. Categorical variables are reported as number and percentages. Area under the curve (AUC) analysis was performed using the trapezoidal method for interstitial glycerol release during the clamp. For comparison of single variables, unpaired Student t-tests were used (or non-parametric equivalents where data were not normally distributed). Where repeated samples were taken repeatedmeasures one-way analysis of variance (ANOVA) was used, incorporating the Dunnett's test for multiple comparisons. The significance level was set at p < 0.05. All analyses were performed using the GRAPHPAD PRISM 5.0 software package.

## **Results**

### **Participant Characteristics**

Participant demographics and clinical characteristics are summarized in Table 1. NASH subjects were significantly older ( $54.4 \pm 2.1$  vs.  $33.1 \pm 2.2$  years; p < 0.0001) and had a higher BMI ( $34.3 \pm 1.0$  vs.  $26.7 \pm 1.0$  kg/m<sup>2</sup>; p < 0.0001) and abdominal fat mass on bioimpedance ( $20.3 \pm 1.5$  vs.  $12.0 \pm 1.5$  kg; p = 0.0011). Of the 16 subjects with NASH, 5 had mild-moderate fibrosis (Kleiner F1-F2) and 9 had advanced fibrosis (F3-F4). NASH subjects had significantly higher serum levels of liver enzymes [alanine transaminase (ALT)  $68.7 \pm 11$  vs.  $18.9 \pm 2.6$  IU/l; p = 0.0001], serum CK-18 M30 levels ( $544 \pm 116$  vs.  $161 \pm 9.8$  IU/l; p = 0.0034) and ELF test ( $9.20 \pm 0.3$  vs.  $7.34 \pm 0.1$ ; p < 0.001); values for all these parameters were within accepted reference ranges in the healthy volunteers.

### Systemic IR

Fasting serum glucose, insulin and homeostatic model assessment insulin resistance (HOMA-IR,  $4.40 \pm 0.8$  vs.  $1.19 \pm 0.2$ ) were significantly higher in patients with NASH (all p < 0.001) (Figure 1A, B). During the two-step hyperinsulinaemic clamp, NASH subjects had significantly lower weight-adjusted glucose infusion rates in response to low-dose ( $1.47 \pm 0.08$  vs.  $3.08 \pm 0.4$  mg/kg/min; p = 0.0008) and high-dose insulin ( $5.80 \pm 0.4$  vs.  $9.14 \pm 0.5$  mg/kg/min; p < 0.0001). In keeping with peripheral (largely muscle) IR, weight-adjusted Gd rates were significantly lower in NASH subjects at low-dose ( $0.85 \pm 0.1$  vs.  $1.76 \pm 0.4$  mg/kg/min; p < 0.05) and high-dose insulin infusions ( $4.55 \pm 0.6$  vs.  $6.10 \pm 0.5$  mg/kg/min; p = 0.05) (Figure 1C).

#### DIABETES, OBESITY AND METABOLISM

**Table 1.** Demographics and clinical parameters of 16 patients with NASH and 15 healthy controls. Values are mean (s.e.), unless stated. All blood parameters were fasting samples. Comparisons of continuous variables were made with unpaired Student's *t*-test, and categorical variables with fisher exact/chi-squared test.

	NASH $(n = 16)$	Controls $(n = 15)$	p-Value
Demographics			
Male sex, n (%)	11 (68.8)	9 (60.0)	0.716
Age (years)	54.4 (2.1)	33.1 (2.2)	< 0.0001
Ethnicity, n (%)			
Caucasian	16 (100)	14 (93.3)	0.484
Asian	0 (0)	1 (6.7)	
Metabolic parameters			
Type 2 diabetes, n (%)	7 (43.8)	0 (0)	0.001
Impaired glucose tolerance, n (%)	3 (18.8)	0 (0)	
Normal glucose tolerance, n (%)	6 (37.5)	15 (100)	
Fasting glucose (mmol/l)	5.34 (0.24)	4.37 (0.067)	0.0008
Fasting insulin (pmol/l)	125.8 (20.8)	43.3 (7.41)	0.0003
HbA1c (%)	5.99 (0.21)	—	—
Pre-study OAD treatment, n (%)	8 (50.0)	0 (0)	0.0024
Pre-study statin treatment, n (%)	7 (43.8)	0 (0)	0.0068
Pre-study anti-hypertensive treatment, n (%)	6 (37.5)	0 (0)	0.0177
BMI (kg/m <sup>2</sup> )	34.3 (1.04)	26.7 (0.95)	< 0.0001
Weight (kg)	100.3 (3.83)	78.5 (3.67)	0.0003
Total fat mass (kg)	35.8 (2.64)	20.2 (1.79)	< 0.0001
Truncal fat mass (kg)	20.3 (1.45)	12.0 (1.51)	0.0011
Systolic BP (mmHg)	129.4 (3.43)	129.3 (2.74)	0.982
Waist circumference (cm)	114.1 (2.87)	85.9 (3.07)	< 0.0001
Total cholesterol (mmol/l)	4.51 (0.20)	4.59 (0.30)	0.891
HDL (mmol/l)	1.11 (0.064)	1.26 (0.11)	0.256
LDL (mmol/l)	3.01 (0.21)	2.73 (0.43)	0.550
Triglycerides (mmol/l)	1.95 (0.26)	1.62 (0.34)	0.438
TSH (µU/l)	2.76 (0.38)	2.01 (0.31)	0.165
Creatinine (µmol/l)	71.3 (3.46)	72.6 (3.47)	0.800
Liver parameters			
AST (IU/l)	55.1 (5.66)	20.4 (1.55)	< 0.0001
ALT (IU/l)	68.7 (10.6)	18.9 (2.57)	0.0001
Alk Phos (IU/l)	155.7 (25.3)	128 (9.64)	0.327
Bilirubin (µmol/l)	13.4 (1.76)	12.0 (1.02)	0.494
Albumin (g/l)	47.1 (0.70)	41.6 (0.77)	< 0.001
Platelets ( $\times 10^9 l^{-1}$ )	203.3 (14.7)	216.1 (10.5)	0.489
CK-18 M30 (IU/l)	543.4 (115.8)	160.9 (9.83)	0.0034
ELF test	9.20 (0.30)	7.34 (0.12)	< 0.0001
Histology parameters (NASH only)			
Median Kleiner fibrosis score*(IQR)	3.0 (1.0-3.75)	—	—
Median NAFLD activity score†(IQR)	4.5 (4.0-5.0)	—	—

Bold are the p-values that are significant at <0.05. ALT, alanine transaminase; AST, aspartate transaminase; Alk phos, alkaline phosphatase; BMI, body mass index; BP, blood pressure; CK-18, cytokeratin-18; ELF, enhanced liver fibrosis; HbA1c, glycated haemoglobin 1c; HDL, high-density lipoprotein; IQR, interquartile range; LDL, low-density lipoprotein; NAFLD, Non-alcoholic fatty liver disease; NASH, non-alcoholic steatohepatitis; OAD, oral anti-diabetic drug; TSH, thyroid stimulating hormone.

\*The Kleiner fibrosis score ranges from 0 to 4, whereby 0 = no fibrosis and 4 = cirrhosis.

 $\dagger$ The NAFLD activity score (scored out of 8) is a sum of steatosis (0–3), hepatocyte ballooning (0–2) and lobular inflammation (0–3).

#### **Hepatic IR**

Although fasting EGP rates were similar in patients with NASH and healthy controls  $(2.14 \pm 0.1 \text{ vs. } 2.15 \pm 0.1 \text{ mg/kg/min};$ p > 0.9; Figure 1D), this was in the context of fasting hyperinsulinaemia (Figure 1B, Table 1); changes were consistent with hepatic IR. The hepatic IR index (= EGP × fasting insulin [31]) was significantly higher in NASH patients (278 ± 52.7 vs. 90.0 ± 14.9 mg/kg/min pmol/ml; p = 0.0024). In addition, low-dose insulin-mediated suppression of EGP was decreased in patients with NASH (Figure 1D), consistent with hepatic IR (% EGP suppression: 41.0 ± 4.3 vs. 70.2 ± 9.5%; p = 0.008). These differences persisted even after removing patients with type 2 diabetes (n = 6) from the NASH cohort (42.2 ± 5.6 vs. 70.2 ± 9.5%; p < 0.05).

#### **Hepatic DNL**

The percentage contribution of DNL to total endogenous palmitate synthesis was variable across all individuals and



**Figure 1.** Subjects with non-alcoholic steatohepatitis (NASH) have significant systemic, muscle and hepatic insulin resistance (IR). Circulating glucose (A) and insulin (B) concentrations during the two-step hyperinsulinaemic euglycaemic clamp. The degree muscle and hepatic insulin sensitivity was determined by glucose disposal (C) and suppression of hepatic glucose production (D), respectively. White bar = controls, black bar = NASH. \*p < 0.05, \*\*p < 0.01, \*\*\*p < 0.001 versus controls.

although higher in NASH subjects compared with controls [median 4.90 (IQR 3.9–5.6) vs. 2.79 (1.2–6.4); p = 0.16] this did not reach significance.

#### Depot-specific adipose tissue IR

Circulating NEFA levels were not different between patients with NASH and healthy controls ( $563 \pm 33$  vs.  $465 \pm 32 \mu mol/l$ ; p = 0.13; Figure 2A). However, taking into account fasting hyperinsulinaemia in patients with NASH, the calculated adipose IR index (fasting NEFA × fasting insulin [31]) was significantly elevated ( $64.4 \pm 9.1$  vs.  $20.5 \pm 3.9$  mmol/l pmol/l; p = 0.0002) in patients with NASH. Insulin infusion significantly suppressed circulating NEFAs in both NASH and control subjects (p < 0.0001 vs. basal NEFA in each group; Figure 2A). In order to determine insulin sensitivity, using regression analysis, the insulin concentrations causing half-maximal suppression of serum NEFA (INS-1/2-max NEFA) were calculated for each subject (Figure 2B). INS-1/2-max NEFA was greater than threefold higher in NASH subjects compared with the controls  $(227 \pm 35 \text{ vs. } 65.2 \pm 14 \text{ pmol/l}; \text{ } \text{p} = 0.0003)$  consistent with adipose tissue IR. The significant difference in INS-1/2max NEFA remained  $(195 \pm 30 \text{ vs. } 65.2 \text{ pmol/l}; \text{ } \text{p} = 0.0002)$ after removing patients with type 2 diabetes (n = 7) from the analysis.

Interstitial glycerol release assessed using microdialysis was used as a direct measure of abdominal SAT function (Figure 3A, B). In the fasting state, the rate of interstitial glycerol release was not different in NASH subjects compared with controls  $(383 \pm 44 \text{ vs. } 286 \pm 40 \,\mu\text{mol/l·h}; p = 0.12)$ . In healthy controls, low-dose insulin infusion (20 mU/m<sup>2</sup>/min) significantly suppressed the rate of interstitial glycerol release (Basal:  $286 \pm 40$  vs. low-dose insulin:  $143 \pm 18 \mu \text{mol/l}\cdot\text{h}$ ; p < 0.001), whereas it did not suppress release in NASH subjects (Basal:  $383 \pm 44$  vs. low-dose insulin:  $379 \pm 43 \,\mu \text{mol/l}\cdot\text{h}$ ; p > 0.05). High dose insulin (100 mU/m<sup>2</sup>/min) suppressed glycerol release in both patients with NASH and in controls, however, the rate of glycerol release remained significantly higher in the NASH subjects compared with controls  $(261 \pm 31)$ vs.  $65.8 \pm 14 \mu \text{mol/l} \cdot \text{h}$ ; p < 0.0001; Figure 3B). Furthermore, the INS-1/2-max glycerol was sixfold higher in the NASH subjects compared with controls (p < 0.0001; Figure 4, summary box). All of the above comparisons remained significant after excluding subjects with type 2 diabetes (n = 7)from the NASH cohort (Figure S2).

#### Serum Adipocytokines and Inflammatory Cytokines

Subjects with NASH had significantly higher fasting circulating levels of TNF- $\alpha$  (p < 0.0001), hs-CRP (p < 0.05), IL-6 (p < 0.05) and CCL-2 (p < 0.05) than controls (Figure 5).



**Figure 2.** Subjects with non-alcoholic steatohepatitis (NASH) have significant global adipose tissue insulin resistance (IR). (A) Circulating non-esterified fatty acid (NEFA) concentrations at basal and hyperinsulinaemic phases of euglycaemic clamp. (B) As a marker of global adipose tissue insulin resistance, the concentration of circulating insulin concentrations (pmol/l) causing 1/2-maximal suppression of circulating NEFA was calculated. White bar = controls, black bar = NASH. \*p < 0.05, \*\*\* p < 0.001 versus controls. ++++ p < 0.001 versus basal phase. NS, non-significant.



**Figure 3.** Non-alcoholic steatohepatitis (NASH) is associated with significant abdominal SAT IR. (A) SAT interstitial fluid concentrations of glycerol during the 2-step hyperinsulinaemic euglycaemic clamp. (B) To determine the rate of lipolysis in subcutaneous adipose tissue (SAT) under basal and hyperinsulinaemic conditions area under the curve (AUC) analysis was performed using the trapezoidal method for interstitial glycerol release. Broken line/white bar = controls, solid line/black bar = NASH. \*\*\*\* p < 0.0001 versus controls; +++ p < 0.001, ++++ p < 0.0001 versus basal phase. NS, non-significant.

Serum adiponectin levels (p = 0.001) were significantly lower in NASH subjects, with a non-significant trend towards higher circulating leptin compared with controls (p = 0.059). The resultant leptin : adiponectin ratio was 2.5-fold higher in NASH subjects than controls ( $3.22 \pm 0.5$  vs.  $1.27 \pm 0.4$ ; p = 0.0032). There were no significant differences in IL-17, resistin and chemotactic cytokines CCL-3, CCL-4 and CCL-5 (RANTES). With the exception of CCL-2 (p = 0.09), differences in TNF- $\alpha$  (p < 0.0001), hs-CRP (p < 0.05), IL-6 (p < 0.05) and adiponectin remained significant after excluding subjects with type 2 diabetes (n = 7) from the NASH cohort (Figure S3).

### Discussion

The data from this study have begun to address the tissuespecific contributions make to global IR seen in patients with NASH. Using novel techniques that have functional readouts of insulin-regulated processes in a tissue-specific manner allows an assessment of the contribution of the liver (EGP and DNL), skeletal muscle (Gd), and adipose tissue (circulating NEFA and adipose microdialysis) to systemic IR. By doing so, we have not only demonstrated significant IR at the level of the liver, muscle and adipose tissue, but also by measuring depot-specific glycerol release, our study represents the first *in vivo* description of dysfunctional abdominal SAT in patients with NASH.



**Figure 4.** Subjects with non-alcoholic steatohepatitis (NASH) have a disproportionate higher degree of insulin resistance (IR) in subcutaneous adipose tissue (SAT) (sixfold vs. controls) compared with whole-body adipose tissue (threefold vs. controls). Line graph representing the concentrations of circulating non-esterified fatty acid (NEFA) (whole-body lipolysis) and interstitial fluid glycerol (SAT-specific lipolysis) in basal, low-dose and high-dose insulin phases of the euglycaemic clamp. Black lines = NASH (mean  $\pm$  s.e.), Grey line = control. Sold line = glycerol levels, broken line = NEFA levels.

We observed significant levels of hepatic and muscle IR in NASH subjects, as represented by impaired insulin-mediated suppression of hepatic glucose production and stimulated muscle Gd (weight-adjusted), respectively. In keeping with previous studies [6,13,32], the level of hepatic and muscle IR remained significant when patients with type 2 diabetes were removed from the analysis. Notably, we only saw a non-significant trend towards higher levels of fasting DNL in NASH subjects compared with healthy controls (4.9 vs. 2.8%; p = 0.16). Even though the low levels of fasting DNL in healthy subjects were consistent with the literature (i.e. <5.0%) [33], our findings in NASH subjects were considerably lower (4.9 vs. 15–24%) than previously reported [11,34]. This might be attributed to sampling DNL in the fasting state only, oral administration of deuterated water (vs. intravenous deuterated tripalmitate [11]) and/or the shorter duration of stable isotope labelling compared with previous reports (14 vs. >96 h [11]). Due to the nature of the stable isotopes incorporated as part of the clamp and the high rates of labelled glucose infusions required to maintain fasting glycaemia, we were unable to assess the rates of DNL associated with hyperinsulinaemia. It is important to note, however, that Donnelly et al. previously

reported that the majority of lipid accumulation in NASH was attributed to adipose-derived NEFA (59%), rather than DNL (26%) [11].

We demonstrated severe adipose tissue dysfunction in patients with NASH using a variety of assessments including adipose IR index, INS-1/2-max NEFA, adipose tissue microdialysis and circulating adipocytokines. The discrepancy between high fasting leptin and low circulating levels of adiponectin provided further evidence of abnormal adipose tissue function. Indeed, a growing body of evidence indicates that the primary defect in NASH subjects occurs in adipose tissue [14], from which triglyceride-derived toxic metabolites including the NEFA pool, impair insulin signalling in both skeletal muscle and liver tissue ('lipotoxicity'). A vicious cycle of hepatic, muscle and adipose tissue dysfunction ensues, leading to development of a pathogenic circulating milieu of high levels of insulin, glucose, NEFA and pro-inflammatory cytokines (e.g. hsCRP, IL-6, TNF- $\alpha$ , and CCL-2), all of which were observed in our patients with NASH.

Traditionally, VAT has been recognized as the major contributor to hepatic IR and lipotoxicity [35], due to its close proximity to the portal vein and concentration



**Figure 5.** Subjects with non-alcoholic steatohepatitis (NASH) have significantly lower levels of fasting adiponectin (A) and higher levels of fasting pro-inflammatory adipocytkines [(B) leptin, (C) high sensitivity C-reactive protein (hs-CRP), (D) tumour necrosis factor alpha (TNF- $\alpha$ ), (E) interleukin-6 (IL-6) and (F) chemokine ligand-2 (CCL-2)/MCP-1]. \*p < 0.05, \*\*p < 0.01, \*\*\*p < 0.001, \*\*\*\*p < 0.001 versus controls.

of inflammatory mediators [16,17]. However, as VAT only contributed to 15-20% of circulating NEFA pool [18,19], researchers proposed that either VAT exerted its effects via other non-NEFA factors including adipocytokines [35] or that abdominal SAT plays an important role in lipotoxicity [36]. Several studies have linked abdominal SAT with IR using euglycaemic clamps in subjects with and without metabolic syndrome [20-23], but our data is one of the first to report depot-specific dysfunction in biopsy-proven NASH subjects. Previous studies in NASH patients have solely relied on circulating NEFA to provide estimates of adipose IR [8,9,12,13,32], which are more reflective of wholebody lipolysis, rather that depot-specific [37]. By directly measuring interstitial fluid concentrations of glycerol, we report novel insights into the degree of abdominal SAT IR and lipolysis in patients with NASH. The greater magnitude of resistance to the anti-lipolytic effect of insulin in SAT (sixfold vs. controls) in comparison to whole-body adipose (threefold vs. controls) in our study may well reflect depotspecific IR, in which abdominal SAT is the major source of lipotoxicity in NASH. Interestingly, using paired adipose and liver biopsies from patients undergoing bariatric surgery Tordjman et al. have recently shown that deep SAT (and not superficial SAT) has an inflammatory profile (i.e. IL-6 gene, macrophage accumulation) similar to VAT in NASH subjects [38].

One hypothesis is that abdominal SAT acts as 'buffer' for excess calorific intake and triglyceride deposition. When SAT fails to match the demand, as might be the case in NASH subjects, adipose hypertrophy, inflammation (via macrophage recruitment via CCL-2) and local IR sequentially develop. The resultant localized excess NEFA, as reported here, can result in an overspill of triglyceride-derived toxic metabolites into VAT and subsequently the liver [39].

The role of ethnicity in adipose IR and NASH has recently been investigated, in which Lomonaco et al. demonstrated no difference in levels of IR (EGP, fasting NEFA) between Hispanic and Caucasian cohorts with NASH, well-matched for adiposity [32]. With the exception of two Italian studies [7,12], very little data exists in well-characterized patients with NASH of western European descent. Adipose IR index in our UK cohort (9.3 mmol/1 $\mu$ U/l) was, however, similar to that previously reported in NASH patients from southern Europe and the USA (8.0–11.9 mmol/1 $\mu$ U/l), all of which were 3–6.6 times higher than their respective healthy controls [8,12,13].

Our study does have limitations, in particular, the metabolic phenotype mismatch between the NASH and 'healthy' controls. This remains a critical challenge in real-world research, due to the high prevalence of obesity and metabolic syndrome in patients with NASH at the time of first presentation. Even though differences in adipose IR remained significant after exclusion of patients with diabetes (Figures S2, S3), we were unable to extrapolate whether our findings were independent of age and measures of adiposity. Gastaldelli et al. have reported that whole-body lipolysis (using adipose-IR index) in NASH is independent of obesity status (defined by BMI) [13], but this requires validation with robust measures of VAT and abdominal SAT mass/volume. In this study, we were unable to directly compare SAT and VAT, as realtime assessment of VAT function is not feasible in human studies. The demonstration that SAT is the dominant source of products of triglyceride hydrolysis in healthy humans [19] and our finding of marked SAT IR may have significant clinico-pathological implications in patients with NASH. This suggests that it is not simply VAT accumulation that is important in driving the pathological process. Lastly, the cross-sectional design of our study did not allow a causal relationship to be determined between dysfunctional SAT and progressive liver disease. Previous longitudinal studies (14 weeks) have shown that diet-induced decreases in MRImeasured VAT correlate with improvements in hepatic lipid content and markers of IR. However, no longitudinal studies have investigated functional changes in VAT or SAT with disease progression or regression after therapeutic intervention [40].

In summary, our study highlights that patients with NASH have marked adipose tissue dysfunction, alongside increased hepatic and muscle IR. In particular, we have drawn attention to the profound levels of IR and lipolysis in abdominal SAT, which appears disproportionate to whole-body adipose. Dysfunctional abdominal SAT likely plays a key role in NASH lipotoxicity, rather than being just a bystander to VAT. Whether this is indeed a tissue-mass effect remains to be investigated. Future prospective studies that are sufficiently powered to enable adjustment of metabolic confounders are now required to investigate the relative contribution of SAT (vs. VAT) in disease progression and the impact of novel interventions in NASH.

### Acknowledgements

This work has been supported by the Wellcome Trust (clinical research training fellowship ref. 089780/Z/09/Z, M. J. A.), Medical Research Council (senior clinical fellowship ref. G0802765, J. W. T.) and the National Institute of Health Research Liver Biomedical Research Unit. Equipment has been provided through Advantage West Midlands Science City. The clinical study was carried out at the NIHR/Wellcome Trust Birmingham Clinical Research Facility. The views expressed are those of the authors and not those of the NHS, NIHR or the funders.

## **Conflict of Interest**

The authors declare no conflicts of interest.

M. J. A., S. G., P. N. N. and J. W. T. contributed to the design of the study protocol. M. J. A., J. M. H., D. H., K. G., S. B. and J. W. T. performed the clamp experiments and generated all the clinical data for the manuscript. M. J. A., J.

M. H. and J. Y. analysed the clinical samples. M. J. A. and J. W. T. performed the statistical analysis. M. J. A. wrote the first draft of the manuscript. All authors contributed to revision of the final manuscript. M. J. A. and J. W. T. are guarantors of the data presented.

### **Supporting Information**

Additional Supporting Information may be found in the online version of this article:

**Figure S1.** Schematic of the study design. All participants underwent a two-step hyperinsulinaemic euglycaemic clamp with stable isotope tracers (13C-glucose, deuterated water) and adipose microdialysis to determine tissue-specific insulin resistance.

**Figure S2.** Subjects with non-alcoholic steatohepatitis (NASH) with and without type 2 diabetes had significant abdominal subcutaneous adipose tissue (SAT) insulin resistance (IR) compared with controls. (A) SAT interstitial fluid concentrations of glycerol during the two-step hyperinsulinaemic euglycaemic clamp. (B) To determine the rate of lipolysis in SAT under basal and hyperinsulinaemic conditions area under the curve (AUC) analysis was performed using the trapezoidal method for interstitial glycerol release. Broken line/white bar = controls, solid grey line/grey bar = NASH without type 2 diabetes, sold black line/black bar = NASH with type 2 diabetes. \*\*\*\* p < 0.0001 versus controls; +p < 0.05, +++p < 0.001, ++++p < 0.0001 versus basal phase. NS, non-significant.

**Figure S3.** Non-diabetic subjects with non-alcoholic steatohepatitis (NASH) have significantly lower levels of fasting adiponectin (A) and higher levels of fasting proinflammatory adipocytkines [(C) high sensitivity C-reactive protein (hs-CRP), (D) tumour necrosis factor alpha (TNF- $\alpha$ ) and (E) interleukin-6 (IL-6)]. Higher levels of (B) leptin and (F) chemokine ligand-2 (CCL-2)/monocyte chemoattractant protein-1 (MCP-1) were seen in non-diabetic subjects with NASH, albeit not achieving significance. \*p < 0.05, \*\*\*\*\* p < 0.0001 versus controls.

Appendix S1. Supplementary methods.

### References

- Anstee QM, Targher G, Day CP. Progression of NAFLD to diabetes mellitus, cardiovascular disease or cirrhosis. Nat Rev Gastroenterol Hepatol 2013; 10: 330–344.
- Charlton MR, Burns JM, Pedersen RA, Watt KD, Heimbach JK, Dierkhising RA. Frequency and outcomes of liver transplantation for nonalcoholic steatohepatitis in the United States. Gastroenterology 2011; 141: 1249–1253.
- Bonora E, Targher G. Increased risk of cardiovascular disease and chronic kidney disease in NAFLD. Nat Rev Gastroenterol Hepatol 2012; 9: 372–381.
- Chitturi S, Abeygunasekera S, Farrell GC et al. NASH and insulin resistance: insulin hypersecretion and specific association with the insulin resistance syndrome. Hepatology 2002; 35: 373–379.
- 5. Marchesini G, Brizi M, Bianchi G et al. Nonalcoholic fatty liver disease: a feature of the metabolic syndrome. Diabetes 2001; **50**: 1844–1850.

- Sanyal AJ, Campbell-Sargent C, Mirshahi F et al. Nonalcoholic steatohepatitis: association of insulin resistance and mitochondrial abnormalities. Gastroenterology 2001; **120**: 1183–1192.
- 7. Bugianesi E, Gastaldelli A, Vanni E et al. Insulin resistance in non-diabetic patients with non-alcoholic fatty liver disease: sites and mechanisms. Diabetologia 2005; **48**: 634–642.
- Lomonaco R, Ortiz-Lopez C, Orsak B et al. Effect of adipose tissue insulin resistance on metabolic parameters and liver histology in obese patients with nonalcoholic fatty liver disease. Hepatology 2012; 55: 1389–1397.
- Ortiz-Lopez C, Lomonaco R, Orsak B et al. Prevalence of prediabetes and diabetes and metabolic profile of patients with nonalcoholic fatty liver disease (NAFLD). Diabetes Care 2012; 35: 873–878.
- Barrows BR, Parks EJ. Contributions of different fatty acid sources to very low-density lipoprotein-triacylglycerol in the fasted and fed states. J Clin Endocrinol Metab 2006; 91: 1446–1452.
- 11. Donnelly KL, Smith CI, Schwarzenberg SJ, Jessurun J, Boldt MD, Parks EJ. Sources of fatty acids stored in liver and secreted via lipoproteins in patients with nonalcoholic fatty liver disease. J Clin Invest 2005; **115**: 1343–1351.
- Musso G, Cassader M, De Michieli F, Rosina F, Orlandi F, Gambino R. Nonalcoholic steatohepatitis versus steatosis: adipose tissue insulin resistance and dysfunctional response to fat ingestion predict liver injury and altered glucose and lipoprotein metabolism. Hepatology 2012; 56: 933–942.
- Gastaldelli A, Harrison SA, Belfort-Aguilar R et al. Importance of changes in adipose tissue insulin resistance to histological response during thiazolidinedione treatment of patients with nonalcoholic steatohepatitis. Hepatology 2009; 50: 1087–1093.
- Cusi K. Role of obesity and lipotoxicity in the development of nonalcoholic steatohepatitis: pathophysiology and clinical implications. Gastroenterology 2012; **142**: 711–725.e716.
- Fabbrini E, Mohammed BS, Magkos F, Korenblat KM, Patterson BW, Klein S. Alterations in adipose tissue and hepatic lipid kinetics in obese men and women with nonalcoholic fatty liver disease. Gastroenterology 2008; 134: 424–431.
- Fontana L, Eagon JC, Trujillo ME, Scherer PE, Klein S. Visceral fat adipokine secretion is associated with systemic inflammation in obese humans. Diabetes 2007; 56: 1010–1013.
- Tordjman J, Poitou C, Hugol D et al. Association between omental adipose tissue macrophages and liver histopathology in morbid obesity: influence of glycemic status. J Hepatol 2009; 51: 354–362.
- Garg A. Regional adiposity and insulin resistance. J Clin Endocrinol Metab 2004; 89: 4206–4210.
- Nielsen S, Guo Z, Johnson CM, Hensrud DD, Jensen MD. Splanchnic lipolysis in human obesity. J Clin Invest 2004; **113**: 1582–1588.
- Abate N, Garg A, Peshock RM, Stray-Gundersen J, Adams-Huet B, Grundy SM. Relationship of generalized and regional adiposity to insulin sensitivity in men with NIDDM. Diabetes 1996; 45: 1684–1693.
- Abate N, Garg A, Peshock RM, Stray-Gundersen J, Grundy SM. Relationships of generalized and regional adiposity to insulin sensitivity in men. J Clin Invest 1995; 96: 88–98.
- 22. Ferreira I, Henry RMA, Twisk JWR, van Mechelen W, Kemper HCG, Stehouwer CDA. Study AGaHL. The metabolic syndrome, cardiopulmonary fitness, and subcutaneous trunk fat as independent determinants of

arterial stiffness: the Amsterdam Growth and Health Longitudinal Study. Arch Intern Med 2005; **165**: 875–882.

- Goodpaster BH, Thaete FL, Simoneau JA, Kelley DE. Subcutaneous abdominal fat and thigh muscle composition predict insulin sensitivity independently of visceral fat. Diabetes 1997; 46: 1579–1585.
- Sanyal AJ, Brunt EM, Kleiner DE et al. Endpoints and clinical trial design for nonalcoholic steatohepatitis. Hepatology 2011; 54: 344–353.
- 25. Kotronen A, Peltonen M, Hakkarainen A et al. Prediction of non-alcoholic fatty liver disease and liver fat using metabolic and genetic factors. Gastroenterology 2009; **137**: 865–872.
- Steele R. Use of C14-glucose to measure hepatic glucose production following an intravenous glucose load or after injection of insulin. Metabolism 1959; 8: 512–519.
- Finegood DT, Bergman RN, Vranic M. Estimation of endogenous glucose production during hyperinsulinemic-euglycemic glucose clamps. Comparison of unlabeled and labeled exogenous glucose infusates. Diabetes 1987; 36: 914–924.
- Hazlehurst JM, Gathercole LL, Nasiri M et al. Glucocorticoids fail to cause insulin resistance in human subcutaneous adipose tissue in vivo. J Clin Endocrinol Metab 2013; **98**: 1631–1640.
- 29. Diraison F, Pachiaudi C, Beylot M. In vivo measurement of plasma cholesterol and fatty acid synthesis with deuterated water: determination of the average number of deuterium atoms incorporated. Metabolism 1996; **45**: 817–821.
- Diraison F, Pachiaudi C, Beylot M. Measuring lipogenesis and cholesterol synthesis in humans with deuterated water: use of simple gas chromatographic/mass spectrometric techniques. J Mass Spectrom 1997; 32: 81–86.
- Gastaldelli A, Cusi K, Pettiti M et al. Relationship between hepatic/visceral fat and hepatic insulin resistance in nondiabetic and type 2 diabetic subjects. Gastroenterology 2007; 133: 496–506.
- Lomonaco R, Ortiz-Lopez C, Orsak B et al. Role of ethnicity in overweight and obese patients with nonalcoholic steatohepatitis. Hepatology 2011; 54: 837–845.
- Timlin MT, Parks EJ. Temporal pattern of de novo lipogenesis in the postprandial state in healthy men. Am J Clin Nutr 2005; 81: 35–42.
- 34. Diraison F, Moulin P, Beylot M. Contribution of hepatic de novo lipogenesis and reesterification of plasma non esterified fatty acids to plasma triglyceride synthesis during non-alcoholic fatty liver disease. Diabetes Metab 2003; 29: 478–485.
- Lebovitz HE, Banerji MA. Point: visceral adiposity is causally related to insulin resistance. Diabetes Care 2005; 28: 2322–2325.
- Miles JM, Jensen MD. Counterpoint: visceral adiposity is not causally related to insulin resistance. Diabetes Care 2005; 28: 2326–2328.
- 37. Karpe F, Dickmann JR, Frayn KN. Fatty acids, obesity, and insulin resistance: time for a reevaluation. Diabetes 2011; **60**: 2441–2449.
- Tordjman J, Divoux A, Prifti E et al. Structural and inflammatory heterogeneity in subcutaneous adipose tissue: relation with liver histopathology in morbid obesity. J Hepatol 2012; 56: 1152–1158.
- 39. Patel P, Abate N. Role of subcutaneous adipose tissue in the pathogenesis of insulin resistance. J Obes. 2013; **2013**: 489187.
- 40. Bosy-Westphal A, Kossel E, Goele K et al. Association of pericardial fat with liver fat and insulin sensitivity after diet-induced weight loss in overweight women. Obesity 2010; **18**: 2111–2117.