

Mid-Pliocene warming

Hall, Jonathan; Jones, Stephen; Jones, Tom Dunkley; Bendle, James

DOI:

[10.5194/egusphere-egu21-220](https://doi.org/10.5194/egusphere-egu21-220)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Hall, J, Jones, S, Jones, TD & Bendle, J 2021, Mid-Pliocene warming: reducing discrepancies between geological archives and climate models in the NE Atlantic and Nordic Seas. in *EGU General Assembly 2021.*, EGU21-220, European Geosciences Union, EGU General Assembly 2021, 19/04/21.
<https://doi.org/10.5194/egusphere-egu21-220>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

EGU21-220

<https://doi.org/10.5194/egusphere-egu21-220>

EGU General Assembly 2021

© Author(s) 2021. This work is distributed under the Creative Commons Attribution 4.0 License.



Mid-Pliocene warming: reducing discrepancies between geological archives and climate models in the NE Atlantic and Nordic Seas

Jonathan Hall, Stephen Jones, Tom Dunkley Jones, and James Bendle

University of Birmingham, School of Geography, Earth and Environmental Sciences, Birmingham, United Kingdom of Great Britain and Northern Ireland (jrh633@student.bham.ac.uk)

The mid-Pliocene Warm Period (mPWP) is the most recent time slice (3.264–3.025 Ma) during which average global surface temperatures were 2–3°C warmer than preindustrial conditions, within the range estimated by the Intergovernmental Panel on Climate Change (IPCC) for the end of the 21st Century. Global mPWP sea surface temperature (SST) compilations indicate enhanced warming in the NE Atlantic and Nordic Seas, with anomalies of >6°C based on alkenone methods (Dowsett et al., 2012). However, this warming far exceeds the more conservative SST estimates (a rise of 2–3°C) predicted by the Pliocene Research, Interpretation and Synoptic Mapping (PRISM) reconstructions and leading climate models (including HadCM3). Here, we present new mid-Pliocene alkenone SST records from four regional drilling sites (IODP Site U1308, DSDP Site 552, ODP Site 642 and ODP Site 907) to further examine the magnitude of warming in the NE Atlantic and Nordic Seas, and to evaluate regional discrepancies between proxy and model SST estimates. We demonstrate mid-Pliocene SSTs peaked up to 21.5°C and 19.7°C in the NE Atlantic and Nordic Seas, respectively, consistent with existing studies (Robinson et al., 2008; Robinson, 2009). However, we reveal the majority of these SST estimates are derived from GC injections of relatively low total alkenone concentrations (<50 ng/μl), which are susceptible to warming biases caused by chromatographic irreversible adsorption (Grimalt et al., 2001). We subsequently filtered and applied a mathematical correction to our new data to rectify for these warming biases, which results in a reduction in mPWP SSTs, by up to 3.2°C, across all four sites. The corrected (and cooler) alkenone SST records indicate the magnitude of warming in the NE Atlantic and Nordic Seas may be significantly less than previously thought, helping to reduce and explain regional discrepancies between proxy- and model-based SST reconstructions.