

Global, regional, and national mortality among young people aged 10-24 years, 1950-2019

GBD 2019 Adolescent Mortality Collaborators

DOI:

[10.1016/s0140-6736\(21\)01546-4](https://doi.org/10.1016/s0140-6736(21)01546-4)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

GBD 2019 Adolescent Mortality Collaborators 2021, 'Global, regional, and national mortality among young people aged 10-24 years, 1950-2019: a systematic analysis for the Global Burden of Disease Study 2019.', *Lancet (London, England)*, pp. 1593-1618. [https://doi.org/10.1016/s0140-6736\(21\)01546-4](https://doi.org/10.1016/s0140-6736(21)01546-4)

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Global, regional, and national mortality among young people aged 10–24 years, 1950–2019: a systematic analysis for the Global Burden of Disease Study 2019



GBD 2019 Adolescent Mortality Collaborators*



Summary

Background Documentation of patterns and long-term trends in mortality in young people, which reflect huge changes in demographic and social determinants of adolescent health, enables identification of global investment priorities for this age group. We aimed to analyse data on the number of deaths, years of life lost, and mortality rates by sex and age group in people aged 10–24 years in 204 countries and territories from 1950 to 2019 by use of estimates from the Global Burden of Diseases, Injuries, and Risk Factors Study (GBD) 2019.

Methods We report trends in estimated total numbers of deaths and mortality rate per 100 000 population in young people aged 10–24 years by age group (10–14 years, 15–19 years, and 20–24 years) and sex in 204 countries and territories between 1950 and 2019 for all causes, and between 1980 and 2019 by cause of death. We analyse variation in outcomes by region, age group, and sex, and compare annual rate of change in mortality in young people aged 10–24 years with that in children aged 0–9 years from 1990 to 2019. We then analyse the association between mortality in people aged 10–24 years and socioeconomic development using the GBD Socio-demographic Index (SDI), a composite measure based on average national educational attainment in people older than 15 years, total fertility rate in people younger than 25 years, and income per capita. We assess the association between SDI and all-cause mortality in 2019, and analyse the ratio of observed to expected mortality by SDI using the most recent available data release (2017).

Findings In 2019 there were 1.49 million deaths (95% uncertainty interval 1.39–1.59) worldwide in people aged 10–24 years, of which 61% occurred in males. 32.7% of all adolescent deaths were due to transport injuries, unintentional injuries, or interpersonal violence and conflict; 32.1% were due to communicable, nutritional, or maternal causes; 27.0% were due to non-communicable diseases; and 8.2% were due to self-harm. Since 1950, deaths in this age group decreased by 30.0% in females and 15.3% in males, and sex-based differences in mortality rate have widened in most regions of the world. Geographical variation has also increased, particularly in people aged 10–14 years. Since 1980, communicable and maternal causes of death have decreased sharply as a proportion of total deaths in most GBD super-regions, but remain some of the most common causes in sub-Saharan Africa and south Asia, where more than half of all adolescent deaths occur. Annual percentage decrease in all-cause mortality rate since 1990 in adolescents aged 15–19 years was 1.3% in males and 1.6% in females, almost half that of males aged 1–4 years (2.4%), and around a third less than in females aged 1–4 years (2.5%). The proportion of global deaths in people aged 0–24 years that occurred in people aged 10–24 years more than doubled between 1950 and 2019, from 9.5% to 21.6%.

Interpretation Variation in adolescent mortality between countries and by sex is widening, driven by poor progress in reducing deaths in males and older adolescents. Improving global adolescent mortality will require action to address the specific vulnerabilities of this age group, which are being overlooked. Furthermore, indirect effects of the COVID-19 pandemic are likely to jeopardise efforts to improve health outcomes including mortality in young people aged 10–24 years. There is an urgent need to respond to the changing global burden of adolescent mortality, address inequities where they occur, and improve the availability and quality of primary mortality data in this age group.

Funding Bill & Melinda Gates Foundation.

Copyright © 2021 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY 4.0 license.

Introduction

The most recent systematic global analysis to focus on mortality in young people aged 10–24 years is now more than a decade old.¹ Since then, there have been huge changes to patterns of health risk, population growth, and improvements in the availability and quality of

mortality estimates. There has also been wider recognition of the importance of adolescent health to global development,^{2,3} and in harnessing the demographic dividend resulting from forecasted population change.⁴ Adolescents are now included within the UN Every Woman Every Child agenda,⁵ the Countdown to 2030

Lancet 2021; 398: 1593–618

This online publication has been corrected. The corrected version first appeared at thelancet.com on February 24, 2022

See [Comment](#) page 1545

*Collaborators are listed at the end of the paper

Correspondence to:
Dr Joseph L Ward, Great Ormond Street Institute of Child Health, University College London, London WC1N 1EH, UK
joseph.ward@ucl.ac.uk

Research in context

Evidence before this study

We searched PubMed and Embase for studies to answer the research question: “What are current global patterns and previous trends in mortality in adolescents and young people”. We used the following terms in titles or abstracts: (“adolescent” OR “young people” AND (“trends” OR “patterns”) AND (“death” OR “mortality”) AND (“world” OR “global” OR “international”). We found one study that focused on global mortality patterns in adolescents (10–24 years), published in 2009. Using data from 2004, this study reported the considerable burden of global adolescent mortality, and analysed geographic variation in outcomes, but did not explore trends over time. Two further studies reported trends in global mortality or years of life lost (YLLs) in adolescents and young people from 1990. As part of wider analyses of patterns of disease and risk factors in people aged 10–24 years, these studies found considerable variation in improvements in mortality and YLLs by country, age group, and sex. We identified other studies that examined adolescent deaths, but these either included smaller age groups (10–14 years, 10–19 years, or 15–24 years), focused on one region or group of countries, or were part of a wider study of mortality in other age groups and were not focused on adolescents. Although these studies provide useful insights into adolescent mortality, they do not provide a comprehensive global analysis of trends and burden of deaths in this age group.

Added value of this study

This is a comprehensive update of global mortality in young people aged 10–24 years, and the first analysis we are aware of to include estimates for 204 countries and territories,

including all members and associate members of WHO. We found huge changes to adolescent mortality over the study period and increasing variation in outcomes. Mortality in people aged 10–24 years is increasing in many regions, particularly in males, and sex differences have widened in most regions of the world. Adolescent mortality rates were strongly associated with level of development. Demographic change and differences in mortality improvements are shifting the global burden of adolescent mortality towards sub-Saharan Africa, however, where communicable and maternal causes of death continue to predominate. We were able to identify countries with good adolescent mortality outcomes relative to their level of socioeconomic development, which can inform global health strategies for this age group.

Implications of all the available evidence

Despite increased recognition of the importance of adolescent health to global development and future economic prosperity, these findings highlight an ongoing failure to adequately respond to health risks during the adolescent years, and of funding levels that remain insufficient. This is compounded by rapid demographic change in low-resource settings where mortality hazards for adolescents remain high. Indirect effects of the COVID-19 pandemic threaten to restrict progress even further. Approaches must be developed to address growing inequities in mortality in this age group, and to focus on regions of the world and causes of death where improvements have stagnated. To monitor progress reliably, improvements in the quality and availability of adolescent health data are also urgently required.

collaboration to track progress towards the Sustainable Development Goals (SDGs),⁶ and the work of the Global Financing Facility.⁷ Despite this progress, there is also growing concern that global adolescent health priorities are still neglected.^{8,9} Global progress to improve health outcomes for young people has been slow, and major challenges remain in addressing social determinants of adolescent health and mortality, such as unmet contraception need, child marriage, and access to quality secondary education.⁸ Furthermore, the COVID-19 pandemic has ongoing consequences on the health of young people that are not fully understood. Although case fatality and morbidity due to COVID-19 is lower in young people than in older adults,¹⁰ this age group is particularly susceptible to indirect effects of the pandemic, with ongoing disruption to education and employment likely to further hinder progress in health.¹¹

Mortality is a fundamental indicator of health¹² and mortality estimates provide a proxy for broader health outcomes with little or no high quality data. This is particularly pertinent for young people, where data scarcity presents a barrier to understanding health needs in many countries,^{13,14} and the process of determining which metrics best capture adolescent health priorities is

ongoing.¹⁵ Earlier work¹ illustrated that global adolescent mortality is greater than previously recognised and that improvements lagged behind those seen in younger children,¹⁶ despite a high proportion of adolescent deaths being from preventable causes.¹⁷

Mortality is included in the indicator set for monitoring future progress in adolescent health. All-cause mortality in people aged 10–19 years is an indicator within the Global Strategy for Women’s, Children’s and Adolescents’ Health 2016–2030,^{5,18} and SDG 3 includes indicators for causes of death that are highly relevant to young people including maternal mortality (death due to complications during or after pregnancy; SDG indicator 3.1.1), suicide (3.4.2), and deaths due to road traffic injury (3.6.1).¹⁹

Agencies that provide adolescent mortality estimates for a broad range of countries include the UN Population Division, Department of Economic and Social Affairs,²⁰ WHO,²¹ US Census Bureau,²² and the UN Inter-agency Group for Child Mortality Estimation.²³ However, the ongoing Global Burden of Diseases, Injuries, and Risk Factors Study (GBD) provides the only source of frequently updated, age-specific mortality estimates (including uncertainty intervals [UIs]) covering almost all countries and territories in the world.¹² Furthermore, the availability

of estimates in GBD 2019 of all-cause mortality from 1950, and cause-specific estimates from 1980,¹² provides a unique opportunity to analyse long-run trends over time, and so identify both successes and areas requiring global and national investment.

We use GBD 2019 estimates¹² to analyse current and long-term trends in mortality in young people aged 10–24 years globally and in 204 countries and territories. We use this age range to capture the social, biological, and neurocognitive transitions that occur during this stage of the life course,²⁴ and use the terms adolescent and young person synonymously. This manuscript was produced as part of the GBD Collaborator Network and in accordance with the GBD Protocol.

Methods

Overview

We used data provided by the Institute for Health Metrics and Evaluation GBD 2019.¹² Estimates were available for number of deaths, years of life lost (YLLs) and mortality rate per 100 000 population with UIs by sex and age group in 204 countries and territories. Mortality estimates for all causes were available from 1950 to 2019, and by cause of death from 1980 to 2019. GBD 2019 estimations are based on primary data from 86 249 sources, including civil registrations, vital statistics, censuses, disease notifications systems, and household surveys. The GBD 2019 capstone papers and appendices describe these methods in detail,¹² including procedures to standardise primary sources, redistribute non-specific or implausible causes of death, adjust for large spikes in mortality due to conflicts or natural disaster, and model estimates for locations that lack primary data. GBD 2019 complies with the Guidelines for Accurate and Transparent Health Estimates Reporting statement.²⁵

GBD 2019 provides estimates for 281 underlying causes of death across a 4-level hierarchy, shown in appendix 1 (table S2). Level 1 consists of three mutually exclusive and collectively exhaustive categories: communicable, maternal, neonatal, and nutritional diseases; non-communicable diseases; and injuries. Level 2 distinguishes these groups into 21 causes of death, (eg, neoplasms), with level 3 and then level 4 specifying underlying cause in greater detail. We report causes at level 2 of this hierarchy, but modified this to examine leading causes of death in young people aged 10–24 years in more detail. Firstly, we separated the level 2 group self-harm and interpersonal violence into two groups (self-harm [ie, suicide] and interpersonal violence and conflict). Secondly, due to the importance of maternal deaths in this age group, we separated the level 2 group maternal and neonatal disorders into its two level 3 causes (maternal disorders and neonatal disorders).

GBD 2019 mortality estimates were available for 204 countries and territories (hereafter referred to as countries), which includes all members and associate members of WHO. Estimates were also available for

21 GBD regions and 7 GBD super-regions (defined by both geography and income status). Here we primarily report outcomes by GBD super-region, with country-level and region-level estimates available in appendix 2. The seven GBD super-regions are central Europe, eastern Europe, and central Asia; Latin America and the Caribbean; southeast Asia, east Asia, and Oceania; north Africa and the Middle East; south Asia; sub-Saharan Africa; and high income. Note that separate from the high-income GBD super-region, there are countries defined as high income by the World Bank in all other GBD super-regions except sub-Saharan Africa and south Asia. However, when we discuss high-income countries in this analysis we are referring to the GBD high-income super-region. The list of countries included in the analysis, their GBD region, GBD super-region and World Bank income classification for 2021 is found in appendix 1 (table S1).

Analyses

We describe current and long-term trends in all-cause mortality rate and number of deaths from 1950 to 2019, by cause of death from 1980 to 2019; and by sex in people aged 10–14 years, 15–19 years, and 20–24 years. We analyse relative risk for mortality between age groups by dividing the mortality rate in people aged 20–24 years and people aged 15–19 years by the mortality rate in adolescents aged 10–14 years. We analyse variation in outcomes by sex by calculating the ratio of male to female all-cause mortality rate. We analyse between-country variation by calculating the ratio for all-cause mortality rate in people aged 10–24 years in the 90th to 10th centile countries globally to account for extreme values in some location-years.

We then analyse the relationship between mortality rate per 100 000 population in 2019 and country development status using the GBD Socio-demographic Index (SDI).²⁶ The SDI is a summary indicator of social and economic conditions that are strongly correlated with health outcomes. An index value between 0 and 1 is defined for three components: average national educational attainment in those older than 15 years, total fertility rate in those younger than 25 years, and lag-distributed income per capita, using the observed minima and maxima over the estimation period for each component to set the scales. The composite SDI is the geometric mean of these indices for each location-year.²⁶ We analyse the strength of the association between SDI and all-cause mortality rate per 100 000 population by use of Spearman's correlation coefficients. We then identified countries with mortality in young people that is lower or higher relative to their level of development. We did this by calculating the expected value of YLLs by age group and sex based solely on SDI for that location-year using a generalised additive model with a Loess smoother on SDI.²⁶ We then calculated the ratio of observed YLLs to the expected value according to country

See Online for appendix 2

For more on GBD 2019 estimation processes see www.healthdata.org

See Online for appendix 1

| | Total deaths | | | Rate per 100 000 people | |
|--------------------------------------------------|--------------|--------------------------|-------------------------|-------------------------|--------------------------|
| | Estimate | 95% uncertainty interval | Percent of global total | Estimate | 95% uncertainty interval |
| Females | | | | | |
| Global | 581 311 | 531 162–634 825 | 100% | 63.99 | 58.47–69.88 |
| Central Europe, eastern Europe, and central Asia | 13 032 | 12 042–14 207 | 2.2% | 36.70 | 33.92–40.01 |
| High income | 19 715 | 19 532–19 913 | 3.4% | 21.28 | 21.09–21.50 |
| Latin America and Caribbean | 33 376 | 29 924–37 078 | 5.7% | 46.19 | 41.41–51.31 |
| North Africa and Middle East | 38 862 | 34 148–45 205 | 6.7% | 49.81 | 43.77–57.95 |
| South Asia | 224 137 | 196 768–253 694 | 38.6% | 88.05 | 77.30–99.67 |
| Southeast Asia, east Asia, and Oceania | 68 295 | 61 737–75 149 | 11.7% | 34.82 | 31.47–38.31 |
| Sub-Saharan Africa | 183 893 | 159 437–213 930 | 31.6% | 102.57 | 88.93–119.33 |
| Males | | | | | |
| Global | 909 678 | 846 675–974 477 | 100% | 95.41 | 88.80–102.20 |
| Central Europe, eastern Europe, and central Asia | 31 263 | 28 845–33 776 | 3.4% | 83.56 | 77.10–90.28 |
| High income | 48 600 | 48 045–49 176 | 5.3% | 49.85 | 49.28–50.44 |
| Latin America and Caribbean | 96 676 | 87 477–107 076 | 10.6% | 130.62 | 118.19–144.67 |
| North Africa and Middle East | 73 335 | 64 813–83 549 | 8.1% | 88.07 | 77.84–100.34 |
| South Asia | 253 999 | 224 664–286 215 | 27.9% | 93.64 | 82.82–105.51 |
| Southeast Asia, east Asia, and Oceania | 157 271 | 141 742–173 199 | 17.3% | 73.31 | 66.07–80.74 |
| Sub-Saharan Africa | 248 533 | 220 401–279 643 | 27.3% | 141.60 | 125.57–159.32 |

Table: Deaths in people aged 10–24 years in 2019 by sex and GBD super-region



Figure 1: Global number of deaths in people aged 10–24 years from 1950 to 2019 by sex and age group

SDI. We calculate YLLs as the sum of each death multiplied by the standard life expectancy at each age using GBD’s standard life table. Note these data were only available from 1990 to 2017.

Finally, we assess variation in progress to reduce mortality across the early life course, comparing trends and current mortality burden in young people aged 10–24 years with infants younger than 1 year and children aged 1–4 years and 5–9 years. We first report annual rates of change between 1990 and 2019 for all-cause mortality rate for each country by 5-year age groups. We used the β coefficient from linear regression models of mortality rate per 100 000 population against time for each country to estimate annual rate of change (expressed as a percentage). We used these models rather than estimates at the start and end of each period to account for large fluctuations in country mortality due to war or natural disasters. We then identified countries with large differences in mortality performance by age group by comparing country-level all-cause mortality percentile in adolescents with that seen in children younger than 5 years, as this age group has been the focus of global programming and experienced good mortality declines in recent years. Finally, we describe how the proportion of deaths in people younger than 25 years that occur in those aged 10–24 years changed between 1950 and 2019. All analyses were done in Stata 16 (StataCorp, College Station TX, USA).

Role of the funding source

The funder of this study had no role in study design, data collection, data analysis, data interpretation, or writing of the report.

Results

The table shows the number of global deaths and mortality rate per 100 000 population in 2019 by sex and GBD super-region for young people aged 10–24 years. Data by age group (10–14 years, 15–19 years, and 20–24 years) can be found in appendix 1 (tables S3–S5), and country-level estimates are available in appendix 2 (tables S1–S233).

Within a total global population of 1.86 billion people aged 10–24 years, there were around 1.49 million deaths in 2019 (95% UI 1.39–1.59). Just under half of these deaths occurred in people aged 20–24 years (692 000, 95% UI 645 000–738 000), a third occurred in people aged 15–19 years (499 000, 465 000–536 000), and a fifth occurred in people aged 10–14 years (299 000, 276 000–325 000). 51% of all people aged 10–24 years in 2019 were male. 61.0% of all deaths in people aged 10–24 years were in males (910 000, 847 000–974 000), with this proportion increasing with age (56.9% of deaths in people aged 10–14 years; 60.5% of deaths in people aged 15–19 years, and 63.2% of deaths in people aged 20–24 years). Most adolescent deaths in both sexes occurred in south Asia. The fewest adolescent deaths in

both sexes were reported in Central Europe, eastern Europe, and central Asia. Mortality rates in 2019 increased with age across adolescence in all regions of the world, particularly in males, but with considerable variation. Within GBD super-regions, the relative risk of mortality in young adulthood (20–24 years) compared with early adolescence (10–14 years) ranged from 6.6 in the high-income GBD super-region in males to 1.9 in north Africa and the Middle East in females.

In 2019, the mortality rate per 100 000 in young people aged 10–14 years ranged from 7.32 (95% UI 6.90–7.78) in Denmark to 187.25 (165.74–217.44) in Central African Republic in males, and from 6.01 (5.79–6.25) in Denmark to 115.73 (107.50–129.23) in Central African Republic in females. The mortality rate in young people age 15–19 years ranged from 22.55 (21.50–23.72) in Denmark to 370.28 (305.42–448.95) in Lesotho in males, and 12.15 (12.02–12.29) in Japan to 248.02 (169.99–348.35) in Lesotho in females. The mortality rate in people aged 20–24 years ranged from 36.24 (35.48–37.05) in Singapore to 559.77 (434.53–670.69) in Lesotho in males and 14.44 (14.05–14.85) in Singapore to 445.22 (303.38–636.53) in Lesotho in females (appendix 1, figures S6–S11; appendix 2, tables S1–S233).

Globally, 32.7% of deaths in young people aged 10–24 years in 2019 were due to transport injuries, unintentional injuries, or interpersonal violence or conflict; 32.1% were due to communicable, nutritional, or maternal causes; 27.0% were due to non-communicable diseases; and 8.2% were due to self-harm. However, there was large variation in the leading causes of adolescent death by GBD super-region, sex, and age. In males aged 10–14 years, the most common cause of death was unintentional injury in all GBD super-regions except high income, where neoplasms were the leading cause of death, and south Asia and sub-Saharan Africa, where enteric infections were the leading cause of death, and North Africa and Middle East where this was transport injuries. In males aged 15–24 years, the most common cause of death was transport injury in all GBD super-regions except Latin America and the Caribbean, where the leading cause of death was interpersonal violence and conflict, and central Europe, eastern Europe and central Asia, where the leading cause of death was self-harm (appendix 1, figures S50–S112).

The leading causes of death in females aged 10–14 years were neoplasms (Latin America and the Caribbean; central Europe, eastern Europe, and central Asia; and high income), unintentional injury (southeast Asia, east Asia, and Oceania), enteric infections (south Asia) and transport injuries (north Africa and the Middle East). In females aged 15–19 years, the leading causes of death were transport injuries (high income; north Africa and the Middle East; and southeast Asia, east Asia, and Oceania), self-harm (south Asia and central Europe, eastern Europe, and central Asia), interpersonal violence and conflict (Latin America and the Caribbean). The most frequent causes of

death in females aged 20–24 years were transport injuries (high income), neoplasms (southeast Asia, east Asia, and Oceania and central Europe, eastern Europe, and central Asia), interpersonal violence and conflict (Latin America and the Caribbean), cardiovascular diseases (north Africa and the Middle East), and self-harm (south Asia). HIV/AIDS and sexually transmitted infections were the leading causes of death in sub-Saharan Africa in females in all adolescent age groups. Although not the leading cause in any GBD super-region, maternal death was still in the three most common causes of death in females aged




Figure 2: Mortality rate per 100 000 population in people aged 10–24 years from 1950 to 2019 by sex and GBD super-region
GBD=Global Burden of Diseases, Injuries, and Risk Factors Study.

20–24 years in sub-Saharan Africa, north Africa and the Middle East, and south Asia, and was the fourth most common cause of death in Latin America and the Caribbean.

Annual global deaths in adolescents aged 10–24 years reduced by 21.7% between 1950 and 2019, during which time the world population in this age group increased by 157% (figure 1; appendix 1, figure S2). The largest decrease in deaths was in females (30.0%, vs 15.3% in males) despite similar population growth (154% in females and 161% in males). There were also large differences in trends in deaths by age group; in children aged 10–14 years, deaths decreased by 37.5% in males and 37.9% in females; in people aged 15–19 years, deaths decreased by 17.8% in males and 27.9% in females; and in people aged 20–24 years, deaths decreased by 26.9% in females and increased by 0.8% in males.

In young people aged 10–14 years, deaths decreased between 1950 and 2019 in all GBD super-regions except sub-Saharan Africa, where deaths increased by 140.6% in males and 144.3% in females. The largest decrease in deaths since 1950 in young people aged 10–14 years were in high-income countries (86.0% reduction in males and 87.3% reduction in females). In males aged 15–24 years, deaths increased in all GBD super-regions except high income; central Europe, eastern Europe, and central Asia; and southeast Asia, east Asia, and Oceania. The greatest increase in deaths in males aged 15–24 years

was in sub-Saharan Africa, where deaths increased by 180.5% in males aged 15–19 years and 218.2% in males aged 20–24 years. The greatest decrease in deaths in males aged 15–24 years was in the high-income GBD super-region, where deaths decreased by 72.6% in males aged 15–19 years and 66.0% in males aged 20–24 years. In females aged 15–24 years, deaths decreased in all GBD super-regions except north Africa and the Middle East and sub-Saharan Africa. The greatest increase in deaths in females aged 15–24 years was in sub-Saharan Africa in both females aged 15–19 years (175.7% increase) and 20–24 years (164.3% increase). The greatest decrease in deaths in females aged 15–24 years was in high-income countries in both females aged 15–19 years (81.0% decrease) and 20–24 years (79.9% decrease).

Figure 2 shows mortality rate per 100 000 population in people aged 10–14 years, 15–19 years, and 20–24 years by sex and GBD super-region between 1950 and 2019. In people aged 10–14 years, global mortality rates reduced by 74.6% in males and 74.3% in females, with the greatest relative change in mortality in high-income countries (reduction of 88.2% in males and 89.0% in females). The GBD super-region with the smallest relative change in mortality rate was sub-Saharan Africa, with reductions of 65.4% in males and 63.9% in females. All other regions had a 72–88% reduction in mortality rate over this period in males and females aged 10–14 years.

Global mortality rate per 100 000 population in people aged 15–19 years decreased by 68.7% in males and 71.7% in females between 1950 and 2019. The region with the greatest relative reduction in mortality in this age group was southeast Asia, east Asia, and Oceania, where the mortality rate declined by 80.1% in males and 86.1% in females. The region with the lowest relative reduction in mortality rate was sub-Saharan Africa, with relative reductions of 57.8% in males and 59.4% in females.

Global mortality rate per 100 000 population in people aged 20–24 years reduced by 63.4% in males and 72.7% in females between 1950 and 2019. Relative declines in mortality were greatest in southeast Asia, east Asia, and Oceania, with reductions of 77.7% in males and 86.4% in females. In males aged 20–24 years, the lowest relative decline was in central Europe, eastern Europe, and central Asia, where mortality rates reduced by 46.1% since 1950. Among females aged 20–24 years, the lowest relative reduction in mortality was seen in sub-Saharan Africa, where mortality rates decreased by 60.7%.

In 2019, the ratio of male to female mortality rate per 100 000 population was 1.2 for people aged 10–14 years, 1.5 for people aged 15–19 years, and 1.7 for people aged 20–24 years (appendix 1, figure S20). Mortality rate per 100 000 people aged 10–24 years was higher in males in all regions of the world except south Asia, where outcomes were similar. Sex differences in mortality were greatest in Latin America and the Caribbean in people aged 20–24 years, where the mortality rate per 100 000 in males





Figure 3: Ratio of mortality rate per 100 000 population in the 90th centile country to 10th centile country in people aged 10–24 years between 1950 and 2019 by sex and age group

was more than three times that in females. The ratio of male to female mortality rate increased between 1950 and 2019 for older adolescents aged 15–24 years in all GBD super-regions except high income and central Europe, eastern Europe, and central Asia, where it has reduced since the mid-1990s.

Inequality in mortality rate per 100 000 population between countries was reported using the ratio of

mortality rate for the highest decile country (90th centile) to lowest decile country (10th centile) in each year. In 2019, variations in mortality by country were greatest in adolescents aged 10–14 years, with the mortality rate in the 90th centile country around 9 times higher than that in the 10th centile country for both males and females. Inequality in outcomes by country has also increased over time for all age groups since 1950, although this



(Figure 4 continues on next page)




Figure 4: Percentage of total deaths by cause in people aged 15–19 years from 1980 to 2019 by sex and GBD super-region
 (A) Females (B) Males. GBD=Global Burden of Diseases, Injuries, and Risk Factors Study.

seems to have decreased in females aged 20–24 years since around 2000 (figure 3).

The percentage of total deaths by cause from 1980 to 2019 in each GBD super-region is shown in figure 4 for young people aged 15–19 years. Bump charts that rank cause of death between 1980 and 2019 for each age group,

sex, and GBD super-region are shown in appendix 1 (figures S50–S112).

We report large reductions in the contribution of communicable and maternal causes to total adolescent deaths between 1980 and 2019 in Latin America and the Caribbean; south Asia; southeast Asia, east Asia, and the Caribbean; and north Africa and the Middle East.




Figure 5: Ratio of observed to expected years of life lost by Sociodemographic Index from all causes in people aged 15–19 years in 2017 from 195 countries

Furthermore, although maternal deaths still contribute a substantial proportion of global deaths in females aged 20–24 years, this has fallen substantially since 1980. Maternal deaths halved as a proportion of total deaths between 1980 and 2019 in southeast Asia, east Asia, and Oceania; north Africa and the Middle East; and central Europe, eastern Europe, and central Asia, with large reductions also seen in south Asia and Latin America and the Caribbean. However, in sub-Saharan Africa, the five most common causes of death in 2019 in females aged 10–24 years were communicable or maternal causes, with communicable causes contributing to three of the top five causes in males. HIV/AIDS and sexually transmitted diseases have been the leading causes of death in this GBD super-region in females aged 15–24 years since the early 1990s, and in females aged 10–14 years since the early 2000s.

In central Europe, eastern Europe, and central Asia, self-harm rose from the third highest cause of death in males aged 15–24 years in 1980 to be the leading cause of death (now contributing to more than 20% of all deaths), and from fourth highest to the leading cause of death in females aged 15–19 years and second highest cause of death in females aged 20–24 years. In males aged 10–14 years, self-harm is now the fourth most common cause of death in this super-region (almost 9% of all deaths). By contrast, in southeast Asia, east Asia, and Oceania, self-harm in females aged 15–24 years fell from the leading cause of death in 1980 (around 15% of all deaths), to fifth highest in females aged 15–19 years and sixth highest in females aged 20–24 years in 2019 (around 7% of all deaths in both age groups). In the high-income GBD super-region, deaths due to substance misuse disorders in young adults aged 20–24 years rose from around 1% of total deaths in 1980 to 14–16% in 2019 in both sexes, with substantial increases also seen in people aged 15–19 years.

Increasing SDI in 2019 was strongly associated with lower all-cause mortality rate per 100 000 people in all

adolescent age groups (appendix 1, table S6). Figure 5 shows the ratio of observed to expected YLLs by SDI from all causes for people aged 15–19 years in 2017. Equivalent figures for people aged 10–14 years and 20–24 years are shown in appendix 1 (figures S26–28), with estimates from 1990 to 2017 for all countries shown in appendix 1 (figures S29–S49). Observed YLLs from all causes in people aged 15–19 years were at least 20% higher than expected by level of SDI in 41 countries, including both Brazil and Pakistan, which collectively contribute to around 10% of deaths in this age group. In people aged 15–19 years, the ratio of observed to expected YLLs in 2017 was highest in Syria (4.02), but from 1990 to 2010 (before the start of the civil war), observed YLLs in Syria were consistently around 30% lower than expected by level of SDI. 84 countries had observed YLLs in people aged 15–19 years at least 20% lower than expected by level of SDI. The countries with the lowest ratio of observed to expected YLLs in this age group were the Maldives (0.30), Spain (0.33), and Singapore (0.35). Other notable countries included China (0.49) and Ethiopia (0.67), which together contribute more than 7% of global deaths in people aged 15–19 years.

Annual percentage decline in mortality rate per 100 000 population in infants younger than 1 year and children and young people aged 1–4 years, 5–9 years, 10–14 years, 15–19 years, and 20–24 years by sex, country, and globally between 1990 and 2019 is shown in figure 6. Rates of decline were highest in children aged 1–4 years and lowest in young people aged 15–24 years. In children aged 1–4 years, global mortality rates decreased by around 2.4% per year since 1990 in both males and females, compared with people aged 15–19 years, in whom mortality rates decreased by 1.3% in males and 1.6% in females. The range in mortality rate change between countries was also greater in adolescents than in younger children (appendix 1, figure S24).




Figure 6: Annual percentage change in mortality rate per 100 000 population in people aged 0–24 years 1990–2019 by sex, age group, and GBD super-region

The dashed line represents global annualised percentage change in mortality rate. Each coloured dot represents the percentage change in a single year for each country, coloured by GBD super-region. Data for Rwanda (both sexes, 15–19 years) and Bosnia and Herzegovina (females aged 1–4 years) are excluded as the range was restricted to 10% annual change. GBD=Global Burden of Diseases, Injuries, and Risk Factors Study.

Mortality rate performance varied within countries for different age groups, particularly in males. Figure 7 shows mortality rate percentile for males younger than 5 years compared with males aged 15–19 years for each country in 2019, with the equivalent figure for females shown in appendix 1 (figure S25). Although in most countries the mortality rate percentiles for these age groups are similar, there are notable exceptions. For example, in Brazil and Venezuela, mortality rates for males younger than 5 years are around the 60th centile but in the 90th centile for males aged 15–19 years. Similarly, mortality rates in Ukraine and Thailand are around the 30th centile for children younger than 5 years, but above the 75th centile for young people aged 15–19 years. By contrast, the mortality rate in India for people younger than 5 years is around the 70th centile, and between the 30th and 40th centiles for those aged 15–19 years.

Figure 8 shows the proportion of global deaths in people aged 0–24 years that occur in people aged 10–24 years. This proportion increased from 9·5% in

1950 to 21·6% by 2019 and increased in all GBD super-regions between 1950 and 2019, with the greatest change seen in Latin America and the Caribbean in males (from 7·5% to 39·2%). In the high-income GBD super-region, deaths in adolescence now account for more than half of all deaths before 25 years of age. In nine countries (Estonia, Thailand, Saudi Arabia, Finland, Puerto Rico, Slovenia, Monaco, Cook Islands, and Andorra), more than 70% of deaths in males younger than 25 years now occur in people aged 10–24 years.

Discussion

Despite marked reductions in total numbers of deaths and mortality rates for adolescents over the past 70 years, improvements have lagged behind those seen in younger children, variation in outcomes between countries has increased, and inequities by sex have worsened. Global numbers of deaths in adolescence remain high, at around 4000 each day, most of which are from preventable causes.




Figure 7: Percentiles of mortality rate per 100 000 population in males aged younger than 5 years and 15-19 years in 2019 by country and GBD super-region
 Size of each mark is proportional to the population of males aged 15-19 years. Due to space constraints, not all countries are labelled. GBD=Global Burden of Diseases, Injuries, and Risk Factors Study.

We found widening variation in all-cause mortality between countries, particularly during early adolescence, and huge differences in the leading causes of death between different regions of the world, as others have reported.^{17,27} Variation in all-cause mortality between countries seems likely to increase further, as population growth in this age group is highest in countries with the worst mortality outcomes (appendix 1, figures S21-S23). Around 20% of people aged 10-24 years live in sub-Saharan Africa, and this is set to rise to a third by 2050,⁴ which presents huge challenges to the improvement of adolescent mortality. Of the ten countries with the highest mortality rates in this age group, eight are in sub-Saharan Africa, and decreases have been far slower than in other GBD super-regions. These changes are shifting the global burden of adolescent mortality towards sub-Saharan Africa, where 29.0% of deaths in adolescents now occur, compared with 8.1% in 1950. Sub-Saharan Africa has already replaced south Asia as the main contributor to global deaths in males aged 10-19 years, and is set to do the same in males aged 20-24 years and females aged 10-19 years if current trends continue.

We found marked sex differences in adolescent mortality, with notably higher mortality and slower rates




Figure 8: Proportion of deaths in people younger than 25 years occurring in adolescents aged 10-24 years in 1950 and 2019
 The size of each plot is proportional to the total number of deaths in people younger than 25 years in that year.

of decline in young men than in young women. Indeed, more adolescent males died in 2019 than adolescent females in 1950, and inequalities in mortality by sex also seem to be widening in many regions of the world. These differences reflect the increasing burden of deaths

due to injuries and violence in this age group, particularly in Latin America and the Caribbean, and the rise in deaths due to substance misuse in high-income settings, which predominantly affect young men. Addressing inequities in access to health services, the social conditions in which young women live, and the impact of gender-based violence are fundamental to improving adolescent health globally and have understandably been the focus of previous programming.¹⁹ However, inequitable gender norms are also damaging to adolescent males, and advancing the health of all adolescents requires action to reduce inequities in outcomes wherever they occur.

Consistent with previous studies,^{14,16,17,27} we found declines in adolescent mortality to have been slower than in younger children, particularly in males (annualised decrease in mortality in males aged 1–4 years almost twice that in males aged 15–19 years, and annualised decrease in mortality in females aged 1–4 years about 50% higher than in females aged 15–19 years).

A previous analysis¹⁶ of long-term mortality trends using WHO estimates found that differences in mortality reduction by age resulted in people aged 15–24 years replacing those aged 1–4 years as the group with the highest mortality burden in many countries. 15–19 years was highlighted as the age group with the slowest mortality decline among people aged 0–19 years using GBD data,²⁷ and Masquelier and colleagues¹⁴ found that the decrease in mortality in children aged 5–9 years exceeded that in young people aged 10–14 years between 1990 and 2016 using estimates from the UN.¹⁴ The continued neglect of this age group is highlighted by several (mostly middle-income) countries such as Brazil and Mexico having good outcomes for children aged 0–4 years but persistently high adolescent mortality (figure 7), which suggests that more adolescent-specific interventions are needed in these settings. Almost a quarter of deaths in people aged 0–24 years now occur in people aged 10–24 years, and this proportion has more than doubled since 1950. In high-income countries, these account for almost half of deaths before 25 years of age, and in countries with particularly high mortality due to violence, adolescent deaths now account for up to three-quarters of early life course mortality.

This poor progress in reducing mortality might reflect the omission of adolescents from most global health investments. Adolescents were largely absent from the Millennium Development Goal (MDG) agenda, and although they will have benefited from public health interventions aimed at other groups, they have not had the accelerated decreases in mortality seen in infants and younger children that is attributed to MDG programming.²⁷ The SDGs include indicators that are highly relevant to young people, but do not provide a comprehensive mechanism to address the unique health needs of this age group,²⁷ and a list of metrics to best capture global adolescent health priorities has not yet

been defined.¹⁵ The 2016 Global Strategy for Women's, Children and Adolescents' Health and other initiatives have increased recognition of the crucial role of adolescents to the sustainable development agenda and global targets for Universal Health Coverage.⁵ However, investment remains inadequate, and specific challenges to improvement of adolescent health outcomes and achieving core Universal Health Coverage indicators in this age group continue to be overlooked.^{8,9}

These factors are compounded by limited evidence for effective adolescent health interventions needed to inform investment.²⁸ A key focus to improve outcomes in this age group should be to address these knowledge gaps and better establish which interventions work, guided by identifying and measuring key indicators that capture adolescent health priorities.¹⁵ Investment in existing evidenced-based actions to prevent causes of death that predominate during adolescence is also likely to improve outcomes, but is currently inadequate. This should include improving water safety; preventing unintentional injuries;²⁹ and targeting key behavioural, legal, and structural risk factors for road traffic deaths.^{30,31} However, global increases in injury prevention spending have been lower than those on other public health interventions³² and progress towards reducing road traffic deaths in line with SDG targets is insufficient.³⁰ Self-harm has emerged as a leading cause of adolescent mortality, and now accounts for around 20% of all deaths in people aged 15–24 years in the high-income and central Europe, eastern Europe, and central Asia GBD super-regions. Self-harm is also the leading cause of death in south Asia for females aged 15–24 years and second highest cause for males aged 20–24 years. Although understanding these trends is complex and solutions to improve outcomes need to be country-specific, investment in evidence-based interventions to improve mental health in this age group^{33,34} and measures to restrict access to firearms and chemicals used in suicide are likely to be beneficial.^{35,36} However, global improvements to the quality and accessibility of mental health services have been slow, disproportionately affecting adolescents and young people.³⁴ Furthermore, strategies to improve outcomes for communicable and maternal causes of death, which still contribute to around a third of global deaths in people aged 10–24 years, also need to be specifically tailored for this age group.³⁷ Although evidence-based interventions around adolescent sexual and reproductive health are available, these are mainly focused on high-income countries, and knowledge gaps remain.³⁸

We found several large spikes in mortality over the study period, particularly in older male adolescents, as a result of violent conflict and natural disasters. Deaths due to interpersonal violence are also impeding progress in reduction of all-cause mortality in males in many countries, particularly in Latin America and the Caribbean where there has been little to no improvement

in all-cause mortality in males aged 15–24 years over the past 20 years. Global strategies to improve adolescent health outcomes must include efforts to mitigate the effect of interpersonal violence and conflicts on young people. Existing humanitarian response guidance highlights specific vulnerabilities of adolescents during natural disasters and conflict.³⁹ However, the evidence base to manage these in low-income and middle-income countries remains weak,⁴⁰ and further work is required to understand adolescent health needs in these situations.

Reducing mortality in young people also requires an understanding of the broader social determinants of adolescent health, and how structural changes such as rapid urbanisation and technological and economic development might affect young people differently from other age groups. The importance of primary education to population health is well described, but national progress in secondary education is also associated with large improvements in all-cause mortality and other important health outcomes for young people.⁴¹ The increasing numbers of adolescents growing up in urban settings could extend opportunities for education, in addition to potential economic benefits for young people and their families. However, rapid, unplanned urbanisation can also increase health risks that are pertinent to adolescents, including those related to injury, separation from family support through migration, exposure to violence, substance misuse, and unsafe employment.^{2,42} The effect of other macro-level health determinants seems to differ across the early life course, with national wealth a weaker predictor of mortality in adolescents than it is in young children.⁴³ Indeed, for some common causes of death in adolescents (eg, road injuries), rapid economic growth can result in a transient increase in mortality, as the introduction of safety legislation and appropriate infrastructure might lag behind rising demand for transportation.^{44,45} By contrast, inequality in income distribution within countries seems to be pervasively harmful throughout the early life course, and thus might be of greater relative importance to outcomes during adolescence.⁴³

Our data are limited by factors inherent in the production of GBD mortality estimates, and by the availability of authoritative mortality data for adolescents. Global coverage of civil registration systems is of varying quality, and progress to improve these systems has been minimal.⁴⁶ Primary data sources for adolescents are particularly scarce, as attempts to develop alternative methods to measure mortality have focused on other groups.¹³ Data availability and accuracy are further impeded by ongoing conflicts and associated migration in many countries. Where data are available, there are often long delays in reporting outcomes. Analysis of the global adolescent mortality burden is therefore reliant on modelled data, and the estimates we report here need to be viewed within that context. Limitations within the

GBD estimation process and the paucity of mortality data for this age group are reflected in wide UIs for many time periods, locations, and causes of death.

Using alternative data sources with different modelling techniques might have provided different results. Estimates for global number of deaths in people aged 10–24 years in 2019 produced by the UN Interagency Group for Child Mortality Estimation were 8–19% higher than GBD 2019 (appendix 1, figures S113–S115), and 10–20% lower than those in the World Population Prospects report.²⁰ Although variation in available mortality estimates for high-income countries have been highlighted,⁴⁷ the main discrepancies are within sub-Saharan Africa, where the future global burden of mortality in adolescents will be concentrated. This further highlights the need to expand primary data collection for adolescent health outcomes in this region.

We used SDI to identify countries with lower or higher adolescent mortality than would be expected from country income per capita, years in education, and fertility younger than 25 years; indicators for socioeconomic development that are particularly relevant for adolescents. However, other disaggregated metrics of development could provide additional insights, and further detailed analyses are required to explore the contribution of trends in key social determinants on adolescent mortality, which are likely to vary in different regions of the world. We report estimates at the country level, and future analyses of adolescent mortality should include sub-national trends. Where these are available within GBD they have wide variation in outcomes. Using level 2 of the GBD cause of death hierarchy allows a description of high-level trends in mortality in young people, but further analyses using more granular cause of death data are needed to increase understanding of the patterns we describe. Analysis of associations between health-care quality and adolescent mortality by use of the Healthcare Access and Quality Index⁴⁸ provided by GBD could provide further insights, and looking beyond mortality and describing key trends in morbidity in young people should also be the focus of future study.

This analysis does not include the impact of the COVID-19 pandemic on young people. Although susceptibility to SARS-CoV-2 in young people is similar to that in older groups,⁴⁹ case fatality has been much lower than in older adults,¹⁰ and the direct effect on mortality in this age group is likely to be small. More concerning are the indirect effects of COVID-19 on the future health of young people, particularly through disruption to health services caused by the pandemic, the impact on opportunities for education and employment and related consequences to health and nutrition, and the exacerbation of existing inequalities.⁵⁰ The ongoing closure of schools and universities in many countries will have lasting effects on the health of young people. Further, this age group are more likely to work in sectors vulnerable to restrictions introduced during the

pandemic than are older adults, and at a greater risk of losing employment and reporting reduced earnings. There is some evidence to suggest that young people have been more affected by worsening mental health than older adults, and concern that suicide risk in early adolescence might have increased.^{51,52} Broader effects of the pandemic to key adolescent health determinants are likely to include those related to economic instability, conflict, and failing to prioritise the climate crisis.⁵³ The extent to which these determinants will affect wellbeing in young people, and prospects to improve health outcomes including mortality, are not understood and warrant further study.

Our findings reveal a persistent failure by policy makers to adequately address global health risks during adolescence or respond to changes in the global burden of early life course mortality. Despite increased understanding of the importance of adolescents to global development, funding remains insufficient and the challenge to improve outcomes is increasing. Investment in this age group builds on health improvements achieved in younger children, will affect future adult health trajectories and those of the next generation,³ and will be an important determinant of future economic development.^{2,17,54} Renewed emphasis is urgently needed to reduce inequities in outcomes in this age group, improve the availability and quality of primary data, and establish mechanisms to use these data to better inform global health policy, focusing on regions of the world where mortality is increasing.

GBD 2019 Adolescent Mortality Collaborators

Joseph L Ward, Peter S Azzopardi, Kate Louise Francis, John S Santelli, Vegard Skirbekk, Susan M Sawyer, Nicholas J Kassebaum, Ali H Mokdad, Simon I Hay, Foad Abd-Allah, Amir Abdoli, Mohammad Abdollahi, Aidin Abedi, Hassan Abolhassani, Lucas Guimarães Abreu, Michael R M Abrigo, Eman Abu-Gharbieh, Abdelrahman I Abushouk, Oladimeji M Adebayo, Victor Adekanmbi, Davoud Adham, Shailesh M Advani, Khashayar Afshari, Anurag Agrawal, Tauseef Ahmad, Keivan Ahmadi, Anwar E Ahmed, Budi Aji, Blessing Akombi-Inyang, Fares Alahdab, Ziyad Al-Aly, Khurshid Alam, Fahad Mashhour Alanezi, Turki M Alanzi, Jacqueline Elizabeth Alcalde-Rabanal, Biresaw Wassihun Alemu, Samar Al-Hajj, Robert Kaba Alhassan, Saqib Ali, Gianfranco Alicandro, Mehran Alijanzadeh, Syed Mohamed Aljunid, Amir Almasi-Hashiani, Nihad A Almasri, Hesham M Al-Mekhlafi, Jordi Alonso, Rajaa M Al-Raddadi, Khalid A Altirkawi, Nelson Alvis-Guzman, Azmeraw T Amare, Saeed Amini, Arya Aminorroaya, Arianna Maeveer L Amit, Dickson A Amugsi, Robert Ancuceanu, Deanna Anderlini, Catalina Liliana Andrei, Sofia Androuti, Fereshteh Ansari, Iman Ansari, Carl Abelardo T Antonio, Davood Anvari, Raziq Anwer, Seth Christopher Yaw Appiah, Jalal Arabloo, Morteza Arab-Zozani, Johan Årnlöv, Malke Asaad, Mehran Asadi-Aliabadi, Ali A Asadi-Pooya, Maha Moh'd Wahbi Atout, Marcel Ausloos, Elvis Korku Avenyo, Leticia Avila-Burgos, Beatriz Paulina Ayala Quintanilla, Getinet Ayano, Yared Asmare Aynalem, Samad Azari, Zelalem Nigussie Azene, Mohammad Hossein Bakhshaei, Shankar M Bakkannavar, Maciej Banach, Palash Chandra Banik, Miguel A Barboza, Suzanne Lyn Barker-Collo, Till Winfried Bärnighausen, Sanjay Basu, Bernhard T Baune, Mohsen Bayati, Neeraj Bedi, Ettore Beghi, Tariku Tesfaye Bekuma, Arielle Wilder Bell, Michelle L Bell, Corina Benjet, Isabela M Bensenor, Abadi Kidanemariam Berhe, Kidanemariam Berhe, Adam E Berman, Akshaya Srikanth Bhagavathula, Nikha Bhardwaj, Pankaj Bhardwaj, Kritika Bhattacharyya, Suraj Bhattarai, Zulfiqar A Bhutta, Ali Bijani,

Boris Bikbov, Antonio Biondi, Tesega Tesega Mengistu Birhanu, Raaj Kishore Biswas, Somayeh Bohlouli, Srinivasa Rao Bolla, Archith Boloor, Rohan Borschmann, Soufiane Boufous, Nicola Luigi Bragazzi, Dejana Braithwaite, Nicholas J K Breitborde, Hermann Brenner, Gabrielle B Britton, Richard A Burns, Sharath Burugina Nagaraja, Zahid A Butt, Florentino Luciano Caetano dos Santos, Luis Alberto Cámera, Ismael R Campos-Nonato, Julio Cesar Campuzano Rincon, Rosario Cárdenas, Giulia Carreras, Juan J Carrero, Felix Carvalho, Joao Mauricio Castaldelli-Maia, Carlos A Castañeda-Orjuela, Giulio Castelpietra, Ferrán Catalá-López, Ester Cerin, Joht Singh Chandan, Hsing-Yi Chang, Jung-Chen Chang, Jaykaran Charan, Vijay Kumar Chattu, Sarika Chaturvedi, Jee-Young Jasmine Choi, Mohiuddin Ahsanul Kabir Chowdhury, Devasahayam J Christopher, Dinh-Toi Chu, Michael T Chung, Sheng-Chia Chung, Flavia M Cicuttini, Traian Vasile Constantin, Vera Marisa Costa, Saad M A Dahlawi, Haijiang Dai, Xiaochen Dai, Giovanni Damiani, Lalit Dandona, Rakhii Dandona, Parnaz Daneshpajouhnejad, Aso Mohammad Darwesh, Claudio Alberto Dávila-Cervantes, Kairat Davletov, Fernando Pio De la Hoz, Diego De Leo, Nikolaos Dervenis, Rupak Desai, Assefa Desalew, Keshab Deuba, Samath Dhamminda Dharmaratne, Govinda Prasad Dhungana, Mostafa Dianatinasab, Diana Dias da Silva, Daniel Diaz, Alireza Didarloo, Shirin Djallalinia, Fariba Dorostkar, Chirag P Doshi, Leila Doshmangir, Kerrie E Doyle, Andre Rodrigues Duraes, Mohammad Ebrahimi Kalan, Sanam Ebtehaj, David Edvardsson, Maha El Tantawi, Islam Y Elgendy, Shaimaa I El-Jaafary, Aisha Elsharkawy, Babak Eshrati, Sharareh Eskandarieh, Saman Esmaeilnejad, Firooz Esmaeilzadeh, Sadaf Esteghamati, Andre Faro, Farshad Farzadfar, Nazir Fattahi, Valery L Feigin, Tomas Y Ferede, Seyed-Mohammad Fereshtehnejad, Eduarda Fernandes, Pietro Ferrara, Irina Filip, Florian Fischer, James L Fisher, Nataliya A Foigt, Morenike Oluwatoyin Folayan, Artem Alekseevich Fomenkov, Masoud Foroutan, Takeshi Fukumoto, Mohamed M Gad, Abhay Motiramji Gaidhane, Silvano Gallus, Teshome Gebre, Ketema Bizuwork Gebremedhin, Gebreamlak Gebremedhn Gebremeskel, Leake Gebremeskel, Assefa Ayalew Gebreslassie, Hailay Abrha Gesesew, Keyghobad Ghadiri, Mansour Ghafourifard, Farhad Ghamari, Ahmad Ghashghaee, Syed Amir Gilani, Elena V Gnedovskaya, Myron Anthony Godinho, Mahaveer Golechha, Srinivas Goli, Philimon N Gona, Sameer Vali Gopalani, Giuseppe Gorini, Michal Grivna, Mohammed Ibrahim Mohialdeen Gubari, Harish Chander Gugrani, Rafael Alves Guimarães, Yuming Guo, Rajeev Gupta, Juanita A Haagsma, Nima Hafezi-Nejad, Teklehaimanot Gereziher Haile, Arvin Haj-Mirzaian, Arya Haj-Mirzaian, Brian J Hall, Randah R Hamadeh, Kanaan Hamagharib Abdullah, Samer Hamidi, Demelash Woldeyohannes Handiso, Asif Hanif, Graeme J Hankey, Hamidreza Haririan, Josep Maria Haro, Ahmed I Hasaballah, Abdiwahab Hashi, Amr Hassan, Soheil Hassanipour, Hadi Hassankhani, Khezir Hayat, Reza Heidari-Soureshjani, Claudiu Herteliu, Fatemeh Heydarpour, Hung Chak Ho, Michael K Hole, Ramesh Holla, Praveen Hoogar, Mostafa Hosseini, Mehdi Hosseinzadeh, Mihaela Hostiu, Sorin Hostiu, Mowafa Househ, Mohamed Hsairi, Tanvir M Huda, Ayesha Humayun, Rabia Hussain, Bing-Fang Hwang, Ivo Iavicoli, Segun Emmanuel Ibitoye, Olayinka Stephen Ilesanmi, Irena M Ilic, Milena D Ilic, Leeberk Raja Inbaraj, Nirun Intarut, Usman Iqbal, Seyed Sina Naghibi Irvani, M Mofizul Islam, Sheikh Mohammed Shariful Islam, Hiroyasu Iso, Rebecca Q Ivers, Mohammad Ali Jahani, Mihajlo Jakovljevic, Amir Jalali, Manthan Dilipkumar Janodia, Tahereh Javaheri, Panniyammakal Jeemon, Ensiyeh Jenabi, Ravi Prakash Jha, Vivekanand Jha, John S Ji, Jost B Jonas, Kelly M Jones, Farahnaz Joukar, Jacek Jerzy Jozwiak, Petur B Juliusson, Mikko Jürisson, Ali Kabir, Zubair Kabir, Leila R Kalankesh, Rohollah Kalhor, Naser Kamyari, Tanuj Kanchan, André Karch, Salah Eddin Karimi, Supreet Kaur, Gbenga A Kayode, Peter Njenga Keiyoror, Nauman Khalid, Mohammad Khammarnia, Maseer Khan, Md Nuruzzaman Khan, Khaled Khatab, Mona M Khater, Mahaliqua Nazli Khatib, Maryam Khayamzadeh, Habibolah Khazaie, Abdullah T Khoja, Christian Kielsing, Young-Eun Kim, Yun Jin Kim, Ruth W Kimokoti, Adnan Kisa, Sezer Kisa, Milka Kivimäki, Ali Koolivand, Soewarta Kosen, Ai Koyanagi, Kewal Krishan, Nuworza Kugbey, G Anil Kumar,

Manasi Kumar, Nithin Kumar, Om P Kurmi, Dian Kusuma, Carlo La Vecchia, Ben Lacey, Dharmesh Kumar Lal, Ratilal Laloo, Qing Lan, Iván Landires, Van Charles Lansingh, Anders O Larsson, Savita Lasrado, Zohra S Lassi, Paolo Lauriola, Paul H Lee, Shaun Wen Huey Lee, James Leigh, Matilde Leonardi, Janni Leung, Miriam Levi, Sonia Lewycka, Bingyu Li, Ming-Chieh Li, Shanshan Li, Lee-Ling Lim, Stephen S Lim, Xuefeng Liu, Stefan Lorkowski, Paulo A Lotufo, Raimundas Lunevicius, Ralph Maddison, Phetole Walter Mahasha, Mokhtar Mahdavi Mahdavi, Morteza Mahmoudi, Azeem Majeed, Afshin Maleki, Reza Malekzadeh, Deborah Carvalho Malta, Abdullah A Mamun, Borhan Mansouri, Mohammad Ali Mansournia, Gabriel Martinez, Jose Martinez-Raga, Francisco Rogerlândio Martins-Melo, Amanda J Mason-Jones, Seyede Zahra Masoumi, Manu Raj Mathur, Pallab K Maulik, John J McGrath, Man Mohan Mehndiratta, Fereshteh Mehri, Peter T N Memiah, Walter Mendoza, Ritesh G Menezes, Endalkachew Worku Mengesha, Atte Meretoja, Tuomo J Meretoja, Tomislav Mestrovic, Bartosz Miazgowski, Tomasz Miazgowski, Irmina Maria Michalek, Ted R Miller, GK Mini, Andreea Mirica, Erkin M Mirrakhimov, Hamed Mirzaei, Maryam Mirzaei, Babak Moazeni, Dara K Mohammad, Shadieh Mohammadi, Abdollah Mohammadian-Hafshejani, Noushin Mohammadifard, Reza Mohammadpourhodki, Shafiu Mohammed, Lorenzo Monasta, Ghobad Moradi, Maziar Moradi-Lakeh, Rahmatollah Moradzadeh, Paula Moraga, Shane Douglas Morrison, Abbas Mosapour, Amin Mousavi Khaneghah, Ulrich Otto Mueller, Moses K Muriithi, Christopher J L Murray, Saravanan Muthupandian, Mehdi Naderi, Ahamarshan Jayaraman Nagarajan, Mohsen Naghavi, Mukhammad David Naimzada, Vinay Nangia, Vinod C Nayak, Javad Nazari, Rawlance Ndejo, Ionut Nego, Ruxandra Irina Nego, Henok Biresaw Netsere, Georges Nguefack-Tsague, Diep Ngoc Nguyen, Huong Lan Thi Nguyen, Jing Nie, Dina Nur Anggraini Ningrum, Chukwudi A Nnaji, Shuhei Nomura, Jean Jacques Noubiap, Christoph Nowak, Virginia Nuñez-Samudio, Felix Akpojene Ogbo, Onome Bright Oghenetega, In-Hwan Oh, Morteza Oladnabi, Andrew T Olagunju, Bolajoko Olubukunola Olusanya, Jacob Olusegun Olusanya, Ahmed Omar Bali, Muktar Omer Omer, Obinna E Onwujekwe, Alberto Ortiz, Adrian Otoi, Nikita Ostavnov, Stanislav S Ostavnov, Simon Øverland, Mayowa O Owolabi, Mahesh P A, Jagadish Rao Padubidri, Keyvan Pakshir, Raffaele Palladino, Adrian Pana, Songhomitra Panda-Jonas, Anamika Pandey, Carlo Irwin Able Pano, Eun-Kee Park, Scott B Patten, Amy E Peden, Veincent Christian Filipino Pepito, Emmanuel K Peprah, Jeevan Pereira, Konrad Pesudovs, Hai Quang Pham, Michael R Phillips, Michael A Piradov, Meghdad Pirsaeheb, Maarten J Postma, Faheem Hyder Pottoo, Hadi Pourjafar, Akram Pourshams, Sergio I Prada, Elisabetta Pupillo, Zahiruddin Quazi Syed, Mohammad Hasan Rabiee, Navid Rabiee, Amir Radfar, Ata Rafiee, Alberto Raggi, Fakher Rahim, Vafa Rahimi-Movaghar, Mohammad Hifz Ur Rahman, Muhammad Aziz Rahman, Kiana Ramezanzadeh, Chhabi Lal Ranabhat, Sowmya J Rao, Wahid Rashedi, Prateek Rastogi, Priya Rathi, David Laith Rawaf, Salman Rawaf, Lal Rawal, Reza Rawassizadeh, Andre M N Renzaho, Negar Rezaei, Nima Rezaei, Mohammad Sadegh Rezaei, Seyed Mohammad Riahi, Jennifer Rickard, Leonardo Roever, Luca Ronfani, Gregory A Roth, Enrico Rubagotti, Susan Fred Rumisha, Godfrey M Rwegerera, Siamak Sabour, Perminder S Sachdev, Basema Sadiq, Ehsan Sadeghi, Sahar Saedi Moghaddam, Rajesh Sagar, Amirhossein Sahebkar, Mohammad Ali Sahraian, S Mohammad Sajadi, Marwa Rashad Salem, Hamideh Salimzadeh, Abdallah M Samy, Juan Sanabria, Milena M Santric-Milicevic, Sivan Yegnanarayana Iyer Saraswathy, Nizal Sarrafzadegan, Arash Sarveazad, Thirunavukkarasu Sathish, Davide Sattin, Deepak Saxena, Sonia Saxena, Silvia Schiavolin, David C Schwebel, Falk Schwendicke, Subramanian Senthilkumar, Sadaf G Sepanlou, Feng Sha, Omid Shafaat, Saeed Shahabi, Amira A Shaheen, Masood Ali Shaikh, Saeed Shakiba, MohammadBagher Shamsi, Mohammed Shannawaz, Kiomars Sharafi, Aziz Sheikh, Sara Sheikhbahaei, B Suresh Kumar Shetty, Peilin Shi, Mika Shigematsu, Jae Il Shin, Rahman Shiri, Kerem Shuval, Soraya Siabani, Inga Dora Sigfusdottir, Rannveig Sigurvinssdottir, Diego Augusto Santos Silva, João Pedro Silva, Biagio Simonetti,

Jasvinder A Singh, Virendra Singh, Abiy H Sinke, Valentin Yurievich Skryabin, Helen Slater, Emma U R Smith, Mohammad Reza Sobhiyeh, Eugene Sobngwi, Amin Soheili, Oluwaseyi Dolapo Somefun, Muluken Bekele Sorrie, Ireneous N Soyiri, Chandrashekhar T Sreeramareddy, Dan J Stein, Mark A Stokes, Agus Sudaryanto, Iyad Sultan, Rafael Tabarés-Seisdedos, Takahiro Tabuchi, Santosh Kumar Tadakamadla, Amir Taherkhani, Animum Tagele Tamiru, Md Ismail Tareque, Kavumpurathu Raman Thankappan, Rekha Thapar, Nihal Thomas, Mariya Vladimirovna Titova, Marcello Tonelli, Marcos Roberto Tovani-Palone, Bach Xuan Tran, Ravensara S Travillian, Alexander C Tsai, Aristidis Tsatsakis, Lorraine Tudor Car, Riaz Uddin, Brigid Unim, Bhaskaran Unnikrishnan, Era Upadhyay, Marco Vacante, Sahel Valadan Tahbaz, Pascual R Valdez, Santosh Varughese, Tommi Juhani Vasankari, Narayanaswamy Venketasubramanian, Paul J Villeneuve, Francesco S Violante, Vasily Vlassov, Theo Vos, Giang Thu Vu, Yasir Waheed, Richard G Wamai, Yafeng Wang, Yanzhong Wang, Yuan-Pang Wang, Ronny Westerman, Nuwan Darshana Wickramasinghe, Ai-Min Wu, Chenkai Wu, Seyed Hossein Yahyazadeh Jabbari, Kazumasa Yamagishi, Yuichiro Yano, Sanni Yaya, Wahid Yazdi-Feyzabadi, Yordanos Gizachew Yeshitila, Paul Yip, Naohiro Yonemoto, Seok-Jun Yoon, Mustafa Z Younis, Taraneh Yousefinezhadi, Chuanhua Yu, Yong Yu, Deniz Yuce, Syed Saoud Zaidi, Sojib Bin Zaman, Mohammad Zamani, Maryam Zamanian, Hadi Zarafshan, Ahmad Zarei, Mikhail Sergeevich Zastrozhin, Yunquan Zhang, Zhi-Jiang Zhang, Xiu-Ju George Zhao, Cong Zhu, George C Patton*, Russell M Viner*
*Joint senior authors.

Affiliations

UCL Great Ormond Street Institute of Child Health (J L Ward MSc, Prof R M Viner PhD), Department of Health Informatics (S Chung PhD), Department of Epidemiology and Public Health (Prof M Kivimäki PhD), Division of Psychology and Language Sciences (M Kumar PhD), University College London, London, UK; Global Adolescent Health Group (P S Azzopardi PhD), Burnet Institute, Melbourne, VIC, Australia; Wardliparingga Aboriginal Research Unit (P S Azzopardi PhD), South Australian Health and Medical Research Institute, Adelaide, SA, Australia; Centre for Adolescent Health (K L Francis MBIostat, R Borschmann PhD), Population Health Theme (Prof G C Patton MD), Murdoch Children's Research Institute, Melbourne, VIC, Australia; Department of Population and Family Health (Prof J S Santelli MD), Department of Psychiatry (Prof M R Phillips MD), Department of Health and Behavior Studies (Prof I D Sigfusdottir PhD), Columbia University, New York, NY, USA; Department of Pediatrics (Prof J S Santelli MD), Columbia University Medical Center, New York, NY, USA; Centre for Fertility and Health (Prof V Skirbekk PhD), Norwegian Institute of Public Health, Oslo, Norway; Department of Paediatrics (Prof S M Sawyer MD), University of Melbourne, Parkville, VIC, Australia; Centre for Adolescent Health (Prof S M Sawyer MD), Murdoch Children's Research Institute, Parkville, VIC, Australia; Department of Anaesthesiology & Pain Medicine (N J Kassebaum MD), Institute for Health Metrics and Evaluation (N J Kassebaum MD, Prof A H Mokdad PhD, Prof S I Hay FMedSci, X Dai PhD, Prof L Dandona MD, Prof R Dandona PhD, Prof S D Dharmaratne MD, Prof V L Feigin PhD, Prof S S Lim PhD, Prof C J L Murray DPhil, Prof M Naghavi MD, G A Roth MD, R S Travillian PhD, Prof T Vos PhD), Department of Health Metrics Sciences, School of Medicine (N J Kassebaum MD, Prof A H Mokdad PhD, Prof S I Hay FMedSci, Prof R Dandona PhD, Prof S D Dharmaratne MD, Prof S S Lim PhD, Prof C J L Murray DPhil, Prof M Naghavi MD, G A Roth MD, Prof T Vos PhD), Division of Cardiology (G A Roth MD), University of Washington, Seattle, WA, USA; Department of Neurology (Prof F Abd-Allah MD, S I El-Jaafary MD, A Hassan MD), Department of Endemic Medicine and Hepatogastroenterology (A Elsharkawy MD), Department of Medical Parasitology (M M Khater MD), Cairo University, Cairo, Egypt; Department of Parasitology and Mycology (A Abdoli PhD), Jahrom University of Medical Sciences, Jahrom, Iran; The Institute of Pharmaceutical Sciences (TIIPS) (Prof M Abdollahi PhD), School of Pharmacy (Prof M Abdollahi PhD), Research Center for Immunodeficiencies (H Abolhassani PhD, Prof N Rezaei PhD), Department of Pharmacology (K Afshari MD), Non-communicable

Diseases Research Center (A Aminorroaya MD, Prof F Farzadfar DSc, N Rezaei PhD, S Saeedi Moghaddam MSc), Tehran Heart Center (A Aminorroaya MD), Multiple Sclerosis Research Center (S Eskandari PhD, Prof M Sahraian MD), Endocrinology and Metabolism Research Center (S Esteghamati MD, N Rezaei PhD), School of Medicine (N Hafezi-Nejad MD), School of Nursing and Midwifery (R Heidari-Soureshjani MSc), Department of Epidemiology and Biostatistics (Prof M Hosseini PhD, M Mansourmia PhD), Pediatric Chronic Kidney Disease Research Center (Prof M Hosseini PhD), Department of Environmental Health Engineering (Prof A Maleki PhD), Digestive Diseases Research Institute (Prof R Malekzadeh MD, Prof A Pourshams MD, H Salimzadeh PhD, S G Sepanlou MD), Metabolomics and Genomics Research Center (F Rahim PhD), Sina Trauma and Surgery Research Center (Prof V Rahimi-Movaghar MD), Faculty of Medicine (S Shakiba MD), Psychiatry and Psychology Research Center (H Zarafshan PhD), Tehran University of Medical Sciences, Tehran, Iran; Department of Orthopaedic Surgery (A Abedi MD), Keck School of Medicine (C P Doshi BA), University of Southern California, Los Angeles, CA, USA; Department of Laboratory Medicine (H Abolhassani PhD), Karolinska University Hospital, Huddinge, Sweden; Department of Pediatric Dentistry (Prof L G Abreu PhD), Department of Maternal and Child Nursing and Public Health (Prof D C Malta PhD), Federal University of Minas Gerais, Belo Horizonte, Brazil; Department of Research (M R M Abrigo PhD), Philippine Institute for Development Studies, Quezon City, Philippines; Department of Clinical Sciences (E Abu-Gharbieh PhD), Sharjah Institute for Medical Research (B Saddik PhD), University of Sharjah, Sharjah, United Arab Emirates; Harvard Medical School (A I Abushouk MD), TH Chan School of Public Health (Prof T W Bärrnighausen MD), Center for Primary Care (S Basu PhD), Department of Global Health and Social Medicine (A W Bell MSW), Division of Cardiology (I Y Elgendy MD), Division of General Internal Medicine (Prof A Sheikh MD), Harvard University, Boston, MA, USA; Department of Medicine (A I Abushouk MD), Department of Entomology (A M Samy PhD), Ain Shams University, Cairo, Egypt; College of Medicine (O M Adebayo MD), Department of Community Medicine (O S Ilesanmi PhD), Department of Medicine (Prof M O Owolabi DrM), University College Hospital, Ibadan, Ibadan, Nigeria; Department of Population Medicine (V Adekanmbi PhD), Cardiff University, Cardiff, UK; School of Health (D Adham PhD), Ardabil University of Medical Science, Ardabil, Iran; Terasaki Institute for Biomedical Innovation, Los Angeles, CA, USA (S M Advani PhD); School of Medicine (S M Advani PhD), Georgetown University, Washington DC, DC, USA; Department of Dermatology (K Afshari MD), University of Massachusetts Medical School, Worcester, MA, USA; Institute of Genomics and Integrative Biology (Prof A Agrawal PhD), Council of Scientific & Industrial Research, Delhi, India; Section of General Internal Medicine (Prof A Agrawal PhD), Baylor College of Medicine, Houston, TX, USA; Department of Epidemiology and Health Statistics (T Ahmad MS), Southeast University, Nanjing, China; Lincoln Medical School (K Ahmadi PhD), Universities of Nottingham & Lincoln, Lincoln, UK; Economics and Rural Development Department (A E Ahmed PhD), University of Gezira, Wad Madani, Sudan; Faculty of Medicine and Public Health (B Aji DrPH), Jenderal Soedirman University, Purwokerto, Indonesia; School of Public Health and Community Medicine (B Akombi-Inyang PhD, Prof R Q Ivers PhD, A E Peden PhD), Transport and Road Safety (TARS) Research Centre (R Biswas MSc, S Boufous PhD), School of Medicine (P K Maulik PhD), School of Optometry and Vision Science (Prof K Pesudovs PhD), University of New South Wales, Sydney, NSW, Australia; Mayo Evidence-based Practice Center (F Alahdab MSc), Mayo Clinic Foundation for Medical Education and Research, Rochester, MN, USA; John T Milliken Department of Internal Medicine (Z Al-Aly MD), Washington University in St Louis, St Louis, MO, USA; Clinical Epidemiology Center (Z Al-Aly MD), Department of Veterans Affairs, St Louis, MO, USA; Murdoch Business School (K Alam PhD), Murdoch University, Perth, WA, Australia; Health Information Management and Technology Department (T M Alanzi PhD), Environmental Health Department (S M A Dahlawi PhD), Forensic Medicine Division (Prof R G Menezes MD), Department of Pharmacology (F H Pottoo PhD), Imam Abdulrahman Bin Faisal University, Dammam, Saudi Arabia (F M Alanezi PhD); Center for Health Systems Research (J E Alcalde-Rabanal PhD, L Avila-Burgos ScD.), Health and Nutrition Research Center (I R Campos-Nonato PhD), National Institute of Public Health, Cuernavaca, Mexico (J Campuzano Rincon PhD); College of Medicine and Health Science (B W Alemu MSc), Department of Public Health (M B Sorrie MPH), Department of Nursing (Y G Yeshitla MSc), Arba Minch University, Arba Minch, Ethiopia; Department of Midwifery (B W Alemu MSc), Arba Minch University, Injibara, Ethiopia; Faculty of Health Sciences - Health Management and Policy (Prof S Al-Hajj PhD), American University of Beirut, Beirut, Lebanon; British Columbia Injury Research Prevention Unit (Prof S Al-Hajj PhD), British Columbia Children's Hospital Research Institute, Vancouver, BC, Canada; Institute of Health Research (R K Alhassan PhD), University of Health and Allied Sciences, Ho, Ghana; Department of Information Systems (S Ali PhD), Sultan Qaboos University, Muscat, Oman; Directorate for Social Statistics and Population Census (G Alicandro PhD), Italian National Institute of Statistics, Rome, Italy; Social Determinants of Health Research Center (M Aljanzadeh PhD), Institute for Prevention of Non-communicable Diseases (R Kalhor PhD), Health Services Management Department (R Kalhor PhD), Qazvin University of Medical Sciences, Qazvin, Iran; Department of Health Policy and Management (Prof S M Aljunid PhD), Kuwait University, Safat, Kuwait; International Centre for Casemix and Clinical Coding (Prof S M Aljunid PhD), National University of Malaysia, Bandar Tun Razak, Malaysia; Department of Epidemiology (Almasi-Hashiani PhD, R Moradzadeh PhD, M Zamarian PhD), Department of Health Services Management (S Amini PhD), Occupational Health Department (F Ghamari PhD), Department of Environmental Health Engineering (A Koolivand PhD), Department of Pediatrics (J Nazari MD), Arak University of Medical Sciences, Arak, Iran; Physiotherapy Department (Prof N A Almasri PhD), The University of Jordan, Amman, Jordan; Medical Research Center (H M Al-Mekhlafi PhD), Epidemiology Department (M Khan MD), Jazan University, Jazan, Saudi Arabia (Prof N Bedi MD); Department of Parasitology (H M Al-Mekhlafi PhD), Sana'a University, Sana'a, Yemen; Research Program of Epidemiology and Public Health (J Alonso MD), Pompeu Fabra University, Barcelona, Spain; Department of Experimental and Health Sciences (J Alonso MD), Biomedical Research Networking Center in Epidemiology and Public Health (CiberESP), Madrid, Spain; Department of Community Medicine (R M Al-Raddadi PhD), King Abdulaziz University, Jeddah, Saudi Arabia; Pediatric Intensive Care Unit (K A Altirkawi MD), King Saud University, Riyadh, Saudi Arabia; Research Group in Hospital Management and Health Policies (Prof N Alvis-Guzman PhD), Universidad de la Costa, Barranquilla, Colombia; Research Group in Health Economics (Prof N Alvis-Guzman PhD), University of Cartagena, Cartagena, Colombia; School of Medicine (A Amare PhD), Robinson Research Institute (Z S Lassi PhD), Centre for Heart Rhythm Disorders (J Noubiap MD), University of Adelaide, Adelaide, SA, Australia; College of Medicine and Health Science (A Amare PhD), Department of Reproductive Health and Population Studies (E W Mengesha MPH), Bahir Dar University, Bahir Dar, Ethiopia; School of Medicine and Public Health (A L Amit BS), Ateneo De Manila University, Manila, Philippines; College of Medicine (A L Amit BS), Department of Health Policy and Administration (C T Antonio MD), Department of Clinical Epidemiology (C A Panelo MD), University of the Philippines Manila, Manila, Philippines; Maternal and Child Wellbeing (D A Amugsi PhD), African Population and Health Research Center, Nairobi, Kenya; Pharmacy Department (Prof R Ancuceanu PhD), Cardiology Department (C Andrei PhD), Urology Department (T Constantin PhD), Internal Medicine Department (M Hostiu PhD), Department of Legal Medicine and Bioethics (S Hostiu PhD), Department of General Surgery (I Negoi PhD), Department of Anatomy and Embryology (R I Negoi PhD), Carol Davila University of Medicine and Pharmacy, Bucharest, Romania; Centre for Sensorimotor Performance (D Anderlini MD), The University of Queensland, Brisbane, Australia; Neurology Department (D Anderlini MD), Royal Brisbane and Women's Hospital, Brisbane, QLD, Australia; Department of Medicine (S Androudi PhD), University of Thessaly, Volos, Greece; Research Center for Evidence Based Medicine (F Ansari PhD), Department of Health Policy and Management (L Doshmangir PhD), Department of Medical Surgical Nursing (M Ghafourifard PhD), School of Nursing and Midwifery (H Hassankhani PhD), School of Management and Medical Informatics (L R Kalankesh PhD), Social Determinants of Health Research Center

(S Karimi PhD), Tabriz University of Medical Sciences, Tabriz, Iran (H Haririan PhD); Razi Vaccine and Serum Research Institute (F Ansari PhD), Agricultural Research, Education, and Extension Organization (AREEO), Tehran, Iran; Medical Students Research Committee (I Ansari MD), Shahed University, Tehran, Iran; Department of Applied Social Sciences (C T Antonio MD), Hong Kong Polytechnic University, Hong Kong, China; Department of Parasitology (D Anvari PhD), Pediatric Infectious Diseases Research Center (Prof M Rezai MD), Mazandaran University of Medical Sciences, Sari, Iran; Department of Parasitology (D Anvari PhD), Iranshahr University of Medical Sciences, Iranshahr, Iran; Department of Pathology (R Anwer PhD), Department of Public Health (A T Khoja MD), Imam Mohammad Ibn Saud Islamic University, Riyadh, Saudi Arabia; Department of Sociology and Social Work (S Appiah PhD), Kwame Nkrumah University of Science and Technology, Kumasi, Ghana; Center for International Health (S Appiah PhD), Ludwig Maximilians University, Munich, Germany; Health Management and Economics Research Center (J Arabloo PhD, S Azari PhD, A Ghashghaeae BSc, M Hosseinzadeh PhD), Preventive Medicine and Public Health Research Center (M Asadi-Aliabadi MSc, B Eshrati PhD, M Moradi-Lakeh MD), Department of Medical Laboratory Sciences (F Dorostkar PhD), Student Research Committee (A Ghashghaeae BSc), Minimally Invasive Surgery Research Center (A Kabir MD), Tehran Institute of Psychiatry (V Rashedi PhD), Colorectal Research Center (A Sarveazad PhD), Iran University of Medical Sciences, Tehran, Iran; Social Determinants of Health Research Center (M Arab-Zozani PhD), Cardiovascular Diseases Research Center (S Riahi PhD), Birjand University of Medical Sciences, Birjand, Iran; Department of Neurobiology, Care Sciences and Society (Prof J Årnlöv PhD), Department of Medical Epidemiology and Biostatistics (Prof J J Carrero PhD), Department of Global Public Health (K Deuba DrPH), Department of Neurobiology (S Fereshtehnejad PhD), Department of Medicine (D K Mohammad PhD), Karolinska Institute, Stockholm, Sweden; School of Health and Social Studies (Prof J Årnlöv PhD), Dalarna University, Falun, Sweden; Department of Plastic Surgery (M Asaad MD), University of Texas, Houston, TX, USA; Epilepsy Research Center (Prof A A Asadi-Pooya MD), Health Human Resources Research Center (M Bayati PhD), Department of Epidemiology (M Dianatinasab MSc), Non-communicable Disease Research Center (Prof R Malekzadeh MD, S G Sepanlou MD), Department of Medical Mycology and Parasitology (Prof K Pakshir PhD), Health Policy Research Center (S Shahabi PhD), Shiraz University of Medical Sciences, Shiraz, Iran; Neurology Department (Prof A A Asadi-Pooya MD), Thomas Jefferson University, Philadelphia, PA, USA; Faculty of Nursing (M M W Atout PhD), Philadelphia University, Amman, Jordan; School of Business (Prof M Ausloos PhD), Department of Health Sciences (P H Lee PhD), University of Leicester, Leicester, UK; Department of Statistics and Econometrics (Prof M Ausloos PhD, Prof C Herteliu PhD, A Mirica PhD, Prof A Otoiui PhD, A Pana MD), Bucharest University of Economic Studies, Bucharest, Romania; Department of International Development (E K Avenyo PhD), Nuffield Department of Population Health (B Lacey PhD), Centre for Tropical Medicine and Global Health (S Lewycka PhD), The George Institute for Global Health (Prof S Yaya PhD), University of Oxford, Oxford, UK; DST/NRF South African Research Chair in Industrial Development (E K Avenyo PhD), University of Johannesburg, Johannesburg, South Africa; The Judith Lumley Centre (B Ayala Quintanilla PhD), School of Nursing and Midwifery (Prof D Edvardsson PhD, M Rahman PhD), School of Psychology and Public Health (M Islam PhD), La Trobe University, Melbourne, VIC, Australia; School of Public Health (G Ayano MSc, T R Miller PhD), School of Physiotherapy and Exercise Science (Prof H Slater PhD), Curtin University, Perth, WA, Australia; Department of Nursing (Y A Aynalem MSc), Debre Berhan University, Debre Berhan, Ethiopia; Department of Reproductive Health (Z N Azene MPH), School of Nursing (H B Netsere MS), Department of Midwifery (A T Tamiru MSc), University of Gondar, Gondar, Ethiopia; Department of Anesthesiology (M Bakhshaei MD), Hamedan University of Medical Sciences, Hamadan, Iran; Department of Forensic Medicine and Toxicology (S M Bakkannavar MD, Prof V C Nayak MD), Kasturba Medical College, Mangalore (R Holla MD, J Padubidri MD, P Rathi MD), Centre for Bio Cultural Studies (CBiCS) (P Hoogar PhD), Manipal College of Pharmaceutical Sciences (Prof M D Janodia PhD), Manipal Academy of Higher Education, Manipal, India (Prof V Jha MD); Department of Hypertension (Prof M Banach PhD), Medical University of Lodz, Lodz, Poland; Polish Mothers' Memorial Hospital Research Institute, Lodz, Poland (Prof M Banach PhD); Department of Non-communicable Diseases (P C Banik MPhil), Bangladesh University of Health Sciences, Dhaka, Bangladesh; Department of Neurosciences (Prof M A Barboza MD), Costa Rican Department of Social Security, San Jose, Costa Rica; School of Medicine (Prof M A Barboza MD), University of Costa Rica, San Pedro, Costa Rica; School of Psychology (Prof S L Barker-Collo PhD), University of Auckland, Auckland, New Zealand; Heidelberg Institute of Global Health (HIGH) (Prof T W Bärnighausen MD, B Moazen MSc), Department of Ophthalmology (Prof J B Jonas MD, S Panda-Jonas MD), Heidelberg University, Heidelberg, Germany; School of Public Health (S Basu PhD, Prof S Saxena MD), Imperial College Business School (D Kusuma DSc), Department of Primary Care and Public Health (Prof A Majeed MD, R Palladino MD, Prof S Rawaf MD), WHO Collaborating Centre for Public Health Education and Training (D L Rawaf MD), Imperial College London, London, UK; Department of Psychiatry (Prof B T Baune PhD), Institute for Epidemiology and Social Medicine (A Karch MD), University of Münster, Münster, Germany; Department of Psychiatry (Prof B T Baune PhD), Melbourne Medical School, Melbourne, VIC, Australia; School of Public Health (Prof N Bedi MD), Dr D Y Patil University, Mumbai, India; Department of Neuroscience (E Beghi MD, E Pupillo PharmD), Department of Environmental Health Sciences (S Gallus DSc), Mario Negri Institute for Pharmacological Research, Milan, Italy; Department of Public Health (T T Bekuma MPH), Wollega University, Nekemte, Ethiopia; Department of Social Services (A W Bell MSW), Tufts Medical Center, Boston, MA, USA; School of the Environment (Prof M L Bell PhD), Yale University, New Haven, CT, USA; Department of Epidemiology and Psychosocial Research (C Benjet PhD), Ramón de la Fuente Muñiz National Institute of Psychiatry, Mexico City, Mexico; Department of Internal Medicine (I M Bensenor PhD), Department of Psychiatry (Prof J Castaldelli-Maia PhD, Y Wang PhD), University of São Paulo, São Paulo, Brazil; College of Medicine and Health Sciences (A K Berhe MSc), Adigrat University, Adigrat, Ethiopia; Department of Nutrition and Dietetics (K Berhe MPH), Department of Nursing (G G Gebremeskel MSc), Department of Pharmacy (L Gebremeskel MSc), Department of Reproductive Health (A Gebreslassie MPH), Department of Epidemiology (H A Gesesew PhD), Department of Medical Microbiology and Immunology (S Muthupandian PhD), Mekelle University, Mekelle, Ethiopia; Department of Medicine (A E Berman MD), Medical College of Georgia at Augusta University, Augusta, GA, USA; Department of Social and Clinical Pharmacy (A S Bhagavathula PharmD), Charles University, Hradec Kralova, Czech Republic; Institute of Public Health (A S Bhagavathula PharmD, Prof M Grivna PhD), United Arab Emirates University, Al Ain, United Arab Emirates; Department of Anatomy (Prof N Bhardwaj MD), Government Medical College Pali, Pali, India; Department of Community Medicine and Family Medicine (P Bhardwaj MD), School of Public Health (P Bhardwaj MD), Department of Pharmacology (J Charan MD), Department of Forensic Medicine and Toxicology (T Kanchan MD), All India Institute of Medical Sciences, Jodhpur, India; Department of Statistical and Computational Genomics (K Bhattacharyya MSc), National Institute of Biomedical Genomics, Kalyani, India; Department of Statistics (K Bhattacharyya MSc), University of Calcutta, Kolkata, India; Department of Global Health (S Bhattarai MD), Global Institute for Interdisciplinary Studies, Kathmandu, Nepal; Centre for Global Child Health (Prof Z A Bhutta PhD), Department of Medicine (V Chattu MD), University of Toronto, Toronto, ON, Canada; Centre of Excellence in Women & Child Health (Prof Z A Bhutta PhD), Aga Khan University, Karachi, Pakistan; Social Determinants of Health Research Center (A Bijani PhD, M A Jahani PhD), Department of Clinical Biochemistry (A Mosapour PhD), Student Research Committee (M Zamani MD), Babol University of Medical Sciences, Babol, Iran; Mario Negri Institute for Pharmacological Research, Ranica, Italy (B Bikbov MD); Department of General Surgery and Medical-Surgical Specialties (Prof A Biondi PhD, M Vacante PhD), University of Catania, Catania, Italy; Planning, Monitoring and Evaluation Directorate (T T M Birhanu MPH), Amhara Public Health Institute, Bahir Dar, Ethiopia; Department of Veterinary Medicine (S Bohlouli PhD), Islamic Azad University,

Kermanshah, Iran; Department of Biomedical Sciences (S Bolla PhD), Nazarbayev University, Nur-Sultan City, Kazakhstan; Department of Internal Medicine (A Bolor MD), Department of Community Medicine (N Kumar MD, R Thapar MD), Department of Forensic Medicine and Toxicology (Prof P Rastogi MD, Prof B K Shetty MD), Kasturba Medical College (Prof B Unnikrishnan MD), Manipal Academy of Higher Education, Mangalore, India; Justice Health Unit (R Borschmann PhD), School of Health Sciences (A Meretoja MD), Department of Pediatrics (Prof G C Patton MD), University of Melbourne, Melbourne, VIC, Australia; University of Genoa, Genoa, Italy (N L Bragazzi PhD); Department of Epidemiology (D Braithwaite PhD), University of Florida, Gainesville, FL, USA; Cancer Population Sciences Program (D Braithwaite PhD), University of Florida Health Cancer Center, Gainesville, FL, USA; Psychiatry and Behavioral Health Department (Prof N J K Breitborde PhD), Department of Psychology (Prof N J K Breitborde PhD), James Cancer Hospital (J L Fisher PhD), Ohio State University, Columbus, OH, USA; Division of Clinical Epidemiology and Aging Research (Prof H Brenner MD), German Cancer Research Center, Heidelberg, Germany; Neuroscience Unit (G B Britton PhD), Institute for Scientific Research and High Technology Services, Panama City, Panama; Gorgas Memorial Institute for Health Studies, Panama City, Panama (G B Britton PhD); Research School of Population Health (R A Burns PhD), Australian National University, Canberra, ACT, Australia; Department of Community Medicine (Prof S Burugina Nagaraja MD), Employee State Insurance Post Graduate Institute of Medical Sciences and Research, Bangalore, India; School of Public Health and Health Systems (Z A Butt PhD), University of Waterloo, Waterloo, ON, Canada; Al Shifa School of Public Health (Z A Butt PhD), Al Shifa Trust Eye Hospital, Rawalpindi, Pakistan; Institute of Microengineering (F Caetano dos Santos PhD), Federal Polytechnic School of Lausanne, Lausanne, Switzerland; Internal Medicine Department (Prof L A Cámara MD), Hospital Italiano de Buenos Aires (Italian Hospital of Buenos Aires), Buenos Aires, Argentina; Board of Directors (Prof L A Cámara MD), Argentine Society of Medicine, Buenos Aires, Argentina (Prof P R Valdez MEd); School of Medicine (J Campuzano Rincon PhD), University of the Valley of Cuernavaca, Cuernavaca, Mexico; Department of Health Care (Prof R Cárdenas DSc), Metropolitan Autonomous University, Mexico City, Mexico; Oncological Network, Prevention and Research Institute (G Gorini MD), Institute for Cancer Research, Prevention and Clinical Network, Florence, Italy (G Carreras PhD); Research Unit on Applied Molecular Biosciences (UCIBIO) (Prof F Carvalho PhD, V M Costa PhD, J P Silva PhD), Laboratory of Toxicology (Prof D Dias da Silva PhD), Associated Laboratory for Green Chemistry (LAQV) (Prof E Fernandes PhD), University of Porto, Porto, Portugal; Colombian National Health Observatory (C A Castañeda-Orjuela MD), National Institute of Health, Bogota, Colombia; Epidemiology and Public Health Evaluation Group (C A Castañeda-Orjuela MD), Department of Public Health (Prof F P De la Hoz PhD), National University of Colombia, Bogota, Colombia; Department of Medicine (G Castelpietra PhD), University of Udine, Udine, Italy; Department of Mental Health (G Castelpietra PhD), Healthcare Agency "Friuli Occidentale", Pordenone, Italy; National School of Public Health (F Catalá-López PhD), Institute of Health Carlos III, Madrid, Spain; Clinical Epidemiology Program (F Catalá-López PhD), Ottawa Hospital Research Institute, Ottawa, ON, Canada; Mary MacKillop Institute for Health Research (Prof E Cerin PhD), Australian Catholic University, Melbourne, VIC, Australia; School of Public Health (Prof E Cerin PhD), Department of Urban Planning and Design (H Ho PhD), Centre for Suicide Research and Prevention (Prof P Yip PhD), Department of Social Work and Social Administration (Prof P Yip PhD), University of Hong Kong, Hong Kong, China; Institute of Applied Health Research (J S Chandan MFPH), University of Birmingham, Birmingham, UK; Institute of Population Health Sciences (H Chang DrPH), National Health Research Institutes, Zhunan Township, Taiwan; College of Medicine (J Chang PhD), National Taiwan University, Taipei, Taiwan; Department of Nursing (J Chang PhD), National Taiwan University Hospital, Taipei, Taiwan; Global Institute of Public Health (GIPH), Thiruvananthapuram, India (V Chattu MD); Research Department (S Chaturvedi PhD), Dr D Y Patil University, Pune, India; Biomedical Informatics (J J Choi PhD), Seoul National University Hospital, Seoul, South Korea; James P Grant School of Public Health (M Chowdhury MPH), BRAC University, Dhaka, Bangladesh; Department of Epidemiology and Biostatistics (M Chowdhury MPH), University of South Carolina, Columbia, SC, USA; Department of Pulmonary Medicine (Prof D J Christopher MD), Department of Endocrinology, Diabetes and Metabolism (Prof N Thomas PhD), Department of Nephrology (Prof S Varughese FRCP), Christian Medical College and Hospital (CMC), Vellore, India; Center for Biomedicine and Community Health (D Chu PhD), VNU-International School, Hanoi, Vietnam; Department of Otolaryngology (M T Chung MD), Wayne State University, Detroit, MI, USA; Health Data Research UK, London, UK (S Chung PhD); School of Public Health and Preventive Medicine (Prof F M Cicuttini PhD, S Li PhD), Department of Epidemiology and Preventive Medicine (Prof Y Guo PhD), The School of Clinical Sciences at Monash Health (S Zaman MPH), Monash University, Melbourne, VIC, Australia; Department of Cardiology (H Dai MD), Central South University, Changsha, China; Department of Mathematics and Statistics (H Dai MD), York University, Toronto, ON, Canada; IRCCS Istituto Ortopedico Galeazzi (Galeazzi Orthopedic Institute IRCCS) (G Damiani MD), Department of Clinical Sciences and Community Health (Prof C La Vecchia MD), University of Milan, Milan, Italy; Department of Dermatology (G Damiani MD), Department of Quantitative Health Science (X Liu PhD), Department of Nutrition and Preventive Medicine (Prof J Sanabria MD), Case Western Reserve University, Cleveland, OH, USA; Health Policy Research (M R Mathur PhD), Department of Research (A Pandey PhD), Public Health Foundation of India, Gurugram, India (Prof L Dandona MD, Prof R Dandona PhD, G Kumar PhD, D K Lal MD); Indian Council of Medical Research, New Delhi, India (Prof L Dandona MD); Department of Pathology (P Daneshpajouhnejad MD), Department of Radiology and Radiological Science (N Hafezi-Nejad MD, O Shafaat MD, S Sheikhabahaei MD), Department of Radiology and Radiological Sciences (A Haj-Mirzaian MD), Department of Health Policy and Management (A T Khoja MD), Johns Hopkins University, Baltimore, MD, USA; Department of Pathology (P Daneshpajouhnejad MD), Isfahan Cardiovascular Research Institute (N Mohammadifard PhD, Prof N Sarrafzadegan MD), Department of Radiology and Interventional Neuroradiology (O Shafaat MD), Isfahan University of Medical Sciences, Isfahan, Iran; Department of Information Technology (A M Darwesh PhD), University of Human Development, Sulaymaniyah, Iraq; Department of Population and Development (C A Dávila-Cervantes PhD), Latin American Faculty of Social Sciences Mexico, Mexico City, Mexico; Health Research Institute (K Davletov PhD), Al Farabi Kazakh National University, Almaty, Kazakhstan; Australian Institute for Suicide Research and Prevention (Prof D De Leo DSc), Griffith University, Mount Gravatt, QLD, Australia; St Paul's Eye Unit (N Dervenis MD), Royal Liverpool University Hospital, Liverpool, UK; Department of Ophthalmology (N Dervenis MD), Aristotle University of Thessaloniki, Thessaloniki, Greece; Division of Cardiology (R Desai MBBS), Atlanta Veterans Affairs Medical Center, Decatur, GA, USA; School of Nursing and Midwifery (A Desalew MSc), Haramaya University, Harar, Ethiopia; National Centre for AIDS and STD Control (K Deuba DrPH), Save the Children, Kathmandu, Nepal; Department of Community Medicine (Prof S D Dharmaratne MD), University of Peradeniya, Peradeniya, Sri Lanka; Department of Microbiology (G P Dhungana MSc), Far Western University, Mahendranagar, Nepal; Department of Epidemiology and Biostatistics (M Dianatinasab MSc), Shahrood University of Medical Sciences, Shahrood, Iran; Center of Complexity Sciences (Prof D Diaz PhD), National Autonomous University of Mexico, Mexico City, Mexico; Faculty of Veterinary Medicine and Zootechnics (Prof D Diaz PhD), Autonomous University of Sinaloa, Culiacán Rosales, Mexico; Department of Community Medicine and Public Health (A Didarloo PhD), Urmia University of Medical Science, Urmia, Iran; Development of Research and Technology Center (S Djalalinia PhD), Ministry of Health and Medical Education, Tehran, Iran; School of Medicine (Prof K E Doyle PhD), Translational Health Research Institute (F A Ogbo PhD), Western Sydney University, Sydney, NSW, Australia; School of Health and Biomedical Sciences (Prof K E Doyle PhD), Royal Melbourne Institute of Technology University, Melbourne, VIC, Australia; School of Medicine (Prof A R Duraes PhD), Federal University of Bahia, Salvador, Brazil; Department of Internal Medicine (Prof A R Duraes PhD), Escola Bahiana de Medicina e Saúde Pública (Bahiana School of Medicine and Public Health), Salvador, Brazil;

Department of Epidemiology (M Ebrahimi Kalan MSc), Florida International University, Miami, FL, USA; Department of Cardiovascular Medicine (S Ebtelah PhD), Mayo Clinic, Rochester, MN, USA; Department of Nursing (Prof D Edvardsson PhD), Umeå University, Umeå, Sweden; Pediatric Dentistry and Dental Public Health Department (Prof M El Tantawi PhD), Alexandria University, Alexandria, Egypt; Division of Cardiology (I Y Elgendy MD), Department of Psychiatry (A C Tsai MD), Massachusetts General Hospital, Boston, MA, USA; Department of Physiology (S Esmailnejad PhD), Department of Clinical Biochemistry (A Mosapour PhD), Tarbiat Modares University, Tehran, Iran; Tehran Medical Sciences Branch (S Esmailnejad PhD), Department of Microbiology (S Valadan Tahbaz PhD), Islamic Azad University, Tehran, Iran; Department of Public Health (F Esmailzadeh PhD), Department of Nutrition and Food Sciences (H Pourjafar PhD), Maragheh University of Medical Sciences, Maragheh, Iran; Department of Psychology (Prof A Faro PhD), Federal University of Sergipe, São Cristóvão, Brazil; Research Center for Environmental Determinants of Health (N Fattahi PhD, Prof M Pirsahab PhD, Prof E Sadeghi PhD, K Sharafi PhD), Infectious Disease Research Center (Prof K Ghadiri MD), Pediatric Department (Prof K Ghadiri MD), Medical Biology Research Center (F Heydarpour PhD), Health Institute (A Jalali PhD), Substance Abuse Prevention Research Center (A Jalali PhD, B Mansouri PhD), Department of Psychiatry (Prof H Khazaie MD), Department of Rehabilitation and Sports Medicine (M Mirzaei MSc), Clinical Research Development Center (M Naderi PhD), Department of Sports Medicine and Rehabilitation (M Shamsi PhD), Department of Health Education and Health Promotion (S Siabani PhD), Department of Vascular and Endovascular Surgery (M Sobhiyeh MD), Kermanshah University of Medical Sciences, Kermanshah, Iran; National Institute for Stroke and Applied Neurosciences (Prof V L Feigin PhD, K M Jones PhD), Auckland University of Technology, Auckland, New Zealand; Third Department of Neurology (E V Gnedovskaya PhD), Research Center of Neurology, Moscow, Russia (Prof V L Feigin PhD, Prof M A Piradov DSc); School of Nursing (T Y Ferede MSc), Hawassa University, Hawassa, Ethiopia; Division of Neurology (S Fereshtehnejad PhD), School of International Development and Global Studies (Prof S Yaya PhD), University of Ottawa, Ottawa, ON, Canada; Research Center on Public Health (P Ferrara MD), University of Milan Bicocca, Monza, Italy; Psychiatry Department (I Filip MD), Kaiser Permanente, Fontana, CA, USA; School of Health Sciences (I Filip MD), AT Still University, Mesa, AZ, USA; Institute of Gerontological Health Services and Nursing Research (F Fischer PhD), Ravensburg-Weingarten University of Applied Sciences, Weingarten, Germany; Institute of Gerontology (N A Foigt PhD), National Academy of Medical Sciences of Ukraine, Kyiv, Ukraine; Department of Child Dental Health (Prof M O Folayan FWACS), Obafemi Awolowo University, Ile-Ife, Nigeria; Department of Cell Biology and Biotechnology (A A Fomenkov PhD), KA Timiryazev Institute of Plant Physiology, Moscow, Russia; Department of Medical Parasitology (M Foroutan PhD), Abadan Faculty of Medical Sciences, Abadan, Iran; Department of Dermatology (T Fukumoto PhD), Kobe University, Kobe, Japan; Department of Cardiovascular Medicine (M M Gad MD), Lerner Research Institute (X Liu PhD), Cleveland Clinic, Cleveland, OH, USA; Gillings School of Global Public Health (M M Gad MD), University of North Carolina Chapel Hill, Chapel Hill, NC, USA; Department of Community Medicine (Prof A M Gaidhane MD, Prof Z Quazi Syed PhD, Prof D Saxena PhD), Global Evidence Synthesis Initiative (Prof M Khatib PhD), Datta Meghe Institute of Medical Sciences, Wardha, India; International Trachoma Initiative (T Gebre PhD), Task Force for Global Health, Decatur, GA, USA; Department of Nursing and Midwifery (K B Gebremedhin MSc), Addis Ababa University, Addis Ababa, Ethiopia; Department of Nursing (G G Gebremeskel MSc, T G Haile MSc), School of Pharmacy (L Gebremeskel MSc), Aksum University, Aksum, Ethiopia; College of Medicine and Public Health (H A Gesesew PhD), Flinders University, Adelaide, SA, Australia; Faculty of Allied Health Sciences (Prof S Gilani PhD), University Institute of Public Health (A Hanif PhD), The University of Lahore, Lahore, Pakistan; Afro-Asian Institute, Lahore, Pakistan (Prof S Gilani PhD); School of Public Health and Community Medicine (M A Godinho MBBS), School of Psychiatry (Prof P S Sachdev MD), University of New South Wales, Kensington, NSW, Australia; Health Systems and Policy Research (M Golechha PhD), Department of Epidemiology (Prof D Saxena PhD), Indian Institute of Public Health, Gandhinagar, India; Australia India Institute New Generation Network (S Goli PhD), UWA Public Policy Institute (S Goli PhD), Medical School (Prof G J Hankey MD), University of Western Australia, Perth, WA, Australia; Department of Exercise and Health Sciences (P N Gona PhD), University of Massachusetts Boston, Boston, MA, USA; Hudson College of Public Health (S V Gopalani MPH), University of Oklahoma Health Sciences Center, Oklahoma City, OK, USA; Department of Health and Social Affairs (S V Gopalani MPH), Government of the Federated States of Micronesia, Palikir, Federated States of Micronesia; Department of Public Health and Preventive Medicine (Prof M Grivna PhD), Charles University, Prague, Czech Republic; Department of Family and Community Medicine (M I M Gubari PhD), University Of Sulaimani, Sulaimani, Iraq; Department of Microbiology (Prof H C Gugnani PhD), Department of Epidemiology (Prof H C Gugnani PhD), Saint James School of Medicine, The Valley, Anguilla; Institute of Tropical Pathology and Public Health (IPTSP) (R A Guimarães MSc), Federal University of Goiás, Goiânia, Brazil; Department of Epidemiology (Prof Y Guo PhD), Binzhou Medical University, Yantai City, China; Department of Preventive Cardiology (Prof R Gupta MD), Eternal Heart Care Centre & Research Institute, Jaipur, India; Department of Medicine (Prof R Gupta MD), Mahatma Gandhi University Medical Sciences, Jaipur, India; Department of Public Health (J A Haagsma PhD), Erasmus University Medical Center, Rotterdam, Netherlands; Department of Pharmacology (A Haj-Mirzaian MD, K Ramezanzadeh PharmD), Obesity Research Center (A Haj-Mirzaian MD), Department of Epidemiology (S Sabour PhD), Injury Prevention and Safety Promotion Research Center (T Yousefinezhadi PhD), Shahid Beheshti University of Medical Sciences, Tehran, Iran (M Khayamzadeh MD); NYU Shanghai, Shanghai, China (B J Hall PhD); Department of Family and Community Medicine (Prof R R Hamadeh PhD), Arabian Gulf University, Manama, Bahrain; College of Law and Political Science (K Hamagharib Abdullah PhD), Diplomacy and Public Relations Department (A Omar Bali PhD), University of Human Development, Sulaimaniyah, Iraq; School of Health and Environmental Studies (Prof S Hamidi DrPH), Hamdan Bin Mohammed Smart University, Dubai, United Arab Emirates; Department of Public Health (D Handiso MPH), Wachemo University, Hossana, Ethiopia; Department of Neurology (Prof G J Hankey MD), Sir Charles Gairdner Hospital, Perth, WA, Australia; Research Unit (J M Haro MD), University of Barcelona, Sant Boi de Llobregat, Barcelona, Spain; Biomedical Research Networking Center for Mental Health Network (CiberSAM), Barcelona, Spain (J M Haro MD); Department of Zoology and Entomology (A I Hasaballah PhD), Al Azhar University, Cairo, Egypt; Department of Public Health (A Hashi PhD, M O Omer MSc), Jigjiga University, Jigjiga, Ethiopia; Gastrointestinal and Liver Diseases Research Center (S Hassanipour PhD, F Joukar PhD), Caspian Digestive Disease Research Center (S Hassanipour PhD, F Joukar PhD), Guilan University of Medical Sciences, Rasht, Iran; Independent Consultant, Tabriz, Iran (H Hassankhani PhD, S N Irvani MD); Institute of Pharmaceutical Sciences (K Hayat MS), University of Veterinary and Animal Sciences, Lahore, Pakistan; Department of Pharmacy Administration and Clinical Pharmacy (K Hayat MS), Xian Jiaotong University, Xian, China; School of Business (Prof C Herteliu PhD), London South Bank University, London, UK; Department of Pediatrics (M K Hole MD), University of Texas Austin, Austin, TX, USA; Clinical Legal Medicine Department (S Hosticiu PhD), National Institute of Legal Medicine Mina Minovici, Bucharest, Romania; College of Science and Engineering (Prof M Househ PhD), Hamad Bin Khalifa University, Doha, Qatar; Faculty of Medicine of Tunis (Prof M Hsairi MPH), University Tunis El Manar, Tunis, Tunisia; School of Public Health (T M Huda PhD), Sydney Medical School (S Islam PhD), Asbestos Diseases Research Institute (J Leigh MD), University of Sydney, Sydney, NSW, Australia; Maternal and Child Health Division (T M Huda PhD, S Zaman MPH), International Centre for Diarrhoeal Disease Research, Bangladesh, Dhaka, Bangladesh; Department of Public Health and Community Medicine (Prof A Humayun PhD), Shaikh Khalifa Bin Zayed Al-Nahyan Medical College, Lahore, Pakistan; School of Pharmaceutical Sciences (R Hussain PhD), University of Science Malaysia, Penang, Malaysia; Department of Occupational Safety and Health (Prof B Hwang PhD), China Medical University, Taichung, Taiwan; Department of Public Health (Prof I Iavicoli PhD, R Palladino MD), University of Naples Federico II, Naples, Italy; Department of Health

Promotion and Education (S E Ibitoye MPH), Department of Community Medicine (O S Ilesanmi PhD), Department of Obstetrics and Gynecology (O B Oghenetega MSc), Department of Medicine (Prof M O Owolabi DrM), University of Ibadan, Ibadan, Nigeria; Faculty of Medicine (I M Ilic PhD, Prof M M Santric-Milicevic PhD), School of Public Health and Health Management (Prof M M Santric-Milicevic PhD), University of Belgrade, Belgrade, Serbia; Department of Epidemiology (Prof M D Ilic PhD), Department of Global Health, Economics and Policy (Prof M Jakovljevic PhD), University of Kragujevac, Kragujevac, Serbia; Division of Community Health and Family Medicine (L R Inbaraj MD), Bangalore Baptist Hospital, Bangalore, India; Department of Health Science (N Intarut PhD), Mahasarakham University, Maha Sarakham, Thailand; College of Public Health (U Iqbal PhD), Graduate Institute of Biomedical Informatics (D N A Ningrum MPH), Taipei Medical University, Taipei, Taiwan; Institute for Physical Activity and Nutrition (S Islam PhD, Prof R Maddison PhD), Department of Psychology (M A Stokes PhD), Deakin University, Burwood, VIC, Australia; Public Health Department of Social Medicine (Prof H Iso MD), Graduate School of Medicine (Prof K Yamagishi MD), Osaka University, Suita, Japan; Institute of Comparative Economic Studies (Prof M Jakovljevic PhD), Hosei University, Tokyo, Japan; Health Informatic Lab (T Javaheri PhD), Department of Computer Science (R Rawassizadeh PhD), Boston University, Boston, MA, USA; Achutha Menon Centre for Health Science Studies (P Jeemon PhD), Sree Chitra Tirunal Institute for Medical Sciences and Technology, Trivandrum, India; Autism Spectrum Disorders Research Center (E Jenabi PhD), Department of Biostatistics (N Kamyari MSc), Department of Midwifery (S Masoumi PhD), Research Center for Molecular Medicine (A Taherkhani PhD), Hamadan University of Medical Sciences, Hamadan, Iran; Department of Community Medicine (R P Jha MSc), Dr Baba Saheb Ambedkar Medical College & Hospital, Delhi, India; Department of Community Medicine (R P Jha MSc), Banaras Hindu University, Varanasi, India; Research Division (P K Maulik PhD), The George Institute for Global Health, New Delhi, India (Prof V Jha MD); Environmental Research Center (J S Ji DSc), Global Health Research Center (C Wu PhD), Duke Kunshan University, Kunshan, China; Nicholas School of the Environment (J S Ji DSc), Duke Global Health Institute (C Wu PhD), Duke University, Durham, NC, USA; Beijing Institute of Ophthalmology (Prof J B Jonas MD), Beijing Tongren Hospital, Beijing, China; Department of Family Medicine and Public Health (J J Jozwiak PhD), University of Opole, Opole, Poland; Department of Health Register Research and Development (Prof P B Juliusson PhD), National Institute of Public Health, Bergen, Norway; Department of Clinical Science (Prof P B Juliusson PhD), Department of Psychosocial Science (Prof S Överland PhD), University of Bergen, Bergen, Norway; Institute of Family Medicine and Public Health (M Jürisson PhD), University of Tartu, Tartu, Estonia; School of Public Health (Z Kabir PhD), University College Cork, Cork, Ireland; University of Texas Health Science Center, San Antonio, TX, USA (S Kaur MD); International Research Center of Excellence (G A Kayode PhD), Institute of Human Virology Nigeria, Abuja, Nigeria; Julius Centre for Health Sciences and Primary Care (G A Kayode PhD), Utrecht University, Utrecht, Netherlands; Open, Distance and eLearning Campus (Prof P N Keiyoro PhD), Department of Psychiatry (M Kumar PhD), School of Economics (M K Muriithi PhD), School of Public Health (R G Wamai PhD), University of Nairobi, Nairobi, Kenya; School of Food and Agricultural Sciences (N Khalid PhD), University of Management and Technology, Lahore, Pakistan; Health Promotion Research Center (M Khammarnia PhD), Zahedan University of Medical Sciences, Zahedan, Iran; Department of Population Science (M Khan PhD), Jatiya Kabi Kazi Nazrul Islam University, Mymensingh, Bangladesh; Faculty of Health and Wellbeing (K Khatab PhD), Sheffield Hallam University, Sheffield, UK; College of Arts and Sciences (K Khatab PhD), Ohio University, Zanesville, OH, USA; The Iranian Academy of Medical Sciences, Tehran, Iran (M Khayamzadeh MD); Department of Psychiatry (C Kieling MD), Federal University of Rio Grande do Sul, Porto Alegre, Brazil; Division of Child & Adolescent Psychiatry (C Kieling MD), Clinical Hospital, Porto Alegre, Brazil; Big Data Department (Y Kim PhD), National Health Insurance Service, Wonju, South Korea; School of Traditional Chinese Medicine (Y Kim PhD), Xiamen University Malaysia, Sepang, Malaysia; Department of Nutrition (R W Kimokoti MD), Simmons University, Boston, MA, USA; School of Health Sciences (Prof A Kisa PhD), Kristiania University College, Oslo, Norway; Department of Global Community Health and Behavioral Sciences (Prof A Kisa PhD), Tulane University, New Orleans, LA, USA; Department of Nursing and Health Promotion (S Kisa PhD), Oslo Metropolitan University, Oslo, Norway; Department of Public Health (Prof M Kivimäki PhD), University of Helsinki, Helsinki, Finland (T J Meretoja MD); Independent Consultant, Jakarta, Indonesia (S Kosen MD); CIBERSAM (A Koyanagi MD), San Juan de Dios Sanitary Park, Sant Boi de Llobregat, Spain; Catalan Institution for Research and Advanced Studies (ICREA), Barcelona, Spain (A Koyanagi MD); Department of Anthropology (K Krishan PhD), Panjab University, Chandigarh, India; University of Environment and Sustainable Development, Somanya, Ghana (N Kugbey PhD); Faculty of Health and Life Sciences (O P Kurmi PhD), Coventry University, Coventry, UK; Department of Medicine (O P Kurmi PhD), Department of Psychiatry and Behavioural Neurosciences (A T Olagunju MD), Population Health Research Institute (T Sathish PhD), McMaster University, Hamilton, ON, Canada; Faculty of Public Health (D Kusuma DSc), University of Indonesia, Depok, Indonesia; National Institute for Health Research (NIHR) Oxford Biomedical Research Centre, Oxford, UK (B Lacey PhD); School of Dentistry (R Laloo PhD), Queensland Brain Institute (Prof J J McGrath MD), School of Health and Rehabilitation Sciences (R Uddin PhD), The University of Queensland, Brisbane, QLD, Australia; Division of Cancer Epidemiology and Genetics (Q Lan PhD), National Cancer Institute, Rockville, MD, USA; Unit of Genetics and Public Health (Prof I Landires MD), Unit of Microbiology and Public Health (V Nuñez-Samudio PhD), Institute of Medical Sciences, Las Tablas, Panama; Department of Public Health (V Nuñez-Samudio PhD), Ministry of Health, Herrera, Panama (Prof I Landires MD); HelpMeSee, New York, NY, USA (Prof V C Lansingh PhD); Mexican Institute of Ophthalmology, Queretaro, Mexico (Prof V C Lansingh PhD); Department of Medical Sciences (Prof A O Larsson PhD), Uppsala University, Uppsala, Sweden; Department of Clinical Chemistry and Pharmacology (Prof A O Larsson PhD), Uppsala University Hospital, Uppsala, Sweden; Department of Otorhinolaryngology (S Lasrado MS), Father Muller Medical College, Mangalore, India; Institute of Clinical Physiology (P Lauriola MD), National Research Council, Pisa, Italy; School of Pharmacy (S W H Lee PhD), Monash University, Bandar Sunway, Malaysia; School of Pharmacy (S W H Lee PhD), Taylor's University Lakeside Campus, Subang Jaya, Malaysia; UO Neurologia, Salute Pubblica e Disabilità (M Leonardi MD, A Raggi PhD, D Sattin PsyD, S Schiavolin MSc), Fondazione IRCCS Istituto Neurologico Carlo Besta (Neurology, Public Health and Disability Unit, Carlo Besta Neurological Institute), Milan, Italy; Center for Youth Substance Abuse Research (J Leung PhD), The University of Queensland, St Lucia, QLD, Australia; Department of Prevention (M Levi PhD), USL Tuscany Center, Firenze, Italy; Department of Health Sciences (M Levi PhD), University of Florence, Florence, Italy; Oxford University Clinical Research Unit (S Lewycka PhD), Wellcome Trust Asia Programme, Hanoi, Vietnam; Department of Sociology (B Li PhD), Shenzhen University, Shenzhen, China; Department of Health Promotion and Health Education (M Li PhD), National Taiwan Normal University, Taipei, Taiwan; Department of Medicine (L Lim MRCP), University of Malaya, Kuala Lumpur, Malaysia; Department of Medicine and Therapeutics (L Lim MRCP), The Chinese University of Hong Kong, Shatin, N.T., China; Institute of Nutritional Sciences (Prof S Lorkowski PhD), Friedrich Schiller University Jena, Jena, Germany; Competence Cluster for Nutrition and Cardiovascular Health (nutriCARD), Jena, Germany (Prof S Lorkowski PhD); Department of Medicine (Prof P A Lotufo DrPH), University of São Paulo, São Paulo, Brazil; Department of General Surgery (Prof R Lunevicius DSc), Liverpool University Hospitals NHS Foundation Trust, Liverpool, UK; Department of Surgery (Prof R Lunevicius DSc), Institute of Population Health Sciences (M R Mathur PhD), University of Liverpool, Liverpool, UK; Grants, Innovation and Product Development Unit (P W Mahasha PhD), Unit on Risk and Resilience in Mental Disorders (Prof D J Stein MD), South African Medical Research Council, Cape Town, South Africa (C A Nnaji MPH); Social Determinants of Health Research Center (M M Mahdavi PhD), Saveh University of Medical Sciences, Saveh, Iran; Radiology and Precision Health Program (M Mahmoudi PhD), Michigan State University, East Lansing, MI, USA; Environmental Health Research Center (Prof A Maleki PhD, S Mohammadi PhD), Social Determinants of Health Research Center (G Moradi PhD), Department of Epidemiology

and Biostatistics (G Moradi PhD), Kurdistan University of Medical Sciences, Sanandaj, Iran; Institute for Social Science Research (A A Mamun PhD), The University of Queensland, Indooroopilly, QLD, Australia; Department of Economics (Prof G Martinez PhD), Autonomous Technology Institute of Mexico, Mexico City, Mexico; Psychiatry Department (J Martinez-Raga PhD), Hospital Universitario Doctor Peset, Valencia, Spain; Department of Medicine (J Martinez-Raga PhD, Prof R Tabarés-Seisdedos PhD), University of Valencia, Valencia, Spain; Campus Caucaia (F R Martins-Melo PhD), Federal Institute of Education, Science and Technology of Ceará, Caucaia, Brazil; Department of Health Sciences (A J Mason-Jones PhD), University of York, York, UK; National Centre for Register-based Research (Prof J J McGrath MD), Aarhus University, Aarhus, Denmark; Neurology Department (Prof M Mehndiratta MD), Janakpuri Super Specialty Hospital Society, New Delhi, India; Department of Neurology (Prof M Mehndiratta MD), Govind Ballabh Institute of Medical Education and Research, New Delhi, India; Nutrition Health Research Center (F Mehri PhD), Iran University of Medical Sciences, Hamadan, Iran; Institute of Human Virology (P T N Memiah DrPH), University of Maryland, Baltimore, MD, USA; Peru Country Office (W Mendoza MD), United Nations Population Fund (UNFPA), Lima, Peru; Neurology Unit (A Meretoja MD), Breast Surgery Unit (T J Meretoja MD), Helsinki University Hospital, Helsinki, Finland; Clinical Microbiology and Parasitology Unit (T Mestrovic PhD), Dr Zora Profozic Polyclinic, Zagreb, Croatia; University Centre Varazdin (T Mestrovic PhD), University North, Varazdin, Croatia; Center for Innovation in Medical Education (B Miazgowski MD), Department of Propeudeutics of Internal Diseases & Arterial Hypertension (Prof T Miazgowski MD), Pomeranian Medical University, Szczecin, Poland (B Miazgowski MD); Woman-Mother-Child Department (I Michalek PhD), Lausanne University Hospital, Lausanne, Switzerland; Pacific Institute for Research & Evaluation, Calverton, MD, USA (T R Miller PhD); Global Institute of Public Health (Prof G Mini PhD), Ananthapuri Hospitals and Research Institute, Trivandrum, India; Women's Social and Health Studies Foundation, Trivandrum, India (Prof G Mini PhD); Internal Medicine Programme (Prof E M Mirrakhimov PhD), Kyrgyz State Medical Academy, Bishkek, Kyrgyzstan; Department of Atherosclerosis and Coronary Heart Disease (Prof E M Mirrakhimov PhD), National Center of Cardiology and Internal Disease, Bishkek, Kyrgyzstan; Research Center for Biochemistry and Nutrition in Metabolic Diseases (H Mirzaei PhD), Kashan University of Medical Sciences, Kashan, Iran; Institute of Addiction Research (ISFF) (B Moazen MSc), Frankfurt University of Applied Sciences, Frankfurt, Germany; Department of Forestry (D K Mohammad PhD), Salahaddin University-Erbil, Erbil, Iraq; Research Institute for Health Development (S Mohammadi PhD), Kurdistan University of Medical Sciences, Sanandaj City, Iran; Department of Epidemiology and Biostatistics (A Mohammadian-Hafshejani PhD), Shahrekord University of Medical Sciences, Shahrekord, Iran; Department of Nursing (R Mohammadpourhodki PhD), Applied Biomedical Research Center (A Sahebkar PhD), Biotechnology Research Center (A Sahebkar PhD), Mashhad University of Medical Sciences, Mashhad, Iran; Health Systems and Policy Research Unit (S Mohammed PhD), Ahmadu Bello University, Zaria, Nigeria; Department of Health Care Management (S Mohammed PhD), Technical University of Berlin, Berlin, Germany; Clinical Epidemiology and Public Health Research Unit (L Monasta DSc, L Ronfani PhD), Burlo Garofolo Institute for Maternal and Child Health, Trieste, Italy; Computer, Electrical, and Mathematical Sciences and Engineering Division (P Moraga PhD), King Abdullah University of Science and Technology, Thuwal, Saudi Arabia; Section of Plastic Surgery (S D Morrison MD), University of Michigan School of Medicine, Ann Arbor, MI, USA; Department of Food Science (Prof A Mousavi Khaneghah PhD), University of Campinas (Unicamp), Campinas, Brazil; Competence Center of Mortality-Follow-Up of the German National Cohort (R Westerman DSc), Federal Institute for Population Research, Wiesbaden, Germany (Prof U O Mueller MD); Center for Population and Health, Wiesbaden, Germany (Prof U O Mueller MD); Saveetha Dental College (S Muthupandian PhD), Saveetha Institute of Medical and Technical Sciences (SIMATS), Chennai, India; Research and Analytics Department (A J Nagarajan M.Tech), Initiative for Financing Health and Human Development, Chennai, India; Department of Research and Analytics (A J Nagarajan M.Tech), Bioinsilico Technologies, Chennai, India; Laboratory of Public Health Indicators Analysis and Health Digitalization (M Naimzada MD, N Otvstavnov BA, S S Otvstavnov PhD), Moscow Institute of Physics and Technology, Dolgoprudny, Russia; Experimental Surgery and Oncology Laboratory (M Naimzada MD), Kursk State Medical University, Kursk, Russia; Suraj Eye Institute, Nagpur, India (V Nangia MD); Department of Disease Control and Environmental Health (R Ndejjo MSc), Makerere University, Kampala, Uganda; Department of General Surgery (I Negoii PhD), Emergency Hospital of Bucharest, Bucharest, Romania; Cardio-Aid, Bucharest, Romania (R I Negoii PhD); College of Medicine and Health Sciences (H B Netsere MS), Bahir Dar University, Gondar, Ethiopia; Department of Public Health (G Nguefack-Tsague PhD), Department of Internal Medicine and Specialties (Prof E Sobngwi PhD), University of Yaoundé I, Yaoundé, Cameroon; Institute for Global Health Innovations (D N Nguyen MA), Faculty of Pharmacy (D N Nguyen MA), Duy Tan University, Da Nang, Vietnam; Institute for Global Health Innovations (H L T Nguyen MPH, H Q Pham MD), Duy Tan University, Hanoi, Vietnam; Department of Sociology & Institute for Empirical Social Science Research (J Nie PhD), Xi'an Jiaotong University, Xi'an, China; Public Health Department (D N A Ningrum MPH), Universitas Negeri Semarang (State University of Semarang), Kota Semarang, Indonesia; School of Public Health and Family Medicine (C A Nnaji MPH), University of Cape Town, Cape Town, South Africa; Department of Health Policy and Management (S Nomura PhD), Keio University, Tokyo, Japan; Department of Global Health Policy (S Nomura PhD), University of Tokyo, Tokyo, Japan; Department of Neurobiology, Care Sciences and Society (C Nowak PhD), Karolinska Institute, Huddinge, Sweden; Department of Preventive Medicine (I Oh PhD), Kyung Hee University, Dongdaemun-gu, South Korea; Gorgan Congenital Malformations Research Center (M Oladnabi PhD), Golestan University of Medical Sciences, Gorgan, Iran; Department of Psychiatry (A T Olagunju MD), University of Lagos, Lagos, Nigeria; Centre for Healthy Start Initiative, Lagos, Nigeria (B O Olusanya PhD, J O Olusanya MBA); Department of Pharmacology and Therapeutics (Prof O E Onwujekwe PhD), University of Nigeria Nsukka, Enugu, Nigeria; Department of Medicine (Prof A Ortiz MD), Autonomous University of Madrid, Madrid, Spain; Department of Nephrology and Hypertension (Prof A Ortiz MD), The Institute for Health Research Foundation Jiménez Díaz University Hospital, Madrid, Spain; Department of Project Management (S S Otvstavnov PhD), Department of Health Care Administration and Economics (Prof V Vlassov MD), National Research University Higher School of Economics, Moscow, Russia; Division of Mental and Physical Health (Prof S Øverland PhD), Norwegian Institute of Public Health, Bergen, Norway; Department of Respiratory Medicine (Prof M P A DNB), Jagadguru Sri Shivarathreeswara Academy of Health Education and Research, Mysore, India; Department of Health Metrics (A Pana MD), Center for Health Outcomes & Evaluation, Bucharest, Romania; Department of Medical Humanities and Social Medicine (Prof E Park PhD), Kosin University, Busan, South Korea; Community Health Sciences (Prof S B Patten PhD), Department of Medicine (Prof M Tonelli MD), University of Calgary, Calgary, AB, Canada; School of Public Health, Medical, and Veterinary Sciences (A E Peden PhD), James Cook University, Townsville, NSW, Australia; Center for Research and Innovation (V F Pepito MSc), Ateneo De Manila University, Pasig City, Philippines; School of Global Public Health (E K Peparah PhD), New York University, New York, NY, USA; Department of Orthopedics (J Pereira MS), Yenepoya Medical College, Mangalore, India; Shanghai Mental Health Center (Prof M R Phillips MD), Shanghai Jiao Tong University, Shanghai, China; University Medical Center Groningen (Prof M J Postma PhD), School of Economics and Business (Prof M J Postma PhD), University of Groningen, Groningen, Netherlands; Dietary Supplements and Probiotic Research Center (H Pourjafar PhD), Alborz University of Medical Sciences, Karaj, Iran; Centro de Investigaciones Clínicas (Clinical Research Center) (S I Prada PhD), Fundación Valle del Lili (Valle del Lili Foundation), Cali, Colombia; Centro PROESA (PROESA Center) (S I Prada PhD), Centro de Investigaciones en Anomalías Congénitas y Enfermedades Raras (Center for Research in Congenital Anomalies and Rare Diseases) (E Rubagotti PhD), Universidad ICESI (ICESI University), Cali, Colombia; Department of Epidemiology (M Rabiee PhD), University of Tehran, Tehran, Iran; Department of Chemistry (N Rabiee MSc), Sharif University of Technology, Tehran, Iran; College of Medicine (A Radfar MD),

University of Central Florida, Orlando, FL, USA; Department of Medicine (A Rafiee MSc), University of Alberta, Edmonton, AB, Canada; Thalassaemia and Hemoglobinopathy Research Center (F Rahim PhD), Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran; Department of Community Medicine (M Rahman PhD), Maharishi Markandeshwar Medical College & Hospital, Solan, India; School of Nursing and Healthcare Professions (M Rahman PhD), Federation University Australia, Berwick, VIC, Australia; Research Department (C L Ranabhat PhD), Policy Research Institute, Kathmandu, Nepal; Health and Public Policy Department (C L Ranabhat PhD), Global Center for Research and Development, Kathmandu, Nepal; Department of Oral Pathology (S Rao MDS), Srinivas Institute of Dental Sciences, Mangalore, India; University College London Hospitals, London, UK (D L Rawaf MD); Academic Public Health England (Prof S Rawaf MD), Public Health England, London, UK; School of Health, Medical and Applied Sciences (L Rawal PhD), CQ University, Sydney, NSW, Australia; School of Medicine (Prof A M N Renzaho PhD), Translational Health Research Institute (Prof A M N Renzaho PhD), Western Sydney University, Campbelltown, NSW, Australia; Network of Immunity in Infection, Malignancy and Autoimmunity (NIIMA) (Prof N Rezaei PhD), Universal Scientific Education and Research Network (USERN), Tehran, Iran; Department of Surgery (J Rickard MD), University of Minnesota, Minneapolis, MN, USA; Department of Surgery (J Rickard MD), University Teaching Hospital of Kigali, Kigali, Rwanda; Department of Clinical Research (L Roever PhD), Federal University of Uberlândia, Uberlândia, Brazil; African Genome Center (E Rubagotti PhD), Mohammed VI Polytechnic University (UM6P), Ben Guerir, Morocco; Malaria Atlas Project (S F Rumisha PhD), Telethon Kids Institute, Perth, Australia; Department of Health Statistics (S F Rumisha PhD), National Institute for Medical Research, Dar es Salaam, Tanzania; Department of Internal Medicine (G M Rwegera MD), University of Botswana, Gaborone, Botswana; Neuropsychiatric Institute (Prof P S Sachdev MD), Prince of Wales Hospital, Randwick, NSW, Australia; Department of Psychiatry (Prof R Sagar MD), All India Institute of Medical Sciences, New Delhi, India; Department of Phytochemistry (Prof S Sajadi PhD), Soran University, Soran, Iraq; Department of Nutrition (Prof S Sajadi PhD), Cihan University-Erbil, Erbil, Iraq; Public Health and Community Medicine Department (M R Salem MD), Cairo University, Giza, Egypt; Department of Surgery (Prof J Sanabria MD), Marshall University, Huntington, WV, USA; Department of Community Medicine (S Y Saraswathy PhD), PSG Institute of Medical Sciences and Research, Coimbatore, India; PSG-FAIMER South Asia Regional Institute, Coimbatore, India (S Y Saraswathy PhD); School of Population and Public Health (Prof N Sarrafzadegan MD), University of British Columbia, Vancouver, BC, Canada; Department of Psychology (D C Schwebel PhD), School of Medicine (Prof J A Singh MD), University of Alabama at Birmingham, Birmingham, AL, USA; Oral Diagnosis, Digital Health and Health Services Research (Prof F Schwendicke PhD), Charité University Medical Center Berlin, Berlin, Germany; Emergency Department (S Senthilkumaran MD), Manian Medical Centre, Erode, India; Center for Biomedical Information Technology (F Sha PhD), Shenzhen Institutes of Advanced Technology, Shenzhen, China; Public Health Division (A A Shaheen PhD), An-Najah National University, Nablus, Palestine; Independent Consultant, Karachi, Pakistan (M A Shaikh MD); Department of Community Medicine (M Shannawaz PhD), BLDE University, Vijayapur, India; Centre for Medical Informatics (Prof A Sheikh MD), University of Edinburgh, Edinburgh, UK; Friedman School of Nutrition Science and Policy (P Shi PhD), Tufts University, Boston, MA, USA; National Institute of Infectious Diseases, Tokyo, Japan (M Shigematsu PhD); College of Medicine (Prof J Shin MD), Yonsei University, Seoul, South Korea; Finnish Institute of Occupational Health, Helsinki, Finland (R Shiri PhD); School of Public Health (K Shuval PhD), University of Haifa, Haifa, Israel; The Cooper Institute, Dallas, TX, USA (K Shuval PhD); School of Health (S Siabani PhD), University of Technology Sydney, Sydney, NSW, Australia; Department of Psychology (Prof I D Sigfusdottir PhD, R Sigurvinsdottir PhD), Reykjavik University, Reykjavik, Iceland; Department of Physical Education (Prof D A S Silva PhD), Federal University of Santa Catarina, Florianópolis, Brazil; Department of Law, Economics, Management and Quantitative Methods (Prof B Simonetti PhD), University of Sannio, Benevento, Italy; WSB University in Gdańsk, Gdańsk, Poland (Prof B Simonetti PhD); Medicine Service (Prof J A Singh MD), US Department of Veterans Affairs (VA), Birmingham, AL, USA; Department of Pulmonary Medicine (Prof V Singh MD), Asthma Bhawan, Jaipur, India; Program Services Unit (A H Sinke MD), Pathfinder International, Addis Ababa, Ethiopia; Department No16 (V Y Skryabin MD), Laboratory of Genetics and Genomics (Prof M S Zastrozhin PhD), Moscow Research and Practical Centre on Addictions, Moscow, Russia; Institute of Bone and Joint Research (IBJR) (E U R Smith PhD), University of Sydney, St Leonards, NSW, Australia; Australian New Zealand Clinical Trials Registry (ANZCTR), NHMRC Clinical Trials Centre (E U R Smith PhD), University of Sydney, Camperdown, NSW, Australia; Department of Endocrinology and Diabetes (Prof E Sobngwi PhD), Yaoundé Central Hospital, Yaoundé, Cameroon; Nursing Care Research Center (A Soheili PhD), Semnan University of Medical Sciences, Semnan, Iran; Demography and Population Studies Department (O D Somefun PhD), University of Witwatersrand, Johannesburg, South Africa; Hull York Medical School (I N Soyiri PhD), University of Hull, Hull City, UK; Division of Community Medicine (C T Sreeramareddy MD), International Medical University, Kuala Lumpur, Malaysia; Department of Nursing (A Sudaryanto MPH), Muhammadiyah University of Surakarta, Surakarta, Indonesia; Pediatric Services (I Sultan MD), King Hussein Cancer Center, Amman, Jordan; Pediatrics (I Sultan MD), University of Jordan, Amman, Jordan; Carlos III Health Institute (Prof R Tabarés-Seisdedos PhD), Biomedical Research Networking Center for Mental Health Network (CiberSAM), Madrid, Spain; Cancer Control Center (T Tabuchi MD), Osaka International Cancer Institute, Osaka, Japan; School of Dentistry and Oral Health (S K Tadakamadla PhD), Griffith University, Gold Coast, QLD, Australia; Department of Population Science and Human Resource Development (Prof M I Tareque PhD), University of Rajshahi, Rajshahi, Bangladesh; Department of Public Health and Community Medicine (Prof K R Thankappan MD), Central University of Kerala, Kasaragod, India; KA Timiryazev Institute of Plant Physiology (M V Titova PhD), Russian Academy of Sciences, Moscow, Russia; Laboratory of Public Health Indicators Analysis and Health Digitalization (M V Titova PhD), Moscow Institute of Physics and Technology, Moscow, Russia; Department of Pathology and Legal Medicine (M R Tovani-Palone PhD), University of São Paulo, Ribeirão Preto, Brazil; Modestum LTD, London, UK (M R Tovani-Palone PhD); Department of Health Economics (B X Tran PhD), Hanoi Medical University, Hanoi, Vietnam; Mbarara University of Science and Technology, Mbarara, Uganda (A C Tsai MD); Department of Medicine (Prof A Tsatsakis DSc), University of Crete, Heraklion, Greece; Lee Kong Chian School of Medicine (L Tudor Car PhD), Nanyang Technological University, Singapore, Singapore; Institute for Physical Activity and Nutrition (R Uddin PhD), Deakin University, Melbourne, VIC, Australia; Department of Cardiovascular, Endocrine-metabolic Diseases and Aging (B Unim PhD), National Institute of Health, Rome, Italy; Amity Institute of Biotechnology (E Upadhyay PhD), Amity University Rajasthan, Jaipur, India; Clinical Cancer Research Center (S Valadan Tahbaz PhD, S Yahyazadeh Jabbari MD), Milad General Hospital, Tehran, Iran; Velez Sarsfield Hospital, Buenos Aires, Argentina (Prof P R Valdez MD); UKK Institute, Tampere, Finland (Prof T J Vasankari MD); Raffles Neuroscience Centre (Prof N Venkatasubramanian MBBS), Raffles Hospital, Singapore, Singapore; Yong Loo Lin School of Medicine (Prof N Venkatasubramanian MBBS), National University of Singapore, Singapore, Singapore; School of Mathematics and Statistics (Prof P J Villeneuve PhD), Carleton University, Ottawa, ON, Canada; Department of Medical and Surgical Sciences (Prof F S Violante MD), University of Bologna, Bologna, Italy; Occupational Health Unit (Prof F S Violante MD), Sant'Orsola Malpighi Hospital, Bologna, Italy; Center of Excellence in Behavioral Medicine (G T Vu BA), Nguyen Tat Thanh University, Ho Chi Minh City, Vietnam; Foundation University Medical College (Prof Y Waheed PhD), Foundation University Islamabad, Islamabad, Pakistan; Cultures, Societies and Global Studies, & Integrated Initiative for Global Health (R G Wamai PhD), Northeastern University, Boston, MA, USA; Department of Epidemiology and Biostatistics (Y Wang BSA, Prof C Yu PhD), School of Medicine (Z Zhang PhD), School of Health Sciences (X G Zhao PhD), Wuhan University, Wuhan, China; School of Population Health and Environmental Sciences (Y Wang PhD), King's College London, London, UK; Department of Community Medicine (N D Wickramasinghe MD), Rajarata University of Sri Lanka, Anuradhapura, Sri Lanka; Department of Orthopaedics (Prof A Wu MD),

Wenzhou Medical University, Wenzhou, China; Research and Development Center for Health Services (Prof K Yamagishi MD), University of Tsukuba, Tsukuba, Japan; Department of Family Medicine and Community Health (Y Yano MD), Duke University, Durham, IL, USA; Health Services Management Research Center (V Yazdi-Fezabadi PhD), Department of Health Management, Policy, and Economics (V Yazdi-Fezabadi PhD), Kerman University of Medical Sciences, Kerman, Iran; Department of Neuropsychopharmacology (N Yonemoto MPH), National Center of Neurology and Psychiatry, Kodaira, Japan; Department of Public Health (N Yonemoto MPH), Juntendo University, Tokyo, Japan; Department of Preventive Medicine (Prof S Yoon PhD), Korea University, Seoul, South Korea; Department of Health Policy and Management (Prof M Z Younis PhD), Jackson State University, Jackson, MS, USA; School of Medicine (Prof M Z Younis PhD), Tsinghua University, Beijing, China; School of Public Health and Management (Y Yu MS), Hubei University of Medicine, Shiyan, China; Cancer Institute (D Yuce MD), Hacettepe University, Ankara, Turkey; Department of Pharmaceutics (S Zaidi PhD), Dow University of Health Sciences, Karachi, Pakistan; Department of Environmental Health Engineering (A Zarei PhD), Gonabad University of Medical Sciences, Gonabad, Iran; Addictology Department (Prof M S Zastrozhin PhD), Russian Medical Academy of Continuous Professional Education, Moscow, Russia; School of Public Health (Y Zhang PhD), Hubei Province Key Laboratory of Occupational Hazard Identification and Control (Y Zhang PhD), Wuhan University of Science and Technology, Wuhan, China; School of Biology and Pharmaceutical Engineering (X G Zhao PhD), Wuhan Polytechnic University, Wuhan, China; Department of Epidemiology, Human Genetics, and Environmental Sciences (C Zhu MPH), University of Texas Health Science Center at Houston, Houston, TX, USA.

Contributors

Joseph L Ward, Peter S Azzopardi, Russell M Viner, and George C Patton conceptualised the paper. Joseph L Ward downloaded the data and created the tables and visualisations. Joseph L Ward, Peter S Azzopardi, Russell M Viner, and George C Patton wrote the first draft. All other authors provided data, developed models, reviewed results, provided guidance on methods, or reviewed and contributed to the manuscript. All authors approved the final version of the manuscript.

Declaration of interests

R Ancuceanu reports consulting fees from AbbVie and AstraZeneca; payment or honoraria for lectures, presentations, speaker's bureaus, manuscript writing or educational events from Sandoz, AbbVie, and B Braun; and support for attending meetings and/or travel from AbbVie and AstraZeneca, all outside the submitted work. J Årnlöv reports payment or honoraria for lectures, presentations, speaker's bureaus, manuscript writing or educational events from AstraZeneca and Novartis; and payment for expert testimony from AstraZeneca and Boehringer Ingelheim, all outside the submitted work. M Ausloos reports grants or contracts from Romanian National Authority for Scientific Research and Innovation, CNDS-UEFISCDI, project number PN-III-P4-ID-PCCF-2016-0084, outside the submitted work. T Bärnighausen reports research grants from the European Union (Horizon 2020 and EIT Health), German Research Foundation (DFG), US National Institutes of Health, German Ministry of Education and Research, Alexander von Humboldt Foundation, Else-Kröner-Fresenius-Foundation, Wellcome Trust, Bill & Melinda Gates Foundation, KfW, UNAIDS, and WHO; consulting fees for KfW on the OSCAR initiative in Vietnam; and participation on a Data Safety Monitoring Board or Advisory Board through the NIH-funded study "Healthy Options" (PIs: Smith Fawzi, Kaaya), Chair, Data Safety and Monitoring Board (DSMB), German National Committee on the "Future of Public Health Research and Education", Chair of the scientific advisory board to the EDCTP Evaluation, Member of the UNAIDS Evaluation Expert Advisory Committee, National Institutes of Health Study Section Member on Population and Public Health Approaches to HIV/AIDS (PPAH), US National Academies of Sciences, Engineering, and Medicine's Committee for the "Evaluation of Human Resources for Health in the Republic of Rwanda under the President's Emergency Plan for AIDS Relief (PEPFAR)", University of Pennsylvania (UPenn) Population Aging Research Center (PARC) External Advisory Board Member; and

Leadership or fiduciary role in other board, society, committee or advocacy group, paid or unpaid as Co-chair of the Global Health Hub Germany (which was initiated by the German Ministry of Health), all outside the submitted work. M Bell reports research funding and payments to institution from Wellcome Trust, NIH, Yale Climate Change and Health Center, Robert Wood Johnson Foundation, Yale Women Faculty Forum, EPA, and High Tide Foundation; honorarium for proposal review from NIH and Johns Hopkins University; honorarium from mentoring program from University of Montana; and travel expenses to give seminars from University of Illinois at Champaign, Johns Hopkins University, Ohio State University, Royal Society London, Atmospheric Chemistry Gordon Research Conference, New York School of Medicine, and Global Research Laboratory (Seoul), Seoul National University, all outside the submitted work. Z Bhutta reports grants or contracts from the Bill & Melinda Gates foundation, outside the submitted work. I Filip reports financial support from Avicenna Medical and Clinical Research Institute, outside the submitted work. B Hall reports stocks that are unrelated to the paper, outside the submitted work. G Hankey reports consulting fees from Bayer for stroke prevention advisory boards; payment or honoraria for lectures, presentations, speakers bureaus, manuscript writing or educational events from the American Heart Association, Medscape, and Bristol Myers Squibb; and participation on a Data Safety Monitoring Board or Advisory Board with AC Immune, all outside the submitted work. C Herteliu reports grants or contracts from Romanian National Authority for Scientific Research and Innovation, CNDS-UEFISCDI, project number PN-III-P4-ID-PCCF-2016-0084, research grant (Oct 2018–Sept 2022) "Understanding and modelling time-space patterns of psychology-related inequalities and polarization", and project number PN-III-P2-2.1-SOL-2020-2-0351, research grant (June 2021–Oct 2021) "Approaches within public health management in the context of COVID-19 pandemic", and from the Ministry of Labour and Social Justice Romania, project number 30/PSCD/2018, research grant (Sept 2018–June 2019) "Agenda for skills Romania 2020–2025", all outside the submitted work. S Islam reports grants or contracts from NHMRC and National Heart Foundation of Australia Fellowships, outside the submitted work. R Ivers reports support for the present manuscript from the National Health and Medical Research Council of NSW through salary funding via a senior research fellowship. V Jha reports grants or contracts from GSK, Baxter Healthcare, and AstraZeneca, outside the submitted work. J Jozwiak reports payment or honoraria for lectures, presentations, speaker's bureaus, manuscript writing or educational events from Teva, Amgen, Synexis, Boehringer Ingelheim, ALAB Laboratories, and Zentiva, all outside the submitted work. C Kieling reports grants or contracts from MQ: Transforming Mental Health, UK Academy of Medical Sciences, UK Royal Academy of Engineering, and US National Institute of Mental Health; and royalties or licences from Manole, outside the submitted work. K Krishan reports non-financial support from UGC Centre of Advanced Study, CAS II, Department of Anthropology, Panjab University, Chandigarh, India, outside the submitted work. S Lorkowski reports grants or contracts from Akcea Therapeutics Germany; consulting fees from Danone, Swedish Orphan Biovitrum (SOBI), and Upfield; payment or honoraria for lectures, presentations, speaker's bureaus, manuscript writing or educational events from Akcea Therapeutics Germany, AMARIN Germany, Amedes Holding, AMGEN, Berlin-Chemie, Boehringer Ingelheim Pharma, Daiichi Sankyo Deutschland, Danone, Hubert Burda Media Holding, Lilly Deutschland, Novo Nordisk Pharma, Roche Pharma, Sanofi-Aventis, SYNLAB Holding Deutschland & SYNLAB Akademie; participation on a Data Safety Monitoring Board or Advisory Board with Akcea Therapeutics Germany, AMGEN, Daiichi Sankyo Deutschland, and Sanofi-Aventis, all outside the submitted work. R Maddison reports grants or contracts from NHMRC Ideas Grant and National Heart Foundation of Australia Vanguard Grant, outside the submitted work. M Mahmoudi reports placement as a co-founder and director of the Academic Parity Movement, a non-profit organisation dedicated to addressing academic discrimination, violence and incivility; and receives royalties/honoraria for his published books, plenary lectures, and licensed patents, outside the submitted work. S Nomura reports grant support for the present manuscript from the Ministry of Education, Culture, Sports, Science and Technology (MEXT). C Nowak

reports employment with Diamyd Medical AB (Stockholm, Sweden) which develops a treatment for type 1 diabetes, outside the submitted work. A Ortiz reports grants or contracts from Sanofi, Mundipharma, and AstraZeneca; and payment or honoraria for lectures, presentations, speakers bureaus, manuscript writing or educational events from Astellas, Astrazeneca, Amicus, Amgen, Fresenius Medical Care, Bayer, Sanofi-Genzyme, Menarini, Kyowa Kirin, Alexion, Otsuka and Vifor Fresenius Medical Care Renal Pharma. C Panelo reports grants or contracts from USAID Philippines as Chief of Party of USAID's ProtectHealth, a financing policy support project assisting the Philippine Government in implementing universal health care with focus on tuberculosis, family planning and HIV, outside the submitted work. M Postma reports leadership or fiduciary role in other board, society, committee or advocacy group, unpaid as member of UK's JCVI. A Radfar reports financial support from Avicenna Medical and Clinical Research Institute. P Sachdev reports an investigator grant from the National Health and Medical Research Council of Australia, paid to the University; participation on a Data Safety Monitoring Board or Advisory Board with the Advisory Committee for Biogen Australia; and leadership or fiduciary role in other board, society, committee or advocacy group, paid or unpaid as honorary director of International Neuropsychiatric Association and executive member of VASCOG; outside the submitted work. J Singh reports consulting fees from Crealta/Horizon, Medisys, Fidia, Two labs Inc, Adept Field Solutions, Clinical Care options, Clearview healthcare partners, Putnam associates, Focus forward, Navigant consulting, Spherix, MedIQ, UBM LLC, Trio Health, Medscape, WebMD, and Practice Point communications; and the National Institutes of Health and the American College of Rheumatology; payment or honoraria for lectures, presentations, speakers bureaus, manuscript writing or educational events from Simply Speaking; support for attending meetings and/or travel from OMERACT, an international organisation that develops measures for clinical trials and receives arm's length funding from 12 pharmaceutical companies, when traveling bi-annually to OMERACT meetings; participation on a Data Safety Monitoring Board or Advisory Board as a member of the FDA Arthritis Advisory Committee; leadership or fiduciary role in other board, society, committee or advocacy group, paid or unpaid, with OMERACT as a member of the steering committee, with the Veterans Affairs Rheumatology Field Advisory Committee as a member, and with the UAB Cochrane Musculoskeletal Group Satellite Center on Network Meta-analysis as a director and editor; and stock or stock options in TPT Global Tech, Vaxart pharmaceuticals, Charlotte's Web Holdings Inc and previously owned stock options in Amarin, Viking, and Moderna pharmaceuticals, all outside the submitted work. H Slater reports grants or contracts awarded for co-developing a digital platform with young people with musculoskeletal pain to support their care from the Washington Department of Health, 2018–2021. D Stein reports personal fees from Lundbeck, Takeda, Johnson & Johnson and Servier, all outside the submitted work. M Stokes reports payment or honoraria for lectures, presentations, speakers bureaus, manuscript writing or educational events from the Autism Teaching Institute; leadership or fiduciary role in other board, society, committee or advocacy group, paid or unpaid as Vice President of Kidsafe Victoria and Board Member of Australasian Society for Autism Research, all outside the submitted work. S Tadakamadla reports grants or contracts from National Health and Medical Research Council, Australia, Early Career Fellowship. A Tsai reports stipend for work as Editor-in-Chief of Social Science and Medicine: Mental Health from Elsevier, Inc. R Uddin is supported by an Alfred Deakin Postdoctoral Research Fellowship, Deakin University, Australia and reports support for attending meetings and/or travel accommodation reimbursement from Deakin University Institute for Physical Activity and Nutrition, all outside of the submitted work. All other authors declare no competing interests.

Data sharing

To download the data used in these analyses, please visit the Global Health Data Exchange at <http://ghdx.healthdata.org/gbd-2019>

Acknowledgments

This Article was supported by the Bill & Melinda Gates Foundation and the National Institute on Ageing of the National Institutes of Health (award P30AG047845). J L Ward is funded by the Medical Research

Council and G C Patton by an NHMRC Fellowship. P Azzopardi is supported by an NHMRC Fellowship GNT1145228. L Abreu would like to thank Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) - Código de Financiamento 001, Conselho Nacional de Desenvolvimento Científico e Tecnológico (404710/2018-2 and 310797/2019-5), and Fundação de Amparo à Pesquisa de Minas Gerais (FAPEMIG). S Aljunid would like to acknowledge the Department of Health Policy and Management, Faculty of Public Health, Kuwait University and International Centre for Casemix and Clinical Coding, Faculty of Medicine, National University of Malaysia for the approval and support to participate in this research project. T Bärnighausen was supported by the Alexander von Humboldt Foundation through the Alexander von Humboldt Professor award, funded by the German Federal Ministry of Education and Research. G Britton is supported by Sistema Nacional de Investigación (SNI) of Secretaría Nacional de Ciencia, Tecnología e Innovación (SENACYT) of Panamá. F Carvalho and E Fernandes are supported by UID/MULTI/04378/2019 and UID/QUI/50006/2019 with funding from FCT/MCTES through national funds. V Costa acknowledges her grant (SFRH/BHD/110001/2015), received by Portuguese national funds through Fundação para a Ciência e Tecnologia (FCT), IP, under the Norma Transitória DL57/2016/CP1334/CT0006. A Fomenkov and M Titova acknowledge support from the state assignment of Ministry of Science and Higher Education of the Russian Federation (theme 121050500047-5). A Gebreslassie acknowledges the Department of Reproductive Health, College of Health Sciences, School of Public Health at Mekelle University. C Herteliu and M Ausloos are partially supported by a grant of the Romanian National Authority for Scientific Research and Innovation, CNDS-UEFISCDI, project number PN-III-P4-ID-PCCF-2016-0084. P Hoogar acknowledges support from the Centre for Bio Cultural Studies (CBiCS), Directorate of Research, Manipal Academy of Higher Education, Manipal and Centre for Holistic Development and Research (CHDR), Kalaghatagi. S Islam is funded by a fellowship from the National Heart Foundation of Australia and NHMRC. M Jakovljevic acknowledges support through the Grant OI 175 014 of the Ministry of Education Science and Technological Development of the Republic of Serbia. P Jeemon received (2015-2021) Wellcome Trust/DBT India Alliance Clinical and Public Health Intermediate Fellowship (IA/CPHI/14/1/501497). M Khan acknowledges the support of Jatiya Kabi Kazi Nazrul Islam University, Bangladesh. C Kieling is a Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq, Brazil) researcher and an Academy of Medical Sciences (UK) Newton advanced fellow. YJ Kim was supported by the Research Management Centre, Xiamen University Malaysia [XMUMRF/2020-C6/ITCM/0004]. K Krishan is supported by the UGC Centre of Advanced Study (Phase II), awarded to the Department of Anthropology, Panjab University, Chandigarh, India. M Kumar acknowledges support from the Fogarty International Center/National Institutes of Health K43 TW010716-04. B Lacey acknowledges support from UK Biobank, University of Oxford. I Landires is member of the Sistema Nacional de Investigación (SNI), supported by the Secretaría Nacional de Ciencia, Tecnología e Innovación (SENACYT), Panama. S Lorkowski acknowledges institutional support from the Competence Cluster for Nutrition and Cardiovascular Health (nutriCARD) Halle-Jena-Leipzig (Germany; German Federal Ministry of Education and Research; grant agreement number 01EA1808A). P Mahasha acknowledges support from Grants, Innovation and Product Development (GIPD) Unit, South African Medical Research Council (SAMRC). A Mason-Jones was supported by the University of York. J McGrath is supported by the Danish National Research Foundation (Niels Bohr Professorship). J McGrath is employed by The Queensland Centre for Mental Health Research which receives core funding from the Department of Health, Queensland Government. S Nomura acknowledges support from the Ministry of Education, Culture, Sports, Science and Technology of Japan (MEXT). J Padubidri acknowledges Manipal Academy of Higher Education, Manipal for their constant support to academic research. S Patten holds the Cuthbertson & Fischer Chair at the University of Calgary. G Patton is supported by an NHMRC research fellowship. M Phillips is supported in part by National Natural Science Foundation of China (NSFC 81761128031). A Raggi, D Sattin and S Schiavolin are supported by a grant from the Italian Ministry of Health (Ricerca Corrente, Fondazione Istituto Neurologico C. Besta,

Linea 4 – Outcome Research: dagli Indicatori alle Raccomandazioni Cliniche). A Samy acknowledges the support from the Egyptian Fulbright Mission Program. A Sheikh is supported by HDRUK. B Shetty acknowledges the research support from Kasturba Medical College, Mangalore, Manipal Academy of Higher Education, Manipal. J Silva acknowledges support from grant number UIDB/04378/2020 from the Applied Molecular Biosciences Unit (UCIBIO), supported through Portuguese national funds via FCT/MCTES. S Tadakamadla acknowledges support from National Health and Medical Research Council Early Career Fellowship, Australia. R Uddin is supported by an Alfred Deakin Postdoctoral Research Fellowship, Deakin University, Australia. B Unnikrishnan acknowledges support from the Manipal Academy of Higher Education, Manipal. S Zaman received a scholarship from the Australian Government research training program (RTP) in support of his academic career. Y Zhang was supported by Science and Technology Research Project of Hubei Provincial Department of Education (Grant Q20201104), Open Fund Project of Hubei Province Key Laboratory of Occupational Hazard Identification and Control (Grant OHIC2020Y01), Outstanding Young and Middle Aged Technology Innovation Team Project of Hubei Provincial Department of Education (Grant T2020003), and the Key Research Center for Humanities and Social Sciences in Hubei Province (Hubei University of Medicine) (Grant 2020ZD001). C Zhu is funded by UTHealth Innovation for Cancer Prevention Research Training Program Pre-doctoral Fellowship (Cancer Prevention and Research Institute of Texas Grant RP160015) which does not necessary endorse this study. The content is solely the responsibility of the authors and does not necessarily represent the official views of the Cancer Prevention and Research Institute of Texas. All other authors received no specific funding for this work.

References

- Patton GC, Coffey C, Sawyer SM, et al. Global patterns of mortality in young people: a systematic analysis of population health data. *Lancet* 2009; **374**: 881–92.
- Patton GC, Sawyer SM, Santelli JS, et al. Our future: a *Lancet* commission on adolescent health and wellbeing. *Lancet* 2016; **387**: 2423–78.
- Patton GC, Olsson CA, Skirbekk V, et al. Adolescence and the next generation. *Nature* 2018; **554**: 458–66.
- Vollset SE, Goren E, Yuan C-W, et al. Fertility, mortality, migration, and population scenarios for 195 countries and territories from 2017 to 2100: a forecasting analysis for the Global Burden of Disease Study. *Lancet* 2020; **396**: 1285–306.
- UN. Survive, thrive, transform. The Global Strategy for Women's, Children's and Adolescent's Health. New York, NY: United Nations; 2015.
- Boerma T, Requejo J, Victora CG, et al. Countdown to 2030: tracking progress towards universal coverage for reproductive, maternal, newborn, and child health. *Lancet* 2018; **391**: 1538–48.
- Claeson M. The Global Financing Facility—towards a new way of financing for development. *Lancet* 2017; **389**: 1588–92.
- Azzopardi PS, Hearps SJC, Francis KL, et al. Progress in adolescent health and wellbeing: tracking 12 headline indicators for 195 countries and territories, 1990–2016. *Lancet* 2019; **393**: 1101–18.
- The Lancet Child & Adolescent Health. Universal health coverage and the forgotten generation. *Lancet Child Adolesc Health* 2019; **3**: 749.
- Verity R, Okell LC, Dorigatti I, et al. Estimates of the severity of coronavirus disease 2019: a model-based analysis. *Lancet Infect Dis* 2020; **20**: 669–77.
- The Lancet Child & Adolescent Health. Growing up in the shadow of COVID-19. *Lancet Child Adolesc Health* 2020; **4**: 853.
- Vos T, Lim SS, Abbafati C, et al. Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* 2020; **396**: 1204–22.
- Patton GC, Azzopardi P. Missing in the middle: measuring a million deaths annually in children aged 5–14 years. *Lancet Glob Health* 2018; **6**: e1048–49.
- Masquelier B, Hug L, Sharrow D, et al. Global, regional, and national mortality trends in older children and young adolescents (5–14 years) from 1990 to 2016: an analysis of empirical data. *Lancet Glob Health* 2018; **6**: e1087–99.
- Guthold R, Moller A-B, Adebayo E, et al. Priority areas for adolescent health measurement. *J Adolesc Health* 2021; **68**: 888–98.
- Viner RM, Coffey C, Mathers C, et al. 50-year mortality trends in children and young people: a study of 50 low-income, middle-income, and high-income countries. *Lancet* 2011; **377**: 1162–74.
- Mokdad AH, Forouzanfar MH, Daoud F, et al. Global burden of diseases, injuries, and risk factors for young people's health during 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2016; **387**: 2383–401.
- WHO. Indicator and monitoring framework for the Global Strategy for Women's, Children's and Adolescents' health (2016–2030). Geneva: World Health Organization, 2016.
- WHO. World health statistics overview 2019: monitoring health for the SDGs, sustainable development goals. Geneva: World Health Organization, 2019.
- UN Department of Economic and Social Affairs. World population prospects 2019 highlights. New York, NY: United Nations, 2019.
- WHO. WHO methods and data sources for global burden of disease estimates 2000–2016. Geneva: World Health Organization, 2018.
- United States Census Bureau. International database. 2021. <https://www.census.gov/programs-surveys/international-programs/about/idb.html> (accessed Jan 29 2021).
- UNICEF. Levels and trends in child mortality 2020. New York, NY: United Nations Inter-agency Group for Child Mortality Estimation, 2020.
- Sawyer SM, Azzopardi PS, Wickremarathne D, Patton GC. The age of adolescence. *Lancet Child Adolesc Health* 2018; **2**: 223–28.
- Stevens GA, Alkema L, Black RE, et al. Guidelines for Accurate and Transparent Health Estimates Reporting: the GATHER statement. *PLoS Med* 2016; **13**: e1002056.
- Dicker D, Nguyen G, Abate D, et al. Global, regional, and national age-sex-specific mortality and life expectancy, 1950–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* 2018; **392**: 1684–735.
- Reiner RC, Olsen HE, Ikeda CT, et al. Diseases, injuries, and risk factors in child and adolescent health, 1990 to 2017: findings from the Global Burden of Diseases, Injuries, and Risk Factors 2017 Study. *JAMA Pediatr* 2019; **173**: e190337.
- Salam RA, Das JK, Lassi ZS, Bhutta ZA. Adolescent health interventions: conclusions, evidence gaps, and research priorities. *J Adolesc Health* 2016; **59**: S88–92.
- WHO. Global report on drowning: preventing a leading killer. Geneva: World Health Organization, 2014.
- WHO. Global status report on road safety 2018. Geneva: World Health Organization, 2018.
- Salam RA, Arshad A, Das JK, et al. Interventions to prevent unintentional injuries among adolescents: a systematic review and meta-analysis. *J Adolesc Health* 2016; **59**: S76–87.
- Gore FM, Bloem PJN, Patton GC, et al. Global burden of disease in young people aged 10–24 years: a systematic analysis. *Lancet* 2011; **377**: 2093–102.
- Das JK, Salam RA, Lassi ZS, et al. Interventions for adolescent mental health: an overview of systematic reviews. *J Adolesc Health* 2016; **59**: S49–60.
- Patel V, Saxena S, Lund C, et al. The *Lancet* Commission on global mental health and sustainable development. *Lancet* 2018; **392**: 1553–98.
- Anestis MD, Selby EA, Butterworth SE. Rising longitudinal trajectories in suicide rates: the role of firearm suicide rates and firearm legislation. *Prev Med* 2017; **100**: 159–66.
- Gunnell D, Fernando R, Hewagama M, Priyangika WD, Konradsen F, Eddleston M. The impact of pesticide regulations on suicide in Sri Lanka. *Int J Epidemiol* 2007; **36**: 1235–42.
- The Lancet Child & Adolescent Health. Children and youth are crucial in the global AIDS response. *Lancet Child Adolesc Health* 2018; **2**: 621.
- Salam RA, Faqqah A, Sajjad N, et al. Improving adolescent sexual and reproductive health: a systematic review of potential interventions. *J Adolesc Health* 2016; **59**: S11–28.
- Sphere Association. The Sphere Handbook: humanitarian charter and minimum standards in humanitarian response, 4th edn. Geneva: Sphere, 2018.

- 40 Lee ACK, Booth A, Challen K, Gardois P, Goodacre S. Disaster management in low- and middle-income countries: scoping review of the evidence base. *Emerg Med J* 2014; **31**: e78–83.
- 41 Viner RM, Hargreaves DS, Ward J, Bonell C, Mokdad AH, Patton G. The health benefits of secondary education in adolescents and young adults: an international analysis in 186 low-, middle- and high-income countries from 1990 to 2013. *SSM Popul Health* 2017; **3**: 162–71.
- 42 Nambiar D, Razzak J, Afsana K, et al. Mental illness and injuries: emerging health challenges of urbanisation in south Asia. *BMJ* 2017; **357**: j1126.
- 43 Ward JL, Viner RM. The impact of income inequality and national wealth on child and adolescent mortality in low and middle-income countries. *BMC Public Health* 2017; **17**: 429.
- 44 Law TH. Factors associated with the relationship between non-fatal road injuries and economic growth. *Transp Policy* 2015; **42**: 166–72.
- 45 Jiang B, Liang S, Peng Z-R, et al. Transport and public health in China: the road to a healthy future. *Lancet* 2017; **390**: 1781–91.
- 46 Mikkelsen L, Phillips DE, AbouZahr C, et al. A global assessment of civil registration and vital statistics systems: monitoring data quality and progress. *Lancet* 2015; **386**: 1395–406.
- 47 Rigby M, Deshpande S, Blair M. Credibility in published data sources. *Lancet* 2019; **393**: 225–26.
- 48 Barber RM, Fullman N, Sorensen RJ, et al. Healthcare Access and Quality Index based on mortality from causes amenable to personal health care in 195 countries and territories, 1990–2015: a novel analysis from the Global Burden of Disease Study 2015. *Lancet* 2017; **390**: 231–66.
- 49 Viner RM, Mytton OT, Bonell C, et al. Susceptibility to SARS-CoV-2 infection among children and adolescents compared with adults: a systematic review and meta-analysis. *JAMA Pediatr* 2021; **175**: 143–56.
- 50 Clark H, Ghebreyesus TA, Albrechtsen A-B, et al. Uniting for adolescents in covid-19 and beyond. *BMJ* 2021; **372**: n719.
- 51 Odd D, Sleaf V, Appleby L, Gunnell D, Luyt K. Child suicide rates during the COVID-19 pandemic in England: real-time surveillance. Bristol: National Child Mortality Database, 2020.
- 52 Pierce M, Hope H, Ford T, et al. Mental health before and during the COVID-19 pandemic: a longitudinal probability sample survey of the UK population. *Lancet Psychiatry* 2020; **7**: 883–92.
- 53 Watts N, Amann M, Arnell N, et al. The 2020 report of The Lancet Countdown on health and climate change: responding to converging crises. *Lancet* 2021; **397**: 129–70.
- 54 Lim SS, Updike RL, Kaldjian AS, et al. Measuring human capital: a systematic analysis of 195 countries and territories, 1990–2016. *Lancet* 2018; **392**: 1217–34.