

Longitudinal analysis of ANA in the Systemic Lupus International Collaborating Clinics (SLICC) inception cohort

Choi, May; Clarke, Ann; Urowitz, Murray; Hanly, John; St Pierre, Yvan; Gordon, Caroline; Bae, Sang-Cheol ; Romero-Diaz, Juanita; Sanchez-Guerrero, Jorge; Bernatsky, Sasha; Wallace, Daniel; Isenberg, David; Rahman, Anisur; Merrill, Joan; Paul, R; Gladman, Dafna; Bruce, Ian; Petri, Michelle; Ginzler, Ellen; Dooley, Mary Anne

DOI:

[10.1136/annrheumdis-2022-222168](https://doi.org/10.1136/annrheumdis-2022-222168)

License:

Creative Commons: Attribution-NonCommercial (CC BY-NC)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Choi, M, Clarke, A, Urowitz, M, Hanly, J, St Pierre, Y, Gordon, C, Bae, S-C, Romero-Diaz, J, Sanchez-Guerrero, J, Bernatsky, S, Wallace, D, Isenberg, D, Rahman, A, Merrill, J, Paul, R, Gladman, D, Bruce, I, Petri, M, Ginzler, E, Dooley, MA, Ramsey-Goldman, R, Manzi, S, Jonsen, A, Alarcon, G, van Vollenhoven, R, Aranow, C, Mackay, M, Ruiz-Irastorza, G, Lim, S, Inanc, M, Kalunian, K, Jacobsen, S, Peschken, CA, Kamen, D, Askanase, A, Buyon, J, Costenbader, KH & Fritzler, MJ 2022, 'Longitudinal analysis of ANA in the Systemic Lupus International Collaborating Clinics (SLICC) inception cohort', *Annals of the Rheumatic Diseases*, vol. 81, no. 8, pp. 1143-1150. <https://doi.org/10.1136/annrheumdis-2022-222168>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

**Longitudinal Analysis of ANA in the Systemic Lupus
 International Collaborating Clinics (SLICC) Inception Cohort**

Journal:	<i>Annals of the Rheumatic Diseases</i>
Manuscript ID	annrheumdis-2022-222168.R1
Article Type:	Extended report
Date Submitted by the Author:	15-Feb-2022
Complete List of Authors:	<p>Choi, May; University of Calgary Cumming School of Medicine Clarke, Ann; University of Calgary Cumming School of Medicine, Division of Rheumatology Urowitz, Murray; University of Toronto, Lupus Clinic, Center for Prognosis Studies in the Rheumatic Diseases, Toronto Western Hospital Hanly, John; Queen Elizabeth II Health Sciences Centre and Dalhousie University, Division of Rheumatology, Department of Medicine and Department of Pathology St-Pierre, Yvan; Research Institute of the McGill University Health Center Gordon, Caroline; Birmingham University Medical School, Rheumatology Research Group Bae, Sang-Cheol; Hanyang University Hospital for Rheumatic Diseases, Rheumatology Romero-Diaz, Juanita; Instituto Nacional de Ciencias Medicas y Nutricion Salvador Zubiran, Immunology and Rheumatology Sanchez-Guerrero, Jorge; Toronto Western Hospital, Center for Prognosis Studies in the Rheumatic Diseases Bernatsky, Sasha; Research Institute of the McGill University Health Centre, Clinical Epidemiology; McGill University Health Centre, Rheumatology Wallace, Daniel; Cedars-Sinai Medical Center, Division of Rheumatology; 3360, Isenberg, David; Centre for Rheumatology Research, Division of Medicine UCL , Division of Medicine, UCL Rahman, Anisur; University College London, Rheumatology Merrill, Joan; Oklahoma Medical Research Foundation Fortin, Paul R.; Centre Hospitalier de l'Université Laval , Medicine - Rheumatology Gladman, Dafna; University of Toronto, Bruce, Ian; University of Manchester, arc Epidemiology Unit Petri, Michelle; Johns Hopkins University, Rheumatology Ginzler, Ellen; SUNY Downstate Medical Center, Medicine/Rheumatology Dooley, Mary Anne; University of North Carolina at Chapel Hill, Medicine Ramsey-Goldman, Rosalind; Northwestern University, Medicine/Rheumatology Manzi, Susan; Allegheny Health Network, Lupus Center of Excellence Jönsen, Andreas; Lund University, Clinical Sciences, Alarcón, Graciela; School of Medicine, The University of Alabama, Department of Medicine, Division of Clinical Immunology and</p>

	<p>Rheumatology van Vollenhoven, Ronald; Amsterdam Rheumatology and Immunology Center, Department of Rheumatology Aranow, Cynthia; The Feinstein Institute for Medical Research, Mackay, Meggan; Feinstein Institute for Medical Research Ruiz-Irastorza, Guillermo; HOSPITAL UNIVERSITARIO CRUCES, Autoimmune Diseases Unit Lim, Sam; Emory University, School of Medicine Inanc, Murat; Istanbul University, Istanbul Medical Faculty, Dept of Internal Medicine, Div of Rheumatology Kalunian , Ken; University of California, Division of Rheumatology, Allergy and Immunology Jacobsen, Søren; Rigshospitalet, Copenhagen University Hospital, Department of Rheumatology - 4242 Peschken, Christine; University of Manitoba Kamen, Diane; Medical University of South Carolina, Askanase, Anca; Columbia University Medical Center Buyon, Jill; New York University School of Medicine, Costenbader, Karen; Brigham and Women's Hospital, Department of Medicine, Div of Rheumatology, Inflammation and Immunity; Harvard Medical School, Medicine Fritzler, Marvin; Health Sciences Centre, Medicine</p>
Keywords:	Systemic Lupus Erythematosus, Autoantibodies, Autoimmunity





I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

Longitudinal Analysis of ANA in the Systemic Lupus International Collaborating Clinics (SLICC) Inception Cohort

¹May Y Choi, ¹Ann E Clarke, ²Murray B Urowitz, ³John G Hanly, ⁴Yvan St. Pierre, ⁵Caroline Gordon, ⁶Sang-Cheol Bae, ⁷Juanita Romero-Diaz, ⁸Jorge Sanchez-Guerrero, ⁹Sasha Bernatsky, ¹⁰Daniel J Wallace, ¹¹David A Isenberg, ¹¹Anisur Rahman, ¹²Joan T Merrill, ¹³Paul R Fortin, ²Dafna D Gladman, ¹⁴Ian N Bruce, ¹⁵Michelle Petri, ¹⁶Ellen M Ginzler, ¹⁷Mary Anne Dooley, ¹⁸Rosalind Ramsey-Goldman, ¹⁹Susan Manzi, ²⁰Andreas Jönsen, ²¹Graciela S Alarcón, ²²Ronald F van Vollenhoven, ²³Cynthia Aranow, ²³Meggan Mackay, ²⁴Guillermo Ruiz-Irastorza, ²⁵Sam Lim, ²⁶Murat Inanc, ²⁷Kenneth C Kalunian, ²⁸Søren Jacobsen, ²⁹Christine A Peschken, ³⁰Diane L Kamen, ³¹Anca Askanase, ³²Jill Buyon, ³³Karen H Costenbader, ¹Marvin J. Fritzler

¹University of Calgary, Cumming School of Medicine, Calgary, Alberta, Canada
²Lupus Program, Centre for Prognosis Studies in The Rheumatic Disease and Schroeder Arthritis Institute, Krembil Research Institute, Toronto Western Hospital, University of Toronto, Toronto, Ontario, Canada;
³Division of Rheumatology, Department of Medicine and Department of Pathology, Queen Elizabeth II Health Sciences Centre and Dalhousie University, Halifax, Nova Scotia, Canada;
⁴Research Institute of the McGill University Health Centre; Montreal, Quebec, Canada
⁵Rheumatology Research Group, Institute of Inflammation and Ageing, College of Medical and Dental Sciences, University of Birmingham, Birmingham, United Kingdom;
⁶Department of Rheumatology, Hanyang University Hospital for Rheumatic Diseases and Hanyang University Institute for Rheumatology, Seoul, Korea;
⁷Instituto Nacional de Ciencias Médicas y Nutrición, Mexico City, Mexico;

- 24 ⁸Mount Sinai Hospital and University Health Network, University of Toronto, Canada,
- 25 ⁹Divisions of Rheumatology and Clinical Epidemiology, McGill University Health Centre;
- 26 ¹⁰Cedars-Sinai/David Geffen School of Medicine at UCLA, Los Angeles, California, USA;
- 27 ¹¹Centre for Rheumatology, Department of Medicine, University College London, United
Kingdom;
- 29 ¹²Department of Clinical Pharmacology, Oklahoma Medical Research Foundation, Oklahoma
City, Oklahoma, USA;
- 31 ¹³Division of Rheumatology, CHU de Québec - Université Laval, Québec City, Canada;
- 32 ¹⁴NIHR Manchester Biomedical Research Centre, Manchester University Hospitals NHS
Foundation Trust, Manchester Academic Health Science Centre, Manchester, Greater
Manchester, UK and Centre for Epidemiology Versus Arthritis, Faculty of Biology, Medicine
and Health, The University of Manchester, UK;
- 36 ¹⁵Division of Rheumatology, Johns Hopkins University School of Medicine, Baltimore,
Maryland, USA;
- 38 ¹⁶Department of Medicine, SUNY Downstate Medical Center, Brooklyn, New York, USA;
- 39 ¹⁷Thurston Arthritis Research Center, University of North Carolina, Chapel Hill, North Carolina,
USA;
- 41 ¹⁸Northwestern University and Feinberg School of Medicine, Chicago, Illinois, USA;
- 42 ¹⁹Allegheny Health Network, Pittsburgh, Pennsylvania, USA;
- 43 ²⁰Lund University, Lund, Sweden
- 44 ²¹Department of Medicine, University of Alabama at Birmingham, Birmingham, Alabama, USA;
- 45 ²²University of Amsterdam, Rheumatology & Immunology Center, Amsterdam, Noord-Holland,
The Netherlands;

1
2
3 47 ²³Feinstein Institute for Medical Research, Manhasset, New York, USA;
4
5 48 ²⁴Autoimmune Diseases Research Unit, Department of Internal Medicine, BioCruces Health
6
7
8 49 Research Institute, Hospital Universitario Cruces, University of the Basque Country, Barakaldo,
9
10 50 Spain;
11
12 51 ²⁵Emory University School of Medicine, Division of Rheumatology, Atlanta, Georgia, USA;
13
14 52 ²⁶Division of Rheumatology, Department of Internal Medicine, Istanbul Medical Faculty,
15
16
17 53 Istanbul University, Istanbul, Turkey;
18
19 54 ²⁷University of California San Diego School of Medicine, La Jolla, California, USA;
20
21 55 ²⁸Department of Rheumatology, Rigshospitalet, Copenhagen University Hospital, , Copenhagen,
22
23
24 56 Denmark;
25
26 57 ²⁹University of Manitoba, Winnipeg, Manitoba, Canada;
27
28 58 ³⁰Medical University of South Carolina, Charleston, South Carolina, USA;
29
30 59 ³¹Hospital for Joint Diseases, New York University, Seligman Centre for Advanced
31
32
33 60 Therapeutics, New York New York, USA;
34
35 61 ³²New York University School of Medicine, New York, NY
36
37 62
38 63 ³³Division of Rheumatology, Inflammation and Immunity, Department of Medicine, Brigham
39
40 64 and Women’s Hospital, Harvard Medical School, Boston, Massachusetts, USA
41
42
43 65

44 66	Manuscript Word Count (max 3000):	3615
45 67	Abstract Word Count (max 250):	250
46 68	Number of Tables:	4
47 69	Number of Figures	2
48 70	Number of Supplemental Tables:	5

71 **Number of Supplemental Figures: 1**

72

73 **Corresponding Author and Reprint Requests:**

74 May Choi, MD, MPH, FRCPC

75 Division of Rheumatology, Department of Medicine

76 Cumming School of Medicine, University of Calgary

77 3230 Hospital Drive NW

78 Calgary, NW T2N 4Z6

79 Email: may.choi@ucalgary.ca

80 Tel: 1-403-955-8733

81

82 **Competing Interests:**

83 Dr. May Choi has received consulting fees from Janssen and MitogenDx (less than \$10,000).

84 Dr. Ann Clarke has received consulting fees from AstraZeneca, BristolMyersSquibb, and Glaxo

85 Smith Kline (less than \$10,000 each).

86 Dr. Costenbader has consulted for or collaborated on research projects with Janssen, Glaxo

87 Smith Kline, Exagen Diagnostics, Eli Lilly, Merck Serono, Astra Zeneca and Neutrolis (less than

88 \$10,000 each).

89 Dr. Gordon has received consulting fees, speaking fees, and/or honoraria from AstraZeneca,

90 Abbvie, Amgen, UCB, GlaxoSmithKline, Merck Serono and BMS (less than \$10,000 each) and

91 grants from UCB. Grants from UCB were not to Dr. Gordon but to Sandwell and West

92 Birmingham Hospitals NHS Trust.

1
2
3 93 Dr. Dafna Gladman received consulting fees, speaking fees, and/or honoraria from
4
5 94 GlaxoSmithKline (less than \$10,000).
6
7
8 95 Dr. Bruce has received consulting fees, speaking fees, and/or honoraria from Eli Lilly, UCB,
9
10 96 Roche, Merck Serono, MedImmune (less than \$10,000 each) and grants from UCB, Genzyme
11
12 97 Sanofi, and GlaxoSmithKline.
13
14
15 98 Dr. Ginzler has paid consultation with investment analysts Guidepoint Global Gerson Lerman
16
17 99 Group.
18
19 100 Dr. Kalunian has received grants from UCB, Human Genome Sciences/GlaxoSmithKline,
20
21 101 Takeda, Ablynx, Bristol-Myers Squibb, Pfizer, and Kyowa Hakko Kirin, and has received
22
23 102 consulting fees from Exagen Diagnostics, Genentech, Eli Lilly, Bristol-Myers Squibb, and
24
25 103 Anthera (less than \$10,000 each).
26
27
28 104 Dr. Fritzler is Director of Mitogen Diagnostics Corporation (Calgary, AB Canada) and a
29
30 105 consultant to Werfen International (Barcelona, Spain), Grifols (Barcelona, Spain), Janssen
31
32 106 Pharmaceuticals of Johnson & Johnson and Alexion Canada (less than \$10,000 each).
33
34
35 107
36
37
38 108 The remainder of the authors have no disclosures.
39
40 109

41
42 110 **Contributorship:**
43
44 111 All authors were involved in the concept and design, data analysis and interpretation, and editing
45
46 112 for intellectual content. MC, AC, MF and KC were involved in manuscript drafting.
47
48

49 113
50
51 114 **Acknowledgements:**
52
53
54
55
56
57
58
59
60

1
2
3 115 The authors are grateful for the technical assistance of Ms. Haiyan Hou, who conducted the
4
5 116 laboratory testing, Ms. Meifeng Zhang (MitogenDx, University of Calgary), Ms. Katherine
6
7 117 Buhler, Ms. Chynace Lambalgen, and the SLICC nurse coordinators.
8
9

10 118

11
12 119 **Grant Support:**
13

14
15 120 The cost of the immunoassay supplies and labor were supported by the Lupus Foundation of
16
17 121 America and MitogenDx.
18

19 122 Dr. May Choi is supported by the Lupus Foundation of America Gary S. Gilkeson Career
20
21 123 Development Award and research gifts in kind from MitogenDx (Calgary, Canada).
22
23

24 124 Dr. Clarke holds The Arthritis Society Research Chair in Rheumatic Diseases at the University
25
26 125 of Calgary.
27

28 126 Dr. Costenbader is supported by NIH K24 AR066109.
29

30
31 127 Dr. Hanly's work was supported by the Canadian Institutes of Health Research (research grant
32
33 128 MOP-88526).
34

35 129 Dr. Caroline Gordon's work was supported by Lupus UK, Sandwell and West Birmingham
36
37 130 Hospitals NHS Trust and the NIHR /Wellcome Trust Clinical Research Facility in Birmingham.
38

39
40 131 Dr. Sang-Cheol Bae's work was supported by the Korea Healthcare technology R & D project,
41
42 132 Ministry for Health and Welfare, Republic of Korea (A120404).
43

44 133 The Montreal General Hospital Lupus Clinic is partially supported by the Singer Family Fund
45
46 134 for Lupus Research.
47

48
49 135 Dr. Rahman and Dr. Isenberg are supported by the National Institute for Health Research
50
51 136 University College London Hospitals Biomedical Research Centre.
52

53
54 137 The Hopkins Lupus Cohort is supported by NIH Grants AR043727 and AR069572
55
56
57
58
59
60

1
2
3 138 Dr. Paul R. Fortin presently holds a tier 1 Canada Research Chair on Systemic Autoimmune
4
5 139 Rheumatic Diseases at Université Laval, and part of this work was done while he was still holding
6
7 140 a Distinguished Senior Investigator of The Arthritis Society.
8
9
10 141 Dr. Bruce is an NIHR Senior Investigator Emeritus and is funded by the National Institute for
11
12 142 Health Research Manchester Biomedical Research Centre and the NIHR/Wellcome Trust
13
14 143 Manchester Clinical Research Facility. The views expressed in this publication are those of the
15
16 144 author(s) and not necessarily those of the NHS, the National Institute for Health Research or the
17
18 145 Department of Health.
19
20
21 146 Dr. Mary Anne Dooley's work was supported by the NIH grant RR00046.
22
23
24 147 Dr. Ramsey-Goldman's work was supported by the NIH (grants 1U54TR001353 formerly
25
26 148 8UL1TR000150 and UL-1RR-025741, K24-AR-02318, and P60AR064464 formerly P60-AR-
27
28 149 48098).
29
30
31 150 Dr. Susan Manzi is supported by grants R01 AR046588 and K24 AR002213
32
33 151 Dr. Ruiz-Irastorza is supported by the Department of Education, Universities and Research of the
34
35 152 Basque Government.
36
37
38 153 Dr. Soren Jacobsen is supported by the Danish Rheumatism Association (A1028) and the Novo
39
40 154 Nordisk Foundation (A05990).
41
42
43 155

44
45 156 **Ethical Approval Information**

46
47 157 The study was approved by the Institutional Review Board at each participating site.
48
49 158

50
51 159 **Data Sharing Statement**
52
53
54
55
56
57
58
59
60

160 All data relevant to the study are included in the article or uploaded as supplementary
161 information.

162

163 **Patient and Public Involvement statement**

164

165 Patients or the public were not involved in the design, or conduct, or reporting, or dissemination
166 plans of our research.

ABSTRACT

Objectives: A perception derived from cross-sectional studies of small SLE cohorts is that there is a marked discrepancy between antinuclear antibody (ANA) assays, which impacts on clinician’s approach to diagnosis and follow-up. We compared three ANA assays in a longitudinal analysis of a large international incident SLE cohort retested regularly and followed for five years.

Methods: Demographic, clinical, and serological data was from 805 SLE patients at enrolment, year 3 and 5. Two HEp-2 indirect immunofluorescence assays (IFA1, IFA2), an enzyme-linked immunosorbent assay (ELISA), and SLE-related autoantibodies were performed in one central laboratory. Frequencies of positivity, titres/units, and IFA patterns were compared using McNemar, Wilcoxon, and kappa statistics, respectively.

Results: At enrolment, ANA positivity ($\geq 1:80$) was 96.1% by IFA1 (median titre 1:1280 [IQR 1:640-1:5120]), 98.3% by IFA2 (1:2560 [IQR 1:640-1:5120]), and 96.6% by ELISA (176.3AU [IQR 106.4-203.5]). At least one ANA assay was positive for 99.6% of patients at enrolment. At year 5, ANA positivity by IFAs (IFA1 95.2%; IFA2 98.9%) remained high, while there was a decrease in ELISA positivity (91.3%, $p<0.001$). Overall, there was $>91\%$ agreement in ANA positivity at all time points and $\geq 71\%$ agreement in IFA patterns between IFA1 and IFA2.

Conclusion: In recent-onset SLE, three ANA assays demonstrated commutability with a high proportion of positivity and titres/units. However, over five years follow-up, there was modest variation in ANA assay performance. In clinical situations where the SLE diagnosis is being considered, a negative test by either the ELISA or HEp-2 IFA may require reflex testing.

Keywords: Antinuclear antibodies, Systemic Lupus Erythematosus, longitudinal, performance, immunoassays, ELISA

INTRODUCTION

Antinuclear antibody (ANA) testing has an integral approach to accurately diagnose and classify SLE (1). A systematic literature review and meta-regression of indirect immunofluorescence assays (IFA) reported high sensitivity (97.8%) for SLE diagnosis at a titer of $\geq 1:80$ (2). This presaged the decision to include a positive ANA at that titer on HEp-2 cell IFA “or an equivalent positive test on other diagnostic platforms” occurring at least once as an entry criterion for the 2019 European League Against Rheumatism/American College of Rheumatology (EULAR/ACR) SLE Classification Criteria (3, 4).

Previous longitudinal examinations of ANA and SLE-related autoantibodies suggest that a patient’s ANA status can change from positive to within the normal range and vice versa during the disease course (2, 5-16). However, these studies have typically been limited to small, single center cohorts with incomplete disease characterization, short follow-up, and/or using outdated assays with conflicting results. The factors influencing changes in ANA have also not been thoroughly studied. Taken together, this has left clinicians with uncertainty about the value and interpretation of ANA testing in making a diagnosis of, or classifying, SLE. In addition, the clinically actionable value of repeat ANA testing once a diagnosis of SLE is established requires clarification (17, 18).

Much of the confusion and debate on the clinical utility of ANA testing in SLE is related to reported variations in HEp-2 IFA assay performance in cross-sectional cohorts (19-22), and some have questioned whether the ANA IFA should continue to be the “gold standard” screening test (23-25). For instance, in a cross-sectional study, Pisetsky et al. tested the same sera using

1
2
3 215 different ANA assays (e.g., IFA, enzyme-linked immunosorbent assay [ELISA], and multiplex
4
5 216 bead assay) (21) and reported that the frequency of an ANA test within normal reference range in
6
7 217 SLE patients with disease duration ranging from 0.1 to 33.4 years varied from 4.9%–22.3%.
8
9 218 Further, it has been proposed that the IFA could be replaced or complemented by newer
10
11 219 generation solid phase multi-analyte immunoassays (SPMAI) such as ELISA and/or addressable
12
13 220 laser bead immunoassays (ALBIA) (24-26). A recent systematic review and meta-regression
14
15 221 analysis of ANA testing in >13,000 SLE patients with disease duration ranging from 0–17 years
16
17 222 reported that only ~2.5% of these patients had an IFA ANA <1:80 (2), although a higher
18
19 223 prevalence of ANA within the normal reference range has been reported in other cohorts
20
21 224 including the Systemic Lupus International Collaborating Clinics (SLICC) Inception Cohort
22
23 225 (6.2% were <1:160 at inception) (27).
24
25
26
27
28
29
30

31 227 The primary goal of this study was to gain a more thorough understanding of ANA detection and
32
33 228 its clinical value by comparing the performance of three currently available ANA assays in a
34
35 229 longitudinal analysis (at least 5 years) of a large multinational SLE inception cohort.
36
37
38
39

40 231 **METHODS**

41
42 232 *Study Population*

43
44 233 Between 1999 and 2011, SLICC (<https://sliccgroup.org>) (28) enrolled 1827 patients fulfilling the
45
46 234 1997 Updated ACR SLE Classification Criteria for definite SLE (29) within 15 months of
47
48 235 diagnosis from 31 medical centres in 11 countries. Sera, clinical and demographic data were
49
50 236 collected at enrolment and annually thereafter. Of the 1827 patients, 1432 (78.4%) were followed
51
52
53 237 for ≥4 years; of these 1432 patients, we included the 805 patients who provided an enrolment
54
55
56
57
58
59
60

and two additional serum samples within five years of enrolment, with the third sample being ≥ 4 years after enrolment. The study was approved by the Institutional Review Board at each participating site. Permission from the SLICC Biological Material and Data Utilization Committee was obtained to access the required data and biobanked serum samples.

ANA and Autoantibody Testing

Aliquots of sera were obtained from the 805 patients in the SLICC Inception Cohort at three time points: 1) enrolment (sample #1), 2) two to four years after enrolment (sample #2), and 3) four to 10 years after enrolment (sample #3). Hereafter, samples #1 – 3 are referred to as enrolment, year 3, and year 5, respectively. Samples were stored at -80°C until required for immunoassays and analyzed centrally at MitogenDx Laboratory (Calgary, Canada). Three US Food and Drug Administration (FDA)-approved and Conformité Européenne (CE) marked ANA tests were used, including two HEp-2 IFA, IFA1 (Bio-Rad Laboratories, Hercules, USA) and IFA2 (NovaLite, Werfen, San Diego, USA), and an ELISA (Werfen, San Diego, USA). In accordance with the manufacturers' directions, a positive test was defined as a titer of $\geq 1:80$ for IFA1 and IFA2 (titre $< 1:80$ is considered normal range) and ≥ 20 absorbance units (AU) for ELISA. IFA1, IFA2, and ELISA were tested on the full patient cohort ($n=805$) sera from all three time points. IFA results (titres and patterns) were initially read by an automated digital IFA microscope and then checked manually by a technologist with 30 years of experience. ANA IFA patterns were classified according to the new International Consensus on ANA Patterns recommendations (<http://www.anapatterns.org/index.php>) (30). Quality control was performed by repeating all ANA results that were within the normal range and a random selection of ANA-positive samples

to ensure inter-test reliability. SLE-related autoantibodies (**Supplemental Table 1**) were also performed on each patient at enrolment, year 3 and 5.

Clinically Defined Samples

Demographic and clinical data (**Supplemental Table 2**) at enrolment included age, sex, disease duration, race/ethnicity, nephritis (fulfilling the ACR criterion for renal disease or based on a renal biopsy), ACR Classification Criteria, Systemic Lupus Erythematosus Disease Activity Index – 2000 (SLEDAI-2K), SLICC/ACR Damage Index (SDI), and medication use (current and ever use of glucocorticoids, antimalarials, and immunosuppressives, including biologics). We also collected longitudinal data on nephritis, SLEDAI-2K, SDI, and medications.

Statistical analysis

Demographic, clinical, and serological characteristics were described using summary statistics. Changes over time in demographic and clinical features were described using differences in means or proportions, with 95% confidence intervals (CI). As our analysis used a subgroup of the larger SLICC cohort based on sera availability, we compared the enrolment characteristics of the 805 patients included in this study with the 627 patients who were followed for ≥ 4 years but were not included as three serial serum samples were unavailable. We also compared the characteristics of the 781 patients providing the third serum sample 4-7 years after enrolment with the 24 patients providing the third serum sample 8-10 years after enrolment.

We assessed the frequency of ANA positivity and titre at each time point. Using the paired McNemar’s test, we calculated changes in ANA positivity between enrolment and year 5 for

each test and the inter-test agreement in ANA positivity between tests at each time point. A histogram with a curve of best fit line was used to plot the changes in distribution of titres and units over time were compared using the Wilcoxon signed rank test for paired data. We examined the frequency of each ANA pattern and how many patients retained their HEp-2 IFA pattern over the three serial samples. ANA patterns were further categorized into three groups: 1) isolated nuclear (AC 1-14, 29), 2) isolated cytoplasmic and/or mitotic (CMP, AC 15-28), and 3) mixed nuclear and CMP patterns. Agreement between IFA1 and IFA2 ANA titres and patterns was assessed using the weighted and unweighted kappa (κ) statistic, respectively. Established SLE-related autoantibody profiles of patients with an ANA result within the normal range on IFA1, IFA2, or ELISA alone, on two of three assays, and on all three assays at enrolment and year 5 were examined to understand which autoantibodies were not being captured by the ANA screening assays. Statistical analysis was performed using Stata 15.1 (StataCorp, College Station, TX, USA).

RESULTS

Study Population

Eight hundred and five SLE patients were included. The mean time from disease diagnosis to enrolment was 0.58 years (standard deviation [SD] 0.49); the mean time between the enrolment and the year 3 sample was 2.8 years (SD 0.8) and between the enrolment and the year 5 sample was 5.0 years (SD 1.1). Patients had a mean age at diagnosis of 35.2 years (SD 13.6), 88.7% (714/805) were female and 47.7% (384/805) were of race/ethnicity other than White (**Table 1**). From enrolment to year 5, the prevalence of lupus nephritis increased by 7.7% [95%CI: 5.7%, 9.7%], mean SLEDAI-2K decreased by 2.3 [95%CI: 1.9, 2.7], and mean SDI increased by 0.52

1
2
3 306 [95%CI: 0.43, 0.62]. There were significantly fewer patients on glucocorticoids (69.6% vs
4
5 307 56.8%, difference -12.8% [95%CI: -16.5%, -9.1%]) and more patients on antimalarials (70.1%
6
7 308 vs 79.4%, difference 9.3% [95%CI: 5.9%, 12.7%]) or immunosuppressants (41.0% vs 50.8%,
8
9 309 difference 9.8% [95%CI: 6.1%, 13.5%]). The frequency of most SLE-related autoantibodies
10
11 310 decreased at year 5.
12
13
14
15 311

16
17 312 The enrolment characteristics of the 805 patients included in our study were similar to the 627
18
19 313 patients who provided ≥ 4 years of data but did not have three available serial serum samples
20
21 314 (**Supplemental Table 3**). However, there was a higher proportion of Asian (18.8% (95%CI:
22
23 315 15.3, 22.2) and lower proportion of Hispanic participants (-20.6% (95%CI: -24.5, -16.8) in the
24
25 316 study cohort compared to the cohort not providing serial samples. The enrolment characteristics
26
27 317 of the 781 patients whose year 5 sample was collected between years 4 and 7 were similar to the
28
29 318 24 patients whose year 5 sample was collected between years 8 and 10 (**Supplemental Table 4**).
30
31 319
32
33
34

35 320 *ANA Positivity and Agreement Among Different Assays Over Time*

36
37 321 At enrolment, the frequency of ANA positivity by IFA1, IFA2, and ELISA was high (96.1%
38
39 322 [95%CI: 94.6-97.3%], 98.3% [95%CI: 97.1-99.0%], and 96.6% [95%CI: 95.2-97.7]),
40
41 323 respectively (**Figure 1**) and 99.6% (802/805) of patients had ≥ 1 positive ANA of $\geq 1:80$. An
42
43 324 additional five (0.6% incremental effect), three (0.5%), and two patients (0.4%) at enrolment,
44
45 325 year 3, and year 5 visits, respectively, would be ANA positive on the ELISA, but within the
46
47 326 normal range for both IFA1 and IFA2. There was no significant change in ANA positivity at
48
49 327 enrolment compared to year 5 for IFA1 or IFA2. However, ANA positivity by ELISA decreased
50
51 328 significantly from enrolment to year 5 (difference -5.3% (95%CI: -7.4, -3.3), $p < 0.001$) such that
52
53
54
55
56
57
58
59
60

91.3% (735/805) of patients were positive by year 5. Notably, 1.2% (10/805) of subjects were within the normal range at all three time points by ELISA compared to 0.9% (7/805) by IFA1 and 0.1% (1/805) by IFA2. At all time points, no patients were classified as being within the normal range if all three of the assays were considered.

Overall, the inter-test agreement for positivity between any pair of assays was >91% (**Table 2**). In cases where there was disagreement between IFA1 and IFA2, there was significant asymmetry (McNemar's test) such that most disagreements were due to more patients with an ANA by IFA1 within the normal range and a positive ANA by IFA2 (-IFA1/+IFA2) rather than a positive ANA by IFA1 and an ANA within the normal range by IFA2 (+IFA1/-IFA2) for all three time points (**Supplemental Table 5**). Regarding the disagreements between IFA1 and ELISA, there was no significant asymmetry until year 5 when there were more cases of disagreement due to +IFA1/-ELISA compared to -IFA1/+ELISA. For disagreements between IFA2 and ELISA, there was significant asymmetry across all time points with more cases of +IFA2/-ELISA than -IFA2/+ELISA.

ANA Titres/Units Among Different Assays Over Time

At enrolment, the median ANA titre/unit for IFA1, IFA2, and ELISA were 1:1280 (interquartile range (IQR) 1:640-1:5120), 1:2560 (IQR 1:640-1:5120), and 176.3 AU (IQR 106.4 AU-203.5 AU), respectively (**Figure 2**). The distribution of ANA titres was skewed to the left for all assays at enrolment (higher proportion of patients with very high ANA titres). Only a small proportion had ANA titres of 1:80 to 1:160 at enrolment (IFA1 10.4% [84/805] and IFA2 8.1% [65/805]). The median titres/units at year 5 were significantly lower compared to enrolment for IFA1

1
2
3 352 (1:640 (IQR 1:320-1:2560), paired Wilcoxon signed rank $p<0.0001$, a change in one dilution
4
5 353 step) and ELISA (157.3 CU (IQR 66.14 CU- 200.65 CU), $p<0.0001$)). There was good
6
7 354 agreement between IFA1 and IFA2 titres at enrolment, 84.9% (95%CI: 82.2-87.3) agreement,
8
9 355 $k=0.49$ (95%CI: 0.45-0.53); at year 3, 81.1% (95%CI: 78.2-83.7%) agreement, $k=0.39$ (95%CI:
10
11 356 0.35-0.43%); and at year 5, 82.0% (95%CI: 79.1-84.6%) agreement, $k=0.41$ (95%CI: 0.37-
12
13 357 0.45%).
14
15
16
17 358

19 359 *ANA Patterns Among Different Assays Over Time*

21 360 The most common ANA IFA pattern was an isolated nuclear staining pattern for IFA1 (62.1%-
22
23 361 68.7%) and IFA2 (59.3%-62.1%) at all visits (**Table 3**). The top three individual IFA patterns for
24
25 362 both IFA1 and IFA2 were AC-1 (homogeneous), AC-4 (nuclear fine speckled), and AC-5
26
27 363 (nuclear large speckled) (**Supplemental Figure 1**). There was fair-to-moderate agreement
28
29 364 between IFA1 and IFA2 ANA IFA staining patterns at enrolment, (74.0% [95%CI 70.7-77.0]
30
31 365 agreement, $\kappa=0.46$ [95%CI 0.39-0.53]), year 3, (71.4% [95%CI 68.0-74.6], $\kappa=0.39$ [95%CI
32
33 366 0.33-0.46]), and year 5, (71.0% [95%CI 67.7-74.2], $\kappa=0.39$ [95%CI 0.33-0.46]).
34
35
36
37 367

40 368 *ANA Patients Within the Normal Range and Seroconversion*

42 369 At enrolment and year 5, 8 and 20 patients were within normal range by IFA1 & ELISA, 3 and 4
43
44 370 patients by ELISA & IFA2, and 8 and 6 patients by IFA1 and IFA2 (**Table 4**). When examining
45
46 371 the autoantibody profiles of patients whose ANA were within normal range at enrolment or year
47
48 372 5, depending on the assay 38.7%-53.8% had no detectable SLE-related autoantibodies. Anti-
49
50 373 Ro52/TRIM21 and anti-SSA/Ro60, the former not detectable by HEp-2 IFA and the latter does
51
52 374 not have a clearly established IFA pattern, were the most frequent autoantibodies detected when
53
54
55
56
57
58
59
60

the ANA test was within normal range. Seroconversion from ANA positive to normal range (titre <1:80) from enrolment to year 5 was observed in 4.8% (39/805) of patients using IFA1, 1.1% (9/805) using IF2, and 8.7% (70/805) using ELISA. The median titre of ANA at enrolment prior to seroconversion was low (IFA1 1:160 [IQR 1:80-1:640]), IFA2 1:320 [IQR 1:160-1:2560], and ELISA 61.5 CU [IQR 20-158]). Among those who were originally anti-dsDNA positive at enrolment (n=273), the frequency of ANA positivity was high at enrolment irrespective of the ANA assay (99.3-100.0%). At year 5, frequency of ANA positivity for these same patients, irrespective of their anti-dsDNA status at year 5, declined slightly using for the IFAs (IFA -2.2%, IFA2 -1.1%) and -4.8% for the ELISA (data not shown).

DISCUSSION

To our knowledge, this is the largest longitudinal, multinational study (805 patients and 2415 serum samples) that compared the performance of different ANA assays in a well-characterized inception cohort of SLE patients. Our study was designed to overcome the limitations of prior reports that studied smaller cohorts and were historical and/or cross-sectional in nature. These data are timely given ANA test positivity is an entry criterion for the 2019 EULAR/ACR classification criteria for SLE (31, 32). We found that, regardless of the assay, almost all patients with recent onset SLE (802/805) had a positive ANA at enrolment on ≥ 1 assay, all were ANA-positive on ≥ 1 assay at least once across the five years, and the mean ANA titres/values were high. However, over the five years, some variation between ANA assay performance was detected, including a statistically significant decrease in ELISA ANA positivity and reduction in titres for IFA1 and ELISA.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

398 It has been suggested that the variation in performance between different ANA assays may be
399 related to differences in laboratory techniques, equipment, inter-observer consistency and
400 reagents (25, 33). However, in our study, all ANAs were performed and interpreted at one
401 central laboratory by a highly experienced (30 years of experience) technician. Even after
402 controlling for the impact of inter-laboratory and inter-observer variation, we still identified
403 some significant inter-assay disagreement. Disagreement between ELISA and IFA is likely
404 primarily due to factors intrinsic to the test platforms themselves. Unlike the IFA, the ELISA
405 contains extracts of cell homogenates augmented by purified proteins derived from native and/or
406 synthetic, recombinant sources (34). The composition of the different ELISA ANA preparations
407 is diverse and dependent on the manufacturer as to which key target autoantigen(s) associated
408 with autoimmune diseases are included and at what concentrations (34). ELISAs may also have
409 decreased detection of ANA because of poor autoantibody binding, as some antigens may also
410 bind to other targets in the same mixture, resulting in a masking effect. Furthermore, many
411 autoantibody targets are components of macromolecular complexes where key epitopes may be
412 hidden or masked (34). A thorough study of the affinity and avidity of the various autoantibodies
413 would add useful understanding to the use of ANA ELISAs.

414

415 Prior studies of more established SLE patients reported that as high as 30% have an ANA below
416 the positive threshold (35). Over time, we observed a decrease in ANA positivity with ELISA, a
417 decrease in ANA titres/values with IFA1 and ELISA, and decreased detection of specific
418 autoantibodies. We postulate that factors such as disease activity and medication exposure
419 influence ANA (36-39). However, the extent to which therapeutic interventions can alter ANA
420 production, especially by long-lived plasma cells, remains to be proven, and the expression of

other autoantibodies can occur following diagnosis, attributed to epitope spreading continuing despite therapy(39).

Our study addresses important questions raised about the ANA in the 2019 EULAR/ACR SLE classification criteria (3, 4, 40), which require an “ever positive” ANA of $\geq 1:80$ by HEp-2 IFA or an equivalent test on another platform as an entry criterion for classification. For example, it is important to note that all subjects had at least one positive ANA at the 1:80 threshold over the five years of follow-up. The new criteria also state that a solid phase assay of at least equivalent performance can be used in place of the HEp-2 IFA, although a precise definition of ‘equivalent performance’ was not specified. Our results show that although some inter-assay disagreement exists between these three assays, >91% of recent-onset SLE patients will have a positive ANA using either HEp-2 IFA or ELISA, although titres decreased by year 5 for IFA1 and the ELISA. As expected from previous reports (20, 41), ELISA had the highest proportion of SLE patients with an ANA within the normal $<1:80$ reference range, and therefore, the ELISA used as a screening test may benefit from judicious reflex testing to the HEp-2 IFA. In turn, since the HEp-2 IFA can be negative when the ELISA is positive, the reciprocal reflex approach could be considered.

Importantly, consistent with other studies and emerging recommendations on ANA testing (20, 41), we demonstrated that a combination of two different ANA assays reduced the proportion of SLE patients with ANAs in the normal range; particularly when IFA2 was combined with ELISA. A combination of all three assays resulted in no patients who had an ANA within normal range at enrollment and two subsequent follow-up visits. This helps shed light on the question of

1
2
3 444 the value of ANA testing to follow the clinical course of SLE, but more detailed follow-up
4
5 445 studies evaluating disease activity and flares at follow-up visits in the context of ANA testing are
6
7 446 still required. Health care providers should be aware of the technical issues for ANA assays used
8
9
10 447 in their jurisdictions and recognize that different ANA assays or simply following
11
12 448 manufacturer's recommended reference ranges might not be optimal in applying ANA testing
13
14 449 results (42, 43). Additional longitudinal studies comparing other ELISAs and SPMAI such as
15
16 450 other multiplex bead immunoassays and emerging ANA technologies are needed.
17
18
19 451
20
21 452 Our study has some important strengths. To our knowledge, this is the largest review of ANA
22
23 453 status in SLE patients with data collected longitudinally and in a protocolized fashion over a
24
25 454 mean follow-up of five years. All ANA testing was conducted in an accredited central laboratory
26
27 455 with stringent quality control. However, we acknowledge some important limitations. First, there
28
29 456 may be a potential selection bias for SLE patients who are ANA positive to be enrolled into the
30
31 457 SLICC cohort compared to patients in conventional clinical care. Second, as enrolment could
32
33 458 occur up to 15 months after diagnosis (although mean disease duration at enrolment was 0.58
34
35 459 years), most patients had already been exposed to ≥ 1 immunomodulatory medication by
36
37 460 enrolment, which could potentially influence the ANA result. Third, although we showed that
38
39 461 demographic and clinical characteristics of the cohort subset with three available serum samples
40
41 462 were largely similar to the remainder of the cohort, our sample included a larger proportion of
42
43 463 Asian and fewer Hispanic participants. While our sample was racially and geographically
44
45 464 diverse, it is not known if our findings are generalizable to other SLE cohorts. Fourth, the
46
47 465 duration of follow-up, although relatively long at five years, does not capture potential
48
49 466 seroconversions or measure assay performance later in the disease. Last, there are >10 different
50
51
52
53
54
55
56
57
58
59
60

ANA immunoassays in use world-wide and our study utilized three. Regrettably, some manufacturers declined to participate in this study. Hence, generalization to all ANA assays is not possible (42, 44).

In conclusion, we demonstrated that early in their disease course almost all adult SLE patients had highly positive ANAs. However, as the disease progressed, we observed increased frequency of ANA within the normal range and decreased ANA titres/values by some assays likely related to differences in assay performance, medication exposure, decreased autoantibody responses over time, and lower disease activity. Combining ANA assays resulted in fewer patients that tested within normal range and no patients who tested within the normal range over the five years with all three assays. A clinical implication of this study is that for patients who have a moderate-to-high suspicion of SLE, especially those early in the disease course but without an established diagnosis, screening on both ELISA and HEp-2 IFA is warranted if one or the other provides results in the normal range. And given the rather modest changes in ANA frequency (and/or titers) observed in this longitudinal study of 5 years follow up, it is difficult to perceive of actionable clinical value of ANA IFA or screening ELISA test results over this time period once the diagnosis of SLE has been established. Since there are differences in the performance characteristics of individual ANA assays, clinicians need to be aware of the performance characteristics of the ANA test that their laboratories use. Future studies testing the comparative performance of other ANA immunoassays over time in large populations will help inform approaches to an earlier and more accurate diagnosis and classification of SLE.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Key Messages:

What is already known about this subject?

- Cross-sectional data of small cohorts suggest significant variation in the performance of antinuclear antibody (ANA) assays from different manufacturers leaving clinicians uncertain about the use or value of ANA testing in making a diagnosis.

What does this study add?

- In a longitudinal analysis of well-characterized patients with incident systemic lupus erythematosus (SLE), almost all SLE patients early in disease had highly positive ANAs and no patients who tested within the normal range over 5 years of follow up with all three assays.
- As the disease evolved over 5 years of follow-up, there was decreased frequency of positive ANAs (above the normal range) and decreased ANA titres by some assays.

How might this impact on clinical practice or future developments?

- In a patient without an established diagnosis of SLE and in whom the clinical suspicion for SLE is moderate to high, both IFA and ELISA should be performed if one or the other provides results in the normal range.

508 TABLES

509 Table 1. Patient characteristics at enrolment and year 5 (n=805)

Characteristic	Enrolment	Year 5	Difference ¹ (95% CI)
Demographic and Clinical			
Mean age at dx, yrs (SD)	35.2 (13.6)		
Female, %	88.7		
Mean disease duration, yrs (SD)	0.58 (0.49)		
Mean number of ACR Criteria without ANA (SD)	3.9 (1.0)		
Ethnicity, %			
Asian	24.3		
African	13.5		
White	52.3		
Hispanic	6.3		
Other ethnicities ²	3.5		
Nephritis ³	28.9	36.6	7.7 (5.7, 9.7)
Mean total SLEDAI-2K (SD) ⁴	5.4 (5.3)	3 (3.5)	-2.3 (-2.7, -1.9)
Mean total SDI (SD) ⁵	0.34 (0.74)	0.86 (1.25)	0.52 (0.43, 0.62)
Medications			
Current, %			
Glucocorticoids	69.6	56.8	-12.8 (-16.5, -9.1)
Antimalarials	70.1	79.4	9.3 (5.9, 12.7)
Immunosuppressants	41.0	50.8	9.8 (6.1, 13.5)
Ever, %			
Glucocorticoids	81.5	87.3	5.8 (4.1, 7.6)
Antimalarials	76.6	91.1	14.4 (11.9, 17)
Immunosuppressants	43.9	66.3	22.5 (19.5, 25.5)
Autoantibodies, %			
dsDNA ⁶	34.2	29.1	-5.1 (-8.7, -1.6)
Ribosomal P	24.3	20	-4.3 (-7.8, -0.9)
Ro52/TRIM21	37.5	37.4	-0.1 (-3.4, 3.2)
SSA/Ro60	42.5	42	-0.5 (-3.7, 2.7)
SSB/La	20.7	16.3	-4.5 (-7.5, -1.5)
Sm	22.7	14.7	-8.1 (-11.1, -5.0)
U1RNP	28.2	23	-5.2 (-8.5, -2.0)
Histones	31.3	22.7	-8.6 (-12.1, -5.0)
Cardiolipin IgG/IgM ⁷	20.5	16.4	-4.1 (-7.7, -0.6)
β2GP1 IgG/IgM ⁷	19.8	12.9	-6.9 (-9.8, -4)
Lupus anticoagulant ⁸	20.6	16.7	-3.9 (-9.8, 2)
Abbreviations: ACR, American College of Rheumatology; ANA, anti-nuclear antibodies; β2GP1, β2-glycoprotein-1; CI, confidence interval; dx, diagnosis; dsDNA, double-stranded DNA; IgG/M, immunoglobulin G/immunoglobulin M; RNP, ribonucleoprotein; SD, standard deviation; SLEDAI-2K, systemic lupus erythematosus disease activity index-2000; SDI, SLICC Damage index; Sm, Smith; TRIM21, Tripartite Motif Protein (TRIM) 21; yrs, years.			
1. Difference between enrolment and year 5 visit;			
2. Other ethnicities include: Native North American, Native Hawaiian or other Pacific Islanders			
3. Nephritis defined as fulfilling the ACR criterion for renal disease or if a renal biopsy was performed prior to cohort entry			
4. Complete data available for n=793 patients			
5. Complete data available for n= 380 as the disease needs to be present for at least 6 months before the SDI can be calculated.			
6. Complete data available for n=798 patients			
7. Complete data available for n= 800			
8. Complete data available for n=282			

Table 2. ANA inter-test percentage agreement among IFA1 (n=805), IFA2 (n=805), and ELISA (n=805)

	Enrolment (%)		Year 3 (%)		Year 5 (%)	
	IFA1	IFA2	IFA1	IFA2	IFA1	IFA2
IFA2	96.4% (94.9 -97.6)		95.2% (93.4-96.5)		95.5% (93.9-96.8)	
ELISA	94.8% (93.0-96.2)	95.7% (94.0-97.0)	91.2% (89.0-93.0)	92.5% (90.5-94.3)	91.4% (89.3-93.3)	91.2% (89.0-93.0)
Abbreviations: ANA, anti-nuclear antibodies; ELISA, enzyme-linked immunosorbent assay; IFA; indirect immunofluorescence assay.						

Table 3. ANA patterns over time with indirect immunofluorescence assay (IFA) 1 (n=805) and IFA2 (n=805)

Pattern	Enrolment n (%)	Year 3 n (%)	Year 5 n (%)	Same ANA Pattern Over 5 years n (%)
IFA 1 Patterns				
Nuclear	481 (62.1)	519 (68.1)	526 (68.7)	305 (37.9)
Cytoplasmic +/- Mitotic	17 (2.2)	18 (2.4)	21 (2.7)	1 (0.1)
Mixed	276 (35.7)	225 (29.5)	219 (28.6)	81 (10.1)
IFA2 Patterns				
Nuclear	491 (62.1)	477 (60.4)	472 (59.3)	273 (33.9)
Cytoplasmic +/- Mitotic	9 (1.1)	6 (0.8)	4 (0.5)	0 (0.0)
Mixed	291 (36.8)	308 (38.8)	320 (40.2)	114 (14.2)
IFA1 and 2 agreement (k)				
Agreement (95%CI)	74.0 (70.7-77.0)*	71.4 (68.0-74.6)*	71.0 (67.7-74.2)*	
Kappa (95%CI)	0.46 (0.39-0.53)	0.39 (0.33-0.46)	0.39 (0.33-0.46)	
Abbreviations: ANA, anti-nuclear antibodies; IFA; indirect immunofluorescence assay. *p<0.0001 using unweighted kappa (k) statistics.				

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

519 **Table 4. Autoantibodies detected in patients with an ANA that was within the normal range on IFA1, IFA2, ELISA, either**
520 **alone, on two or all three assay at enrolment and year 5***
521

	ELISA		IFA1		IFA2		IFA1&ELISA		ELISA and IFA2		IFA1&IFA2		All three assays	
% Autoantibodies	Enrolment (N=27)	Year 5 (N=70)	Enrolment (n=31)	Year 5 (n=39)	Enrolment (N=14)	Year 5 (N=9)	Enrolment (N=8)	Year 5 (N=20)	Enrolment (N=3)	Year 5 (N=4)	Enrolment (N=8)	Year 5 (N=6)	Enrolment (N=3)	Year 5 (N=3)
None detected	44.4	45.7	38.7	53.8	42.9	44.4	62.5	65.0	66.7	50.0	50.0	50.0	66.7	66.7
dsDNA ¹	7.7	5.7	6.7	5.1	0.0	11.1	0.0	0.0	0.0	0.0	0.0	16.7	0.0	0.0
Ribosomal P	3.7	11.4	6.5	10.3	7.1	11.1	0.0	10.0	0.0	25.0	0.0	16.7	0.0	33.3
Ro52/TRIM21	11.1	21.4	22.6	20.5	21.4	11.1	0.0	20.0	0.0	25.0	0.0	0.0	0.0	0.0
SSA/Ro60	7.4	12.9	25.8	10.3	21.4	11.1	0.0	5.0	0.0	0.0	12.5	0.0	0.0	0.0
SSB/La	7.4	7.1	0.0	5.1	0.0	0.0	0.0	5.0	0.0	0.0	0.0	0.0	0.0	0.0
Sm	3.7	4.3	6.5	2.6	0.0	11.1	0.0	0.0	0.0	25.0	0.0	0.0	0.0	0.0
U1RNP	3.7	7.1	0.0	5.1	0.0	0.0	0.0	5.0	0.0	0.0	0.0	0.0	0.0	0.0
Histones	0.0	10.0	0.0	2.6	7.1	11.1	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
*Patients who were within the normal range for ANA at enrolment are not necessarily the same patients at year 5 and vice versa. Abbreviations: ANA, anti-nuclear antibodies; β2GP1, β2-glycoprotein-1; dsDNA, double-stranded DNA; ELISA, enzyme-linked immunosorbent assay, IFA; indirect immunofluorescence assay; IgG/M, immunoglobulin G/immunoglobulin M; RNP, ribonucleoprotein; Sm, Smith; TRIM21, TRIPartite Motif protein (TRIM) 21. ¹ dsDNA was measured at enrolment for only 26 patients on ELISA, 13 on IFA2, and 2 on both who tested within the normal range for ANA														

522
523

Only

References:

1. Choi MY, Fritzler MJ. Autoantibodies in SLE: prediction and the p value matrix. *Lupus*. 2019;28(11):1285-93.
2. Leuchten N, Hoyer A, Brinks R, Schoels M, Schneider M, Smolen J, et al. Performance of Antinuclear Antibodies for Classifying Systemic Lupus Erythematosus: A Systematic Literature Review and Meta-Regression of Diagnostic Data. *Arthritis Care Res (Hoboken)*. 2018;70(3):428-38.
3. Aringer M, Costenbader K, Daikh D, Brinks R, Mosca M, Ramsey-Goldman R, et al. 2019 European League Against Rheumatism/American College of Rheumatology Classification Criteria for Systemic Lupus Erythematosus. *Arthritis Rheumatol*. 2019;71(9):1400-12.
4. Aringer M, Brinks R, Dorner T, Daikh D, Mosca M, Ramsey-Goldman R, et al. European League Against Rheumatism (EULAR)/American College of Rheumatology (ACR) SLE classification criteria item performance. *Ann Rheum Dis*. 2021.
5. Heller CA, Schur PH. Serological and clinical remission in systemic lupus erythematosus. *J Rheumatol*. 1985;12(5):916-8.
6. Paz E, Adawi M, Lavi I, Mussel Y, Mader R. Antinuclear antibodies measured by enzyme immunoassay in patients with systemic lupus erythematosus: relation to disease activity. *Rheumatology international*. 2007;27(10):941-5.
7. Frodlund M, Wettero J, Dahle C, Dahlstrom O, Skogh T, Ronnelid J, et al. Longitudinal anti-nuclear antibody (ANA) seroconversion in systemic lupus erythematosus: a prospective study of Swedish cases with recent-onset disease. *Clin Exp Immunol*. 2020;199(3):245-54.
8. Faria AC, Barcellos KS, Andrade LE. Longitudinal fluctuation of antibodies to extractable nuclear antigens in systemic lupus erythematosus. *J Rheumatol*. 2005;32(7):1267-72.
9. Arbuckle MR, McClain MT, Rubertone MV, Scofield RH, Dennis GJ, James JA, et al. Development of autoantibodies before the clinical onset of systemic lupus erythematosus. *N Engl J Med*. 2003;349(16):1526-33.
10. Hassan AB, Lundberg IE, Isenberg D, Wahren-Herlenius M. Serial analysis of Ro/SSA and La/SSB antibody levels and correlation with clinical disease activity in patients with systemic lupus erythematosus. *Scand J Rheumatol*. 2002;31(3):133-9.
11. Tench CM, Isenberg DA. The variation in anti-ENA characteristics between different ethnic populations with systemic lupus erythematosus over a 10-year period. *Lupus*. 2000;9(5):374-6.
12. Praprotnik S, Bozic B, Kveder T, Rozman B. Fluctuation of anti-Ro/SS-A antibody levels in patients with systemic lupus erythematosus and Sjogren's syndrome: a prospective study. *Clin Exp Rheumatol*. 1999;17(1):63-8.
13. Fisher DE, Reeves WH, Wisniewolski R, Lahita RG, Chiorazzi N. Temporal shifts from Sm to ribonucleoprotein reactivity in systemic lupus erythematosus. *Arthritis Rheum*. 1985;28(12):1348-55.
14. Barada FA, Jr., Andrews BS, Davis JSt, Taylor RP. Antibodies to Sm in patients with systemic lupus erythematosus. Correlation of Sm antibody titers with disease activity and other laboratory parameters. *Arthritis Rheum*. 1981;24(10):1236-44.
15. Acosta-Merida A, Isenberg DA. Antinuclear antibodies seroconversion in 100 patients with lupus. *Clin Exp Rheumatol*. 2013;31(4):656.

16. Gisca E, Duarte L, Farinha F, Isenberg DA. Assessing outcomes in a lupus nephritis cohort over a 40-year period. *Rheumatology (Oxford)*. 2021;60(4):1814-22.
17. Fritzler MJ. Choosing wisely: Review and commentary on anti-nuclear antibody (ANA) testing. *Autoimmun Rev*. 2016;15(3):272-80.
18. Raissi TC, Hewson C, Pope JE. Repeat Testing of Antibodies and Complements in Systemic Lupus Erythematosus: When Is It Enough? *J Rheumatol*. 2018;45(6):827-34.
19. Bossuyt X, De Langhe E, Borghi MO, Meroni PL. Understanding and interpreting antinuclear antibody tests in systemic rheumatic diseases. *Nat Rev Rheumatol*. 2020;16(12):715-26.
20. Bizzaro N, Brusca I, Previtali G, Alessio MG, Daves M, Platzgummer S, et al. The association of solid-phase assays to immunofluorescence increases the diagnostic accuracy for ANA screening in patients with autoimmune rheumatic diseases. *Autoimmun Rev*. 2018;17(6):541-7.
21. Pisetsky DS, Spencer DM, Lipsky PE, Rovin BH. Assay variation in the detection of antinuclear antibodies in the sera of patients with established SLE. *Ann Rheum Dis*. 2018;77(6):911-3.
22. Chevrier M, Jordan J, Schreiter J, Benson J, editors. Comparative analysis of anti-nuclear antibody testing using blinded replicate samples reveals variability between commercial testing laboratories (abstract). *Arthritis & rheumatology*; 2016.
23. Kumagai S, Hayashi N. Immunofluorescence--still the 'gold standard' in ANA testing? *Scand J Clin Lab Invest Suppl*. 2001;235:77-83.
24. Perez D, Gilburd B, Azoulay D, Shovman O, Bizzaro N, Shoenfeld Y. Antinuclear antibodies: Is the indirect immunofluorescence still the gold standard or should be replaced by solid phase assays? *Autoimmun Rev*. 2018;17(6):548-52.
25. Meroni PL, Chan EK, Damoiseaux J, Andrade LEC, Bossuyt X, Conrad K, et al. Unending story of the indirect immunofluorescence assay on HEp-2 cells: old problems and new solutions? *Ann Rheum Dis*. 2019;78(6):e46.
26. Claessens J, Belmondo T, De Langhe E, Westhovens R, Poesen K, Hue S, et al. Solid phase assays versus automated indirect immunofluorescence for detection of antinuclear antibodies. *Autoimmun Rev*. 2018;17(6):533-40.
27. Choi MY, Clarke AE, St Pierre Y, Hanly JG, Urowitz MB, Romero-Diaz J, et al. Antinuclear Antibody-Negative Systemic Lupus Erythematosus in an International Inception Cohort. *Arthritis Care Res (Hoboken)*. 2019;71(7):893-902.
28. Isenberg D, Ramsey-Goldman R, Gladman D, Hanly J. The Systemic Lupus International Collaborating Clinics (SLICC) group--It was 20 years ago today. *Lupus*. 2011;20(13):1426-32.
29. Hochberg MC. Updating the American College of Rheumatology revised criteria for the classification of systemic lupus erythematosus. *Arthritis and rheumatism*. 1997;40(9):1725.
30. Damoiseaux J, von Muhlen CA, Garcia-De La Torre I, Carballo OG, de Melo Cruvinel W, Francescantonio PL, et al. International consensus on ANA patterns (ICAP): the bumpy road towards a consensus on reporting ANA results. *Auto Immun Highlights*. 2016;7(1):1.
31. Aringer M, Costenbader KH, Dorner T, Johnson SR. Importance of high-quality ANA testing for SLE classification. Response to: 'Role of ANA testing in the classification of patients with systemic lupus erythematosus' by Pisetsky et al. *Ann Rheum Dis*. 2019;annrhumdis-2019-216337.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 611 32. Pisetsky DS, Rovin BH, Lipsky PE. New Perspectives in Rheumatology: Biomarkers as
612 Entry Criteria for Clinical Trials of New Therapies for Systemic Lupus Erythematosus: The
613 Example of Antinuclear Antibodies and Anti-DNA. *Arthritis Rheumatol.* 2017;69(3):487-93.
614 33. Mahler M, Meroni PL, Bossuyt X, Fritzler MJ. Current concepts and future directions for
615 the assessment of autoantibodies to cellular antigens referred to as anti-nuclear antibodies. *J*
616 *Immunol Res.* 2014;2014:315179.
617 34. Olsen NJ, Choi MY, Fritzler MJ. Emerging technologies in autoantibody testing for
618 rheumatic diseases. *Arthritis Res Ther.* 2017;19(1):172.
619 35. Pisetsky DS, Thompson DK, Wajdula J, Diehl A, Sridharan S. Variability in Antinuclear
620 Antibody Testing to Assess Patient Eligibility for Clinical Trials of Novel Treatments for Systemic
621 Lupus Erythematosus. *Arthritis Rheumatol.* 2019;71(9):1534-8.
622 36. Wohrer S, Troch M, Zwerina J, Schett G, Skrabbs C, Gaiger A, et al. Influence of rituximab,
623 cyclophosphamide, doxorubicin, vincristine and prednisone on serologic parameters and clinical
624 course in lymphoma patients with autoimmune diseases. *Ann Oncol.* 2007;18(4):647-51.
625 37. Prado MS, Dellavance A, Rodrigues SH, Marvulle V, Andrade LEC. Changes in the result
626 of antinuclear antibody immunofluorescence assay on HEp-2 cells reflect disease activity status
627 in systemic lupus erythematosus. *Clin Chem Lab Med.* 2020;58(8):1271-81.
628 38. Ng KP, Cambridge G, Leandro MJ, Edwards JC, Ehrenstein M, Isenberg DA. B cell
629 depletion therapy in systemic lupus erythematosus: long-term follow-up and predictors of
630 response. *Ann Rheum Dis.* 2007;66(9):1259-62.
631 39. Pisetsky DS, Lipsky PE. New insights into the role of antinuclear antibodies in systemic
632 lupus erythematosus. *Nat Rev Rheumatol.* 2020;16(10):565-79.
633 40. Aringer M, Dörner T, Leuchten N, Johnson S. Toward new criteria for systemic lupus
634 erythematosus—a standpoint. *Lupus.* 2016;25(8):805-11.
635 41. Bizzaro N. Can solid-phase assays replace immunofluorescence for ANA screening? *Ann*
636 *Rheum Dis.* 2020;79(3):e32.
637 42. Fritzler MJ, Wiik A, Fritzler ML, Barr SG. The use and abuse of commercial kits used to
638 detect autoantibodies. *Arthritis Res Ther.* 2003;5(4):192-201.
639 43. Fritzler MJ. The antinuclear antibody test: last or lasting gasp? *Arthritis Rheum.*
640 2011;63(1):19-22.
641 44. Wiik AS, Bizzaro N. Missing links in high quality diagnostics of inflammatory systemic
642 rheumatic diseases: It is all about the patient! *Auto Immun Highlights.* 2012;3(2):35-49.
643




Figure 1. ANA positivity among IFA1 (n=805), IFA2 (n=805) and ELISA (n=805) at enrolment, year 3 and year 5. There is a break in the y-axis between 40% and 90% to enhance the readability of the graph from 90-100%.

162x117mm (300 x 300 DPI)




Figure 2. Distribution of ANA titres at enrolment and year 5 visit for A) indirect immunofluorescence 1 (IFA1) (n=805), B) IFA2 (n=805) and C) enzyme-linked immunosorbent assay (ELISA) (n=805). Lines represent the curve of best fit.

149x190mm (300 x 300 DPI)

Supplemental Table 1. SLE-related Autoantibodies performed

Autoantibody	Assay and Cut-offs
Anti-dsDNA	Anti-dsDNA positivity and titers were detected by a chemiluminescence immunoassay (CLIA) (Werfen, San Diego, USA). A cut-off of ≥ 27 chemiluminescence units (CU) was utilized, where 27-35 (CU) was indeterminate (borderline), and >35 was positive.
Other SLE-specific autoantibodies	Performed using ALBIA (FIDIS Connective13: TheraDiag, Paris, France) on a Luminex 200 flow luminometer (Luminex, Austin, USA) focussing on SLE-related analytes that included ribosomal P, Ro52/Tripertite Motif Protein 21 (TRIM21), SSA/Ro60, SSB/La, Sm, U1-RNP, and histones. A cut-off of >40 median fluorescence units (MFU) was considered positive.
Anti-phospholipid antibodies	Anti-phospholipid antibodies including IgG and IgM anticardiolipin and anti- $\beta 2$ -glycoprotein-1 were measured using ELISA (Werfen, San Diego, USA). Using the revised Sapporo antiphospholipid syndrome classification criteria (1), a cut-off of >40 units for IgG/IgM anti-cardiolipin was considered medium to high positive while a cut-off of ≥ 20 units ($>99^{\text{th}}$ percentile) was positive for IgG/IgM anti- $\beta 2$ -glycoprotein-1 ($\beta 2\text{GP1}$) (1). All autoantibodies were measured at MitogenDx except for lupus anticoagulant, which was measured at Oklahoma Medical Foundation (Oklahoma City, OK) as previously described (2).

1. Miyakis S, Lockshin MD, Atsumi T, Branch DW, Brey RL, Cervera R, et al. International consensus statement on an update of the classification criteria for definite antiphospholipid syndrome (APS). J Thromb Haemost. 2006;4(2):295-306.

2, Hanly J, Urowitz M, Siannis F, Farewell V, Gordon C, Bae S, et al. Autoantibodies and neuropsychiatric events at the time of systemic lupus erythematosus diagnosis: results from an international inception cohort study. Arthritis & Rheumatism: Official Journal of the American College of Rheumatology. 2008;58(3):843-53.

Supplemental Table 2. Clinically defined samples

Demographic and Clinical Variables	Definition
Age	Years at diagnosis
Sex	Female or male
Race/ethnicity	Asian: Chinese, Filipino, Japanese, Korean, other Asians; African descendants: African, Caribbean; Hispanic: Hispanics only; White: North American, Indian – sub-continent, other Caucasians; Other races/ethnicities: Native North American, Native Hawaiian or other Pacific Islanders, others
Presence of nephritis	Based on renal biopsy or fulfillment of the renal item of the ACR Classification criteria
American College of Rheumatology (ACR) criteria	Number of specific ACR criteria fulfilled
SLE Disease Activity Index (SLEDAI-2K) score	Disease activity measured by global SLEDAI-2K score and its individual components grouped to represent the following organ systems: Neurological: seizures, psychosis, organic brain syndrome, visual disturbance, cranial nerve disorder, lupus headache, CVA Mucocutaneous: vasculitis, rash, alopecia, mucosal ulcers Musculoskeletal: arthritis, myositis Renal: urinary casts, hematuria, proteinuria, pyuria Serositis: pleurisy, pericarditis Constitutional: fever Immunological: low complement, increased DNA binding Hematological: thrombocytopenia, leukopenia
Medications	Any use of oral or parental glucocorticoids, antimalarials, immunosuppressive agents (methotrexate, azathioprine, mycophenolate, cyclophosphamide, cyclosporine, and biologics) at or prior to enrollment.

Supplemental Table 3. Patient characteristics at enrolment comparing patients included in this study and the remaining SLICC pts providing at least 4 yrs of clinical data, but for whom 3 serial samples were not available

Characteristic	Study Cohort n=805	Remainder of SLICC n=627	Difference (95% CI) ¹
Mean age at dx, yrs (SD)	35.2 (13.6)	34.1 (13.0)	1.1 (-0.3, 2.4)
Female, %	88.7	88.5	0.2 (-3.1, 3.5)
Mean disease duration, yrs (SD)	0.58 (0.49)	0.49 (0.35)	0.1 (0.05, 0.14)
Mean (SD) number of ACR Criteria excluding the ANA	3.9 (1)	4 (1.1)	-0.1 (-0.2, 0)
Ethnicity, %			
Asian	24.3	5.6	18.8 (15.3, 22.2)
African	13.5	18.3	-4.8 (-8.6, -1)
White	52.3	45.1	7.2 (2, 12.4)
Hispanics	6.3	27	-20.6 (-24.5, -16.8)
Other ethnicity ²	3.5	4	-0.5 (-2.5, 1.5)
Nephritis ³	28.9	33.3	-4.4 (-9.2, 0.5)
Mean total SLEDAI-2K (SD)	5.4 (5.3)	5.2 (5.46)	0.2 (-0.4, 0.8)
Mean total SDI (SD)	0.34 (0.74)	0.35 (0.8)	-0.01 (-0.13, 0.11)
Current medications, %			
Glucocorticoids	69.6	71	-1.4 (-6.2, 3.4)
Antimalarials	70.1	67.3	2.8 (-2.1, 7.6)
Immunosuppressants	41	43.1	-2.1 (-7.2, 3.1)
Medications ever, %			
Glucocorticoids	81.5	82.1	-0.6 (-4.7, 3.4)
Antimalarials	76.6	73.5	3.1 (-1.4, 7.6)
Immunosuppressants	43.9	46.3	-2.4 (-7.6, 2.8)
Autoantibodies, %			
Lupus anticoagulant	19.7	19.8	-0.1 (-5.6, 5.3)

Abbreviations: ACR, American College of Rheumatology; ANA, anti-nuclear antibodies; CI, confidence interval; x, diagnosis; SD, standard deviation; SLEDAI-2K, systemic lupus erythematosus disease activity index-2000; SDI, SLICC Disease index; yrs, years.

1. Difference between study cohort and remainder of SLICC cohort

2. Other ethnicities include: Native North American, Native Hawaiian or other Pacific Islanders

3. Nephritis defined as fulfilling the ACR criterion for renal disease or if a renal biopsy was performed prior to cohort entry

Supplemental Table 4. Patient characteristics at enrolment comparing patients providing 3rd sample between 4 and 7 yrs of follow-up and pts providing 3rd sample between 8 and 10 yrs of follow-up. Bolded indicates statistically significant $p < 0.05$.

Characteristic	Patients without delayed year 5 visit N=781	Patients with delayed year 5 visit N=24	Difference (95% CI) ¹
Demographic and Clinical			
Mean age at dx, yrs (SD)	35.2 (13.6)	32.8 (14.1)	2.4 (-3.3, 8.1)
Female, %	88.7	87.5	1.2 (-12.2, 14.6)
Mean disease duration, yrs (SD)	0.58 (0.49)	0.76 (0.63)	-0.18 (-0.44, 0.07)
Meeting ACR Criteria without ANA, %	3.9 (1)	3.9 (1.02)	0 (-0.4, 0.4)
Ethnicity, %			
Asian	24.6	16.7	7.9 (-7.3, 23.1)
African	13.6	12.5	1.1 (-12.4, 14.5)
Caucasian	52.4	50	2.4 (-17.9, 22.7)
Hispanic	6.1	12.5	-6.4 (-19.7, 7)
Other ethnicities ²	3.3	8.3	-5 (-16.1, 6.1)
Nephritis ³	28.9	29.2	-0.2 (-18.7, 18.2)
Mean total SLEDAI-2K (SD)	5.4 (5.3)	4.3 (4.03)	1.1 (-0.6, 2.7)
Mean total SDI (SD) ⁴	0.33 (0.73)	0.67 (0.89)	-0.34 (-0.85, 0.17)
Medications			
Current, %			
Glucocorticoids	69.4	75	-5.6 (-23.2, 12)
Antimalarials	70.2	66.7	3.5 (-15.6, 22.6)
Immunosuppressants	41	41.7	-0.7 (-20.7, 19.3)
Ever, %			
Glucocorticoids	81	95.8	-14.8 (-23.2, -6.3)
Antimalarials	77	66.7	10.3 (-8.8, 29.4)
Immunosuppressants	43.8	45.8	-2 (-22.3, 18.2)
Autoantibodies, %			
DsDNA ⁵	34.4	25	9.4 (-8.3, 27)
Ribosomal P	24.7	12.5	12.2 (-1.4, 25.8)
Ro52/TRIM21	37.3	45.8	-8.6 (-28.8, 11.6)
SSA/Ro60	42.6	37.5	5.1 (-14.5, 24.8)
SSB/La	21	12.5	8.5 (-5, 22)
Sm	23.2	8.3	14.8 (3.4, 26.3)
U1RNP	28.4	20.8	7.6 (-9, 24.1)
Histones	31.8	16.7	15.1 (-0.2, 30.4)
Cardiolipin IgG/IgM ⁶	20.6	16.7	4 (-11.2, 19.1)
β 2GP1 IgG/IgM ⁶	20.1	8.3	11.8 (0.4, 23.2)

Lupus anticoagulant ⁷	20.1	5.3	14.8 (4.3, 25.3)
----------------------------------	------	-----	-------------------------

Abbreviations: ACR, American College of Rheumatology; ANA, anti-nuclear antibodies; β 2GP1, β 2-glycoprotein-1; CI, confidence interval; dx, diagnosis; dsDNA, double-stranded DNA; IgG/M, immunoglobulin G/immunoglobulin M; RNP, ribonucleoprotein; SD, standard deviation; SLEDAI-2K, systemic lupus erythematosus disease activity index-2000; SDI, SLICC Disease index; TRIM21, Tripartite Motif Protein (TRIM) 21; yrs, years.

1. Difference between enrolment and year 5 visit
2. Other ethnicities include: Native North American, Native Hawaiian or other Pacific Islanders
3. Nephritis defined as fulfilling the ACR criterion for renal disease or if a renal biopsy was performed prior to cohort entry
4. Data available for n= 368, 781, 780 at enrolment, year 3, and year 5 respectively
5. Data available for n= 777, 778, 778 at enrolment, year 3, and year 5 respectively
6. Data available for n= 776, 781, and 781, at enrolment, year 3, and year 5 respectively
7. Data available for n= 647, 469, and 288, at enrolment, year 3, and year 5 respectively

Supplemental Table 5. Pairwise conflicting results between IFA1, IFA2, and ELISA (total n at each time point = 805)

	+IFA1 -IFA2 (n)	-IFA1 +IFA2 (n)	p-value	+IFA1 -ELISA (n)	-IFA1 +ELISA (n)	p-value	+IFA2 -ELISA (n)	-IFA2 +ELISA (n)	p-value
Enrolment	6	23	<0.05	19	23	NS	24	11	<0.05
Year 3	4	35	<0.001	43	28	NS	53	7	<0.001
Year 5	3	33	<0.001	50	19	<0.001	66	5	<0.001

Abbreviations: ANA, anti-nuclear antibodies; ELISA, enzyme-linked immunosorbent assay; IFA; indirect immunofluorescence assay; NS, non-significant. **p<0.001, *p<0.05 for significant asymmetry using McNemar's Test

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 **Supplemental Figure 1. Frequency of IFA1 (A) and IFA2 (B) ANA on HEp-2 ICAP patterns AC0-AC29, matrix at enrolment,**
49 **year 3, and year 5. ** Represents AC- 2-3, 6, 9, 11-18, 22-26, 28-29, * represent AC-0, 2-3, 6, 9, 11-18, 22-23, 25-29. Refer to the**
50 **ICAP website (www.anapatterns.org) for detailed descriptors for each AC pattern. For IFA1 or IFA2, the most common**
51 **patterns: AC-4 represents nuclear fine speckled, AC-5 nuclear large speckled, AC-1 nuclear homogeneous, AC-19 cytoplasmic**
52 **dense fine speckled, AC-20 cytoplasmic fine speckled, AC-matrix is matrix, AC-10 punctate nucleolar, AC-7 few discrete**
53 **nuclear dots, AC-21 cytoplasmic reticular/anti-mitochondrial antibodies, AC-8 homogeneous nucleolar, AC-24 centrosome,**
54 **AC-27 intracellular bridge, AC-0 negative/no pattern observed.**
55

56

