

Association between wood and other biomass fuels and risk of low birthweight in Uganda

Eputai, Joshua; Woolley, Katherine E; Bartington, Suzanne E; Thomas, G Neil

DOI:

[10.3390/ijerph19074377](https://doi.org/10.3390/ijerph19074377)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Eputai, J, Woolley, KE, Bartington, SE & Thomas, GN 2022, 'Association between wood and other biomass fuels and risk of low birthweight in Uganda: a cross-sectional analysis of 2016 Uganda demographic and health survey data', *International Journal of Environmental Research and Public Health*, vol. 19, no. 7, 4377. <https://doi.org/10.3390/ijerph19074377>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



Article

Association between Wood and Other Biomass Fuels and Risk of Low Birthweight in Uganda: A Cross-Sectional Analysis of 2016 Uganda Demographic and Health Survey Data

Joshua Eputai^{1,2}, Katherine E. Woolley¹ , Suzanne E. Bartington¹ and G. Neil Thomas^{1,*}

¹ Institute of Applied Health Research, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK; jxe078@alumni.bham.ac.uk (J.E.); kew863@student.bham.ac.uk (K.E.W.); s.bartington@bham.ac.uk (S.E.B.)

² Department of Nursing, Faculty of Health Sciences, Busitema University, Mbale P.O. Box 1460, Uganda

* Correspondence: g.n.thomas@bham.ac.uk

Abstract: In utero exposure to household air pollution (HAP) from polluting cooking fuels has been linked to adverse pregnancy outcomes including low birthweight (LBW). No previous study in Uganda has attempted to investigate the association between the different types of biomass cooking fuels and LBW. This study was conducted to investigate the association between wood and other biomass cooking fuel use with increased risk of LBW, using the 2016 Uganda Demographic and Health Survey for 15,270 live births within five years prior to interview. LBW, defined as birthweight of <2500 g, was estimated from maternal recall and health cards. Association between household exposure to the different solid biomass cooking fuels and LBW was determined using multivariable logistic regression. Biomass cooking fuels were used in 99.6% of the households, with few (0.3%) using cleaner fuels and 0.1% with no cooking, while the prevalence of LBW was 9.6% of all live-births. Although the crude analysis suggested an association between wood fuel use and LBW compared to other biomass and kerosene fuel use (AOR: 0.82; 95% CI: 0.67–1.00), after adjusting for socio-demographic and obstetric factors, no association was observed (AOR: 0.94; 95% CI: 0.72–1.22). LBW was significantly more likely among female neonates (AOR: 1.32 (95% CI: 1.13–1.55) and neonates born to mothers living in larger households (AOR: 1.03; 95% CI: 1.00–1.07). LBW was significantly less likely among neonates delivered at term (AOR: 0.39; 95% CI: 0.31–0.49), born to women with secondary or tertiary level of education (AOR: 0.80; 95% CI: 0.64–1.00), living in households with a higher wealth index (AOR: 0.69; 95% CI: 0.50–0.96), Eastern (AOR: 0.76; 95% CI: 0.59–0.98) and Northern (AOR: 0.75; 95% CI: 0.57–0.99) regions. The study findings suggest inconclusive evidence regarding the association between the use of wood compared to other biomass and kerosene cooking fuels and risk of LBW. Given the close observed association between socioeconomic status and LBW, the Ugandan government should prioritize public health actions which support female education and broader sustainable development to improve household living standards in this setting.

Keywords: biomass cooking fuels; low birthweight; pregnancy outcomes; Uganda; household air pollution



Citation: Eputai, J.; Woolley, K.E.; Bartington, S.E.; Thomas, G.N. Association between Wood and Other Biomass Fuels and Risk of Low Birthweight in Uganda: A Cross-Sectional Analysis of 2016 Uganda Demographic and Health Survey Data. *Int. J. Environ. Res. Public Health* **2022**, *19*, 4377. <https://doi.org/10.3390/ijerph19074377>

Academic Editors: Carla Viegas and Elisabete Carolino

Received: 18 February 2022

Accepted: 31 March 2022

Published: 5 April 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Globally, about 2.8 billion (36%) people rely on kerosene and biomass fuels (charcoal, coal, wood, animal dung, and agricultural residues), referred to collectively as biomass fuels, for cooking and heating [1,2]. In Asia, significant progress in transitioning to cleaner fuels (liquid petroleum gas [LPG], electricity) has been seen, but little progress has been made in Africa, where the proportion of solid fuel use has only decreased slightly from 90% in 1990 to 84% in 2020 [1]. The limited decrease in proportion of biomass cooking fuel use in Africa is offset by an absolute increase in the population that solely relies on polluting cooking fuels due to population growth [1]. By 2025, the use of biomass fuels for cooking is projected to exceed one billion people in Africa [1]. Biomass fuels often

undergo incomplete combustion, producing high levels of household air pollution (HAP), which includes particulate matter (PM) and carbon monoxide (CO) at levels that frequently exceed the World Health Organization Global Air Quality Guidelines (WHO-AQGs) [2].

Globally, exposure to HAP results in approximately four million deaths annually, 4.8% of all disability-adjusted life years, and 7.7% of global mortality [3], and is associated with increased risk of upper and lower respiratory tract infections, asthma, cataracts, lung cancer, chronic obstructive pulmonary disease, tuberculosis, and cardiovascular diseases [3,4]. In addition, women and girls are predisposed to musculoskeletal injuries, sexual violence, animal bites, insect-borne disease, and widening of gender inequities related to the gathering of biomass fuels [5].

Women and children under five years of age are more vulnerable to the adverse effects of HAP, due to gender-patterned cooking roles which lead to greater HAP exposure among females, while children aged under five years are often kept in close proximity to their mothers during cooking periods [6]. Exposure to HAP in utero, a particularly vulnerable stage of fetal development [6], has been associated with increased risk of adverse pregnancy outcomes including increased risk of low birthweight (LBW), preterm birth, small for gestational age, spontaneous abortion, stillbirth, and neonatal mortality [4,7,8]. Globally, 90% of 4 million neonatal deaths annually occur in developing countries, including Uganda [9]. LBW prevalence is disproportionately high (14%) in Sub-Saharan Africa [10], and is one of the leading causes of neonatal morbidity and mortality in Sub-Saharan Africa [9].

Studies and meta-analyses have identified an association between exposure to biomass compared to cleaner fuels (LPG, electricity) with increased risk of LBW [4,7,11,12]. However, only a limited number of studies from Africa were included in the meta-analyses [4,7,11,12], a region with a high burden of LBW [10], and predominant use of polluting fuels [1]. The studies from Africa were limited to relatively few countries, including Ghana [13], Nigeria [14–17], Malawi [18], Ethiopia [19–21], and Zimbabwe [22]. The African studies were limited to hospital-based settings [21] with a reliance on the woman's subjective recall of the birth weight or size of the child [14,18,22]. The studies did not address the effect of outdoor or indoor place of cooking on LBW risk [13–15,17,23], or generally control for important confounders [18]. Moreover, there remains limited understanding of the relationship between different biomass fuel types and risk of LBW, which is of importance in Sub-Saharan Africa where access to cleaner fuels remains limited. In Uganda, the only two studies focusing on HAP that were included in the review were limited to respiratory symptoms only [24–26]. To the best of our knowledge, no study in Uganda has investigated the impact of different types of biomass fuels on the risk of LBW.

Our study was conducted in Uganda, a low-income country in Sub-Saharan Africa [27], where 97% of the households predominantly use biomass fuels for cooking [28]. Uganda, with a population of about 40 million people, has a gross domestic product (GDP) per capita of 958.19 USD [27]. The maternal mortality rate is 336 per 100,000 live births with a neonatal mortality rate of 27 per 1000 live births [28]. The high gender inequality in Uganda means that household activities such as cooking and collecting firewood are usually undertaken by women [29]. About 21% of the population in Uganda live below the poverty line, with poor housing and under-ventilated living conditions experienced by over half (56%) of its population. Although at a national level ambitious energy policy targets have been set to promote access and adoption of cleaner fuels by 2040, the lack of innovative implementation strategies to scale up adoption of cleaner fuel alternatives may have greatly restricted real-world realization of these policy goals [30].

The low proportion of households using cleaner fuels, corresponding high use of wood fuel, and relatively high utility of switching from raw to processed biomass fuels (e.g., wood to charcoal) suggests the need to investigate the relative risk of LBW if households used other biomass fuels rather than wood. The aim of this study was therefore to determine the association between the use of wood compared to other types of biomass and kerosene

cooking fuels and the risk of LBW in Uganda, using the population-based 2016 Uganda Demographic and Health Survey (UDHS).

2. Materials and Methods

2.1. Study Design, Setting, and Data Source

The UDHS (2016) is a cross-sectional population-based national dataset funded by the U.S. Agency for International Development, with the birth recode (a file produced by DHS where each observation is an individual birth within the last five years) and relevant variables from the individual recode being extracted for this study [28]. The birth recodes contained birth history data, while individual recode (each observation is every woman within the survey) provided information on socio-demographic and household characteristics [28]. A two-stage stratified sampling methodology was employed to randomly select a representative sample [28]. Any woman residing in the selected household of reproductive age (15–49 years) was interviewed and asked to report their birth history (including live and still births) for the five years preceding the survey. A total of 19,588 households and 18,506 women were surveyed [28], with response rates of 97% (18,506/19,088) for the individual (women's) dataset and 67% (10,429/15,522) of the birth records respectively. Additional information regarding the UDHS has been described elsewhere [28].

2.2. Study Population

Singleton live births, occurring at term (≥ 37 weeks gestation) and/or pre-term (< 37 weeks gestation) which occurred in the last five years (2012–2016) from the time of interview were included in the study. Multiple births were excluded from the analysis because of the high risk of LBW among multiple pregnancies [31].

2.3. Modifications to the Wealth Index

The wealth index provided by DHS is calculated through principal component analysis (PCA), including assets, toilet facility, drinking water sources, cooking fuel, and house construction as predictor variables [32], with the final variable containing wealth quintiles (lowest, low, middle, high, and highest). As cooking fuel was the exposure of interest within this study, the wealth index (categorized as low, second, middle, fourth, and higher) was recalculated using the methods provided by the DHS [33] in SPSS [34] to remove cooking fuel to prevent circularity [35].

2.4. Exposure Variables

Self-reported main household cooking fuel was categorized into cleaner fuels (LPG), electricity, biogas, and no cooking), and biomass and kerosene fuels (kerosene, coal/lignite, charcoal, wood, straw/shrubs/grass, agricultural crop, animal dung). Households, where cooking was not done, were classified as using cleaner fuel because it was assumed that HAP levels in these households would be comparable to households that used cleaner fuels. The exposure variable was the biomass or kerosene cooking fuels, categorized into wood and other biomass and kerosene fuels (kerosene, charcoal, straw/shrubs/grass, agricultural crop, animal dung).

2.5. Outcome Variables

The outcome variable was LBW defined as a birthweight of less than 2500 g, obtained from either the health card (34%) or the maternal recall of child's weight at birth (66%).

2.6. Covariates

Covariates were identified from the literature [21,36–38] as those potentially associated with HAP or LBW. The covariates from household and contextual characteristics included age of the household head, access to electricity, place of residence (urban, rural), geographical region (central, east, north, west), household smoking status (yes, no), place of cooking

(in the main house, separate house, outdoors) and wealth index (low, second, middle, fourth, or highest). The 15 sub-regions in Uganda were categorized into four regions which included central, east, west, and north, which are defined based on ethnicity, poverty index, and geographical location [39]. Information from the respondents included age (15–19, 20–34, 35–49 years), level of education (no or primary education, secondary or tertiary). Pregnancy-related maternal covariates considered included parity (primigravida or multi-gravida), birth order (continuous variable), sex of the baby (male or female), and body mass index (BMI) (<18.5 or ≥ 18.5 kg/m²). Circumstantial pregnancy-related covariates included mode of delivery (spontaneous vaginal delivery or caesarian section), the timing of first antenatal care visit (ANC) (≤ 5 or >5 months), number of ANC visits (≥ 4 or <4), sulphadoxine-pyrimethamine (SP) (yes or no), birth interval (<24 months or ≥ 24 months), iron-folate supplementation (yes or no), deworming during pregnancy (yes or no), birth interval, based on WHO categorization, of less than 24 months or birth intervals of ≥ 24 months was used [40]. BMI, measured in kg/m², was categorized as low when BMI was <18.5 or normal when BMI was ≥ 18.5 [41], as there is a higher risk of LBW for BMI of less than 18.5 [42,43].

2.7. Data Analysis

Categorical variables were summarized using frequencies and proportions. Skewed continuous variables were summarized using the median and inter-quartile range (IQR), while normally distributed continuous variables were summarized using means and standard deviations. Bivariate and multivariate logistic regression, using survey commands to adjust for the complex sampling structure, was deployed to determine the association between exposure to wood and other forms of polluting fuels and LBW. The odds ratios (OR), 95% confidence intervals (95% CI), and *p*-values were reported. Clinically relevant variables, those with a *p*-value was less than 0.2, and variables without high levels of missing, in the bivariate analyses were included in the multivariate logistic regression model. Missing values were handled by case-wise deletion.

Sensitivity analyses were undertaken to ensure robustness of study findings and to further investigate confounding factors (e.g., BMI) that could not be accounted for in the main analysis due to a large proportion of missing data (Figure 1). Further stratified analyses were undertaken according to residence (rural or urban), cooking location (indoor, outdoor), and maternal BMI (≤ 18.5). Multivariable linear regression model was performed using birthweight as a continuous variable. Stata software (version 16.1) [44] was used to analyze the data.




Figure 1. Description of sensitivity analysis.

2.8. Ethical Approval and Authorisation

USAID obtained ethical approvals from the relevant authorities in Uganda to collect the data. Permission was obtained from the USAID to gain access to the anonymized and aggregated freely available dataset from the DHS online data archive [45].

3. Results

3.1. Descriptive Statistics

Out of 15,270 births recorded in the last five years, birthweights were available for 10,267 births (67.2%), with 986 births (9.6%) classified as LBW (Table 1). Of these births, cleaner fuels (LPG, biogas & electricity) were used in 0.3% ($n = 48$) households, while biomass fuels and kerosene were used in 99.6% ($n = 15,209$) of the households. Overall 11% ($n = 2168$) of the household respondents reported cooking to be performed inside the main residential house, of which 68% ($n = 1478$) did not have a separate room as a kitchen. The dominant household cooking fuel was mostly charcoal in urban areas ($n = 1880$, 68.2%) and wood in rural areas ($N = 6272$, 83.7%) (Figure 2). Cleaner fuels (LPG, electricity) were only observed for 39 births (1.4%) in urban and 10 births (0.1%) in rural areas. Risk of LBW was significantly associated with female sex (56% vs. 49%), pre-term delivery (28% vs. 13%), less than four antenatal visits (41% vs. 34%), a lower prevalence of sulphadoxine-pyrimethamine (77% vs. 81%) or deworming treatment (60% vs. 65%) (Table 2).

Table 1. Household characteristics by low birthweight classification of study participants.

	Birthweight < 2500 g	Birthweight \geq 2500 g	<i>p</i> -Value
	N (%)	N (%)	
	N = 986	N = 9281	
Cooking fuel			0.062
Wood	720 (73.3)	6393 (69.3)	
Other polluting fuels	262 (26.7)	2831 (30.7)	
Missing (%)	4 (0.4)	56 (0.6)	
Cooking location			0.942
In the house	100 (10.2)	967 (10.4)	
In a separate building	622 (63.1)	5774 (62.3)	
Outdoors	264 (26.8)	2521 (27.2)	
Missing (%)	264 (26.8)	0 (0.0)	
Household smoking			0.863
Yes	862 (87.5)	8137 (87.7)	
No	124 (12.5)	1144 (12.3)	
Type of place of residence			0.293
Urban	247 (25.0)	2522 (27.2)	
Rural	739 (75.0)	6759 (72.8)	
Region			0.096
Central	277 (28.1)	2719 (29.3)	
East	250 (25.3)	2354 (25.4)	
North	259 (26.3)	2071 (22.3)	
West	200 (20.3)	2137 (23.0)	
Electricity			0.011
Yes	700 (74.3)	6169 (69.0)	
No	243 (25.7)	2772 (31.0)	
Missing (%)	43 (4.4)	339 (3.7)	
Number of household members (listed)			0.568
Median (IQR)	6.0 (4.0, 8.0)	5.0 (4.0, 7.0)	
Combined wealth index			0.003
Low	248 (25.1)	1851 (19.9)	
Second	193 (19.6)	1662 (17.9)	
Middle	182 (18.4)	1632 (17.6)	
Fourth	158 (16.1)	1804 (19.4)	
Highest	205 (20.8)	2332 (25.1)	
Maternal characteristics			
Respondent's current age			<0.001
Median (IQR)	26.0 (22.0, 32.0)	27.0 (23.0, 32.0)	
Mother's education			0.010
No education/Primary	697 (70.7)	6072 (65.4)	
Secondary/higher	289 (29.3)	3208 (34.6)	

Table 1. Cont.

	Birthweight < 2500 g	Birthweight ≥ 2500 g	p-Value
	N (%)	N (%)	
	N = 986	N = 9281	
Mother’s BMI (kg/m²)			0.223
<18.5	19 (6.2)	253 (8.3)	
≥18.5	296 (93.8)	2795 (91.7)	
Missing (%)	670 (68.0)	6233 (67.2)	

Footnotes: IQR = Inter quartile range.




Figure 2. Description of cooking fuel use in rural and urban settings. Other biomass and kerosene = kerosene, charcoal, straw/shrubs/grass, agricultural crop, animal dung.

Table 2. Birth characteristics by low birthweight classification of study participants.

Variable	Birthweight < 2500 g	Birthweight ≥ 2500 g	p-Value
	N (%)	N (%)	
	N = 986	N = 9281	
Parity			0.327
Once	170 (17.2)	1466 (15.8)	
More than once	816 (82.8)	7814 (84.2)	
Birth order number			
Median (IQR)	2.0 (1.0, 5.0)	3.0 (2.0, 5.0)	
Sex of child			<0.001
Male	438 (44.4)	4761 (51.3)	
Female	548 (55.6)	4520 (48.7)	
Birth Interval			0.576
≤24 months	824 (83.5)	7681 (82.8)	
>24 months	162 (16.5)	1599 (17.2)	
Duration of pregnancy			<0.001
Pre-term	274 (27.8)	1184 (12.8)	
Term	711 (72.2)	8097 (87.2)	

Table 2. Cont.

Variable	Birthweight < 2500 g	Birthweight ≥ 2500 g	p-Value
	N (%)	N (%)	
	N = 986	N = 9281	
Timing of first ANC visits			0.066
<5 months gestation	387 (65.1)	3896 (60.8)	
≥5 months gestation	207 (34.9)	2509 (39.2)	
Missing (%)	391 (39.7)	2875 (31.0)	
Number of ANC visits			0.003
<4	252 (41.4)	2206 (34.3)	
≥4	357 (58.6)	4225 (65.7)	
Missing (%)	377 (38.3)	2849 (30.7)	
Place of delivery			0.131
Health facility	906 (92.5)	8655 (93.9)	
Home	73 (7.5)	561 (6.1)	
Missing (%)	7 (0.7)	64 (0.7)	
Delivery by caesarean section			0.763
No	895 (91.4)	8479 (91.8)	
Yes	84 (8.6)	760 (8.2)	
Missing (%)	7 (0.7)	42 (0.4)	
Iron supplementation			0.131
No	66 (10.8)	562 (8.7)	
Yes	545 (89.2)	5906 (91.3)	
Missing (%)	375 (38.0)	2812 (30.3)	
Sulphadoxine-pyrimethamine			0.014
Yes	467 (76.8)	5251 (81.4)	
No	141 (23.2)	1199 (18.6)	
Missing (%)	378 (38.4)	2831 (30.5)	
Deworming			0.021
Yes	364 (59.7)	4179 (65.2)	
No	246 (40.3)	2226 (34.8)	
Missing (%)	376 (38.1)	2876 (31.0)	

3.2. Association between Type of Biomass Cooking Fuels with LBW

In the bivariate analysis, a borderline association was observed between LBW in households using wood cooking compared to other biomass fuels (OR: 0.82; 95% CI: 0.67–1.00) (Table 3). However, after adjustment for socio-demographic and obstetric factors, the association was no longer significant (AOR: 0.94; 95% CI: 0.72–1.22). A number of the covariates were associated with LBW risk, including the size of household, where for every additional household member the odds ratio of LBW increased by 3% (AOR: 1.03; 95% CI: 1.00–1.07). Having a higher level of maternal education was associated with lower odds ratio of LBW in other polluting fuel-cooking (AOR: 0.80; 95% CI: 0.64–1.00) households compared to wood-cooking households, as was being female (AOR: 1.32; 95% CI: 1.13–1.55) compared to being male. Pregnancies that lasted longer than 37 weeks were also associated with lower odds of LBW (AOR: 0.39; 95% CI: 0.31–0.49) compared to pregnancies that were less than 37 weeks gestation. Likewise, the fourth highest wealth index was associated with lower odds of LBW compared to the lowest wealth index (AOR: 0.69; 95% CI: 0.5–0.96)

Table 3. Unadjusted and adjusted analysis for the association between LBW and type of biomass cooking fuel.

	Unadjusted Analysis			Adjusted Analysis (N = 9863)		
	UOR	95% CI	p-Value	AOR	95% CI	p-Value
Household Characteristics						
Cooking fuel						
Wood	Ref.			Ref.		
Other polluting fuels	0.82	(0.67, 1.00)	0.053	0.94	(0.72, 1.22)	0.646
Cooking location						
In the house	Ref.			Ref.		
In a separate building	1.04	(0.75, 1.43)	0.818	0.99	(0.71, 1.36)	0.928
Outdoors	1.01	(0.71, 1.43)	0.959	0.94	(0.67, 1.30)	0.691
Household smoking						
Yes	Ref.			Ref.		
No	1.02	(0.82, 1.27)	0.863	0.92	(0.73, 1.15)	0.470
Type of place of residence						
Urban	Ref.			Ref.		
Rural	1.12	(0.91, 1.37)	0.294	0.90	(0.70, 1.15)	0.400
Region						
Central	Ref.			Ref.		
East	1.04	(0.83, 1.30)	0.726	0.76	(0.59, 0.98)	0.035
North	1.23	(0.98, 1.54)	0.076	0.75	(0.57, 0.99)	0.042
West	0.92	(0.71, 1.19)	0.525	0.82	(0.62, 1.06)	0.134
Electricity						
Yes	Ref.			Ref.		
No	1.30	(1.06, 1.58)	0.011	0.94	(0.73, 1.22)	0.655
Number of household members (listed)						
	1.01	(0.99, 1.04)	0.418	1.03	(1.00, 1.07)	0.027
Combined wealth index						
Low	Ref.			Ref.		
Second	0.87	(0.70, 1.08)	0.214	0.92	(0.71, 1.18)	0.500
Middle	0.83	(0.66, 1.04)	0.108	0.89	(0.67, 1.19)	0.435
Fourth	0.66	(0.51, 0.84)	0.001	0.69	(0.50, 0.96)	0.027
Highest	0.66	(0.50, 0.85)	0.002	0.73	(0.50, 1.08)	0.120
Maternal characteristics						
Respondent's current age						
	0.98	(0.97, 0.99)	0.001	0.99	(0.97, 1.01)	0.403
Mother's education						
No education/Primary only	Ref.			Ref.		
Secondary only/higher	0.78	(0.65, 0.94)	0.010	0.80	(0.64, 1.00)	0.050
Mother's BMI (Kg/m²)						
<18.5	Ref.			Ref.		
≥18.5	1.37	(0.82, 2.29)	0.225			
Birth characteristics						
Parity						
Once	Ref.			Ref.		
More than once	0.90	(0.73, 1.11)	0.327	1.18	(0.89, 1.55)	0.249
Birth order number						
	0.95	(0.92, 0.99)	0.008	0.96	(0.90, 1.02)	0.164
Sex of child						
Male	Ref.			Ref.		
Female	1.32	(1.13, 1.53)	<0.001	1.32	(1.13, 1.55)	0.001
Birth Interval						
≤24 months	Ref.			Ref.		
>24 months	0.95	(0.78, 1.15)	0.576	0.95	(0.77, 1.16)	0.597
Duration of pregnancy						
Pre-term	Ref.			Ref.		
term	0.38	(0.31, 0.46)	<0.001	0.39	(0.31, 0.49)	<0.001
Timing of ANC first visits						
<5 months gestation	Ref.			Ref.		
≥5 months gestation	0.83	(0.68, 1.01)	0.066			

Table 3. Cont.

	Unadjusted Analysis			Adjusted Analysis (N = 9863)		
	UOR	95% CI	p-Value	AOR	95% CI	p-Value
Number of ANC visits						
<4 times	Ref.					
≥4 times	0.74	(0.61, 0.90)	0.003			
Place of delivery						
Health facility	Ref.			Ref.		
Home	1.25	(0.94, 1.67)	0.131	1.21	(0.90, 1.62)	0.211
Delivery by caesarean section						
No	Ref.			Ref.		
Yes	1.05	(0.78, 1.41)	0.763	1.16	(0.87, 1.56)	0.313
Iron supplementation						
No	Ref.					
Yes	0.78	(0.57, 1.08)	0.132			
Sulphadoxine-pyrimethamine						
Yes	Ref.					
No	1.32	(1.06, 1.65)	0.014			
Deworming						
Yes	Ref.					
No	1.27	(1.04, 1.55)	0.021			

Footnotes: UOR = unadjusted odds ratio, AOR = adjusted odds ratio, 95% CI = 95% confidence interval, Ref = reference group.

3.3. Sensitivity Analyses

Place of residence, cooking location, and adjustment for maternal BMI in sensitivity analyses all had no observed association of LBW between births from wood cooking compared to other biomass cooking households (Table 4). Birthweight was explored as a continuous variable, with births from other polluting fuels-cooking households having a 29.43 g higher (95% CI: −38.15, 76.01) birthweight compared to wood cooking, however, this was not statistically significant ($p = 0.393$). In the sensitivity analyses, which included the addition of all available confounders including mothers BMI, the timing of ANC visit, and number of ANC visits, no significant association (AOR = 0.78 (95% CI: 0.46, 1.31) was found among households where wood compared to other biomass fuels were used.

Table 4. Exploratory and sensitivity analysis.

	AOR	95% CI	p-Value
Place of residence			
Urban (N = 2274)	1.09	(0.65, 1.83)	0.745
Rural (N = 7589)	0.85	(0.61, 1.18)	0.322
Cooking location			
Indoor (N = 7190)	1.04	(0.77, 1.42)	0.786
Outdoor (N = 2682)	0.77	(0.49, 1.19)	0.235
Adjusting for normal maternal BMI			
BMI ≤ 18 (N = 2655)	0.92	(0.59, 1.44)	0.726
Adjusting for all confounders			
N = 2167	0.78	(0.46, 1.31)	0.348
Birthweight as a continuous variable			
N = 9863	29.43	(−38.15, 97.01)	0.393

Footnotes: AOR = adjusted odds ratio, 95% CI = 95% confidence interval. Adjusted for cooking location, household smoking, type of place of residence, region, electricity, wealth index, respondents current age, mother's education, mothers BMI, parity, birth order, sex of child, birth interval, duration of pregnancy, the timing of ANC visit, number of ANC visits, place of delivery and mode of delivery.

4. Discussion

This population-based study of 15,270 live-births and a high response rate (97%) in Uganda was conducted to determine the association between wood compared to kerosene and other biomass cooking fuels with the risk of LBW. No evidence of an association was observed between the different types of biomass and kerosene fuels and risk of LBW (AOR:

0.94; 95% CI: 0.72, 1.22) after adjustment for confounding factors. In the sensitivity analyses which considered the associated effect of residence (rural, urban), cooking location (indoor, outdoor), maternal BMI, and birthweight as a continuous dependent variable with risk of LBW, findings similar to the main analyses were found.

LBW was observed to occur among almost 10% of all deliveries which, although lower than the 14% previous estimates for Sub-Saharan Africa [10], reflects a major public health concern. LBW, which may partly result from HAP exposure, is one of the leading causes of neonatal mortality in developing countries [9], which therefore calls for further research, innovations, and policies to reduce the health burden of LBW brought by use of biomass cooking fuels.

To the best of our knowledge, this was the first study in Uganda to determine the association between the different types of biomass fuels and LBW, a setting where biomass fuels were used by almost all (99.6%) of the population; limiting the possibility of comparing biomass fuels to cleaner fuels. The DHS collects robust, nationally representative data, however, there were a large (32.8%) number of missing birthweights recorded. However, 67.2% of birthweights were recorded totaling 15,270 live-births. Birthweights obtained by maternal recall may have been subject to recall bias, but this was reduced by limiting the data collection period to births within the last five years. Although there was no evidence of an effect in our study and the reference cooking fuel differs to that of other studies, they are similar to those obtained in single-country analyses investigating the difference between biomass and cleaner fuels in Malawi (AOR, 1.29 (95% CI: 0.34; 4.48) [18]), Ethiopia (AOR, 1.3 (95% 0.9–1.9) [20]), Nigeria (−0.09 (95% CI: −0.31, 0.10) [17]), Bangladesh (AOR 0.96 (95% CI: 0.81–1.13) [46]), and India (AOR: 1.07(95% CI: 0.94–1.22) [47]). On the contrary, findings in this study were inconsistent with the population-based study from Zimbabwe which showed biomass fuel use was associated with a 175 g lower birthweight (95% CI: −300 to −50 g) [22]. While solid biomass fuels significantly increased the risk of LBW by 35% in a meta-analysis of 19 studies, the association was not significant for a sub-group analysis of studies from Africa [7]. Similarly, a systematic review of DHS studies found that two studies out of five from African countries (Malawi and Zimbabwe) cited no evidence of an association between biomass cooking fuels and LBW, which was consistent with findings in this study [48]. In African settings like Uganda where biomass cooking fuels are almost universally used and the consequent small proportion that use cleaner fuels, direct measures of exposure to HAP may be more suited to investigate the effect of HAP on maternal and child health.

There is evidence for the biological plausibility of LBW risk associated with in-utero HAP exposure including PM_{2.5}, CO, and aromatic hydrocarbons [12,49]. Increased risk of LBW has been observed among infants exposed to second-hand smoke and HAP exposure in animals [7,12,50]. Exposure to polluting fuels is proposed to cause LBW by reducing fetal growth [7,49]. Reduced fetal growth occurs through impaired maternal lung function, fetal toxicity, endocrine disruption, pathological placental changes, oxidative processes, and reduced oxygen and micronutrients supply to the fetus [7,12,16,49,51]. The difference in pollutant levels between wood and other biomass may not differ or reduce enough to produce an effect, along with households often switching between biomass fuel types. In addition, the WHO has stated in the updated AQGs that there is no safe level of PM [2] and levels of CO remain high [2]; therefore, barriers should be broken down to allow access to cleaner alternatives.

Charcoal was predominantly used in urban settings, and wood was mostly used in rural settings, but no difference in odds of LBW risk was observed between rural and urban areas. In settings where biomass fuels are mostly used and access to cleaner fuels remains limited, feasible mitigation measures should be promoted to reduce exposure to HAP among vulnerable women and children under five years of age [24]. The mitigation measures such as cooking outdoors, keeping children away from cooking places, opening doors and windows, improving housing and ventilation conditions, use of cured wood or charcoal, and building kitchens with chimneys might play a vital role in modifying fuel user

behaviors, the environment, and the source of HAP [5,52]. These addressable mitigation measures may not only reduce exposure to HAP but would also create awareness and facilitate uptake of long-term active measures such as scaling up the adoption of cleaner fuels [53]. In the long term, transitioning to cleaner fuels should be prioritized as the ultimate solution to reducing the impact of HAP.

Although improved cookstoves can still emit levels of PM_{2.5} and CO which exceed WHO-AQGs [54], a reduced risk of LBW has been observed with the use of improved cookstoves compared to traditional stoves in a systematic review and meta-analysis [55]. Government can prioritize the adoption of improved cookstoves as the first step in transitioning to cleaner fuels [5]. The promotion of cleaner fuels requires government commitment, strong energy fuel policies, subsidies, and tax exemptions on modern fuels [5]. Programs should prioritize pregnant women to reduce the adverse effects of HAP on pregnancy [56].

Proxy measures may be reliable and valid ways of estimating HAP exposure, however, the inability to use direct measures of HAP may have resulted in misclassification bias and inaccurate estimation of HAP. Further DHS data collection should endeavor to check birthweight against health card and capture use of secondary cooking fuels, stove type, alongside fuel and stove stacking practices [47]. Future studies should ideally use a direct measurement of personal exposure to HAP to investigate the association between biomass fuels and LBW. Direct measures of HAP would help determine an exposure-response relationship. Causality could not be determined in this study, therefore, robust study designs such as prospective cohort studies or if feasible, randomized controlled trials should be used to infer causality. We recommend exploration of other sources of HAP on LBW risk, including outdoor burning of garbage in the household including burning of solid biomass fuels and wastes from the roadsides and use of polluting fuels for lighting.

5. Conclusions

In this study, almost all (99.6%) study households in Uganda used biomass and kerosene fuels for domestic cooking, with LBW observed among 9.6% of singleton live births. After adjusting for household, maternal and demographic confounding factors, no evidence of an association was observed between risk of LBW and the use of wood fuel compared to other biomass or kerosene cooking fuels. Although previous evidence has indicated reduced risks of LBW with cleaner fuels, this requires further investigation, as the very low proportion of LPG, electricity or biogas use meant a comparison of risks associated with biomass fuels could not be undertaken in this present study. Achieving energy policy targets for adoption of cleaner fuels by 2040 in Uganda remains a key policy priority given the existing high reliance upon polluting biomass fuels. Public health initiatives to reduce LBW prevalence should seek to reduce socio-economic inequity and increase levels of female education in this context.

Author Contributions: Conceptualization, J.E., K.E.W. and G.N.T.; methodology, J.E., K.E.W. and G.N.T.; formal analysis, J.E. and K.E.W.; writing—original draft preparation, J.E.; writing—review and editing, K.E.W., S.E.B. and G.N.T.; supervision, K.E.W. and G.N.T. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The data presented in this study are available on request from the corresponding author.

Acknowledgments: We are grateful to Demographic and Health Surveys for granting access to the UDHS data. We are grateful to Raoul Reulen and Gavin Rudge for their review comments during the development of the protocol and thesis.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Stoner, O.; Lewis, J.; Martínez, I.L.; Gumy, S.; Economou, T.; Adair-Rohani, H. Household cooking fuel estimates at global and country level for 1990 to 2030. *Nat. Commun.* **2021**, *12*, 5793. [CrossRef] [PubMed]
2. WHO. Particulate matter (PM2.5 and PM10), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide. In *WHO Global Air Quality Guidelines*; World Health Organization: Geneva, Switzerland, 2021.
3. Smith, K.R.; Bruce, N.; Balakrishnan, K.; Adair-Rohani, H.; Balmes, J.; Chafe, Z.; Dherani, M.; Hosgood, H.D.; Mehta, S.; Pope, D. Millions dead: How do we know and what does it mean? Methods used in the comparative risk assessment of household air pollution. *Annu. Rev. Public Health* **2014**, *35*, 185–206. [CrossRef] [PubMed]
4. Lee, K.K.; Bing, R.; Kiang, J.; Bashir, S.; Spath, N.; Stelzle, D.; Mortimer, K.; Bularga, A.; Doudesis, D.; Joshi, S.S. Adverse health effects associated with household air pollution: A systematic review, meta-analysis, and burden estimation study. *Lancet Glob. Health* **2020**, *8*, e1427–e1434. [CrossRef]
5. Avis, W.M.S.; Singh, A. Air Pollution Exposure in Low Income Households in Kampala. ASAP East Africa Vulnerability Scoping Study No. 6. University of Birmingham: Birmingham, UK, 2018. Available online: <https://assets.publishing.service.gov.uk> (accessed on 11 February 2022).
6. Okello, G.; Devereux, G.; Semple, S. Women and girls in resource poor countries experience much greater exposure to household air pollutants than men: Results from Uganda and Ethiopia. *Environ. Int.* **2018**, *119*, 429–437. [CrossRef] [PubMed]
7. Amegah, A.K.; Quansah, R.; Jaakkola, J.J. Household air pollution from solid fuel use and risk of adverse pregnancy outcomes: A systematic review and meta-analysis of the empirical evidence. *PLoS ONE* **2014**, *9*, e113920. [CrossRef]
8. Younger, A.; Alkon, A.; Harknett, K.; Louis, R.J.; Thompson, L.M. Adverse birth outcomes associated with household air pollution from unclean cooking fuels in low-and middle-income countries: A systematic review. *Environ. Res.* **2021**, *204*, 112274. [CrossRef] [PubMed]
9. Oza, S.; Lawn, J.E.; Hogan, D.R.; Mathers, C.; Cousens, S.N. Neonatal cause-of-death estimates for the early and late neonatal periods for 194 countries: 2000–2013. *Bull. World Health Organ.* **2014**, *93*, 19–28. [CrossRef]
10. Blencowe, H.; Krusevec, J.; De Onis, M.; Black, R.E.; An, X.; Stevens, G.A.; Borghi, E.; Hayashi, C.; Estevez, D.; Cegolon, L. National, regional, and worldwide estimates of low birthweight in 2015, with trends from 2000: A systematic analysis. *Lancet Glob. Health* **2019**, *7*, e849–e860. [CrossRef]
11. Misra, P.; Srivastava, R.; Krishnan, A.; Sreenivaas, V.; Pandav, C.S. Indoor air pollution-related acute lower respiratory infections and low birthweight: A systematic review. *J. Trop. Pediatrics* **2012**, *58*, 457–466. [CrossRef]
12. Pope, D.P.; Mishra, V.; Thompson, L.; Siddiqui, A.R.; Rehfuess, E.A.; Weber, M.; Bruce, N.G. Risk of low birth weight and stillbirth associated with indoor air pollution from solid fuel use in developing countries. *Epidemiol. Rev. Environ. Health Perspect.* **2010**, *32*, 70–81. [CrossRef]
13. Amegah, A.K.; Jaakkola, J.J.; Quansah, R.; Norgbe, G.K.; Dzodzomenyo, M. Cooking fuel choices and garbage burning practices as determinants of birth weight: A cross-sectional study in Accra, Ghana. *J. Environ. Health* **2012**, *11*, 78. [CrossRef]
14. Adeyemi, R.A.; Zewotir, T.; Ramroop, S. Semiparametric multinomial ordinal model to analyze spatial patterns of child birth weight in Nigeria. *Int. J. Environ. Res. Public Health* **2016**, *13*, 1145. [CrossRef] [PubMed]
15. Alexander, D.A.; Northcross, A.; Karrison, T.; Morhasson-Bello, O.; Wilson, N.; Atalabi, O.M.; Dutta, A.; Adu, D.; Ibigbami, T.; Olamijulo, J. Pregnancy outcomes and ethanol cook stove intervention: A randomized-controlled trial in Ibadan, Nigeria. *Environ. Int.* **2018**, *111*, 152–163. [CrossRef]
16. Arinola, G.O.; Dutta, A.; Oluwole, O.; Olopade, C.O. Household air pollution, levels of micronutrients and heavy metals in cord and maternal blood, and pregnancy outcomes. *Int. J. Environ. Res. Public Health Engl.* **2018**, *15*, 2891. [CrossRef] [PubMed]
17. Roberman, J.; Emeto, T.I.; Adegboye, O.A. Adverse birth outcomes due to exposure to household air pollution from unclean cooking fuel among women of reproductive age in Nigeria. *Int. J. Environ. Res. Public Health* **2021**, *18*, 634. [CrossRef] [PubMed]
18. Milanzi, E.B.; Namacha, N.M. Maternal biomass smoke exposure and birth weight in Malawi: Analysis of data from the 2010 Malawi Demographic and Health Survey. *Malawi Med. J.* **2017**, *29*, 160–165. [CrossRef]
19. Admasie, A.; Kumie, A.; Worku, A. Association of household fuel type, kitchen characteristics and house structure with child size at birth in Wolaita Sodo, Southern Ethiopia. *Open Public Health J.* **2018**, *11*, 298–308. [CrossRef]
20. Kanno, G.G.; Anbesse, A.T.; Shaka, M.F.; Legesse, M.T.; Andarge, S.D. Investigating the effect of biomass fuel use and kitchen location on maternal report of birth size: A Cross-Sectional Analysis of 2016 Ethiopian Demographic Health Survey data. *Public Health Pract.* **2021**, *2*, 100211. [CrossRef]
21. Demelash, H.; Motbainor, A.; Nigatu, D.; Gashaw, K.; Melese, A. Risk factors for low birth weight in Bale zone hospitals, South-East Ethiopia: A case-control study. *BMC Pregnancy Childbirth* **2015**, *15*, 264. [CrossRef] [PubMed]
22. Mishra, V.; Dai, X.; Smith, K.R.; Mika, L. Maternal exposure to biomass smoke and reduced birth weight in Zimbabwe. *Ann. Epidemiol.* **2004**, *14*, 740–747. [CrossRef]
23. Weber, E.; Adu-Bonsaffoh, K.; Vermeulen, R.; Klipstein-Grobusch, K.; Grobbee, D.E.; Browne, J.L.; Downward, G.S. Household fuel use and adverse pregnancy outcomes in a Ghanaian cohort study. *J. Reprod. Health* **2020**, *17*, 29. [CrossRef] [PubMed]
24. Woolley, K.E.; Bagambe, T.; Singh, A.; Avis, W.R.; Kabera, T.; Weldetinsae, A.; Mariga, S.T.; Kirenga, B.; Pope, F.D.; Thomas, G.N. Investigating the association between wood and charcoal domestic cooking, respiratory symptoms and acute respiratory infections among children aged under 5 Years in Uganda: A Cross-sectional analysis of the 2016 Demographic and Health Survey. *Int. J. Environ. Res. Public Health* **2020**, *17*, 3974. [CrossRef]

25. Coker, E.; Katamba, A.; Kizito, S.; Eskenazi, B.; Davis, J.L. Household air pollution profiles associated with persistent childhood cough in urban Uganda. *Environ. Int.* **2020**, *136*, 105471. [CrossRef] [PubMed]
26. Van Gemert, F.; Chavannes, N.; Nabadda, N.; Luzige, S.; Kirenga, B.; Eggermont, C.; de Jong, C.; van der Molen, T. Impact of chronic respiratory symptoms in a rural area of sub-Saharan Africa: An in-depth qualitative study in the Masindi district of Uganda. *Prim. Care Respir. J.* **2013**, *22*, 300–305. [CrossRef] [PubMed]
27. The World Bank Group. Uganda | Data. Available online: <https://data.worldbank.org/country/uganda> (accessed on 8 March 2021).
28. Uganda Bureau of Statistics (UBOS); ICF. *Uganda Demographic and Health Survey 2016*; UBOS: Kampala, Uganda; ICF: Rockville, MD, USA, 2018.
29. Uganda Human Development Reports. Available online: http://hdr.undp.org/sites/all/themes/hdr_theme/country-notes/UGA.pdf (accessed on 1 September 2021).
30. Ministry of Energy and Mineral Development. The Renewable Energy Policy for Uganda. 2007. Available online: https://www.era.go.ug/index.php/download-repo/doc_details/209-era-ceo-s-keynote-speech-at-the-era-getfit-uganda-solar-pv-award-event?tmpl=component (accessed on 1 September 2021).
31. Vohr, B.R.; Tyson, J.E.; Wright, L.L.; Perritt, R.L.; Li, L.; Poole, W.K.; Network, N.N.R. Maternal age, multiple birth, and extremely low birth weight infants. *J. Pediatrics* **2009**, *154*, 498–503.e492. [CrossRef] [PubMed]
32. Rutstein, S.O.; Johnson, K. *The DHS Wealth Index*; DHS Comparative Reports No. 6.; ORC Macro: Calverton, MD, USA, 2004.
33. Rutstein, S.O. *Steps to Constructing the New DHS Wealth Index*; ICF International: Rockville, MA, USA, 2015.
34. Corp, I.S. *IBM SPSS Statistics for Windows. Version 27.0*; IBM: Armonk, NY, USA, 2020.
35. Tusting, L.S.; Bottomley, C.; Gibson, H.; Kleinschmidt, I.; Tatem, A.J.; Lindsay, S.W.; Gething, P.W. Housing improvements and malaria risk in sub-Saharan Africa: A multi-country analysis of survey data. *PLoS Med.* **2017**, *14*, e1002234. [CrossRef]
36. Manyeh, A.K.; Kukula, V.; Odonkor, G.; Ekey, R.A.; Adjei, A.; Narh-Bana, S.; Akpakli, D.E.; Gyapongi, M. Socioeconomic and demographic determinants of birth weight in southern rural Ghana: Evidence from Dodowa Health and Demographic Surveillance System. *BMC Pregnancy Childbirth* **2016**, *16*, 160. [CrossRef]
37. Dahlui, M.; Azahar, N.; Oche, O.M.; Aziz, N.A. Risk factors for low birth weight in Nigeria: Evidence from the 2013 Nigeria Demographic and Health Survey. *Glob. Health Action* **2016**, *9*, 28822. [CrossRef]
38. Ahammed, B.; Maniruzzaman, M.; Ferdousi, F.; Abedin, M.M.; Hossain, M.T. Socioeconomic and demographic factors associated with low birth weight in Nepal: Data from 2016 Nepal demographic and health survey. *Soc. Health Behav.* **2020**, *3*, 158. [CrossRef]
39. Poverty Maps of Uganda-Uganda Bureau of Statistics. Available online: https://www.ubos.org/wp-content/uploads/publications/02_2020Poverty_Map_report_Oct_2019.pdf (accessed on 1 September 2021).
40. World Health Organization. *Report of a WHO Technical Consultation on Birth Spacing: Geneva, Switzerland 13–15 June 2005*; World Health Organization: Geneva, Switzerland, 2007.
41. World Health Organization. *Global Nutrition Targets 2025: Low Birth Weight Policy Brief*; World Health Organization: Geneva, Switzerland, 2014.
42. Mahumud, R.A.; Sultana, M.; Sarker, A.R. Distribution and determinants of low birth weight in developing countries. *J. Prev. Med. Public Health* **2017**, *50*, 18. [CrossRef]
43. McDonald, S.D.; Han, Z.; Mulla, S.; Beyene, J. Overweight and obesity in mothers and risk of preterm birth and low birth weight infants: Systematic review and meta-analyses. *BMJ* **2010**, *341*, c3428. [CrossRef] [PubMed]
44. StataCorp. *Stata Statistical Software: Release 16*; StataCorp LLC.: College Station, TX, USA, 2019.
45. The Demographic and Health Survey Program (DHS) The DHS Program US-AID. Available online: <https://dhsprogram.com/> (accessed on 20 December 2021).
46. Khan, M.N.; Islam, M.M.; Islam, M.R.; Rahman, M.M. Household air pollution from cooking and risk of adverse health and birth outcomes in Bangladesh: A nationwide population-based study. *Environ. Health Perspect.* **2017**, *16*, 57. [CrossRef] [PubMed]
47. Sreeramareddy, C.T.; Shidhaye, R.R.; Sathiakumar, N. Association between biomass fuel use and maternal report of child size at birth—an analysis of 2005–06 India Demographic Health Survey data. *BMC Public Health* **2011**, *11*, 403. [CrossRef] [PubMed]
48. Odo, D.B.; Yang, I.A.; Knibbs, L.D. A systematic review and appraisal of epidemiological studies on household fuel use and its health effects using demographic and health surveys. *Int. J. Environ. Res. Public Health* **2021**, *18*, 1411. [CrossRef] [PubMed]
49. Fullerton, D.G.; Bruce, N.; Gordon, S.B. Indoor air pollution from biomass fuel smoke is a major health concern in the developing world. *Trans. R. Soc. Trop. Med. Hyg.* **2008**, *102*, 843–851. [CrossRef]
50. Boy, E.; Bruce, N.; Delgado, H. Birth weight and exposure to kitchen wood smoke during pregnancy in rural Guatemala. *Environ. Health Perspect.* **2002**, *110*, 109–114. [CrossRef] [PubMed]
51. Amegah, A.K.; Jaakkola, J.J. Household air pollution and the sustainable development goals. *Bull. World Health Organ.* **2016**, *94*, 215. [CrossRef] [PubMed]
52. Woolley, K.E.; Dickinson-Craig, E.; Bartington, S.E.; Oludotun, T.; Kirenga, B.; Mariga, S.T.; Kabera, T.; Coombe, A.; Pope, F.D.; Singh, A. Effectiveness of interventions to reduce household air pollution from solid biomass fuels and improve maternal and child health outcomes in low-and middle-income countries: A systematic review protocol. *Syst. Rev.* **2021**, *10*, 73–90. [CrossRef]
53. Barnes, B.R. Behavioural change, indoor air pollution and child respiratory health in developing countries: A review. *Int. J. Environ. Res. Public Health Engl.* **2014**, *11*, 4607–4618. [CrossRef]
54. Rosenthal, J.; Quinn, A.; Grieshop, A.P.; Pillarisetti, A.; Glass, R.I. Clean cooking and the SDGs: Integrated analytical approaches to guide energy interventions for health and environment goals. *J. Energy Sustain. Dev.* **2018**, *42*, 152–159. [CrossRef]

55. Woolley, K.E.; Dickinson-Craig, E.; Lawson, H.L.; Sheikh, J.; Day, R.; Pope, F.D.; Greenfield, S.M.; Bartington, S.E.; Warburton, D.; Manaseki-Holland, S. Effectiveness of interventions to reduce household air pollution from solid biomass fuels and improve maternal and child health outcomes in low-and middle-income countries: A systematic review and meta-analysis. *Indoor Air* **2022**, *32*, e12958. [[PubMed](#)]
56. Vakalopoulos, A.; Dharmage, S.C.; Dharmaratne, S.; Jayasinghe, P.; Lall, O.; Ambrose, I.; Weerasooriya, R.; Bui, D.S.; Yasaratne, D.; Heyworth, J. Household air pollution from biomass fuel for cooking and adverse fetal growth outcomes in rural Sri Lanka. *Int. J. Environ. Res. Public Health* **2021**, *18*, 1878. [[CrossRef](#)] [[PubMed](#)]