

Asteroseismology of the red-giant hosts KOI-3886 and iota Draconis

Campante, Tiago; Li, Tanda; Ong, J. M. Joel; Corsaro, Enrico; Hill, Michelle L.; Lillo-box, Jorge; Bedding, Timothy R.; Bossini, Diego; Brandt, Timothy D.; Breton, Sylvain N.; Buzasi, Derek L.; Chaplin, Bill; Cunha, Margarida S.; García, Rafael A.; Hon, Marc; Huber, Daniel; Jiang, Chen; Kane, Stephen R.; Kayhan, Cenk; Kuszlewicz, James

DOI:
[10.5281/zenodo.5127814](https://doi.org/10.5281/zenodo.5127814)

License:
Creative Commons: Attribution (CC BY)

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (Harvard):
Campante, T, Li, T, Ong, JMJ, Corsaro, E, Hill, ML, Lillo-box, J, Bedding, TR, Bossini, D, Brandt, TD, Breton, SN, Buzasi, DL, Chaplin, B, Cunha, MS, García, RA, Hon, M, Huber, D, Jiang, C, Kane, SR, Kayhan, C, Kuszlewicz, J, Mathur, S, Pereira, F, Santos, NC, Serenelli, AM & Stello, D 2021, Asteroseismology of the red-giant hosts KOI-3886 and iota Draconis. in *TESS Science Conference 2: Posters from the TESS Science Conference II (TSC2), held virtually Aug 2 - 6 2021.*, 5127814, Zenodo, TESS Science Conference II, 2/08/21. <https://doi.org/10.5281/zenodo.5127814>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

ASTEROSEISMOLOGY OF THE RED-GIANT HOSTS KOI-3886 AND ι DRACONIS

T. L. Campante[†], T. Li, J. M. J. Ong, E. Corsaro, M. L. Hill, J. Lillo-Box, T. R. Bedding, D. Bossini, T. D. Brandt, S. N. Breton, D. L. Buzasi, W. J. Chaplin, M. S. Cunha, R. A. García, M. Hon, D. Huber, C. Jiang, S. R. Kane, C. Kayhan, J. S. Kuszlewicz, S. Mathur, F. Pereira, N. C. Santos, A. Serenelli, and D. Stello


[†] E-mail: tiago.campante@astro.up.pt

ABSTRACT

Kepler asteroseismology has played an important role in the characterization of host stars and their planetary systems. Target selection biases, however, meant that this synergy would remain mostly confined to main-sequence stars. The advent of *TESS* has since lifted this restriction, enabling the systematic search for transiting planets around seismic giants, as well as revisiting previously known evolved hosts (mostly in radial-velocity systems) using asteroseismology.


Here, we present the detailed asteroseismology of two high-luminosity red-giant branch hosts, KOI-3886 and ι Draconis. KOI-3886, observed by *Kepler* over 4 years and later by *TESS* over 1 sector (27.4 days), has been a longtime candidate host [1]. ι Dra, observed by *TESS* over 5 sectors, hosts a planet (ι Dra b) in a highly eccentric ~ 511 -day-period orbit [2].

PHOTOMETRY




The top panel shows a segment (Quarter 9) of the long-cadence *Kepler* light curve of KOI-3886, optimized for asteroseismology [3]. The bottom panel shows a segment (Sectors 22–24) of the *TESS* light curve of ι Dra, acquired at a 2-minute cadence. To deal with the latter target’s saturated nature, a large custom aperture had to be adopted [4].

GLOBAL OSCILLATION PARAMETERS



Power spectral density (PSD) of KOI-3886 (left) and ι Dra (right). Power spectra are shown in gray (smoothed version in black). Solid red curves are fits to the background [5], consisting of two Harvey-like profiles (blue dot-dashed curves) and a white noise offset (yellow dot-dashed line). Solar-like oscillations can be seen at $\sim 50 \mu\text{Hz}$ and $\sim 40 \mu\text{Hz}$, respectively. We measured the large frequency separation, $\Delta\nu$, and the frequency of maximum oscillation amplitude, ν_{max} , using a range of automated methods [6]. Consolidated values for KOI-3886 (ι Dra) are $\Delta\nu = 4.60 \pm 0.20 \mu\text{Hz}$ and $\nu_{\text{max}} = 46.9 \pm 0.3 \mu\text{Hz}$ ($\Delta\nu = 4.02 \pm 0.02 \mu\text{Hz}$ and $\nu_{\text{max}} = 38.4 \pm 0.5 \mu\text{Hz}$).


INDIVIDUAL MODE FREQUENCIES



We extracted individual mode frequencies from the the power spectrum of each star using the FAMED pipeline [7]. A total of 34 (23) modes of angular degree $\ell = 0, 1, 2$, and 3 were extracted across 8 (7) radial orders for KOI-3886 (ι Dra).

The outcome of the *peak-bagging* process is shown above for the four central radial orders of KOI-3886. The green curve is a smoothing of the power density by an amount equivalent to the average radial-mode linewidth found by FAMED. The sloping dashed line represents the local background. Extracted individual mode frequencies are tagged according to their pressure radial order (n_p) and angular degree (ℓ), with color bands indicating their 3σ uncertainties.


DETAILED MODELING



A grid-based inference procedure was used to determine the fundamental stellar properties. It consisted in fitting the individual mode frequencies, along with a set of classical constraints, following the approach of [8], without considering interpolation.


The échelle diagram displays the observed frequencies and the model frequencies corresponding to a representative best-fitting model of KOI-3886. Circles, triangles, and squares indicate modes of angular degree $\ell = 0$ (radial), $\ell = 1$ (dipole), and $\ell = 2$ (quadrupole), respectively. We plotted a range of g-dominated $\ell = 1$ model frequencies per order (with symbol size scaled as $I^{-0.5}$, I being the mode inertia).

A PLANET IMPOSTOR



Asteroseismology, together with a set of follow-up observations and light curve analyses, helped reveal the planet candidate around KOI-3886 (A component) as a false positive and reinterpreting the system as an eclipsing brown dwarf in a hierarchical triple [9].

AN ELUSIVE COMPANION



The precise ($\sim 6\%$) seismic mass derived for ι Dra was used to enhance the precision of the known planet’s parameters [4]. Moreover, the combination of continued radial-velocity (RV) monitoring with *Hipparcos* and *Gaia* astrometry led to the detection of an additional long-period companion, its mass estimate lying on the border of the planet and brown dwarf regimes.

REFERENCES

- [1] Rowe, J. F., Coughlin, J. L., Antoci, V., et al. 2015, ApJS, 217, 16
- [2] Frink, S., Mitchell, D. S., Quirrenbach, A., et al. 2002, ApJ, 576, 478
- [3] García, R. A., Hekker, S., Stello, D., et al. 2011, MNRAS, 414, L6
- [4] Hill, M. L., Kane, S. R., Campante, T. L., et al. 2021, submitted to AJ
- [5] Corsaro, E., & De Ridder, J. 2014, A&A, 571, A71
- [6] Campante, T. L., Corsaro, E., Lund, M. N., et al. 2019, ApJ, 885, 31
- [7] Corsaro, E., McKeever, J. M., Kuszlewicz, J. S., et al. 2020, A&A, 640, A130
- [8] Li, T., Bedding, T. R., Christensen-Dalsgaard, J., et al. 2020, MNRAS, 495, 3431
- [9] Lillo-Box, J., Ribas, Á., Montesinos, B., et al. 2021, accepted for publication in A&A (arXiv:2106.05011)