UNIVERSITY^{OF} BIRMINGHAM University of Birmingham Research at Birmingham

Prevalence and determinants of oral health conditions and treatment needs among slum and non-slum urban residents

Osuh, Mary E.; Oke, Gbemisola A.; Lilford, Richard J.; Owoaje, Eme; Harris, Bronwyn; Taiwo, Olalekan John; Yeboah, Godwin; Abiona, Taiwo; Watson, Samuel I.; Hemming, Karla; Quinn, Laura; Chen, Yen-fu; Trevisan, Maurizio *DOI*:

10.1371/journal.pgph.0000297

License: Creative Commons: Attribution (CC BY)

Document Version Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Osuh, ME, Oke, GA, Lilford, RJ, Owoaje, E, Harris, B, Taiwo, OJ, Yeboah, G, Abiona, T, Watson, SI, Hemming, K, Quinn, L, Chen, Y & Trevisan, M (ed.) 2022, 'Prevalence and determinants of oral health conditions and treatment needs among slum and non-slum urban residents: evidence from Nigeria', *PLOS Global Public Health*, vol. 2, no. 4, e0000297. https://doi.org/10.1371/journal.pgph.0000297

Link to publication on Research at Birmingham portal

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



Citation: Osuh ME, Oke GA, Lilford RJ, Owoaje E, Harris B, Taiwo OJ, et al. (2022) Prevalence and determinants of oral health conditions and treatment needs among slum and non-slum urban residents: Evidence from Nigeria. PLOS Glob Public Health 2(4): e0000297. https://doi.org/10.1371/ journal.pgph.0000297

Editor: Maurizio Trevisan, VinUniversity, VIET NAM

Received: November 8, 2021

Accepted: March 6, 2022

Published: April 22, 2022

Copyright: © 2022 Osuh et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: The data underlying the results presented in the study are available from the University of Ibadan Urban Health Repository (https://trd.ui.edu.ng/health).

Funding: National Institute for Health Research (NIHR) Global Health Research Unit for Improving Health in Slums, grant number (16/136/87) using UK aid from the UK Government to support global health research: MEO, EO, BH, OJT, GY, SW, RJL, YFC. NIHR Applied Research Collaboration (ARC) West Midlands: RJL, LQ. NIHR doctoral research **RESEARCH ARTICLE**

Prevalence and determinants of oral health conditions and treatment needs among slum and non-slum urban residents: Evidence from Nigeria

Mary E. Osuh^{1,2}*, Gbemisola A. Oke², Richard J. Lilford³, Eme Owoaje⁴, Bronwyn Harris¹, Olalekan John Taiwo⁵, Godwin Yeboah⁶, Taiwo Abiona⁴, Samuel I. Watson³, Karla Hemming³, Laura Quinn³, Yen-Fu Chen¹

Division of Health Sciences, Warwick Medical School, University of Warwick, Coventry, United Kingdom,
Faculty of Dentistry, Department of Periodontology and Community Dentistry, College of Medicine,
University of Ibadan, Ibadan, Oyo State, Nigeria, 3 Faculty of Public Health, Department of Community
Medicine, College of Medicine, University of Ibadan, Ibadan, Oyo State, Nigeria, 4 Institute of Applied Health
Research, College of Medical and Dental Sciences, The University of Birmingham, Birmingham, United
Kingdom, 5 Faculty of Social Sciences, Department of Geography, University of Ibadan, Ibadan, Oyo State,
Nigeria, 6 Warwick Information and Digital Group, University of Warwick, Coventry, United Kingdom

* Mary.Osuh@warwick.ac.uk, meosuh2@gmail.com

Abstract

Oral diseases constitute a neglected epidemic in Low and Middle-Income Countries (LMICs). An understanding of its distribution and severity in different settings can aid the planning of preventive and therapeutic services. This study assessed the oral health conditions, risk factors, and treatment needs among adult residents in the slum and compared findings with non-slum urban residents in Ibadan, Nigeria. The Multistage sampling was used to select adult (>18-years) residents from a slum and a non-slum urban sites. Information sought from participants included dietary habits, oral hygiene practices, and the use of dental services. Oral examinations were performed in line with WHO guidelines. Associations were examined using logistic regression. Mediation analysis was undertaken using generalized structural equation modeling. The sample comprised 678 slum and 679 nonslum residents. Median age in slum vs non-slum was 45 (IQR:32-50) versus 38 (IQR:29-50) years. Male: female ratio was 1:2 in both sites. Prevalence of oral diseases (slum vs non-slum sites): dental caries (27% vs 23%), gingival bleeding (75% vs 53%) and periodontal pocket (23% vs 16%). The odds of having dental caries were 21% higher for the slum dwellers compared to non-slum residents (OR = 1.21, 95% CI:0.94 to 1.56); and 50% higher for periodontal pocket (OR = 1.50, 95%CI: 1.13 to 1.98), after adjusting for age and sex. There was little evidence that tooth cleaning frequency mediated the relationship between place of residence and caries (OR = 0.95, 95%CI: 0.87 to 1.03 [indirect effect], 38% mediated) or periodontal pocket (OR = 0.95, 95%CI: 0.86 to 1.04, 15% mediated). Thirty-five percent and 27% of residents in the slum and non-slum sites respectively required the "prompt and urgent" levels of treatment need. Oral diseases prevalence in both settings are high and the prevalence was generally higher in the slum with correspondingly higher levels of prompt fellowship: LQ. NIHR Evidence Synthesis Programme, grant number (14/25/05): YFC The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

and urgent treatment needs. Participants may benefit from targeted therapeutic and health promotion intervention services.

Introduction

Slum settlements provide homes for about 1 billion world population [1,2]. Slums are characterized by crowded, unhealthy places with a high risk of infection and injury [3,4]. Slum residents are often marginalized and have limited access to basic services [1,5]. Empirical evidence points to higher disease burdens among slum dwellers compared to their rural and urban counterparts [6–9]. This has led many to suggest that slums should be identified and studied separately from other types of urban settings lest the plight of slum dwellers gets lost in urban averages [1,7,10]. The health of slum dwellers has been a major concern to government and researchers as slum dwellers appeared to have comparatively greater health challenges compared to non-slum dwellers within the same urban continuum [6–8]. That said, the determinants of disease between slum and non-slum settings might be less clear-cut when it comes to non-communicable diseases. Whereas there is little doubt that children fare particularly badly in slums as compared to non-slum city areas, the situation for adults is more nuanced. Evidence points to a higher prevalence of smoking [11,12], obesity, and diabetes mellitus [13] among slum dwellers than residents of other parts of the city, although the prevalence of high blood pressure appeared similar in both settings [13].

Oral health is growing in significance as an integral part of health and exerts significant effects on individuals, communities, and the wider society [14-18]. For long, oral health is recognized to be an essential component of overall health and well-being [18-20]. Currently, oral diseases affect an estimated 3.5 billion people globally and untreated dental caries is one of the most prevalent non-communicable diseases [14,21]. The economic burden of oral diseases is considerable [22-24]. From the 2010 Global Burden of Disease Study, the global economic burden of dental diseases amounted to \$442 billion yearly, of which \$298 billion (4.6% of global health expenditure) was attributed to direct treatment costs and \$144 billion to indirect costs in terms of productivity losses due to caries, periodontitis, and tooth loss [22]. We recently carried out a systematic review of population-based surveys of oral health among adult urban residents in LMICs and found only three eligible studies dealing with oral health in slums: one study in Ahmedabad, India compared oral health across slum versus non-slum urban locations [25] and the remaining two from Bangladesh and Indore city, central India examined slum populations exclusively [9,26]. None was found for sub-Saharan African countries. This is in spite of the fact that oral health issues are regarded as a substantial population health burden, especially for the disadvantaged and poor population groups in both the developing and developed countries [27,28]. We, therefore, undertook this study to compare oral health and the determinants of oral health across slum and non-slum city areas. It is worth noting that major risk factors for caries and periodontal disease which include unhealthy sugary foods, tobacco use, excessive alcohol consumption, and poor hygiene practices, are also risk factors for other non-communicable diseases such as cardiovascular diseases, cancer, chronic respiratory disease, and diabetes [14,15,29-31].

In this paper we compared oral health in a slum vs. a non-slum area with the following aims: 1) To find out whether and to what extent differences in oral conditions may occur across slum vs. non-slum areas. 2) To find out if any difference between slum and non-slum areas could simply be attributed to age and gender differences. 3)To examine known

determinants (risk factors) of oral diseases by location. 4) To examine the effect of each risk factor on oral conditions net of age, gender, and location. 5) To examine the causal component of risk factors associated with the location identified through the above analyses.

We hypothesized that the determinants of oral health might pull in opposite directions with slum dwellers taking less cariogenic diet than non-slum city inhabitants while the non-slum residents exhibited better dental hygiene and had better access to dental health care [25,32–36].

Methods

Design

The study was a descriptive cross-sectional survey with an analytical comparison between slum and non-slum sites in Ibadan, the capital city of Oyo State, South-Western Nigeria. A purposive sampling method was deployed in the selection of the slum (Idikan) and the nonslum urban (Okeado) sites. The sites were selected based on similar ethnic and religious profiles.

Ethics

Ethical approvals for the research protocol were obtained from the Biomedical and Scientific Research Ethics Committee (BSREC: 37/18-19) of the University of Warwick and the Oyo State Research Ethics Review Committee (AD 13/479/1247) of Ibadan, Nigeria (<u>S1</u> and <u>S2</u> Files).

Sample size calculation

Sample size calculations were made to ensure sufficient statistical power for assessing the prevalence of oral diseases and for comparing the proportions of oral disease between slum and non-slum sites [37,38]. The estimates of prevalence for oral disease outcomes used were 12.8% for dental caries and 75.4% for periodontal disease [39,40]. For estimating oral disease prevalence, the level of confidence was set at 95% and the desired level of precision was 5 percentage points. For comparison between slum and non-slum sites, a difference of 10% was assumed with an alpha of 0.05 and 80% power. Based on the largest sample size suggested by the calculations, a minimum sample size of 650 people in each slum and non-slum setting and a total of 1,300 was required.

Mapping of study sites for easy sampling

The boundaries of each site were determined together with the National Population Commission (NPopC) staff (a commission vested with the duty of publishing information or data on population for various purposes in Nigeria) who were familiar with the communities. The site boundaries were identified on Google Map from where structures were delineated and digitized using high-resolution satellite imagery from the DigitalGlobe. A building-footprint database was created for each site using this high-resolution satellite imagery. A total of 1,923 and 3711 structures were extracted from the slum and the non-slum areas, respectively.

Sampling strategy

We used a multistage sampling technique. First, we selected 1000 building structures each for slum and non-slum sites from the structures identified from satellite imagery described above using the random selection algorithm in Quantum GIS software. Second, we located the buildings using the Geographic Positioning System (GPS) technology aided by the local internet

network. Teams of researchers created an inventory of all households within each selected building, then selected one household per building at random using the OpenDataKit (ODK) client application. Lastly, we obtained an inventory of all adults in each selected household from which we selected one eligible adult participant at random, again using ODK software.

Inclusion and exclusion criteria

To be eligible, participants had to be at least 18 years of age and must be residents in the area for at least one year. Visitors, guests, house staff, and chronically ill patients, were excluded from the study.

Data collection exercise

Following the random selection of an eligible household member to be included in the study, a data collection team visited the selected household to meet the selected person. If the selected adult was available, he/she was enrolled in the survey and when not available, a second and up to a third visit (if needed) was made. During the meeting, the number of visits before a successful meeting was noted, then detailed information about the project was shared, and clarifications were provided, where needed. The participant was asked about their willingness to participate in the research and if yes, they were invited to sign the consent form. After obtaining the participant's signature, a good location with adequate light and sitting arrangement was identified in the home environment with the assistance of the participant. The participant was then seated on a regular chair and the questionnaire was administered by the trained recording clerk while the dentist prepared for the oral examination (light source—headlamp, fresh tray of sterile and sealed set of examination instruments, wooden spatula, sterile gloves, and face mask). After the administration of the questionnaire, oral examination was performed by the dentist, and findings were entered into the prepared tablets by the recording clerk. Following the oral examination, if a participant was found to have dental problems that required urgent attention, they were referred to the University College Hospital with a unique identifier that entitled them to subsidized cost of care. Others who did not require urgent dental treatment benefitted from oral hygiene practice reinforcement and education.

On the few occasions (8 times) that the participants declined to partake in the research exercise, the reasons for declining were noted and the process was discontinued. Thereafter, another eligible member within the same household was randomly selected as a replacement for the person that declined.

At the end of the data collection exercise, each participant received a tube of toothpaste and a toothbrush as a token of appreciation for their time in participation in the study.

Instrument validation

The questionnaire for this study was an adaptation of a questionnaire from the National Institute for Health Research—Global Health Research Unit (NIHR-GHRU) [41] and the WHO adult oral health questionnaire [42]. The face validity of the adapted questionnaire was checked by two experts in the field of dental public health.

Training of data collection teams

Ten teams, each consisting of a qualified dentist and an experienced recording clerk, were trained to collect data. The activities of each team were supervised and monitored by the first author. All team members were trained in understanding and interpretation of the question-naire as well as the application of criteria and codes for various oral diseases and appropriate

recording in line with the WHO standards [42]. The dentists were fresh graduates of the University of Ibadan and were specially trained at the University College Hospital (UCH) dental clinic, to elicit and record findings on oral examination in line with the WHO set criteria [42]. The WHO's Community Periodontal Index (CPI) probe (a specially designed, lightweight metallic probe) for the assessment of periodontal status and disposable materials [42] were used in the training of the research team members. Examiners' interrater variability was tested for dental caries, periodontal pocket formation and other oral health conditions, in which 10 examiners interpreted 10 different scenarios for both oral health outcomes.

Pre-test

A pre-test of the study was conducted at the Abadina community of the University of Ibadan using 50 randomly selected buildings and households. The exercise enabled verification of the veracity of information obtained from satellite imagery by a physical on-the-spot visit of the 50 buildings (5 buildings per team). The exercise also enabled the collection of other useful information on the QField, an Android-based GIS for collecting field data. The processes of sampling study participants using the ODK software were tested. The duration of data collection activities (average time and range; including both questionnaire and oral examination), as well as the recording of information into the tablet computer, were verified. Ambiguity in the questionnaire design was also identified and promptly reviewed.

Questionnaire

Participants responded to a structured and standardized questionnaire which was intervieweradministered. Information collected included participants' socio-demographic information, period of residence, the experience of a dental problem within the last 12-month period, selfassessment of oral health status, dental hygiene practices (frequency of tooth cleaning, use of aids for oral hygiene, use of toothpaste containing fluoride), utilization of dental services (dental visits, reasons for dental visits), consumption of sugary foods and drinks, use of tobacco, and consumption of alcohol. The questionnaire was translated into the local (Yoruba) language and back-translated in line with WHO recommendations.

Oral examination

Oral examinations were performed by the dentist in each team. The WHO's Community Periodontal Index (CPI) probe and disposable materials [42] were deployed for the oral examination. Mouth examination was carried out with the aid of visual and tactile senses. The teeth were examined systematically from the first to the fourth quadrant. Diagnosis of oral conditions was made in accordance with the WHO criteria for oral health surveys [42] and the findings were recorded using the WHO standardized oral health assessment format which was prepared into the tablet computers. The following information was obtained from all participants: dentition status (carious teeth, missing due to caries and filled teeth for the reason of caries), periodontal status (gingival bleeding and pocket formation); loss of attachment, and intervention urgency [42]. The lifetime caries experience—DMFT (sum of decayed, missing, and filled teeth) [42], was derived for each participant, and the presence and stages of periodontal diseases (gingival bleeding, periodontal pocket, or periodontal attachment loss) were recorded [43]. Sterilization of all clinical dental instruments was done at the end of each day's exercise at the Primary Oral Health Care Centre, University College Hospital (UCH), Ibadan, based in Idikan.

Data analysis

Demographic characteristics and oral health outcomes were described by place of residence (slum and non-slum). For categorical variables, counts and percentages were reported. For numeric variables, means and standard deviations or medians and interquartile ranges were reported where appropriate. The frequency of cariogenic food consumption was determined by response to 8 items whose scores were added to construct a continuous variable reporting a sum score for each individual (range 8–48). The mean score was derived and used as a cut-off to produce binary output of less and more frequent intake [44,45]. Alcohol intake within the preceding 30 days was also determined using the mean number of drinks by those who drank as the cut-off to produce a binary output into; moderate and excessive alcohol intake [46].

Our objective was to explore the possible reasons for any differences in oral health outcomes between people living in the slum vs. those in the non-slum area. To this end, we first examined the relationship between place of residence and each of the two main oral health outcomes (dental caries, and periodontal pocket formation).

We chose dental caries and periodontal pocket formation because of their well-known competing risk for tooth loss in the adult population [47–50].

Second, we determined whether particular risk factors such as alcohol use, tobacco use, diet, and oral hygiene habits had a relationship with the place of residence and oral health outcomes. This is because these risk factors independently or collectively play significant roles in the development of oral diseases [51–56]. To establish causal pathways, we predetermined whether certain risk factors (diet and oral hygiene habits) acted as mediators. For example, we explored whether a cariogenic diet mediates the relationship between place of residence and dental caries. These risk factors were chosen as potential mediators for analysis on the basis of our dental knowledge as well as their mediating roles in other situations for example tooth brushing as a mediator between cognitive impairment and oral health outcome [57] and nutrition as a mediator between oral and systemic disease [58].

Associations between oral health outcomes, place of residence, and risk factors were explored using logistic regression models, unadjusted and adjusted for age and sex. Risk factors considered were alcohol intake, tobacco use, cariogenic diet, and frequency of tooth cleaning. Odds ratios, 95% confidence intervals, and p-values were reported. Mediation analysis was performed using generalized structural equation modeling (*gsem* command in Stata). The total effect, direct effect, and indirect effect were calculated using the nonlinear combination command (*nlcom*). The proportion of the relationship mediated was calculated by taking the proportion of the indirect effect divided by the total effect. The effect values from the non-linear combination command were then transformed to odds ratios by taking the exponential. All effects were reported with 95% confidence intervals and p-values. Percentage agreement and Gwet's AC1 index were used to measure interrater variability of researchers for diagnosing the oral health conditions. The statistical analysis was carried out using IBM SPSS version 26 and Stata version 16.1.

Results

Sample characteristics

A total of 1,357 participants were included, evenly distributed between the slum and non-slum residence. Overall, there were 245 (36%) males in slum residences and 234 (35%) males in non-slum residences (Table 1). The average age was higher in the slum residences; the median age was 45 years (interquartile range [IQR]: 32 to 50 years) compared to a median of 38 (IQR: 29–50) in non-slum residences. Oral health perceptions and hygiene practices, and utilization

Characteristic	Slum (n = 678)	Non-slum (n = 679)
Male	245 (36.1)	234 (34.5)
Age-median (IQR)	45 (32 to 60)	38 (29 to 50)
Age		
<35 years	192 (28.3)	260 (38.3)
Adult age group (35–44 years)	127 (18.7)	186 (27.4)
45-54 years	114 (16.8)	109 (16.1)
55–64 years	102 (15.0)	61 (9.0)
Elderly age group (65–74 years)	88 (13.0)	40 (5.9)
75 years and above	55 (8.1)	23 (3.4)
Current marital status		
Married	437 (64.5)	458 (67.5)
Living together but not married	9 (1.3)	13 (1.9)
Divorced or separated	34 (5.0)	7 (1.0)
Widowed	110 (16.2)	50 (7.4)
Never married and never lived together	88 (13.0)	151 (22.2)
Highest educational level		
No education	175 (25.8)	42 (6.2)
Primary school	179 (26.4)	61 (9.0)
Secondary school	285 (42.0)	301 (44.3)
Post-secondary school	32 (4.7)	183 (27.0)
University education	7 (1.0)	92 (13.5)
Currently working	533 (78.6)	552 (81.3)
Length of time of residence in the neighbourhood		
1–10 years	102 (15.0)	203 (29.9)
11–20 years	103 (15.2)	80 (11.8)
>20years	473 (69.8)	396 (58.3)
Wealth quintile/status (SES)		
1st quintile (lowest)	231 (34.1)	40 (5.9)
2nd quintile (lower)	167 (24.6)	105 (15.5)
3rd quintile (middle)	157 (23.2)	114 (16.8)
4th quintile (higher)	97 (14.3)	175 (25.8)
5th quintile (highest)	26 (3.8)	245 (36.1)
Main drinking water source		
Piped/ tap water	31 (4.6)	47 (6.9)
Borehole	74 (10.9)	283 (41.7)
Well/ spring/ tanker or cart supply	5 (0.7)	8 (1.2)
Rainwater	64 (9.4)	10 (1.5)
Sachet/ bottled water	504 (74.3)	331 (48.7)

Table 1. Baseline characteristics by place of residence.

IQR: Inter quartile range; SES: Socio-economic status.

Number and percentages reported unless stated otherwise.

https://doi.org/10.1371/journal.pgph.0000297.t001

of dental services are summarized in <u>S1 Table</u>. A lower percentage of those in slum residences had seen a dentist (17% versus 24%), used a toothbrush for tooth cleaning (82% versus 98%), and brushed at least twice daily (24% versus 27%) compared to non-slum residents.

Oral health conditions

Prevalence of the common oral health conditions among the overall study sample was: dental caries—25%; gingival bleeding—64%; pocket formation—19%; and dental trauma—26%. Other findings include: missing teeth—14%; filled teeth– 2%; attachment loss—10% (S2 Table). Regarding the levels of treatment needed or intervention urgency: very few participants (4%) had no need for dental treatment. Sixty-four percent of the overall study sample required the "preventive or routine" level of dental treatment, while 31% had the "prompt" and "immediate/ urgent" levels of treatment need (S2 Table). Examiners' assessment for oral health conditions revealed substantial, moderate, and almost perfect levels of interrater agreements for the three common oral health conditions respectively: Dental caries—77% agreement (Gwet's index: 0.71), periodontal pocket formation—66% agreement (Gwet's index: 0.49), dental trauma—96% agreement (Gwet's index: 0.94); gingival attachment loss- 67% agreement (Gwet's index: 0.50; enamel fluorosis- 80% agreement (Gwet's index: 0.74); dental erosion- 86% agreement (Gwet's index: 0.84).

Associations between the place of residence and oral health conditions. The number and prevalence of oral health conditions among the participants, according to their residential settings, are shown in S2 Table: Dental caries (27% versus 23%) and periodontal pocket formation (23% vs 16%) were more prevalent in slum residences than non-slum residences (Table 2 and Fig 1). Even after adjusting for age and sex, the odds of having dental caries were estimated to be 21% higher for people who lived in slums compared to non-slum residences (OR = 1.21, 95% CI: 0.94 to 1.56); and 50% higher (OR = 1.50, 95% CI: 1.13 to 1.98) for periodontal pocket. The number of participants with no need for dental treatment was 3% and 5% in slum and non-slum sites respectively. The type of treatment required in both sites was mostly preventive/ routine dental treatment and this comprised 62% versus 67% of residents in the slum and the non-slum setting respectively. Prompt and urgent levels of treatment were required for 35% (slum) versus 28% (non-slum) of participants (S2 Table).

Associations between the place of residence and risk factors for oral health conditions. Alcohol intake for participants was low in both slum and non-slum residence: moderate (10% versus 13%) and excessive (2% versus 4%). Tobacco use was slightly higher in slum compared to non-slum residences (13% versus 9%), whereas higher consumption of cariogenic diet was slightly lower in slum compared to non-slum residences (44% versus 50%). After adjusting for age and sex, the odds of having a higher consumption of cariogenic diet was 10% lower (OR = 0.90, 95% CI:0.72 to 1.13) in slum residences compared to non-slum residences. A slightly lower percentage of participants cleaned their teeth at least twice daily in slum residences compared to non-slum residences (24% versus 27%), After adjusting for age and sex, the odds of tooth cleaning at least twice a day were 19% lower (OR = 0.81, 95% CI: 0.63 to 1.04) in slums compared to non-slum residences (Table 2 and Fig 1).

Association between risk factors and oral health conditions. Associations between oral health conditions and potential risk factors, including cariogenic diet, alcohol use, tobacco use, and frequency of tooth cleaning were reported unadjusted and adjusted for age group and sex (S3 and S4 Tables). The odds of having dental caries were 28% higher (OR = 1.28, 95% CI:0.97 to 1.69) for people who cleaned their teeth at least twice daily compared to less than twice daily, adjusting for age and sex. The odds of having periodontal pocket formations was 103% higher (OR = 2.03, 95% CI:0.97 to 4.22) for people who had an excessive intake of alcohol compared to no alcohol intake and 28% higher (OR = 1.28, 95% CI:0.95 to 1.73) for people who cleaned their teeth at least twice daily compared to less.

Characteristic	Slum (n = 678)	Non-slum (n = 679)	Unadjusted odds ratio (95% CI) p-value	Adjusted odds ratio* (95% CI) p-value
Oral health conditions			-	
Dental caries	181 (26.7)	154 (22.7)	1.24 (0.97 to 1.59) p = 0.087	1.21 (0.94 to 1.56) p = 0.145
Periodontal pocket formation	153 (22.6)	105 (15.5)	1.59 (1.21 to 2.10) p = 0.001	1.50 (1.13 to 1.98) p = 0.005
Risk factors				
Alcohol intake**				
Didn't drink alcohol (last 30 days)	594 (87.6)	570 (84.0)	Reference	Reference
Moderate intake	69 (10.2)	85 (12.5)	0.78 (0.56 to 1.09) 0.147	0.75 (0.52 to 1.08) p = 0.127
Excessive intake	15 (2.2)	24 (3.5)	0.60 (0.31 to 1.15) 0.126	0.68 (0.34 to 1.34) p = 0.265
Ever used tobacco	89 (13.1)	63 (9.3)	1.48 (1.05 to 2.08) p = 0.025	1.59 (1.11 to 2.30) p = 0.012
More frequent cariogenic food consumption	295 (43.5)	338 (49.8)	0.78 (0.63 to 0.96) p = 0.021	0.90 (0.72 to 1.13) p = 0.365
Tooth cleaning (at least twice daily)	163 (24.0)	184 (27.1)	0.85 (0.67 to 1.09) p = 0.197	0.81 (0.63 to 1.04) p = 0.099

Table 2. Logistic regression models to explore associations between the place of residence and 1) oral health conditions; 2) risk factors, unadjusted and adjusted for age group and sex.

* Adjusted age group and sex.

** Multinomial logistic regression used and relative-risk ratios reported.

https://doi.org/10.1371/journal.pgph.0000297.t002

Validity check for logistic regression. Assumptions of logistic regression were checked for each of the models reported above. All models had binary outcomes; little or no multicollinearity was found between independent variables; the influence of outliers did not affect the results and the observations were independent.

Mediation analysis. *Mediation via tooth cleaning frequency.* The odds of having dental caries were estimated to be 22% higher for people who lived in slums compared to non-slum residents (OR = 1.22, 95% CI:0.95 to 1.58); and 51% higher (OR = 1.51, 95% CI:1.14 to 2.01) for periodontal pocket formation, after adjusting for tooth cleaning frequency, age and sex (direct effects, Fig 2 and Table 3). The odds of tooth cleaning at least twice a day were 19% lower (OR = 0.81, 95% CI:0.63 to 1.04) in slums compared to non-slum residences, adjusting for age and sex (indirect effect). The odds of having dental caries were 30% higher (OR = 1.30, 95% CI:0.98 to 1.71) and periodontal pocket formation was 31% higher (OR = 1.31, 95% CI:0.96 to 1.77) for cleaning teeth at least twice daily compared to less, adjusting for place of residence, age, and sex (indirect effect).

Mediation via cariogenic diet. The odds of having dental caries were estimated to be 21% higher for people who lived in slums compared to non-slum residents (OR = 1.21, 95% CI:0.94 to 1.56); and 49% higher (OR = 1.49, 95% CI:1.12 to 1.97) for periodontal pocket formation,



Fig 1. Distribution of key survey data (demographic characteristics, oral health conditions and risk factors) by slum versus non-slum site.

https://doi.org/10.1371/journal.pgph.0000297.g001



Fig 2. Casual pathways between place of residence and oral health conditions (dental caries and periodontal pocket formation) with frequency of tooth cleaning as a potential mediator, controlling for age group and sex.

https://doi.org/10.1371/journal.pgph.0000297.g002

after adjusting for cariogenic diet, age and sex (direct effects). The odds of having a more frequent cariogenic diet were 10% lower (OR = 0.90, 95% CI:0.72 to 1.13) for people in slum compared to non-slum residents. The odds of having dental caries was 2% higher (OR = 1.02, 95% CI:0.79 to 1.32) and periodontal pocket formation was 41% lower (OR = 0.59, 95% CI:0.44 to 0.78) for having a more frequent cariogenic diet compared to less, adjusting for place of residence, age and sex (indirect effect).

There was no evidence cariogenic diet mediated the relationship between place of residence and dental caries (OR = 1.00, 95% CI:0.97 to 1.02 [total indirect effect], estimated 1% mediated) or periodontal pocket formation (OR = 1.06, 95% CI:0.93 to 1.19 [total indirect effect], estimated 12% mediated; see Fig 3 and Table 4).

Table 3. Generalized structural equation modeling to explore potential mediator frequency of tooth cleaning on the relationship between place of residence and oral health outcomes.

Effect	Odds ratio (95% CI) p-value
Dental caries as outcome	<u> </u>
Direct effect	
Place of residence \rightarrow Dental caries	1.22 (0.95 to 1.58) p = 0.123
Indirect effects	
Place of residence \rightarrow Tooth cleaning frequency	0.81 (0.63 to 1.04) p = 0.096
Tooth cleaning frequency \rightarrow Dental caries	1.30 (0.98 to 1.71) p = 0.067
Total indirect effect	0.95 (0.87 to 1.03) p = 0.224
Total effect	
Place of residence \rightarrow Dental caries	1.16 (0.88 to 1.51) p = 0.288
Percentage mediated	38%
Periodontal pocket formation as outcome	
Direct effect	
Place of residence \rightarrow Periodontal pocket formation	1.51 (1.14 to 2.01) p = 0.004
Indirect effects	
Place of residence \rightarrow Tooth cleaning frequency	0.81 (0.63 to 1.04) p = 0.096
Tooth cleaning frequency \rightarrow Periodontal pocket formation	1.31 (0.96 to 1.77) p = 0.085
Total indirect effect	0.95 (0.86 to 1.04) p = 0.225
Total effect	
Place of residence \rightarrow Periodontal pocket formation	1.43 (1.06 to 1.92) p = 0.017
Percentage mediated	15%

Analysis adjusted for age group and sex.

https://doi.org/10.1371/journal.pgph.0000297.t003

Discussion

Our study has established that the prevalence of common dental diseases in the two communities studied ranged from moderate to very high for dental caries and periodontal disease respectively. Complete absence of oral diseases was detected in only about 4% of participants who were adjudged to require no form of dental treatment. The high prevalence of some oral diseases such as caries and periodontal disease is comparable to reports from available literature worldwide [59–61]. The fact that these diseases are chronic, silent, and slowly progressive





Fig 3. Causal pathways between place of residence and oral health outcomes (dental caries and periodontal pocket formation) with cariogenic diet as a potential mediator, controlling for age group and sex.

https://doi.org/10.1371/journal.pgph.0000297.g003

in nature [14-18], often renders them prone to neglect [62,63]. Given the economic effects of dental disease and their effect on general health and well-being [22-24], more attention should be paid to early detection and treatment of dental diseases [23].

Magnifying the already heavy oral disease burden is that the slum residents recorded higher prevalence rates relative to their non-slum counterparts, although the difference was modest. The observed pattern of distribution would seem to mimic the general distribution pattern of non-communicable diseases and risk factors associated with ill-health such as diabetes mellitus, smoking and obesity, which were reported as preponderant in disadvantaged communities in previous studies [6–8].

Effect	Odds ratio (95% CI) p-value
Dental caries as outcome	
Direct effect	
Place of residence \rightarrow Dental caries	1.21 (0.94 to 1.56) p = 0.144
Indirect effects	
Place of residence \rightarrow Cariogenic diet	0.90 (0.72 to 1.13) p = 0.364
Cariogenic diet \rightarrow Dental caries	1.02 (0.79 to 1.32) p = 0.853
Total indirect effect	1.00 (0.97 to 1.02) p = 0.855
Total effect	
Place of residence \rightarrow Dental caries	$\begin{array}{c} 1.21 \\ (0.93 \text{ to } 1.56) \\ p = 0.149 \end{array}$
Percentage mediated	1%
Periodontal pocket formation as outcome	
Direct effect	
Place of residence \rightarrow Periodontal pocket formation	1.49 (1.12 to 1.98) p = 0.006
Indirect effects	
Place of residence \rightarrow Cariogenic diet	0.90 (0.72 to 1.13) p = 0.364
Cariogenic diet \rightarrow Periodontal pocket formation	0.59 (0.44 to 0.78) p<0.001
Total indirect effect	1.06 (0.93 to 1.19) p = 0.380
Total effect	
Place of residence \rightarrow Periodontal pocket formation	1.57 (1.15 to 2.14) p = 0.004
Percentage mediated	12%

Table 4. Generalized structural equation modeling to explore potential mediator cariogenic diet on the relationship between place of residence and dental outcomes.

Analysis adjusted for age group and sex.

https://doi.org/10.1371/journal.pgph.0000297.t004

According to the WHO, dental caries is the most common non-communicable diseases worldwide and is a major public health problem, globally [64]. The global overall prevalence of dental caries for all ages combined as reported in the global burden of disease study 2010 is approximately 35% [65]. In comparison, caries prevalence from our study population is 25%. This lower prevalence may be explained in part by the age of our study participants which is predominantly younger, since the disease is known to increase with age [42]. Other likely determinants of the lower prevalence rates recorded in our study include the habit of

consuming less of free sugar [66], fluoride use [67], as well as indigenous caries prevention measures [68]. Further studies to investigate the role and importance of each factor in disease aetiology for our study population should enhance efforts at controlling the same. The difference in the prevalence of caries observed between the slum and the non-slum locations was not significant, even after adjusting for age and gender. However, when the estimated prevalence of dental caries from our study of 27% and 23% in the slum and non-slum urban settings respectively, were compared with that reported from a similar study in India [25], which reported 61% and 47% in the respective settings, the difference was marked. Whereas in our study, participants were selected at random information on how India's study participants were selected and this methodological factor may explain the marked difference in findings between our study and the Indian study.

Access to clean drinking water, especially with respect to its fluoride content, plays a significant role in caries prevention. The main source of fluoride is water, while other sources include food, industrial exposure, drugs, and cosmetics [69]. The WHO, specified a fluoride concentration level of 0.5 mg/L (equivalent to 0.5 parts per million [ppm]) in drinking water for its beneficial effects in the prevention of caries and a tolerance limit of 1.5 mg/L [28,70]. In a tropical environment like Nigeria, a fluoride level of 0.3-0.6 ppm in drinking water was recommended depending on fluoride ingestion from other sources [71]. Although our study did not assess the fluoride concentration in participants' drinking water, available studies revealed that the majority of drinking water sources (58-75%) in all but one LGA in Nigeria contained 0.3 mg/L or less of fluoride [71]. A recent study conducted in south-west Nigeria reported a smaller proportion (48%) of drinking water with low fluoride from similar sources: ground water- well or borehole [71,72]. Access to clean drinking water is generally a challenge for a large proportion of the Nigerian population but the situation is worse in the slums [73,74]. The non-recognition of slums in official discourses often limit their consideration in the planning of essential public services such as water [74]. The aforementioned may explain why the commonest source of drinking water in both the slum and non-slum areas in our study is packaged water (sachet/ bottle water) followed by borehole. Available reports have confirmed these water sources as clean drinking water sources with adequate fluoride concentration [71,75]. Furthermore, fluoride toothpaste is recommended for the prevention of caries, and is usually advised in conjunction with good oral hygiene [76]. The majority of our study participants reported use of fluoridated toothpaste in cleaning their mouth. Most of the commercially available toothpastes in Nigeria are reported to have the standard recommended concentration of 1000-1500 ppm [72,77,78]. This range is considered both beneficial against caries and safe against dental fluorosis for any age if toothpaste was the individual's sole source of fluoride. The combination of access to clean/ safe drinking water source and the use of fluoridated toothpaste by the majority of our study participants probably make up for the low fluoride content available in the regions' ground water. Therefore, there may be justification to shift focus to other causes of caries such as dietary [79], cleaning habits and other risk factors [80].

More than two-thirds of the entire population in our study had periodontal disease. This comprised gingival bleeding of 64%; periodontal pocket formation of 19% and an attachment loss of 10%. Previous studies in different settings, inclusive of slums have utilized various criteria in the assessment of periodontal disease prevalence. These include the Community Periodontal Index of Treatment Needs (CPITN), Clinical Attachment Level (CAL), and/or probing depth (PD). The reported prevalence rates varied but were equally high [25,26,81,82]. Our findings that periodontal disease affected close to three-quarters of the slum dwellers and about half of the non-slum residents further buttresses the heavy burden of periodontal disease being borne by populations in LMICs particularly those residing in slums. However, our findings were again dissimilar to those reported in the study in India, in which the non-slum dwellers had a higher

prevalence of gingivitis than slum dwellers while having an equal prevalence for gingival bleeding [25]. This may be explained by their purposively selected small (300) population and that their participants were drawn from across the ages (children and adults). The difference in the prevalence of periodontal pocket formation observed between the slum and the non-slum locations in our study were statistically significant, and remained so, even after adjusting for age and sex. Overall, literature on population oral health surveys among adult residents of LMICs are sparse and more data/studies will be needed to make an adequate comparison.

In terms of the oral healthcare needs, participants in both settings had high needs for oral healthcare, and most of the needs related to preventive/routine treatments. The high treatment need observed in this study is similar to other studies which reported high prevalence of oral diseases [9,26]. However, when the level of treatment required was compared between the two residential locations, more of the slum dwellers required urgent level of care, while more of the non-slum residents required the preventive/routine dental treatments, relative to their counterparts respectively.

In tackling the oral health needs of the populace, access to basic publicly funded oral health services should be central [83,84]. In Nigeria as well as other LMICs, oral health is largely considered less important than the general health, and receives little attention in terms of health care planning and service delivery [85]. Oral health services are delivered in all health care levels (primary, secondary and tertiary) in Nigeria. The primary level of oral care is concerned with the prevention of dental diseases and management of dental emergencies, however, the services are available at only a few primary health care centres (PHCCs) nationwide [79,83,85]. General treatment of oral disorders is provided at the secondary care facilities: private dental clinics, general hospitals, institutions run by faith-based organisations, federal medical centres and armed forces hospitals. Specialised oral health services including treatment of oral diseases and patient rehabilitation are provided at the tertiary (teaching hospitals) [83,85]. Traditional oral health care providers are also available in most regions as well as few non-governmental organisations, all providing oral health services to the people [17,85,86]. The primary level of oral care, is considered as the veritable means to ensure easy access to oral health service to people in all locations, due to its widespread location in the country [83-85], but it has not succeeded in delivering on its service mandate to the people due to persistent challenges (manpower and other resources). Therefore, most oral health services are directed towards the provision of curative and rehabilitative care from the secondary and tertiary levels of care [85]. These levels of care are available mainly in the big cities and largely through out of pocket payment [17]. To date the inclusion of oral health into existing PHCs has continued to suffer setbacks in terms of facilities and capacity building [79,85]. Consequently, a number of independent initiatives targeted at charting a direction for Primary Oral Health Care were set up in Nigeria's dental schools, but these are too few to meet the current oral health need, they lack adequate manpower and are largely funded from out of pocket [79,85]. The Nigeria's National Health Insurance Scheme (NHIS) has a mandate, through various prepayment systems, to design and implement a social health insurance scheme that can facilitate easier access to affordable and available quality health care services and achieve a universal health coverage (UHC) [83]. But this equally has its own challenges which range from poor financing of the health system by governments, an absence of prepayment schemes, and a growing population of poorly paid people who reside majorly in slums [83,85]. As such, the NHIS billing system concerning oral health is largely considered only at the level of secondary care [85,87]. These challenges in accessing oral health care may also explain the low use of professional dental services observed among our study participants. A revision of the position of oral health services on the NHIS should advocate to include the coverage of preventive oral health care services as recommended by the WHO [85,88].

Tooth brushing, at least twice daily is the professionally recommended routine that should contribute to individual oral hygiene [89,90]. We found that only a small proportion of our study participants engaged in twice daily mouth cleaning routine across both settings. The indigenous norms, cultural practices, and habits native to the people as well as their limited exposure to correct information from school and through the media may have played a role [68,91]. The low prevalence of mouth cleaning was in stark contrast to findings from a national oral health survey conducted in Malawi which reported a much higher prevalence of 75% for brushing teeth at least twice daily [92]. About 40% in their study reported brushing their teeth three or more times a day. This however, did not seem to reduce the caries prevalence (49%) reported among their adult populations.

In advance of our data analysis, we hypothesized that different determinants would be at play in slum vs. non-slum areas. Previous work had suggested that diets in slums may be less cariogenic than in non-slum urban areas [66,93-95], while at the same time we suspected that hygiene practice and access to dental care may be poorer in slum areas. In the mediation analysis of the association between residential places (slum vs non-slum) and oral health conditions (dental caries and periodontal pocket formation), we found little evidence of either the frequency of tooth cleaning or cariogenic diet being potential mediators. This was unexpected given the established influence of these factors on oral health. The lack of a mediation effect may partly reflect the relatively small observed differences in tooth cleaning practice, cariogenic diet consumption, and dental caries between slum and non-slum populations. Previous studies have linked the pattern of sugar consumption [96-98] and tooth brushing [99,100] with periodontal disease. The unexpected finding in our study may be due to the effect of unknown confounders which require further investigation. They may also indicate issues related to self-reported measures of cariogenic diet consumption and tooth cleaning practice, and the grouping of exposure for these variables. For example, how the respondents' selfreported brushing frequencies approximate their actual tooth brushing behavior is not clear. In addition, the act of tooth brushing itself is complex as it combines many other variables such as the duration of brushing, the design and quality of the brush, the brushing method and the toothpaste used [99–102]. We have used brushing at least twice daily as the threshold for defining higher versus lower frequency of tooth cleaning, but a previous systematic review suggested the effect might be similar even with brushing at least once daily [99]. Maintenance of good oral hygiene is the foundation on which most oral diseases can be prevented [103]. However, achieving good oral hygiene involves more than brushing the teeth twice daily. A combination of quality in the act of tooth brushing itself [99-102], with the right cleaning frequency, is considered essential and plays a significant role in achieving the level of hygiene that can be effective against oral diseases [89,91]. Moreover, the role of professional tooth cleaning at regular intervals in inhibiting oral diseases should not be overlooked [54,104]. Data from our study suggested that the majority of both the slum and the non-slum residents had never visited a dentist similar to other studies [68,105]. Lastly, our study is cross-sectional and one may indeed require longitudinal studies to properly assess a causal effect. On the whole, our hypotheses regarding key determinants of oral health conditions in slum versus non-slum urban settings were not supported by the data that we observed.

Study limitations

Our study has certain limitations. First, the study was carried out in only one slum and one non-slum area, and so the findings may not be generalizable to all slums. Second, the cross-sectional design precludes causal inferences, hence only associations can be drawn. However, our study has some major strengths: To the best of our knowledge, it is the first community

oral health study in the LMICs to deploy the use of GIS software with GPS technology to select representative samples from randomly selected households and buildings in dense urban areas such as the slums. In the slums, it is rarely feasible to conduct a comprehensive door-to-door survey of every individual or household unit, that can ensure representativeness [41,106]. The adoption of this approach in our study is therefore an innovation among oral health studies.

Additionally, we utilized the latest in the manual series of WHO oral health survey methods [42] in the assessment of public oral health status. The methods were recommended for all oral health surveys to enhance easy comparison of results among populations globally. Lastly, our study is the second and currently, the largest of its kind in the LMICs to compare oral health survey findings related to the prevalence of oral diseases and their determinants among adults residing in the slum and non-slum settings, making a significant contribution to the currently sparse evidence base.

Conclusion

Oral disease prevalence is high in both the slum and the non-slum urban locations, and the prevalence is generally higher in the slum relative to the non-slum setting. We did not find clear evidence of the mediation effect of cariogenic diet consumption and tooth brushing frequency on the relationship between place of residence and oral health conditions. More than a third and more than a quarter of the slum and the non-slum residents respectively required the prompt and urgent levels of dental treatments. The participants may benefit from interventions targeted at therapeutic, preventive, and oral health promotion services.

Supporting information

S1 Table. Oral health perceptions, hygiene practices, and utilization of dental services by place of residence.

(DOCX)

S2 Table. Oral health conditions following dental examination. (DOCX)

S3 Table. Logistic regression models exploring associations between dental caries and risk factors.

(DOCX)

S4 Table. Logistic regression models exploring associations between periodontal pocket formation and risk factors. (DOCX)

S1 File. Biomedical and Scientific Research Ethics Committee full approval letter. (PDF)

S2 File. Oyo state Research Ethics Review Committee full approval letter. (PDF)

Acknowledgments

Our sincere appreciations go to the other members of the Improving Health in Slums Collaborative who are not on the authors' list for their assistances in no small measure to ensure the success of this project. All the study participants- residents of the slum and the non-slum communities are deeply appreciated for their huge support and cooperation that saw a seamless data collection process. The staff of the National Population Commission (NPoPC) in Nigeria are also well appreciated for the support and assistance rendered during the mapping exercises for the boundaries of the project study sites.

Author Contributions

- **Conceptualization:** Mary E. Osuh, Gbemisola A. Oke, Richard J. Lilford, Eme Owoaje, Bronwyn Harris, Olalekan John Taiwo.
- **Data curation:** Mary E. Osuh, Bronwyn Harris, Samuel I. Watson, Laura Quinn, Yen-Fu Chen.

Formal analysis: Mary E. Osuh, Gbemisola A. Oke, Richard J. Lilford, Eme Owoaje, Bronwyn Harris, Taiwo Abiona, Samuel I. Watson, Karla Hemming, Laura Quinn, Yen-Fu Chen.

Funding acquisition: Richard J. Lilford.

Investigation: Mary E. Osuh.

Methodology: Mary E. Osuh, Gbemisola A. Oke, Richard J. Lilford, Eme Owoaje, Bronwyn Harris, Olalekan John Taiwo, Godwin Yeboah, Yen-Fu Chen.

Project administration: Mary E. Osuh, Olalekan John Taiwo, Yen-Fu Chen.

Software: Mary E. Osuh, Godwin Yeboah, Samuel I. Watson.

Supervision: Gbemisola A. Oke, Richard J. Lilford, Eme Owoaje, Bronwyn Harris, Yen-Fu Chen.

Validation: Mary E. Osuh, Bronwyn Harris, Yen-Fu Chen.

Visualization: Mary E. Osuh.

Writing - original draft: Mary E. Osuh, Laura Quinn.

Writing – review & editing: Mary E. Osuh, Gbemisola A. Oke, Richard J. Lilford, Eme Owoaje, Bronwyn Harris, Olalekan John Taiwo, Godwin Yeboah, Taiwo Abiona, Samuel I. Watson, Karla Hemming, Laura Quinn, Yen-Fu Chen.

References

- 1. Ezeh A, Oyebode O, Satterthwaite D, Chen Y-F, Ndugwa R, Sartori J, et al. The health of people who live in slums 1 The history, geography, and sociology of slums and the health problems of people who live in slums. The Lancet. 2016; 389(10068):547–58.
- 2. Saglio-Yatzimirsky M-C. Dharavi: from mega-slum to urban paradigm. e-book edition ed. London: Routledge 2021 1 February 2021. 379 p.
- Simon FR, Adeoke A, Adewale B. Slum Settlements Regeneration in Lagos Mega-city: an Overview of a Waterfront Makoko Community. International Journal of Education and Research 2013; 1(3):1– 16.
- 4. Pat-Mbano E, Nwadiaro E. The rise of urban slum in Nigeria: implications on the urban landscape. International Journal of Development and Management Review (INJODEMAR) 2012; 7:257–69.
- Riley LW, Ko AI, Unger A, Reis MG. Slum health: diseases of neglected populations. BMC international health and human rights. 2007; 7:2. https://doi.org/10.1186/1472-698X-7-2 PMID: 17343758
- 6. Mberu BU, Haregu TN, Kyobutungi C, Ezeh AC. Health and health-related indicators in slum, rural, and urban communities: a comparative analysis. Global Health Action. 2016; 9(1):33163.
- Kyobutungi C, Ziraba AK, Ezeh A, Yé Y. The burden of disease profile of residents of Nairobi's slums: Results from a Demographic Surveillance System. Population Health Metrics. 2008; 6(1):1. https://doi. org/10.1186/1478-7954-6-1 PMID: 18331630
- Marlow MA, Maciel ELN, Sales CMM, Gomes T, Snyder RE, Daumas RP, et al. Tuberculosis DALY-Gap: Spatial and Quantitative Comparison of Disease Burden Across Urban Slum and Non-slum Census Tracts. Journal of Urban Health. 2015; 92(4):622–34. https://doi.org/10.1007/s11524-015-9957-0 PMID: 25840553

- Airen B, Dasar P, Nagarajappa S, Kumar S, Jain D, Warhekar S. Dentition status and treatment need in urban slum dwellers in Indore city, Central India. Journal of Indian Association of Public Health Dentistry. 2014; 12(3):163–6.
- Lilford RJ, Oyebode O, Satterthwaite D, Melendez-Torres GJ, Chen YF, Mberu B, et al. Improving the health and welfare of people who live in slums. The Lancet. 2017; 389(10068):559–70. <u>https://doi.org/ 10.1016/S0140-6736(16)31848-7 PMID: 27760702</u>
- Khan MMH, Khan A, Kraemer A, Mori M. Prevalence and correlates of smoking among urban adult men in Bangladesh: slum versus non-slum comparison. BMC Public Health. 2009; 9(1):149. <u>https:// doi.org/10.1186/1471-2458-9-149 PMID: 19463157</u>
- Gupta V, Yadav K, Anand K. Patterns of tobacco use across rural, urban, and urban-slum populations in a north Indian community. Indian journal of community medicine: official publication of Indian Association of Preventive & Social Medicine. 2010; 35(2):245–51.
- Snyder RE, Rajan JV, Costa F, Lima HC, Calcagno JI, Couto RD, et al. Differences in the prevalence of non-communicable disease between slum dwellers and the general population in a large urban area in Brazil. Tropical medicine and infectious disease. 2017; 2(3):47.
- 14. WHO. Oral health: Key facts. The World Health Organization; 2020.
- Peres MA, Macpherson LMD, Weyant RJ, Daly B, Venturelli R, Mathur MR, et al. Oral diseases: a global public health challenge. The Lancet. 2019; 394(10194):249–60. <u>https://doi.org/10.1016/S0140-6736(19)31146-8 PMID: 31327369</u>
- 16. Varenne B. WHO Oral Health Fact Sheet. Geneva 27, Switzerland: Malmo University; 2012.
- Akpata ES. Oral health in Nigeria. International Dental Journal. 2004; 54:361–6. https://doi.org/10. 1111/j.1875-595x.2004.tb00012.x PMID: 15631097
- 18. Petersen P. Improvement of oral health in Africa in the 21st century—the role of the WHO Global Oral Health Programme. Developing Dentistry. 2004; 5:9–20.
- Glick M, Williams DM, Kleinman DV, Vujicic M, Watt RG, Weyant RJ. A new definition for oral health developed by the FDI World Dental Federation opens the door to a universal definition of oral health. British dental journal. 2016; 221(12):792. https://doi.org/10.1038/sj.bdj.2016.953 PMID: 27981999
- Hummel J, Philips K, Holt B, Hayes C. Oral Health: An Essential Component of Primary Care- White Paper Seattle, WA. Qualis Health. In: Health Q, editor. Seattle WA: Safety net medical home initiative; 2015.
- Disease GBD, Injury I, Prevalence C, Looker K. Global, regional, and national indicence, prevalence, and years lived with disability for 355 diseases and injuries for 195 countries, 1990–2017. a systematic analysis for the Global Burden of Disease Study 2017. 2018; 392(10159):1789–858.
- Listl S, Galloway J, Mossey PA, Marcenes W. Global Economic Impact of Dental Diseases. Research Report: Clinical. Journal of Dental Research. 2015; 94:1355–61. <u>https://doi.org/10.1177/</u> 0022034515602879 PMID: 26318590
- Tonetti MS, Jepsen S, Jin L, Otomo-Corgel J. Impact of the global burden of periodontal diseases on health, nutrition and wellbeing of mankind: A call for global action. Journal of clinical periodontology. 2017; 44(5):456–62. https://doi.org/10.1111/jcpe.12732 PMID: 28419559
- Botelho J, Machado V, Leira Y, Proença L, Chambrone L, Mendes JJ. Economic burden of periodontitis in the United States of America and Europe–an updated estimation. Journal of Periodontology. 2021.
- 25. Patel AB, Shah RR, Ramanuj VB. Comparative study of oral hygienic practices and oral health status among people residing in urban and urban slum of Ahmedabad municipal corporation. Int J Community Med Public Health. 2017; 4(6):2181–5.
- Hannan MA, Chowdhury MT, Khan MA, Chowdhury AF, Shahidullah KM, Saha AK, et al. Prevalence of Gingivitis, Plaque accumulation and Decayed, Missing and Filled Teeth among slum population in Bangladesh. Bangladesh Medical Research Council Bulletin. 2014; 40(2):47–51. https://doi.org/10. 3329/bmrcb.v40i2.25182 PMID: 26415339
- Bernabe E, Marcenes W, Hernandez CR, Bailey J, Abreu LG, Alipour V, et al. Global, Regional, and National Levels and Trends in Burden of Oral Conditions from 1990 to 2017: A Systematic Analysis for the Global Burden of Disease 2017 Study. Journal of Dental Research 2020; 99(4):362–73. https://doi.org/10.1177/0022034520908533 PMID: 32122215
- Petersen PE, Bourgeois D, Ogawa H, Estupinan-Day S, Ndiaye C. The global burden of oral diseases and risks to oral health. Bulletin of the World Health Organization. 2005; 83:661–9. <u>https://doi.org// S0042-96862005000900011</u> PMID: 16211157
- 29. Lee J-H, Oh J-Y, Youk T-M, Jeong S-N, Kim Y-T, Choi S-H. Association between periodontal disease and non-communicable diseases: A 12-year longitudinal health-examinee cohort study in South

Korea. Medicine. 2017; 96(26):e7398–e. https://doi.org/10.1097/MD.00000000007398 PMID: 28658175

- Kantovitz KR, Pascon FM, Rontani RMP, Gaviao MBD, Pascon FM. Obesity and dental caries—A systematic review. Oral Health & Preventive Dentistry. 2006; 4(2). PMID: <u>16813143</u>
- Schulze MB, Manson JE, Ludwig DS, Colditz GA, Stampfer MJ, Willett WC, et al. Sugar-sweetened beverages, weight gain, and incidence of type 2 diabetes in young and middle-aged women. JAMA. 2004; 292(8):927–34. https://doi.org/10.1001/jama.292.8.927 PMID: 15328324
- 32. Deprivation Locker D. and oral health: a review. Community Dent Oral Epidemiol. 2000; 28(3):161–9. https://doi.org/10.1034/j.1600-0528.2000.280301.x PMID: 10830642
- 33. Gautam D, Vikas J, Amrinder T, Rambhika T, Bhanu K. Evaluating dental awareness and periodontal health status in different socioeconomic groups in the population of Sundernagar, Himachal Pradesh, India. Journal of International Society of Preventive & Community Dentistry. 2012; 2(2):53. https://doi. org/10.4103/2231-0762.109367 PMID: 24478968
- Sanders AE, Slade GD, Turrell G, John Spencer A, Marcenes W. The shape of the socioeconomic– oral health gradient: implications for theoretical explanations. Community Dentistry and Oral Epidemiology. 2006; 34(4):310–9. https://doi.org/10.1111/j.1600-0528.2006.00286.x PMID: 16856951
- 35. Gautam DK, Vikas J, Amrinder T, Rambhika T, Bhanu K. Evaluating dental awareness and periodontal health status in different socioeconomic groups in the population of Sundernagar, Himachal Pradesh, India. J Int Soc Prev Community Dent. 2012; 2(2):53–7. <u>https://doi.org/10.4103/2231-0762.109367</u> PMID: 24478968
- **36.** Aikins E, Braimoh O. Utilization of dental services among civil servants in Port Harcourt, Nigeria. Journal of Dental Research and Review. 2015; 2(2):62–6.
- Taherdoost H. Determining Sample Size; How to Calculate Survey Sample Size. International Journal of Economics and Management Systems. 2017; 2:236–9.
- Bartlett JE, Kotrlik JW, Higgins CC. Organizational Research: Determining Organizational Research: Determining Appropriate Sample Size in Survey Research. Information Technology, Learning, and Performance Journal. 2001; 19(1):43–50.
- Lawal F, Alade O. Dental caries experience and treatment needs of an adult female population in Nigeria. African Health Sciences. 2017; 17:905. https://doi.org/10.4314/ahs.v17i3.34 PMID: 29085419
- **40.** Umoh AO, Azodo CC. Prevalence of gingivitis and periodontitis in an adult male population in Nigeria. Nigerian Journal of Basic and Clinical Sciences. 2012; 9(2):65–9.
- Bakibinga P, Kabaria C, Kyobutungi C, Manyara A, Mbaya N, Mohammed S, et al. A protocol for a multi-site, spatially-referenced household survey in slum settings: methods for access, sampling frame construction, sampling, and field data collection. BMC Medical Research Methodology. 2019; 19(1):109. https://doi.org/10.1186/s12874-019-0732-x PMID: 31146676
- 42. Petersen P, Baez RJ. Oral Health Surveys Basic Methods. 2013:1–137.
- 43. Wiebe CB, Putnins EE. The periodontal disease classification system of the American Academy of Periodontology-an update. Journal-Canadian Dental Association. 2000; 66(11):594–9. PMID: 11253351
- 44. Nurelhuda N, Malde M, Ahmed M, TA T, Nurelhuda N. Correlation between caries experience in Sudanese school children and dietary habits, according to a food frequency questionnaire and a modified 24-hr recall method. African Journal of Food, Agriculture, Nutrition and Development. 2013; 13(2).
- 45. Cade J, Thompson R, Burley V, Warm D. Development, validation and utilisation of food-frequency questionnaires–a review. Public Health Nutrition. 2002; 5(4):567–87. <u>https://doi.org/10.1079/PHN2001318 PMID: 12186666</u>
- 46. Tjønneland A, Grønbæk M, Stripp C, Overvad K. Wine intake and diet in a random sample of 48763 Danish men and women. The American Journal of Clinical Nutrition. 1999; 69(1):49–54. <u>https://doi.org/10.1093/ajcn/69.1.49 PMID: 9925122</u>
- 47. Akhter R, Hassan NMM, Aida J, Zaman KU, Morita M. Risk indicators for tooth loss due to caries and periodontal disease in recipients of free dental treatment in an adult population in Bangladesh. Oral Health & Preventive Dentistry. 2008; 6(3).
- Susin C, Oppermann RV, Haugejorden O, Albandar JM. Tooth loss and associated risk indicators in an adult urban population from south Brazil. Acta Odontologica Scandinavica. 2005; 63(2):85–93. https://doi.org/10.1080/00016350510019694 PMID: 16134547
- Silva Junior MF, Batista MJ, de Sousa MdLR. Risk factors for tooth loss in adults: A population-based prospective cohort study. Plos One. 2019; 14(7):e0219240. https://doi.org/10.1371/journal.pone. 0219240 PMID: 31329623

- Gilbert GH, Shelton BJ, Chavers LS, Bradford Jr. EH. Predicting Tooth Loss During a Population-Based Study: Role of Attachment Level in the Presence of Other Dental Conditions. Journal of Periodontology. 2002; 73(12):1427–36. https://doi.org/10.1902/jop.2002.73.12.1427 PMID: 12546092
- Pitiphat W, Merchant AT, Rimm EB, Joshipura KJ. Alcohol Consumption Increases Periodontitis Risk. Journal of Dental Research. 2003; 82(7):509–13. <u>https://doi.org/10.1177/154405910308200704</u> PMID: 12821709
- Peres MA, Sheiham A, Liu P, Demarco FF, Silva AER, Assunção MC, et al. Sugar Consumption and Changes in Dental Caries from Childhood to Adolescence. Journal of Dental Research. 2016; 95 (4):388–94. https://doi.org/10.1177/0022034515625907 PMID: 26758380
- Bernabé E, Vehkalahti MM, Sheiham A, Aromaa A, Suominen AL. Sugar-sweetened beverages and dental caries in adults: A 4-year prospective study. Journal of Dentistry. 2014; 42(8):952–8. https://doi. org/10.1016/j.jdent.2014.04.011 PMID: 24813370
- Bellini HT, Arneberg P, Von Der Fehr FR. Oral hygiene and caries. Acta Odontologica Scandinavica. 1981; 39(5):257–65. https://doi.org/10.3109/00016358109162287 PMID: 7039213
- Andlaw RJ. Oral hygiene and dental caries—a review. International dental journal. 1978; 28(1):1–6. PMID: 346493
- Winn DM. Tobacco Use and Oral Disease. Journal of Dental Education. 2001; 65(4):306–12. PMID: 11336115
- Lee KH, Plassman BL, Pan W, Wu B. Mediation Effect of Oral Hygiene on the Relationship Between Cognitive Function and Oral Health in Older Adults. Journal of Gerontological Nursing. 2016; 42 (5):30–7. https://doi.org/10.3928/00989134-20151218-03 PMID: 26716459
- 58. Ritchie CS, Joshipura K, Hung H-C, Douglass CW. Nutrition as a mediator in the relation between oral and systemic disease: associations between specific measures of adult oral health and nutrition outcomes. Critical reviews in oral biology & medicine. 2002; 13(3):291–300. <u>https://doi.org/10.1177/</u> 154411130201300306 PMID: 12090466
- Kassebaum N, Bernabé E, Dahiya M, Bhandari B, Murray C, Marcenes W. Global burden of untreated caries: a systematic review and metaregression. Journal of Dental Research. 2015; 94(5):650–8. https://doi.org/10.1177/0022034515573272 PMID: 25740856
- Watt RG, Petersen PE. Periodontal health through public health-the case for oral health promotion. Periodontology 2000. 2012; 60(1):147–55. <u>https://doi.org/10.1111/j.1600-0757.2011.00426.x</u> PMID: 22909112
- Kaste L, Gift H, Bhat M, Swango P. Prevalence of incisor trauma in persons 6 to 50 years of age: United States, 1988–1991. Journal of Dental Research. 1996; 75(2_suppl):696–705. <u>https://doi.org/10.1177/002203459607502S09</u> PMID: 8594093
- Varenne B, Petersen PE, Fournet F, Msellati P, Gary J, Ouattara S, et al. Illness-related behaviour and utilization of oral health services among adult city-dwellers in Burkina Faso: evidence from a household survey. BMC Health Services Research. 2006; 6:164. <u>https://doi.org/10.1186/1472-6963-6-164</u> PMID: 17192172
- Okunseri C, Born D, Chattopadhyay A. Self-reported dental visits among adults in Benin City, Nigeria. International Dental Journal. 2004; 54:450–6. <u>https://doi.org/10.1111/j.1875-595x.2004.tb00303.x</u> PMID: 15633502
- 64. World Health Organization. Sugars and dental caries. World Health Organization; 2017.
- Marcenes W, Kassebaum NJ, Bernabé E, Flaxman A, Naghavi M, Lopez A, et al. Global burden of oral conditions in 1990–2010: a systematic analysis. J Dent Res. 2013; 92(7):592–7. https://doi.org/ 10.1177/0022034513490168 PMID: 23720570
- 66. Ismail AI, Tanzer JM, Dingle JL. Current trends of sugar consumption in developing societies. Community Dentistry and Oral Epidemiology. 1997; 25(6):438–43. https://doi.org/10.1111/j.1600-0528.1997. tb01735.x PMID: 9429817
- Malago J, Makoba E, Muzuka AN. Fluoride levels in surface and groundwater in Africa: a review. Am J Water Sci Eng. 2017; 3(1):1–17.
- Olusile AO, Adeniyi AA, Orebanjo O. Self-rated oral health status, oral health service utilization, and oral hygiene practices among adult Nigerians. BMC Oral Health. 2014; 14:140. <u>https://doi.org/10. 1186/1472-6831-14-140 PMID: 25427860</u>
- Hardwick K, Barmes D, Writer S, Richardson LM. International Collaborative Research on Fluoride. Journal of Dental Research. 2000; 79(4):893–904. <u>https://doi.org/10.1177/00220345000790040301</u> PMID: 10831090
- 70. World Health Organization. Guidelines for drinking-water quality: second addendum. Vol. 1, Recommendations. 2008.

- Akpata ES, Danfillo I, Otoh E, Mafeni J. Geographical mapping of fluoride levels in drinking water sources in Nigeria. African health sciences. 2009; 9(4). PMID: 21503173
- 72. Ibiyemi O, Ibiyemi S. Fluoride concentrations and labeling information on adult toothpastes from Nigeria and the United Kingdom. African Journal of Medicine and Medical Sciences. 2021; 50(1):107–15.
- Aliu IR, Akoteyon IS, Soladoye O. Living on the margins: Socio-spatial characterization of residential and water deprivations in Lagos informal settlements, Nigeria. Habitat International. 2021; 107:102293.
- Akpabio EM, Wilson N-AU, Essien KA, Ansa IE, Odum PN. Slums, women and sanitary living in South-South Nigeria. Journal of Housing and the Built Environment. 2021; 36(3):1229–48. https://doi. org/10.1007/s10901-020-09802-z PMID: 33424521
- **75.** Ani F, Akaji E, Uguru N, Ndiokwelu E. Fluoride content of commercial drinking water and carbonated soft drinks available in Southeastern Nigeria: dental and public health implications. Niger J Clin Pract. 2020; 23:65–70. https://doi.org/10.4103/njcp.njcp_248_19 PMID: 31929209
- 76. Toumba KJ, Twetman S, Splieth C, Parnell C, van Loveren C, Lygidakis NA. Guidelines on the use of fluoride for caries prevention in children: an updated EAPD policy document. European Archives of Paediatric Dentistry. 2019; 20(6):507–16. <u>https://doi.org/10.1007/s40368-019-00464-2</u> PMID: 31631242
- Cury JA, Tenuta LMA. Evidence-based recommendation on toothpaste use. Brazilian oral research. 2014; 28:1–7. https://doi.org/10.1590/S1806-83242014.50000001 PMID: 24554097
- Walsh T, Worthington HV, Glenny AM, Appelbe P, Marinho VC, Shi X. Fluoride toothpastes of different concentrations for preventing dental caries in children and adolescents. Cochrane database of systematic reviews. 2010(1). https://doi.org/10.1002/14651858.CD007868.pub2 PMID: 20091655
- 79. Oke G. Oral Health care at Primary Health care Level (Section Two). In Primary Mental, Oral, Eye, and Ear Care in Nigeria. Nigerian Health Review 2007: Primary Health Care in Nigeria: 30 Years afer Alma Ata. National Health Review. Abuja, Nigeria: Health Reform Foundation of Nigeria (HERFON); 2007. p. 226–34.
- Okeigbernen SA, Ibiyemi O. Prevention of Dental Caries in Nigeria: A Narrative Review of Strategies and Recommendations from 1999 to 2019. J Int Soc Prev Community Dent. 2020; 10(3):240–5. https://doi.org/10.4103/jispcd.JISPCD 423 19 PMID: 32802768
- **81.** Jaafar N, Hakim H, Mohd Nor NA, Mohamed A, Saub R, Esa R, et al. Is the burden of oral diseases higher in urban disadvantaged community compared to the national prevalence? BMC Public Health. 2014; 14(3):S2.
- Nazir M, Al-Ansari A, Al-Khalifa K, Alhareky M, Gaffar B, Almas K. Global Prevalence of Periodontal Disease and Lack of Its Surveillance. The Scientific World Journal. 2020; 2020:2146160. <u>https://doi.org/10.1155/2020/2146160</u> PMID: 32549797
- 83. Gaines J, Malumfashi D, Omenka S, Oti N. TAPI-Desk Study. 2020.
- Northridge ME, Kumar A, Kaur R. Disparities in Access to Oral Health Care. Annual Review of Public Health. 2020; 41(1):513–35. https://doi.org/10.1146/annurev-publhealth-040119-094318 PMID: 31900100
- Adeniyi A, Sofola O, Kalliecharan R. An appraisal of the oral health care system in Nigeria. International Dental Journal. 2012; 62:292–300. https://doi.org/10.1111/j.1875-595X.2012.00122.x PMID: 23252586
- 86. Oke GA, Bankole OO, Denloye OO, Danfillo IS, Enwonwu CO. Traditional and emerging oral health practices in parts of Nigeria. Odonto-stomatologie tropicale = Tropical dental journal. 2011; 34 (136):35–46. PMID: 22457991
- Adeniyi A, Onajole A. The National Health Insurance Scheme (NHIS): a survey of knowledge and opinions of Nigerian dentists' in Lagos. African Journal of Medicine and Medical Sciences. 2010; 39(1):29– 35. PMID: 20632669
- Helderman WvP, Benzian H. Implementation of a Basic Package of Oral Care: towards a reorientation of dental NGOs and their volunteers. International dental journal. 2006; 56(1):44–8. https://doi.org/10. 1111/j.1875-595x.2006.tb00073.x PMID: 16515012
- Ganss C, Schlueter N, Preiss S, Klimek J. Tooth brushing habits in uninstructed adults—frequency, technique, duration and force. Clinical Oral Investigations. 2009; 13(2):203–8. <u>https://doi.org/10.1007/</u> s00784-008-0230-8 PMID: 18853203
- **90.** Attin T, Hornecker E. Tooth brushing and oral health: how frequently and when should tooth brushing be performed? Oral Health & Preventive Dentistry. 2005; 3(3). PMID: <u>16355646</u>
- Bashiru BO, Anthony IN. Oral self-care practices among university students in Port Harcourt, Rivers State. Nigerian medical journal: journal of the Nigeria Medical Association. 2014; 55(6):486–9. https://doi.org/10.4103/0300-1652.144703 PMID: 25538367

- 92. Msyamboza KP, Phale E, Namalika JM, Mwase Y, Samonte GC, Kajirime D, et al. Magnitude of dental caries, missing and filled teeth in Malawi: National Oral Health Survey. BMC Oral Health. 2016; 16 (1):1–6. https://doi.org/10.1186/s12903-016-0190-3 PMID: 26956884
- **93.** Sheiham A. Changing Trends in Dental Caries. International Journal of Epidemiology. 1984; 13 (2):142–7. https://doi.org/10.1093/ije/13.2.142 PMID: 6376384
- Mazengo MC, Tenovuo J, Hausen H. Dental caries in relation to diet, saliva and cariogenic microorganisms in Tanzanians ot selected age groups. Community Dentistry and Oral Epidemiology. 1996; 24(3):169–74. https://doi.org/10.1111/j.1600-0528.1996.tb00836.x PMID: 8871014
- Enwonwu CO. Socio-economic factors in dental caries prevalence and frequency in Nigerians. Caries Research. 1974; 8(2):155–71. https://doi.org/10.1159/000260104 PMID: 4525913
- Moreira ARO, Batista RFL, Ladeira LLC, Thomaz EBAF, Alves CMC, Saraiva MC, et al. Higher sugar intake is associated with periodontal disease in adolescents. Clinical Oral Investigations. 2021; 25 (3):983–91. https://doi.org/10.1007/s00784-020-03387-1 PMID: 32519237
- Lula EC, Ribeiro CC, Hugo FN, Alves CM, Silva AA. Added sugars and periodontal disease in young adults: an analysis of NHANES III data. The American Journal of Clinical Nutrition. 2014; 100 (4):1182–7. https://doi.org/10.3945/ajcn.114.089656 PMID: 25240081
- Sidi AD, Ashley FP. Influence of Frequent Sugar Intakes on Experimental Gingivitis. Journal of Periodontology. 1984; 55(7):419–23. https://doi.org/10.1902/jop.1984.55.7.419 PMID: 6379142
- 99. Kumar S, Tadakamadla J, Johnson N. Effect of toothbrushing frequency on incidence and increment of dental caries: a systematic review and meta-analysis. Journal of Dental Research. 2016; 95 (11):1230–6. https://doi.org/10.1177/0022034516655315 PMID: 27334438
- **100.** Frandsen A. Mechanical oral hygine practices. Dental Plaque Control Measures and Oral Hygine Practices. 1986:93–116.
- 101. Ashley P. Toothbrushing: why, when and how? Dental Update. 2001; 28(1):36–40. https://doi.org/10. 12968/denu.2001.28.1.36 PMID: 11819948
- 102. Asadoorian J. Tooth brushing. Canadian Journal of Dental Hygiene. 2006; 40:1–14.
- 103. Bakdash B. Current patterns of oral hygiene product use and practices. Periodontology 2000. 1995; 8 (1):11–4. https://doi.org/10.1111/j.1600-0757.1995.tb00041.x PMID: 9567942
- 104. Isidor F, Karring T. Long-term effect of surgical and non-surgical periodontal treatment. A 5-year clinical study. Journal of Periodontal Research. 1986; 21(5):462–72. https://doi.org/10.1111/j.1600-0765. 1986.tb01482.x PMID: 2946847
- **105.** Adeniyi AA, Oyapero A. Predisposing, enabling and need factors influencing dental service utilization among a sample of adult Nigerians. Population Medicine. 2020; 2(December):1–9.
- Levy PS, Lemeshow S. Sampling of populations: methods and applications: John Wiley & Sons; 2013.