

What is the most appropriate method for the measurement of the range of motion in the lumbar spine and how does surgical fixation affect the range of movement of the lumbar spine in adolescent idiopathic scoliosis?

Hartley, Laura; Zappalà, Mattia; Ehiogu, Uzo; Heneghan, Nicola R; Gardner, Adrian

DOI:

[10.1186/s13643-022-02077-1](https://doi.org/10.1186/s13643-022-02077-1)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Hartley, L, Zappalà, M, Ehiogu, U, Heneghan, NR & Gardner, A 2022, 'What is the most appropriate method for the measurement of the range of motion in the lumbar spine and how does surgical fixation affect the range of movement of the lumbar spine in adolescent idiopathic scoliosis? A systematic review protocol', *Systematic Reviews*, vol. 11, no. 1, 208, pp. 7. <https://doi.org/10.1186/s13643-022-02077-1>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.


If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

PROTOCOL

Open Access



What is the most appropriate method for the measurement of the range of motion in the lumbar spine and how does surgical fixation affect the range of movement of the lumbar spine in adolescent idiopathic scoliosis? A systematic review protocol

Laura Hartley¹ , Mattia Zappalà², Uzo Ehiogun¹, Nicola R. Heneghan² and Adrian Gardner^{1,3*}

Abstract

Introduction: Adolescent idiopathic scoliosis (AIS) is a three-dimensional rotational change in the normal shape of the spine which affects children aged 10 to 18 years. Both the condition and its management can have significant impact on functional ability. Currently, expected restriction in spinal motion is experience based, rather than evidence based, and discussions to inform patient expectations pre-operatively can be difficult. The aim of this review is to evaluate the evidence pertaining to measurement of spinal motion and whether this is altered following surgery, dependent on the anatomical level of surgical fixation in AIS.

Methods/analysis: This protocol is reported in line with both PRISMA-P and informed by the COSMIN methodology. Electronic databases will be searched using a two-stage search strategy. The first stage will identify and evaluate the methods used to assess spinal motion. The second stage aims to evaluate the change in spinal motion using these methods based on anatomical level of fixation following surgery along with the measurement properties of those methods, to include the validity, reliability and responsiveness of the methods. Two reviewers will independently screen the search results against eligibility criteria, extract the data and assess the quality of the included studies. Any disputes between the reviewers will be resolved with a third independent reviewer. Data may be pooled where possible; however, this is not expected. The overall strength of the body of evidence will then be assessed using the Grading of Recommendations, Assessment, Development and Evaluation (GRADE) approach.

Patient and public involvement: Patients and members of the public will not be consulted in the production of this review, although the review was conceived based on the experiences of the authors when managing this patient population and a need to address patient expectations in pre-operative planning.

*Correspondence: adrian.gardner@nhs.net

¹ The Royal Orthopaedic Hospital NHS Foundation Trust, The Woodlands, Bristol Road South, Northfield, Birmingham B31 2AP, UK
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Ethics, dissemination and data availability: No ethical approval required. The final review will be submitted to peer-reviewed journals for publication and disseminated publicly. The datasets used and/or analysed in this review will be available from the corresponding author on reasonable request.

Systematic review registration: PROSPERO registration number.

CRD42021282264.

Keywords: Adolescent idiopathic scoliosis, Scoliosis, Lumbar motion, Range of motion, Spinal motion, Function, Surgery, Surgical intervention, Instrumentation, Fusion, Fixation

Strengths and limitations

- A two-stage search strategy will be used to identify current methods of assessing spinal motion and then, using the methods identified in stage 1, identify evidence regarding how spinal motion is affected with different levels of surgical fixation in AIS.
- This protocol has been designed in line with COSMIN methodology.
- Two independent reviewers have been used to extract data, assess quality and perform the analysis.
- Excluding studies published in languages other than English may result in some relevant studies being excluded.

Introduction

Scoliosis is defined as an abnormal lateral curvature of the spine with rotational deformity and a Cobb angle greater than 10 degrees in the coronal plane [1]. In the absence of an identifiable underlying cause and when diagnosed between the ages of 10 and 18 years, it is known as adolescent idiopathic scoliosis (AIS) [2]. AIS is the most common form of scoliosis, affecting up to 4% of teenagers with a roughly equal sex distribution [1]. ‘Idiopathic’ refers to the unclear aetiology of this condition however, hormonal imbalance, asymmetric growth and muscle imbalance are thought to be contributory [3]. Up to 30% of those affected have a positive family history indicating a genetic component [3]. Generally, AIS curves will progress during the rapid growth that accompanies puberty and most curves will eventually slow their progression by the time skeletal maturity is reached [1–4]. Some curves continue to progress into adulthood [1].

Whilst untreated AIS does not increase mortality rates [4], when compared with those without scoliosis, most teenagers with AIS function at or near normal levels [4]. Severe curves may impair functional ability and cause pain [4, 5]. In those affected, the potential caregiver and financial burden can be significant and, therefore, the economic impact of AIS is greater than in healthy age-matched peers [5].

The management of AIS primarily aims to prevent progression through minimising deformity alongside maximising function [6]. Patients with a Cobb angle of 25–30° prior to skeletal maturity, or less than 45° after, may be managed with radiographic surveillance [7]. Similarly, bracing has been shown to be effective in reducing the need for surgical intervention [8]. Surgical intervention aims to correct the deformity, prevent future progression and improve functional ability [6]. It should be considered for those with curves greater than 45° prior to skeletal maturity and those with progression beyond 45° after skeletal maturity is achieved [9].

Many methods exist for determining the anatomical inferior level at which instrumentation should extend. The decision takes into account various factors including curve type, Cobb angle, pre-operative mobility (a systematic review of which method of assessment is most reliable and accurate is underway by another group [10]) and the likely post-operative restriction in movement [11–13]. Restriction in spinal motion is a recognised and expected outcome of corrective surgery for deformity in AIS [4]. When discussing surgical intervention, information regarding the reduction in spinal motion post-operatively is usually experience-based rather than evidence based. It can be difficult to predict how much movement may be lost and therefore what post-operative activities an individual may experience difficulty with [11]. Recent studies have highlighted that the restriction in motion post-operatively is greater with increasing overall number of levels of fixation [12, 13] and with instrumentation that extends to and caudal to L3/4 [12–14]. Over time, abnormal or increased segmental motion in the adjacent unfused spinal levels post fixation, thought to lead to degeneration, and may be responsible for ongoing pain [14–18]. Avoidance of instrumenting the inferior lumbar spine has been recommended where possible to reduce the risk of post-operative pain and preserve as much spinal motion as possible [19–22].

Current methods for assessing the post-operative spinal shape [23] are useful to assess the level of correction achieved and detect complications such as metalwork failure; however, they are limited by their inability to quantify parameters related to spinal motion and have

been cited to be unreliable, insensitive and non-specific, especially in the context of ongoing pain [24–27]. Alternative strategies exist, including the use of skin based motion trackers [17] and fingertip-to-floor distance [14]; however, these have also been found to be inaccurate and limited by their inability to pinpoint precise anatomical areas of abnormal motion within the spine. Quantitative fluoroscopy (QF) has shown promise in the assessment and identification of potential therapeutic targets in chronic, non-specific lower back pain (CNSLBP) [24–28]. Several studies have been carried out assessing the validity of QF both in healthy controls and individuals with CNSLBP, where it has been shown to be both reliable and valid [24, 29–31]. QF also has the added benefit of lower overall radiation exposure compared to lumbar spine radiography [32]. However, QF has not yet been investigated in AIS. To our knowledge, no studies exist comparing pre-operative spinal motion directly with post-operative spinal motion in AIS and whilst studies exist looking at the level of function post-operatively based on instrumentation levels, the results are mixed [14, 16, 19].

Aims

Our aims with this study are twofold. Firstly is to collate and evaluate current methods of assessing spinal motion in AIS. Secondly is to compare these identified methods for measuring pre- to post-operative spinal motion based on the level of instrumentation along with the measurement properties of these methods. Combined, this data will allow the assessment of methods for the measurement, and any post-operative loss of movement.

Design and methods

This protocol has been designed according to the Consensus-based Standards for the Selection of Health Measurement Instruments (COSMIN) methodology for systematic reviews [33] and is reported in line with Preferred Reporting Items for Systematic Review and Meta-Analysis Protocol (PRISMA-P) [34]. The design and methods have been informed through a collaboration between experts in physiotherapy, rehabilitation and scoliosis management. It has also been registered in the International Prospective Register of Systematic Reviews (PROSPERO ID – CRD42021282264).

The proposed methodology has a two-stage approach—in stage 1, scoping searches will be conducted to identify both historic and contemporary methods of spinal motion assessment in AIS. In stage 2, searches will be conducted for studies that evaluate the effects of surgery on spinal motion based on the anatomical levels of instrumentation.

Stage 1: Identifying and evaluating the methods used to assess spinal motion

Eligibility criteria

Population Studies assessing spinal motion in the age group defined as AIS [2] (i.e. aged 10 to 18 years inclusive) will be included.

Outcome Any study that includes assessment of spinal motion will be included. No restrictions will be applied on the method of assessment of spinal motion.

Study design Any study design reporting quantitative data will be included, e.g. randomised clinical trials, cohort, observational and case studies. There will be no limitation placed on geographical location. Studies published in languages other than English will be excluded.

Search strategy

The search strategy has been developed following scoping searches and discussions with experts (subject specific, methodological and a medical librarian). It will involve systematic searches of electronic databases with structured search blocks. The search will be completed by one reviewer (LH). Stage 1 search blocks will contain terms relevant to the population of interest (patients defined as having AIS [2]) and the outcome of interest (spinal motion, any parameter, measured by any method). Unformatted terms may also be used—these are free text terms that require appropriate formatting, prefixing and/or suffixing for searching the relevant database. An example of the search strategy used can be found in Additional file 1.

The title and abstracts of generated studies will be screened independently by two reviewers (LH and MZ). Studies will be categorised as relevant, irrelevant or unsure. Any studies considered irrelevant based on title and abstract by both reviewers will be excluded at this stage. Where both reviewers are unsure or disagree, a third arbitrating reviewer (AG) will make the final decision. Studies deemed relevant to the study question in stage 1 based on abstract and title will proceed to data extraction; the full text articles will be reviewed and categorised into relevant, irrelevant and unsure. For relevant studies, the method used to assess spinal motion will be recorded along with the outcomes. Irrelevant studies will be excluded. Studies where the reviewers are unsure will be reviewed by a third arbitrating reviewer (AG) and, if deemed relevant, will proceed to data extraction. This first stage is expected to yield a list of methods used to evaluate spinal motion,

along with outcomes, merits and disadvantages of each method.

Stage 2: Evaluating change in spinal motion post-operatively dependent on the anatomical levels of instrumentation

Eligibility criteria

Population Patients aged 10 to 18 years, diagnosed with AIS, managed with surgical intervention of any type, will be included. Animal studies will not be considered.

Outcome Any study that includes the assessment of post-operative spinal motion based on level of instrumentation will be included. No restrictions will be applied on method of assessment of post-operative spinal motion or to the mode of instrumentation used.

Study design The study design is as described in the stage 1 inclusion criteria.

Information sources

The search strategy for stage 2 will involve systematic searches of electronic databases, trial registries, grey literature and experts in the field. An electronic search will be performed through PubMed, Scopus, Web of Science, Cochrane database, EMBASE, MedLine and Ovid bibliographic databases from inception to the date of the last search. Other sources will be searched for grey literature, ProQuest for dissertations and meeting abstracts through Scopus, Web of Science and pertinent websites. Reference lists of relevant studies and systematic reviews will be searched as well.

Search strategy

Searches of the electronic databases will be conducted in a similar manner to stage 1 using structured search blocks in order to identify studies that measure changes in spinal motion following surgery using the methods identified in stage 1. The search will be conducted by one reviewer (LH). To identify appropriate key words, in addition to Medical Subject Headings (MeSH terms), popular and commonly used phrases stated in relation to the literature will be used. The search strategy will be developed in MedLine, then the same search strategy applied to the other databases. The search will be conducted using terms relevant to the population of interest (patients with AIS between 10 and 18 years old inclusive) and method of assessing spinal motion identified in stage 1 with the limits detailed above applied. An example of the search strategy used can be found in Additional file 2.

Selection process

Two authors will independently perform the initial title and abstract screening (LH and MZ). The studies will be organised into three groups-relevant, irrelevant and unsure. Studies categorised as 'irrelevant' based on title and abstract by both reviewers will be excluded from the study. Where studies are categorised as 'unsure' by both reviewers or where there is a disagreement, there will be a third review by an arbitrator (AG). Once a list of studies to be considered has been generated, searching for full text articles will take place. Both reviewers will then review the full text of the eligible studies and organise them into 'relevant', 'irrelevant' and 'unsure' groups. The two lists will then be compared and any non-conformities (particularly where both reviewers are unsure) discussed with the third arbitrating reviewer (AG) in the same manner as in stage 1. Relevant studies will proceed to critical appraisal and data extraction. Irrelevant studies will be excluded. This will yield a list of studies relevant to the study question to be included in the review.

Critical appraisal of studies and risk of bias

Critical appraisal of the studies will be undertaken using the COSMIN Risk of Bias tool checklists relevant to the individual study types to assess the methodological quality of each article [34]. The full texts will be appraised by two reviewers independently (LH and MZ). Studies of sufficient quality will then proceed to data extraction and synthesis.

Data management

Eligible search records will be imported into Mendeley Reference Management software (London, UK). If required data is missing from the full text papers, or are incomplete or unclear, enquiries will be sent to the authors.

Data items

From each paper, the following data will be extracted.

Study and participant details

Reference, year of publication, geographical location of study, age, sex, sample size, curve type, curve severity and pattern, surgical intervention including but not limited to type of instrumentation, number of levels fused, most superior (upper instrumented vertebra or UIV) or inferior (lowest instrumented vertebra or LIV), complications

Method of post-operative spinal motion measurement	Name of method, outcomes obtained (including but not limited to rotational stability, motion sharing inequality and motion sharing variance), assessment of method (including but not limited to measures of validity, reliability, sensitivity and specificity)
Method properties	<p><i>Validity</i> (including type of validity, descriptive statistics, comparator/predictor outcome, hypothesis, missing value, confidence intervals, sample size and validation results)</p> <p><i>Reliability</i> (including type, descriptive statistics, time intervals, reliability coefficients, measurement errors, number of repeated measurements and sample size)</p> <p><i>Responsiveness</i> (including method of testing, whether hypothesis tested, distribution or anchor based, follow-up length, severity of curve prior to surgery, aetiology if available, intervention modality)</p> <p><i>Interpretability</i> (including distribution of scores within the population studied)</p> <p><i>Feasibility</i> (including type and ease of surgical intervention, patient comprehension, cost of intervention, required equipment if stated, availability in different settings, applicability of study sample to other populations with the same diagnosis)</p>

Data synthesis

COSMIN guidelines for systematic reviews will be followed for the synthesis of the data extracted [34]. Data on the methods of the assessment of the loss of spinal motion caused by surgery, its measurement properties, interpretability and feasibility will be presented in an overview table for ease of interpretation. Method properties will be evaluated as per COSMIN methodology against the criteria for good measurement properties and determined as being 'sufficient', 'insufficient' or 'indeterminate' [34]. Following completion of the overview table, the results of each study will be compared based on the level of instrumentation. If the studies show sufficient methodological and clinical homogeneity, the results will be combined. Pooling of results will only occur where there is sufficient comparison between patient groups studied including similar age range, sex distribution, curve type and severity, similar surgical interventions made (e.g. anterior stabilisation only, posterior stabilisation only or a combination of the two) and where there is comparison between the method of measuring post-operative spinal mobility over a similar length of follow-up. From the scoping searches, these authors do not anticipate that any pooling will occur and

that, consequently, a narrative synthesis of data will be reported. If appropriate, subgroup analysis can be carried out at this stage (e.g. by sex, curve severity, surgical intervention).

The recommendation of a method for the evaluation of the change in spinal motion following scoliosis surgery will depend on the properties, interpretability and feasibility of the methods described, along with an assessment of confidence in the evidence using the Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach [35, 36]. As per the COSMIN guidelines, only methods with sufficient validity and at least low-quality evidence for sufficient internal consistency will be recommended [37].

Discussion and implications of the review

The primary goals of the surgical management of AIS are to correct the spinal deformity and to prevent future progression. A side effect of this surgery, however, is a potential reduction in spinal motion post-operatively. Surgeons recognise that both AIS and its management can impact on spinal motion and therefore an individuals' function in daily life [12, 14–17, 20]. Understanding the impact of different anatomical levels of instrumentation is crucial to clinical practice. A review looking at how differing anatomical levels of surgical fixation affect post-operative range of motion is therefore justified and this protocol aims to provide a transparent framework for a comprehensive overview and critical appraisal of the current literature. The authors expect that this work will benefit those that treat AIS, around the expected reduction in spinal motion following surgery and facilitating evidence-based discussions with patients pre-operatively. Additionally, the review will present sufficient quality evidence regarding the methods of assessing spinal motion to influence a change in the methods currently employed.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13643-022-02077-1>.

Additional file 1. Example Search Strategy Stage 1: methods of assessment of spinal motion.

Additional file 2. Example Search Strategy Stage 2: evaluating post-operative spinal motion based on the level of instrumentation.

Acknowledgements

Not applicable.

Amendments, report writing and dissemination

Any amendments to the protocol will be documented with reference to saved searches and analysis methods recorded in bibliographic databases for data collection and synthesis. Any amendments will also be noted in the final systematic review report. Writing of the systematic report will follow current PRISMA guidelines. The final article will be disseminated in an open-access

journal. The findings will also be disseminated in various relevant proceedings such as conferences, congresses, seminars and symposia.

Authors' contributions

All authors were involved in the conceptualisation and design of this protocol. LH and MZ drafted the manuscript. AG, UE and NH reviewed the manuscript and provided guidance on methodology, design, topic and analysis. All authors subsequently reviewed and commented on every draft of the protocol. All authors reviewed, contributed to and approved the final manuscript.

Author's information

Not applicable.

Funding

No funding was received for this work.

Availability of data and materials

The datasets used and/or analysed in this review are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

Ethical approval is not deemed necessary for the purpose of systematic review. Patients and the general public will not be consulted in the production of the review. Consent to participate is not needed.

Consent for publication

Patient consent is not required for this research or its publication.

Competing interests

No competing interests have been declared.

Author details

¹The Royal Orthopaedic Hospital NHS Foundation Trust, The Woodlands, Bristol Road South, Northfield, Birmingham B31 2AP, UK. ²School of Sport, Exercise and Rehabilitation Sciences, University of Birmingham, Birmingham, UK. ³Institute of Clinical Sciences, University of Birmingham, Birmingham, UK.

Received: 22 November 2021 Accepted: 18 September 2022

Published online: 30 September 2022

References

- Konieczny MR, Senyurt H, Krauspe R. Epidemiology of adolescent idiopathic scoliosis. *J Child Orthop*. 2013;7(1):3–9.
- James JJ. Idiopathic scoliosis: the prognosis, diagnosis and operative indication related to curve patterns and the name of onset. *J Bone Joint Surg Br*. 1954;36-B(1):36–49.
- Kikanloo SR, Tarpada SP, Cho W. Etiology of adolescent idiopathic scoliosis: a literature review. *Asian Spine J*. 2019;13(3):519–26.
- Asher MA, Burton DC. Adolescent idiopathic scoliosis: natural history and long-term treatment effects. *Scoliosis*. 2006;1:2.
- Spinal Deformity BMUS: The burden of musculoskeletal diseases in the United States <https://www.boneandjointburden.org/fourth-edition/iib0/spinal-deformity> Accessed 21 Sept 2021
- Choudhry MN, Ahmad Z, Verma R. Adolescent idiopathic scoliosis. *Open Orthop*. 2016;10:143–54.
- SRS Education Resource Centre Text Book, Knowledge Domain III: Non-Operative Considerations online <https://educationresources.srs.org/adolescent-idiopathic-scoliosis/idiopathic-scoliosis> Accessed 10 Nov 2021
- Weinstein SL, Dolan LA, Wright JG, Dobbs MB. Effects of bracing in adolescents with idiopathic scoliosis. *N Eng J Med*. 2013;369:1512–21.
- Horne JP, Flannery R, Usman S. Adolescent idiopathic scoliosis: diagnosis and management. *Am Fam Physician*. 2014;89(3):193–8.
- Jalalabadi MK, Chan A, Le L, Pachêco-Pereira C, Lou E (registered on PROSPERO 2017). Accuracy and reliability of radiographic methods on spinal flexibility to predict surgical outcomes in AIS patients: a systematic review https://www.crd.york.ac.uk/prospéro/display_record.php?RecordID=57960. Accessed 21 Sept 2021.
- Fabricant P, Admoni S, Green DW, Ipp LS, Widmann RF. Return to athletic activity after posterior spinal fusion for adolescent idiopathic scoliosis; analysis of independent predictors. *J Pediatr Orthop*. 2012;32:259–65.
- Trobisch P, Ducoffe AR, Lonner BS, Errico TJ. Choosing fusion levels in adolescent idiopathic scoliosis. *J Am Acad Orthop Surg*. 2013;21(9):519–28.
- Kim DH, Hyun SJ, Kim KJ. Selection of fusion level for adolescent idiopathic scoliosis: selective fusion vs postoperative decompensation. *J Koren Neurosurg Soc*. 2021;64(4):473–85.
- Uehara M, Takahashi J, Ikegami S, Kuraishi S, Futatsugi T, Oba H, et al. Correlation of lower instrumented vertebra with spinal mobility and health-related quality of life after posterior spinal fusion for adolescent idiopathic scoliosis. *Clin Spine Surg*. 2019;32(7):326–9.
- Breen A, Breen A. Uneven intervertebral motion sharing is related to disc degeneration and is greater in patients with chronic, non-specific low back pain: an in vivo, cross-sectional cohort comparison of intervertebral dynamics using quantitative fluoroscopy. *Eur Spine J*. 2018;27(1):145–53.
- Louer C, Yaszay B, Cross M, Bartley CE, Bastrom TP, Shah SA, et al. Ten-year outcomes of selective fusions for adolescent idiopathic scoliosis. *J Bone Joint Surg Am*. 2019;101(9):761–70.
- Wilk B, Karol LA, Johnston CE, Colby S, Haideri N. The effect of scoliosis fusion on spinal motion: a comparison of fused and non-fused patients with idiopathic scoliosis. *Spine (Phila PA 1976)*. 2006;31(3):309–14.
- Nohara A, Kawakami N, Seki K, Tsuji T, Ohara T, Saito T, et al. The effects of spinal fusion on lumbar disc degeneration in patients with adolescent idiopathic scoliosis: a minimum 10-year follow-up. *Spine Deform*. 2015;3(5):462–8.
- Lenke LG, Bridwell KH, Baldus C, Blanke K, Schoenecker PL. Ability of Cotrel-Dubousset instrumentation to preserve distal lumbar motion segments in adolescent idiopathic scoliosis. *J Spine Disord*. 1993;6(4):339–50.
- Betz RR, Harms J, Clements DH, Lenke LG, Lowe TG, Shuffeberger HL, et al. Comparison of anterior and posterior instrumentation for correction of adolescent thoracic idiopathic scoliosis. *Spine (Phila PA 1976)*. 1999;24(3):225–39.
- Gingsburg HH et al. Longitudinal study of back pain in postoperative idiopathic scoliosis: long-term follow-up phase IV. Paper presented at the Scoliosis Research Society 30th Annual Meeting, September 1995, Asheville, NC. 1995
- Eardley-Harris, Munn Z, Geiroba T (registered on PROSPERO 2015). The effectiveness of selective thoracic fusion for treating adolescent idiopathic scoliosis: a systematic review protocol https://www.crd.york.ac.uk/prospéro/display_record.php?RecordID=32771 Accessed 21 Sept 2021
- Yang X, Xu H, Li M, Gu S, Fang X, Wang J, et al. Clinical and radiographic outcomes of the treatment of adolescent idiopathic scoliosis with segmental pedicle screws and combined local autograft and allograft bone for spinal fusion: a retrospective case series. *BMC Musculoskelet Disord*. 2010;11:159.
- Mellor FE, Muggleton JM, Bagust J, Mason W, Thomas PW, Breen AC. Mid-lumbar lateral flexion stability measured in healthy volunteers by in vivo fluoroscopy. *Spine*. 2009;34(22):811–7.
- Breen A, Breen A. Dynamic interactions between lumbar intervertebral motion segments during forward bending and return. *J Biomech*. 2020;102:109603.
- Dvorák J, Panjabi MM, Chang DG, Theiler R, Grob D. Functional radiographic diagnosis of the lumbar spine Flexion-extension and lateral bending. *Spine (Phila PA 1976)*. 1991;16(5):562–71.
- Panjabi M, Chang D, Dvorák J. An analysis of errors in kinematic parameters associated with in vivo functional radiographs. *Spine (Phila PA 1976)*. 1992;17(2):200–5.
- Mayer RS, Chen IH, Lavender SA, Trafimow JH, Andersson GB. Variance in the measurement of sagittal lumbar spine range of motion among examiners, subjects and instruments. *Spine (Phila PA 1976)*. 1995;20(13):1489–93.
- Breen A, Hemming R, Mellor FE, Breen A. Intrasubject repeatability of in vivo intervertebral motion parameters using quantitative fluoroscopy. *Eur Spine J*. 2019;28(2):450–60.
- Breen A, Claerhout E, Hemming R, Ayer R, Breen A. Comparison of intra-subject repeatability of quantitative fluoroscopy and static radiography

- in the measurement of lumbar intervertebral flexion translation. *Sci Rep.* 2019;9(1):19253.
31. Mellor FE, Thomas PW, Thompson P, Breen AC. Proportional lumbar spine inter-vertebral motion patterns: a comparison of patients with chronic, non-specific low back pain and healthy controls. *Eur J Spine.* 2014;23:2059–67.
 32. Mellor FE, Thomas P, Breen A. Moving back: The radiation dose received from lumbar spine quantitative fluoroscopy compared to lumbar spine radiographs with suggestions for dose reduction. *Radiography.* 2014;20(3):251–7.
 33. Mokkink LB, Prinsen CAC, Patrick DL, Alonso J, Bouter LM, de Vet HCW, Terwee CB. COSMIN manual for systematic reviews of PROMs user manual p1–78. 2018. https://www.cosmin.nl/wp-content/uploads/COSMIN-syst-review-for-PROMs-manual_version-1_feb-2018-1.pdf. Accessed 16 Sept 2021.
 34. Mokkink LB, Boers M, van der Vleuten C, Patrick DL, Alonso J, Bouter LM, et al. COSMIN Risk of Bias tool to assess the quality of studies on reliability and measurement of error of outcome measurement instrument user manual p1–70. 2021. https://www.cosmin.nl/wp-content/uploads/user-manual-COSMIN-Risk-of-Bias-tool_v4_JAN_final.pdf. Accessed 21 Sept 2021
 35. Guyatt G, Oxman AD, Akl EA, Kunz R, Vist G, Brozek J, et al. GRADE guidelines: 1. Introduction – GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol.* 2011;64(4):383–94.
 36. Balshem H, Helfand M, Schunemann HJ, Oxman AD, Kunz R, Brozek J, et al. GRADE guidelines: 3 Rating the quality of evidence. *J Clin Epidemiol.* 2011;64(4):401–6.
 37. Moher D, Shamseer L, Clarke M, Ghersi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev.* 2016;4(1):1.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

