

# Type I interferon rapidly restricts infectious hepatitis C virus particle genesis

Meredith, Luke W.; Farquhar, Michelle J.; Tarr, Alexander W.; Mckeating, Jane A.

DOI:  
[10.1002/hep.v60.6](https://doi.org/10.1002/hep.v60.6)

License:  
Creative Commons: Attribution (CC BY)

*Document Version*  
Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*  
Meredith, LW, Farquhar, MJ, Tarr, AW & Mckeating, JA 2014, 'Type I interferon rapidly restricts infectious hepatitis C virus particle genesis', *Hepatology*, vol. 60, no. 6, pp. 1891-1901. <https://doi.org/10.1002/hep.v60.6>

[Link to publication on Research at Birmingham portal](#)

**Publisher Rights Statement:**  
Eligibility for repository : checked 1/12/2014

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# Type I Interferon Rapidly Restricts Infectious Hepatitis C Virus Particle Genesis

Luke W. Meredith,<sup>1</sup> Michelle J. Farquhar,<sup>1</sup> Alexander W. Tarr,<sup>2</sup> and Jane A. McKeating<sup>1,3</sup>

**Interferon-alpha (IFN $\alpha$ ) has been used to treat chronic hepatitis C virus (HCV) infection for over 20 years with varying efficacy, depending on the infecting viral genotype. The mechanism of action of IFN $\alpha$  is not fully understood, but is thought to target multiple stages of the HCV lifecycle, inhibiting viral transcription and translation leading to a degradation of viral RNA and protein expression in the infected cell. IFN $\alpha$  induces the expression of an array of interferon-stimulated genes within minutes of receptor engagement; however, the impact of these early responses on the viral lifecycle are unknown. We demonstrate that IFN $\alpha$  inhibits the genesis of infectious extracellular HCV particles within 2 hours of treating infected cells, with minimal effect on the intracellular viral burden. Importantly, this short duration of IFN $\alpha$  treatment of infected cells significantly reduced cell-free and cell-to-cell dissemination. The secreted viral particles showed no apparent change in protein content or density, demonstrating that IFN $\alpha$  inhibits particle infectivity but not secretion rates. To investigate whether particles released from IFN $\alpha$ -treated cells have a reduced capacity to establish infection we used HCV lentiviral pseudotypes (HCVpp) and demonstrated a defect in cell entry. Using a panel of monoclonal antibodies targeting the E2 glycoprotein, we demonstrate that IFN $\alpha$  alters glycoprotein conformation and receptor utilization. **Conclusion:** These observations show a previously unreported and rapid effect of IFN $\alpha$  on HCV particle infectivity that inhibits *de novo* infection events. Evasion of this response may be a contributing factor in whether a patient achieves early or rapid virological response, a key indicator of progression to sustained virological response or clearance of viral infection. (HEPATOLOGY 2014;60:1891-1901)**

**H**epatitis C virus (HCV) infection is a major cause of chronic liver disease that affects 170 million people worldwide, leading to cirrhosis and hepatocellular carcinoma. For many years the only available therapy was pegylated type I interferon alpha (IFN $\alpha$ ) and ribavirin that cured less than 50% of cases in a genotype-dependent manner, prompting the development of direct-acting antiviral agents (DAAs) targeting the viral replicase.<sup>1</sup> The approval of HCV protease and polymerase inhibitors and combination treatment with IFN has led to a significant increase in patient response rates, providing a new standard of care.<sup>2</sup>

Type I IFN stimulates the expression of a number of interferon-stimulated genes (ISGs) that inhibit HCV

genome translation and replication<sup>3,4</sup> and protect neighboring uninfected cells from viral infection. Once IFN pathways are activated the downstream response has enormous breadth, as noted by the many hundreds of ISGs. In a recent screen 380 ISGs were tested for their ability to inhibit genotype 2a HCV replication and at least 25 genes were identified that reduced viral replication.<sup>5</sup> Zhao et al.<sup>6</sup> reported that silencing 60 ISGs prior to IFN treatment revealed no individual gene knockdown that could rescue the inhibitory effect of IFN on HCV replication. This finding is consistent with the widely accepted hypothesis that the IFN system works in a combinatorial fashion, with multiple ISGs contributing to antiviral responses with no single ISG serving as a “magic bullet.”

Abbreviations: DAA, direct-acting antiviral; HCV, hepatitis C virus; HCVcc, HCV cell-culture virus; HCVpp, HCV pseudoparticle; IFN $\alpha$ , interferon-alpha; ISG, interferon-stimulated gene; mAbs, monoclonal antibodies.

From the <sup>1</sup>Viral Hepatitis Research Group, Centre for Human Virology, University of Birmingham, Birmingham, UK; <sup>2</sup>School of Molecular Medical Sciences and the Nottingham Digestive Diseases Centre Biomedical Research Unit, University of Nottingham, Queen's Medical Centre, Nottingham, UK; <sup>3</sup>NIHR Liver Biomedical Research Unit, University of Birmingham, Birmingham, UK.

Received April 17, 2014; accepted July 24, 2014.

Supported by the MRC (G0801976) and by EU “HEPACUTE” (FP7) programme.

Mathematical modeling of HCV replication kinetics following treatment has provided valuable insights into the mechanism of action of antiviral agents.<sup>7,8</sup> HCV RNA declines in a biphasic pattern after IFN therapy, with the initial phase representing a decrease in viral production from infected cells, while the slower and more variable phase is attributed to the loss of infected cells by way of the stimulated immune system. Importantly, there are multiple outcomes observed following IFN therapy, including rapid, partial, and nonresponse, where a rapid virological response is frequently associated with a cure even more than the commonly used IL-28B polymorphisms.<sup>9</sup> Recent modeling studies suggest that for maximal efficacy IFN needs to inhibit infectious virus production and *de novo* transmission events.<sup>10,11</sup> However, the majority of reports studying the impact of IFN on HCV replication have not analyzed these steps in the viral lifecycle.

Reports of IFN-mediated signaling generally study cellular responses within minutes of receptor engagement. However, studies investigating the antiviral effects of IFN measure HCV RNA and proteins several days posttreatment. We demonstrate that treating HCV-infected Huh-7 cells with IFN $\alpha$  for 1 hour reduces the infectivity of extracellular particles with no detectable effect on particle secretion. We failed to detect soluble antiviral effectors released from IFN-treated cells and purified particles show reduced infectivity, demonstrating a direct effect of IFN $\alpha$  on extracellular HCV particles. Studies with lentiviral HCV pseudoparticles (HCVpp) demonstrate that IFN alters E2 glycoprotein conformation that limits receptor-dependent internalization. This is the first report of such a rapid and direct effect of IFN $\alpha$  to reduce particle specific infectivity in the absence of any detectable impact on viral RNA load, raising questions on the interpretation of early phase IFN responses that are solely based on viral RNA measurements.

## Materials and Methods

**Cell Lines and Antibodies.** Huh-7 or Huh-7.5 cells were propagated in Dulbecco's modified Eagle's medium (DMEM) / 10% fetal bovine serum and 1%

nonessential amino acids. Anti-CD81 2s131 and anti-SR-BI (gift from Pfizer) were previously reported.<sup>30</sup> A panel of rodent<sup>31</sup> and human monoclonal antibodies (mAbs) mapping to linear or conformation-dependent HCV E2 epitopes were previously reported.<sup>32,33</sup> Immunoglobulin was isolated from the sera of normal or chronically infected HCV patients by protein G-conjugated Sepharose beads as previously reported.<sup>12</sup>

**ISG Analysis.** HCV-infected cells were treated with IFN $\alpha$  and cellular RNA extracted using RNAeasy Mini-prep Kits (Qiagen). ISG expression was quantified using the Human Interferon Response PCR Array and analyzed using RT<sup>2</sup> Profiler Analysis software (SABiosciences).

**Sodium Dodecyl Sulfate-Polyacrylamide Gel Electrophoresis (SDS-PAGE) and Western Blotting.** Cells were harvested in lysis buffer containing protease inhibitors, clarified by centrifugation (20,000g, 10 minutes), and protein concentration quantified by BCA (Pierce). Normalized protein lysates or virus pellets resuspended in Laemmli buffer were separated by SDS-PAGE, transferred to PVDF membranes, and probed for HCV-E2 (mAb 3/11) or HCV-Core (mAb C750, Thermo Scientific) and expression quantified by enhanced chemiluminescence and densitometric analysis using ImageJ software.

**HCVcc Infectivity, Density Analysis, and Transmission Assay.** Plasmids encoding HCV SA13/JFH, J6/JFH, or H77/JFH were used to generate RNA and electroporated into Huh-7.5 cells.<sup>12</sup> IFN $\alpha$  was added to cells for defined time periods followed by thorough washing and harvesting of extracellular media. Infected cells were fixed with paraformaldehyde, stained for viral antigen expression with mAbs specific for NS5A (9E10), Core (C750), or E2 glycoprotein (9/75, 3/11, 6/53) and isotype-matched Alexa-488 conjugated IgG. Viral antigen expressing cells were enumerated by fluorescent-activated cell sorter (FACS) and protein expression assessed by measuring mean fluorescent intensity (MFI). Extracellular media was collected, concentrated through 100,000 MWCO concentrators (Sartorius), loaded onto 10-60% Iodixanol gradients, and centrifuged at 100,000g for 16 hours at 4°C. Fractions were collected and density assessed by weight, infectivity by inoculating naïve Huh-7.5 cells,

Address reprint requests to: Jane A. McKeating, Ph.D., NIHR Liver Biomedical Research Unit, University of Birmingham, Birmingham, UK. E-mail: j.a.mckeating@bham.ac.uk; fax: +44 121 414 3599.

Copyright © 2014 by the Authors. HEPATOLOGY published by Wiley Periodicals, Inc., on behalf of the American Association for the Study of Liver Diseases. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

View this article online at [wileyonlinelibrary.com](http://wileyonlinelibrary.com).

DOI 10.1002/hep.27333

Potential conflict of interest: Nothing to report.

and HCV RNA by quantitative reverse-transcription polymerase chain reaction (qRT-PCR).

HCV-infected cells were labeled with fluorescent cell-tracker CMFDA and cocultured with an equal number of naïve Huh-7 cells at  $2.5 \times 10^4$  cells/cm<sup>2</sup>. Extracellular virus is neutralized with anti-HCV Ig or control Ig (150 µg/mL) and transmission stopped at indicated times by anti-CD81 (2s131, 10 µg/mL).<sup>12</sup> Cells were harvested and culture media analyzed for infectious virus, expressed as the number of focus-forming units (FFU)/mL. Newly infected target cells were identified by staining for HCV NS5A and quantified by flow cytometry.

**HCVpp Infectivity, Glycoprotein Conformation, and Neutralization.** Luciferase reporter pseudoparticles expressing HCV envelope glycoproteins (HCVpp), or no-glycoprotein controls, were generated in 293T cells as previously reported.<sup>31</sup> HCVpp-293T producer cells were treated with IFN $\alpha$  for 1 hour, unbound IFN removed, and the cells incubated for 1 hour prior to collecting extracellular media to measure infectious HCVpp and staining cells for E2 expression and quantification by flow cytometry. All anti-E2 mAbs were used at subsaturating concentrations. HCVpp stocks were incubated with anti-E2 mAbs or polyclonal anti-HCV Ig for 1 hour prior to infecting Huh-7.5 cells. Receptor neutralization was performed by adding anti-SR-BI, anti-CD81, or isotype control antibody to Huh-7.5 cells for 1 hour prior to infection.

HCV E2 expression in p24 normalized HCVpp containing supernatants was determined by enzyme-linked immunosorbent assay (ELISA). Briefly, Immulon plates (Thermo) were coated with GNA (Sigma, UK) and emipigen-lysed HCVpp preparations added for 2 hours at room temperature. Bound E2 was detected with a panel of anti-E2 antibodies and HRP-conjugated anti-rat Ig. For HIV-1 p24 quantification, MaxiSorp plates (Nalge Nunc) coated with anti-p24 antibody D7324 were used to capture HCVpp (0.5% Tween-20 / 0.1% casein / 1.44M NaCl / 250 mM Tris-Cl, pH 7.5) and bound antigen detected with biotinylated rabbit anti-p24 and streptavidin-HRP (BD Pharmingen). Bound HRP-conjugates were quantified colorimetrically after reaction with a TMB substrate (BioFX).


## Results

**IFN $\alpha$  Rapidly Induces an Antiviral State.** While the clinical impact of IFN therapy on HCV replication kinetics has been extensively studied, our understanding of early IFN-signaling events in the virus lifecycle is limited. Using a commercial IFN-selected PCR array,

we confirmed a significant induction of multiple ISGs within 1 hour of treating HCV-infected Huh-7 cells with IFN $\alpha$  (Fig. 1A). To assess whether this early response has an impact on virus replication, infected cells were treated with varying doses of IFN $\alpha$  for 1 hour and virus infectivity measured. The treated cells were washed extensively to remove exogenous IFN $\alpha$  prior to harvesting samples to ensure minimal carryover when measuring viral infectivity. We noted a dose-dependent decrease in extracellular infectivity with no observable effect on the intracellular pool of infectious virus (Fig. 1B). Importantly, the infectivity of extracellular virus harvested from IFN-treated cells was not restored by pretreating Huh-7 cells with Jak/Stat inhibitor (InSolution JAK Inhibitor 1) (Fig. 1B). Extracellular media from IFN-treated Huh-7 cells had no detectable antiviral activity, suggesting a lack of soluble antiviral factors. To investigate whether IFN affects particle infectivity, we purified the extracellular virus by ultracentrifugation and demonstrated a significant reduction in the infectivity of particles (Fig. 1C). In parallel, the supernatant from ultracentrifuged IFN-treated samples had no effect on HCV infection of naïve Huh-7.5 cells (Fig. 1D), demonstrating that IFN reduces HCV particle infectivity. Finally, we demonstrated that IFN reduces the level of extracellular infectious virus of chimeric strains expressing genotype 1 (H77), genotype 2 (J6), or genotype 5 (SA13) structural proteins (Fig. 1E).

The observation that IFN $\alpha$  reduces extracellular HCV infectivity could be explained by the reduced secretion of particles and/or a loss in particle infectivity. To determine whether IFN $\alpha$  affects HCV particle secretion we measured viral-encoded core, envelope glycoprotein E2, and genomic RNA levels in infected cells and in purified extracellular particles. This short-term IFN $\alpha$  treatment had no detectable effect on intracellular or extracellular particulate viral protein (Fig. 2A) or RNA expression (Fig. 2B), suggesting that IFN $\alpha$  has minimal effects on virus secretion. HCV assembly is dependent on the VLDL pathway and extracellular lipoviral particle infectivity has been reported to associate with particle density. IFN $\alpha$  had minimal impact on HCV particle density, with untreated particles showing a peak of infectious virus at a density of 1.09 g/L compared to 1.08 g/L for IFN $\alpha$ -treated cells and no significant change in viral RNA levels (Fig. 2C). In summary, these results show that IFN $\alpha$  reduces the infectivity of extracellular HCV particle while having minimal impact on secretion or particle density.

**IFN $\alpha$  Rapidly Prevents De Novo HCV Infection.** To investigate the impact of this rapid IFN-mediated response on infectious particle genesis on viral








Fig. 2. IFN $\alpha$  does not perturb HCV particle composition or density. HCV (SA13/JFH)-infected Huh-7.5 cells were treated with IFN $\alpha$  (1,000 IU/mL) for 1 hour, washed thoroughly, and the extracellular media collected for a further 1 hour. HCV Core and E2 expression in extracellular purified virus and cell lysates was assessed by western blot and densitometric imaging (expressed in arbitrary light units, ALU) (A) while HCV RNA was quantified by qRT-PCR (B). The lipid density of the treated and untreated virus was assessed using 10-60% iodixanol gradient centrifugation: the graphs depict the infectivity, HCV RNA, and density of each fraction from IFN-treated or untreated cells (C).




Fig. 3. IFN $\alpha$  inhibits cell-free and cell-to-cell HCV transmission. HCV (SA13/JFH)-infected Huh-7.5 cells were treated with IFN $\alpha$  (1,000 IU/mL) for 1 hour, washed thoroughly to remove IFN $\alpha$ , labeled with CMFDA, and cocultured with naïve Huh-7.5 cells. One hour post-coculture anti-HCV Ig was added to neutralize extracellular virus and at indicated timepoints anti-CD81 was added to block all subsequent transmission events. Twenty-four hours post-coculture the levels of the infectious virus in the extracellular media were measured (A) and *de novo* infection events determined by flow cytometry (B). The results represent the mean and standard deviation of three experiments, and significance calculated using Student *t* tests (\* $P$  < 0.05, \*\* $P$  < 0.01, \*\*\* $P$  < 0.005) comparing treated sample to untreated controls. To assess whether IFN $\alpha$  induced any viral protein degradation during this time period, CMFDA-labeled infected cells were stained for HCV Core or NS5A (C) and the data presented as mean fluorescent intensity (MFI).

dissemination, we used a recently developed single-cycle infection assay that quantifies *de novo* infection events.<sup>12</sup> This assay uses the neutralizing capacity of anti-HCV Ig to limit extracellular virus infectivity to discriminate between cell-free and cell-to-cell infection events. Thus, we were able to study the temporal effect(s) of IFN $\alpha$  on HCV-infected cells and to discriminate from bystander effects on naïve cells. HCV SA13/JFH infected Huh-7.5 cells were treated for 1 hour with IFN $\alpha$  (1,000 IU/mL), washed thoroughly to remove exogenous IFN, and labeled with CMFDA prior to coculturing with naïve Huh-7.5 cells in the presence or absence of neutralizing anti-HCV Ig. At indicated timepoints, viral dissemination was stopped by adding an antibody targeting the CD81 receptor (2s131, 10  $\mu$ g/mL), then incubated for a further 24 hours prior to harvesting cells for flow cytometric detection of newly infected tar-

get cells (NS5A<sup>+</sup>/CMFDA<sup>+</sup>) and collection of extracellular media to measure virus infectivity.

As expected, we noted significant reductions in the level of extracellular infectious virus in the IFN-treated samples at all timepoints (Fig. 3A). IFN $\alpha$  treatment of infected cells for 1 hour significantly reduced cell-free and cell-to-cell dissemination routes within the first 2 hours of coculture and this effect was maintained for the duration of the assay (Fig. 3B). We noted a minimal effect of IFN $\alpha$  on the expression of HCV Core or NS5A in infected cells within the first 8 hours following treatment (Fig. 3C). These observations highlight a new role for IFN to limit the genesis of infectious HCV particles and inhibit *de novo* transmission events.

**IFN $\alpha$  Perturbs HCV Glycoprotein-Dependent Particle Infectivity.** To ascertain whether HCV particles released from IFN $\alpha$ -treated cells are defective at




Fig. 4. IFN $\alpha$ -treated HCVpp show defective entry and increased sensitivity to anti-SR-BI neutralization. 293T cells expressing pseudoparticles containing the indicated viral envelopes were treated with IFN $\alpha$  (1,000 IU/mL) for 1 hour, washed thoroughly, incubated for 1 hour, and the extracellular media collected and used to inoculate naïve Huh-7.5 cells (A). The levels of HCVpp-H77 associated E2 were measured on p24-normalized particle preparations by ELISA (B). The effect of IFN $\alpha$  (1,000 IU/mL, 1 hour) on the infectivity of HCVpp expressing a range of genotype 1 glycoproteins (C). HCVpp-H77 were allowed to infect Huh-7.5 cells preincubated for 1 hour with anti-SR-BI (D) or anti-CD81 (E) and infection assessed 72 hours later by measuring luciferase activity. Results are the mean and standard deviation of three assays with duplicate samples where the significance of differences between sensitivities to neutralization at each timepoint was assessed using paired *t* tests (\**P* < 0.05, \*\**P* < 0.01, \*\*\**P* < 0.005). IFN $\alpha$  had comparable effects on the sensitivity of HCVcc (H77/JFH) to neutralization by anti-SR-BI. Data are presented as IC<sub>50</sub> values (F).

entering target cells, we used lentiviral pseudoviruses expressing HCV glycoproteins to measure the impact of IFN $\alpha$  on receptor-dependent particle entry. HCVpp (strain H77)-expressing cells were treated with IFN $\alpha$  (1,000 IU/mL) for 1 hour, washed thoroughly, and extracellular media collected to quantify viral infectivity. As controls, we evaluated the effect of IFN $\alpha$  on the infectivity of pseudoviruses expressing Lassa virus

and Murine Leukemia virus (MLV) envelope glycoproteins. IFN $\alpha$  reduced the infectivity of HCVpp by 60% but had minimal effect on Lassa or MLVpp infectivity (Fig. 4A). Importantly, IFN $\alpha$  had no detectable effect on the level of HCVpp-incorporated E2 glycoprotein (Fig. 4B), suggesting a similar phenotype to that observed for HCVcc, where intact pseudoparticles are released but are unable to enter cells. To assess the






Fig. 5. IFN $\alpha$  perturbs HCV glycoprotein conformation. HCVpp-H77 producer 293T cells were treated with IFN $\alpha$  (1,000 IU/mL) for 1 hour, washed thoroughly, incubated for a further hour, and the reactivity of a panel of mAbs targeting linear E1E2 epitopes or a control polyclonal anti-HCV Ig to cell surface expressed antigen or extracellular HCVpp measured by flow cytometry or ELISA, respectively (A). A panel of human antibodies targeting conformation-dependent E2 epitopes (1  $\mu$ g/mL) were tested for their ability to bind HCVpp-H77-expressing 293T cells or to neutralize extracellular virus following IFN $\alpha$  (1,000 IU/mL, 1 hour) treatment (B). Human antibody anti-E2 (1  $\mu$ g/mL) neutralization of HCVcc (H77-JFH) released from IFN $\alpha$  (1,000 IU/mL, 1 hour) treated or untreated cells (C). Results are the mean and standard deviation of three experiments and significance was calculated using paired *t* tests (\**P* < 0.05, \*\**P* < 0.01, \*\*\**P* < 0.005).

breadth of this phenotype we assessed the impact of IFN $\alpha$  treatment on the infectivity of pseudoparticles expressing a panel of genotype 1 HCV glycoproteins. Of note, five of the 10 envelopes showed a significant reduction in particle infectivity, whereas IFN $\alpha$  had no effect on the infectivity of the remaining pseudoparticles (Fig. 4C). We next investigated the receptor usage of HCVpp released from IFN $\alpha$ -treated or untreated cells. HCVpp-H77 released from IFN $\alpha$ -treated cells showed an increased sensitivity to neutralization by anti-SR-BI, showing an median inhibitory concentration (IC<sub>50</sub>) of 0.06  $\mu$ g/mL compared to 0.9  $\mu$ g/mL for untreated virus, but no change to anti-CD81 (Fig. 4D,E), suggesting that virus released from IFN-treated cells had a reduced dependency on SR-BI for entry. We observed a similar reduction in the sensitivity of HCVcc (H77/JFH) released from IFN-treated cells to neutralization by anti-SR-BI (Fig. 4F).

**IFN $\alpha$  Alters HCV E2 Glycoprotein Conformation.** Since SR-BI has been reported to bind antigenic region A<sup>13</sup> of E2 glycoprotein, including the hypervariable region 1 (HVR1), these results suggest that IFN may promote conformational changes within the glycoprotein. To test this hypothesis we evaluated the effect of IFN $\alpha$  on the reactivity of a panel of antibodies targeting linear and conformation-dependent E2 epitopes. Anti-E2 antibodies were assessed for their ability to bind IFN $\alpha$ -treated or untreated HCVpp-producer cells by flow cytometry or to bind extracellular HCVpp by ELISA. IFN $\alpha$  reduced the binding of antibodies targeting HVR1 and region A (6/16, 6/82A, 9/86A, 2/69A), region B (7/16B, 6/1A), but had no significant effect on those binding region C (Fig. 5A). Further studies with a number of human anti-E2 antibodies showed that IFN $\alpha$  reduced the binding and neutralizing activity of antibody AR2A that recognizes a conformation-dependent epitope for HCVpp (Fig. 5B). We confirmed that IFN reduced the sensitivity of HCVcc (H77/JFH) to neutralization by AR2A and had modest effects on AR1A and L1 neutralization (Fig. 5C). Given the low levels of glycoprotein expression at the surface of HCVcc-infected cells, we were unable to evaluate antibody binding to the infected cell surface. The results confirm that IFN $\alpha$  induces a conformational change in the viral E2 glycoprotein that limits particle entry.

## Discussion

We show that IFN $\alpha$  induces a rapid response in HCV-infected hepatoma cells that limits the genesis of infectious extracellular particles with minimal effect on

the intracellular viral burden. This short 2-hour duration of IFN treatment significantly reduced *de novo* transmission events, including cell-free and cell-cell dissemination. There was limited evidence for IFN-dependent cellular retention of infectious particles, consistent with a previous report that tetherin, an interferon-stimulated protein, has minimal impact on HCV infectivity.<sup>14</sup> Previous studies reported that IFN $\alpha$  reduced viral RNA and protein expression in HCV-infected Huh-7 cells or those supporting HCV replicons within 12-16 hours.<sup>4,15</sup> However, our studies focus on the first 2 hours following IFN treatment, highlighting a new role for IFN to limit particle infectivity prior to reducing viral RNA and protein expression. Given the difficulties in measuring the infectivity of primary HCV strains, clinical studies measuring IFN efficacy rely on genomic RNA measurements and may underscore the impact of early IFN doses. A recent model of HCV replication kinetics that measured intracellular RNA and infectivity concluded that the NS5A inhibitor daclatasvir has two modes of action: to block viral RNA synthesis and virion assembly/secretion.<sup>16</sup> The authors commented that analysis of HCV kinetic data following IFN $\alpha$  treatment with this new model suggested that IFN $\alpha$  reduces HCV assembly/secretion in addition to its role in slowing intracellular HCV RNA replication. Our data are consistent with this interpretation and highlights the importance of studying particle infectivity, i.e., the ratio of particle infectivity to genomic RNA, allowing one to discriminate the effect of drugs on virus assembly, secretion, and infectivity.

Our recently published single-cycle infection assay<sup>12</sup> enabled us to assess the “early” effects of IFN $\alpha$  on HCV dissemination. By using a short treatment time targeted specifically at infected cells, we demonstrate that IFN $\alpha$  inhibits both cell-free and cell-cell dissemination routes. We confirmed that this effect on particle infectivity and transmission occurred significantly earlier than degradation of viral proteins and is not explained by secreted antiviral factors. Since HCV is a lipid-rich particle that depends on the very low density lipoprotein (VLDL) secretion machinery for assembly and secretion,<sup>17,18</sup> we assessed the density of viral particles released from IFN $\alpha$ -treated cells and saw no obvious changes in particle density. IFN $\alpha$  treatment had no significant impact on the level of viral-associated core or E2 glycoprotein, demonstrating that intact lipoviral particles with reduced infectivity are secreted from IFN-treated cells.

The HCVpp system measures glycoprotein-dependent entry and enabled us to address the role of

IFN $\alpha$  on the virus internalization process. IFN treatment of HCVpp 293T producer cells reduced the infectivity of extracellular virus with minimal effect on E2 incorporation into particles. Previous reports have shown that IFN $\alpha$  inhibits HIV-1 particle infectivity by reducing gp120 incorporation into the nascent particles<sup>19</sup>; however, this is unlikely to explain the effect of IFN $\alpha$  on HCVpp infectivity. IFN $\alpha$  had no detectable effect on the infectivity of pseudoviruses expressing MLV or Lassa envelopes, suggesting that this effect is specific to HCV. It is interesting to note the varying effect of IFN $\alpha$  on the infectivity of pseudoparticles expressing a panel of genotype 1 patient-derived clones. HCVpp and HCVcc released from IFN $\alpha$ -treated cells showed an increased sensitivity to the neutralizing activity of an antibody targeting the cellular receptor SR-BI but not CD81, suggesting that IFN $\alpha$  modulates E2 glycoprotein conformation and this may affect receptor utilization and particle entry. Previous studies have shown that region A-HVR of the viral E2 glycoprotein binds SR-BI<sup>20-22</sup>; we therefore used a panel of mAbs targeting linear and conformational E2 epitopes to study the effect of IFN $\alpha$  on glycoprotein conformation. We noted significant differences in the ability of several mAbs recognizing linear and a conformation-dependent epitope (AR2A) to bind viral glycoproteins or to neutralize virus infectivity following IFN $\alpha$  treatment.

This phenotype is comparable to reports showing that treatment of HCV-infected cells with glycosidase inhibitors led to the incorporation of misfolded glycoproteins into nascent particles.<sup>23,24</sup> However, glycosidase inhibitors reduced the infectivity of intracellular and extracellular HCV, while IFN only reduced extracellular virus infectivity, suggesting that HCV glycoproteins undergo a late maturation step that is required for maximal particle infectivity.<sup>25,26</sup> Maturation of HIV-1 glycoproteins has been shown to “mask” neutralizing epitopes behind a glycan shield,<sup>27,28</sup> and recent studies suggest a similar model for HCV. Glycosylation plays an important role in defining E2 conformation, with up to eight glycan sites reported to be critical for protein folding.<sup>29</sup> We failed to detect any significant change in the mobility of virion associated E2 from IFN-treated virus by SDS-PAGE or sensitivity to glycosidase inhibitors (data not shown), suggesting that IFN $\alpha$ -mediated effects on HCV E2 conformation are not glycan-specific.

The results shown here have a particular significance as therapies move to IFN-free regimens of DAAs such as protease and polymerase inhibitors. The ability of

IFN $\alpha$  to prevent cell-free and cell-to-cell spread may contribute to the efficacy of IFN in responsive patients. Entry inhibitors have long been suggested as useful adjunct therapies to current treatments, as they would significantly impair virus dissemination and aid the clearance of infected cells. It is possible that short treatments with IFN $\alpha$  may continue to play a significant role in patients treated with DAAs, by preventing viral dissemination.

**Acknowledgment:** We thank Mansun Law for anti-E2 mAbs AR1A and AR2A; Mats Persson for anti-E2 mAbs 1:7, A8 and L1; Charlie Rice for Huh-7.5 cells and anti-NS5 9E10; Takaji Wakita and Jens Bukh for HCV JFH-1 and SA13/JFH infectious molecular clones. We also thank Peter Balfe and Samantha Lissauer for critical reading of the article.

## References

1. Scheel TK, Rice CM. Understanding the hepatitis C virus life cycle paves the way for highly effective therapies. *Nat Med* 2013;19:837-849.
2. Pawlotsky JM. New hepatitis C therapies: the toolbox, strategies, and challenges. *Gastroenterology* 2014;146:1176-1192.
3. Wang C, Pflugheber J, Sumpter R Jr, Sodora DL, Hui D, Sen GC, et al. Alpha interferon induces distinct translational control programs to suppress hepatitis C virus RNA replication. *J Virol* 2003;77:3898-3912.
4. Dahari H, Sainz B Jr, Perelson AS, Uprichard SL. Modeling subgenomic hepatitis C virus RNA kinetics during treatment with alpha interferon. *J Virol* 2009;83:6383-6390.
5. Schoggins JW, Wilson SJ, Panis M, Murphy MY, Jones CT, Bieniasz P, et al. A diverse range of gene products are effectors of the type I interferon antiviral response. *Nature* 2011;472:481-485.
6. Zhao H, Lin W, Kumthip K, Cheng D, Fusco DN, Hofmann O, et al. A functional genomic screen reveals novel host genes that mediate interferon-alpha's effects against hepatitis C virus. *J Hepatol* 2012;56:326-333.
7. Neumann AU, Lam NP, Dahari H, Gretch DR, Wiley TE, Layden TJ, et al. Hepatitis C viral dynamics in vivo and the antiviral efficacy of interferon-alpha therapy. *Science* 1998;282:103-107.
8. Perelson AS, Herrmann E, Micol F, Zeuzem S. New kinetic models for hepatitis C virus. *HEPATOLOGY* 2005;42:749-754.
9. Fried MW, Hadziyannis SJ, Shiffman ML, Messinger D, Zeuzem S. Rapid virological response is the most important predictor of sustained virological response across genotypes in patients with chronic hepatitis C virus infection. *J Hepatol* 2011;55:69-75.
10. Dahari H, Shudo E, Ribeiro RM, Perelson AS. Mathematical modeling of HCV infection and treatment. *Methods Mol Biol* 2009;510:439-453.
11. Dahari H, Perelson AS. Hepatitis C virus RNA kinetics: drug efficacy and the rate of HCV-infected cells loss. *World J Gastroenterol* 2007;13:3020-3021.
12. Meredith LW, Harris HJ, Wilson GK, Fletcher NF, Balfe P, McKeating JA. Early infection events highlight the limited transmissibility of hepatitis C virus in vitro. *J Hepatol* 2013;58:1074-1080.
13. Scarselli E, Ansuini H, Cerino R, Roccasecca RM, Acali S, Filocamo G, et al. The human scavenger receptor class B type I is a novel candidate receptor for the hepatitis C virus. *EMBO J* 2002;21:5017-5025.
14. Ye L, Wang X, Li J, Liu J, Ramirez SH, Wu J, et al. Tetherin has negligible activity in restricting hepatitis C virus in hepatocytes. *Innate Immun* 2012;18:398-405.

15. Guo JT, Bichko VV, Seeger C. Effect of alpha interferon on the hepatitis C virus replicon. *J Virol* 2001;75:8516-8523.
16. Guedj J, Dahari H, Rong L, Sansone ND, Nettles RE, Cotler SJ, et al. Modeling shows that the NS5A inhibitor daclatasvir has two modes of action and yields a shorter estimate of the hepatitis C virus half-life. *Proc Natl Acad Sci U S A* 2013;110:3991-3996.
17. Syed GH, Amako Y, Siddiqui A. Hepatitis C virus hijacks host lipid metabolism. *Trends Endocrinol Metab* 2010;21:33-40.
18. Targett-Adams P, Boulant S, Douglas MW, McLauchlan J. Lipid metabolism and HCV infection. *Viruses* 2010;2:1195-1217.
19. Gendelman HE, Baca LM, Kubrak CA, Genis P, Burrous S, Friedman RM, et al. Induction of IFN-alpha in peripheral blood mononuclear cells by HIV-infected monocytes. Restricted antiviral activity of the HIV-induced IFN. *J Immunol* 1992;148:422-429.
20. Dao Thi VL, Dreux M, Cosset FL. Scavenger receptor class B type I and the hypervariable region-1 of hepatitis C virus in cell entry and neutralisation. *Expert Rev Mol Med* 2011;13:e13.
21. Callens N, Ciczora Y, Bartosch B, Vu-Dac N, Cosset FL, Pawlotsky JM, et al. Basic residues in hypervariable region 1 of hepatitis C virus envelope glycoprotein e2 contribute to virus entry. *J Virol* 2005;79:15331-15341.
22. Keck ZY, Xia J, Wang Y, Wang W, Krey T, Prentoe J, et al. Human monoclonal antibodies to a novel cluster of conformational epitopes on HCV E2 with resistance to neutralization escape in a genotype 2a isolate. *PLoS Pathog* 2012;8:e1002653.
23. Chapel C, Garcia C, Bartosch B, Roingeard P, Zitzmann N, Cosset FL, et al. Reduction of the infectivity of hepatitis C virus pseudoparticles by incorporation of misfolded glycoproteins induced by glucosidase inhibitors. *J Gen Virol* 2007;88:1133-1143.
24. Chapel C, Samuel E, Compain P, Blu J, Alotte C, Bartosch B, et al. Novel glucosidase inhibitors with anti-HBV and anti-HCV activity. *J Hepatol* 2009;50:S343-S343.
25. McQuade TJ, Tomasselli AG, Liu L, Karacostas V, Moss B, Sawyer TK, et al. A synthetic HIV-1 protease inhibitor with antiviral activity arrests HIV-like particle maturation. *Science* 1990;247:454-456.
26. Joyner AS, Willis JR, Crowe JE, Aiken C. Maturation-induced cloaking of neutralization epitopes on HIV-1 particles. *PLoS pathogens* 2011;7.
27. Teeraputon S, Louisirirojchanakul S, Auewarakul P. N-linked glycosylation in C2 region of HIV-1 envelope reduces sensitivity to neutralizing antibodies. *Viral Immunol* 2005;18:343-353.
28. Quinones-Kochs MI, Buonocore L, Rose JK. Role of N-linked glycans in a human immunodeficiency virus envelope glycoprotein: effects on protein function and the neutralizing antibody response. *J Virol* 2002;76:4199-4211.
29. Goffard A, Callens N, Bartosch B, Wychowski C, Cosset FL, Montpellier C, et al. Role of N-linked glycans in the functions of hepatitis C virus envelope glycoproteins. *J Virol* 2005;79:8400-8409.
30. Brimacombe CL, Grove J, Meredith LW, Hu K, Syder AJ, Flores MV, et al. Neutralizing antibody-resistant hepatitis C virus cell-to-cell transmission. *J Virol* 2011;85:596-605.
31. Hsu M, Zhang J, Flint M, Logvinoff C, Cheng-Mayer C, Rice CM, et al. Hepatitis C virus glycoproteins mediate pH-dependent cell entry of pseudotyped retroviral particles. *Proc Natl Acad Sci U S A* 2003;100:7271-7276.
32. Giang E, Dorner M, Prentoe JC, Dreux M, Evans MJ, Bukh J, et al. Human broadly neutralizing antibodies to the envelope glycoprotein complex of hepatitis C virus. *Proc Natl Acad Sci U S A* 2012;109:6205-6210.
33. Johansson DX, Voisset C, Tarr AW, Aung M, Ball JK, Dubuisson J, et al. Human combinatorial libraries yield rare antibodies that broadly neutralize hepatitis C virus. *Proc Natl Acad Sci U S A* 2007;104:16269-16274.