

Depression and anxiety are associated with hypertension five years later in a cohort of late middle aged men and women

Ginty, Annie; Carroll, Douglas; Roseboom, TJ; Phillips, Anna; De Rooij, SR

DOI:

[10.1038/jhh.2012.18](https://doi.org/10.1038/jhh.2012.18) IF 2.18

License:

None: All rights reserved

Document Version

Early version, also known as pre-print

Citation for published version (Harvard):

Ginty, A, Carroll, D, Roseboom, TJ, Phillips, A & De Rooij, SR 2013, 'Depression and anxiety are associated with hypertension five years later in a cohort of late middle aged men and women', *Journal of Human Hypertension*, vol. 27, pp. 187-190. <https://doi.org/10.1038/jhh.2012.18> IF 2.18

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This is a pre-copy-editing, author-produced PDF of an article accepted for the above publication following peer review. The definitive publisher-authenticated version is available online at the above DOI.

Eligibility checked for repository: December 2014

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Author's pre-print not final published version. Please cite as Ginty, A.T., Carroll, D., Roseboom, T.J., Phillips, A.C., & de Rooij, S.R. (2013). Depression and anxiety are associated with hypertension five years later in a cohort of late middle aged men and women. *Journal of Human Hypertension*, 27, 187-190.
<http://dx.doi.org/10.1038/jhh.2012.18>

Depression and anxiety are associated with hypertension five years later in a cohort of late middle aged men and women.

Running head: Depression, anxiety, and hypertension

The study was performed at the Department of Clinical Epidemiology and Biostatistics, Academic Medical Centre, University of Amsterdam, Amsterdam, the Netherlands

Annie T. Ginty, MSc¹, Douglas Carroll, PhD¹, Tessa J. Roseboom, PhD², Anna C. Phillips, PhD¹, and Susanne R. de Rooij, PhD²

¹School of Sport and Exercise Sciences, University of Birmingham, Birmingham, United Kingdom; ²Department of Clinical Epidemiology and Biostatistics, Academic Medical Centre, University of Amsterdam, Amsterdam, the Netherlands

Address correspondence to: Susanne R. de Rooij, Department of Clinical Epidemiology and Biostatistics, Amsterdam Medical Centre, University of Amsterdam, Meibergdreef 9, PO Box 22660, 1100 DD Amsterdam, The Netherlands. Tel: **31 20 56 65810; Fax: **31 20 69 12683; E-mail: s.r.derooij@amc.uva.nl

Word count: 2755

Author's pre-print not final published version. Please cite as Ginty, A.T., Carroll, D., Roseboom, T.J., Phillips, A.C., & de Rooij, S.R. (2013). Depression and anxiety are associated with hypertension five years later in a cohort of late middle aged men and women. *Journal of Human Hypertension*, 27, 187-190.

<http://dx.doi.org/10.1038/jhh.2012.18>

Tables: 1

Figures: 0

ABSTRACT

Objective: To examine the association between symptoms of depression and anxiety and hypertension status. **Methods:** Participants ($n = 455, 238$ women) were drawn from the Dutch Famine Birth Cohort Study. In 2002-2004, they attended a clinic assessment during which socio-demographics, anthropometrics, and health behaviours were measured. Symptoms of depression and anxiety were measured using the Hospital Anxiety and Depression Scale. In 2008-2009 participants completed a questionnaire which asked, among other things, whether they ever had a physician diagnosing them as suffering from hypertension. **Results:** In separate regression models that initially adjusted for age and then additionally for sex, SES, smoking, alcohol consumption, antidepressive medication (in the case of HADS depression), anxiolytic medication (in the case of HADS anxiety), whether or not participants were exposed to the Dutch famine *in utero*, BMI, and waist:hip ratio, both depression and anxiety were positively associated with hypertension status. Those who met the criterion for possible clinical depression and anxiety were also more likely to be hypertensive, although the association between anxiety caseness and hypertension was not statistically significant in the fully adjusted regression model. **Conclusion:** Symptoms of depression and anxiety were associated with hypertension status assessed five years later, although the mechanisms underlying these associations remain to be determined.

Key words: anxiety, depression, hypertension

Author's pre-print not final published version. Please cite as Ginty, A.T., Carroll, D., Roseboom, T.J., Phillips, A.C., & de Rooij, S.R. (2013). Depression and anxiety are associated with hypertension five years later in a cohort of late middle aged men and women. *Journal of Human Hypertension*, 27, 187-190.

<http://dx.doi.org/10.1038/jhh.2012.18>

BMI = body mass index; **C** = coefficient of contingency; **CI** = confidence intervals; **HADS** =

Hospital Anxiety and Depression Scale; **ISEI** = International Socio-Economic Index; **OR** = odds

ratio; **SES** = socio-economic status

Author's pre-print not final published version. Please cite as Ginty, A.T., Carroll, D., Roseboom, T.J., Phillips, A.C., & de Rooij, S.R. (2013). Depression and anxiety are associated with hypertension five years later in a cohort of late middle aged men and women. *Journal of Human Hypertension*, 27, 187-190.
<http://dx.doi.org/10.1038/jhh.2012.18>

INTRODUCTION

A number of recent studies testify to positive cross-sectional and/or prospective associations between major depressive disorder and hypertension [1–3] and between generalised anxiety disorder and hypertension [1]. However, evidence of an association between questionnaire assessed symptoms of depression and anxiety and hypertension comes from older studies and is not wholly consistent. The National Health and Nutrition Examination Survey [4] and the Framingham study [5] observed that high levels of symptoms of depression and anxiety were associated with an increased risk of developing hypertension, whereas the Coronary Artery Risk Development in Young Adults study [6] found no consistent relationship between symptoms of depression and anxiety and hypertension. Accordingly, we judged it timely to re-visit this issue and data collected as part of the Dutch Famine Birth Cohort Study [7] permitted re-examination of the association between symptoms of depression and anxiety and hypertension status. We hypothesised, on the balance of the previous evidence, that elevated depression and anxiety symptom levels would be associated with an increased risk for hypertension.

METHOD

Participants

Participants were selected from the Dutch Famine Birth Cohort, which comprises men and women who were born in Amsterdam, the Netherlands, between November 1943 and February 1947. The Dutch Famine Birth Cohort Study was designed to investigate the potential consequences of prenatal exposure to famine on health in later life. One can therefore suggest

Author's pre-print not final published version. Please cite as Ginty, A.T., Carroll, D., Roseboom, T.J., Phillips, A.C., & de Rooij, S.R. (2013). Depression and anxiety are associated with hypertension five years later in a cohort of late middle aged men and women. *Journal of Human Hypertension*, 27, 187-190.
<http://dx.doi.org/10.1038/jhh.2012.18>

that population characteristics may hamper generalization of the present study results. However, this is very unlikely as health differences have mainly been found in the group of people exposed to famine in early gestation which only comprises 8% of the total study sample. The selection procedures and subsequent loss to follow up have been described in detail elsewhere [8,9]. In 2002-2004, 740 members of the cohort provided data at a clinic assessment, performed by a trained research nurse. In 2008-2009, 455 of those who had attended the clinic returned a completed questionnaire package, which among other things, asked about diagnosed illness. The study was approved by the local Medical Ethics Committee, carried out in accordance with the Declaration of Helsinki, and with the adequate understanding and written consent of the participants.

Measures

In 2002-2004, socio-demographics, life style, and anthropometrics were assessed. Socio-economic status (SES) was measured using the ISEI (International Socio-Economic Index)-92, which is based on the participant's or their partner's occupation, whichever has the higher status [10]. Measured values in the ISEI-92 scale ranged from 16 (low status, for example a cleaning person) to 87 (high status, for example a lawyer). Participants were asked whether they were current smokers, ex-smokers, or had never smoked. They were also asked how many alcoholic drinks of different types they consumed per week, and an overall units per week measure was derived. Participants had to indicate whether they were taking antidepressant and/or anxiolytic medication. Height was measured twice using a fixed or portable stadiometer and weight twice using Seca and portable Tefal scales. Body mass index (BMI) was computed in kg/m^2 from the

Author's pre-print not final published version. Please cite as Ginty, A.T., Carroll, D., Roseboom, T.J., Phillips, A.C., & de Rooij, S.R. (2013). Depression and anxiety are associated with hypertension five years later in a cohort of late middle aged men and women. *Journal of Human Hypertension*, 27, 187-190.

<http://dx.doi.org/10.1038/jhh.2012.18>

Averages of the two height and weight measurements. Waist and hip circumference was measured in duplicate with a flexible tape measure and an average taken. Waist was measured midway between the costal margin and the iliac crest and hip at the widest point over the buttocks to compute the waist:hip ratio.

Symptoms of depression and anxiety were measured using the Hospital Anxiety and Depression Scale (HADS) [11]. The HADS is a well-recognised assessment instrument that comprises 14 items, seven measuring depression and seven measuring anxiety. The depression subscale emphasises anhedonia and excludes somatic items. Items are scored on a 4-point scale, 0 to 3; the higher the score, the greater the depression and anxiety. The HADS has good concurrent validity [12,13], performs well as a psychiatric screening device [13,14], and boasts acceptable psychometric properties; for example, a Cronbach's α of .90 for the depression items and .93 for the anxiety items has been reported [15] and test-retest reliability coefficients as high as .85 for depression and .84 for anxiety have been found [13]. Possible caseness for both depression and anxiety is defined by scores ≥ 8 .

The 2008-2009 questionnaire asked participants whether or not they had ever received a diagnosis of hypertension from a physician. The mean (SD) temporal lag between this assessment and the measurement of depression and anxiety was 5.5 (0.6) years.

Statistical Analyses

Author's pre-print not final published version. Please cite as Ginty, A.T., Carroll, D., Roseboom, T.J., Phillips, A.C., & de Rooij, S.R. (2013). Depression and anxiety are associated with hypertension five years later in a cohort of late middle aged men and women. *Journal of Human Hypertension*, 27, 187-190.

<http://dx.doi.org/10.1038/jhh.2012.18>

Preliminary analyses of differences between participants with and without a diagnosis of hypertension were conducted using χ^2 and ANOVA. The main analysis was by logistic regression, where depression and anxiety served as the independent variables in separate models with hypertension status as the dependent variable. The initial models adjusted only for age at clinic assessment, whereas subsequent models additionally adjusted for sex, SES, smoking, alcohol consumption, antidepressive medication (in the case of HADS depression), anxiolytic medication (in the case of HADS anxiety), whether or not participants were exposed to the Dutch famine *in utero*, BMI, and waist:hip ratio. These covariates were chosen because they have been implicated in depression, anxiety, and/or hypertension. Depression and anxiety were first tested as continuous variables, i.e. actual HADS scores, and subsequently as categorical variables, i.e. HADS caseness.

RESULTS

The summary characteristics of the cohort by hypertension status are presented in Table I. Two hundred and nineteen (48%) of the participants reported a diagnosis of hypertension. HADS depression and anxiety scores were highly correlated, $r(433) = .68, p < .001$, as were HADS depression and anxiety caseness, $C(\text{coefficient of contingency})(433) = .43, p < .001$. In regression models adjusting only for age, both depression, $OR = 1.13, 95\%CI 1.05 - 1.21, p = .001$, and anxiety scores, $OR = 1.11, 95\%CI 1.05 - 1.18, p = .001$, were associated with future hypertension status; the greater the symptoms of depression and anxiety the greater the likelihood that the particular person would have been diagnosed with hypertension five years later. These associations remained statistically significant in the fully adjusted models: for

Author's pre-print not final published version. Please cite as Ginty, A.T., Carroll, D., Roseboom, T.J., Phillips, A.C., & de Rooij, S.R. (2013). Depression and anxiety are associated with hypertension five years later in a cohort of late middle aged men and women. *Journal of Human Hypertension*, 27, 187-190.

<http://dx.doi.org/10.1038/jhh.2012.18>

symptoms of depression, OR = 1.12, 95%CI 1.04 – 1.21, $p = .004$, and for symptoms of anxiety, OR = 1.09, 95%CI 1.02 – 1.17, $p = .01$. The covariates significantly associated with hypertension in these models were BMI and smoking; smokers, $p = .03$ and $p = .04$ respectively, and those who were leaner, $p < .001$ in both cases, were less likely to be hypertensive. These analyses were repeated using cut-off scores of ≥ 8 to identify possible clinical depression and anxiety. In the age adjusted models, both depression caseness, OR = 2.34, 95%CI 1.14 – 4.79, $p = .02$, and anxiety caseness, OR = 1.73, 95%CI 1.08 – 2.77, $p = .02$, were positively associated with hypertension status. With full adjustment, the association between depression caseness and hypertension remained significant, OR = 2.45, 95%CI 1.02 – 4.95, $p = .04$. However, the association between anxiety caseness and hypertension was attenuated to non-significance, OR = 1.50, 95%CI 0.89 – 2.52, $p = .13$. As before, the outcomes remained unchanged when we additionally adjusted for whether or not participants had been exposed to the Dutch famine. Again, of the covariates entered in these models, only BMI and smoking status were significantly related to hypertension.

DISCUSSION

Almost half the current sample reported that a physician had diagnosed them as hypertensive. This is a higher prevalence than that reported in other studies [1,3], but could reflect the age of the sample who were all over 60 at the time of the questionnaire assessment of hypertension status as well as the close medical attention paid to this cohort. It certainly suggests, though, that there was little undiagnosed hypertension, a concern when relying on self-reports of physician diagnosis [1]. As hypothesised, participants with high levels of depression and anxiety

Author's pre-print not final published version. Please cite as Ginty, A.T., Carroll, D., Roseboom, T.J., Phillips, A.C., & de Rooij, S.R. (2013). Depression and anxiety are associated with hypertension five years later in a cohort of late middle aged men and women. *Journal of Human Hypertension*, 27, 187-190.

<http://dx.doi.org/10.1038/jhh.2012.18>

symptoms were more likely to be hypertensive when questioned five years later. Hypertensive participants were also more likely to meet the criterion for possible clinical caseness at the earlier assessment, although the association between anxiety caseness and hypertension was not statistically significant in the fully adjusted regression model. Overall, then our results are in line with those from other studies comparing hypertension in those varying in symptoms of depression and anxiety [4,5] and those with and without Major Depressive Disorder [1-3] and with and without Generalised Anxiety Disorder [1].

There are several possible pathways through which depression and anxiety might contribute to the development of hypertension. The two most cited are unhealthy behaviours and physiological dysregulation. However, in the present analyses, the associations between symptoms of depression and anxiety and hypertension were still evident after adjustment for two of the more prominent unhealthy behaviours, smoking and high levels of alcohol consumption. Indeed, smoking in the current study was negatively associated with hypertension, a not uncommon finding [1,16]. With regard to physiological dysregulation, altered activity of the hypothalamic-pituitary-adrenal axis has been observed in approximately 50% of depressed patients [17], and this, in turn, may increase the risk of hypertension. However, cortisol was measured in the present study during the clinic assessment and the addition of basal cortisol to our regression models left the outcomes unaltered. One other possibility is reverse causation, such that increased symptoms of depression and anxiety are reactions to a diagnosis of hypertension, a chronic and ultimately life-threatening condition; the majority of the sample (71%) was diagnosed hypertensive prior to the clinic visit. However, HADS depression, 3.60

Author's pre-print not final published version. Please cite as Ginty, A.T., Carroll, D., Roseboom, T.J., Phillips, A.C., & de Rooij, S.R. (2013). Depression and anxiety are associated with hypertension five years later in a cohort of late middle aged men and women. *Journal of Human Hypertension*, 27, 187-190.

<http://dx.doi.org/10.1038/jhh.2012.18>

versus 3.30, $p = .57$, and anxiety, 5.98 versus 5.73, $p = .66$, scores were similar for those who had been diagnosed before the clinic visit and those who had been diagnosed subsequently.

The present study is not without limitations. First, from observational data it is impossible to determine causality and the direction of causality. However, as indicated above, reverse causation would seem to be an unlikely explanation. Nevertheless, the issue of confounding is ever present in observational studies [18]. Although we did adjust statistically for a broad range of potential confounders, including whether or not participants were exposed to the Dutch famine, residual confounding as a consequence of poorly measured or un-measured variables cannot be wholly discounted. Second, we relied on a questionnaire measure of symptoms rather than formal diagnoses of major depressive and anxiety disorders. However, our findings are very much in accord with those from studies that did examine hypertension in the context of psychiatric diagnoses [1-3]. Third, the relative seniority of our participants and the consequently high prevalence of diagnosed hypertension could be regarded as limiting the generalisability of our findings. However, given that similar associations have emerged in middle-aged cohorts [1-3], it would appear that the relationship between depression and anxiety and hypertension transcends the age of the sample studied. Finally, this study relied on self-report of a physician diagnosis of hypertension. However, the high prevalence of hypertension suggests that there were unlikely to be many false negatives. Further, other studies in the field have also used self-report of hypertension [2,3].

Author's pre-print not final published version. Please cite as Ginty, A.T., Carroll, D., Roseboom, T.J., Phillips, A.C., & de Rooij, S.R. (2013). Depression and anxiety are associated with hypertension five years later in a cohort of late middle aged men and women. *Journal of Human Hypertension*, 27, 187-190.
<http://dx.doi.org/10.1038/jhh.2012.18>

In conclusion, the present analyses indicated that symptoms of depression and anxiety were associated with hypertension status assessed five years later. These associations would not seem to be accounted for by unhealthy behaviour, nor would they appear a function of reverse causation, where increased symptoms of depression and anxiety are reactions to a diagnosis of hypertension. Further studies are needed to unravel the mechanisms underlying what appear to be fairly robust associations between psychological status and hypertension.

Author's pre-print not final published version. Please cite as Ginty, A.T., Carroll, D., Roseboom, T.J., Phillips, A.C., & de Rooij, S.R. (2013). Depression and anxiety are associated with hypertension five years later in a cohort of late middle aged men and women. *Journal of Human Hypertension*, 27, 187-190.
<http://dx.doi.org/10.1038/jhh.2012.18>

Acknowledgements

We thank the participants for their willing cooperation. The present study was supported by the Netherlands Heart Foundation (grant number 2007B083) and the European Science Foundation (EUROSTRESS) / Netherlands Organization for Scientific Research. These funding sources had no role in study design, data collection, analysis and interpretation of the data.

Competing Interest Statement

All authors have completed the Unified Competing Interest form. The authors have no competing interests to report.

Author's pre-print not final published version. Please cite as Ginty, A.T., Carroll, D., Roseboom, T.J., Phillips, A.C., & de Rooij, S.R. (2013). Depression and anxiety are associated with hypertension five years later in a cohort of late middle aged men and women. *Journal of Human Hypertension*, 27, 187-190.
<http://dx.doi.org/10.1038/jhh.2012.18>

References

- [1] Carroll D, Phillips AC, Gale CR, Batty D. Generalized anxiety and major depressive disorders, their comorbidity and hypertension in middle-aged men. *Psychosom Med*, 2010; 72: 16-9.
- [2] Patten SB, Beck CA, Kassam A, Williams JVA, Barbui C, Metz LM. Long-term medical conditions and major depression: strength of association in the general population. *Can J Psychiatry* 2005; 50:195-202.
- [3] Patten SB, Williams JVA, Lovorato DH, Campbell NRC, Eliasziw M, Campbell TS. Major depression as a risk factor for high blood pressure: Epidemiologic evidence from a national longitudinal study. *Psychosom Med* 2009; 71: 273-9.
- [4] Jonas BS, Franks P, Ingram DD. Are symptoms of anxiety and depression risk factors for hypertension? Longitudinal evidence from the National Health and Nutrition Examination Survey I Epidemiologic Follow-up Study. *Arch Fam Med* 1997;6:43-9.
- [5] Markovitz JH, Matthews KA, Kannel WB, Cobb JL, D'Agostino RB. Psychological predictors of hypertension in the Framingham Study. Is there tension in hypertension? *JAMA* 1993; 270:2439-43.
- [6] Yan LL, Liu K, Matthews KA, Daviglius ML, Ferguson TF, Kiefe CI. Psychosocial factors and risk of hypertension: the Coronary Artery Risk Development in Young Adults (CARDIA) study. *JAMA* 2003; 290:2138-48.
- [7] Roseboom T, de Rooij S, Painter R. The Dutch famine and its long-term consequences for adult health. *Early Hum Dev* 2006; 82:485-91.
- [8] Ravelli AC, van der Meulen JH, Michels RP, Osmond C, Barker DJ, Hales CN, Bleker OP. Glucose tolerance in adults after prenatal exposure to famine. *Lancet* 1998; 351:173-7.
- [9] Painter RC, Roseboom TJ, Bossuyt PM, Osmond C, Barker DJ, Bleker OP. Adult mortality at age 57 after prenatal exposure to the Dutch famine. *Eur J Epidemiol* 2005; 20:673-6.
- [10] Bakker B, Sieben I. Maten voor prestige, social-economische status en sociale klasse voor de standard beroepenclassificatie 1992. *Sociale Wetenschappen* 1997; 40:1-22.
- [11] Zigmond AS, Snaith RP. The Hospital Anxiety and Depression Scale. *Acta Psychiatr Scand* 1983; 67:361-70.

Author's pre-print not final published version. Please cite as Ginty, A.T., Carroll, D., Roseboom, T.J., Phillips, A.C., & de Rooij, S.R. (2013). Depression and anxiety are associated with hypertension five years later in a cohort of late middle aged men and women. *Journal of Human Hypertension*, 27, 187-190.
<http://dx.doi.org/10.1038/jhh.2012.18>

- [12] Bramley PN, Easton AME, Morley S, Snaith RP. The differentiation of anxiety and depression by rating scales. *Acta Psychiatr Scand* 1988; 77:133-8.
- [13] Herrmann, C. International experiences with the Hospital Anxiety and Depression Scale- a review of validation data and clinical results. *J Psychosom Res* 1997; 42:17-41.
- [14] Bjelland I, Dahl AA, Haug TT, Neckelmann D. The validity of the Hospital Anxiety and Depression Scale: an updated literature review. *J Psychosom Res* 2002; 52:69-77.
- [15] Moorey S, Greer S, Watson M, Gorman C, Rowden L, Tunmore R, Robertson B, Bliss J. The factor structure and factor stability of the Hospital Anxiety and Depression Scale in patients with cancer. *Br J Psychiatry* 1991; 158:255-9.
- [16] Green MS, Jucha E, Luz Y. Blood pressure in smokers and non-smokers: epidemiologic findings. *Am Heart J* 1986; 111:932-40.
- [17] Brown ES, Varghese FP, McEwen BS. Association of depression with medical illness: does cortisol play a role? *Biol Psychiatry* 2004; 55:1-9.
- [18] Christenfeld, NJ, Sloan RP, Carroll D, Greenland S. Risk factors, confounding and the illusion of statistical control. *Psychosom Med* 66, 737-43.

Author's pre-print not final published version. Please cite as Ginty, A.T., Carroll, D., Roseboom, T.J., Phillips, A.C., & de Rooij, S.R. (2013). Depression and anxiety are associated with hypertension five years later in a cohort of late middle aged men and women. *Journal of Human Hypertension*, 27, 187-190. <http://dx.doi.org/10.1038/jhh.2012.18>

Table I. Descriptive characteristics of those with and without a diagnosis of hypertension.

	Hypertensive (n = 219)		Normotensive (n = 236)		P
	Mean	SD	Mean	SD	
Age at clinic assessment (years)	58.16	0.88	58.28	0.94	.16
Socio-economic status (ISEI-92)	49.82	13.88	50.88	13.79	.42
Body mass index (kg/m ²)	30.42	5.04	27.08	3.94	<.001
Waist:hip ratio	93.69	9.00	91.89	8.58	.03
Alcohol consumption (units/week)	9.54	16.53	9.97	12.56	.76
HADS depression score	3.66	3.44	2.64	2.43	<.001
HADS anxiety score	5.99	3.55	4.85	2.97	<.001
	n (%)		n (%)		P
Sex					.23

Author's pre-print not final published version. Please cite as Ginty, A.T., Carroll, D., Roseboom, T.J., Phillips, A.C., & de Rooij, S.R. (2013). Depression and anxiety are associated with hypertension five years later in a cohort of late middle aged men and women. *Journal of Human Hypertension*, 27, 187-190. <http://dx.doi.org/10.1038/jhh.2012.18>

Male	98 (45)	119 (55)	
Female	121 (51)	117 (49)	
Smoking			.13
Current smokers	44 (42)	61 (58)	
Ex-smokers	83 (46)	97 (54)	
Never smoker	91 (54)	78 (46)	
Exposed to famine <i>in utero</i>	90 (41)	99 (42)	.22
Medication			
Antidepressive	14 (6)	12 (5)	.55
Anxiolytic	18 (8)	18 (8)	.90
Caseness			
Depression	25 (12)	12 (5)	.01
Anxiety	54 (26)	37 (16)	.02
