

Ecological consequences of historic moorland 'improvement'

Rowney, Francis; Fyfe, Ralph; Anderson, Philip ; Barnett, Robert; Blake, William ; Daley, Tim ; Head, Katie; MacLeod, Alison; Mathews, Ian ; Smith, David

DOI:

[10.1007/s10531-022-02479-6](https://doi.org/10.1007/s10531-022-02479-6)

License:

Other (please specify with Rights Statement)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Rowney, F, Fyfe, R, Anderson, P, Barnett, R, Blake, W, Daley, T, Head, K, MacLeod, A, Mathews, I & Smith, D 2022, 'Ecological consequences of historic moorland 'improvement'', *Biodiversity and Conservation*.
<https://doi.org/10.1007/s10531-022-02479-6>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

This version of the article has been accepted for publication, after peer review (when applicable) and is subject to Springer Nature's AM terms of use, but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: [http://dx.doi.org/\[insert DOI\]](http://dx.doi.org/[insert DOI])

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Ecological consequences of historic moorland ‘improvement’

Francis M. Rowney^{1*}, Ralph M. Fyfe^{1*}, Philip Anderson¹, Robert Barnett², William Blake¹,
Tim Daley¹, Katie Head¹, Alison MacLeod³, Ian Matthews⁴, David N. Smith⁵

1 School of Geography, Earth and Environmental Sciences, University of Plymouth

2 Department of Geography, University of Exeter

3 Department Geography and Environmental Science, University of Reading

4 Department of Geography, Royal Holloway University of London

5 Department of Classics, Ancient History and Archaeology, University of Birmingham

*corresponding authors

Francis M. Rowney

francis.rowney@plymouth.ac.uk

ORCID: 0000-0002-7986-5351

Ralph M. Fyfe

ralph.fyfe@plymouth.ac.uk

Declarations

Funding: This work was supported by the Moorland Restoration and Improvement Project coordinated by Southwest Water, Exmoor National Park Authority and Natural England, and a Leverhulme Trust grant (RPG-2020-045).

Conflicts of Interest: The authors have no conflicts of interest to declare that are relevant to the content of this article.

Availability of data and materials: Data have been archived on University of Plymouth's PEARL repository.

Code availability: Relevant code to ensure reproducibility has been included in the article.

Author contributions: Francis M. Rowney and Ralph M. Fyfe analysed the data and led the writing of the manuscript; Ralph M. Fyfe, Tim Daley, and David N. Smith collected samples in the field; Ralph M. Fyfe, Francis M. Rowney, Philip Anderson, Robert Barnett, William Blake, Katie Head, Alison Macleod, Ian Matthews and David N. Smith produced the proxy data. All authors contributed to the drafts and gave approval for publication.

Acknowledgements

Lee Bray is thanked for his help in the field and, along with Rob Wilson-North, for discussions of aspects of the results. Katy Flowers (RHUL) is thanked for her assistance with tephra sample processing. We are grateful to the anonymous reviewers for their contributions to improving the paper.

Abstract

Upland peatlands are nationally and internationally important habitats that can provide a range of ecosystem services, but many are considered degraded by human activities. On Exmoor, (South West England, UK) restoration activities are often aimed at reversing the effects of nineteenth century agricultural ‘improvement’ schemes, the effects of which are not yet fully understood. To develop this understanding, long-term ecological context is essential. We used sub-fossil pollen, plant macrofossils, testate amoebae, insects, coprophilous fungal spores and charcoal to study ecological conditions and disturbance regimes over the last ~7700 years at a site in Exmoor National Park (‘Ricksy Ball’). Multivariate analyses were used to explore changes in ecological communities over time and a range of techniques were used to establish the chronology. During the last ~7700 years, anthropogenic disturbance regimes (burning, grazing, drainage) have varied through time, reflecting changing land use and management, the effects of which are evident in vegetation (pollen, plant macrofossils) and microbial (testate amoebae) communities. In particular, a combination of drainage and high-intensity grazing appears to have substantially altered local ecology during the nineteenth century, indicated by increases in coprophilous fungal spores and the loss of *Sphagnum* and associated biota. This occurred in the context of more gradual, centennial-scale declines in *Sphagnum* and microbial biovolumes. We provide a range of reference conditions and show that the moorland has been influenced by land management changes for millennia, and this may have been most pronounced during the nineteenth century. There is no single, readily identifiable, ‘stable’ pre-drainage baseline.

Keywords: Peatlands, Palaeoecology; Pollen; Testate amoebae; Multi-proxy; Restoration

Introduction

Uplands provide nationally and internationally important habitats, particularly in areas with spatially-extensive peatlands (Reed et al. 2009). Much recent attention has been placed on the protection and restoration of upland peatlands, owing to their range of potential ecosystem services, including regulatory services such as climate change mitigation (through greenhouse gas sequestration), provisioning services (*e.g.* water supply) and cultural services (*e.g.* recreation and preservation of heritage assets) (Grand-Clement et al. 2013). Peatland vegetation and ecological function are important in the provision of these services (Belyea and Malmer 2004; Ritson et al. 2016). Many peatlands in the UK and Ireland are considered degraded as a result of anthropogenic activities including burning, peat cutting and drainage (Holden et al. 2007; Bonn et al. 2016). In the UK, drainage has historically been used in moorland ‘agricultural improvement’ schemes, particularly during the nineteenth century following parliamentary enclosure acts (Whyte 2006) and in the post-war period (1940s to late 1970s) in response to national food supply concerns (Dallimer et al. 2009; Lobleby and Winter 2009). Drainage ditches reduce vegetation diversity on blanket peat (Gatis et al. 2016), which may be exacerbated when drainage is used to facilitate increased grazing: long-term data suggest this may have led to the expansion and dominance of purple moor-grass (*Molinia caerulea*) across uplands (Chambers et al. 2007). Future climatic changes may also have significant effects on peatland functioning, particularly in regions such as the southwest of the UK where peatlands lie in climatically-marginal situations (Gallego-Sala et al. 2010). Peatland restoration activities are often aimed at reversing the historical effects of drainage (Bonn et al. 2016), and so understanding these effects, particularly from a long-term perspective, is important. This includes understanding the extent of historical *Sphagnum* losses, associated changes in broader ecological communities (vegetation, microbes, etc), and the relative importance of other disturbance processes.

Exmoor is an upland region with a long history of pre- and post-1940s drainage (Riley and Wilson-North 2001). Landscape-scale drainage and management for agriculture on Exmoor began in the 1820s following the sale of the Royal Forest of Exmoor (*c.* 60 km²) to the Knight family in 1818 (Orwin and Sellick 1970; Riley and Wilson-North 2001; Hegarty and Wilson-North 2014), and further drainage took place during the mid-twentieth century. Restoration work aimed at reversing the effects of drainage on Exmoor (*e.g.* blocking nineteenth and twentieth century drainage ditches to raise local water tables: ‘re-wetting’) has been on-going since 2010, with pre- and post-restoration monitoring to understand responses in hydrological and biological systems, and how these may relate to carbon flux (Grand-Clement et al. 2015; Gatis et al. 2016). The aim of the present research is to understand the ecological consequences of historic moorland ‘improvement’ by providing long-term context, and to assess the extent to which pre-drainage conditions can be used to evaluate the success of on-going restoration works.

This paper presents a detailed multi-proxy palaeoecological study of an area of moorland that is subject to on-going restoration (*e.g.* ditch-blocking) and monitoring on Exmoor, southwest Britain (**Fig. 1**). Palaeoecological datasets offer unparalleled potential for understanding the long-term dynamics of peatland systems (Davies and Bunting 2010; Gearey and Fyfe 2016; Marcisz et al. 2022), and provide valuable insights to restoration projects (Chambers et al. 2007; Blundell and Holden 2015; McCarroll et al. 2016). In this case, palaeoecological study provides the context of historical *Sphagnum* losses, how drainage may have influenced wider ecologies, including the potential formation of novel communities, and the additional importance of grazing and burning. We find that the moorland’s ecology has changed with land use over millennia, and that nineteenth century ecological changes (probably related to both drainage and high-intensity grazing) were particularly notable.

Materials and Methods

Sampling Site

Sampling was undertaken in 2012 in the Aclands catchment on Exmoor (0.179 km²), an area subject to on-going restoration work (since 2014) and monitoring (Grand-Clement et al. 2015; Gatis et al. 2016). A key aim of this is to raise water tables, restoring hydrological function and associated ecosystem services (*e.g.* carbon sequestration, enhanced water quality, flood prevention) (Grand-Clement et al. 2015). Peat depths range from 0.33 to 2 m and there are 7.5 km of drainage ditches arranged in a herring-bone pattern (Grand-Clement et al. 2015) (Fig. 1). These drains direct water in the headwaters of the River Barle across an area of deeper peat ('Ricksey Ball'), approximately 190 m long and up to 80 m wide. A 1.9 m peat sequence was extracted from Ricksey Ball (-3.808° longitude, 51.131° latitude WGS84; elevation: 421 m OD) using monolith tins (open section, upper 1.1 m) and a closed-chamber corer (lower 0.8 m). The sampling site is located within the extensively drained area.

Chronology

Chronological control of the sequence was established using a suite of methods.

Samples of peat 1 cm thick were used for AMS (accelerator mass spectrometry) radiocarbon analysis, with dating undertaken on the humic acid fraction to avoid contamination by penetration of younger carbon from higher levels (Piotrowska et al. 2011).

The core was scanned for tephra horizons at 5 cm resolution, with precise depths established through shard counts from contiguous 1 cm thick samples where necessary. Tephra samples were prepared for chemical analysis following Dugmore, Larsen, & Newton (1995) and Blockley et al. (2005). Further details are given in Supplementary Information.

Contiguous 1 cm samples were taken from the upper part of the section for fallout radionuclide (unsupported ^{210}Pb , ^{137}Cs and ^{241}Am) based sediment dating techniques (Appleby, 2001) at the University of Plymouth Consolidated Radioisotope Facility (CoRiF). Further details are given in Supplementary Information. The Constant Rate of Supply (CRS) model (Appleby and Oldfield 1978; Appleby 2001) was applied to construct core chronologies from activity concentrations.

The analysis of major and trace elements within the upper 1.1 m of the core was undertaken using a portable Niton X-Ray Fluorescence (XRF) elemental analyser. Major pollutants (particularly lead (Pb)) and lithogenic elements (*e.g.* zirconium (Zr) and titanium (Ti)) were measured (as parts per million: ppm). Ratios between pollutants (Pb) and lithogenic elements (Ti) have been successful elsewhere in identifying increased atmospheric deposition associated with known industrial pollution histories (Mighall et al. 2002), under the assumption that increases in ratios indicate excess atmospheric Pb not otherwise visible in the raw inventory.

Based on the results of the above analyses, an age-depth model was constructed for the sequence. Radiocarbon dates were calibrated using the IntCal20 calibration curve (Reimer et al. 2020). Using the R (R Core Team 2021) packages ‘rbacon’ (Blaauw and Christen 2011; Blaauw 2021a) and ‘clam’ (Blaauw 2010, 2021b), a range of age-depth models were tested. Three radiocarbon dates (UBA-24401, UBA-24402, UBA-24403) that gave age estimates younger than underlying tephras were not used in estimating interpolated ages (automated in ‘rbacon’). Whilst radiocarbon dates of peat are generally reliable, there are a number of potential sources of error (Piotrowska et al. 2011; Välimäki et al. 2014). Results were broadly similar and the simplest model was chosen: ages between directly dated depths were estimated by linear interpolation using ‘clam’ (Blaauw 2010, 2021b) and the following code:

162 clam(coredir="clam_runs", core="RicksyBall", cc=1, outliers=c(7,8,10),
163 depthseq=print(depths), type=1, its=10000)

164

165 *Palaeoecology*

166 *Plant macrofossils*

167 Material for plant macrofossil analysis was extracted following Mauquoy, Hughes, & van
168 Geel (2010) methods. Sub-samples of peat (2 cm contiguous intervals) were disaggregated
169 and wet sieved to 125 μm and examined using low-power (x10-50) microscopy. Where
170 epidermal material or *Sphagnum* leaves were present, an additional representative sample (50
171 *Sphagnum* leaves) was identified to sub-genus level using high-power microscopy (x100-
172 400).

173

174 *Testate amoebae*

175 Testate amoebae analysis followed Booth, Lamentowicz, & Charman (2010). Sub-samples of
176 sediment (1 cm^3) were disaggregated in potassium hydroxide, wet-sieved to isolate
177 particulate fractions (>15 to <300 μm), mounted in glycerol and identified using high-power
178 (x400-600) microscopy (using Charman, Hendon, & Woodland (2000)). A known amount of
179 an exotic spike (*Lycopodium* spores) (Stockmarr 1971)) was added to facilitate calculations
180 of test concentrations. One hundred tests were counted per sample, which is demonstrably
181 representative (Payne and Mitchell 2009). Count data were converted to influx (number of
182 individuals $\text{yr}^{-1} \text{cm}^{-3}$) using test concentrations and sediment accumulation rates from the age-
183 depth model. Testate amoeba biovolumes were estimated based on morphometric
184 measurements derived from the literature (see Supplementary Table 1) and test shape
185 ('hemispheric', 'ovoid', 'cylindrical' or 'saucer') according to equations in Mitchell (2004)

and Gilbert, Amblard, Bourdier, & Francez (1998). Total biovolume is the sum for all taxa, which was then scaled by influx rates to give biovolume values per year per cubic centimetre of sediment ($\mu\text{m yr}^{-1} \text{ cm}^{-3}$). Overall testate amoebae biovolume ($\mu\text{m}^3 \text{ yr}^{-1}$) for each sample was estimated by scaling the biovolume information by the influx of each taxon.

Pollen and coprophilous fungal spores

Pollen and coprophilous fungal spore sample preparation followed Moore, Webb, & Collinson (1991). One cm^3 sub-samples were disaggregated in potassium hydroxide, silicates were removed using hydrofluoric acid, cellulose was digested using an acetolysis reaction (acetic anhydride and sulphuric acid), samples were dehydrated with alcohols and mounted in silicon oil for identification using high-power (x400-600) microscopy. Three hundred land pollen grains were identified from each sub-sample, a sufficient number for characterising assemblages (Djamali and Cilleros 2020). Coprophilous fungal spores (per 100 *Lycopodium* spores) were counted on pollen slides to 168.5 cm core depth. *Sporormiella*-type spores may occur as lone cells or chains of cells, and counts were based on the number of chains plus the number of lone cells.

Insects

Contiguous 10 cm samples of ~10 litres were taken from the open peat section, adjacent to the monolith tin samples. Samples were sieved (300 μm) before undergoing paraffin (kerosene) flotation (Kenward et al. 1980). Insect remains were sorted and identified using low-power (x15-45) microscopy. Where achievable, they were identified to species-level by direct comparison to modern specimens (Gorham and Girling insect collections, University of

Birmingham). Identifiable insect remains are mostly from Coleoptera (beetles). Taxonomy follows Lucht (1987). Coleoptera have been assigned to ecological groupings where possible, following a simplified version of Robinson's (Robinson 1981, 1983) scheme. Dung, open habitat and moorland ecological groupings are calculated as percentages of the total number of terrestrial species, as opposed to the entire fauna.

Multivariate analyses

To explore the extent and directions of change in ecological communities over time, and the taxa associated with these changes, unconstrained ordination techniques were used with both testate amoeba (influx rates) and pollen assemblages (relative abundances), representing more local and more regional ecological conditions, respectively (*n.b.* pollen and testate abundances were treated as percentages and influx rates, respectively, in correspondence with Figures 3 and 6, to aid understanding). Datasets were filtered prior to analysis to include only common taxa (occurring in at least 50% of samples) and abundances were square root transformed (to facilitate analyses based on the direction and relative magnitude of changes in abundances). Testate amoebae were insufficiently preserved below 94 cm (~403 cal BP), and so multivariate analyses of testate amoebae and pollen assemblages focus on the upper 94 cm to facilitate comparison.

Detrended correspondence analyses (DCA) were performed on each dataset to determine environmental gradient lengths and all gradient lengths were <2.5 standard deviations. On this basis (following Legendre & Birks (2012)), a linear method (principal components analysis: PCA) was selected to explore trends in the composition of pollen assemblages, but this was determined as inappropriate for testate amoebae assemblages. For these, non-metric multidimensional scaling (NMDS) analysis was used instead. This uses rank-based

dissimilarity between samples, and is suitable for non-Gaussian data (Oksanen et al. 2019).
Analyses were performed using the R package ‘vegan’ (Oksanen et al. 2019).

Results

Chronology

A summary of the age-depth model is given in Fig. 2 and uncalibrated, calibrated and modelled ages for dated depths are given in Supplementary Table 2. This model is used to provide date estimates for key transitions in palaeoecological sequences, and to calculate influx rates. Dates are given in calibrated years before present (cal BP), where 1950 CE is ‘present’, and in some cases, calendar years (BCE/CE) are also given. Further details of chronological results are given in Supplementary Information.

Palaeoecology

Assemblage changes and their implications for local environmental changes are summarised here, and further details are given in Tables 1, 2 and 3, and in Supplementary Information.

Pollen and coprophilous fungal spores

The results of pollen and fungal spore analysis from 49 samples describe vegetation changes in the local and wider area around Ricksey Ball, and grazing activity in the local area (Fig. 3). As a general rule-of-thumb, herbaceous (and wetland) pollen types are likely to derived more local sources (<100 m) than pollen from woody, dryland taxa (Binney et al. 2005; Blaus et al. 2020). The pollen has been visually divided into five local pollen assemblage zones (lpaz), to

aid the description of key periods of continuity and change (see Table 1 for detailed
 descriptions). Mixed open woodland conditions are represented in the lowermost zone (zone
 RIB 1paz1, 190.5-162.5 cm, from ~7703 to 3340 cal BP). At 162.5 cm (~3340 cal BP)
 woodland taxa decline and are replaced by Poaceae and ‘improved’ grassland taxa.
 Coprophilous fungal spores increase in concentration (Fig. 3), suggesting increased herbivore
 density. The start of the third pollen zone at 114.5 cm (~595 cal BP / 1355 CE) is marked by
 declining Poaceae and increases in woody heathland shrubs and some trees. Coprophilous
 fungal spores are also present in this zone (Fig. 3). From 94.5 cm (~410 cal BP / 1540 CE)
 the pollen assemblages are dominated by Poaceae and associated grassland taxa, with
Calluna vulgaris declining to trace levels. In this zone, *Sphagnum* is present in substantial
 quantities for the first time. Coprophilous fungal spores are present at low levels, but with a
 substantial increase in *Sordaria*-type at 8.5 cm (Fig. 3). In the uppermost samples (from 6.5 cm:
 52 cal BP / 1898 CE) Poaceae and indicators of improved grassland decrease, replaced with
 taxa more associated with ‘poorer’ moorland and levels of *Sphagnum* decline to very low
 amounts. Coprophilous fungal spore concentrations increase during this period (Fig. 3).

Principal components analysis (PCA) of common pollen taxa in the upper parts of the
 sequence (from 92.5 cm) (Fig. 4) supports this description of vegetation change. The first
 axis represents 40.0% of variance, and the second axis represents 28.3% of variance. The
 major assemblage change in this portion of the sequence is in the upper 8.5 cm (~61 cal BP /
 1889 CE; min: 1830 CE, max: 1939 CE), where both Poaceae and *Sphagnum* decline relative
 to a number of herbaceous taxa (e.g. *Plantago lanceolata*-type, *Rumex acetosa*-type,
 Cyperaceae). Prior to this, there are continual fluctuations which appear to represent
 relatively minor changes in Poaceae and *Sphagnum*, though there is some suggestion that
 older samples (92.5 cm to 60.5cm, ~403 to 291 cal BP / 1547 to 1659 CE) may have a
 tendency towards slightly higher Poaceae and lower *Sphagnum* relative abundances.

280

281 *Plant macrofossils*

282 Plant macrofossil analysis (48 samples) describes the character of local (*i.e.* on-site)
283 vegetation at Rickys Ball (Fig. 5). There are four macrofossil zones (mz: see Table 2), which
284 are briefly described here. The earliest zone is dominated by monocotyledon remains.
285 Charcoal is abundant in every sample in this zone and increases around 160 cm (~2950 cal
286 BP). From 114 cm (~588 cal BP / 1362 CE), *Sphagnum* is recorded consistently, though at
287 low levels within an assemblage that continues to be dominated by monocotyledon remains.
288 From 70 cm (~325 cal BP / 1625 CE) *Sphagnum* remains increase significantly, but above
289 10 cm (~69 cal BP / 1881 CE) the samples are all dominated by monocotyledon remains,
290 with no *Sphagnum* present.

291

292 *Testate amoebae*

293 Testate amoebae were only preserved in concentrations sufficient for the development of
294 reliable counts in the upper 94 cm of the sampled section. Analysis therefore focussed on 49
295 samples from the upper section of the core (~ the last 500 years) (Fig. 6, Table 3). The most
296 notable assemblage change occurs at 8.5 cm depth (~61 cal BP / 1889 CE), and prior to this
297 the assemblage is largely dominated by species indicative of ‘hummocky’
298 (microtopographically complex) bog surface conditions. There is a general decrease in testate
299 biovolume (*i.e.* a shift to smaller species) from around 40.5 cm (~210 cal BP / 1740 CE),
300 suggesting decreasing community metabolic rates (reduced capacity in the food web to
301 process energy) (Koenig et al. 2018; Marcisz et al. 2020). Above 8.5 cm (~61 cal BP / 1889
302 CE), the taxa indicative of ‘hummocky’ conditions that were previously dominant are largely

absent, and the assemblage becomes dominated by *Diffflugia pristis*-type. These are small testates that are indicators of moderate bog surface wetness, but generally not standing water (Woodland et al. 1998; Charman et al. 2000; Swindles et al. 2009). The trend towards decreasing biovolume is continued, asides from a brief peak at 4.5 cm (~43 cal BP / 1907 CE), indicating continued loss of larger species, which tend to be poorer colonisers and are less able to overcome perturbations (Marcisz et al. 2020).

Non-metric multidimensional scaling (NMDS) analysis (based on testate amoeba influx values) further illustrates the significance of the shift in testate amoeba assemblages from 10.5 cm (Fig. 7). This is shown on the first axis (NMDS1) and demonstrates a move towards assemblages characterised by *Diffflugia pristis*-type, *Cyclopyxis arcelloides*-type and *Pseudodiffflugia fulva*, possibly representing a novel stable state. This analysis indicates that earlier in the sequence (below 8.5 cm, before ~61 cal BP / 1889 CE), abundances of individual species fluctuated, but there were no similarly notable long-term changes in assemblage characteristics, suggesting a previous state of dynamic equilibrium.

Insects

Seven insect samples were analysed, from peat taken in 10 cm thick spits between 100 and 30 cm depth (~429 to 162 cal BP / 1521 to 1788 CE) (Supplementary Table 3, Fig. 8). Relative to pollen, palaeoentomological data typically represent more local conditions (Smith et al. 2010). Beetles (Coleoptera) represent the majority of insect remains recovered, with true bugs (Hemiptera) and flies (Diptera) also present, but rare (see Supplementary Table 3, nomenclature for Coleoptera follows Lucht 1987). Phytophagous beetle host plants (predominantly from Koch (1989a, b, 1992)) are also presented on Supplementary Table 3 (plant taxonomy follows Stace (2010)).

There is little change in insect assemblages through the sequence (100 cm to 30 cm) and the taxa present are characteristic of upland bogs (see Supplementary Table 3, Fig. 8), indicating that this habitat has existed in the area for the entirety of the period represented.

Discussion

Pre-nineteenth century ecological history

The palaeoecological record presented here covers a substantial part of the Holocene, with the lowest sections deposited earlier than 7000 cal BP (5000 BCE), during the Mesolithic. Plant macrofossils indicate that local, peat-forming vegetation communities during this period were dominated by monocots (grasses and sedges). *Alnus*, *Corylus*, *Betula*, *Quercus* and *Salix* pollen indicates the presence of mixed woodlands nearby, and their absence in macrofossil samples indicates that this was distal to the sampling site. Similar to previous studies (Merryfield and Moore 1974; Fyfe 2012), from around 3000 cal BP (1000 BCE) woodlands declined and there was a general increase in heathland (*e.g. Calluna vulgaris*) and disturbed grassland (*e.g. Plantago lanceolata*, *Potentilla*) taxa. As at other sites across the region, coincidently increasing quantities of charcoal suggests that increasing openness was driven by enhanced burning regimes (Fyfe et al. 2003, 2018) and coprophilous fungal spores suggest that grazing animals may have also played a role.

From the fourteenth century CE (112.5 cm: ~568 cal BP / 1382 CE), *Sphagnum* moss became established locally (demonstrated by spores and plant macrofossils), increasing substantially during the seventeenth century CE (68.5 cm: ~319 cal BP / 1631 CE). This was accompanied by microbial (*e.g. Archerella flavum*) and faunal (*e.g. Plateumaris discolor*) wetland taxa. Increasing *Plantago lanceolata* pollen and decreasing coprophilous fungal spore influx

during this period suggests a change in local disturbance regimes. Fungal taxa like *Sporormiella* and *Podospora* are specialist coprophiles (Baker et al. 2013; Perrotti and van Asperen 2018), whereas *P. lanceolata* is an adaptable, ruderal generalist that may be favoured by various disturbances, including moderate grazing (Mook et al. 1989; Reudler et al. 2013). This suggests that local grazing pressure may have been reduced (e.g. fewer grazing animals and/or modified annual grazing patterns), possibly associated with increasing surface wetness. Around the eighteenth century (from 40.5 cm, ~210 cal BP / 1740 CE), *Sphagnum* spores and overall testate amoeba biovolume both began general long-term declines, suggesting a gradual decrease in *Sphagnum*-rich peatbog habitats, though these habitats appear to have persisted locally, as indicated by *Sphagnum* macrofossils.

Ecological impacts of nineteenth century moorland ‘improvement’

The most rapid ecological change recorded appears to have occurred during the late nineteenth century (from ~10.5 cm, ~70 cal BP / 1880 CE, 95% confidence range: 1822-1931 CE). *Sphagnum* moss almost disappeared locally (indicated by plant macrofossils) and declined substantially in the wider landscape (indicated by spores), with vegetation communities becoming dominated by grasses. Testate amoeba assemblages became dominated by *Diffflugia pristis*-type, indicators of moderate surface wetness (but not standing water) (Woodland et al. 1998; Charman et al. 2000; Swindles et al. 2009), largely replacing species indicative of ‘hummocky’ bog surface conditions, suggesting a simplification of local microtopography. These changes in floral and microbial communities indicate a rapid change in the landscape, consistent with nineteenth century drainage on Exmoor (Hegarty and Wilson-North 2014). A notable increase in influx rates of all coprophilous fungal spore types (*Sordaria*-type, *Sporormiella*-type and *Podospora*-type) and a decline in *Plantago lanceolata*

also occurred at the same time. Whilst *P. lanceolata* may be favoured by various disturbances, including moderate grazing pressure (Mook et al. 1989), repeated defoliation under heavy grazing pressure may conceivably reduce fitness (Reudler et al. 2013), suggesting the initiation of a high-intensity grazing regime.

The precise timing of this ecological shift is difficult to determine on a decadal scale, due to uncertainty inherent in age-depth modelling (e.g. 10.5 cm ~70 cal BP / 1880 CE, 95% confidence range: 1822-1931 CE, see Fig. 2), but occurrence during Knight family ownership of the Royal Forest of Exmoor (1818-1886) seems likely. This was a period of deliberate social, economic and agricultural change (Orwin and Sellick 1970; Hegarty and Wilson-North 2014), with relevance for modern Exmoor (e.g. Riley 2019). During the mid to late nineteenth century thousands of Scottish Blackface and Northumbrian Cheviot sheep, which could be pastured on Exmoor throughout the year, were introduced and areas of peat were drained and managed to ‘improve’ the land for this purpose (Orwin and Sellick 1970; Hegarty and Wilson-North 2014). Notably, a flock of Cheviot sheep were brought to Cornham Farm (adjacent to Ricksy Ball, see Fig. 1), in 1873 (Orwin and Sellick 1970). This is strikingly consistent with the palaeoecological evidence for the disappearance of *Sphagnum*, intensification of grazing and associated changes in vegetation and microbial communities in the upper 10.5 cm of the sequence and provides the most likely explanation.

Implications for modern landscape management

The nature of local and regional ecology, and the functioning of the moorland, has been influenced by successive changes in management practices, including different fire, grazing and drainage regimes. Our palaeoecological analyses provide a range of reference conditions

397 from the past, as well as revealing the ecological consequences of these catchment
398 management practices.

399 Palaeoecological analyses demonstrate that the post-drainage ecology of Ricksy Ball is
400 characterised by novel communities that do not have past analogues. In vegetation
401 assemblages, the loss of *Sphagnum* is the most conspicuous difference pre- and post-
402 drainage, but more subtle changes in *Plantago lanceolata* (decline), monocots (increase) and
403 *Calluna vulgaris* (increase) are also evident. Microbial (testate amoebae) assemblages are
404 strikingly different post-drainage, shifting from a relatively diverse community of species
405 associated with ‘hummocky’ bog surfaces, to a community lacking in those species and
406 dominated by a small number of taxa. Sub-fossil insect assemblages were only recovered
407 from pre-drainage contexts, but these can be compared to a modern entomological survey
408 from Ricksy Ball (undertaken in 2009, prior to restoration works) (Boyce 2010): notably, the
409 most abundant taxon in sub-fossil samples, *Plateumaris discolor* (associated with a range of
410 marshland plants, often in acid conditions (Cox 2007)), was lacking from the modern survey.
411 Collectively, this evidence suggests that post-drainage Ricksy Ball was distinct from pre-
412 drainage in terms of both biota and microhabitat diversity. This is a common pattern amongst
413 UK uplands that underwent major disturbances (e.g. drainage, peat-cutting, high-intensity
414 grazing, nitrogen deposition) during the nineteenth and twentieth centuries, subsequently
415 becoming characterised by low *Sphagnum* abundance and mono-dominant surface vegetation
416 types (e.g. *Molinia caerulea* (Poaceae), *Calluna vulgaris*) (Chambers et al. 2013, 2017;
417 Davies 2016). As these periods of major disturbance typically occurred prior to the initiation
418 of long-term ecological monitoring studies (e.g. UK Countryside Survey (UK Centre for
419 Ecology & Hydrology 2022)), this highlights the utility of palaeoecological data in
420 understanding systems targeted for restoration (Davies and Bunting 2010; Chambers et al.
421 2017; Chambers 2022; Marcisz et al. 2022).

There is an opportunity for mutual learning between ecology and palaeoecology if organisms studied as palaeoenvironmental proxies are incorporated into peatland restoration monitoring programmes. In particular, these could be enhanced by the incorporation of testate amoebae (Davis and Wilkinson 2004; Payne 2013; Swindles et al. 2016), which have been shown here to be sensitive bioindicators of changing catchment conditions. They are important in peatland microbial communities, often occupying high trophic levels (Jassey et al. 2013), and they are a food source for invertebrates (*e.g.* chironomid larvae) (Mieczan et al. 2015), providing a link between microbial and faunal food webs. The reduction in testate amoebae biovolume over the last 500 years suggests decreasing community metabolic rates (reduced capacity in the food web to process energy). Given the trophic position of testate amoebae, this may be associated with changes in other parts of the trophic system (*e.g.* invertebrate biomass), but more work is required to understand this aspect of peatland ecosystem function (Swindles et al. 2016).

On Exmoor, there is considerable interest amongst certain stakeholder groups in ‘reversing’ the effects of nineteenth century land management practices, particularly peatland drainage (Grand-Clement et al. 2015). A key aim of this is to raise water tables, restoring hydrological function and associated ecosystem services (*e.g.* carbon sequestration, enhanced water quality, flood prevention). This is achieved through practical interventions such as ditch-blocking, which may promote the re-growth of *Sphagnum* and increase local biodiversity and carbon storage (Grand-Clement et al. 2013, 2015). Whether the resultant ecological communities are likely to be taxonomically comparable to past communities remains unclear, though the relative importance of this may depend on different stakeholder interests, and which points in time are compared to the present. For example, peatlands with a mixture of surface vegetation functional types (*e.g.* ericaceous shrubs and grasses) are likely to be more

efficient for carbon sequestration than those with mono-dominant types (particularly grass) (Ward et al. 2013), but both situations have existed in the past (see Figures 4 and 5).

There is no single, readily identifiable, ‘stable’ pre-drainage ecological baseline. The system is inherently dynamic and successive changes in management practices over centuries and millennia (including different fire, grazing and drainage regimes) have continually modified the nature of local and regional ecology, and the functioning of the moorland. There is also little evidence for post-drainage communities reverting to pre-drainage states within the timescale of the record (*n.b.* sampling was undertaken prior to restoration works). In this context, knowledge of the past derived from palaeoecological data acts as “a guide rather than a template” (Higgs et al. 2014), providing a range of reference conditions and revealing the consequences of past management practices, rather than specifying a single target for restoration.

Conclusions

Palaeoecological analyses reveal varied management practices (burning, grazing, drainage) and associated ecological responses over the last eight millennia at Rickys Ball (Exmoor), and major changes in vegetation and microbial communities have been broadly coincident over the last five hundred years. During the nineteenth century, a combination of drainage and increased grazing intensity appears to have substantially altered the local ecology, resulting in the loss of *Sphagnum* and associated macro- and microbiota. This occurred in the context of more gradual, centennial-scale declines in *Sphagnum* and microbial biovolumes.

There is no single, readily identifiable, ‘stable’ pre-drainage baseline. We provide a range of reference conditions and reveal the consequences of past management practices, showing that

the moorland's character and ecological functioning have been influenced by changes in land management for millennia, and this may have been most pronounced during the nineteenth century.

References

- Appleby PG (2001) Chronostratigraphic Techniques in Recent Sediments. In: Last WM, Smol JP (eds) Tracking Environmental Change Using Lake Sediments. Volume 1: Basin Analysis, Coring, and Chronological Techniques. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp 171–203
- Appleby PG, Oldfield F (1978) The calculation of lead-210 dates assuming a constant rate of supply of unsupported ^{210}Pb to the sediment. *Catena* 5:1–8.
[https://doi.org/10.1016/S0341-8162\(78\)80002-2](https://doi.org/10.1016/S0341-8162(78)80002-2)
- Baker AG, Bhagwat SA, Willis KJ (2013) Do dung fungal spores make a good proxy for past distribution of large herbivores? *Quat Sci Rev* 62:21–31.
<https://doi.org/10.1016/j.quascirev.2012.11.018>
- Belyea LR, Malmer N (2004) Carbon sequestration in peatland: patterns and mechanisms of response to climate change. *Glob Chang Biol* 10:1043–1052.
<https://doi.org/10.1111/j.1529-8817.2003.00783.x>
- Binney HA, Waller MP, Bunting MJ, Armitage RA (2005) The interpretation of fen carr pollen diagrams: The representation of the dry land vegetation. *Rev Palaeobot Palynol* 134:197–218. <https://doi.org/10.1016/j.revpalbo.2004.12.006>
- Blaauw M (2021a) rbacon: Age-Depth Modelling using Bayesian Statistics

491 Blaauw M (2021b) clam: Classical Age-Depth Modelling of Cores from Deposits.

492 Blaauw M (2010) Methods and code for “classical” age-modelling of radiocarbon sequences.

493 Quat Geochronol 5:512–518. <https://doi.org/10.1016/j.quageo.2010.01.002>

494 Blaauw M, Christen JA (2011) Flexible paleoclimate age-depth models using an

495 autoregressive gamma process. Bayesian Anal 6:457–474. [https://doi.org/10.1214/11-](https://doi.org/10.1214/11-BA618)

496 BA618

497 Blaus A, Reitalu T, Gerhold P, et al (2020) Modern Pollen–Plant Diversity Relationships

498 Inform Palaeoecological Reconstructions of Functional and Phylogenetic Diversity in

499 Calcareous Fens. Front Ecol Evol 8:1–22. <https://doi.org/10.3389/fevo.2020.00207>

500 Blockley SPE, Pyne-O’Donnell SDF, Lowe JJ, et al (2005) A new and less destructive

501 laboratory procedure for the physical separation of distal glass tephra shards from

502 sediments. Quat Sci Rev 24:1952–1960. <https://doi.org/10.1016/j.quascirev.2004.12.008>

503 Blundell A, Holden J (2015) Using palaeoecology to support blanket peatland management.

504 Ecol Indic 49:110–120. <https://doi.org/10.1016/j.ecolind.2014.10.006>

505 Bonn A, Allott T, Evans M, et al (2016) Peatland Restoration and Ecosystem Services:

506 Science, Policy and Practice. Cambridge University Press, Cambridge

507 Booth RK, Lamentowicz M, Charman DJ (2010) Preparation and analysis of testate amoebae

508 in peatland palaeoenvironmental studies. Mires Peat 7:1–7

509 Bowes AC (2006) Exmoor Blanket Bog Inventory and Resoration Plan for English Nature.

510 University of Calgary

511 Boyce D (2010) Invertebrate survey of Exmoor MIRE blanket bog sites, 2009. Report to

512 Exmoor National Park Authority.

513 Chambers FM (2022) The use of paleoecological data in mire and moorland conservation.
514 Past Glob Chang Mag 30:16–17. <https://doi.org/10.22498/pages.30.1.16>

515 Chambers FM, Cloutman EW, Daniell JRG, et al (2013) Long-term ecological study
516 (palaeoecology) to chronicle habitat degradation and inform conservation ecology: An
517 exemplar from the Brecon Beacons, South Wales. Biodivers Conserv 22:719–736.
518 <https://doi.org/10.1007/s10531-013-0441-4>

519 Chambers FM, Crowle A, Daniell J, et al (2017) Ascertaining the nature and timing of mire
520 degradation: using palaeoecology to assist future conservation management in Northern
521 England. AIMS Environ Sci 4:54–82. <https://doi.org/10.3934/environsci.2017.1.54>

522 Chambers FM, Mauquoy D, Gent A, et al (2007) Palaeoecology of degraded blanket mire in
523 South Wales: Data to inform conservation management. Biol Conserv 137:197–209.
524 <https://doi.org/10.1016/j.biocon.2007.02.002>

525 Charman DJ, Hendon D, Woodland W (2000) The identification of testate amoebae
526 (Protozoa: Rhizopoda) in peats. Quaternary Research Association

527 Cox M (2007) Atlas of the seed and leaf beetles of Britain and Ireland. Pisces Publications,
528 Newbury

529 Dallimer M, Tinch D, Acs S, et al (2009) 100 years of change: Examining agricultural trends,
530 habitat change and stakeholder perceptions through the 20th century. J Appl Ecol
531 46:334–343. <https://doi.org/10.1111/j.1365-2664.2009.01619.x>

532 Davies AL (2016) Late Holocene regime shifts in moorland ecosystems: high resolution data
533 from the Pennines, UK. Veg Hist Archaeobot 25:207–219

534 Davies AL, Bunting MJ (2010) Applications of Palaeoecology in Conservation. 44:54–67

535 Davis SR, Wilkinson DM (2004) The conservation management value of testate amoebae as
 536 “restoration” indicators: speculations based on two damaged raised mires in northwest
 537 England. *The Holocene* 14:135–143. <https://doi.org/10.1191/0959683604hl696rp>
 538 Djamali M, Cilleros K (2020) Statistically significant minimum pollen count in Quaternary
 539 pollen analysis; the case of pollen-rich lake sediments. *Rev Palaeobot Palynol*
 540 275:104156. <https://doi.org/10.1016/j.revpalbo.2019.104156>
 541 Dugmore AJ, Larsen G, Newton AJ (1995) Seven tephra isochrones in Scotland. *The*
 542 *Holocene* 5:257–266. <https://doi.org/10.1177/095968369500500301>
 543 Fyfe RM (2012) Bronze Age landscape dynamics: Spatially detailed pollen analysis from a
 544 ceremonial complex. *J Archaeol Sci* 39:2764–2773.
 545 <https://doi.org/10.1016/j.jas.2012.04.015>
 546 Fyfe RM, Brown AG, Rippon SJ (2003) Mid- to late-Holocene vegetation history of Greater
 547 Exmoor, UK: Estimating the spatial extent of human-induced vegetation change. *Veg*
 548 *Hist Archaeobot* 12:215–232. <https://doi.org/10.1007/s00334-003-0018-3>
 549 Fyfe RM, Ombashi H, Davies HJ, Head K (2018) Quantified moorland vegetation and
 550 assessment of the role of burning over the past five millennia. *J Veg Sci* 29:393–403.
 551 <https://doi.org/10.1111/jvs.12594>
 552 Gallego-Sala A V., Clark JM, House JJ, et al (2010) Bioclimatic envelope model of climate
 553 change impacts on blanket peatland distribution in Great Britain. *Clim Res* 45:151–162.
 554 <https://doi.org/10.3354/cr00911>
 555 Gatis N, Luscombe DJ, Grand-Clement E, et al (2016) The effect of drainage ditches on
 556 vegetation diversity and CO₂ fluxes in a *Molinia caerulea*-dominated peatland.
 557 *Ecohydrology* 9:407–420. <https://doi.org/10.1002/eco.1643>

558 Gearey BR, Fyfe RM (2016) Peatlands as knowledge archives. In: Bonn A, Allott T, Evans
 559 M, et al. (eds) Peatland Restoration and Ecosystem Services: Science, Policy and
 560 Practice. Cambridge University Press, Cambridge

561 Gilbert D, Amblard C, Bourdier G, Francez AJ (1998) The microbial loop at the surface of a
 562 peatland: Structure, function, and impact of nutrient input. *Microb Ecol* 35:83–93.
 563 <https://doi.org/10.1007/s002489900062>

564 Grand-Clement E, Anderson K, Smith D, et al (2015) New approaches to the restoration of
 565 shallow marginal peatlands. *J Environ Manage* 161:417–430.
 566 <https://doi.org/10.1016/j.jenvman.2015.06.023>

567 Grand-Clement E, Anderson K, Smith D, et al (2013) Evaluating ecosystem goods and
 568 services after restoration of marginal upland peatlands in South-West England. *J Appl*
 569 *Ecol* 50:324–334. <https://doi.org/10.1111/1365-2664.12039>

570 Hegarty C, Wilson-North R (2014) The Archaeology of Hull Farming on Exmoor. English
 571 Heritage, Swindon

572 Higgs E, Falk DA, Guerrini A, et al (2014) The changing role of history in restoration
 573 ecology. *Front Ecol Environ* 12:499–506. <https://doi.org/10.1890/110267>

574 Holden J, Gascoign M, Bosanko NR (2007) Erosion and natural revegetation associated with
 575 surface land drains in upland peatlands. *Earth Surf Process Landforms* 32:1547–1557.
 576 <https://doi.org/10.1002/esp.1476>

577 Jassey VEJ, Meyer C, Dupuy C, et al (2013) To What Extent Do Food Preferences Explain
 578 the Trophic Position of Heterotrophic and Mixotrophic Microbial Consumers in a
 579 Sphagnum Peatland? *Microb Ecol* 66:571–580. [https://doi.org/10.1007/s00248-013-](https://doi.org/10.1007/s00248-013-0262-8)
 580 0262-8

581 Kenward H, Hall A, Jones A (1980) A tested set of techniques for the extraction of plant and
 582 animal macrofossils from water-logged archaeological deposits. *Sci Archaeol* 22:3–15

583 Koch K (1989a) *Die Käfer Mitteleuropas. Ökologie, 1.* Goecke & Evers, Krefeld

584 Koch K (1989b) *Die Käfer Mitteleuropas, Ökologie 2.* Goecke & Evers, Krefeld

585 Koch K (1992) *Die Käfer Mitteleuropas. Ökologie 3.* Goecke & Evers, Krefeld

586 Koenig I, Mulot M, Mitchell EAD (2018) Taxonomic and functional traits responses of
 587 *Sphagnum* peatland testate amoebae to experimentally manipulated water table. *Ecol*
 588 *Indic* 85:342–351. <https://doi.org/10.1016/j.ecolind.2017.10.017>

589 Legendre P, Birks HJB (2012) From Classical to Canonical Ordination. In: Birks HJB, Lotter
 590 AF, Juggins S, Smol JP (eds) *Tracking Environmental Change Using Lake Sediments:*
 591 *Data Handling and Numerical Techniques.* Springer, London, pp 201–248

592 Lobley M, Winter M (2009) “Born out of crisis”: Assessing the Legacy of the Exmoor
 593 Moorland Management Agreements. *Rural Hist* 20:229–247.
 594 <https://doi.org/10.1017/S0956793309990069>

595 Lucht WH (1987) *Die Käfer Mitteleuropas (Katalog).* Krefeld: Goecke and Evers

596 Marcisz K, Czerwiński S, Lamentowicz M, et al (2022) How paleoecology can support
 597 peatland restoration. *Past Glob Chang Mag* 30:12–13.
 598 <https://doi.org/10.22498/pages.30.1.12>

599 Marcisz K, Jassey VEJ, Kosakyan A, et al (2020) Testate Amoeba Functional Traits and
 600 Their Use in Paleoecology. *Front Ecol Evol* 8:.
 601 <https://doi.org/10.3389/fevo.2020.575966>

602 Mauquoy D, Hughes P, van Geel B (2010) A protocol for plant macrofossil analysis of peat

603 deposits. Mires Peat 7:

604 McCarroll J, Chambers FM, Webb JC, Thom T (2016) Using palaeoecology to advise
605 peatland conservation: An example from West Arkengarthdale, Yorkshire, UK. J Nat
606 Conserv 30:90–102. <https://doi.org/10.1016/j.jnc.2016.02.002>

607 Merryfield DL, Moore PD (1974) Prehistoric human activity and blanket peat initiation on
608 Exmoor. Nature 250:439–441. <https://doi.org/10.1038/250439a0>

609 Mieczan T, Niedźwiecki M, Tarkowska-Kukuryk M (2015) Effects of rotifers, copepods and
610 chironomid larvae on microbial communities in peatlands. Eur J Protistol 51:386–400.
611 <https://doi.org/10.1016/j.ejop.2015.06.010>

612 Mighall TM, Abrahams PW, Grattan JP, et al (2002) Geochemical evidence for atmospheric
613 pollution derived from prehistoric copper mining at Copa Hill, Cwmystwyth, mid-
614 Wales, UK. Sci Total Environ 292:69–80. [https://doi.org/10.1016/S0048-](https://doi.org/10.1016/S0048-9697(02)00027-X)
615 [9697\(02\)00027-X](https://doi.org/10.1016/S0048-9697(02)00027-X)

616 Mitchell EAD (2004) Response of testate amoebae (Protozoa) to N and P fertilization in an
617 Arctic Wet Sedge Tundra. Arctic, Antarct Alp Res 36:78–83.
618 [https://doi.org/10.1657/1523-0430\(2004\)036\[0078:ROTAPT\]2.0.CO;2](https://doi.org/10.1657/1523-0430(2004)036[0078:ROTAPT]2.0.CO;2)

619 Mook JH, Haeck J, Van Der Toorn J, Van Tienderen PH (1989) Comparative demography of
620 *Plantago*. I. Observations on eight populations of *Plantago lanceolata*. Acta Bot Neerl
621 38:67–78. <https://doi.org/10.1111/j.1438-8677.1989.tb01913.x>

622 Moore P, Webb J, Collinson ME (1991) Pollen Analysis. Blackwell, Oxford

623 Oksanen J, Blanchet F, Friendly F, et al (2019) vegan: Community Ecology Package. R
624 package version 2.5-6.

625 Orwin CS, Sellick RJ (1970) *The Reclamation of Exmoor Forest*, Second Edi. David and
 626 Charles, Newton Abbot

627 Payne RJ (2013) Seven reasons why protists make useful bioindicators. *Acta Protozool*
 628 52:105–113. <https://doi.org/10.4467/16890027AP.13.0011.1108>

629 Payne RJ, Mitchell EAD (2009) How many is enough? Determining optimal count totals for
 630 ecological and palaeoecological studies of testate amoebae. *J Paleolimnol* 42:483–495.
 631 <https://doi.org/10.1007/s10933-008-9299-y>

632 Perrotti AG, van Asperen EN (2018) Dung fungi as a proxy for megaherbivores:
 633 opportunities and limitations for archaeological applications. *Veg Hist Archaeobot* 0:1–
 634 12. <https://doi.org/10.1007/s00334-018-0686-7>

635 Piotrowska N, Blaauw M, Mauquoy D, Chambers FM (2011) Constructing deposition
 636 chronologies for peat deposits using radiocarbon dating. *Mires Peat* 7:1–14.
 637 <https://doi.org/10.1111/j.1365-2486.2009.01920.x>

638 R Core Team (2021) *R: A language and environment for statistical computing*.

639 Reed MS, Bonn A, Slee W, et al (2009) The future of the uplands. *Land use policy* 26:204–
 640 216. <https://doi.org/10.1016/j.landusepol.2009.09.013>

641 Reimer PJ, Austin WEN, Bard E, et al (2020) The IntCal20 Northern Hemisphere
 642 Radiocarbon Age Calibration Curve (0–55 cal kBP). *Radiocarbon* 62:725–757.
 643 <https://doi.org/10.1017/RDC.2020.41>

644 Reudler JH, Honders SC, Turin H, Biere A (2013) Trade-offs between chemical defence and
 645 regrowth capacity in *Plantago lanceolata*. *Evol Ecol* 27:883–898.
 646 <https://doi.org/10.1007/s10682-012-9609-8>

647 Riley H (2019) The Landscape of the Knights on Exmoor: A Case Study for the Exmoor
648 Mires Partnership

649 Riley H, Wilson-North R (2001) The field archaeology of Exmoor. English Heritage,
650 Swindon

651 Ritson JP, Bell M, Brazier RE, et al (2016) Managing peatland vegetation for drinking water
652 treatment. *Sci Rep* 6:1–9. <https://doi.org/10.1038/srep36751>

653 Robinson M (1981) The use of ecological groupings of Coleoptera for comparing sites. In:
654 Jones M, Dimbleby G (eds) *The Environment of Man: the Iron Age to the Anglo-Saxon*
655 *Period*. B.A.R. British Series 87, Oxford, pp 251–286

656 Robinson M (1983) Arable/Pastoral Ratios From Insects? In: Jones M (ed) *Integrating the*
657 *Subsistence Economy* (British Archaeological Reports, International Series 181). British
658 Archaeological Reports, Oxford, pp 19–47

659 Smith D, Whitehouse NJ, Bunting MJ, Chapman H (2010) Can we characterise “openness” in
660 the Holocene palaeoenvironmental record? Modern analogue studies of insect faunas
661 and pollen spectra from Dunham Massey deer park and Epping Forest, England. *The*
662 *Holocene* 20:215–229. <https://doi.org/10.1177/0959683609350392>

663 Stace C (2010) *New Flora of the British Isles*, 3rd Editio. Cambridge University Press,
664 Cambridge

665 Stockmarr J (1971) Tablets with spores used in absolute pollen analysis. *Pollen et Spores*
666 XIII:615–621

667 Swindles GT, Charman DJ, Roe HM, Sansum PA (2009) Environmental controls on peatland
668 testate amoebae (Protozoa: Rhizopoda) in the North of Ireland: Implications for

669 Holocene palaeoclimate studies. *J Paleolimnol* 42:123–140.
670 <https://doi.org/10.1007/s10933-008-9266-7>

671 Swindles GT, Green SM, Brown L, et al (2016) Evaluating the use of dominant microbial
672 consumers (testate amoebae) as indicators of blanket peatland restoration. *Ecol Indic*
673 69:318–330. <https://doi.org/10.1016/j.ecolind.2016.04.038>

674 UK Centre for Ecology & Hydrology (2022) UKCEH Countryside Survey.
675 <https://countrysidesurvey.org.uk/>

676 Välliranta M, Oinonen M, Seppä H, et al (2014) Unexpected Problems in AMS 14C Dating of
677 Fen Peat. *Radiocarbon* 56:95–108. <https://doi.org/10.2458/56.16917>

678 Ward SE, Ostle NJ, Oakley S, et al (2013) Warming effects on greenhouse gas fluxes in
679 peatlands are modulated by vegetation composition. *Ecol Lett* 16:1285–1293.
680 <https://doi.org/10.1111/ele.12167>

681 Whyte I (2006) Parliamentary enclosure and changes in landownership in an upland
682 environment: Westmorland, c. 1770-1860. *Agric Hist Rev* 54:240–256

683 Woodland WA, Charman DJ, Sims PC (1998) Quantitative estimates of water tables and soil
684 moisture in Holocene peatlands from testate amoebae. *The Holocene* 8:261–273.
685 <https://doi.org/10.1191/095968398667004497>

686




Fig. 1 Location map showing Rickys Ball and other notable sites nearby. Grey shading shows the boundary of the former Royal Forest. New and previously measured (Bowes 2006) peat depths.





Fig. 2 Chronological summary. *A*: Full age-depth model (with 95% confidence intervals); *B*: XRF results; *C*: Focussed view of the upper 55cm of the age-depth model (with 95% confidence intervals). Dates with red crosses were not used in the model

697



698

699 *Fig. 3 Pollen and spore diagram. Values are expressed as percentage of total land pollen, total land pollen plus spores (fern*
700 *and moss spores) or influx rates (coprophilous fungal spores)*






Fig. 4 PCA of common (present in >50% samples) pollen and spore taxa. Upper diagram shows taxa, lower diagram shows samples

704



705

706 *Fig. 5 Plant macrofossil diagram. Values of peat components are percentage estimates of total composition for each sample.*
707 *Sphagnum leaf counts as count data rather than proportional*

708




Fig. 6 Testate amoebae diagram. Tests were not preserved below 94 cm at sufficient concentrations to be counted





Fig. 7 NMDS of common (present in >50% samples) testate amoeba taxa. Upper diagram shows taxa, lower diagram shows samples




Fig. 8 Summary of key proxy indicators for the top 1 m of the section

Pollen Zone	Depth / Age	Description
RIB lpaz1	190.5 to 162.5 cm ~5753 to 1390 BCE (~7703 to 3340 cal BP)	The greatest relative abundances of arboreal taxa, including <i>Corylus avellana</i> -type, <i>Betula</i> , <i>Quercus</i> and <i>Salix</i> . High levels of Pteropsida suggest a fern understory in open woodland. Various herbaceous taxa are recorded, including Poaceae, Cyperaceae, <i>Filipendula</i> and <i>Galium</i> .
RIB lpaz2	162.5 to 114.5 cm ~1390 BCE to 1355 CE (~3340 to 595 cal BP)	Decline in <i>Corylus avellane</i> -type and <i>Betula</i> , which are briefly replaced by <i>Alnus glutinosa</i> , followed by an increase (doubling) in Poaceae relative abundances. Most other woodland taxa (and Pteropsida) also decrease. Herbaceous grassland taxa increase, including <i>Potentilla</i> -type and <i>Plantago lanceolata</i> at low, but continuous levels. <i>Sporormiella</i> -type and <i>Sordaria</i> -type spores increase in concentration at 152.5 cm (~1782 cal BP / 168 CE) (Fig. 3), suggesting increased herbivore density.
RIB lpaz3	114.5 to 94.5 cm ~1355 to 1540 CE (~595 to 410 cal BP)	Decrease in Poaceae and increases in <i>Calluna vulgaris</i> , <i>Alnus glutinosa</i> and <i>Corylus avellana</i> -type. Herbaceous grassland taxa (<i>Plantago lanceolata</i> , <i>Potentilla</i> -type) also decline slightly. Continued presence of coprophilous fungi: <i>Sporormiella</i> -type presence becomes less consistent, <i>Sordaria</i> -type becomes more consistent and there is a spike in <i>Podospora</i> -type.
RIB lpaz4	94.5 to 6.5 cm ~1540 to 1898 CE (~410 to 52 cal BP)	Poaceae dominates this zone, whilst <i>Corylus avellana</i> -type, <i>Alnus glutinosa</i> and <i>Calluna vulgaris</i> relative abundances decline to <1%. Improved grassland taxa, notably <i>Plantago lanceolata</i> , increase to their highest levels. <i>Sphagnum</i> abundance increases substantially, peaking at around 40.5 cm (~210 cal BP / 1740 CE) before undergoing a general decreasing trend. <i>Sordaria</i> -type spores are present at low concentrations, and <i>Sporormiella</i> -type and <i>Podospora</i> -type spores are largely absent <i>Sordaria</i> -type increase substantially at 8.5 cm.
RIB lpaz5	6.5 to 0 cm ~1898 to 2012 CE	Poaceae and improved grassland taxa (<i>Plantago lanceolata</i>) decrease and are replaced with taxa more associated with

	(~52 to -62 cal BP)	‘poorer’ moorland (<i>e.g. Galium</i> , Rubiaceae). <i>Sphagnum</i> declines to very low abundances. <i>Sporormiella</i> -type, <i>Sordaria</i> -type and <i>Podospora</i> -type increase substantially, followed by near absence in the uppermost sample.
722		
723		
724		
725		
726		
727		
728		
729		
730		
731		
732		
733		
734		
735		
736		
737		
738		
739		
740		
741		
742		
743		
744		
745		

Macro Zone	Depth / Age	Description
RIB mz1	188 to 114 cm ~5363 BCE to 1362 CE (~7313 to 588 cal BP)	Dominated by monocotyledon remains (some leaves, but mostly root material). Charred material is abundant. Some unidentified organic material (UOM) is present up to about 150 cm.
RIB mz2	114 to 70 cm ~1362 to 1624 CE (~588 to 326 cal BP)	An abrupt decline in charcoal, and the first consistent presence of <i>Sphagnum</i> , though at low abundances, and the monocotyledon remains continue to dominate.
RIB mz3	70 to 10 cm ~1624 to 1881 CE (~326 to 69 cal BP)	Marked by a substantial increase in <i>Sphagnum</i> remains (~25% relative abundance). Leaf counts indicate two main <i>Sphagnum</i> types: <i>Sphagnum</i> section <i>Cuspidata</i> and <i>Sphagnum pulchrum</i> . Lower samples also include <i>Polytrichum commune</i> .
RIB mz4	10 to 0 cm ~1881 to 2012 CE (~69 to -62 cal BP)	Assemblages dominated by monocotyledon remains. <i>Sphagnum</i> almost absent. Slight increase in UOM.

747

748

749

750

751

752

753

754

755

756

757

758

759

Zone	Depth / Age	Description
Lower	94.5 to 8.5 cm	Assemblage is largely dominated by <i>Archerella flavum</i> , <i>Hyalosphenia papilo</i> , <i>Nebela flabellehum</i> and <i>Nebela militaris</i> : species indicative of ‘hummocky’ (microtopographically complex) bog surface conditions (Charman et al. 2000), alongside a wide range of other taxa. General decrease in testate biovolume (<i>i.e.</i> a shift to smaller species) from around 40.5 cm (~210 cal BP / 1740 CE). This suggests decreasing metabolic rates (reduced capacity in the food web to process energy) in microbial communities (Koenig et al. 2018; Marcisz et al. 2020).
	~1540 to 1889 CE (~410 to 61 cal BP)	
Upper	8.5 to 0 cm ~1889 to 2012 CE (~61 to -62 cal BP)	Assemblage becomes dominated by <i>Diffflugia pristis</i> -type: small testates that generally indicate moderate surface wetness (but not standing water) (Woodland et al. 1998; Charman et al. 2000; Swindles et al. 2009), and loses species indicative of ‘hummocky’ (microtopographically complex) surface conditions (<i>e.g.</i> <i>Archerella flavum</i> (Charman et al. 2000)). This suggests a simplification of the bog surface. <i>Cyclopyxis arcelloides</i> -type, a species complex that may indicate a range of relative wetness conditions (Charman et al. 2000), continues as a dominant part of the assemblage. The trend towards decreasing biovolume is continued, asides from a brief peak at 4.5 cm (~43 cal BP / 1907 CE), indicating continued loss of larger species, which tend to be poorer colonisers and are less able to overcome perturbations (Marcisz et al. 2020).