

Prevalence and length of hospital stay in patients with aortic valve disease is lower amongst South Asians

Thakker, Clare; Gollop, Nicholas D.; Carter, Paul; Lavu, Deepthi; Uppal, Hardeep; Chandran, Suresh; Potluri, Rahul

DOI:

[10.1016/j.ijcard.2014.09.040](https://doi.org/10.1016/j.ijcard.2014.09.040)

License:

Other (please specify with Rights Statement)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Thakker, C, Gollop, ND, Carter, P, Lavu, D, Uppal, H, Chandran, S & Potluri, R 2014, 'Prevalence and length of hospital stay in patients with aortic valve disease is lower amongst South Asians', *International Journal of Cardiology*, vol. 177, no. 1, pp. 34-36. <https://doi.org/10.1016/j.ijcard.2014.09.040>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

NOTICE: this is the author's version of a work that was accepted for publication in International Journal of Cardiology. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in International Journal of Cardiology, Vol 177, Issue 1, November 2014, DOI: 10.1016/j.ijcard.2014.09.040.

Eligibility for repository checked November 2014

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Accepted Manuscript


Prevalence and length of hospital stay in patients with Aortic Valve Disease is lower amongst South Asians

Clare Thakker, Nicholas D. Gollop, Paul Carter, Deepthi Lavu, Hardeep Uppal, Suresh Chandran, Rahul Potluri

PII: S0167-5273(14)01742-2
DOI: doi: [10.1016/j.ijcard.2014.09.040](https://doi.org/10.1016/j.ijcard.2014.09.040)
Reference: IJCA 18816

To appear in: *International Journal of Cardiology*

Received date: 31 July 2014
Accepted date: 16 September 2014



Please cite this article as: Thakker Clare, Gollop Nicholas D., Carter Paul, Lavu Deepthi, Uppal Hardeep, Chandran Suresh, Potluri Rahul, Prevalence and length of hospital stay in patients with Aortic Valve Disease is lower amongst South Asians, *International Journal of Cardiology* (2014), doi: [10.1016/j.ijcard.2014.09.040](https://doi.org/10.1016/j.ijcard.2014.09.040)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Prevalence and length of hospital stay in patients with Aortic Valve Disease is lower amongst South Asians

International Journal of Cardiology - Short Communication

Clare Thakker ¹	BSc
Nicholas D Gollop ²	MBBCh
Paul Carter ³	BSc
Deepthi Lavu ⁴	MBBS
Hardeep Uppal ⁴	MRCPsych
Suresh Chandran ⁵	FRCP
Rahul Potluri ⁴	MRCP

1 – School of Clinical Medicine, University of Cambridge, Cambridge, UK

2 – Department of Medicine, Norfolk and Norwich University Hospital, Norwich, UK

3 – The Medical School, University of Birmingham, Birmingham, UK

4 – ACALM Study Unit in collaboration with Aston Medical School, Aston University, Birmingham, UK

5 – Department of Acute Medicine, North Western Deanery, UK

Running Title: Aortic Valve Disease, prevalence and length of hospital stay

Key word(s): Aortic valve stenosis, Length of Hospital Stay, Ethnic, South Asian

Conflict of interest statement: No conflicts of interest to declare No specific funding was received in relation to this article.

Correspondence to: Dr Rahul Potluri, Honorary Clinical Lecturer in Cardiology, Aston University, Email: rahulpotluri@outlook.com

Word Count: 962 words

Dear Editor,

Aortic valve stenosis is the most common lesion of heart valves in European and American populations with calcific aortic valve stenosis present in 2-7% of the population aged >65 years¹ and severe stenosis present in 3% over 75 years². Aortic valve stenosis can have serious consequences such as increased perioperative morbidity and mortality³ and a mortality of 75% in symptomatic patients with severe aortic stenosis⁴. Ethnicity may influence the prevalence of the condition as well as its progression; congenital bicuspid aortic valve, calcific aortic stenosis and pathological calcification are all more prevalent in black patients⁵. Progression of aortic valve calcification has also been reported to be greater in black patients⁶, although other studies have also demonstrated progression in this ethnic group to be lower⁷ or not affected⁸. There is therefore a limited and conflicting evidence base on the effect of ethnicity, with many previous studies excluding Asian patients, although such studies could highlight important differences risk of the condition and its complications.

To better understand the incidence and management of aortic valve stenosis in the United Kingdom (UK) we investigated the prevalence, LOS and its predictors in black and ethnic minority patients with aortic valve stenosis in the North West of England.

We examined LOS and ethnic variations using completely anonymous data of adult patients admitted across seven hospitals across Greater Manchester (North West England) between 2000 and 2013 using the ACALM (Algorithm of Comorbidities, Associations, Length of stay and Mortality) study protocol, which uses ICD-10 diagnosis and OPCS-4 procedure codes to trace patients. The Caucasian population served as the reference ethnic group. The black and ethnic minority populations were defined as South Asian, Afro-Caribbean, Oriental and mixed race ethnicity. Data on the LOS, age, gender, ethnicity, co-morbidities and type of admission was available for all patients. The presence of a co-morbidity was defined by the presence of any of the top eight causes of mortality in the UK: ischemic heart disease, heart failure, cerebrovascular disease, lung cancer, breast cancer, dementia, chronic kidney disease, and chronic obstructive airways disease. All analyses were performed using SPSS version 20.0 and Microsoft Office Excel 2007. Such methodology has been previously used by our group and others⁹⁻¹⁴.

During the study period, there were 929465 overnight admissions of which 4764 (0.51%) were coded for aortic valve stenosis. Patients of ethnic minority origin constituted 3.2% of aortic valve stenosis admissions and LOS was significantly shorter ($p<0.05$) in South Asian patients (but not in other ethnic minority groups)

compared to Caucasian patients with aortic valve stenosis after analysis by means of a multi-variant logistic regression model accounting for variations in age, gender and co-morbidity. Moreover, the mean age of South Asian patients with aortic valve stenosis was notably lower than those of Caucasian origin (56 v 73 years). Interestingly, aortic valve stenosis is more prevalent in Caucasian admissions than those of ethnic minority origin (Table 1).

The precise reasons for shorter LOS in South Asian patients with aortic valve stenosis are currently unknown but these findings are consistent with similar findings in previous work looking at LOS in patients with diabetes in Birmingham, UK⁹, patients with acute pulmonary embolism¹⁰ and myocardial infarction¹² in Greater Manchester, UK. Moreover, there is previous evidence to suggest that ethnic minorities may receive less support at discharge and that they may be discharged prematurely¹⁵. These potential inequalities require further investigation ideally via prospective cohort studies with solid quality outcomes such as adverse event rate and mortality. A limitation of our discussion is that it is based on the assumption that short LOS is related to recovery or hospital discharge as we have no data on mortality. Furthermore, we have no data on the proportion of patients undergoing aortic valve replacement and this has been shown to produce a variety of complications and different LOS according to risk factors such as age and gender^{16,17}: an important point to consider given that the South Asian admissions for aortic valve stenosis were notably younger than Caucasian admissions.

Our data suggests that aortic valve stenosis is more prevalent in Caucasian admissions than those of ethnic minority origin. In accordance with this, a previous study of a United States (US) population demonstrated a lower frequency of bicuspid aortic valve and reduced aortic dimensions in African Americans compared to Caucasians¹⁸ and in another study calcified aortic disease was 69% lower amongst African Americans^{7,8}. However, the current literature on this subject is conflicting, with reports of a greater risk of aortic valve disease in this patient group⁵⁻⁸. The evidence base on the influence of ethnicity is also limited, especially with respect to Asian patients with aortic valve disease, for whom the results of the present study are particularly relevant. The reasons for race being a modifier in aortic valve disease are unclear but previous studies have identified possible risk factors for degenerative aortic valve disease including, male sex, active smoking status and a history of hypertension¹⁹. It is possible that these risk factors may be present at different rates in Caucasian and ethnic minority populations. Moreover, recent studies have suggested that genetic loci associated with aortic valve stenosis may occur at a slightly higher frequency in the Caucasian population than in other ethnic groups²⁰. However, it is difficult to draw conclusions on the prevalence of aortic valve stenosis as it is

often asymptomatic and diagnosis via echocardiogram is often made relatively late. Thus, access to health care will affect the apparent prevalence in different populations and other studies have shown this can be influenced by age and sex²¹ as well as ethnic origin²².

We believe our study highlights the need for more research into the reasons for and methods for reduction of health inequalities in multi-ethnic populations with a view to improving care for patients with aortic valve stenosis.

Table 1 – Prevalence and characteristics of admissions for patients with aortic valve stenosis according to ethnic group

Ethnicity	Aortic valve stenosis per 1000 admissions	N (%)	Mean Age (years)	M:F Ratio	% admitted as emergency	% with co-morbidity	Mean Length of Stay (days)	ODDs ratio for length of stay*
All groups	4.91	4564 (100.0)	72	1.2:1	61.7	64.7	7.7	-
Caucasian	5.65	4053 (88.8)	73	1.1:1	62.7	65.2	13.7	1
South Asian	1.07	78 (1.7)	56	2.4:1	71.8	67.9	8.0	0.982 (0.968–0.996)
Afro-Caribbean	0.60	16 (0.4)	69	0.6:1	62.5	68.8	9.7	0.975 (0.915-1.038)
Oriental	1.02	6 (0.1)	68	2.0:1	50.0	66.7	14.0	1.010 (0.966-1.056)
Mixed	0.68	5 (0.1)	74	F=0	40.0	60.0	1.3	0.662 (0.331-1.325)
Other	1.92	39 (0.9)	69	0.8:1	71.8	71.8	12.1	1.003 (0.981-1.025)
Unknown	4.68	367 (8.0)	72	1.2:1	47.4	58.0	15.1	1.006 (0.999-1.013)
*adjusted for age, sex, co-morbidity, # statistically significant, p<0.05								

Research was performed with completely anonymous data and conformed with the local ethical review/research and development policies.

References

1. Lung B, Baron G, Butchart EG, Delahaye F, Gohlke-Bärwolf C, Levang OW, et al. A prospective survey of patients with valvular heart disease in Europe: The Euro Heart Survey on Valvular Heart Disease. *Eur Heart J*. 2003 Jul;24(13):1231–43.
2. Lindroos M, Kupari M, Heikkilä J, Tilvis R. Prevalence of aortic valve abnormalities in the elderly: an echocardiographic study of a random population sample. *J Am Coll Cardiol*. 1993 Apr;21(5):1220–5.
3. Brown J, Morgan-Hughes NJ. Aortic stenosis and non-cardiac surgery. *Contin Educ Anaesth Crit Care Pain*. 2005 Feb 1;5(1):1–4.
4. Ross J, Braunwald E. Aortic Stenosis. *Circulation*. 1968 Jul 1;38(1S5):V–61–V–67.
5. Novaro GM, Houghtaling PL, Gillinov AM, Blackstone EH, Asher CR. Prevalence of mitral valve prolapse and congenital bicuspid aortic valves in black and white patients undergoing cardiac valve operations. *Am J Cardiol*. 2013 Mar 15;111(6):898–901. doi: 10.1016/j.amjcard.2012.11.051. Epub 2012 Dec 28. PubMed PMID: 23276473.
6. Mohler ER, Sheridan MJ, Nichols R, Harvey WP, Waller BF. Development and progression of aortic valve stenosis: atherosclerosis risk factors--a causal relationship? A clinical morphologic study. *Clin Cardiol*. 1991 Dec;14(12):995–9. PubMed PMID: 1841025.
7. Novaro GM, Katz R, Aviles RJ, Gottdiener JS, Cushman M, Psaty BM, Otto CM, Griffin BP. Clinical factors, but not C-reactive protein, predict progression of calcific aortic-valve disease: the Cardiovascular Health Study. *J Am Coll Cardiol*. 2007 Nov 13;50(20):1992–8. Epub 2007 Oct 29. PubMed PMID: 17996566.
8. Owens DS, Katz R, Takasu J, Kronmal R, Budoff MJ, O'Brien KD. Incidence and progression of aortic valve calcium in the Multi-ethnic Study of Atherosclerosis(MESA). *Am J Cardiol*. 2010 Mar 1;105(5):701–8. doi: 10.1016/j.amjcard.2009.10.071. PubMed PMID: 20185020; PubMed Central PMCID: PMC2829478.
9. Kimani KN, Potluri R, Natalwala A, Ghosh S, Heun R, Narendran P. Length of hospital stay is shorter in black and ethnic minority patients with diabetes. *Diabet Med J Br Diabet Assoc*. 2012 Jun;29(6):830–1.
10. Smith SF, Gollop ND, Uppal H, Chandran S, Potluri R. Length of hospital stay is shorter in South Asian patients with acute pulmonary embolism. *Heart Asia*. 2014 Jan 1;6(1):1–2.
11. Ciputra RN, Sembiring YE, Prawoto OL, Khouw N, Baig M, Uppal H, Chandran S, Potluri R. Length of stay in hospital is longer in ethnic minority patients after coronary artery bypass surgery. *Int J Cardiol*. 2014 Apr 1;172(3):e434–5. doi: 10.1016/j.ijcard.2013.12.182. Epub 2014 Jan 8. PubMed PMID: 24456872
12. Khouw N, Wasim M, Aziz A, Uppal H, Chandran S, Potluri R. Length of hospital stay is shorter in South Asian patients with myocardial infarction. *Int J Cardiol*. 2014 Feb 1;171(2):e54–55.
13. Green B, Gollop ND, Baig M, Uppal H, Chandran S, Potluri R. Prolonged length of hospital stay in Far-East Asian and Afro-Caribbean patients with cardiomyopathy. *Int J Cardiol*. 2014 Jun 28. pii: S0167-5273(14)01079-1. doi: 10.1016/j.ijcard.2014.06.012. [Epub ahead of print] PubMed PMID: 25064199.
14. Potluri R, Baig M, Mavi JS, Ali N, Aziz A, Uppal H, Chandran S. The role of angioplasty in patients with acute coronary syndrome and previous coronary artery bypass grafting. *Int J Cardiol*. (IN PRESS)
15. Carers National Association. "You Can Take Him Home Now": Carer's Experience of Hospital Discharge. June 2001 [Internet]. 2001. Available from: <http://www.carersuk.org/professionals/resources/research-library/item/517-you-can-take-him-home-now-carers-experiences-of-hospitaldischarge>
16. Wang P, Acker MA, Bilello M, Melhem ER, Stambrook E, Ratcliffe SJ, et al. Sex, aging, and preexisting cerebral ischemic disease in patients with aortic stenosis. *Ann Thorac Surg*. 2010 Oct;90(4):1230–5.

17. Craver JM, Weintraub WS, Jones EL, Guyton RA, Hatcher CR Jr. Predictors of mortality, complications, and length of stay in aortic valve replacement for aortic stenosis. *Circulation*. 1988 Sep;78(3 Pt 2):185–90.
18. Chandra S, Lang RM, Nicolarsen J, Gayat E, Spencer KT, Mor-Avi V, et al. Bicuspid Aortic Valve: Inter-racial difference in frequency and aortic dimensions: Impact of race in adult BAV in a single center retrospective observational study. *Jacc Cardiovasc Imaging*. 2012 Oct;5(10):981–9.
19. Stewart BF, Siscovick D, Lind BK, Gardin JM, Gottdiener JS, Smith VE, et al. Clinical Factors Associated With Calcific Aortic Valve Disease. *J Am Coll Cardiol*. 1997 Mar 1;29(3):630–4.
20. Thanassoulis G, Campbell CY, Owens DS, Smith JG, Smith AV, Peloso GM, et al. Genetic associations with valvular calcification and aortic stenosis. *N Engl J Med*. 2013 Feb 7;368(6):503–12.
21. Bach DS, Radeva JI, Birnbaum HG, Fournier A-A, Tuttle EG. Prevalence, referral patterns, testing, and surgery in aortic valve disease: leaving women and elderly patients behind? *J Heart Valve Dis*. 2007 Jul;16(4):362–9.
22. Atkinson M, Clark M, Clay D, et al Coventry: Centre for Health Services Studies, University of Warwick, 2001. Systematic review of ethnicity and health service access for London. 2001.