UNIVERSITY BIRMINGHAM University of Birmingham Research at Birmingham

Atypical interoception as a common risk factor for psychopathology

Brewer, Rebecca; Murphy, Jennifer; Bird, Geoffrey

DOI: 10.1016/j.neubiorev.2021.07.036

License: Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Brewer, R, Murphy, J & Bird, G 2021, 'Atypical interoception as a common risk factor for psychopathology: a review', *Neuroscience and biobehavioral reviews*, vol. 130, pp. 470-508. https://doi.org/10.1016/j.neubiorev.2021.07.036

Link to publication on Research at Birmingham portal

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Contents lists available at ScienceDirect



Neuroscience & Biobehaviora Review:

Neuroscience and Biobehavioral Reviews

journal homepage: www.elsevier.com/locate/neubiorev

Atypical interoception as a common risk factor for psychopathology: A review

chopathology are highlighted.



Rebecca Brewer^a, Jennifer Murphy^a,*, Geoffrey Bird^{b,c}

^a Department of Psychology, Royal Holloway, University of London, United Kingdom

^b Department of Experimental Psychology, University of Oxford, United Kingdom

^c Department of Psychology, Institute of Psychiatry, Psychology and Neuroscience, King's College London, United Kingdom

A R T I C L E I N F O A B S T R A C T Keywords: Interoception Interoception Scomorbidity A B S T R A C T The inadequacy of a categorial approach to mental health diagnosis is now well-recognised, with many authors, diagnostic manuals and funding bodies advocating a dimensional, trans-diagnostic approach to mental health research. Variance in interoception, the ability to perceive one's internal bodily state, is reported across diagnostic boundaries, and is associated with atypical functioning across symptom categories. Drawing on behavioural and neuroscientific evidence, we outline current research on the contribution of interoception to numerous cognitive and affective abilities (in both typical and clinical populations), and describe the interoceptive atypicalities seen in a range of psychopathology, as well as the ways in which interoception may differ across

1. Structure of the paper

Debate as to the utility of a psychiatric nosology, such as that contained in the Diagnostic and Statistical Manual (5th edition; DSM-5; American Psychiatric Association, 2013), for research and health care delivery, is far from resolved. While diagnostic manuals such as DSM-5 now acknowledge that a continuous diagnostic approach may be favourable, substantial diversion from the categorical approach to diagnosis has not yet been achieved. Opponents of attempts to develop a classification of mental disorders point to the cross-disorder nature of many symptoms such as working memory impairments or emotion regulation difficulties (Cuthbert and Insel, 2013), and to the high rates of comorbidity and co-occurrence (Hasin and Kilcoyne, 2012; Kessler, Chui, et al., 2005) within psychiatry, such that half of those who meet diagnostic criteria for one disorder also meet diagnostic criteria for another disorder. Responses to this challenge tend to adopt one of two approaches. First, in the approach adopted by the National Institute of Health's Research Domain Criteria (RDoC), a collection of clinicians and basic scientists agree on a number of areas of psychological functioning, for example within domains such as positive and negative valence systems, cognitive systems, social processes, and arousal and regulatory mechanisms. Within this approach, traditional diagnostic categories are

ignored and each area of functioning is itself the unit of analysis (e.g. a study may seek to determine the genetic basis of trait anxiety, regardless of whether this anxiety is displayed by individuals with Generalised Anxiety Disorder or Autism Spectrum Disorder). Similarly, the Hierarchical Taxonomy of Psychopathology (HiTOP) provides a dimensional classification system with which to diagnose psychopathology, again identifying a range of continua representing variation within the typical and clinical populations, which may be used to classify patients (Kotov et al., 2017). In a related approach (e.g. Krueger et al., 1998), factor analysis has been used to uncover a higher-order structure among diagnostic symptoms. Original work suggested a two-factor structure, made up of an Internalising factor (predisposing towards mood and anxiety disorders) and an Externalising factor (predisposing towards substance abuse and antisocial disorders), which was later supplemented by a third Thought Disorder factor (predisposing towards psychotic symptoms; Kotov, Chang, et al., 2011; Kotov, Ruggero, et al., 2011). Several recent studies have extended this conceptualisation, however, demonstrating the existence of a single higher-order factor, namely the 'P Factor', representing lesser-to-greater severity of psychopathology with associated disruption in neural circuitry (Caspi et al., 2014; Lahey et al., 2012; Selzam et al., 2018). Although differing in their methodological approach, all of these conceptualisations suggest that

clinical presentations. A number of important areas for further research on the role of interoception in psy-

* Corresponding author at: Department of Psychology, Royal Holloway, University of London, Egham, Surrey, TW20 0EX, United Kingdom. *E-mail address:* Jennifer.murphy@rhul.ac.uk (J. Murphy).

https://doi.org/10.1016/j.neubiorev.2021.07.036

Received 16 December 2020; Received in revised form 9 July 2021; Accepted 31 July 2021 Available online 3 August 2021 0149-7634/© 2021 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0y). there are similarities in symptoms across psychopathology, and that impairments in a specific psychological process may lead to a range of symptoms that are shared across traditional diagnostic categories.

One such psychological process that has clear relevance within these frameworks is interoception, the perception of, interpretation of, or attention to, one's internal bodily state (note that henceforth 'internal state' refers to bodily states). A number of authors have previously highlighted the role that atypical interoception may play in a range of mental health conditions (Barrett and Simmons, 2015; Bonaz et al., 2021; Khalsa and Lapidus, 2016; Khalsa et al., 2018; Murphy, Brewer, et al., 2017; Paulus and Stein, 2010; Quadt et al., 2018; Tsakiris and Critchley, 2016), and we have previously proposed that interoceptive atypicalities may represent a common vulnerability factor for psychopathology (Brewer and Bird, 2019; Brewer et al., 2016a; Murphy, Geary, et al., 2017). Despite consensus in the literature regarding the relevance of interoception for psychopathology, and the recent publication of a number of reviews relevant to interoception, no papers to date have provided a detailed review of the evidence surrounding the relationship between interoceptive abilities and mental health across a broad range of clinical conditions. In this paper, we review the evidence that atypical interoception may represent a key risk factor for the development of psychopathology, across a range of traditional diagnostic categories.

As an introduction to the topic, we first define interoception, and discuss its measurement and neural basis. Subsequently, in order to show that interoception is a sufficiently basic process for atypicalities to affect multiple symptom domains, we describe the contribution of interoception to physical and mental health. In particular, we highlight the contribution of interoception to multiple aspects of emotional (and socio-emotional) processing, and for multiple aspects of learning and decision-making. We then review evidence for the presence of interoceptive impairment in multiple clinical conditions. Drawing these two sections together, we critically assess the available evidence suggesting that interoceptive impairments at least contribute to, or are responsible for, a range of symptoms across clinical conditions. This latter point includes explanation of how interoceptive deficits can explain heterogeneity within conditions and commonalities across clinical conditions. We argue that many symptoms are a product of atypical interoception, rather than consequences of individual disorders themselves. For example, it might be the case that a particular impairment associated with interoceptive deficits (e.g., emotion processing difficulties) seen in many, but not all, individuals diagnosed with eating disorders is due to the fact that a significant proportion of this population has atypical interoception (which may not be a product of the eating disorder per se). If true, the presence or absence of atypical interoception would explain the variance within the eating disorders population with respect to emotion processing. Furthermore, the fact that atypical interoception is seen in several other clinical conditions means that emotion processing difficulties should also be seen in other conditions besides eating disorders.

If our conjecture is correct, and atypical interoception confers a general susceptibility to psychopathology, then it is useful to address what may cause atypical interoception, particularly in the context of how individuals develop typical interoception. Furthermore, it is useful to consider the potential for intervention when interoception becomes problematic. These questions are addressed before we consider some outstanding questions fundamental to the idea that atypical interoception contributes to symptoms across conditions and suggest ways in which these might be addressed. Note that, in line with dimensional approaches to psychopathology, the current paper discusses interoception within the context of a wide range of conditions, including neurodevelopmental disorders such as autism, and neurodegenerative disorders, such as dementia.

2. Introduction to Interoception

2.1. What is Interoception?

On a surface level it is easy to define interoception; it is the ability to perceive the internal state of one's own body. However, such a seemingly simple definition masks uncertainty in specifying exactly how to define 'internal', and raises issues relating to the distinction between perception and recognition of such states. Whilst early definitions included visceral (internal) sensations only, more recent definitions include signals that are not visceral (e.g., affective touch) but which are processed using similar neural pathways as other interoceptive signals (see Khalsa and Lapidus, 2016; Chen et al., 2021). Thus, contemporary definitions of interoception include signals sent either via lamina I of the spinal and trigeminal dorsal horn to the anterior insula and anterior cingulate cortex (Craig, 2002) or via cranial nerves to the nucleus of the solitary tract (Critchley and Harrison, 2013); see Section 2.3 for a detailed description of the neural basis of interoception). Accordingly, signals relating to hunger, satiety, itch, thirst, muscular effort, bladder, gastrointestinal, respiratory, cardiac, temperature, blood (PH, glucose level), vasomotor flush, air hunger, shudder, sensual touch, genital sensation, bruising, headache, broken bones and many more visceral sensations are typically considered to be interoceptive signals (Khalsa and Lapidus, 2016), though this remains debated (Ceunen et al., 2016; Critchley and Garfinkel, 2017). Anterior insula has also been implicated in the processing of smell (Plailly et al., 2007), suggesting that olfaction could also be considered interoceptive under broader definitions. Similarly, taste appears to be represented in both the anterior dorsal insula, as well as more posteriorly (Small, 2010). Debate also exists as to whether proprioception (the sense of the position of one's body, particularly the position of the limbs in relation to the trunk) should also be considered interoceptive (Cameron, 2001; Vaitl, 1996); although information originates from inside the body, it is processed by distinct neurological systems, in particular the posterior insula (Bottini et al., 2001; Fasold et al., 2008; Ferrè et al., 2012; Petit and Beauchamp, 2003; Zu Eulenburg et al., 2013).

Further debate concerns the level of the cognitive representation required in order for interoception to have occurred (e.g. Khalsa et al., 2018). At one level, any instance of homeostatic control involves interoception. For example, the release of insulin in response to hyperglycemia relies on the detection of excess glucose in the blood. Such detection fulfils the definition of interoception even though this process may occur without the individual's conscious awareness, and various computational models have been put forward to explain the role of interoception in maintaining homeostasis (Petzschner et al., 2021). A higher degree of conscious awareness is associated with the perception of interoceptive signals. At this level, interoception has occurred when the individual is consciously aware that their interoceptive state has changed. Crucially, it is possible to distinguish this from a still higher level in which the individual is able to recognise the interoceptive signal explicitly. An individual with conscious perception, but without explicit recognition, of interoceptive signals may be aware that their internal state has changed, but be unable to label that state. For example, they may be aware that they are experiencing an unusual internal state, but be unaware of which specific interoceptive change they are experiencing. At the highest level of recognition, an individual would be able to detect that their state has changed, and to recognise and label their new interoceptive state (e.g., as heat, hunger, etc.). This characterisation is not without controversy, however, especially when applied to emotional internal states, when the specific theory of emotion one subscribes to plays a role. For example, under Schachter and Singer's (1962) model of emotion, determining whether one is hungry or hot may be achieved on the basis of interoceptive signals alone, but distinguishing between the emotions of surprise and fear, for example, may rely on the integration of interoceptive cues and the cognitive evaluation of situational cues. Under such a model, it may be possible for an

individual with intact interoceptive awareness to fail to recognise their own emotional internal states beyond a coarse valence level if they lack the experience or ability necessary to evaluate situational cues from the environment.

Beyond these (already problematic) definitions of interoception, separate components of interoception have been defined. Garfinkel and Critchley (2013) proposed a tripartite model of interoception comprising of three facets of interoceptive ability: 1) interoceptive sensitivity, one's accuracy of perceiving one's internal state (for example, how accurately an individual can count their heartbeats when instructed to do so), 2) interoceptive sensibility, an individual's self-reported beliefs regarding their attention to and accuracy of perceiving internal signals and 3) interoceptive awareness, a metacognitive measure reflecting the level of correspondence between an individual's true accuracy (interoceptive sensitivity) and their self-rated interoceptive sensibility. Following this initial description a number of variants of this model have been proposed (Khalsa et al., 2018; Murphy, Catmur, and Bird, 2018; Murphy, Catmur, and Bird, 2019); for example, one recent model highlights a need to distinguish between one's interoceptive accuracy and one's propensity to attend to interoceptive information, along both subjectively reported and objectively assessed axes (Murphy, Catmur, and Bird, 2018; Murphy, Catmur, and Bird, 2019). Previously available measures of interoception (see Section 2.2) led interoceptive accuracy to be measured most frequently using objective, performance-based tasks, and interoceptive attention to be measured using self-report measures. However, assuming both interoceptive attention and interoceptive accuracy can be measured using subjective and objective measures, Murphy, Catmur, and Bird (2018, 2019) proposed a 2 \times 2 dimensional structure of interoception, reflecting both what is measured (accuracy versus attention) as well as how it is measured (self-report versus objective performance) (see Fig. 1). Within this 2×2 structure, interoceptive awareness, as defined by Garfinkel and Critchley (2013), can be seen as the degree to which self-report and accuracy measures correlate with each other, either along the accuracy or the attention dimension. The propensity to use internal signals in one's daily life may also be an additional interoceptive dimension (Murphy, Catmur, and Bird, 2019), and this may be a function of interoceptive accuracy, attention and metacognitive interoceptive awareness combined.

Notably, as described above, there are multiple levels of representation at which interoception can occur (e.g. implicit homeostasis, conscious perception of a signal without recognition of the specific signal, recognition without the requirement for a verbal label, and verbal labelling of the signal), and tasks tend to tap into different levels of this hierarchy. For example, threshold tasks simply require participants to detect the presence or absence of a sensation, interoceptive load tasks require participants to determine whether there has been a change in signal strength, while questionnaire measures require individuals to reflect on distinct verbally labelled interoceptive states. The difficulties identifying relationships between different interoceptive tasks may, therefore, be exacerbated by tasks tapping into different levels of representation as well as different dimensions of interoception (attention, accuracy and metacognitive abilities) and different measurement types (objective performance or beliefs).

2.2. How is interoception measured?

The measurement of interoception is generally acknowledged to be the biggest difficulty within the field (for a review, see Quigley et al., 2021). Studies of interoceptive accuracy have relied almost exclusively upon measures of heartbeat counting (HCT) or discrimination (also referred to as 'heartbeat detection') (HDT). In heartbeat counting procedures, participants are asked to count their heartbeats without physically measuring their heartbeat over a series of intervals (typically between 3 and 6 time intervals in the range of \sim 25 to 100 s) (Dale and Anderson, 1978; Schandry, 1981). The difference between an objective measurement and the participants' estimate is taken as a measure of interoceptive sensitivity. Variants of this task also exist where counting procedures are replaced with tapping in time with one's heartbeat (Ludwick-Rosenthal and Neufeld, 1985). In heartbeat discrimination procedures, participants are asked to determine whether visual or auditory stimuli are presented synchronously or asynchronously with their heartbeat. Whilst cardiac sensitivity tasks are extensively used, and in some respects reliable, (Brener and Kluvitse, 1988; Jones, 1994; Wildman and Jones, 1982), their suitability for research has been questioned. First, approximately 40 % of typical individuals are not able to consciously perceive their heartbeat (Khalsa, Rudrauf, Sandesara, et al., 2009) making these tasks unsuitable for quantifying interoceptive sensitivity in individuals with poor interoception. While this issue has been addressed using infusions of isoproterenol (a beta-adrenergic agonist) to increase intensity of cardiac (and respiratory) signals (e.g. Hassanpour et al., 2018; Khalsa, Rudrauf, Sandesara, et al., 2009), this technique is invasive and so not always feasible to use. Second, heartbeat may be perceived via (exteroceptive) touch receptors due to the vibration of the chest wall, with factors such as the individual's percentage of body fat (Rouse et al., 1988), systolic blood pressure (O'Brien



Fig. 1. 2×2 dimensional structure of interoception, distinguishing between different measurement types (beliefs or performance) and interoceptive abilities (accuracy or attention). Metacognitive ability (referred to as interoceptive awareness; Garfinkel and Critchley, 2013, or interoceptive insight; Khalsa et al., 2018) is shown by the black arrows, reflecting the extent to which beliefs and performance are related, either within the accuracy or attention ability. Figure adapted from Murphy, Catmur, and Bird (2018) and Murphy, Catmur, and Bird (2019).

et al., 1998) resting heart rate, and heart rate variability (Knapp-Kline and Kline, 2005) potentially affecting the extent to which heartbeat is perceived via this route. Indeed, whilst such influence of physiology may not be a limitation *per se* (as presumably such factors act upon one's ability to perceive cardiac signals across all situations), physiological factors may impact on the above tasks of interoceptive sensitivity specifically; for example, individual differences in resting heartrate may alter the number of hits and misses in the heartbeat counting task. Likewise, elevated blood pressure may influence the time at which external signals are perceived as synchronous with one's heartbeat in the heartbeat discrimination task (O'Brien et al., 1998). At present the mechanism by which these physiological factors influence performance on tasks of cardiac interoceptive accuracy remains unknown.

Whilst the factors described above may affect performance in any cardiac-based measure of interoceptive accuracy, there are important considerations for each task specifically. The heartbeat counting task is also strongly influenced by an individual's beliefs regarding their own or the average resting heart rate (Brener and Ring, 2016; Murphy, Millgate, et al., 2018; Ring and Brener, 1996; Ring et al., 2015; Windmann et al., 1999). Indeed, beliefs about one's heartbeat can alter performance on a range of tasks; false feedback about heart rate, for example, has been found to alter perception of one's effort during exercise (Iodice et al., 2019). The extent to which beliefs about heart beat affect interoceptive accuracy estimates in the heartbeat counting task may depend on the task instructions given; whilst early instructions encouraged participants to 'estimate' if they could not feel their heartbeat (Brener and Ring, 2016; Schandry, 1981) more recent approaches advocate a response of zero in this instance (Murphy, Brewer, et al., 2018) given evidence that the instructions given to participants may alter the pattern of results obtained (Desmedt et al., 2020, 2018; Ehlers et al., 1995). Equally problematic, is the fact that the task is often administered in the absence of a control task, meaning that an individual's performance may be influenced by other factors (e.g., attention, motivation) that are not interoceptive. Whilst newer research has advocated the use of a timing control task (Ainley et al., 2014), it is likely that this time estimation is not an adequate control for the factors affecting performance on counting based measures of interoception (Desmedt et al., 2020). Notably, it has also been argued that the psychometric properties of the HCT are less than ideal; for example, performance on the HCT appears to differ across different time intervals and a weak correlation is often observed between the participants' actual and reported heartbeats (Zamariola et al., 2018). Such evidence has been taken to suggest that individuals generally display poor perception of cardiac signals (Zamariola et al., 2018). Importantly, Zamariola et al. (2018) also found that the correlation between reported and actual number of heartbeats did not differ across individuals in the first and second quintile of HCT performers, suggesting that the task may also not be sensitive to differences in cardiac accuracy in better performers (but see Ainley et al. (2020)). These factors, combined with the general factors affecting cardiac-based measures of interoceptive accuracy, lead to concerns about the reliability and validity of this task. Indeed, the few studies examining the test-retest reliability of the heartbeat counting task over longer durations suggest fairly poor temporal stability in children and adults (Ferentzi, Drew, et al., 2018; Koch and Pollatos, 2014; Murphy, Cheesman, et al., 2019). Whilst this may suggest that differing degrees of interoceptive accuracy are a product of an individual's state, rather than a stable trait (Wittkamp et al., 2018), it is possible that temporal changes in this task may be due to changes across time in physiological or psychological factors that are not interoceptive.

In contrast to the heartbeat counting task, research suggests that the heartbeat discrimination task is less influenced by beliefs (Phillips et al., 1999), making this a preferred method for quantifying interoceptive accuracy. However, this task is not without limitations; for example, a high number of trials are required to gain precise estimates of ability (Kleckner et al., 2015), and concerns have been raised over the difficulty of this task, meaning that it may not be suitable for examining

interoceptive accuracy at the lower range of ability (Brener and Ring, 2016). Further, although Ring and Brener (2018) argue that humans are able to judge very precisely (within 20 ms) whether two stimuli in different modalities are simultaneous (Zampini et al., 2005), a sizeable portion of variance in this task appears to be accounted for by the ability to determine synchronicity of (non-interoceptive) cross-modal signals (Knapp et al., 1997). This problem is exacerbated by the fact that individuals perceive heartbeats at different locations within the body, each of which is associated with a different temporal delay (Brener and Kluvitse, 1988; Christopher Ring and Brener, 1992). Control tasks assessing these abilities are rarely employed (although for an exception see Garfinkel et al., 2016). Variations also exist as to the exact methodology employed for the heartbeat discrimination task; whilst two-alternative forced choice procedures are used most frequently, it has been argued that such methods do not account for individual differences in the delay at which individuals perceive an external signal to be synchronous with their heartbeat, and that lengthier procedures (for example the method of constant stimuli) may be more appropriate (Brener and Ring, 2016). A recently developed task addresses these issues; in the Phase Adjustment Task (Plans et al., 2020), participants are required to adjust the phase relationship between a tone and their heartbeat in order to achieve synchrony. This task is quicker to complete than improved versions of the heartbeat discrimination task (such as the method of constant stimuli), and allows participants to identify any delay between the tone and their heartbeat as 'synchronous'; meaning it both accounts for the individual differences in delay length perceived as synchronous that cause problems for standard versions of the heart beat discrimination task, and makes it especially useful in clinical populations who may struggle with longer procedures or who require a high level of precision in their responses. In addition, in contrast to the HCT, non-interoceptive participants cannot perform well on the task simply by knowing their resting heart rate. The relationship with psychopathology is yet to be investigated, however, making future research utilising this task in clinical groups a priority.

Surprisingly, scores on the heartbeat counting and discrimination tasks often correlate only modestly, if at all (Forkmann et al., 2016; Hickman et al., 2020; Kandasamy et al., 2016; Knoll and Hodapp, 1992; Phillips et al., 1999; Ring and Brener, 2018; Schulz et al., 2013; Weisz et al., 1988). Whilst this may reflect the unreliability of the measures, it is possible that different abilities are quantified by each task (Khalsa and Lapidus, 2016). Such a conjecture is supported by evidence that different neural activation has been associated with each task (Schulz, 2016) and that certain perturbations (e.g., stress) have differential effects on performance across these tasks (Schulz et al., 2013). Such differences may largely reflect task demands; while the heartbeat counting task (at least in theory) quantifies an individual's ability to perceive their heartbeat, the discrimination procedure requires the individual to integrate an internal signal with an external signal (Couto et al., 2015). Given questions over the validity and specificity of certain measures of cardiac interoception, recent efforts have focused on the development of new measures of cardiac interoceptive accuracy (e.g., Plans et al., 2020).

Beyond tasks measuring interoception from heartbeat perception, a small number of studies have used a measure of gastric distension; Whitehead and colleagues, for example, assessed participants' ability to detect whether stomach contractions coincided with an exteroceptive light stimulus, as well as their ability to control their gastric motility, and observed a positive relationship between perception of cardiac and gastric signals (Whitehead and Drescher, 1980). Similarly, detection rates and intensity ratings of colon distension have been investigated (Hölzl et al., 1996) and Zaman et al. (2016) measured ratings of esophageal distension intensity, identifying thresholds for initial perception, discomfort and pain. The invasive nature of these tests, however, means that they have not been used extensively. An alternative assessment of gastric interoception is the water load test, which involves participants consuming water until the point of perceived fullness (e.g. Herbert, Muth, et al., 2012; Koch et al., 2000). An adapted version of this task (the Two Step Water Load Test; WLT-II) uses a two-stage procedure, whereby participants first drink until perceived satiation, and then to maximum fullness, in an attempt to control for stomach capacity (Van Dyck et al., 2016). It is assumed that interoceptive ability is negatively associated with volume of water consumed (or the proportion of water consumed in step one relative to step two in WLT-II). As such, participants' ability to perceive their heartbeat was negatively correlated with amount of water consumed in the original water load task (Herbert, Muth, et al., 2012). FMRI has also been used to assess interoceptive cortex activity while participants attend to sensations from the stomach (Simmons et al., 2013), experience proximal and distal distension of the oesophagus (Aziz et al., 2000), or attempt to regulate their gastric responses to a virtual rollercoaster (Li et al., 2017). These studies implicated interoceptive cortex, but do not provide an objective measure of interoceptive accuracy. Electrogastrography has also been used to assess gastric activity non-invasively, with activity being linked to neural responses in interoceptive regions (Rebollo et al., 2018) and cardiac perception (Herbert, Muth, et al., 2012), for example. See Wolpert et al. (2020) for a review of electrogastrogram use.

Tests of respiratory effort have also been used to assess interoceptive ability. A respiratory resistance threshold task, for example, involves participants breathing through a circuit with varying levels of resistance, induced by filters, and determining when resistance is present (Garfinkel et al., 2016; Harver et al., 1993). Participants' ability to detect resistance is taken as a measure of respiratory interoceptive accuracy. Respiratory tasks have also been used that require participants to breathe air that varies in CO_2 concentration, and rate the frequency ('faster breathing') and volume ('deeper breathing') of their respiration using a tickbox to indicate the presence of each perturbation (Bogaerts et al., 2005; van den Bergh et al., 2004). Interoceptive accuracy is inferred from the strength of the relationship between subjective ratings and objective respiratory frequency and volume.

While the cardiac, respiratory and gastric tests described above are the most commonly used measures used to explicitly assess interoceptive ability, as a wide variety of signals are considered to be interoceptive in nature, we would suggest that a range of existing perceptual sensitivity measures are in fact also measures of interoceptive ability. These would include tests of taste (Murphy, Catmur, and Bird, 2018; Stevens et al., 1995), hunger, thirst and satiety (Harshaw, 2008), oral temperature (Guest et al., 2007), muscular effort (Herbert, Ulbrich, et al., 2007; Murphy, Catmur, and Bird, 2018), pain (e.g. de Zwaan et al., 1996; Pollatos, Dietel, et al., 2015), and (depending on one's definition of interoception; Khalsa and Lapidus, 2016) interoception may also be involved in the existing measures of perception of proprioceptive signals (e.g. Moberg, 1983; Ponzo et al., 2018, 2019), and balance (e.g. Evkaya et al., 2019). Notably, established explicit measures of interoception, such as heartbeat perception, correlate with some of these tasks. Participants with better perception of their heartbeats, for example, appear to be more sensitive to their cardiac signals, and exert less physical effort, in a free cycling situation than those with poorer heartbeat perception (Herbert et al., 2007). Individuals with higher cardiac accuracy are also more sensitive to, and less tolerant of, pain (Pollatos et al., 2012).

The addition of these tests to the existing limited battery of tests of interoceptive ability would greatly reduce the over-reliance on tests of heartbeat perception, and serve to overcome the limitations of the methods described above. Similarly, this would allow comparison of abilities across interoceptive domains, and a more reliable measure of overall interoceptive ability. There is an obvious need to assess interoceptive abilities across a range of domains, as it is thus far unclear whether or not interoception is a unitary construct, or one that consists of multiple interoceptive domains (Vaitl, 1996). Some studies have found moderate correlations between perception of different internal states, such as heartbeat and gastric distension (Herbert, Muth, et al., 2012; Van Dyck et al., 2016; Whitehead and Drescher, 1980), or between rectal distention and both cold perception and heat-induced pain

(Horing et al., 2013). Increasing hunger through fasting also appears to increase cardiac accuracy, suggesting potential overlap in the processing of these interoceptive signals (Herbert, Herbert, et al., 2012). Conversely, many studies have found non-significant relationships between performance across different interoceptive domains. Evidence suggests, for example, that cardiac and pain perception (Werner, Duschek, Mattern, and Schandry, 2009) and cardiac and respiratory abilities (Garfinkel et al., 2016; Harver et al., 1993; Nicholson et al., 2019; Pollatos et al., 2016; although see Steptoe and Noll, 1997) may dissociate. Early evidence also found that detection of high blood pressure, sweaty hands, and shortness of breath were not associated (Steptoe and Vögele, 1992). The most comprehensive investigation of multiple interoceptive domains within the same participants found no relationships between performance across the domains of cardiac, gastric, taste, pain, proprioceptive, and vestibular perception (Ferentzi, Bogdány, et al., 2018). Taken together, these findings indicate that one's performance on a single interoceptive task may not reflect one's ability to perceive or recognise other internal signals. This is consistent with evidence that intracranial stimulation of different insula regions is associated with distinct internal sensations (Stephani et al., 2011). It is also important to assess abilities across interoceptive signals as some bodily signals may simply be easier to detect than others, potentially because interoceptive signals vary highly in terms of their timescales and amplitudes (Khalsa et al., 2018). Perception of heartbeats for example, has been suggested to be easier than perception of other visceral signals (Kollenbaum et al., 1996).

As described above, each of the existing tasks assessing interoceptive accuracy has its own limitations. In addition to these task-specific limitations is a task-general problem regarding variation in signal strength. While individuals can vary in their ability to perceive a given signal accurately, signals arising within the body are also not consistent across individuals or across time within the same individual. One may perform poorly at tasks of interoceptive accuracy owing to poor perception and/ or interpretation of a signal, or owing to this signal being particularly weak. Indeed, cardiac perception accuracy has been found to increase following an increase in cardiac signal strength, for example that induced by isoproterenol infusions or exercise (Jones and Hollandsworth, 1981; Khalsa, Rudrauf, Sandesara, et al., 2009). In contrast to tasks assessing exteroceptive perception, it is very difficult to control the stimulus to be perceived in interoceptive tasks. Even in tasks where an interoceptive signal is manipulated, the experimenter can only control the stimulus inducing the interceptive change, rather than this signal change itself. Interoceptive tasks therefore tend to rely on measuring (rather than controlling) interoceptive signals objectively where possible, but the extent to which it is possible to measure internal signals varies across interoceptive domain; it is far easier to measure cardiac signals accurately than hunger or satiety, for example. This difficulty leaves the conclusions of many studies on individual differences in interoception open to interpretation, as it is possible that atypical performance is due to differences in signal strength, rather than in interoceptive accuracy. Where psychopathology is concerned, this may be a particular issue as many clinical conditions are associated with co-occurring physical atypicalities, making it feasible that interoceptive signal strength varies with mental health. Where relationships between psychopathology and interoceptive task performance are observed, therefore, these may be driven by atypical processing of the interoceptive signal, or differences in the signal itself. Future work should aim to manipulate and control across participants, or at least measure and account for, interoceptive signal strength where possible.

Beyond measures of interoceptive accuracy (sensitivity under Garfinkel and Critchely's (2013) model), a number of self-report measures have been used to assess self-reported interoception (interoceptive sensibility under Garfinkel and Critchely's (2013) model). These include the Body Perception Questionnaire (BPQ; Porges, 1993), the Multidimensional Assessment of Interoceptive Awareness (MAIA; Mehling et al., 2012), the Body Consciousness Questionnaire (Miller et al., 1981), the interoception subscale of the Eating Disorder Inventory (Garner et al., 1983), the Body Awareness Questionnaire (Shields et al., 1989), the Interoception Sensory Questionnaire (Fiene et al., 2018), and the Self Awareness Questionnaire (Longarzo et al., 2015). These measures are limited, however, by the fact that subjective perception of interoceptive attention is often confounded with the extent to which interoceptive signals are present within the individual (BPO), or by assessing multiple facets of interoception, for example interoceptive accuracy and attention (MAIA and BCQ). Note that confidence ratings are often used to quantify self-reported interoception in tasks of interoceptive accuracy (e.g., the heartbeat counting or discrimination procedures) but these are often uncorrelated with self-report questionnaires of interoception (Garfinkel et al., 2015; Murphy et al., 2020). While efforts have been made to assess the relationship between self-report questionnaire measures and objective measures of interoception, findings have varied across studies.

Notably, these findings are likely to vary depending on whether selfreport measures assess interoceptive accuracy or attention. While the majority of interoceptive questionnaire measures focus on interoceptive attention alone, or conflate interoceptive accuracy and attention (e.g. Fiene et al., 2018; Mehling et al., 2012; Shields et al., 1989), self-report measures of interoceptive accuracy have recently been developed, namely the Interoception Confusion Questionnaire (Brewer et al., 2016a) and the Interoceptive Accuracy Scale (Murphy et al., 2020). While these different self-report interoceptive accuracy measures have been found to correlate with each other, neither appears to be associated with self-reported interoceptive attention (Murphy et al., 2020). Further, while weak or non-significant correlations have been found between self-reported interoceptive attention and objectively measured interoceptive accuracy (Ferentzi, Drew, et al., 2018; Garfinkel et al., 2016, 2015; Meessen et al., 2016; Murphy et al., 2019; Whitehead et al., 1977), self-reported interoceptive accuracy appears to correlate with performance on the heartbeat counting task (Murphy et al., 2020), lending some support to the importance of distinguishing between interoceptive accuracy and attention and a 2×2 dimensional structure of interoception. We therefore recommend that future research includes both measures of interoceptive attention (e.g., the IATS; Gabriele, Spooner, Brewer, & Murphy, 2020) and accuracy (e.g., the IAS) in order to distinguish between these two dimensions of self-reported interoception. Similarly, it is necessary that future work aims to develop not only improved objective measures of interoceptive accuracy, but also objective measures of interoceptive attention (for example experience sampling procedures; Murphy, Catmur, et al., 2019).

While questions remain concerning the unitary or fractionated nature of interoception across internal signals (e.g., cardiac, gastric, and respiratory channels), it is difficult to determine whether clinical symptoms are associated with a general interoceptive impairment, or atypicalities in specific interoceptive domains. While it may be the case that interoception is a unitary ability, and that observed dissociations are due to differences in task demands, it is also possible that different interoceptive channels, or clusters of channels, exist, and that these show different associations with distinct clinical profiles. Further work is required in order to differentiate between these possibilities.

2.3. The physiological and neurological basis of Interoception

As discussed briefly above, recent definitions of interoception tend to class bodily signals as interoceptive if they are sent via lamina 1 to the AI or ACC (Craig, 2002), or via cranial nerves (vagus and glossopharyngeal) to the nucleus of the solitary tract (Critchley and Harrison, 2013). These pathways are discussed in more detail in this section. It should be noted that multiple humoral pathways also exist that convey certain interoceptive signals (thirst, blood sugar) though it is beyond the scope of this paper to discuss these in detail (see Critchley and Harrison, 2013). For a recent, more thorough, review of the neural circuitry of interoception, see Berntson and Khalsa (2021).

Small diameter A δ and C fibres innervate all tissues in the body, and project sympathetic afferents monosynaptically to lamina 1 of the spinal and trigeminal dorsal horn (the most dorsal section of the spinal cord) (Panneton, 1991; Woolf and Fitzgerald, 1983). Lamina 1 neurons are divided into physiologically, morphologically and chemically distinct classes, selective to modality, and responsive to various physical sensations (Andrew and Craig, 2001; Craig et al., 2001; Han et al., 1998; Light and Willcockson, 1999; Yu et al., 1999). From here, projections exist to the parabrachial nucleus (directly and via A1 and A2 catecholamine cell groups), and the hypothalamus via the A1 group (Craig, 2002). Parasympathetic afferents concerned with the heart and digestive tract (vagus nerve) and taste, cardiac, and general visceral sensory information (glossopharyngeal nerve), on the other hand, project to the nucleus of the solitary tract (NTS), which also projects to the parabrachial nucleus, periaqueductal grey (PAG) and hypothalamus. The parabrachial nucleus is the major brain region involved in integration of the homeostatic afferents from the wide range of interoceptive cues (Craig, 1995), and projects to the medial and basal ventral medial nuclei of the thalamus (VMb) (Krout and Loewy, 2000; Krukoff et al., 1993; Saper, 2002). Parabrachial projections also connect to the PAG, known as the mesencephalic homeostatic motor centre of the brain. The PAG is involved in guiding goal-directed activity (autonomic, neuroendocrine and behavioural) in order to maintain homeostasis (Canteras and Swanson, 1992; Saper, 2002). The hypothalamus, which is the diencephalic homeostatic motor centre, plays a similar role (Canteras and Swanson, 1992). The NTS also projects to the VMb, and a phylogenetically new pathway (only present in primates) projects from Lamina 1 neurons to thalamic nuclei; the ventral caudal portion of the medial dorsal nucleus (MDvc), the posterior ventral medial nucleus (VMpo) and the VMb. VMpo and VMb pathways project to the dorsal section of the insula (the limbic sensory cortex), which is bi-directionally connected to the ACC, OFC (which is centrally involved in processing emotion and reward for decision making (Bechara et al., 2000)), hypothalamus and amygdala. The MDvc, on the other hand, projects directly to the ACC. Both the ACC and the insula project descending connections, in order to control the homeostatic integration sites in the brainstem, as well as projecting to the AI. These projections to the AI are again phylogentically new, being specific to primates.

Whilst the posterior dorsal insula appears to provide a cortical representation of interoceptive sensations, and is activated in response to a wide range of internal signals (e.g., temperature, pain, itch, affective touch; Craig, 2002), this information is then relayed to the mid insula. The mid insula also receives information from other regions, for example the amygdala, hypothalamus and the secondary somatosensory cortex, the latter conveying non-homeostatic sensory information, and is thought to be where information is integrated (Ceunen et al., 2016; Craig, 2008). Like the posterior insula, the mid insula often shows activation in response to several internal states (Craig, 2002). However, it is in the AI that bodily states (generally parasympathetic and sympathetic in the left and right hemispheres respectively (Craig, 2005, 2002,2009)) are 're-represented', allowing for interoception (subjective experience of these states) (Craig, 2005, 2011, 2002, 2009). Projections to the ACC then allow for appropriate motor responses to the subjective feelings represented in AI to be selected and completed, in line with proposals that the AI and ACC are complimentary limbic sensory (subjective feeling) and motor (response) (Craig, 2002), or input and output (Medford and Critchley, 2010) areas, respectively. The two cortical areas tend to be activated simultaneously (e.g. Medford and Critchley, 2010), consistent with their strong anatomical (Augustine, 1996; Moisset et al., 2010) and functional (Cauda et al., 2011; Sridharan et al., 2008; Taylor et al., 2009) connectivity. The AI and ACC are implicated in a wide range of interoceptive states (Ibañez et al., 2010), including hunger and satiety (Del Parigi et al., 2002), heat (Craig et al., 2000; Olausson et al., 2005), thirst (de Araujo et al., 2003; Farrell et al., 2006), itch (Ikoma et al., 2006; Mochizuki et al., 2007), affective touch (Francis et al., 1999; Gordon et al., 2013), gastric distension of the stomach

(Stephan et al., 2003; Vandenbergh et al., 2005), distension of the bladder (Jarrahi et al., 2015), oesophagus and rectal stimulation (Coen and Gregory, 2007; Eickhoff et al., 2006; Hobday et al., 2001), taste (Kinomura et al., 1994; Small, 2010), pain (Derbyshire, 2003; Peyron et al., 2000), and fatigue (Caseras et al., 2008), as well as changes in heart rate (Critchley et al., 2000), blood pressure (Harper et al., 2000), respiration (Liotti et al., 2001; Pattinson et al., 2009), and glucose levels (Allport et al., 2004). Insula and ACC activation have also been associated with suppression of natural urges involving interoceptive processes, for example suppression of breathing and voiding (Banzett et al., 2000; Kuhtz-Buschbeck et al., 2005; Seseke et al., 2006). The insula is also involved in processing body movement and proprioception, which are classed as interoceptive states under some definitions (e.g. Cameron, 2001; Vaitl, 1996), although the posterior insula is more commonly implicated than the anterior insula (Bottini et al., 2001; Farrer et al., 2003; Farrer & Frith, 2002; Fasold et al., 2008; Ferrè et al., 2012; Karnath and Baier, 2010; Mazzola et al., 2012; Petit and Beauchamp, 2003; Zu Eulenburg et al., 2013). Crucially, the re-representation of states into a meta-representation in AI is associated with subjective awareness of these states. While the objective temperatures of stimuli (non-painful and painful) are represented in the dorsal posterior insula, for example, the subjective ratings of these stimuli correlate best with AI activity (Craig et al., 2000; Kong et al., 2006). Similarly, AI activation has been found to correlate with subjective fullness (relating to gastric distension) (Stephan et al., 2003), subjective orgasm quality in females (Ortigue et al., 2007), subjective unpleasantness of dyspnea (von Leupoldt et al., 2008), and bladder distension (Jarrahi et al., 2015), and AI activity (and volume) is associated with individuals' ability to perceive their own heart beats (Critchley, Wiens, Rotshtein, Ohman, and Dolan, 2004). This study also found involvement of the AI and ACC in interoceptive attention (when participants were attending to timing of heart beats, relative to attending to pitch of auditory tones). AI activity has also been associated with subjectively reported fatigue following inflammation (Harrison et al., 2009).

Studies of patients with insula lesions have generally observed interoceptive difficulties (see Ibañez et al., 2010 for a review). The specific location of the lesion appears to play a role in the resulting impairment, both in terms of the interoceptive domains affected, and whether deficits are observed at the level of perception or recognition (Jones et al., 2010) Further, stimulation of the insula in humans has been found to elicit unpleasant sensations in the throat, mouth and nose (Krolak-Salmon et al., 2003), changes in heart rate (Oppenheimer et al., 1992), inhibition of respiration, gastrointestinal motility, abdominal sensations and nausea (Penfield and Faulk, 1955). rTMS stimulation of the anterior insula has also been found to cause reduced interoceptive accuracy in both the cardiac and respiratory domains (Pollatos et al., 2016, but see Coll, Penton, et al., 2017). While much evidence therefore suggests that the insula is integral to consciously representing feeling states, it is worth noting that some evidence indicates that interoceptive states, such as itch, tickle, pain and temperature, as well as emotions, may be experienced even following bilateral insula lesion (Damasio et al., 2013). Damasio and colleagues therefore emphasise the role of brain stem, thalamic, hypothalamic, and somatosensory regions (alongside interoceptive cortices) in representing these states. Similarly, ACC and AI lesions do not appear to impair the perception of heart beats (Khalsa, Rudrauf, Feinstein, and Tranel, 2009), although it is possible that exteroceptive signals (for example due to vibration of the chest wall) may contribute to this ability. Studies in typical populations also question the role of the AI; for example, a recent meta-analysis observed that the posterior insula, and other regions not including the AI, were most commonly associated with attention to cardiac signals (Schulz, 2016). Such evidence suggests that further work is required to elucidate the exact mapping of neural activity to facets of interoception across interoceptive domains. Overall, it appears that the AI and ACC are involved in interoception, but subcortical and somatosensory regions also contribute to the representation of internal states, and that different areas may support different aspects of interoceptive ability (see Section 2.1). Indeed, whilst the above model appears to propose a sequential hierarchical process from spinal, vagal and glossopharyngeal afferents to the AI (with the AI supporting the perception of internal states) and ACC it should be noted that there is a large degree of interaction (or 'crosstalk') between levels, from higher to lower areas, and between channels conveying interoceptive and exteroceptive information (Critchley and Garfinkel, 2017; Critchley and Harrison, 2013). As such, it appears that multiple structures contribute towards interoception, with the anterior insula and anterior cingulate cortex playing a crucial role in the perception and conscious awareness of these signals.

3. The relevance of interoception to mental health

Interoception plays a role in a wide range of psychological functions (Quigley et al., 2021). Our central thesis is that atypical interoception is associated with impairments in a number of domains of functioning, and that such impairments characterise a broad range of psychiatric and neurological conditions. Such a conjecture has been raised previously (Barrett and Simmons, 2015; Khalsa and Lapidus, 2016; Khalsa et al., 2018; Murphy, Brewer, et al., 2017; Paulus and Stein, 2010; Quadt et al., 2018; Tsakiris and Critchley, 2016) either implicating interoception in specific disorders, or psychiatry more generally. Building on these, here we provide an in-depth overview regarding the proposed role of interoception in a range of abilities that relate to mental health. The following sections outline the role of interoception in two major domains of functioning of relevance to mental health; emotional processing and learning and decision-making.

3.1. The role of interoception in typical functioning: emotional processing

Interoception has been relatively understudied within psychology and cognitive neuroscience, certainly in comparison to exteroceptive senses such as vision and audition. The one domain in which interoception has a long history of study, however, is that of emotional processing (Pace-Schott et al., 2019). Historical debates between the James-Lange (James, 1894; Lange, 1885) and Cannon-Bard (Bard, 1928; Cannon, 1929) positions centred on whether awareness of physiological arousal (i.e. interoception of arousal) in and of itself constituted an emotion, or whether cognitive appraisal of the individual's situation was also necessary. Most modern theories are a variant on the Schacter-Singer (Schachter and Singer, 1962) model (see also Cantril and Hunt, 1932), which suggests that emotions are a combination of awareness of physiological arousal and cognitive appraisal of contextual cues. As such, interoception of arousal plays a central role in the experience of emotion, with interoception necessary for accurate detection of emotional signals, and judgements of emotional intensity, which are both necessary to identify one's emotional state (e.g. Bechara and Naqvi, 2004). The necessity of intact interoceptive ability for typical emotional functioning is further highlighted when one considers second-order effects of interoception on affective processes such as emotional memory (see Section 3.2), emotion regulation and moral reasoning. Empirical evidence supports the necessity of interoception for all aspects of emotional processing. Much of this evidence utilises an individual differences approach to demonstrate that, across individuals, interoceptive accuracy is correlated with emotional lability (Rainer Schandry, 1981), emotion regulation (Füstös et al., 2012; Kever et al., 2015), arousal focus (Barrett et al., 2004), and emotional intensity (Füstös et al., 2012; Herbert et al., 2011; Pollatos, Herbert, et al., 2007; Pollatos, Traut--Mattausch, et al., 2007; Wiens et al., 2000).

In support of a role for interoception in emotion, much evidence suggests that better perception of internal signals leads to more intense experience of one's own emotions (e.g. Ferguson and Katkin, 1996; Pollatos, Gramann, et al., 2007; Pollatos, Traut-Mattausch, et al., 2007; Wiens et al., 2000; although see Zamariola, Luminet, et al., 2019) and greater arousal focus when recounting one's emotional experiences

(Barrett et al., 2004). Higher interoceptive accuracy (rather than autonomic reactivity) also predicts decreased tolerance of (and therefore increased sensitivity to) pain (Pollatos et al., 2012), although the separate sensory and affective components of pain (Fernandez and Turk, 1992) complicate this finding. Depression, associated with less intense emotions, has also consistently been associated with reduced physiological reactivity to positive stimuli (Bylsma et al., 2008; Sloan and Sandt, 2010), which presumably makes perception of this physiological reaction more difficult. Further, specific internal states are associated with particular emotions; for example, associations exist between disgust and cardiac and gastric activity (Harrison, Gray, Gianaros, and Critchley, 2010), anger and increased heart rate and temperature (Ekman et al., 1983; Ray et al., 2008), fear and increased heart rate and blood pressure (Ekman et al., 1983; Schwartz et al., 1981), and surprise and increased skin conductance and decreased blood volume pulse (change in blood volume per heart beat) (Jang et al., 2015). In line with this, individuals tend to perceive specific emotional states as similar to specific non-emotional (interoceptive) internal states (Brewer et al., 2016a), and there is consistency among individuals in where within the body specific emotions (Nummenmaa et al., 2014) and non-emotional interoceptive signals (Nummenmaa et al., 2018) are experienced. Where interoceptive accuracy itself is concerned, as well as leading to more intense emotional experiences, evidence suggests that those with increased cardiac accuracy are more likely to physically express emotion (Ferguson and Katkin, 1996). Overall, it seems that perception of non-affective interoceptive states is associated with perfection of affective states, supporting the hypothesis that interoception and emotion are intrinsically linked.

With respect to regulation of one's own emotions, it has been argued that interoceptive accuracy is of specific benefit when individuals use a reappraisal strategy to regulate emotion. Reappraisal is an especially effective and adaptive form of emotion regulation in which negative situations are cognitively re-appraised such that they are viewed as either neutral or positive in order to limit the negative affect they induce (Gross and John, 2003). Füstös et al. (2012) reasoned that individuals with better interoception should be better able to regulate their emotions through reappraisal, owing to earlier identification of negative states (enabling earlier intervention to prevent the escalation and consolidation of negative states), and more precise individuation of emotional states (enabling more targeted, and therefore effective, intervention (Barrett et al., 2001)). These authors provided evidence for this hypothesis in two studies. They first demonstrated that both the magnitude of the P3 ERP component, which reflects the intensity of emotional experience induced by an external stimulus (Pollatos, Gramann, et al., 2007), and participants' subjective experience of emotion, were predicted by interoceptive accuracy. Their second study demonstrated that interoceptive accuracy predicted the effectiveness of emotional reappraisal in reducing the P3 magnitude, physiological arousal, and subjective experience in response to negative stimuli. Interestingly, reduced interoceptive accuracy may increase the likelihood of suppression, a less adaptive form of emotion regulation (in which emotion-consistent behaviours and/or thoughts are inhibited), owing to the fact that less perceptible interoceptive signals are likely to be easier to inhibit. The increased use of suppression strategies over reappraisal strategies is thought to lead to a variety of negative physical and mental health outcomes (Cutuli, 2014; Gross and John, 2003; John and Gross, 2004).

Beyond neurological measures, research has tended to utilise selfreport measures of emotion regulation. Reports of difficulties decreasing one's negative emotions in interviews analysed using Interpretative Phenomenological Analysis, for example, were associated with self-reported interoceptive difficulties (Zamariola, Frost, et al., 2019). Further, individuals with better cardiac interoceptive accuracy report using both cognitive reappraisal and suppression emotion regulation strategies more frequently than those with poorer cardiac accuracy (Kever et al., 2015; Pollatos, Matthias, et al., 2015). In line with this,

Weiss and colleagues found that, in a sample of typical individuals and those with multisomatoform disorders, higher interoceptive accuracy was associated with better self-reported emotion regulation in the 'frustration tolerance' and 'affect differentiation and affect tolerance' subscales of a self-regulation scale (Weiss, Sack, Henningsen, and Pollatos, 2014). A similar relationship between interoception and emotion regulation has been observed in developmental samples. In a sample of 9-16 year olds, heartbeat counting task performance was negatively correlated with self-reported maladaptive emotion regulation, but not associated with self-reported adaptive emotion regulation (De Witte et al., 2016). In 4-6 year old children, performance on an emotion regulation vignette task was also related to interoceptive accuracy scores on a cardiac perception task, although it should be noted that this child-friendly interoceptive measure reflected the degree of over- or under-estimation of change in heartrate across two conditions, so is not comparable to the more commonly used heartbeat counting task (Schaan et al., 2019a). In line with these findings are reports of emotion regulation and interoceptive difficulties within a sample of obese individuals relative to those with lower weight (Willem et al., 2019), and poorer metacognitive interoceptive awareness (although higher interoceptive accuracy) being associated with more emotional eating (Young et al., 2017). Taken together, these results suggest that interoception may play an important role in the regulation of one's own emotions. Inconsistent with this, however, is the finding that neither self-reported interoception (BAQ) nor cardiac interoceptive accuracy was associated with mood following negative mood induction, or self-reported emotion regulation (Zamariola, Luminet, et al., 2019). Notably, however, mood per se does not directly assess emotion regulation; individuals who experienced an emotion more intensely may also be more effective emotion regulators, thereby reducing the emotion to a greater extent. Future work should therefore aim to assess emotion regulation more directly, for example through change in emotional response, and its relationship with interoceptive measures.

Interoception is not only relevant to self-focussed emotional processing. Evidence also links interoceptive abilities to one's responsiveness to others' emotions. For example, Terasawa et al. (2014) demonstrated that those with better interoceptive accuracy were more likely to report experiencing emotional responses to images of others' emotional expressions. Similarly, both accuracy of expression recognition and facial mimicry of avatars' emotional expressions, which may prompt empathic responding (Bird and Viding, 2014; Coll, Viding, et al., 2017), appear to be higher in those with better cardiac interoceptive accuracy (Chick et al., 2019). In adolescents, those with good cardiac interoceptive accuracy also appear to recognise others' facial expressions of sadness and fear better than those with poor interoceptive accuracy (Georgiou et al., 2018). Further, judgements of the intensity of others' naturalistic disgusted and painful facial expressions have been positively associated with cardiac interoceptive accuracy in a female sample, although accuracy distinguishing these expressions was unrelated to interoceptive accuracy (Dirupo et al., 2020). Increased empathy in those with higher interoceptive awareness may explain why higher interoceptive accuracy has also been found to correlate positively with altruism (Piech et al., 2017). Findings have been mixed, however, with Ainley et al. (2015) finding a non-significant association between HCT performance and self-reported empathy, perspective taking, or performance on the Reading the Mind in the Eyes Task, although this task included recognition of non-emotional mental states alongside emotional states.

Subjective ratings of the intensity and valence of one's own emotional responses to others' emotional expressions have also been associated negatively with systolic blood pressure (Pury et al., 2004), while accuracy of facial expression recognition has been found to correlate negatively with systolic and diastolic blood pressure, as well as total peripheral resistance (McCubbin et al., 2011). Similarly, heartbeat evoked potentials (an electroencephalography component thought to reflect the cortical processing of heartbeat sensations (Schandry et al., 1986; for a meta-analysis see Coll et al., 2020) vary in response to different observed facial expressions (Marshall et al., 2018). Where appraisal of others' emotions is concerned, others' facial disgust and fear expressions are rated as more intense when presented at the systole (ventricle contraction) phase of the cardiac cycle than at the diastole phase (when the ventricles relax and fill with blood) (Garfinkel et al., 2014; Gray et al., 2012). Cardiac timing of accuracy feedback has also been found to influence the speed with which good heartbeat perceivers learn the names of fearful faces, with feedback at systole being more effective for learning than feedback at diastole (Pfeifer et al., 2017). For a more in depth discussion of this literature, see Critchley and Garfinkel (2015) and Garfinkel and Critchley (2016). Overall, this evidence suggests that both internal signals themselves, and one's interpretation of these signals, relate to processing of others' emotions.

Beyond behavioural responses, evidence suggests strong links between others' emotions and one's own physiological signals and neural responses in interoceptive cortex. Heart rate, for example, appears to change differentially in response to different emotions expressed by others, and these heart rate changes are associated with activation of brain regions involved in interoception, such as the amygdala, brainstem, OFC, AI, and ACC (Critchley et al., 2005; Gray et al., 2012). Similarly, responses of the AI and ACC to oesophageal stimulation appear to be affected by emotional context, whereby neural responses to oesophageal stimulation are greatest when intensely fearful facial stimuli are presented simultaneously (relative to less fearful or neutral facial expressions; Phillips, Gregory, et al., 2003). Further, neural responses to fearful faces are larger in the amygdala, insula and ACC, when the face is presented at systole than diastole (Garfinkel et al., 2014), and responses of the left OFT to disgusted faces, and the periaqueductal grey area to a range of emotional faces, are more strongly coupled to reductions in heart rate when stimuli are presented at systole than diastole (Gray et al., 2012). Non-invasive carotid artery stimulation also appears to decrease neural responses in the amygdala, hippocampus, thalamus and temporal fusiform areas while rating the intensity of fearful and neutral expressions (though increased lateral occipital and decreased temporal pole activity were observed while viewing the stimuli; Makovac et al., 2015). Further, carotid artery stimulation in this study led to higher ratings of emotional expression intensity, and this increase for fearful faces was associated with a decrease in activation in a number of brain areas including the insula, amygdala, and periaqueductal grey area. Others' emotions therefore give rise to or alter interoceptive signals in observers and vice versa. As interoception is involved in one's own emotional experience, it is likely that one's internal state when encountering another's emotion will affect one's emotional response to the other's emotion.

Findings such as these are consistent with a broad class of models that posit that one's own emotions, and by inference the awareness of one's own emotions, play a causal role in the recognition, understanding, and response to the emotions of others. Such models can be distinguished on the basis of the role that one's emotions is assumed to play in processing those of the other (developmental or 'on-line'). Under the former group of theories, infants learn to associate their own interoceptive states with their associated facial and vocal expression in others (through caregivers producing the associated expressions, for example), leading to accurate recognition of those states and an appropriate empathic response (Bird and Viding, 2014; Gergely and Watson, 1996; Heyes and Bird, 2007; Quattrocki and Friston, 2014). If interoception is impaired (either absent or noisy), then learning signals will be degraded, making associations between internal and external signals problematic, and leading to delayed or otherwise atypical learning. Within the 'on-line' group are 'shared network' and 'embodied cognition' models, in which another's emotional state is mirrored, shared, or embodied in the self, and one's own affective system is used to understand and recognise the state of the other (e.g. Barsalou et al., 2003; Gallese, 2001; Goldman and Sripada, 2005; Niedenthal et al., 2005; Preston and de Waal, 2001). Interoception is perhaps of even greater importance in these models;

mirroring another's state is futile if one cannot then perceive that state in the self, and use it to understand the state of the other.

Evidence consistent with the role of interoception in one's own affective processing also comes from neuroimaging and neuropsychological studies of interoceptive cortex. Obviously these structures are not solely concerned with interoception, and therefore evidence of affective impairment following damage to these areas can only ever suggest rather than confirm a role for interoception, but it is a useful literature to briefly survey nonetheless. The available literature from neuroimaging studies is replete with examples of activation of insula and ACC (interoceptive cortices) in affective tasks, suggesting functional overlap of the two processes. For example, both the AI and ACC are involved in processing one's own emotions (Craig, 2009; Phillips, Drevets, et al., 2003), and activated during processing of others' emotions (Calder and Young, 2005; Deng et al., 2013; Wicker et al., 2003; Zaki et al., 2012). Terasawa and colleagues identified numerous brain regions, namely the anterior insula, medial frontal cortex, lingual gyrus, temporo-parietal junction, and some brain stem regions, that were activated both in a condition where participants were asked to evaluate their current emotional state and in a condition where they evaluated their bodily signals, without an emphasis on emotion (Terasawa et al., 2012). In line with this, direct electrical stimulation of the insula has been found to increase recognition of facial expressions of anger, although this did not affect recognition of other basic emotions (Motomura et al., 2019). Fukushima et al. (2011) also demonstrated that cortical responses to one's own heartbeats also predicted self-reported empathic traits. Similarly, a meta-analysis of the literature on empathy for pain revealed that the insula and ACC are the most reliably active structures in response to another's pain. Finally, a recent multi-level kernel density analysis of fMRI studies found substantial overlap between emotional and interoceptive processing in bilateral insula, subgenual ACC, medial anterior temporal lobe, ventral mPFC, left basal ganglia, fusiform gyrus, and occipital cortex (Adolfi et al., 2017).

Lesion studies, especially studies of focal insula lesions, are not plentiful, and there is a resultant paucity of evidence for the role of interoceptive cortex. There are some reports of a reduction in the intensity of emotional experience following insula lesions (Berntson et al., 2011; Borg et al., 2013), but impairments in recognition and experience seem more common for disgust than other emotions. There is also evidence that insula lesions can result in the loss of the affective quality of pain; the patient can still 'feel' pain, but does not recognise it as aversive ('asymbolia'; Berthier et al., 1988; Greenspan et al., 1999). This may be analogous to the amusia reported after insula lesions, whereby patients remain able to hear music, but lose their appreciation of its affective qualities (Griffiths et al., 2004; Habib et al., 1995). Anterior insula lesions have also been associated with impaired recognition of pain in others, and reduced empathic arousal caused by pain in another (Gu et al., 2012). In line with this, in a meta-analysis of studies on patients with fronto-insulo-temporal lesions, patients displayed both interoceptive and emotional impairments relative to healthy control participants (Adolfi et al., 2017). There is also evidence of both emotional and interoceptive impairments in frontotemporal dementia, where insula damage is common (Hobson et al., 2019). Interoceptive cortex lesions therefore appear to impair processing of one's emotional, as well as non-emotional, internal states. It should be noted, however, that there is some evidence suggesting that there may be other routes to emotion processing; as with non-emotional interoceptive abilities, typical emotional experiences have been reported in an individual with bilateral insula damage (Damasio et al., 2013).

3.1.1. The role of interoception in emotional processing: evidence from alexithymia

Given the importance of interoception for aspects of emotional functioning, research has begun to examine the relationship between interoceptive ability and alexithymia, a subclinical condition traditionally characterised by difficulties identifying and describing feelings, difficulties distinguishing emotional states from other bodily states, and a tendency to allocate attention to external rather than internal stimuli (Apfel and Sifneos, 1979; Nemiah et al., 1976). While the cognitive aspects of alexithymia (difficulties identifying emotions) have been most commonly studied, it is worth noting that multiple types of alexithymia have been described, characterising individuals experiencing impairment at either the affective level only, where physiological signals themselves are atypical (Type III), the cognitive level only (Type II), or in both affect and cognition (Type I; Bermond, 1997; Moormann et al., 2008). Alexithymia can be seen as a neurodevelopmental condition in the majority of cases, occurring in the absence of neurological trauma, but can also be acquired following traumatic brain injury (Henry et al., 2006; Wood and Williams, 2007; Hogeveen et al., 2016). Although originally developed within a psychodynamic framework, the last few decades have seen alexithymia investigated by experimental psychologists, psychiatrists, and cognitive neuroscientists. While originally defined in terms of emotional deficits, alexithymia was first observed in individuals with psychosomatic disorders (Sifneos, 1973), and we have recently proposed that many individuals with alexithymia experience a generalised interoceptive impairment, rather than one specific to emotional processing (Brewer et al., 2016a; Brewer et al., 2016b). Indeed, alexithymia may be driven by interoceptive impairments in some individuals, and language or executive functioning impairments in others (Hobson et al., 2018, 2019; Murphy et al., 2018).

The conceptual links between alexithymia and interoception are clear; in Section 2.1 we outlined how interoception could be defined at a number of levels, and impairment at each of the conscious levels would affect emotional (as well as non-emotional) internal states, constituting alexithymia. Impaired conscious perception of a change of interoceptive state is likely to both result in, and be a result of, reduced attention to internal stimuli, and increased attention to external stimuli is thought to be a feature of alexithymia. Further, impaired conscious recognition of interoceptive signals would result in difficulties distinguishing emotional states from other, non-emotional interoceptive states such as hunger, and also result in a difficulty using interoceptive signals to distinguish between emotional states. Self-report alexithymia scales, such as the Toronto Alexithymia Scale (TAS-20; Bagby et al., 1994) and the Bermond-Vorst Alexithymia Questionnaire (BVAQ; Vorst and Bermond, 2001), therefore indirectly assess interoception, albeit specifically in the emotional domain.

There is direct and indirect evidence for alexithymia being closely associated with interoceptive atypicalities beyond the emotional domain. Most direct are studies that have employed objective tests of interoceptive abilities. A number of studies have utilised the most widely-used measure of interoceptive accuracy, the Heartbeat Counting Task, in order to investigate interoceptive ability as a function of alexithymia. Some evidence suggests that increased alexithymia was associated with decreased interoceptive accuracy; specifically that those with higher levels of alexithymia were less accurate at detecting their heartbeat (Herbert et al., 2011; Murphy, Brewer, et al., 2018; Näring and van der Staak, 1995; Shah, Hall, Catmur, and Bird, 2016). In line with this, reductions in alexithymia have been associated with increases in cardiac perception accuracy following contemplative mental training (Bornemann and Singer, 2017). While Scarpazza et al. (2017), found the opposite pattern of results (that higher alexithymia was associated with better cardiac accuracy), it should be noted that this study did not include the standard control task for the Heartbeat Counting Task, meaning results should be interpreted with caution. Additionally, some studies have not observed relationships between cardiac accuracy and alexithymia in either direction using either the heartbeat counting task (e.g. Christensen et al., 2018; Nicholson et al., 2018, 2019; Zamariola et al., 2018) or discrimination task (Mul et al., 2018; Mulcahy et al., 2019). While this evidence argues against the association between inteorception and alexithymia, some of these mixed findings may also reflect the limitations of cardiac-based measures described in Section 2.2, or pocion (e.g. Murphy, Brewer, Coll, et al., 2019; Nicholson et al.,

2018). More recently, studies have investigated the relationship between alexithymia and interoceptive tasks outside of the cardiac domain. In one study, for example, alexithymia was found to correlate negatively with perception of taste, and muscular effort and positively with increased reliance on external cues for gauging respiratory output (Murphy, Catmur, and Bird, 2018). Similarly, another study reported that alexithymia was associated with reduced perception of thermal sensation (Borhani et al., 2017). However, not all studies are consistent with a relationship between alexithymia and interoceptive accuracy; for example, a recent study did not find a relationship between alexithymia and respiratory output or cardiac perception in a sample of autistic and neurotypical adults (Nicholson et al., 2019). However, differences in task administration of the respiratory output task (in comparison to its original description; Murphy, Catmur, and Bird, 2018) limit the conclusions that can be drawn from this study. Nevertheless, some evidence suggests no relationship between alexithymia and interoception across other domains (e.g., pain, affective touch, gastric perception; (Borhani et al., 2017; Jones et al., 2005). Indeed, a recent meta-analysis noted no simple relationship between alexithymia and objectively measured interoceptive accuracy (Trevisan et al., 2019). However, the results of this meta-analysis are difficult to interpret; most included studies utilised the HCT and the inclusion or absence of a control task was not accounted for. Overall, these data emphasise a need for further study of the relationship between alexithymia and interoception across a range of populations and methodologies.

Individuals with high alexithymia also seem less able to report their state of arousal accurately (Gaigg et al., 2016; Nandrino et al., 2012; Stone and Nielson, 2001). In Gaigg et al.'s study, participants were required to report their degree of emotional arousal in response to a series of photographs, while objective measures of arousal including heart rate and galvanic skin response were recorded. Alexithymia scores predicted interoceptive accuracy, whereby increasing alexithymia predicted an increasing degree of disconnect between the subjective report of arousal level, and that measured objectively. Similarly, decoupling between self-reported emotional experience and autonomic responses has been observed in those with alexithymia (Connelly and Denney, 2007; Eastabrook et al., 2013; Martin and Pihl, 1986; Newton and Contrada, 1994; Papciak et al., 1985; Pollatos, Werner, et al., 2011; Stone and Nielson, 2001).

Beyond objective measures of interoception, individuals with alexithymia have also been found to self-report impaired non-affective interoception, although findings differ according to whether interoceptive accuracy or attention is measured. Where interoceptive attention alone is measured (using the Body Perception Questionnaire; Porges, 1993), results have been mixed, with findings suggesting no correlation (Murphy et al., 2020), or a positive correlation (Betka et al., 2018; Ernst et al., 2014) with alexithymia. Self-reported interoceptive accuracy, on the other hand, has been shown to correlate negatively with alexithymia (Brewer et al., 2016a; Murphy, Catmur, et al., 2019). Relationships between alexithymia and other measures of self-reported interoception, such as the Multidimensional Assessment of Interoceptive Awareness (Mehling et al., 2012), the Self-Awareness Questionnaire (Longarzo et al., 2015), the Interoceptive Awareness Questionnaire (van den Bergh, Bogaerts, Walentynowicz & Van Diest, 2012) and the Body Awareness Questionnaire (Shields et al., 1989), have also been observed (Longarzo et al., 2015; Muir et al., 2017; Mul et al., 2018; Zamariola et al., 2018), but as these measures include items assessing both attention and accuracy, it is difficult to interpret these relationships. In line with these mixed findings, a meta-analysis of 66 independent samples concluded that the measures used to assess interoception unsurprisingly affect the relationship with alexithymia (Trevisan et al., 2019). Although Trevisan et al. found a negative relationship between alexithymia and interoceptive ability over a range of subjective measures, a complex pattern of results emerged. Negative correlations were observed between alexithymia and both self-reported interoceptive accuracy and attention assessed by MAIA, while a positive correlation was

seen between alexithymia and self-reported attention assessed by BPQ. These different relationships may be attributable to the fact that the BPQ and MAIA may assess negative and positive (e.g. mindfulness) aspects of interoception, respectively (Mehling, 2016).

Less direct evidence in support of a relationship between alexithymia and interoception adds to these findings. For example, individuals with alexithymia are delayed in seeking medical treatment in response to acute myocardial infarction, for example (Carta et al., 2013; Kenyon et al., 1991). Similarly, those who self-report high levels of alexithymia consume caffeine (Lyvers et al., 2014), alcohol (Thorberg et al., 2009), and other substances (de Haan et al., 2014; Taylor et al., 1990) more erratically than those with low self-reported alexithymia, suggesting difficulties perceiving the effects of these substances on one's internal state. Indeed, recent evidence suggests that the relationship between self-reported interoception and alcohol consumption is mediated by alexithymia (Betka et al., 2018). Finally, alexithymia is associated with impaired abilities known to rely upon interoception, such as emotion regulation (Stasiewicz et al., 2012), emotion recognition (Grynberg et al., 2012), empathy (e.g. Enzi et al., 2016; Gleichgerrcht et al., 2015; Moriguchi et al., 2007a; Silani et al., 2008), reward processing (Bibby and Ferguson, 2011; Goerlich et al., 2016; Morie et al., 2016) and decision-making, for example in the Iowa Gambling task (Ferguson et al., 2009; Kano et al., 2011) (see Section 3.2 for a more detailed description of the role of interoception in decision-making). Notably, certain items on the TAS-20 are likely to assess interoceptive difficulties (e.g., "I have physical sensations that even doctors don't understand", "I am often puzzled by sensations in my body", "I don't know what's going on inside me").

Further support for the association between alexithymia and interoception comes from atypicalities in the structure and function of interoceptive cortex (AI and ACC) in alexithymia, which seem relatively specific to these regions, rather than diffuse across the brain (e.g. Borsci et al., 2009; Deng et al., 2013; Frewen et al., 2006; Goerlich-Dobre et al., 2014; Grabe et al., 2014; Heinzel et al., 2010; Ihme et al., 2013; Jongen et al., 2014; Kano and Fukudo, 2013; Kano et al., 2007, 2003; Moriguchi et al., 2007b; Paradiso et al., 2008; Reker et al., 2010; Silani et al., 2008; Zhang et al., 2011). Evidence for the role of the AI in particular for the ontogenesis of alexithymia was provided by a large-scale (N = 129) study of patients with traumatic brain injury (Hogeveen et al., 2016). Voxel-based lesion-symptom mapping was used to identify areas in the brain in which the degree of damage correlated with alexithymia severity. These analyses identified the degree of AI damage as a significant predictor of levels of alexithymia, with damage to the ACC approaching significance. In line with this, individuals with alexithymia also appear to exhibit atypical neural responses to interoceptive processes in these regions; higher alexithymia has been associated with greater insula and ACC activation in response to colonic distension (Kano et al., 2007), and increased insula activity but decreased ACC activity during interoceptive attention (Wiebking and Northoff, 2015). While studies have identified both positive and negative relationships between alexithymia and interoceptive cortex structure and function (Craig, 2009), likely owing to methodological differences, these regions remain the most commonly implicated in studies of the neural bases of alexithymia.

Taken together with the findings presented in Section 3.1 above, there is substantial evidence that interoception contributes to both typical and atypical emotional experience. Interoception appears to contribute to one's own emotional experience, the ability to recognise and label one's own emotions, and the ability to recognise and empathise with others' emotions. Where interoception is atypical, this has been associated with impairments in these emotional processes, including alexithymia. It can therefore be concluded that interoception is a fundamental process for typical emotion processing. Section 5.1 details how interoceptive atypicalities may, unsurprisingly, be related to the emotional difficulties that are observed in a number of clinical conditions.

3.2. The role of interoception in typical functioning: learning and decision-making

It is intuitive that learning - particularly operant conditioning, in which a behaviour is associated with punishment or reward - relies on interoception. Operant conditioning requires the individual to perceive signals pertaining to punishment and reward, which are intrinsically interoceptive. The link is especially clear when one considers that poor interoception may be characterised not just by a failure to perceive any internal signal, but also by a failure to recognise and distinguish between interoceptive signals. Miscategorisation of interoceptive information may result in 1) a noisy or inconsistent learning signal, 2) systematic misclassification of the internal source of the punishment signal (e.g. pain as hunger), or 3) classification of a specific interoceptive signal which typically elicits a negative evaluation (e.g. pain, starvation, or muscle damage), as a positive reward signal (see Bevins and Besheer, 2014 for a thorough review of such ideas).

The association of interoception with motivation and decisionmaking is also intuitive. Classic theories of decision-making hold that, given two or more choices, an organism calculates the value of each, and selects the option with the highest value (e.g. Edwards, 1954; Bentham 1784-1832; Mill, 1773-1836). For many physiologically relevant stimuli (such as food and water), the value of the outcome associated with the stimulus depends on one's present interoceptive state. The more dehydrated one is, for example, the higher the value of water. It is therefore necessary to accurately perceive one's internal state in order to determine the value of these stimuli. Even more fundamentally, value itself may be perceived as an interoceptive signal. This may be in a specific sense (such as the bodily changes associated with an attractive mate), or in a general sense of 'wanting' (Besheer et al., 2004; Bevins, 2009; Pittenger and Bevins, 2013). Furthermore, under certain theories, decision-making is guided by stored representations of the bodily consequences of stimuli and responses, such as in Damasio's (1994) Somatic Marker Hypothesis. These stored representations ('somatic markers'), provided they can be perceived, are a further source of information when calculating the value of options.

A role for interoception in learning, motivation and decision-making was recognised by Pavlov (1849–1936) in his seminal work on classical conditioning. Pavlov noted that interoceptive states can enter into associations themselves, as either conditioned or unconditioned stimuli, and can also affect the acquisition and expression of learning by acting as contextual cues or occasion setters. Such ideas have been developed by Paulus and colleagues (Paulus, 2007; Paulus and Stein, 2010; Paulus and Stewart, 2014; Paulus et al., 2009), including their relevance to various psychiatric conditions (see Section 3.5), but there has been little direct investigation of the role of interoception in learning in humans.

One study that directly assessed the relationship between interoceptive accuracy and learning and decision-making was that of Werner, Jung, Duschek, and Schandry (2009), who found that higher scores on the heartbeat counting task predicted better performance on the Iowa Gambling Task, which relies on the ability to learn which of four options are advantageous, and which are disadvantageous. Interestingly, while this result has been replicated in typical adults, the opposite pattern has been observed in those with panic disorder (Wölk et al., 2014). Findings of a positive association between interoceptive accuracy and gambling task performance provide an intriguing explanation for a group of typical individuals described by Bechara and Damasio (2002), who displayed increased autonomic arousal when selecting disadvantageous options (indicating implicit knowledge of which options produced positive and negative outcomes), but continued to select these options. It may be the case that these individuals, owing to poor interoceptive accuracy, were less able to perceive these bodily cues, and therefore less able to use them to aid their decision-making. Data consistent with this hypothesis were obtained by Dunn and colleagues (Dunn, Galton, et al., 2010), who used a modified version of the Iowa Gambling Task to demonstrate that when arousal cues favoured adaptive choices, those

with better interoception made better choices, yet when arousal cues favoured maladaptive choices, individuals with better interoception made worse choices than those with poor interoception. Similarly, cardiac interoceptive accuracy has been found to correlate positively with the size of framing effects in a gambling task (the extent to which one's decision to gamble is affected by whether identical outcomes are described as losses or gains) (Shah, Catmur, and Bird, 2016). Rejection rates in the ultimatum game (in which participants are required to decide whether to accept or reject proposals from a partner concerning division of monetary rewards) have also been found to be predicted by electrodermal responses in those with good interoceptive accuracy, but not in those with poor interoceptive accuracy (Dunn et al., 2012). One's ability to perceive interoceptive signals therefore appears to affect the extent to which internal cues are utilised in decision-making processes. Attention to interoceptive signals may also affect behaviour in the ultimatum game; listening to the timing of one's own heart beats has been found to increase feelings of unfairness in response to unfair offers, and increase the likelihood of making unfair offers to other players (Lenggenhager et al., 2013).

The relationship between interoception and specifically risky decision-making has also been investigated. Previous work demonstrated that individual differences in physiological arousal correlate with individual differences in loss aversion (the overweighting of losses with respect to equal gains) during risky decision-making (Sokol-Hessner et al., 2009). Sokol-Hessner and colleagues (Sokol-Hessner et al., 2014) predicted that the degree to which these physiological signals of arousal can be perceived (interoceptive accuracy) is likely to modulate the impact of arousal on loss aversion during risky decision-making. Results confirmed their prediction, demonstrating that individuals with better interoceptive accuracy were more loss-averse, providing direct evidence for an impact of interoception on risky decision-making. These findings are consistent with the suggestion that the increased levels of risky decision-making during adolescence (e.g. Hoorn et al., 2016; Smith et al., 2011; Steinberg, 2007) may be attributable to reduced interoceptive abilities during this stage (Murphy, Brewer, et al., 2017). A recent study found that individuals with higher self-reported interoceptive attention (measured by the BPQ) were more conservative in a task where they filled a virtual body with air to earn money, at the risk of the body being filled to explosion (leading to the loss of the monetary reward), although this effect was not observed when the virtual item being filled was a balloon (Salvato et al., 2019). The severity of the risk conveyed by the task may modulate findings, as a simple probability discounting measure of risk-taking (in which the choice is between small certain or large uncertain gains) was not associated with BPQ scores in one study (Herman et al., 2018). Interoceptive accuracy in the cardiac domain also appears to be related to risky decision-making in naturalistic scenarios; financial traders have been found to have higher accuracy than matched control participants, and interoceptive accuracy within the trader sample was positively related to profitability and years of experience (Kandasamy et al., 2016).

Further evidence for the role of interoceptive awareness during learning and decision-making was provided by Katkin et al. (2001) who showed that the acquisition of fear conditioning with masked stimuli was dependent upon interoception. When masked images were associated with the delivery of an electric shock, individuals with high interoceptive accuracy were able to use the physiological arousal response associated with the stimulus to predict the likelihood of shock, whereas those low in interoceptive accuracy were not.

Improved learning by those with high interoceptive accuracy would be expected to result in improved recall. Consistent with this suggestion, interoceptive accuracy has been positively associated with recognition memory for previously presented affective stimuli (Pollatos and Schandry, 2008; Werner et al., 2010). Additional evidence for a role for interoceptive accuracy in memory was provided by Garfinkel et al. (2013), who showed that the phase of the cardiac cycle during which a stimulus was presented was a significant predictor of subsequent memory, and that this effect was modulated by interoceptive accuracy. Similarly, higher heart rate and higher interoceptive accuracy on the HCT have been associated with performance on a prospective memory task, in which participants were required to identify targets while performing a concurrent 2-back task (Umeda et al., 2016).

Internal signal changes such as cardiac deceleration, and increased pupil dilation and skin conductance, also tend to follow (e.g. Bastin et al., 2017; Hajcak et al., 2003; Łukowska et al., 2018) and even precede (Bury et al., 2019) errors, and recent work suggests interoceptive abilities are associated with error monitoring. In Sueyoshi and colleagues' study, interoceptive accuracy, as measured by the heartbeat counting task, was associated with the degree of post-error behavioural slowing (assessed by reaction times), thought to be driven by an affective signal associated with error (Sueyoshi et al., 2014). These results indicate that, by influencing the degree to which the error-related affective signal can be perceived, interoceptive ability affects feedback-related behaviour. Furthermore, interoceptive accuracy correlated with the size of the Pe event-related potential, an electrophysiological signal reflecting the later portion of the error-related neural signal, allowing the impact of interoceptive ability to be located in time. Finally, both alexithymia and interoceptive accuracy have been associated with the degree to which one engages cognitive control to prevent errors in tasks inducing response conflict (de Galan et al., 2014).

While there is not an abundance of work directly testing the impact of interoception on learning, two bodies of work provide indirect evidence for this association, by demonstrating that common neural areas represent interoceptive information, and are involved in learning. The first of these combines computational modelling of learning with cognitive neuroscience methods, in order to map the neural correlates of learning parameters. In these studies, individuals perform a learning task, and computational models are fit to participants' data to derive learning parameters such as reward prediction errors and risk estimates. These parameters are then regressed against neural data to identify activity that covaries with learning parameters. Studies using this technique have demonstrated that activity in areas subserving interoception, including the insula and ACC, show patterns of activity consistent with the representation of risk (formally the variance associated with the outcome of a particular stimulus) and risk and reward prediction errors (Asahi et al., 2006; Mizuhiki et al., 2012; Paulus, Rogalsky, et al., 2003; Preuschoff et al., 2008; Xiang et al., 2013; Xue et al., 2010). Additionally, insula and ACC activation have been found to increase with higher degrees of uncertainty concerning the outcome of one's decision (Critchley et al., 2001), and those with higher levels of intolerance of uncertainty exhibit greater anterior insula activation when reward is uncertain (Gorka et al., 2016). Anterior insula activity has been associated with risk-free choices, and during risk-aversion mistakes (when an individual chooses not to take a risky decision, when it would have been advantageous) (Kuhnen and Knutson, 2005), and with individuals' risk-taking propensity during risky decision-making; those with higher levels of harm avoidance and neuroticism exhibited greater insula activation (Paulus, Rogalsky, et al., 2003). Decision urgency (the increasing desire to make a decision as time passes) has also been associated with insula activation (van Maanen et al., 2016). The degree of insula activation during decision-making may also be a function of interoceptive accuracy; one study found that right insula activation was associated with Iowa Gambling Task performance only in individuals who performed well on the heartbeat counting task, and not in those with poor cardiac perception (Werner et al., 2013). It is worth noting, of course, that co-activation of large, differentiated neural structures during interoception and representation of learning parameters does not necessarily indicate that the same neurons, or even populations of neurons, are involved in both processes.

A similar caveat should be applied to the second body of work, which reports the impact of damage to interoceptive cortex on learning in both human and non-human animals. A full examination of this literature is beyond the scope of this paper, but several authors have reported effects of insula lesions on learning and decision-making. Clark and colleagues, for example, reported that patients with insula lesions were less able to adjust their behaviour in a risky decision-making task, in response to changing levels of risk (Clark et al., 2008). Such an observation was supported by Weller et al. (2009), who observed that patients with insular lesions adopted lower risk decision-making strategies, even in situations in which risky decisions were optimal. Disadvantageous choices on the Iowa Gambling task in lesion patients have also been reported by Bar-On et al. (2003) although it should be noted that the patient sample was heterogeneous, with those with insula lesions being grouped with those with ventromedial and amygdala lesions. The hypothesised role of the insula in learning is also supported by non-human animal studies, in which an experimental reduction in serotonin in the anterior insula impairs conditioned taste avoidance, while reduction of serotonin in posterior insula impairs the acquisition of conditioned disgust reactions (Tuerke et al., 2012). Lesions to the rat insula have also been associated with reduced encoding of the incentive value of instrumentally conditioned behaviour to obtain food; while control rats performed fewer food-obtaining actions following low food deprivation than following high deprivation, lesioned rats did not alter their behaviour across these conditions (Balleine and Dickinson, 2000). Non-human animal studies also imply a role of the insula in risky decision-making. Ishii et al. (2012), for example, demonstrated that temporary inactivation of rat anterior insula cortex resulted in a reduction of risky choices across two separate gambling tasks. Similarly, insula lesions have been found to lead to reductions in performance on an adaptation of the Iowa Gambling Task in rats who previously performed optimally (Daniel et al., 2017).

Overall, the evidence presented in this section suggests a role for interoception in learning and decision making, across a range of tasks. As learning and decision-making impairments have been observed across a range of clinical conditions, these may again be explained by interoceptive atypicalities. Section 5.2 discusses the relationship between interoceptive atypicality and impairments in these domains, and their relevance to psychopathology.

4. Increased prevalence of interoceptive impairments in clinical conditions

Sections 3.1 and 3.2 briefly surveyed the literature linking interoception to two major domains of functioning (emotional processing, and learning and decision-making). Having provided this evidence for the importance of interoception in typical functioning, the following sections detail evidence of interoceptive impairment in several psychiatric and neurological conditions.

4.1. Evidence of interoceptive impairment in psychiatric and neurological conditions

Atypical interoception is ubiquitous across psychiatric and neurological conditions. While certain conditions have long been associated with atypical interoception, such as Feeding and Eating Disorders (EDs) including anorexia Nervosa, Bulimia Nervosa, and obesity (Eshkevari et al., 2014; Fassino et al., 2004; Kinnaird et al., 2020; Klabunde et al., 2013; Lapidus et al., 2020; Lattimore et al., 2017; Lutz et al., 2019; Pollatos et al., 2008; although see e.g. Ambrosecchia et al., 2017; Kinnaird et al., 2020), the full range of conditions impacted by interoceptive atypicalities is only just being realised (Barrett and Simmons, 2015; Brewer et al., 2016a; Brewer, Happé, et al., 2015). There is evidence of increased prevalence of atypical interoception in anxiety and panic disorders (Ehlers and Breuer, 1992; Paulus and Stein, 2010; Adrián Yoris et al., 2015), alcohol and substance abuse (Jakubczyk et al., 2019; Naqvi and Bechara, 2010; Paulus and Stewart, 2014; Paulus et al., 2009; Verdejo-Garcia et al., 2012), depression (Aaronson et al., 2017; Dunn et al., 2007; Furman et al., 2013; Harshaw, 2015; Paulus and Stein,

2010; Pollatos et al., 2009), somatoform disorders (Schaefer et al., 2012), Autism Spectrum Disorder (ASD; Garfinkel et al., 2016a; Hatfield et al., 2017; Mul et al., 2018; Nicholson et al., 2019), Attention-Deficit/Hyperactivity Disorder (ADHD; Kutscheidt et al., 2019), Obsessive Compulsive Disorder (OCD; Lazarov et al., 2010; Schultchen et al., 2019; see also Stern, 2014 for a discussion), schizophrenia (Ardizzi et al., 2016), depersonalisation/derealisation disorder (Schulz et al., 2015; Sedeño et al., 2014; but see Michal et al., 2014), personality disorders (Mussgay et al., 1999; although see Hart et al., 2013), and those with high levels of psychopathic traits (Nentjes et al., 2013). Notably, evidence for atypical interoception across each of these clinical groups has been mixed, with reports of increased, decreased and typical interoceptive abilities sometimes reported across the same condition. Such discrepancies are likely owing to individual differences in interoception within each population as well as difficulties with the measurement of interoception (see Section 2.2). The majority of studies investigating interoception in clinical groups have utilised measures of cardiac, respiratory and gastric internal signals, but evidence also exists for impairments of the perception of interoceptive signals such as taste and itch in clinical conditions (e.g. Caccavale et al., 2016; Kazour et al., 2017; Kinnaird et al., 2018).

Atypical interoception and high alexithymia have also been observed in neurological conditions with similar symptom profiles, such as Frontotemporal dementia (behavioural variant; García-Cordero et al., 2016; Salvato et al., 2018; semantic variant; Sturm and Levenson, 2011), Alzheimer's Disease (García-Cordero et al., 2016), Parkinson's Disease (Assogna et al., 2012; Ricciardi et al., 2016), and Multiple Sclerosis (Chahraoui et al., 2014; Prochnow et al., 2011). A number of medical conditions have also been associated with interoceptive difficulties (see Section 5.3.7). It is worth noting that much of the work in clinical populations has relied on measures of interoceptive accuracy in the cardiac domain, or self-reported interoceptive attention. Future work should therefore aim to characterise each population in terms of interoceptive accuracy, attention, and metacognitive awareness.

Notably, the relationship observed between interoceptive accuracy and anxiety disorders and OCD is often in the opposite direction to that observed in other conditions. While the majority of clinical conditions are associated with reduced interoceptive accuracy, interoceptive accuracy has been found to be increased in anxiety and panic disorder (for reviews, see Domschke et al., 2010; Paulus and Stein, 2010) and in OCD (Yoris et al., 2017). This finding has not been consistent, however, indicating heterogeneity among anxiety patients with regard to interoceptive abilities (De Pascalis et al., 1984: Van Der Does et al., 2000) with some recent studies suggesting that higher anxiety is associated with lower interoceptive accuracy (Ewing et al., 2017; Garfinkel et al., 2016b; Garfinkel et al., 2016). Relatedly, elevated interoceptive accuracy when observed in those with anxiety disorders may be specific to the cardiac interoceptive domain, as respiratory interoceptive accuracy appears to be reduced in these individuals (Bogaerts et al., 2005; van den Bergh et al., 2004). Potentially, cardiac signals are particularly salient for (or preferentially attended to by) individuals with anxiety or panic disorders, leading to improved performance on this task specifically, rather than generally superior interoceptive accuracy (Ehlers, 1993). Relatedly, there is debate concerning the extent to which cognitive processes in anxiety disorder contribute to improved interoception; it is possible that anxiety and panic are associated with increased preoccupation with, atypical prediction of, and/or catastrophising of, internal signals, or superior detection of prediction errors, rather than increased accuracy in detecting these signals per se (Anderson and Hope, 2009; Clark, 1986; Ludewig et al., 2005; Paulus and Stein, 2006, 2010). Consistent with this conjecture, there is evidence to suggest that anxiety is characterised by high levels of attention to internal signals (as measured by the BPO) relative to one's cardiac interoceptive accuracy (Garfinkel et al., 2016b; Palser et al., 2018). Indeed, interoceptive exposure therapy is often used clinically to reduce attention to interoceptive cues in those with anxiety and panic disorders (Boettcher et al., 2016; Boswell et al.,

2013). Alternatively, internal cues may be stronger in those with panic/anxiety (e.g., higher arousal; Hoehn-Saric and Mcleod, 2000; Paulus and Stein, 2010). It is currently unclear, however, whether this arousal follows the initial misinterpretation of the interoceptive cue, or proceeds it and therefore aids detection. The recent development of an interoceptive framework emphasising the distinction between interoceptive accuracy and attention, both where self-report and behavioural measures are concerned (Murphy, Catmur, and Bird, 2018; Murphy et al., 2020) is likely to elucidate the relationship between interoception and anxiety in future research.

Beyond behavioural studies, interoceptive cortex atypicalities are also common in clinical samples. Atypical structure and function of the insula and ACC have been observed in those with eating disorders (Frank, 2015; Uher et al., 2004), ASD (Simms et al., 2009; Uddin and Menon, 2009), OCD (Menzies, Chamberlain, Laird et al., 2008), anxiety disorders (Paulus and Stein, 2006; Rosso et al., 2010; Holzschneider and Mulert, 2011), depression and bipolar disorder (Drevets et al., 2008; Vizueta et al., 2012; L. Wang et al., 2012), schizophrenia (Bouras et al., 2001; Reid et al., 2010; Shepherd et al., 2012; Wylie and Tregellas, 2010), substance abuse/addiction disorders (Luijten et al., 2014; Ma et al., 2011; Naqvi and Bechara, 2009, 2010; Zhou et al., 2011) and PTSD (Garfinkel and Liberzon, 2009). Indeed, a large-scale meta-analysis of brain morphology across six distinct psychiatric disorders identified left and right insula and dorsal anterior cingulate as the only areas of grey matter loss common to all disorders (Goodkind et al., 2015). Again, while this evidence is only indirect, it is consistent with the idea that interoceptive deficits are a common feature of psychiatric conditions.

Notably, while a linear relationship between interoceptive impairment and disorder severity is often assumed, it may also be the case that a quadratic relationship exists, whereby either atypically low or atypically high interoceptive abilities are associated with psychopathology, or where atypical interoception may be more likely to be observed in individuals with moderate rather than severe psychopathology (Dunn et al., 2007). The nature of the relationship may also differ, of course, across clinical conditions. Further, while much research suggests an association between interoceptive abilities and psychopathology, this conclusion is based upon correlational findings. Longitudinal studies, as well as intervention studies, are required to determine whether interoceptive impairment causally affects disorder development. Similarly, whether the causal relationship between interoception and psychopathology is identical across disorders requires further investigation. As discussed in Section 7.2, the differences across disorders may be partially attributable to interoceptive impairment at different developmental stages, or in different interoceptive domains or dimensions. Whether specific atypicalities in interoception lead to particular disorders, and whether disorder symptoms can cause or exacerbate interoceptive impairment remains to be determined. It is possible, for example, that some disorders lead one to suppress internal signals; those with eating disorders may attempt to attend away from signals of hunger, thirst, and satiety, while those with depression may attempt to reduce negative affect by suppressing internal signals associated with aversive emotions. On the other hand, reduced attention towards, or accuracy of, detection of internal signals could contribute to flattened affect in depression, or the aversiveness of food restriction in eating disorders.

4.1.1. Interoceptive impairment in psychiatric and neurological conditions: Evidence from alexithymia

As alexithymia is closely associated with interoceptive impairment (see Section 3.1.1), the high number of psychological conditions that cooccur with alexithymia is also suggestive of impaired interoception across clinical populations. While this evidence is indirect, it is worthy of discussion while more direct evidence is still to be accumulated across clinical populations. Increased prevalence of alexithymia has been observed in eating disorders (Bourke et al., 1992; Cochrane et al., 1993; Jimerson et al., 1994; Rozenstein et al., 2011), ASD (Berthoz and Hill,

2005; Hill et al., 2004; Kinnaird et al., 2019), schizophrenia (Henry et al., 2010; Heshmati et al., 2010), depression (Honkalampi et al., 2001), post-traumatic stress disorder (Frewen et al., 2008; Yehuda et al., 1997), substance abuse (Dorard et al., 2008; Mann et al., 1995), somatisation/Medically Unexplained Symptoms (De Gucht and Heiser, 2003), chronic fatigue syndrome (van de Putte et al., 2007) and panic and anxiety disorders (Cox et al., 1995; De Berardis et al., 2008; Marchesi et al., 2005). Indeed, a recent study linked alexithymia to a range of self-reported psychopathological symptoms in a community sample of children and adolescents (Weissman et al., 2020). High levels of alexithymia in non-clinical developmental samples have been associated with high levels of anxiety (Sayin et al., 2007), depression and interpersonal problems (Joybari, 2014), aggressive behaviour (Manninen et al., 2011) and delinquency (Zimmermann, 2006). Notably, while the increased alexithymia levels in anxiety appear to be at odds with the increased interoceptive accuracy often observed in this population, the relationship between interoceptive abilities and anxiety appears to be complicated (see Section 4). Perhaps alexithymia is common following a range of interoceptive atypicalities, including atypically high and low interoceptive accuracy and attention. Alexithymia levels are also elevated in those with some neurological conditions, such as frontotemporal dementia (Sturm and Levenson, 2011), and medical conditions, such as HIV (McIntosh et al., 2014), diabetes (Mnif et al., 2014; Topsever et al., 2006), and Multiple Sclerosis (Bodini et al., 2008; Chahraoui et al., 2014; Prochnow et al., 2011).

Consistent with the suggestion that alexithymia is characterised by general interoceptive impairment, the clinical disorders that have been associated with poor interoception overlap largely with those that cooccur with alexithymia. It is also interesting to note that in none of the studies listed above was alexithymia observed in every member of the clinical samples studied, and not all members of the clinical samples showed atypical interoception. This suggests that interoceptive impairment is not a universal symptom of those clinical conditions with which it has been associated, and that individual differences in interoceptive ability are likely to exist in clinical populations (as they do in the typical population), and these individual differences in interoception may covary with alexithymia. It is worth noting that, while interoceptive impairments are likely to give rise to alexithymia, alexithymia may also result from linguistic impairments (Hobson et al., 2018, 2019), so evidence for interoceptive atypicalities using alexithymia measures should be treated as indirect.

5. The link between interoceptive impairment and clinical symptoms across disorders

A claim consistent with the evidence above is that interoception constitutes the 'P' Factor of psychopathology; that atypical interoception underlies various clinical symptoms seen across a multitude of psychiatric (and neurological) conditions. Of course, one does not need to agree with the existence of the P Factor to acknowledge the possible role of interoception across diagnostic and symptoms categories, it may be that different patterns of atypical interoception (heightened, reduced) across the domains (cardiac, gustatory, respiratory etc) and dimensions (accuracy, attention) of interoceptive processing result in a myriad of distinct conditions. Both positions are consistent with the evidence reviewed above: In Sections 3.1 and 3.2, we argued that interoception plays a significant role in typical cognition, particularly within the affective domain, and in learning and decision-making. In Section 4, we argued that atypical interoception is observed in a multitude of psychiatric and neurological conditions. Evidence for the link between interoception and emotional and learning and decisionmaking processes makes it plausible that at least some of the symptoms common to these conditions have a unitary, interoceptive basis, though the exact pattern of interoceptive impairments may dissociate across disorders (see Section 7.2). In the following sections, we outline existing evidence for this claim within selected domains of function. This

review of evidence is not exhaustive, nor is it intended to cover all symptom domains. Instead, it is intended to demonstrate the range of common psychological difficulties that may be (partly) caused, or exacerbated, by atypical interoception.

5.1. Interoceptive impairment and affective deficits

As described in Section 3.1, almost all aspects of emotion processing rely on interoception. One would therefore expect affective impairment in all conditions characterised by interoceptive impairment. Major Depressive Disorder is the clinical condition most closely associated with affective deficits, so it is unsurprising that depression presence and severity (Dunn et al., 2007; Harshaw, 2015; Pollatos et al., 2009) and symptoms of depression such as anhedonia (Dunn, Stefanovitch, et al., 2010; Lackner and Fresco, 2016), are associated with interoceptive impairment (though the relationship between interoception and depression may be non-linear; Dunn et al., 2007; see Sections 4.1 and 7.3). Indeed, stimulation of the vagus nerve (one of the core interoceptive neural pathways) has been successfully used as a means of treating depression (Aaronson et al., 2017). Evidence also suggests that affective impairments exist in disorders beyond depression; emotional difficulties such as emotion recognition, emotion regulation, and empathy deficits have been observed in a wide range of conditions, including Autism Spectrum Disorder (Harms et al., 2010; Harmsen, 2019; Weiss, Thomson, and Chan, 2014), ADHD (Friedman et al., 2003), Feeding and Eating Disorders (Brewer, Cook, et al., 2015; Brewer et al., 2019; Harrison, Tchanturia, and Treasure, 2010; Kerr-Gaffney et al., 2019), Schizophrenia (Bonfils et al., 2016, 2017; Edwards et al., 2002; Kimhy et al., 2016; O'Driscoll et al., 2014), post-traumatic stress disorder (PTSD; Palgi et al., 2017; Plana et al., 2014; Villalta et al., 2018), OCD (Cain et al., 2015; Daros et al., 2014; Yap et al., 2018), depression (Bourke et al., 2010; Harshaw, 2015; Schreiter et al., 2013), and Huntington's Disease (Trinkler et al., 2017). Although individuals with these conditions are more likely to exhibit emotional difficulties than typical individuals, such difficulties are not universal. Under our hypothesis, affective difficulties are more likely to be explained by the presence and severity of co-occurring interoceptive impairment, than of the disorder itself.

This hypothesis has been examined in our lab using a series of studies in which affective ability is examined in individuals with a clinical diagnosis (of ASD or ED for example) with varying degrees of alexithymia (likely an indicator of interoceptive impairment; see Section 3.1.1), and a non-clinical control group matched for alexithymia. Using this design, we have observed that impairments thought to be core characteristics of ASD, such as reduced empathic response of the insula (Bird et al., 2010) and impaired recognition of emotional facial expressions (Cook, Brewer, et al., 2013; Oakley et al., 2016), are in fact due to co-occurring alexithymia and, by inference, potentially interoceptive impairment. A similar pattern of results have been observed in a neurotypical sample, whereby alexithymia largely explained the relationship between autistic traits and facial emotion recognition (Bothe et al., 2019). The ability to produce typical emotional facial expressions (Trevisan et al., 2016), and to recognise emotion from vocal cues (Heaton et al., 2012), also appear to be explained by alexithymia, rather than by ASD presence or severity itself. Similar findings have been observed in other clinical populations; again using groups matched according to alexithymia, we found that the emotion recognition impairment often reported in individuals with EDs is due to co-occurring alexithymia, rather than ED presence or severity itself (Brewer, Cook, et al., 2015). Alexithymia has also been linked to emotion recognition deficits in a sample of adolescents with EDs (Zonnevijlle-Bender et al., 2002). Alexithymia also seems to predict emotion recognition in individuals with Huntington's Disease (Trinkler et al., 2017) and both emotion recognition and empathy deficits in those with traumatic brain injury (McDonald et al., 2011; Neumann et al., 2014). High levels of empathic personal distress have been associated with high alexithymia,

rather than ED presence or severity (Brewer et al., 2019), and empathic impairments (Maurage et al., 2011) and emotion regulation difficulties (Stasiewicz et al., 2012) appear to be predicted by alexithymia in those with alcohol addictions. Alexithymia also appears to contribute to the relationship between self-reported empathy and both autistic and schizoptypal traits in the non-clinical population (Aaron et al., 2015). It has also been suggested that high prevalence of alexithymia may account for negative experience of emotions, such as feeling 'flooded' by emotions or difficulties accepting one's own emotions (Edel et al., 2010), in those with ADHD. This may be due to alexithymia reducing one's ability to label emotional states, which has been shown to reduce the aversive nature of the state (Foland-Ross et al., 2010; Lieberman et al., 2007; Mazefsky and White, 2014). Finally, Martin (1985) hypothesised that the observed relationship between alexithymia and stress-related illness emerges as alexithymia leads to difficulties recognising that a situation is stressful, leading to prolonged exposure and therefore and increased physiological stress response.

Where interoception has been assessed using cardiac measures, it has been found to predict a number of affective abilities that are often impaired in clinical conditions, as described in Section 3.1. Interoceptive accuracy is associated, for example, with social anxiety (in anticipation of public speaking; Stevens et al., 2011), emotional lability (Wiens, 2005; Wiens et al., 2000), emotion regulation (Füstös et al., 2012; Kever et al., 2015), emotional memory (Pollatos and Schandry, 2008; Werner et al., 2010), emotional stability (Schandry, 1981), pain perception (Werner et al., 2009), and the intensity of one's emotional experiences (Füstös et al., 2013; Herbert et al., 2010; Pollatos, Herbert, et al., 2007; Pollatos, Traut-Mattausch, et al., 2007; Wiens et al., 2000). It is therefore possible that, where these abilities are atypical in clinical populations, interoceptive deficits may contribute to these impairments. While very little research thus far has examined the relationship between interoceptive abilities and emotional deficits in clinical populations, those that have support this hypothesis.

In an autistic sample, self-reported interoceptive attention (assessed by the MAIA) was associated with both self-reported and behaviourally measured empathy, and this relationship was mediated by alexithymia (Mul et al., 2018). This is consistent with the idea that atypical interoception may give rise to difficulties recognising one's own emotions, which in turn leads to difficulties empathising with others' emotions, although the correlational design prevents conclusions regarding causality from being drawn. Better metacognitive interoceptive awareness has also been associated with more accurate recognition of others' vocal emotion in autistic individuals (Mulcahy et al., 2019). Across a sample of autistic and neurotypical adults, interoceptive trait prediction error (ITPE; the relative difference between z-transformed BPQ scores and accuracy scores, both on the heartbeat counting task and heartbeat discrimination task) was negatively related to self-reported empathy (Garfinkel et al., 2016b). While ITPE was interpreted as indicating overor under-estimation of one's interoceptive abilities, this is complicated by the fact that the BPQ may be better characterised as a measure of interoceptive attention than accuracy. In this study, individuals with autism had increased interoceptive attention (BPQ scores) and decreased interoceptive cardiac accuracy, and individuals with this pattern of interoceptive tendencies had reduced self-reported empathy. Reduced interoceptive accuracy (assessed with the heartbeat counting task) and impaired recognition of facial emotion relative to typical controls have also been observed in a sample of individuals with somatoform disorder (Pollatos, Herbert, et al., 2011). Although correlations between these two measures were not reported, emotion recognition was found to correlate with heart rate variability (high frequency normalised units). Finally, individuals with schizophrenia report atypical bodily locations when asked where their emotions are experienced on a body map (Torregrossa et al., 2019). While this study did not utilise an explicit interoceptive measure, this suggests atypical links between internal representations and emotional experience. Therefore, while the relationship between interoception and emotional cognition is

under-researched, early evidence does implicate interoception in some of the emotional and social difficulties experienced within clinical populations.

5.2. Interoceptive impairment and learning, reward and decision-making deficits

Section 3.2 describes evidence suggesting a role for interoception in learning and decision-making. This work suggests that interoceptive impairment common to a number of clinical conditions may underlie reports of learning-related impairment in those populations. Learning and decision-making impairments have been observed, for example, in ASD (Friston et al., 2013; Johnson et al., 2006; Luke et al., 2011; Pellicano and Burr, 2012), Major Depression (Porter et al., 2003), schizophrenia (Waltz et al., 2007), EDs (Brogan et al., 2010), OCD (Joel et al., 2005), PTSD (Jovanovic et al., 2012), and anxiety (Castaneda et al., 2008). Empirical tests of the hypothesis that these impairments are associated with interoception in clinical groups are scarce, although some studies have addressed the issue directly. Furman et al. (2013), for example, found that among patients with depression, those reporting difficulties with decision-making had poorer interoceptive accuracy (measured by the Heartbeat Counting Task) than patients not reporting problems with decision-making, and a healthy control group. Interestingly, Shah and colleagues found that, although interoceptive accuracy was positively correlated with the size of emotional framing effects while making gambling decisions in typical individuals, this relationship was not present in autistic individuals (Shah, Catmur, and Bird, 2016). It is therefore essential that future work explicitly investigates the role of interoception in multiple learning and decision-making contexts across psychiatric and neurodevelopmental conditions.

Several authors have suggested that interoception is of central importance in drug seeking, maintenance of drug use, and withdrawal behaviours, which are thought to be a product of atypical learning and decision-making (Gray and Critchley, 2007; Naqvi and Bechara, 2009; Paulus et al., 2009; Paulus, 2007; Schmidt et al., 2013; Verdejo-Garcia et al., 2012). While recent models focus on the role of interoception with respect to craving, earlier accounts argued for a role for interoceptive states as occasion-setters, conditioned stimuli, or reinforcers (see Verdejo-Garcia et al., 2012 for a review). According to recent models, interoceptive abilities may govern the extent to which withdrawal-related states of anxiety and panic are perceived, or the extent to which the effect of certain drugs is perceived, and impacts craving (Schmidt et al., 2013). These models are supported by evidence for a role for interoceptive signals in cravings; smokers' cravings, for example, appear to be reduced by anaesthesia of respiratory airways (Rose et al., 1985, 1984).

In their review, Verdejo-Garcia et al. (2012) highlight that several models of addiction already posit a role for interoception. Under classical conditioning models of addiction, whereby drugs and drug-related stimuli act as conditioning stimuli for the hedonic, withdrawal, and craving aspects of drug use and addiction, individuals with high interoceptive abilities may be more likely to form associations, or be less able to ignore cravings, than those with poor perception of interoceptive signals (see also Verdejo-Garcia and Bechara, 2009; Navqi and Bechara, 2010). It has also been argued that typical individuals who are better at predicting interoceptive sensations are better at regulating their cravings for unhealthy food (Kruschwitz et al., 2019; although results may be due to memory for interoceptive states in this study rather than interoceptive predictions). Another prominent model highlights how the change in addictive behaviour over time may result in part from interoceptive changes (Paulus et al., 2009). Central to this idea is that the reward value of a stimulus (how pleasurable it is perceived to be), is a function of the effect it has on the body's current internal state, relative to an ideal state (Cabanac, 1971). It is claimed that drug addiction results in alterations of the body's current internal state, and the representation of the ideal body state, such that the hedonic effects of drugs

become less intense, and the craving and withdrawal effects become more intense, resulting in drug use changing from being impulsive to being compulsive. Goldstein et al. (2009) extend this approach, highlighting how interoceptive dysfunction may impair detection and recognition of interoceptive signals, which in turn affects emotional awareness, and potentially explains the tendency for some addicts to deny their addiction. While all of these models assign important roles to interoception, as highlighted by Vardejo-Garcia et al., a number of outstanding questions remain. For example, the mechanism underlying the relationship between interoception and addiction is unclear, especially as both high and low interoceptive abilities have been observed. While it is possible that a quadratic relationship between interoceptive ability and addiction susceptibility exists (whereby low interoceptive accuracy leads to misidentification of substances as more rewarding than is typical, and high interoceptive accuracy and/or attention lead to stronger craving sensations, for example), it may also be the case that interoceptive abilities change, or play different roles, across the course of addictions. For example, whilst poor interoceptive accuracy has been implicated in the initial development of addiction, it has also been suggested that high interoceptive accuracy is involved in the maintenance of addition (Verdejo-Garcia et al., 2012). It is also likely that the relationship between interoception and addiction varies depending on whether attention or accuracy is measured. Therefore, it is important for future research to consider whether it is the attention towards interoceptive signals, the recognition of such signals, or their regulation, that is of most importance for addiction, as well as the role of interoception in the vulnerability for, development of, and maintenance of addiction.

Research into the impact of insula lesions provides evidence consistent with the hypothesis that interoception underlies craving in addiction and learning-related abnormalities in other disorders. It should be acknowledged that the fact that interoception relies on the insula, and that the insula is damaged in these studies, does not necessarily mean that interoception itself is impaired. However, the studies may be informative as to the impact of atypical interoception. The seminal finding in the human literature was that of Navqi and colleagues (Naqvi et al., 2007), who demonstrated that insula lesion patients found it easier to stop smoking, reporting that they no longer experienced cravings, or that cravings were greatly reduced following their lesion. This finding was replicated in prospective studies by Suner-Soler et al. (2012) and Gaznick and colleagues (Gaznick et al., 2014), although not in a shorter prospective study (Bienkowski et al., 2010). Similarly, inactivation of the insula in rats reduced drug-seeking behaviour (Contreras et al., 2007). That the insula is the neural basis for reward and cravings (Paulus, 2007) is further supported by the results of several imaging studies, which demonstrate that insula activity is correlated with craving-associated stimuli (Brody et al., 2002; Dom et al., 2005; Kilts et al., 2001, 2004; Naqvi and Bechara, 2009), and craving intensity (G.-J. Wang et al., 1999). Similarly, insula and anterior cingulate activation have been found to be atypical in individuals with addiction disorders during decision-making and risk-taking (Gowin et al., 2014; Hoffman et al., 2008; London et al., 2005; Nestor et al., 2011; Paulus, Hozack, et al., 2003). Animal studies also support the role of the insula in addiction. Contreras et al. (2007), for example, demonstrated that inactivation of the insula in rats interfered with a previously learned place preference associated with amphetamine delivery. Interestingly, insula lesions are also linked to a reduction in the cognitive distortions linked to pathological gambling. While typical individuals exhibit an increase in gambling motivation in response to near-misses, and overestimate the frequency of one of two equally probable outcomes after a run of the alternate outcome, these effects are increased in those with problematic gambling behaviour (Fortune and Goodie, 2012; Michalczuk et al., 2011). Patients with insula lesions, however, do not show these gambling-related cognitive distortions (Clark et al., 2014). This effect is consistent with neuroimaging work identifying insula involvement in near-miss situations (Akitsuki et al., 2003; Clark et al., 2009; Shao et al., 2013; Xue et al., 2010), and the involvement of the insula in

risky decision-making reported in Section 3.2.

As argued in Section 3.1.1, we suggest that alexithymia may be indicative of a general failure of interoception, and there is an increased prevalence of alexithymia in almost all psychiatric conditions (see Section 4.1.1). Alexithymia research may therefore add to our understanding of the relationship between interoception and reward and decision-making, where addictive behaviours are concerned. In individuals with alcohol addiction, for example, alexithymia is positively associated with duration of alcohol use and scores on the Michigan Alcoholism Screening Test (Uzun, 2003), and use severity (Bruce et al., 2012; Thorberg et al., 2009), including severity of alcohol-related behaviours following treatment (Cleland et al., 2005). Similarly, alexithymia predicts faster relapse, and lower levels of engagement with treatment in individuals with addiction disorders (Cleland et al., 2005; Stasiewicz et al., 2012). Further, Speranza and colleagues observed that, in individuals with Eating Disorders or Addictive Disorders, high degrees of alexithymia predicted high levels of depression, which in turn predicted addictive behaviours (Speranza et al., 2004). Alexithymia is also positively related to craving in alcoholics (Junghanns et al., 2005), smokers (Sutherland et al., 2013) and methamphetamine addicts (Saladin et al., 2012). This may be due to increased interoceptive attention in alexithymia (Betka et al., 2018; Christensen et al., 2018; Ernst et al., 2014; Longarzo et al., 2015; Zamariola et al., 2018), although findings have been mixed.

A small but growing literature exists on the impact of alexithymia on more explicit learning and decision-making. Individuals with alexithymia, for example, have been found to make riskier choices on the Iowa gambling task than those with low alexithymia levels, especially when less information was available concerning their previous cumulative performance (Ferguson et al., 2009). Kano and colleagues conducted a PET study and found that those with alexithymia made less advantageous choices on the same task, and exhibited reduced interoceptive cortex (insula and ACC) activation during the task (Kano et al., 2011). Alexithymia also predicted performance on a version of the Iowa Gambling Task in pathological gamblers (Aïte et al., 2014), although not in a sample of patients with de novo Parkinson's Disease (Poletti et al., 2011). In Shah, Catmur, and Bird (2016) study reported above, higher alexithymia was associated with smaller emotional framing effects in a gambling task, but this was only observed in the typical, and not the ASD, sample. Recent evidence also suggests that individuals with alexithymia are more likely that those without alexithymia to select immediate smaller rewards relative to delayed larger rewards, although this tendency was also associated with higher interoceptive accuracy, making interpretation of the underlying mechanisms difficult (Scarpazza et al., 2017). Finally, alexithymia appears to be associated with weaker fear conditioning, indicated by lower skin conductance in response to conditioned stimuli, and faster extinction of this response in those with high, relative to those with medium and low, levels of alexithymia (Starita et al., 2016). While the literature on learning and decision-making in alexithymia is highly heterogeneous in terms of methodology, evidence thus far implicates alexithymia in difficulties with these processes, potentially due to atypical interoception.

5.3. Other domains potentially affected by interoceptive impairment

Sections 3.5.1 and 3.5.2 have described the way in which atypical interoception is associated with common impairments in clinical populations, with a focus on the emotional domain, and learning and decision-making. Notably, however, a number of other symptoms frequently observed in clinical groups may also be explained by interoceptive impairment. These are addressed in the following sections.

5.3.1. Sensory processing

Atypical sensory processing may be explained by interoceptive impairment, and is particularly relevant to ASD. Atypical sensory responsivity to the environment is now included as a diagnostic criterion

for ASD in the DSM-5, and individuals with ASD often experience these sensory atypicalities (Ben-Sasson et al., 2009; Bettelheim, 1967; Crane et al., 2009; Green et al., 2016; Kanner, 1943; Lane et al., 2014, 2010; Leekam et al., 2007; Marco et al., 2011; Tomchek and Dunn, 2007; Zachor and Ben-Itzchak, 2014). Many of these sensory domains can be considered interoceptive; ASD is associated with hyper- or hypo-sensitivity to taste, smell, and tactile stimulation (of which affective touch is interoceptive, as it is carried by interoceptive C fibres (Vallbo et al., 1999; Wessberg et al., 2003)); see DuBois et al. (2016) and Hatfield et al. (2017) for reviews. Atypical processing of affective touch may also be associated with hyper- or hypo-sensitivity to particular fabrics or textures. Findings concerning sensitivity to tactile stimuli have been relatively inconsistent across ASD samples (Cascio et al., 2008; Tanidir et al., 2007; Tavassoli et al., 2006), in line with the suggestion that they arise due to impaired interoception (present in some but not all individuals with ASD), rather than ASD itself. Those with ASD also sometimes exhibit hypersensitivity to exteroceptive signals, such as visual and auditory stimuli (Green et al., 2016; Lane et al., 2014, 2010; Leekam et al., 2007: Marco et al., 2011; Tomchek and Dunn, 2007). While these stimuli are not interoceptive, it is possible that poor interoception in some individuals with ASD leads to relative increases in the salience of exteroceptive stimuli, and therefore hypersensitivity. Alternatively, it may be the case that increased arousal in response to salient exteroceptive stimuli, such as intense light stimuli, may be incorrectly classified as interoceptive signals, such as pain.

It is possible that sensory atypicalities contribute to some of the core symptoms of ASD (Quattrocki and Friston, 2014), as sensory atypicalities have been associated with restricted and repetitive behaviours (Chen et al., 2009; Foss-Feig et al., 2012) and need for sameness (Wigham et al., 2015), as well as social and communication deficits (Foss--Feig et al., 2012). Similarly, Ben-Sasson et al. (2009), found that sensory responsiveness is associated with overall ASD symptom severity. Indeed, a recent study specifically investigated the relationship between interoceptive abilities and core features of ASD. Palser et al. (2019) found that social affective difficulties in autistic children were related to overconfidence in the heartbeat discrimination task, while restricted and repetitive behaviours were associated with higher heartbeat discrimination accuracy. This latter result may be driven by task features, such as the requirement to attend to multiple, repeated stimuli in the heartbeat discrimination task, which may be linked to restricted interests and repetitive behaviours. Alternatively, these findings may indicate a complex relationship between restricted interests/repetitive behaviours and the balance between interoceptive and exteroceptive attention or accuracy, and the extent to which these cues are integrated. Relatedly, autism appears to affect HCT performance differently at different intervals, with autistic children numerically outperforming neurotypical children at long intervals (Schauder et al., 2014). Autistic traits may, therefore, be linked to particularly high or low interoceptive abilities in different task contexts. Taken together, these studies suggest that interoception and sensory atypicalities may link specifically to core autistic symptoms. As sensory atypicalities are not universal in those with ASD, however, interoceptive deficits likely represent one of many potential risk factors for ASD traits.

Although sensory hyper- and hypo- responsiveness are more common in ASD than in other neurodevelopmental disorders (Baranek et al., 2006; Ben-Sasson et al., 2009; Leekam et al., 2007), it is worth noting that sensory atypicalities have also been observed in individuals with intellectual disability and other neurodevelopmental disorders (Green et al., 2003, 2016; Lane et al., 2012; Tomchek and Dunn, 2007; Watling et al., 2001), with a systematic review suggesting that atypical olfaction may be a biomarker for a range of psychiatric disorders (Schecklmann et al., 2013). Consistent with the hypothesis that poor interoception is associated with a range of clinical symptoms, it is possible that interoceptive impairment contributes to atypical sensory processing across multiple clinical groups.

5.3.2. Pain sensitivity

Beyond sensory responsiveness in developmental disorders, hyperand hypo-sensitivity to pain have been reported in a number of clinical populations, such as depression (Hermesdorf et al., 2016), schizophrenia (Stubbs et al., 2015), eating disorders (Papezová et al., 2005; Yamamotova et al., 2017), autism (Riquelme et al., 2016), and OCD (Hezel et al., 2012). As pain is, by most contemporary definitions (Craig, 2002; Khalsa et al., 2018), an interoceptive signal, these atypicalities are unsurprising within the context of our proposal. It is likely that widespread interoceptive impairments across the clinical population contribute to atypical perception and recognition of pain in multiple psychiatric conditions. While pain sensitivity in individuals with chronic pain conditions has been associated with alexithymia (Glaros and Lumley, 2005; Lumley et al., 2002), little research has investigated the relationship between pain hyper- and hypo-sensitivity and other interoceptive difficulties in these clinical groups. In individuals with medically unexplained symptoms (painful sensations in the absence of an organic underlying cause), cardiac interoceptive accuracy has been negatively associated with symptom severity (Schaefer et al., 2012), and symptoms appear to improve following cardiac perception training (Schaefer et al., 2014). This population has also been hypothesised to exhibit increased interoceptive attention (Barsky, 1992), although this hypothesis has not been empirically tested.

5.3.3. Repetitive behaviours and compulsions

Repetitive behaviours, which are a core diagnostic feature of both ASD and OCD, are also common in Eating Disorders (e.g. repeated bingeing and/or purging behaviours) and addiction disorders (repeated addictive behaviours), and may be explained by atypical interoception. It is possible that difficulties perceiving or recognising the aversive consequences of maladaptive behaviours, and associating these consequences with the action, contribute to their maintenance. As behaviours are frequently learnt through conditioning processes, atypical processing of reward and punishment signals that are associated with a given behaviour means that those with poor interoception will be slower or less able to learn to repeat rewarded behaviours, and avoid punished behaviours. In the case of compulsions, it is possible that individuals with poor interoception misattribute the source of negative internal states, and experience an urge to perform compulsions in an attempt to reduce these signals. If, for example, one is preoccupied with cleanliness and contamination, one may classify any internal state of anxiety as arising due to cleanliness being compromised, and compulsively clean in order to reduce the aversive internal state. Forming inaccurate associations between internal states and their external triggers may, therefore, maintain compulsive behaviours. Equally, it is possible that these behaviours are explained by reduced recognition of the reward signals typically associated with novelty (e.g. Bardo et al., 1996; Pierce et al., 1990), causing the individual to engage in more predictable, and less novel, familiar, behaviours. Compulsions and repetitive behaviours may also arise owing to difficulties suppressing urges, which may be perceived as interoceptive signals themselves. Indeed, suppression of natural urges such as breathing and voiding involves interoceptive cortex activation (Banzett et al., 2000; Kuhtz-Buschbeck et al., 2005; Seseke et al., 2006). Multiple disorders are associated with difficulties suppressing urges, such as OCD, tics and Tourette syndrome, hyperactivity and impulse control disorders, and addictions, and these conditions have been associated with atypical interoception, as described above. Interestingly, the premonitory urge to tic in individuals with Tourette Syndrome has been found to be positively associated with cardiac interoceptive accuracy and self-reported interoceptive attention (Ganos and Garrido, 2015; Rae et al., 2019). While the specific nature of the relationship may differ across populations, interoceptive atypicalities may account for repetitive behaviours, compulsions, and difficulties suppressing urges across a range of clinical conditions.

5.3.4. Self-other distinction

Recent theoretical work has suggested that interoception (in combination with exteroception; see Park and Blanke, 2019; Park and Tallon-Baudry, 2014) is crucial for developing an accurate representation of the self (Seth, 2013; Quattrocki and Friston, 2014). The implication that one's own interoceptive experiences are being experienced by another body, for example, has been found to produce illusory distortion of one's self-representation (e.g. Aspell et al., 2013; Botvinick and Cohen, 1998). Deficits in self-other distinction (the ability to represent the self as a distinct individual, separate from others) are observed in some clinical groups, and may again be explained by interoceptive deficits. The Rubber Hand Illusion of body ownership, for example, in which individuals experience a rubber hand as their own following stroking of their own hand while viewing simultaneous stroking of the rubber hand, is weaker in those with high levels of interoception, arguably due to more accurate representations of the self (Suzuki et al., 2013; Tsakiris et al., 2011; although see Crucianelli et al., 2018). Similarly, representations of one's own face appear to be more malleable (assessed by the enfacement illusion; Sforza et al., 2010) in those with poor interoceptive accuracy (Tajadura-Jiménez and Tsakiris, 2014). Furthermore, lesions to the right posterior insula (an area associated with attention to interoceptive sensations; Schulz, 2016) have been associated with somatoparaphrenia (in which one's limbs or other body parts are not perceived as belonging to oneself). Taken together, these findings suggest that individual differences in interoception are associated with those in self-other distinction.

Where clinical groups are concerned, a reduced sense of self and distorted boundaries between representations of the self and others have long been implicated in, and are core features of, disorders such as schizophrenia (van der Weiden et al., 2015) and depersonalisation/derealisation disorder (American Psychiatric Association, 2013). A number of other psychological conditions have also been associated with atypical representations of self and other, however, and it is possible that interoceptive differences account for these atypicalities and a range of related symptoms/behaviours across various clinical groups. It is possible that interoception-related atypicalities in self-other distinction, for example, explain the delusions of control and auditory hallucinations experienced by some individuals with schizophrenia, bipolar disorder, and depersonalisation/derealisation disorder (American Psychiatric Association, 2013). Delusions and hallucinations such as these are likely to arise owing to poor self-other distinction, such that one's own behavioural commands are not represented accurately, leading to reduced precision of predictions concerning the consequences of these motor commands, and the perception that these behaviours were generated by an external agent (Frith, 2012; Seth et al., 2012). Evidence suggests, for example, that individuals with auditory hallucinations and passivity experiences perceive self-generated tactile sensations to be as intense as other-generated sensations, while self-generated stimuli are perceived as less intense than other-generated sensations by control participants (Blakemore et al., 2000), likely due to reduced attenuation of sensory information, caused by reduced precision of predictions based on motor efferents. Similarly, those with schizophrenia experience the Rubber Hand Illusion more strongly than those without (Peled et al., 2000), and exhibit atypical somatosensory evoked potentials during this illusion (Peled et al., 2003), although see Cornielle and Lush's (2021) recent discussion of the potential role of demand characteristics in this task. Recognition of images of the self also appears to be impaired in schizophrenia (Ferroni et al., 2019; Sandsten et al., 2020).

Further, contemporary models proposing that accurate interoception is crucial for representing the self as distinct from others (Quattrocki and Friston, 2014; Seth, 2013), suggest that poor interoception would lead to unintended mirroring of others, for example echopraxia and echolalia (Quattrocki and Friston, 2014). While evidence has been mixed, many have argued for atypical imitation in individuals with ASD (see Hamilton, 2013 for a review). Recent evidence suggests that autistic individuals have a tendency to over-imitate (Sowden et al., 2015), in line with increased prevalence of echolalia and echopraxia in this population (Baltaxe and Simmons, 1975; Cunningham and Dixon, 1961; Grossi et al., 2013; Jordan, 1993; Kanner, 1946; Koegel et al., 1982). Consistent with the high rates of alexithymia and interoceptive impairment in ASD, a relationship has been observed between both interoceptive accuracy (Ainley et al., 2014) and alexithymia (Sowden et al., 2016), and the ability to inhibit automatic imitation in typical individuals. Poor interoception, and in turn poor self-other distinction may, therefore, explain atypical imitation in those with ASD. Notably, echolalia and echopraxia have also been observed in a range of clinical groups, including those with aphasia (Grossi et al., 1991; Trojano et al., 1988), Tourette syndrome (Eisenberg et al., 1959; Michael, 1957) and intellectual disability (Critchley, 1964), and children who later develop schizophrenia (Bearden et al., 2000).

Beyond echolalia and echopraxia, reduced distinction between the self and other may also be responsible for the increased levels of emotion contagion and personal distress often observed in clinical groups (overimitation in the affective domain). Emotion contagion (sharing another's emotion) often follows observation of an emotional state in another, and can lead to personal distress if this 'mirrored' internal state is negative. Intact ability to distinguish between the self and other then allows the observer to attribute the emotional state to the observed individual, thereby reducing personal distress, and giving rise to empathic concern (Bird and Viding, 2014; Coll, Viding, et al., 2017; Davis, 1980; Decety and Lamm, 2006; Moriguchi et al., 2007b). Interestingly, personal distress (a marker of emotion contagion) is often increased in clinical conditions, while other facets of empathy (e.g., empathic concern) are often decreased in these groups. Self-reported personal distress has been found to be atypically high, for example, in those with eating disorders (Beadle et al., 2013; Duchesne et al., 2012; Guttman and Laporte, 2002), schizophrenia (Bonfils et al., 2017), Tourette syndrome (Eddy et al., 2015), OCD (Kang et al., 2012), and depression (Banzhaf et al., 2018). As interoception is likely to play a role in self-other distinction, increased personal distress (due to decreased self-other distinction) across multiple clinical populations may arise owing to interoceptive difficulties in these groups.

5.3.5. Further social impairments

Social impairments (beyond emotion recognition and empathy difficulties) are common in many clinical populations, with ASD being the most notable; difficulties with social interaction and communication are a core diagnostic feature of ASD (American Psychiatric Association, 2013). One social impairment that has been consistently documented in those with ASD is reduced attention to social stimuli, such as faces (Dawson et al., 2005; Klin et al., 2002; Nakano et al., 2010; Riby and Hancock, 2008). Quattrocki and Friston (2014) attributed this to interoceptive impairment, suggesting that if one struggles to perceive the interoceptive cues associated with a face, such as warmth and satiety during infancy, one will not learn the affective value of faces, and therefore experience reduced motivation to attend to them. Further, direct eye gaze has been described as inducing a range of negative internal states, such as pain, dizziness, headaches, nausea, and overheating (Trevisan et al., 2017), which may be due to misclassification of these associated internal cues. Decreased motivation to interact with others is also likely to exacerbate the communication difficulties experienced by those with ASD, via reduced opportunity for receiving social feedback on one's communication.

A further social difficulty that is often observed in ASD is Theory of Mind (ToM) or mentalising, referring to the ability to infer another's mental states (e.g. thoughts, feelings, and beliefs) (Brunsdon and Happé, 2014; Frith et al., 1991). Ondobaka and colleagues proposed that one's understanding of one's own internal states is likely to assist in inferring others' internal states, which in turn may assist with determining their mental state (e.g. the observed individual is cold, therefore they want to go indoors) (Ondobaka et al., 2017). Empirical evidence relating ToM

abilities to interoceptive abilities is scarce, but Shah et al. (2017) tested this hypothesis in typical individuals using the Movie for Assessment of Social Cognition, a commonly utilised ToM task. While they found that cardiac interoceptive accuracy was positively associated with performance on items relating to others' emotional states, it was unrelated to performance on ToM items that did not require emotional inference. Notably, this task does not include items that require inference of non-emotional internal states (mental states to be inferred were either emotional or unrelated to interoceptive signals). Further evidence relating to the role that interoception may play in ToM is clearly required, but it is possible that interoceptive abilities are involved in inference of others' states where interoceptive or emotional signals are involved, but not when mental states do not relate to internal (including emotional) signals. While this theory has not been tested in clinical groups (beyond emotion recognition), interoceptive difficulties may explain some of the difficulties with inference of others' internal states in a range of populations.

Beyond ASD, social difficulties have been observed in a number of psychological disorders, such as eating disorders (Cardi et al., 2018), schizophrenia (Green et al., 2015), bipolar disorders (Wolf et al., 2010), and depression (Romera et al., 2010). These may stem, at least in part, from the emotional impairments that are common in these populations. In individuals with Eating Disorders, for example, alexithymia predicts decreased self-reported social skills (Beales and Dolton, 2000). More research is required, however, into the nature of the relationship between interoception, emotional abilities, and social abilities outside of the emotional domain, especially relating to the causal nature of relationships across disorder populations. If interoceptive impairment follows development of a given disorder (as may be the case with FTD and AIDS, for example), it remains to be determined whether the disorder causes both interoceptive difficulties and social difficulties, or whether interoceptive impairment leads in turn to further emotional and social deficits. Similarly, it is unclear whether interoceptive impairment contributes directly to the development of both psychopathology and emotional deficits, or whether interoception leads to further emotional and social difficulties, and in combination these impairments lead to disorder development. It is likely that there are multidirectional relationships between interoception, psychopathology and socio-emotional abilities, but these may vary according to clinical diagnosis, making investigation of these associations across populations and across time necessary.

5.3.6. Eating behaviours

One area of psychopathology in which the role of interoception is clear is atypical eating behaviours. Clearly, difficulties detecting signals of hunger and satiety, and distinguishing them from other internal cues, would lead to difficulties determining appropriate quantities of food (Bruch, 1973). This is consistent with Herbert, Blechert, Hautzinger, Matthias, & Herbert's (2013) finding that those with higher interoceptive accuracy show more intuitive eating, an adaptive form of eating in response to internal cues of hunger and satiety. While the relationship between reduced or increased eating and perception or recognition of hunger and satiety cues is intuitive, atypical eating also refers to behaviours such as avoidance of particular tastes, flavours, and textures, and abnormal eating rituals and routines. While eating atypicalities are most strongly associated with Feeding and Eating Disorders (American Psychiatric Association, 2013), they are also common in individuals with ASD, (Martins et al., 2008; Schreck et al., 2004), and depression, where changes in weight and appetite are a diagnostic criterion (American Psychiatric Association, 2013). Notably, while ASD and depression frequently co-occur with EDs (Marucci et al., 2018), many individuals with ASD or depression without co-occurring EDs still exhibit atypical eating behaviours that do not necessarily stem from weight and shape concerns or difficulties processing hunger or satiety, such as dislike of particular food textures or flavours, and eating rituals in ASD (Sharp et al., 2013; Spek et al., 2019), or appetite changes in

depression, potentially linked to reduced reward (Coccurello, 2019). Reward-processing models of EDs have also been proposed; whereby food may be less rewarding for individuals who restrict calorific intake, but strongly associated with reward in those who binge eat (see Monteleone et al., 2018 for a review). Relatedly, atypical eating behaviour has also been related to emotional difficulties; emotional overeating is characterised by eating in response to negative emotions, potentially due to poor distinction between internal cues of hunger and of other states including emotions (Koch and Pollatos, 2014). Indeed, Koch and Pollatos (2014) found that emotional overeating is associated with interoceptive accuracy in overweight children, although in this longitudinal study emotional overeating predicted cardiac accuracy a year later, rather than vice versa, suggesting a complex and potentially bidirectional relationship. Importantly, taste, smell, and reward cues, can be seen as interoceptive signals, so the relationship between interoception and eating may be driven by processing of these cues, in addition to hunger, thirst and satiety signals.

5.3.7. Physical health

As described above, much research suggests a link between interoception and mental health. While physical health has received less attention in this field, interoception may play a role in the development of physical health conditions. Poor interoceptive accuracy has been observed, for example, in individuals with asthma (Khosravani et al., 2016), diabetes (Pauli et al., 1991), and congenital heart disease (Rietveld et al., 2004). Heartbeat evoked potentials following resuscitation after cardiac arrest have also been found to be larger in survivors than non-survivors at 6 month follow-up (Schulz et al., 2018), and correlate with cardiac responses in patients with cardiac dysfunction (Gray et al., 2007). While interoception has not been specifically examined in a wide range of physical illnesses, the fact that high levels of alexithymia have also been observed in conditions such as human immunodeficiency virus (HIV) (McIntosh et al., 2014), diabetes (Mnif et al., 2014; Topsever et al., 2006), multiple sclerosis (Bodini et al., 2008; Chahraoui et al., 2014; Prochnow et al., 2011), and cancer (although evidence is limited; De Vries et al., 2012) raises the possibility that interoceptive deficits may be common across a range of conditions. Poor interoceptive accuracy has also been linked to being overweight (Herbert and Pollatos, 2014). Interoception also appears to be associated with health-related decision-making, such as intuitive eating (Herbert et al., 2013) and regulation of physical load during activity (Herbert et al., 2007).

Research is lacking into whether interoception explains physical health problems that are common in those with psychological conditions, but several conditions warrant investigation. In particular, gastrointenstinal difficulties are often observed in ASD (Klukowski et al., 2015; Wakefield et al., 2000; White, 2003; Williams et al., 2011) and irritable bowel syndrome commonly co-occurs with anxiety and depression (Fond et al., 2014; Tosic-Golubovic et al., 2010). It is possible that a variety of physical health symptoms arise due to interoceptive difficulties in clinical populations.

5.3.8. Sleep

Disrupted sleep is commonly reported across a variety of psychiatric conditions, such as ASD (Elrod and Hood, 2015; Klukowski et al., 2015), depression (Alvaro et al., 2013; Shanahan et al., 2014), anxiety (Alvaro et al., 2013; Shanahan et al., 2014), OCD (Díaz-Román et al., 2015), PTSD (Kobayashi et al., 2007), and schizophrenia (Chan et al., 2017). It is likely that interoception plays a role in sleep, as the vagus nerve (a core pathway for the transmission of interoceptive information; Craig, 2002, 2009) plays a key role in sleep and wakefulness (Leichnetz, 1972; Peñaloza Rojas, 1964), and sleep is accompanied by changes in bodily physiology, such as heart rate, blood pressure, and temperature (see Orem and Barnes, 2012). Whilst a new proposal, recent evidence supports a link between poor interoception and poor self-reported sleep quality across disorder groups, but not in the non-clinical control group (Ewing et al., 2017). Interestingly, in the non-clinical group, a positive

association was observed between reported sleep difficulties and interoceptive accuracy, in line with findings that insomnia is associated with greater heartbeat evoked potential amplitudes (Wei et al., 2016). Furthermore, poor sleep quality is observed across multiple psychiatric conditions (Freeman et al., 2017) including depression, anxiety and alexithymia (e.g. Bauermann et al., 2008; Bazydlo et al., 2001; Engin et al., 2010; Papadimitriou and Linkowski, 2005; Tsuno et al., 2005). More recent evidence suggests that both depression and alexithymia uniquely contribute towards poor self-reported sleep quality in population samples even after controlling for self-reported anxiety symptoms (Murphy, Wulff, et al., 2018). However, the causal relationship between interoception and sleep quality remains unclear; whilst it is possible that poor interoception results in risk of disrupted sleep, it is equally possible that poor sleep quality impacts negatively upon interoception (Ewing et al., 2017; Harshaw, 2015).

6. What gives rise to poor interoception? The development of typical interoception and how it may go awry

The evidence that interoception plays a fundamental role in a range of abilities has prompted interest in the stability of interoception across development, and how early interoceptive deficits may contribute to later difficulties. Converging evidence from measures of interoception, as well as alexithymia, indicate variability across the lifespan that coincides with established risk periods for the development of psychopathology and changes in higher order cognition. In particular, fluctuations in interoception may be especially common in adolescence and in late adulthood (see Murphy, Brewer, et al., 2017 for a review).

While interoceptive abilities appear to develop early in life (Maister et al., 2017), it is thus far unclear how this developmental process occurs. Further work is required, for example, into understanding how infants learn to distinguish multiple (and constantly changing) internal signals from each other, and how they learn to associate particular signals or combinations of signals with appropriate verbal labels. It is likely that caregivers assist infants in this process by labelling internal states using contextual cues. Children's behaviour, including infant cries, after a period without food or sleep, for example, may be labelled by caregivers as indicative of feeling hungry and tired, respectively. Learnt associations between internal states and these verbal labels may then allow the child to form representations of hunger and tiredness as distinct internal states. Difficulties perceiving internal cues would clearly hamper this process, as little interoceptive information would be available to be associated with verbal labels. Early difficulties perceiving interoceptive changes are therefore likely to exacerbate interoceptive difficulties later in life, as the individual would have less clearly defined categorical representations of internal states. Notably, if it is the case that linguistic input plays a role in the development of internal state representations, as has been proposed for emotional states (Barrett, 2006), interoceptive difficulties may also emerge in childhood following typical perception of internal signals during infancy. If, for example, caregivers are less able to correctly identify the child's internal states, or provide less verbal information about that state, the child would have less opportunity to refine their representation of that state, leading to intergenerational transfer of interoceptive impairment and alexithymia. Similarly, if an individual has intact interoceptive processing but limited verbal ability, they may not benefit from others providing verbal labels for their internal states (see Hobson et al. (2019) and (Way et al., 2007) for reviews of the role of language abilities in the development of interoceptive difficulties and alexithymia). While the relationship between parent and child interoception has not been explicitly investigated, parental anterior insula activation while watching video recordings of parent-infant interactions (their child and others' children) has been found to predict their reports of their child's somatic symptoms six years later, suggesting that parent interoception may be associated with child interoception (Abraham et al., 2019).

As in adulthood, interoception during childhood has predominantly

been examined in the cardiac domain. Echoing findings from adulthood, evidence supports the association between interoceptive accuracy in childhood and physical and mental health (Eley et al., 2004; Georgiou et al., 2015) and emotional abilities (Schaan et al., 2019a). These findings are supported by a similar relationship between alexithymia and mental health outcomes in children (Housiaux et al., 2010; Rieffe et al., 2006). Interoceptive ability in 6-11 year olds, however, appears equivalent to that in adults (Koch and Pollatos, 2014), indicating that if interoception develops over time, it may do so during early childhood years. Whilst the factors that contribute towards individual differences in the development and stability of interoceptive ability across childhood remain unclear, twin studies of cardiac interoception in childhood indicate that a substantial portion of the variance is accounted for by nonshared environmental factors (Elev et al., 2007), a similar heritability pattern to that observed for alexithymia in adulthood (Jørgensen et al., 2007). Indeed, more recent evidence suggests that a substantial portion of stability in interoceptive accuracy between the ages of 8 and 10 years is also accounted for by non-shared environmental factors (factors that make individuals within a family different). Whilst low stability was observed in this study, changes in interoception between the ages of 8-10 years were generally driven by improvements in interoceptive accuracy with age (Murphy, Cheesman, et al., 2019). Whilst the non-shared environmental factors that contribute to individual differences, stability and change in interoceptive accuracy have not been systematically identified, evidence that childhood adversity is associated with poor mental health (Kaplow and Widom, 2007; Macmillan et al., 2001; Repetti et al., 2002), atypical interoceptive abilities (Schaan et al., 2019b), and structural changes in interoceptive regions (Teicher et al., 2014) suggests that early negative experiences may contribute towards atypical interoception (see Murphy, Brewer, et al., 2017). Indeed, increased rates of alexithymia in adulthood have been associated with a number of negative experiences in childhood, such as physical or sexual abuse, emotional neglect, and poor family expressiveness (Aust et al., 2013; Berenbaum, 1996; Kench and Irwin, 2000).

While adolescence (the period of transition from childhood to adulthood) is associated with a number of positive changes, such as the emergence of higher-order cognitive abilities (e.g., Blakemore and Choudhury, 2006), this developmental period is also associated with increased risk taking, accidents, and susceptibility to psychopathology (Crone et al., 2016; Giedd et al., 2008; Kessler, Berglund, et al., 2005; Steinberg, 2007). Consistent with the proposal that both atypical learning and decision-making and the emergence of psychopathology are underpinned by interoceptive impairment, adolescence is associated with elevated rates of alexithymia (e.g. Gatta et al., 2014; Säkkinen et al., 2007, although see Honkalampi et al., 2009; Joukamaa et al., 2007), particularly at the earlier stages of this period (Gatta et al., 2014; Karukivi et al., 2014; Säkkinen et al., 2007). This spike in alexithymia prevalence in adolescence is consistent with the idea that interoceptive impairment represents a risk factor for, or is causally associated with, the development of psychopathology during this period of development (Murphy, Brewer, et al., 2017; Murphy, Viding, et al., 2019).

Whilst few studies have explicitly investigated interoception during this period, interoception appears to be related to a range of factors such as emotional abilities (Georgiou et al., 2018), body image (Todd et al., 2019), and health in adolescence, as in other stages of development. Atypical insula activation to interoceptive signals, for example, has been reported in adolescents with obesity (Delgado-Rico et al., 2013; Mata et al., 2015) and those with substance use disorders (Berk et al., 2015; Migliorini et al., 2013). Little research, however, has investigated developmental change in neural activity evoked by interoceptive signals in the typical population, across childhood, adolescence and adulthood. Whilst May et al. (2014) observed greater bilateral posterior insula activation in adolescents than adults in response to affective touch, this was not specific to the interoceptive signal, as similar increases were observed for non-affective touch. 6-17 year olds have been found to activate similar neural substrates to adults while performing the heartbeat counting task, such as the insula, medial prefrontal cortex and inferior parietal lobule (Klabunde et al., 2019). In this study, age was positively associated with activation in the anterior cingulate cortex, the orbitofrontal cortex and the left prefrontal cortex, although it should be noted that these findings are based on a sample of 11 participants across a broad age range. In a larger sample, activation patterns appeared to change with age differently across neural regions, with age being positively and linearly related to dorsal anterior activation, but nonlinearly related to ventral anterior insular activation, in response to interoceptive signals (Li et al., 2017). Neither of these studies directly compared neural activation to that in adult participants, however, meaning further work is needed to determine the full developmental trajectory. In a sample of neurotypical and autistic individuals spanning a larger age range (8-54 years), quadratic and linear relationships were observed between age and neural activation while performing the heartbeat counting task in different subdivisions of the insula, and linear relationships between age and activation in a broader interoceptive network, although age was unrelated to behavioural task performance (Failla et al., 2020). Interoceptive accuracy does appear to be associated with heartbeat evoked potentials in adolescence, as in adulthood (Mai et al., 2018), but again work is needed that explicitly follows the developmental trajectory throughout childhood and into adulthood. It is also worth noting that these studies have used cross-sectional designs to investigate effects of age, so longitudinal designs following development within the same individuals are clearly required in order to determine trajectories accurately.

Another stage where interoceptive ability may fluctuate is late adulthood, the period from 65 years of age until the end of life (Levinson, 1978). Like adolescence, this transitional period is associated with a number of cognitive and physical changes that may be underpinned by changes in interoception (see Whitbourne, 2012). For example, increasing age is associated with physical difficulties, such as increased risk of dehydration (e.g. Silver, 1990), as well as social and emotional difficulties, including poor emotion recognition (Ruffman et al., 2008) and risky decision-making (Sparrow and Spaniol, 2016), which have been linked to, or are supported by, interoception (Dunn, Stefanovitch, et al., 2010; Füstös et al., 2012; Rainer Schandry, 1981; Sokol-Hessner et al., 2014; Terasawa et al., 2014; Werner, Jung, Duschek, and Schandry, 2009; Wiens et al., 2000). Consistent with the proposal that interoceptive changes during this period may contribute to these difficulties (see Murphy, Brewer, et al., 2017), aging is associated with both a decline in interoceptive accuracy on cardiac tests (Khalsa, Rudrauf, and Tranel, 2009; Murphy, Geary, et al., 2017 although see Ainley et al., 2012; Mikkelsen et al., 2019) as well as increased rates of alexithymia (Joukamaa et al., 1996; Mattila et al., 2006; Paradiso et al., 2008 although see Gunzelmann et al., 2002). Impairments have also been observed in late adulthood in perceiving other interoceptive signals, such as pain, temperature, hydration and taste (Clark and Mehl, 1971; Gagliese, 2009; Silver, 1990; Stevens et al., 1995). Further, at the neural level, Paradiso et al. (2008) observed that increased age was associated with decreased volume of the anterior cingulate cortex. Whilst few studies have investigated the consequences of poor interoception in the later stages of life, a body of evidence indicates that elevated alexithymia in older adults is associated with poor physical and mental health (Bamonti et al., 2010; Hintistan et al., 2013; Waldstein et al., 2002). However, more recent research found that whilst emotion recognition abilities showed a small association with interoceptive accuracy in individuals age 20-90 years, interoceptive accuracy did not uniquely predict emotion recognition abilities after multiple cognitive and affective factors were controlled for (Murphy, Millgate, et al., 2019).

7. Outstanding questions

Taken together, existing evidence suggests that interoceptive abilities are closely related to typical functioning, for example in the domains of emotional processing, learning and decision-making, with atypical interoception being associated with impairments in these domains. Importantly, atypical interoception has been observed across a range of clinical conditions, and related to a number of transdiagnostic symptoms, in line with the suggestion that atypical interoception may be a general risk factor for the development or maintenance of psychopathology. A number of outstanding questions remain, however, that require additional testing in order to fully comprehend the role of interoception in psychopathology. The following sections outline these unresolved issues, and highlight priorities for future research.

7.1. Is interoception a unitary ability?

Whether interoceptive abilities are unitary, i.e., whether an individual with good cardiac perception will necessarily have good respiratory, gustatory, or temperature perception, or whether interoceptive abilities can dissociate in a signal-dependent manner, is a crucial question that remains to be answered. While we have thus far treated interoception as a unitary ability, it is likely that distinct interoceptive signals are processed independently, and multiple signals across domains are integrated to create a representation of the body's internal state as a whole (Ouigley et al., 2021). Research into the factor structure of interoceptive ability has been limited thus far by the high degree of reliance on cardiac measures of interoceptive abilities, but as discussed in Section 2.2, recent efforts have been made to produce tasks that span a number of interoceptive signals. While findings have been mixed, these initial investigations do not suggest that interoceptive ability is a unitary construct. It is important to note, however, that the tasks utilised in these studies are not always directly comparable; they often vary substantially in their trial and response formats, and place differential demands on working memory and attention. Indeed, the low correlations that have been observed between tasks that measure interoception within the same domain (e.g. between the heartbeat counting task and the heartbeat discrimination task; (Hickman et al., 2020) suggest that variation in task requirements can strongly influence relationships between tasks. Additionally, while a number of interoceptive domains have now been tested, there are numerous domains in which valid tests have not yet been developed. Future work is clearly required in this area, including further development of tasks assessing interoceptive abilities across a wide range of domains, which are ideally matched in terms of task demands. Once appropriate measures are available, it will be possible to determine the relationships between interoceptive abilities (both in typical and clinical populations), and whether training abilities in one domain leads to improvements in others.

Even if it is the case that perception of all interoceptive signals cannot be explained by a single ability, a small number of interoceptive clusters may exist, whereby processing of a given interoceptive signal is associated with processing of some, but not all, other interoceptive signals. If this is the case, the nature of clusters and how these are formed remains to be seen. Potentially, clusters could be based on the fibres that carry different signals (such as A delta fibres carrying signals of cold, touch, and first pain, C fibres carrying signals of second pain, warmth, itch, and muscle ache, and the vagus nerve carrying cardiac, respiratory information and gastric information (Craig, 2002)), or on the neurophysiological pathways that signals follow before being represented in the AI (e.g. A delta and C fibres projecting to lamina 1 of the spinal horn, and parasympathethic afferents such as the vagus and glossopharangeal nerve projecting directly to the nucleus of the solitary tract (Craig, 2002). Alternatively, clusters may represent psychological relationships, such as signals relaying information concerning threat (e.g. cardiac and respiratory cues; Diest et al., 2009) forming a distinct cluster.

Whether interoception is a unitary ability will have a number of implications for both research and clinical practice. First, it will determine whether interventions for those with interoceptive impairment can be standardised across individuals, or must be tailored to each individual's needs. While a number of clinical groups have previously been characterised by 'interoceptive impairment', evidence for

dissociations among interoceptive abilities necessitates further investigation into multiple interoceptive domains in each clinical population. It may be the case that specific patterns of interoceptive impairment are common to a given disorder. Individuals with EDs, for example, may experience particular impairment in the gastric domain, while individuals with anxiety disorders may only struggle to perceive signals relating to threat, such as cardiac and respiratory cues. This seems a possibility, as many of the disorder symptoms described throughout this report are linked to particular aspects of interoception. This proposal is complicated, however, by the observation of cardiac difficulties (most commonly assessed by the HCT) in a wide range of clinical conditions. Further research is clearly required to determine whether interoceptive profiles differ across and within clinical groups. Interventions aiming to improve interoception-related behaviours may be required to begin with a full interoceptive profile for each individual. If some interoceptive abilities do cluster, however, then interventions targeting one ability may be sufficient to improve others within that cluster, and reduce all associated disorder symptoms.

Secondly, determining the conceptual structure of interoception will identify whether previous findings in a particular interoceptive domain (in particular, from studies relying on cardiac perception) are applicable to other interoceptive abilities. While cardiac perception has been associated with multiple abilities, such as emotion processing (Ferguson and Katkin, 1996; Pollatos, Gramann, et al., 2007; Wiens et al., 2000; Critchley, Wiens, Rotshtein, Ohman, and Dolan, 2004), and learning and decision-making (Sokol-Hessner et al., 2014; Werner, Jung, Duschek, and Schandry, 2009), as the relationship between interoceptive domains is unclear, it cannot be assumed that recognition of other interoceptive signals also relates to these abilities. From a theoretical perspective, understanding the factor structure of interoceptive ability will aid in resolving the current controversy within the literature concerning the definition of interoception (Khalsa and Lapidus, 2016). While some define interoception in terms of neurophysiological pathways, and others in terms of the location of the origin of a signal, an understanding of the interoceptive signals that cluster together may lead to a more nuanced definition, encompassing multiple distinct domains. This relates to the debate over whether proprioception should be considered interoceptive. While early theories did discuss proprioceptive signals as within the definition of interoception (Cameron, 2001; Damasio, 2003; Vaitl, 1996), more recent work has excluded these signals (Craig, 2002, 2003; Critchley and Harrison, 2013). Proprioceptive information (similarly to exteroceptive information) is relayed by larger-diameter afferents than signals currently considered to be interoceptive, which project to a distinct area of the dorsal horn. An argument could be made for the inclusion of proprioception in the concept of interoception, however, based on the fact that these signals arise internally, and are processed by the insula (although more posteriorly to other interoceptive signals) (Bottini et al., 2001; Fasold et al., 2008; Ferrè et al., 2012; Mazzola et al., 2012; Petit and Beauchamp, 2003; Zu Eulenburg et al., 2013). Further, perception of peripersonal space appears to be associated with interoceptive accuracy (Ardizzi and Ferri, 2018). Determining which interoceptive cues are associated with each other would allow the umbrella term of 'interoception' to be expanded, with distinct clusters within this domain being clearly defined. Indeed, other signals beyond proprioception that are not currently considered interoceptive, but share a number of characteristics with interoception (e.g., olfaction) may also be included under a broader definition, with specified sub-domains. It is worth noting that if further sensory domains are included in the definition of interoception, interoceptive impairment may help to explain an even broader range of characteristics or symptoms across clinical conditions. If, for example, proprioception is treated as interoceptive, interoceptive difficulties could account for atypical biomotion and kinematics in those with ASD (Cook, Blakemore, and Press, 2013; Cook, 2016), and may link interoceptive difficulties to developmental coordination disorder, characterised by difficulties acquiring and executing coordinated motor skills (American Psychiatric

Association, 2013).

As well as the need to consider separable interoceptive domains (i.e., cardiac, respiratory, gastric, etc.), it is also necessary to determine separable interoceptive dimensions, as these may also relate differently to other abilities and psychopathology. As discussed above, we have recently proposed that one's objective accuracy in perceiving internal states, when explicitly aiming to do so, is distinct from one's attention towards interoceptive signals (Murphy et al., 2020; Murphy, Catmur, et al., 2019; Murphy, Brewer, et al., 2019). The latter may also be separable from one's propensity to utilise or rely on these states to guide behaviour in everyday life, but this is yet to be empirically tested. The notion of metacognitive awareness of one's own interoceptive abilities as described above (Garfinkel and Critchley, 2013) can be applied to each of these dimensions separately, in that an individual could have good or poor metacognitive understanding of their own accuracy, as well as of the degree to which they attend to internal signals (Murphy, Catmur, et al., 2019). Again, whether these dimensions relate to each other will inform understanding of whether interoception is a unitary ability.

If these dimensions (accuracy and attention, and their respective metacognitive abilities) are not strongly related to each other, it is possible that they relate differentially to psychological disorders. Whilst individuals with depression, for example, may have typical attention towards interoceptive information, they may struggle to perceive signals accurately. Alternatively, individuals with anxiety disorder may attend more to internal signals, while autistic individuals may attend to them less, leading to differences in the tendency to use these cues even if those with ASD and anxiety do not differ in their ability to objectively perceive internal signals. Of course, these dimensions may also interact or be causally related; reduced attention to internal signals, for example, may lead to reduced accuracy, owing to fewer opportunities to learn about one's internal signals. Profiles of abilities and difficulties across these interoceptive dimensions, as well as across interoceptive domains, therefore, should be examined across different clinical populations.

Research into the relationship between interoceptive accuracy and attention, and metacognitive awareness of each, is in its early stages. Much of this work is also limited by reliance upon comparing self-report measures of interoceptive attention with objective measures of interoceptive accuracy, making it difficult to determine relationships between them (Murphy et al., 2019). However, distinctions between these dimensions do appear to exist. Subjective reports of interoceptive attention, for example using the Body Perception Questionnaire and Body Awareness Questionnaire, do not appear to correlate well with objective interoceptive accuracy measures e.g. the Heartbeat Counting Task (e.g. Critchley, Wiens, Rotshtein, Öhman, and Dolan, 2004; Ferentzi, Drew, et al., 2018; Garfinkel et al., 2015), while self-reported interoceptive accuracy measures (questionnaires and confidence ratings) appear to correlate with each other as well as with objective measures of interoceptive accuracy, although the cardiac perception task used appears to affect results (Forkmann et al., 2016; Garfinkel et al., 2015; Murphy, Brewer, et al., 2019). Metacognitive awareness of one's own accuracy (also termed interoceptive insight) also appears to be weakly associated with objective accuracy in the cardiac domain, although again discrepancies across the objective cardiac tasks have been observed (Forkmann et al., 2016; Garfinkel et al., 2015; Murphy, Brewer, et al., 2019). Where clinical groups are concerned, research is lacking concerning the stability of the relationships between these different dimensions across different disorders, and the relationship between these dimensions and disorder symptoms. Early evidence does suggest, however, that these dimensions may be separately related to clinical symptoms. In autistic individuals, for example, a pattern of increased self-reported interoceptive attention but decreased objective cardiac accuracy has been observed, with the discrepancy predicting anxiety severity (Garfinkel et al., 2016b). The existence and impact of such discrepancies across different clinical conditions therefore warrants further scrutiny.

Whether interoceptive ability is a unitary construct also relates to evidence for multiple forms of alexithymia. Although many see alexithymia as primarily reflecting cognitive impairment in the representation or labelling of emotions (e.g. Luminet et al., 2006; Suslow and Junghanns, 2002), others have argued for the central role of affective difficulties, characterised by decreased ability to experience emotions (e.g. Bermond, 1997; Vorst and Bermond, 2001). The term 'Type I' alexithymia has been used to describe individuals who experience impairment in both the affective and cognitive domains, whereas 'Type II' alexithymia defines those with impairment only in the cognitive domain (Bermond, 1997). More recently, Moorman and colleagues also identified 'Type III' alexithymics, with affective but not cognitive impairment (Moormann et al., 2008). Impairment in the affective and cognitive domains has been associated, unsurprisingly, with different underlying neural atypicalities (Goerlich-Dobre et al., 2015). While the majority of research focuses on the cognitive dimension, research into the affective dimension of alexithymia is becoming more common, and these two dimensions are likely to relate differently to interoceptive abilities

In particular, it is possible that a similar distinction relates to interoception; that we can separate interoception at the perceptual level (comparable to the affective domain of alexithymia), from interoception at the cognitive level, involving representation, recognition and verbal labelling of interoceptive states. Some evidence does exist for a distinction between interoceptive perception and interoceptive recognition. Lesions to the left and right insula, for example, are associated with different consequences for taste perception and recognition; while perception is only impaired in the ipsilateral domain following either lesion, recognition is impaired ipsilaterally following right insula lesions, but bilaterally following left insula lesions (Cereda et al., 2002; Pritchard et al., 1999). Similarly, dissociations have been found in patients with insula lesions concerning olfactory stimuli; patients have been observed who struggle to label smells (as sweet or non-sweet), despite remaining able to distinguish perceptually between smells (Stevenson, Miller and Thayer, 2008). Similarly, patients with insula lesions have been observed who are able to perceive signals of pain, but no longer categorise these as painful (Berthier et al., 1988; Greenspan et al., 1999). These types of distinctions are similar to evidence suggesting that individuals with cognitive alexithymia are unimpaired at perceiving differences between emotional facial expressions, but impaired at classifying them, or labelling them as a particular emotion (Cook et al., 2013). There is clearly a need to distinguish between perception and recognition of internal states, despite the fact that these terms are often used interchangeably, or confounded, in the existing literature. Future work is required to determine whether these separable levels of interoceptive abilities relate differentially to psychopathology, or to social and emotional outcomes in the typical population.

A final component of the debate concerning whether interoception is unitary relates to the developmental origins of interoceptive impairment. While research into alexithymia indicates that developmental and acquired routes exist (Henry et al., 2006; Hogeveen et al., 2016; Lemche et al., 2004; Wood and Williams, 2007), little research has focused on this explicitly in the domain of non-emotional interoception. Interoceptive deficits following lesions, especially to the AI and ACC, are well documented (Ibañez et al., 2010), but the nature of the development of interoceptive difficulties in the absence of brain trauma requires further investigation, as described in Section 6. In particular, it is important to understand whether interoceptive impairments differ in their nature depending on whether they are neurodevelopmental, develop in response to environmental triggers, or develop in response to brain trauma. Similarly, whether interoceptive impairment is a stable trait, or a dynamic state requires investigation. Debate exists concerning whether alexithymia is a state or trait phenomenon; while many have observed stability over time (e.g. Mikolajczak and Luminet, 2006; Picardi et al., 2005) even in clinical populations (Berthoz and Hill, 2005; de Timary et al., 2008; Martínez-Sánchez et al., 2003; Saarijärvi et al.,

2006; Salminen et al., 2006; Stingl et al., 2008), others argue that severity may vary, especially with disorder symptoms in clinical populations (e.g. Honkalampi et al., 2004, 2001; Marchesi et al., 2005). Similarly, stability appears to differ depending on severity of alexithymia (de Haan et al., 2012), as well as across clinical populations (de Timary et al., 2008; Luminet et al., 2001; Rufer et al., 2004; Saarijärvi et al., 2006; Stingl et al., 2008). Overall, it appears that while alexithymia scores can change (low absolute stability), relative stability is fairly high (see Lumley et al., 2007 for a review). Where explicit measures of interoception have been used, while some degree of stability in interoceptive accuracy has been observed (e.g. Ferentzi, Drew, et al., 2018; Herbert et al., 2011; Koch and Pollatos, 2014; Pollatos, Traut-mattausch, et al., 2007; Van Dyck et al., 2016), situational factors can influence interoceptive accuracy, at least in the respiratory and cardiac domains (Bogaerts et al., 2005; van den Bergh et al., 2004; Wittkamp et al., 2018). Self-reported interoceptive attention (Ferentzi, Drew, et al., 2018; Murphy et al., 2020; Shields et al., 1989) and accuracy (Murphy, Brewer, et al., 2019) have also shown good test-retest reliability. Thus far, however, interoceptive abilities have not been monitored over a substantial time period, at least in adulthood. In childhood, modest correlations in HCT performance over one or two vears have been observed (Koch and Pollatos, 2014; Murphy, Cheesman, et al., 2019). Assessing and comparing the development and stability of interoceptive abilities across multiple interoceptive domains and dimensions will help to determine whether interoception is a unitary or fractionated construct.

7.2. If interoception accounts for symptom commonalities across disorders, what are differences between clinical groups attributable to?

Thus far, we have argued that atyipcal interoception may lead to multiple manifestations of psychopathology, and that similar difficulties across disorder groups may be explained by a common underlying interoceptive impairment. Despite the substantial overlap in symptoms and transdiagnostic markers across clinical populations, however, there are clearly differences in the behavioural and neural profiles associated with distinct disorders. If it is the case that poor interoception is a contributing factor to psychopathology, the causes of varying disorder outcomes require further investigation. Notably, clinician bias may affect the attribution of formal diagnostic labels (e.g. Schwartz et al., 2019), and there is an emphasis on the importance of individual case formulation, rather than relying on diagnostic categories, in clinical psychology (e.g. Bakker, 2019; Persons, 2008). This section is therefore focused on understanding the differences between individuals, and groups of individuals with similar manifestations of psychopathology, regardless of formal diagnosis.

As discussed above, clinical groups may be distinguished by impairment in separate domains or dimension of interoception, if interoceptive ability is not unitary. Relatedly, differences in the processing stage at which interoceptive impairment occurs may distinguish clinical populations, for example whether atypicalities exist in the physiological signal itself, perception (consciously or subconsciously) of the signal, or identification (labelling/recognition) of the signal (see e.g. Khalsa et al., 2018). Alternatively, it may be the case that the disorder that an individual develops is determined (at least in part) by the developmental stage at which interoceptive impairment is experienced. It is also possible that compensatory strategies for interoceptive impairment may be utilised differentially by clinical populations, leading to different patterns of difficulties. Notably, 'state' and 'trait' forms of interoceptive impairment may exist, whereby some individuals may be faced with these difficulties throughout their life, whereas others may experience impairments during specific periods only, potentially associated with changes in disorder sympomatology. Finally, demographic group and environmental influences, for example sex, ethnicity, education, social economic status, parenting styles, exposure to trauma, and cultural context, may also influence the specific manifestation of psychopathology (e.g. Breslau et al., 2005; Murphy, Viding, et al., 2019; Sayed et al., 2015; South and Krueger, 2011; Wadsworth and Achenbach, 2005). Further research should therefore investigate not only the interoceptive similarities across disorders, but also each of these potential explanations for varied behavioural and neurological manifestations of poor interoception, and its consequences, across clinical groups. Similarly, variation in interoceptive abilities within a particular disorder population may be responsible for the heterogeneity observed within that diagnostic category.

While we propose interoceptive impairment as a general risk factor, and even a potential candidate for the P factor of psychopathological susceptibility should it exist, we acknowledge that interoceptive abilities vary within a given clinical population, and that interoceptive impairment is not universal in psychological disorders. While cross-disorder symptom similarity may result from a common interoceptive impairment in a large portion of individuals, specific symptoms that are central to a particular disorder are presumably responsible for the majority of differences between diagnostic categories. It is likely that, while interoceptive difficulties are associated with a tendency to develop a psychological disorder, additional routes to disorder development (both genetic and environmental in nature) also exist, and contribute to the differences between, as well as within, clinical groups. The relative contribution of interoceptive and additional difficulties across clinical populations is therefore also an important area for subsequent investigation.

7.3. What is the precise nature of the relationship between interoception and psychopathology?

The main claim of the current paper is that atypical interoception causally increases the risk of psychopathology. The majority of the existing work on interoception and psychopathology has relied, however, on correlational designs and the relationship between interoception and clinical symptoms is likely to be complex, potentially varying across conditions. While it seems plausible that atypical interoception may lead to the development of a number of disorders such as depression, anxiety, eating disorders, OCD, PTSD, etc., the relationship may be bidirectional. For example, difficulties recognising internal signals of hunger and satiety may contribute to development of eating disorders, and presence of an eating disorder may lead to deliberate suppression of these signals, for example to reduce distress caused by hunger when calorific intake is being reduced, leading to reduced interoception attention, and in turn, accuracy. It may be the case that in some individuals, atypical interoception causes disorder development, while in others psychopathology leads to changes in interoception. It is likely that in many individuals, both of these processes are at play simultaneously, whereby the presence and severity of symptoms exacerbates interoceptive difficulties and vice versa. It is also important to consider neurodevelopmental disorders, which are lifelong conditions present from, or before, birth. In autism, for example, it is unlikely that interoception causes development of this condition. Autism may lead to interoceptive atypicalities, but it is also possible that both autism and interoceptive difficulties arise from widespread neural atypicality, making their co-occurrence highly likely. A similar explanation has been proposed for the co-occurrence between alexithymia and autism (Bird and Cook, 2013). While atypical interoception seems likely to cause psychopathology, testing the direction of the relationship through experimental manipulation (and longitudinal studies to some extent) across a range of clinical populations is clearly required.

Another aspect of the relationship between interoception and psychopathology that requires further investigation is its linearity. While it is often assumed that less accurate interoception, or lower attention to internal signals, leads to poorer mental health outcomes, this relationship has also been observed in the opposite direction, for example in the case of anxiety. This raises the possibility of a quadratic relationship between interoceptive abilities and psychopathology in certain conditions, which remains to be explicitly tested. Of course, this relationship could also vary across interoceptive dimensions and domains, so future work should aim to investigate this possibility using multiple measures of interoception.

7.4. Is intact interoception required online, or only during development?

Much of the evidence described thus far suggests that interoception is necessary for typical functioning in a number of domains (in particular, those of emotion processing and learning and decision-making), and that interoceptive impairment can lead to psychopathological outcomes. What remains to be seen, however, is whether interoception is only essential during development, for example while learning the affective value of particular cues, or whether it is crucial to continue to utilise interoception online. While interoceptive cues are clearly used continually, it may be that learnt associations subsequently allow cognitive processing to occur without interoceptive information. If, for example, one has associated a facial expression of happiness with internal cues to happiness, it is unclear whether the facial expression could be recognised in the absence of interoceptive signals, as exteroceptive cues (physical properties of the face) have previously been associated with the label 'happiness'. Simulationist models that emphasise 'embodiment' as important for emotion recognition (e.g. Barsalou et al., 2003; Gallese, 2001, 2003; Goldman and Sripada, 2005; Niedenthal et al., 2005; Preston and de Waal, 2001) would suggest that online interoceptive cues are necessary for this process. One way in which to determine the online contribution of internal cues would be to train or impede interoception in typical adults, and assess changes in other domains such as emotion recognition, or disorder symptom severity. Similarly, inducing internal states, for example during an emotion recognition task, and determining the extent to which internal signals bias perception and cognition, would add to our understanding of the online use of internal signals.

One indication that online interoception may not be required is the observation of intact emotional and interoceptive abilities in patients with insula and ACC lesions during adulthood (Damasio et al., 2013; Feinstein et al., 2016). Instead of suggesting that interoceptive cortex is not required for interoception, these findings may indicate that, assuming typical development, interoceptive information is not required online later in life. Potentially, if one has learnt to associate particular external stimuli with internal states, one can infer these internal states using these associations even in the absence of recognisable interoceptive information. If interoception is not required online, the possibility is raised that a sensitive period may exist during development, in which interoceptive learning is crucial. If an individual does not learn to associate interoceptive cues with exteroceptive cues, such as facial expressions and verbal labels, during this period, they may struggle to form these associations later in life. Similarly, if one does not attend to or perceive internal signals during a sensitive period, learning to do so subsequently may be particularly challenging. Whether interoceptive abilities develop during a sensitive period will have consequences for the efficacy of interoceptive training later in life; if a sensitive period exists, interoceptive training in adults identified as experiencing interoceptive impairment may be less effective than training during childhood. Sensitive periods may also occur later in life, for example during puberty, pregnancy and menopause, when physical changes may disrupt interoceptive signals, and perception of these signals must be adapted or relearnt (Murphy, Viding, et al., 2019). Whether these periods exist requires investigation, and will impact on our understanding of the importance of interoceptive interventions at specific developmental stages. The role of interoception online, and potential critical periods, may also vary across skill domains, for example emotion recognition, reward processing, and learning and decision-making, making it crucial to investigate this question across a range of skills.

7.5. How can interoceptive abilities be improved?

If it is the case that poor interoception contributes to multiple clinical conditions and impairments in numerous aspects of cognition, an obvious aim should be to improve these abilities where they are deficient. While a number of attempts have been made to improve interoception, it remains to be determined whether improvements are long-lasting, generalisable across domains, and indeed helpful for reducing psychopathology (see Weng et al. (2021) for a review). Once questions concerning the nature of the relationship between interoception, psychopathology, cognition, and the structure of interoception, have been answered, determining the most efficient and effective interventions will be possible.

Thus far, attempts have been made to train interoception using heartbeat perception training, whereby feedback leads to improved perception of one's heartbeats. Schandry and Weitkunat (1990), for example, demonstrated that auditory feedback presented 130 ms after the EKG-R wave (such that it is perceived in synchrony with the heartbeat) increased both cardiac accuracy, and heartbeat evoked potentials on a subsequent heartbeat perception task, indicating increased interoceptive accuracy in the cardiac domain. More recently, Schaefer et al. (2014) have used a similar procedure, involving presenting participants with visual cues 200 ms following the R wave, and a subsequent training phase where participants were required to press a button in synchrony with a particular heart beat (e.g. in three beats' time), and were provided with immediate feedback on their accuracy. Schaefer et al. found that this training procedure improved heartbeat perception in patients with medically unexplained symptoms. What remains to be seen is whether training in the cardiac domain can improve interoceptive abilities in other domains of interoception (e.g., perception and recognition of signals relating to respiration, hunger, temperature, pain, etc.). This type of training has also not been compared to control training conditions, meaning neither its specificity nor its generalisability has been investigated. Clearly, this is likely to depend on whether or not interoception is a unitary ability, so cross-domain training studies will help to elucidate the structure of interoception as well as whether single domain interventions are likely to be useful clinically. Further, whether this type of training alters attention to, or propensity to use, internal signals, or simply interoceptive accuracy when explicitly instructed to attend to these signals, requires investigation. A greater understanding of the relationship between psychopathology and potentially distinct interoceptive domains and dimensions will inform the development of effective interventions.

Beyond explicit interoceptive training, it is often assumed that meditation may improve interoception, although evidence has been mixed. Meditation experience appears to be associated with increased volume of the insula (Hernández et al., 2016) and ACC (Boccia et al., 2015), and functional activation of these interoceptive regions is observed during meditation tasks (Boccia et al., 2015; Farb et al., 2013). Behavioural evidence has been mixed; while some studies have found that meditation experience is unrelated to cardiac accuracy (Khalsa et al., 2008; Melloni et al., 2013; Parkin et al., 2014), a recent study suggested that cardiac accuracy was improved following contemplative practice over a period of three months (Bornemann and Singer, 2017). The same contemplative practice has been found to reduce self-reported alexithymia scores in those with low levels of alexithymia (Bornemann and Singer, 2017). Experienced meditators also exhibit increased levels of coherence between subjective and objective measures of emotional arousal (Sze et al., 2010). Notably, improved respiratory perception has been observed in meditators, possibly due to the fact that meditation focuses more on respiration than on cardiac signals (Daubenmier et al., 2013). A recent meta-analysis, however, did not find compelling evidence for an association between meditation and interoceptive accuracy, although only studies investigating the cardiac domain were included (Khalsa et al., 2020). Again, determining whether a fractionated structure of interoception exists will shed light upon the most

appropriate form of interoceptive training for individuals with specific interoceptive difficulties.

The distinction between interoceptive accuracy and attention is again relevant to the relationship between interoception and meditation or other forms of interoceptive training. The body-focused nature of meditation makes it likely that this practice leads to increased attention to, and use of, internal cues rather than increased interoceptive accuracy (Garland et al., 2012; Grossman, 2015), although increased attention may of course lead, in turn, to increased accuracy. Indeed, meditation has been found to increase self-reported interoceptive attention, as measured by the MAIA (Bornemann et al., 2015), and confidence and perceived ease in completing the heartbeat discrimination task (Khalsa et al., 2008). It is important to utilise measures that distinguish between these dimensions of interoception, to enable interventions to be tailored more effectively to individuals' interoceptive impairments. Importantly, whether interventions alter individuals' attention to, or accuracy perceiving, internal cues may affect their efficacy differentially in different clinical populations.

As studies explicitly training interoceptive abilities are scarce, it is also worth briefly examining the literature on changes in alexithymia. Attempts have been made to reduce alexithymia through therapeutic intervention, with group therapy sessions seeming to reduce selfreported alexithymia in a range of clinical groups, such as those with heart disease (Beresnevaite, 2000) and bulimia nervosa (de Groot et al., 1995), and panic disorder (Michael Rufer et al., 2010). Indeed, a review of studies reporting the effect of psychological interventions on alexithymia found that reductions in alexithymia are common, in particular when interventions specifically targeted alexithymia (Cameron et al., 2014). Whether changes in interoceptive accuracy or attention underlie this reduction remains to be investigated. As the alexithymia construct is explicitly related to emotional states, it may be the case that improving general interoceptive abilities can reduce alexithymia, but that interventions specifically aimed at reducing alexithymia are not sufficient to improve interoceptive skills more generally. If this is the case, training in the interoceptive domain may be a more effective strategy for improving recognition of both emotional and non-emotional internal states.

While future research should aim to determine how interoceptive abilities across multiple domains and dimensions can be improved, it is worth noting that not all individuals or disorder groups may benefit from interoceptive training. As discussed above, atypically high interoceptive accuracy or attention may be detrimental to one's mental health, in a similar way to atypically low interoceptive abilities. Determining the precise nature of the relationship between interoception and psychopathology, both in terms of directionality and linearity, will therefore be essential in order to develop the most beneficial interventions.

8. Conclusion

The current paper has reviewed evidence suggesting that interoception contributes to typical functioning, in terms of emotional abilities, reward-processing, learning and decision making, as well as findings indicating that atypical interoception is associated with impairments in these abilities. As many of these difficulties are shared by clinical conditions, we presented evidence for impaired interoception in a wide variety of these populations, and described possible mechanisms linking atypical interoception to disorder symptomatology. While a number of outstanding questions remain, it seems that interoceptive deficits represent a general risk factor for the development of psychopathology, perhaps even representing a plausible candidate for the P factor of psychopathology (Brewer et al., 2016b, 2016a; Murphy, Brewer, et al., 2017), whereby difficulties developing interoceptive abilities early in life, or changes in interoceptive abilities later in life, may predispose one to developing a number of psychological and neurological disorders. Future empirical work should investigate the

factors that determine which manifestations of psychopathology occur following atypical interoception, and whether distinct interoceptive domains and dimensions are associated with distinct clinical outcomes.

Acknowledgements

RB is supported by a New Investigator Grant from the Medical Research Council (MR/S003509/1).

References

- Aaron, R.V., Benson, T.L., Park, S., 2015. Investigating the role of alexithymia on the empathic deficits found in schizotypy and autism spectrum traits. Pers. Individ. Dif. 77, 215–220.
- Aaronson, S.T., Sears, P., Ruvuna, F., Bunker, M., Conway, C.R., Dougherty, D.D., et al., 2017. A 5-Year observational study of patients with treatment-resistant depression treated with vagus nerve stimulation or treatment as usual: comparison of response, remission, and suicidality. Am. J. Psychiatry 174 (7), 640–648.
- Abraham, E., Hendler, T., Zagoory-sharon, O., Feldman, R., 2019. Interoception sensitivity in the parental brain during the first months of parenting modulates children's somatic symptoms six years later: the role of oxytocin. Int. J. Psychophysiol. 136, 39–48.
- Adolfi, F., Couto, B., Richter, F., Decety, J., Lopez, J., Sigman, M., et al., 2017. Convergence of interoception, emotion, and social cognition: a twofold fMRI metaanalysis and lesion approach. Cortex 88, 124–142.
- Ainley, V., Tajadura-Jiménez, A., Fotopoulou, A., Tsakiris, M., 2012. Looking into myself: changes in interoceptive sensitivity during mirror self-observation. Psychophysiology 49 (11), 1672–1676.
- Ainley, V., Brass, M., Tsakiris, M., 2014. Heartfelt imitation: high interoceptive awareness is linked to greater automatic imitation. Neuropsychologia 60, 21–28.
- Ainley, V., Maister, L., Tsakiris, M., 2015. Heartfelt empathy? No association between interoceptive awareness, questionnaire measures of empathy, reading the mind in the eyes task or the director task. Front. Psychol. 6, 554.
- Ainley, V., Tsakiris, M., Pollatos, O., Schulz, A., Herbert, B.M., 2020. Comment on "Zamariola et al. (2018), Interoceptive Accuracy Scores are Problematic: evidence from Simple Bivariate Correlations"— the empirical data base, the conceptual reasoning and the analysis behind this statement are misconceived and do not support. Biol. Psychol. 152, 107870.
- Aïte, A., Barrault, S., Cassotti, M., Borst, G., Bonnaire, C., Houdé, O., et al., 2014. The impact of alexithymia on pathological gamblers' decision making: a preliminary study of gamblers recruited in "sportsbook" casinos. Cogn. Behav. Neurol. 27 (2), 59–67.
- Akitsuki, Y., Sugiura, M., Watanabe, J., Yamashita, K., Sassa, Y., 2003. Contextdependent cortical activation in response to financial reward and penalty: an eventrelated fMRI study. Neuroimage 19, 1674–1685.
- Allport, L.E., Butcher, K.S., Baird, T.A., MacGregor, L., Desmond, P.M., Tress, B.M., et al., 2004. Insular cortical ischemia is independently associated with acute stress hyperglycemia. Stroke 35 (8), 1886–1891.
- Alvaro, P.K., Roberts, R.M., Clinical, M., Harris, J.K., 2013. A systematic review assessing bidirectionality between sleep disturbances, anxiety, and depression. Sleep 36 (7), 1059–1068.
- Ambrosecchia, M., Ardizzi, M., Russo, E., Ditaranto, F., Speciale, M., Vinai, P., et al., 2017. Interoception and autonomic correlates during social interactions. Implications for anorexia. Front. Hum. Neurosci. 11, 219.
- American Psychiatric Association, 2013. Diagnostic and Statistical Manual of Mental Disorder, 5th ed. American Psychiatric Association, Arlington, VA.
- Anderson, E.R., Hope, D.A., 2009. The relationship among social phobia, objective and perceived physiological reactivity, and anxiety sensitivity in an adolescent population. J. Anxiety Disord. 23 (1), 18–26.
- Andrew, D., Craig, A.D., 2001. Spinothalamic lamina I neurons selectively sensitive to histamine: a central neural pathway for itch. Nat. Neurosci. 4 (1), 72–77.
- Apfel, R.J., Sifneos, P.E., 1979. Alexithymia: concept and measurement. Psychother. Psychosom. 32 (1–4), 180–190.
- Ardizzi, M., Ferri, F., 2018. Interoceptive influences on peripersonal space boundary. Cognition 177 (July 2017), 79–86.
- Ardizzi, M., Ambrosecchia, M., Buratta, L., Ferri, F., Peciccia, M., Donnari, S., et al., 2016. Interoception and positive symptoms in schizophrenia. Front. Hum. Neurosci. 10 (July), 379.
- Asahi, T., Uwano, T., Eifuku, S., Tamura, R., Endo, S., Ono, T., Nishijo, H., 2006. Neuronal responses to a delayed-response delayed-reward go/nogo task in the monkey posterior insular cortex. Neuroscience 143 (2), 627–639.
- Aspell, J.E., Heydrich, L., Marillier, G., Lavanchy, T., Herbelin, B., Blanke, O., 2013. Turning body and self inside out: visualized heartbeats alter bodily selfconsciousness and tactile perception. Psychol. Sci. 24 (12), 2445–2453.
- Assogna, F., Palmer, K., Pontieri, F.E., Pierantozzi, M., Stefani, A., Gianni, W., et al., 2012. Alexithymia is a non-motor symptom of Parkinson disease. Am. J. Geriatr. Psychiatry 20 (2), 133–141.
- Augustine, J.R., 1996. Circuitry and functional aspects of the insular lobe in primates including humans. Brain Res. Rev. 22 (3), 229–244.
- Aust, S., Härtwig, E.A., Heuser, I., Bajbouj, M., 2013. The role of early emotional neglect in alexithymia. Psychol. Trauma Theory Res. Pract. Policy 5 (3), 225–232.

Aziz, Q., Thompson, D.G., Ng, V.W.K., Hamdy, S., Sarkar, S., Brammer, M.J., et al., 2000. Cortical processing of human somatic and visceral sensation. J. Neurosci. 20 (7), 2657–2663.

Bagby, R.M., Parker, J.D.A., Taylor, G.J., 1994. The twenty-item Toronto Alexithymia Scale - I. Item selection and cross-validation of the factor structure. J. Psychosom. Res. 38 (1), 23–32.

Bakker, G.M., 2019. A new conception and subsequent taxonomy of clinical psychological problems. BMC Psychol. 7 (1), 46.

Balleine, B.W., Dickinson, A., 2000. The effect of lesions of the insular cortex on instrumental conditioning: evidence for a role in incentive memory. J. Neurosci. 20 (23), 8954–8964.

Baltaxe, C.A.M., Simmons I.I.I., J.Q., 1975. Lanuage in childhood psychosis: a review. J. Speech Hear. Disord. 40, 439–458.

Bamonti, P.M., Heisel, M.J., Topciu, R.A., Franus, N., Talbot, N.L., Duberstein, P.R., 2010. Association of alexithymia and depression symptom severity in adults 50 years of age and older. Am. J. Geriatr. Psychiatry 18 (1), 51–56.

Banzett, R.B., Mulnier, H.E., Murphy, K., Rosen, S.D., Wise, R.J., Adams, L., 2000. Breathlessness in humans activates insular cortex. Neuroreport 11 (10), 2117–2120.

Banzhaf, C., Hoffmann, F., Kanske, P., Fan, Y., Walter, H., Spengler, S., et al., 2018. Interacting and dissociable effects of alexithymia and depression on empathy. Psychiatry Res. 270 (February), 631–638.

- Baranek, G.T., David, F.J., Poe, M.D., Stone, W.L., Watson, L.R., 2006. Sensory Experiences Questionnaire: discriminating sensory features in young children with autism, developmental delays, and typical development. J. Child Psychol. Psychiatry 47 (6), 591–601.
- Bard, P., 1928. A diencephalic mechanism for the expression of rage with special reference to the sympathetic nervous system. Am. J. Physiol. 84, 490–515.Bardo, M.T., Donohew, R.L., Harrington, N.G., 1996. Psychobiology of novelty seeking

and drug seeking behavior. Behav. Brain Res. 77 (1–2), 23–43. Bar-On, R., Tranel, D., Denburg, N.L., Bechara, A., 2003. Exploring the neurological

substrate of emotional and social intelligence. Brain 126, 1790–1800. Barrett, L.F., 2006. Are Emotions Natural Kinds? Perspect. Psychol. Sci. 1 (1), 28–58.

Barrett, L.F., 2000. Ale Emotions Natural Knits: Perspect, Psychol. Sci. 1 (1), 28–36. Barrett, L.F., Simmons, W.K., 2015. Interoceptive prediction in the brain. Nat. Rev. Neurosci. 16 (7), 419–429.

Barrett, L.F., Gross, J., Christensen, T.C., Benvenuto, M., 2001. Knowing what you're feeling and knowing what to do about it: mapping the relation between emotion differentiation and emotion regulation. Cogn. Emot. 15 (6), 713–724.

Barrett, L.F., Quigley, K.S., Bilss-Moreau, E., Aronson, K.R., 2004. Interoceptive sensitivity and self-reports of emotional experience. J. Pers. Soc. Psychol. 87 (5), 684–697.

- Barsalou, L.W., Niedenthal, P.M., Barbey, A.K., Ruppert, J.A., 2003. Social embodiment. In: Psychology of Learning and Motivation - Advances in Research and Theory, vol. 43, pp. 43–92.
- Barsky, A.J., 1992. Amplification, somatization, and the somatoform disorders. Psychosomatics: J. Consultation and Liaison Psychiatry 33 (1), 28–34.

Bastin, J., Deman, P., David, O., Gueguen, M., Benis, D., Minotti, L., et al., 2017. Direct recordings from human anterior insula reveal its leading role within the errormonitoring network. Cereb. Cortex 27 (bastin), 1545–1557.

Bauermann, T.M., Parker, J.D.A., Taylor, G.J., 2008. Sleep problems and sleep hygiene in young adults with alexithymia. Pers. Individ. Dif. 45, 318–322.

Bazydlo, R., Lumley, M.A., Roehrs, T., 2001. Alexithymia and polysomnographic measures of sleep in healthy adults. Psychosom. Med. 63 (1), 56–61.

Beadle, J.N., Paradiso, S., Salerno, A., McCormick, L.M., 2013. Alexithymia, emotional empathy, and self-regulation in anorexia nervosa. Ann. Clin. Psychiatry 25 (2), 107–120.

Beales, D.L., Dolton, R., 2000. Eating disordered patients: personality, alexithymia, and implications for primary care. Br. J. Gen. Pract. 50 (450), 21–26.

Bearden, C.E., Rosso, I.M., Houister, J.M., Sanchez, L.E., Hadley, T., Cannon, T.D., 2000. A prospective cohort study of childhood behavioral deviance and language

abnormalities as predictors of adult schizophrenia. Schizophr. Bull. 26 (2), 395–410. Bechara, Antoine, Damasio, H., 2002. Decision-making and addiction (part I): impaired activation of somatic states in substance dependent individuals when pondering

decisions with negative future consequences. Neuropsychologia 40 (10), 1675–1689. Bechara, Antoine, Naqvi, N., 2004. Listening to your heart: Interoceptive awareness as a gateway to feeling. Nat. Neurosci. 7 (2), 102–103.

Bechara, A., Damasio, H., Damasio, A.R., 2000. Emotion, decision making and the orbitofrontal cortex. Cereb. Cortex 10 (3), 295–307.

Ben-Sasson, A., Hen, L., Fluss, R., Cermak, S.A., Engel-Yeger, B., Gal, E., 2009. A metaanalysis of sensory modulation symptoms in individuals with autism spectrum disorders. J. Autism Dev. Disord. 39 (1), 1–11.

Berenbaum, H., 1996. Childhood abuse, alexithymia and personality disorder. J. Psychosom. Res. 41 (6), 585–595.

Beresnevaite, M., 2000. Exploring the benefits of group psychotherapy in reducing alexithymia in coronary heart disease patients: a preliminary study. Psychother. Psychosom. 69 (3), 117–122.

Berk, L., Stewart, J.L., May, A.C., Wiers, R.W., Davenport, P.W., Paulus, M.P., Tapert, S. F., 2015. Under pressure: adolescent substance users show exaggerated neural processing of aversive interoceptive stimuli. Addiction 110 (12), 2025–2036.

(Eds.), The (non)Expression of Emotions in Health and Disease. Tilburg University Press, Tilburg, The Netherlands, pp. 115–130.

Berntson, G.G., Khalsa, S.S., 2021. Neural circuits of interoception. Trends Neurosci. 44 (1), 17–28.

Berntson, G.G., Norman, G.J., Bechara, A., Tranel, D., Bruss, J., Cacioppo, J.T., 2011. The insula, the amygdala and evaluative processes. Psychol. Sci. 22 (1), 80–86.

Neuroscience and Biobehavioral Reviews 130 (2021) 470-508

Berthier, M., Starkstein, S., Leiguarda, R., 1988. Asymbolia for pain: a sensory-limbic disconnection syndrome. Ann. Neurol. 24 (1), 41–49.

Berthoz, S., Hill, E.L., 2005. The validity of using self-reports to assess emotion regulation abilities in adults with autism spectrum disorder. Eur. Psychiatry 20 (3), 291–298.

Besheer, J., Palmatier, M.I., Metschke, D.M., Bevins, R.A., 2004. Nicotine as a signal for the presence or absence of sucrose reward: a Pavlovian drug appetitive conditioning preparation in rats. Psychopharmacology 172 (1), 108–117.

Betka, S., Pfeifer, G., Garfinkel, S., Prins, H., Bond, R., Sequeira, H., et al., 2018. How Do Self-Assessment of Alexithymia and Sensitivity to Bodily Sensations Relate to Alcohol Consumption? Alcohol. Clin. Exp. Res. 42 (1), 81–88.

Bettelheim, B., 1967. Empty Fortress. Simon & Schuster., New York.

Bevins, R.A., 2009. Altering the motivational function of nicotine through conditioning processes. The Motivational Impact of Nicotine and Its Role in Tobacco Use. Springer US., pp. 111–129

Bevins, Rick A., Besheer, J., 2014. Interoception and learning: import to understanding and treating diseases and psychopathologies. ACS Chem. Neurosci. 5, 624–631.

Bibby, P.A., Ferguson, E., 2011. The ability to process emotional information predicts loss aversion. Pers. Individ. Dif. 51 (3), 263–266.

Bienkowski, P., Zatorski, P., Baranowska, A., Ryglewicz, D., Sienkiewicz-Jarosz, H., 2010. Neuroscience Letters Insular lesions and smoking cessation after first-ever ischemic stroke: A 3-month follow-up. Neurosci. Lett. 478 (3), 161–164.

Bird, G., Cook, R., 2013. Mixed emotions: the contribution of alexithymia to the emotional symptoms of autism. Transl. Psychiatry 3 (7), e285.

Bird, G., Viding, E., 2014. The self to other model of empathy: providing a new framework for understanding empathy impairments in psychopathy, autism, and alexithymia. Neurosci. Biobehav. Rev. 47, 520–532.

Bird, G., Silani, G., Brindley, R., White, S., Frith, U., Singer, T., 2010. Empathic brain responses in insula are modulated by levels of alexithymia but not autism. Brain 133 (5), 1515–1525.

Blakemore, S.-J., Choudhury, S., 2006. Brain development during puberty: state of the science. Dev. Sci. 9 (1), 11–14.

- Blakemore, S.J., Smith, J., Steel, R., Johnstone, C.E., Frith, C.D., 2000. The perception of self-produced sensory stimuli in patients with auditory hallucinations and passivity experiences: evidence for a breakdown in self-monitoring. Psychol. Med. 30 (5), 1131–1139.
- Boccia, M., Boccia, M., Piccardi, L., Guariglia, P., 2015. The meditative mind: a comprehensive Meta- analysis of MRI studies the meditative mind: a comprehensive meta-analysis of MRI studies. Comput. Biomed. Res. 2015 (June), 1–11.

Bodini, B., Mandarelli, G., Tomassini, V., Tarsitani, L., Pestalozza, I., Gasperini, C., et al., 2008. Alexithymia in multiple sclerosis: Relationship with fatigue and depression. Acta Neurol. Scand. 118 (1), 18–23.

Boettcher, H., Brake, C.A., Barlow, D.H., 2016. Origins and outlook of interoceptive exposure. J. Behav. Ther. Exp. Psychiatry 53, 41–51.

Bogaerts, K., Notebaert, K., Van Diest, I., Devriese, S., De Peuter, S., Van Den Bergh, O., 2005. Accuracy of respiratory symptom perception in different affective contexts. J. Psychosom. Res. 58 (6), 537–543.

Bonaz, B., Lane, R.D., Oshinsky, M.L., Kenny, P.J., Sinha, R., Mayer, E.A., Critchley, H.D., 2021. Diseases, disorders, and comorbidities of interoception. Trends Neurosci. 44 (1), 39–51.

Bonfils, K.A., Lysaker, P.H., Minor, K.S., Salyers, M.P., 2016. Affective empathy in schizophrenia: a meta-analysis. Schizophr. Res. 175 (1–3), 109–117.

Bonfils, K.A., Lysaker, P.H., Minor, K.S., Salyers, M.P., 2017. Empathy in schizophrenia: A meta-analysis of the Interpersonal Reactivity Index. Psychiatry Res. 249, 293–303.

Borg, C., Bedoin, N., Peyron, R., Bogey, S., Laurent, B., Thomas-antérion, C., et al., 2013. Impaired emotional processing in a patient with a left posterior insula-SII lesion Impaired emotional processing in a patient with a left posterior insula-SII lesion. Neurocase 19 (6), 592–603.

Borhani, K., Làdavas, E., Fotopoulou, A., Haggard, P., 2017. "Lacking warmth": alexithymia trait is related to warm-specific thermal somatosensory processing. Biol. Psychol. 128 (January), 132–140.

Bornemann, B., Singer, T., 2017. Taking time to feel our body: Steady increases in heartbeat perception accuracy and decreases in alexithymia over 9 months of contemplative mental training. Psychophysiology 54 (3), 469–482.

Bornemann, B., Herbert, B.M., Mehling, W.E., Singer, T., 2015. Differential changes in self-reported aspects of interoceptive awareness through 3 months of contemplative training. Front. Psychol. 6 (JAN), 1–13.

Borsci, G., Boccardi, M., Rossi, R., Rossi, G., Perez, J., Bonetti, M., Frisoni, G.B., 2009. Alexithymia in healthy women: a brain morphology study. J. Affect. Disord. 114 (1–3), 208–215.

Boswell, J.F., Farchione, T.J., Sauer-Zavala, S., Murray, H.W., Fortune, M.R., Barlow, D. H., 2013. Anxiety sensitivity and interoceptive exposure: a transdiagnostic construct and change strategy. Behav. Ther. 44 (3), 417–431.

Bothe, E., Palermo, R., Rhodes, G., Burton, N., Jeffery, L., 2019. Expression recognition difficulty is associated with social but not attention - to - detail autistic traits and reflects both alexithymia and perceptual difficulty. J. Autism Dev. Disord. (0123456789).

Bottini, G., Karnath, H.O., Vallar, G., Sterzi, R., Frith, C.D., Frackowiak, R.S., Paulesu, E., 2001. Cerebral representations for egocentric space: functional-anatomical evidence from caloric vestibular stimulation and neck vibration. Brain 124 (6), 1182–1196.

Botvinick, M., Cohen, J.D., 1998. Rubber hand 'feels' what eyes see. Nature 391 (February), 756.

Bouras, C., Kövari, E., Hof, P.R., 2001. Anterior cingulate cortex pathology in schizophrenia and bipolar disorder. Acta Neuropathol. 102, 373–379.

Bourke, M.P., Taylor, G.J., Parker, J.D., Bagby, R.M., 1992. Alexithymia in women with anorexia nervosa. A preliminary investigation. Br. J. Psychiatry 161 (2), 240–243. Bourke, C., Douglas, K., Porter, R., 2010. Processing of facial emotion expression in major depression: a review. Aust. N. Z. J. Psychiatry 44 (8), 681–696.

Brener, J., Kluvitse, C., 1988. Heartbeat detection: judgments of the simultaneity of external stimuli and heartbeats. Psychophysiology 25 (5), 554–561.

Brener, J., Ring, C., 2016. Towards a psychophysics of interoceptive processes: the measurement of heartbeat detection. Philos. Trans. Biol. Sci. 371 (1708), 20160015.

- Breslau, J., Kendler, K.S., Su, M., Gaxiola-Aguilar, S., Kessler, R., 2005. Lifetime risk and persistence of psychiatric disorders across ethnic groups in the United States. Psychol. Med. 35, 317–327.
- Brewer, R., Bird, G., 2019. Disordered social cognition: alexithymia and interoception. In: Workman, L., Reader, W., Barkow, J.H. (Eds.), The Cambridge Handbook of Evolutionary Perspectives on Human Behavior. Cambridge University Press, pp. 436–448.
- Brewer, R., Cook, R., Cardi, V., Treasure, J., Bird, G., 2015. Emotion recognition deficits in eating disorders are explained by co-occurring alexithymia. R. Soc. Open Sci. 2 (1), 140382.
- Brewer, R., Happé, F., Cook, R., Bird, G., 2015. Commentary on "Autism, oxytocin and interoception.": alexithymia, not Autism Spectrum disorders, is the consequence of interoceptive failure. Neurosci. Biobehav. Rev. 56, 348–353.
- Brewer, R., Cook, R., Bird, G., 2016a. Alexithymia: a general deficit of interoception. R. Soc. Open Sci. 3 (10), 150664.
- Brewer, R., Cook, R., Bird, G., 2016b. Shared interoceptive representations: the case of alexithymia. In: Obhi, S.S., Cross, E.S. (Eds.), Shared Representations. Cambridge Social Neuroscience Series. Cambridge University Press, Cambridge, UK.
- Brewer, R., Cook, R., Cardi, V., Treasure, J., Catmur, C., Bird, G., 2019. Alexithymia explains increased empathic personal distress in individuals with and without eating disorders. Q. J. Exp. Psychol. 72 (7), 1827–1836.
- Brody, A.L., Mandelkern, M.A., London, E.D., Childress, A.R., Lee, G.S., Bota, R.G., et al., 2002. Brain metabolic changes during cigarette craving. Arch. Gen. Psychiatry 59, 1162–1172.
- Brogan, A., Hevey, D., Pignatti, R., 2010. Anorexia, bulimia, and obesity: shared decision making deficits on the Iowa Gambling Task (IGT). J. Int. Neuropsychol. Soc. 16 (4), 711–715.
- Bruce, G., Curren, C., Williams, L., 2012. Addictive Behaviors Alexithymia and alcohol consumption: the mediating effects of drinking motives. Addict. Behav. 37 (3), 350–352.
- Bruch, H., 1973. Eating Disorders: Obesity, Anorexia Nervosa, and the Person Within. Basic Books, New York, NY.
- Brunsdon, V.E.A., Happé, F., 2014. Exploring the "fractionation" of autism at the cognitive level. Autism 18 (1), 17–30.
- Bury, G., García Huesca, M., Bhattacharya, J., Herrojo Ruiz, M., 2019. Cardiac afferent activity modulates early neural signature of error detection during skilled performance. Neuroimage 199, 704–717.
- Bylsma, L.M., Morris, B.H., Rottenberg, J., 2008. A meta-analysis of emotional reactivity in major depressive disorder. Clin. Psychol. Rev. 28 (4), 676–691.
- Cabanac, M., 1971. Physiological role of pleasure. Science 173 (4002), 1103–1108.
 Caccavale, S., Bove, D., Bove, R.M., 2016. Skin and brain: itch and psychiatric disorders.
 Giornale Italiano Di Dermatologia e Venereologia: Organo Ufficiale. Soc. Ital.
- Dermatol. Sifilogr. Sezioni Interprov. Soc. Ital. Dermatol. Sifilogr. 151 (5), 525–529.
 Cain, N.M., Ansell, E.B., Simpson, H.B., Pinto, A., 2015. Interpersonal functioning in obsessive–Compulsive personality disorder. J. Pers. Assess. 97, 90–99.
- Calder, A.J., Young, A.W., 2005. Understanding the recognition of facial identity and
- facial expression. Nat. Rev. Neurosci. 6 (8), 641–651Cameron, O.G., 2001. Interoception: the inside story A model for psychosomatic processes. Psychosom. Med. 63 (5), 697–710.
- Cameron, K., Ogrodniczuk, J., Hadjipavlou, G., 2014. Changes in alexithymia following psychological intervention: a review. Harv. Rev. Psychiatry 22 (3), 162–178.
- Cannon, W.B., 1929. Bodily Changes in Pain, Hunger, Fear and Rage. Appleton, New York, NY.
- Canteras, N.S., Swanson, L.W., 1992. The dorsal premammillary nucleus: an unusual component of the mammillary body. Proc. Natl. Acad. Sci. U.S.A. 89 (21), 10089–10093.
- Cantril, H., Hunt, W.A., 1932. Emotional effects produced by the injection of adrenalin. Am. J. Psychol. 44 (2), 300–307.
- Cardi, V., Tchanturia, K., Treasure, J., 2018. Premorbid and illness-related social difficulties in eating disorders: an overview of the literature and treatment developments. Curr. Neuropharmacol. 16, 1122–1130.
- Carta, M.G., Sancassiani, F., Pippia, V., Bhat, K.M., Sardu, C., Meloni, L., 2013. Alexithymia is associated with delayed treatment seeking in acute myocardial infarction. Psychother. Psychosom. 82 (3), 190–192.
- Cascio, C., McGlone, F., Folger, S., Tannan, V., Baranek, G., Pelphrey, Ka., Essick, G., 2008. Tactile perception in adults with autism: A multidimensional psychophysical study. J. Autism Dev. Disord. 38 (1), 127–137.
- Caseras, X., Mataix-Cols, D., Rimes, K.A., Giampietro, V., Brammer, M., Zelaya, F., et al., 2008. The neural correlates of fatigue: an exploratory imaginal fatigue provocation study in chronic fatigue syndrome. Psychol. Med. 38 (7), 941–951.
- Caspi, A., Houts, R.M., Belsky, D.W., Goldman-Mellor, S.J., Harrington, H., Israel, S., et al., 2014. The p factor: One general psychopathology factor in the structure of psychiatric disorders? Clin. Psychol. Sci. 2 (2), 119–137.
- Castaneda, A.E., Tuulio-Henriksson, A., Marttunen, M., Suvisaari, J., Lönnqvist, J., 2008. A review on cognitive impairments in depressive and anxiety disorders with a focus on young adults. J. Affect. Disord. 106 (1–2), 1–27.
- Cauda, F., D'Agata, F., Sacco, K., Duca, S., Geminiani, G., Vercelli, A., 2011. Functional connectivity of the insula in the resting brain. NeuroImage 55 (1), 8–23.
- Cereda, C., Ghika, J., Maeder, P., Bogousslavsky, J., 2002. Strokes restricted to the insular cortex. Neurology 59, 1950–1955.

- Ceunen, E., Vlaeyen, J.W.S., Van Diest, I., 2016. On the origin of interoception. Front. Psychol. 7, 743.
- Chahraoui, K., Duchene, C., Rollot, F., Bonin, B., Moreau, T., 2014. Longitudinal study of alexithymia and multiple sclerosis. Brain Behav. 4 (1), 75–82.
- Chan, M.-S., Chung, K.-F., Yung, K.-P., Yeung, W.-F., 2017. Sleep in schizophrenia: a systematic review and meta-analysis of polysomnographic findings in case-control studies. Sleep Med. Rev. 32, 69–84.
- Chen, Y.H., Rodgers, J., McConachie, H., 2009. Restricted and repetitive behaviours, sensory processing and cognitive style in children with autism spectrum disorders. J. Autism Dev. Disord. 39 (4), 635–642.
- Chen, W.G., Schloesser, D., Arensdorf, A.M., Simmons, J.M., Cui, C., Valentino, R., et al., 2021. The emerging science of interoception: sensing, integrating, interpreting, and regulating signals within the self. Trends Neurosci. 44 (1), 3–16.
- Chick, C.F., Rounds, J.D., Hill, A.B., Anderson, A.K., 2019. My body, your emotions: viscerosomatic modulation of facial expression discrimination. Biol. Psychol., 107779
- Christensen, J.F., Gaigg, S.B., Calvo-Merino, B., 2018. I can feel my heartbeat: dancers have increased interoceptive accuracy. Psychophysiology 55, e13008.
- Clark, D.M., 1986. A cognitive approach to panic. Behav. Res. Ther. 24 (4), 461–470.
- Clark, W.C., Mehl, L., 1971. Thermal pain: a sensory decision theory analysis of the effect of age and sex on d', various response criteria, and 50% pain threshold. J. Abnorm. Psychol. 78 (2), 202–212.
- Clark, L., Bechara, A., Damasio, H., Aitken, M., Sahakian, B., Robbins, T., 2008. Differential effects of insular and ventromedial prefrontal cortex lesions on risky decision-making, Brain 131 (Pt 5), 1311–1322.
- Clark, L., Lawrence, A.J., Astley-jones, F., Gray, N., 2009. Article gambling near-misses enhance motivation to gamble and recruit win-related brain circuitry. Neuron 61 (3), 481–490.
- Clark, L., Studer, B., Bruss, J., Tranel, D., Bechara, A., 2014. Damage to insula abolishes cognitive distortions during simulated gambling. Proc. Natl. Acad. Sci. U.S.A. 111 (16), 6098–6103.
- Cleland, C., Magura, S., Foote, J., Rosenblum, A., Kosanke, N., 2005. Psychometric properties of the Toronto Alexithymia Scale (TAS-20) for substance users. J. Psychosom. Res. 58 (3), 299–306.
- Coccurello, R., 2019. Anhedonia in depression symptomatology: appetite dysregulation and defective brain reward processing. Behav. Brain Res. 372, 112041.
- Cochrane, C.E., Brewerton, T.D., Wilson, D.B., Hodges, E.L., 1993. Alexithymia in the eating disorders. Int. J. Eat. Disord. 14 (2), 219–222.
- Coen, S., Gregory, L., 2007. Reproducibility of human brain activity evoked by esophageal stimulation using functional magnetic resonance imaging. Am. J. Physiol. Gastrointest. Liver Physiol. 293 (1), 188–197.
- Coll, M.-P., Penton, T., Hobson, H., 2017. Important methodological issues regarding the use of transcranial magnetic stimulation to investigate interoceptive processing: A Comment on Pollatos et al. (2016). Philos. Trans. Biol. Sci. 372, 20170046.
- Coll, M.P., Viding, E., Rütgen, M., Silani, G., Lamm, C., Catmur, C., Bird, G., 2017. Are we really measuring empathy? Proposal for a new measurement framework. Neurosci. Biobehav. Rev. 83 (October), 132–139.
- Coll, M.-P., Hobson, H., Bird, G., Murphy, J., 2020. Systematic Review and Meta-analysis of the Relationship Between the Heartbeat-evoked Potential and Interoception.
- Connelly, M., Denney, D.R., 2007. Regulation of emotions during experimental stress in alexithymia. J. Psychosom. Res. 62 (6), 649–656.
- Contreras, M., Ceric, F., Torrealba, F., 2007. Inactivation of the interoceptive insula disrupts drug craving and malaise induced by Lithium. Science 811 (5850), 655–659.
- Cook, R., Brewer, R., Shah, P., Bird, G., 2013. Alexithymia, not autism, predicts poor recognition of emotional facial expressions. Psychological Science 24 (5), 723–732.

Cook, J., 2016. From movement kinematics to social cognition: the case of autism. Philos. Trans. Biol. Sci. 371, 20150372.

- Cook, J.L., Blakemore, S.-J., Press, C., 2013. Atypical basic movement kinematics in autism spectrum conditions. Brain 136 (Pt 9), 2816–2824.
- Cook, R., Brewer, R., Shah, P., Bird, G., 2013. Alexithymia, not autism, predicts poor recognition of emotional facial expressions. Psychol. Sci. 24 (5), 723–732.
- Cornielle, O., Lush, P., 2021. Sixty Years after Orne's American Psychologist Article: a Conceptual Analysis of "Demand Characteristics.".
- Couto, B., Adolfi, F., Sedeño, L., Salles, A., Canales-Johnson, A., Alvarez-Abut, P., et al., 2015. Disentangling interoception: insights from focal strokes affecting the perception of external and internal milieus. Front. Psychol. 6, 503.
- Cox, B.J., Bourdeau, D., Company, S., 1995. Alexithymia in panic disorder and social phobia. Compr. Psychiatry 36 (3), 195–198.
- Craig, A.D.(Bud)., 1995. Distribution of brainstem projections from spinal lamina I neurons in the cat and the monkey. J. Comp. Neurol. 361 (2), 225–248.
- Craig, A.D., 2002. How do you feel? Interoception: the sense of the physiological condition of the body. Nat. Rev. Neurosci. 3 (8), 655–666.
- Craig, A.D., 2003. Interoception: the sense of the physiological condition of the body. Curr. Opin. Neurobiol. 13 (4), 500–505.
- Craig, A.D.(Bud), 2005. Forebrain emotional asymmetry: A neuroanatomical basis? Trends Cogn. Sci. 9 (12), 566–571.
- Craig, A.D.(Bud), 2008. Interoception and emotion: a neuroanatomical perspective. In: Lewis, M., Haviland-Jones, J.M., Feldman-Barrett, L. (Eds.), Handbook of Emotion, 3rd ed. Guildford Press, New York, NY, pp. 272–288.
- Craig, A.D., 2009. How do you feel-now? The anterior insula and human awareness. Nat. Rev. Neurosci. 10 (1), 59–70.
- Craig, A.D. (Bud)., 2011. Significance of the insula for the evolution of human awareness of feelings from the body. Ann. N. Y. Acad. Sci. 1225 (1), 72–82.
- Craig, A.D., Chen, K., Bandy, D., Reiman, E.M., 2000. Thermosensory activation of insular cortex. Nat. Neurosci. 3 (2), 184–190.

Crane, L., Goddard, L., Pring, L., 2009. Sensory processing in adults with autism spectrum disorders. Autism 13 (3), 215–228.

- Critchley, M., 1964. The neurology of psychotic speech. Br. J. Psychiatry 110, 353–364. Critchley, Hugo D., Garfinkel, S.N., 2015. Interactions between visceral afferent signaling and stimulus processing. Front. Neurosci. 9, 286.
- Critchley, Hugo D., Garfinkel, S.N., 2017. Interoception and emotion. Curr. Opin. Psychol. 17, 7–14.
- Critchley, Hugo D., Harrison, N.A., 2013. Visceral influences on brain and behavior. Neuron 77 (4), 624–638.
- Critchley, H.D., Corfield, D.R., Chandler, M.P., Mathias, C.J., Dolan, R.J., 2000. Cerebral correlates of autonomic cardiovascular arousal: a functional neuroimaging investigation in humans. J. Physiol. 523 (1), 259–270.
- Critchley, H.D., Mathias, C.J., Dolan, R.J., 2001. Neural activity in the human brain relating to uncertainty and arousal during anticipation. Neuron 29 (2), 537–545.
- Critchley, Hugo D., Wiens, S., Rotshtein, P., Ohman, A., Dolan, R.J., 2004. Neural systems supporting interoceptive awareness. Nat. Neurosci. 7 (2), 189–195.
- Critchley, Hugo D., Rotshtein, P., Nagai, Y., O'Doherty, J., Mathias, C.J., Dolan, R.J., 2005. Activity in the human brain predicting differential heart rate responses to emotional facial expressions. NeuroImage 24 (3), 751–762.
- Crone, E.A., van Duijvenvoorde, A.C.K., Peper, J.S., 2016. Annual Research Review: neural contributions to risk-taking in adolescence – developmental changes and individual differences. J. Child Psychol. Psychiatry 57 (3), 353–368.
- Crucianelli, L., Krahé, C., Jenkinson, P.M., Fotopoulou, A., 2018. Interoceptive ingredients of body ownership: affective touch and cardiac awareness in the rubber hand illusion. Cortex 104, 180–192.
- Cunningham, M.A., Dixon, C., 1961. A study of the language of an autistic child. J. Child Psychol. Psychiatry 2, 193–202.
- Cuthbert, B.N., Insel, T.R., 2013. Toward the future of psychiatric diagnosis: the seven pillars of RDoC. BMC Med. 11, 126.
- Cutuli, D., 2014. Cognitive reappraisal and expressive suppression strategies role in the emotion regulation: an overview on their modulatory effects and neural correlates. Front. Syst. Neurosci. 8, 175.
- Dale, A., Anderson, D., 1978. Information variables in voluntary control and classical conditioning of heart rate: field dependence and heart-rate perception. Percept. Mot. Skills 41 (1), 79–85.
- Damasio, A.R., 1994. Descartes' Error: Emotion, Reason, and the Human Brain. Putnam, NY.
- Damasio, Antonio, 2003. Feelings of emotion and the self. Ann. N. Y. Acad. Sci. 1001, 253–261.
- Damasio, A., Damasio, H., Tranel, D., 2013. Persistence of feelings and sentience after bilateral damage of the insula. Cereb. Cortex 23 (4), 833–846.
- Daniel, M.L., Cocker, P.J., Lacoste, J., Mar, A.C., Houeto, J.L., Belin-Rauscent, A., Belin, D., 2017. The anterior insula bidirectionally modulates cost-benefit decisionmaking on a rodent gambling task. Eur. J. Neurosci. 46, 2620–2628.
- Daros, A.R., Zakzanis, K.K., Rector, N.A., 2014. A quantitative analysis of facial emotion recognition in obsessive – compulsive disorder. Psychiatry Res. 215, 514–521.
 Daubenmier, J., Sze, J., Kerr, C.E., Kemeny, M.E., Mehling, W., 2013. Follow your
- Daubenmier, J., Sze, J., Kerr, C.E., Kemeny, M.E., Mehling, W., 2013. Follow you breath: respiratory interoceptive accuracy in experienced meditators. Psychophysiology 50 (8), 777–789.
- Davis, M.H., 1980. A multidimensional approach to individual differences in Empahty. Catalog of Selected Document in Psychol. 40 (7), 3480.
- Dawson, G., Webb, S.J., McPartland, J., 2005. Understanding the nature of face processing impairment in autism: insights from behavioral and electrophysiological studies. Dev. Neuropsychol. 27 (3), 425–458.
- de Araujo, I.E.T., Kringelbach, M.L., Rolls, E.T., McGlone, F., 2003. Human cortical responses to water in the mouth, and the effects of thirst. J. Neurophysiol. 90 (3), 1865–1876.
- De Berardis, D., Campanella, D., Nicola, S., Gianna, S., Alessandro, C., Chiara, C., et al., 2008. The impact of alexithymia on anxiety disorders: a review of the literature. Curr. Psychiatry Rev. 4 (2), 80–86.
- de Galan, M., Sellaro, R., Colzato, L.S., Hommel, B., 2014. Conflict adaptation is predicted by the cognitive, but not the affective alexithymia dimension. Front. Psychol. 5, 768.
- de Groot, J.M., Rodin, G., Olmsted, M.P., 1995. Alexithymia, depression, and treatment outcome in bulimia nervosa. Compr. Psychiatry 36 (1), 53–60.
- De Gucht, V., Heiser, W., 2003. Alexithymia and somatisation: quantitative review of the literature. J. Psychosom. Res. 54, 425–434.
- de Haan, H., Joosten, E., Wijdeveld, T., Boswinkel, P., van der Palen, J., De Jong, C., 2012. Alexithymia is not a stable personality trait in patients with substance use disorders. Psychiatry Res. 198 (1), 123–129.
- de Haan, H.A., van der Palen, J., Wijdeveld, T.G.M., Buitelaar, J.K., De Jong, C.A.J., 2014. Alexithymia in patients with substance use disorders: state or trait? Psychiatry Res. 216 (1), 137–145.
- De Pascalis, V., Alberti, M.L., Pandolfo, R., 1984. Anxiety, perception, and control of heart rate. Percept. Mot. Skills 59 (1), 203–211.
- de Timary, P., Luts, A., Hers, D., Luminet, O., 2008. Absolute and relative stability of alexithymia in alcoholic inpatients undergoing alcohol withdrawal: relationship to depression and anxiety. Psychiatry Res. 157 (1–3), 105–113.
- De Vries, A.M.M., Forni, V., Voellinger, R., Stiefel, F., 2012. Alexithymia in cancer patients: review of the literature. Psychother. Psychosom. 81, 79–86.
- De Witte, N.A.J., Sütterlin Braet, C., Mueller, S.C., 2016. Getting to the heart of emotion regulation in youth: the role of interoceptive sensitivity, heart rate variability, and parental psychopathology. PLoS One 11 (10), e0164615.

- de Zwaan, M., Biener, D., Bach, M., Wiesnagrotzki, S., Stacher, G., 1996. Pain sensitivity, alexithymia, and depression in patients with eating disorders: are they related? J. Psychosom. Res. 41 (1), 65–70.
- Decety, J., Lamm, C., 2006. Human empathy through the Lens of social neuroscience. Sci. World J. 6, 1146–1163.
- Del Parigi, A., Gautier, J., Chen, K., Salbe, A.D., Ravussin, E., Tataranni, P.A., 2002. Neuroimaging and obesity. Ann. N. Y. Acad. Sci. 967, 389–397.
- Delgado-Rico, E., Soriano-Mas, C., Verdejo-Román, Río-Valle, J.S., Verdejo-García, A., 2013. Decreased insular and increased midbrain activations during decision-making under risk in adolescents with excess weight. Obesity 21 (8), 1662–1668.
- Deng, Y., Ma, X., Tang, Q., 2013. Brain response during visual emotional processing: an fMRI study of alexithymia. Psychiatry Res. 213 (3), 225–229.
- Derbyshire, S.W.G., 2003. A systematic review of neuroimaging data during visceral stimulation. Am. J. Gastroenterol. 98 (1), 12–20.
- Desmedt, O., Luminet, O., Corneille, O., 2018. The heartbeat counting task largely involves non-interoceptive processes: evidence from both the original and an adapted counting task. Biol. Psychol. 138, 185–188.
- Desmedt, O., Corneille, O., Luminet, O., Murphy, J., Bird, G., Maurage, P., 2020. Contribution of time estimation and knowledge to heartbeat counting task performance under original and adapted instructions. Biol. Psychol. 154 (May), 107904.
- Díaz-Román, A., Perestelo-Pérez, L., Buela-Casal, G., 2015. Sleep in obsessive-compulsive disorder: a systematic review and meta-analysis. Sleep Med. 16 (9), 1049–1055.
- Diest, I., Van, Bradley, M.M., Guerra, P., Bergh, O.Van Den, Lang, P.J., 2009. Fearconditioned respiration and its association to cardiac reactivity. Biol. Psychol. 80, 212–217.
- Dirupo, G., Corradi-Dell'Acqua, C., Kashef, M., Debbané, M., Badoud, D., 2020. The role of interoception in understanding others' affect. Dissociation between superficial and detailed appraisal of facial expressions. Cortex 130, 16–31.
- Dom, G., Sabbe, B., Hulstijn, W., Van Den Brink, W., 2005. Substance use disorders and the orbitofrontal cortex Systematic review of behavioural decision-making and neuroimaging studies. Br. J. Psychiatry 187, 209–220.
- Domschke, K., Stevens, S., Pfleiderer, B., Gerlach, A.L., 2010. Interoceptive sensitivity in anxiety and anxiety disorders: an overview and integration of neurobiological findings. Clin. Psychol. Rev. 30 (1), 1–11.
- Dorard, G., Berthoz, S., Haviland, M.G., Phan, O., Corcos, M., Bungener, C., 2008. Multimethod alexithymia assessment in adolescents and young adults with a cannabis use disorder. Compr. Psychiatry 49 (6), 585–592.
- Drevets, W.C., Price, J.L., Furey, M.L., 2008. Brain structural and functional abnormalities in mood disorders: implications for neurocircuitry models of depression. Brain Struct. Funct. 213 (1–2), 93–118.
- DuBois, D., Ameis, S.H., Lai, M.-C., Casanova, M.F., Desarkar, P., 2016. Interoception in autism Spectrum disorder: a review. Int. J. Dev. Neurosci. 52, 104–111.
- Duchesne, M., de Oliveira Falcone, E.M., de Freitas, S.R., D'Augustin, J.F., Marinho, V., Appolinario, J.C., 2012. Assessment of interpersonal skills in obese women with binge eating disorder. J. Health Psychol. 17 (7), 1065–1075.
- Dunn, B.D., Dalgleish, T., Ogilvie, A.D., Lawrence, A.D., 2007. Heartbeat perception in depression. Behav. Res. Ther. 45 (8), 1921–1930.
- Dunn, B.D., Galton, H.C., Morgan, R., Evans, D., Oliver, C., Meyer, M., et al., 2010. Listening to your heart. How interoception shapes emotion experience and intuitive decision making. Psychol. Sci. 21 (12), 1835–1844.
- Dunn, B.D., Stefanovitch, I., Evans, D., Oliver, C., Hawkins, A., Dalgleish, T., 2010. Can you feel the beat? Interoceptive awareness is an interactive function of anxiety- and depression-specific symptom dimensions. Behav. Res. Ther. 48 (11), 1133–1138.
- Dunn, B.D., Evans, D., Makarova, D., White, J., Clark, L., 2012. Gut feelings and the reaction to perceived inequity: The interplay between bodily responses, regulation, and perception shapes the rejection of unfair offers on the ultimatum game. Cogn. Affect. Behav. Neurosci. 12 (3), 419–429.
- Eastabrook, J.M., Lanteigne, D.M., Hollenstein, T., 2013. Decoupling between physiological, self-reported, and expressed emotional responses in alexithymia. Pers. Individ. Dif. 55 (8), 978–982.
- Eddy, C.M., Macerollo, A., Martino, D., Cavanna, A.E., 2015. Interpersonal reactivity differences in Tourette syndrome. Psychiatry Res. 228, 932–935.
- Edel, M.A., Rudel, A., Hubert, C., Scheele, D., Brüne, M., Juckel, G., Assion, H.J., 2010. Alexithymia, emotion processing and social anxiety in adults with ADHD. Eur. J. Med. Res. 15, 403–409.
- Edwards, W., 1954. The theory of decision making. Psychol. Bull. 51 (4), 380-417.
- Edwards, J., Jackson, H.J., Pattison, P.E., 2002. Erratum to "Emotion recognition via facial expression and affective prosody in schizophrenia: A methodological review" [Clinical Psychology Review 22 (2002) 789–832]. Clin. Psychol. Rev. 22 (8), 1267–1285.
- Ehlers, A., 1993. Interoception and panic disorder. Adv. Behav. Res. Ther. 15, 3–21. Ehlers, A., Breuer, P., 1992. Increased cardiac awareness in panic disorder. J. Abnorm. Psychol. 101 (3), 371–382.
- Ehlers, A., Breuer, P., Dohn, D., Fiegenbaum, W., 1995. Heartbeat perception and panic disorder: Possible explanations for discrepant findings. Behav. Res. Ther. 33, 69–76.
- Eickhoff, S.B., Lotze, M., Wietek, B., Amunts, K., Enck, P., Zilles, K., 2006. Segregation of visceral and somatosensory afferents: an fMRI and cytoarchitectonic mapping study. NeuroImage 31 (3), 1004–1014.
- Eisenberg, L., Ascher, E., Kanner, L., 1959. A clinical study of Gilles de la Tourette's disease (maladie des tics) in children. Am. J. Psychiatry 115 (8), 715–723.
- Ekman, P., Levenson, R.W., Friesen, W.V., 1983. Autonomic nervous system activity distinguishes among emotions. Science 221 (4616), 1208–1210.
- Eley, T.C., Stirling, L., Ehlers, A., Gregory, A.M., Clark, D.M., 2004. Heart-beat perception, panic/somatic symptoms and anxiety sensitivity in children. Behav. Res. Ther. 42 (4), 439–448.

- Eley, T.C., Gregory, A.M., Clark, D.M., Ehlers, A., 2007. Feeling anxious: a twin study of panic/somatic ratings, anxiety sensitivity and heartbeat perception in children. J. Child Psychol. Psychiatry 48 (12), 1184–1191.
- Elrod, M., Hood, B., 2015. Sleep differences among children with autism spectrum disorders and typically developing peers: a meta-analysis. J. Dev. Behav. Pediatr. 36, 166–177.
- Engin, E., Keski, G., Dulgerler, S., Bilge, A., 2010. Anger and alexithymic characteristics of the patients diagnosed with insomnia: a control group study. J. Psychiatr. Ment. Health Nurs. 17, 692–699.
- Enzi, B., Amirie, S., Brüne, M., 2016. Empathy for pain-related dorsolateral prefrontal activity is modulated by angry face perception. Exp. Brain Res. 234 (11), 3335–3345. Ernst, J., Böker, H., Hättenschwiler, J., Schüpbach, D., Northoff, G., Seifritz, E.,
- Grimm, S., 2014. The association of interoceptive awareness and alexithymia with neurotransmitter concentrations in insula and anterior cingulate. Soc. Cogn. Affect. Neurosci. 9 (6), 857–863.
- Eshkevari, E., Rieger, E., Musiat, P., Treasure, J., 2014. An investigation of interoceptive sensitivity in eating disorders using a heartbeat detection task and a self-report measure. Eur. Eat. Disord. Rev. 22, 383–388.
- Evkaya, A., Karadag-saygi, E., Karali, D., Giray, E., 2019. Gait & Posture Validity and reliability of the Dynamic Gait Index in children with hemiplegic cerebral palsy *. Gait Posture 75 (May 2019), 28–33.
- Ewing, D.L., Manassei, M., Gould van Praag, C., Philippides, A.O., Critchley, H.D., Garfinkel, S.N., 2017. Sleep and the heart: interoceptive differences linked to poor experiential sleep quality in anxiety and depression. Biol. Psychol. 127, 163–172.
- Failla, M.D., Bryant, L.K., Heflin, B.H., Mash, L.E., Schauder, K., Davis, S., et al., 2020. Neural correlates of cardiac interoceptive focus across development: implications for social symptoms in autism Spectrum disorder. Autism Res. 13 (6), 908–920.
- Farb, N.A.S., Segal, Z.V., Anderson, A.K., 2013. Mindfulness meditation training alters cortical representations of interoceptive attention. Soc. Cogn. Affect. Neurosci. 8 (1), 15–26.
- Farrell, M.J., Egan, G.F., Zamarripa, F., Shade, R., Blair-West, J., Fox, P., Denton, D.A., 2006. Unique, common, and interacting cortical correlates of thirst and pain. Proc. Natl. Acad. Sci. U.S.A. 103 (7), 2416–2421.
- Farrer, C., Frith, C.D., 2002. Experiencing oneself vs another person as being the cause of an action: the neural correlates of the experience of agency. NeuroImage 15 (3), 596–603.
- Farrer, C., Franck, N., Georgieff, N., Frith, C.D., Decety, J., Jeannerod, M., 2003. Modulating the experience of agency: a positron emission tomography study. NeuroImage 18 (2), 324–333.
- Fasold, O., Heinau, J., Trenner, M.U., Villringer, A., Wenzel, R., 2008. Proprioceptive head posture-related processing in human polysensory cortical areas. NeuroImage 40 (3), 1232–1242.
- Fassino, S., Pierò, A., Gramaglia, C., Abbate-Daga, G., 2004. Clinical, psychopathological and personality correlates of interoceptive awareness in anorexia nervosa, bulimia nervosa and obesity. Psychopathology 37 (4), 168–174.
- Feinstein, J.S., Khalsa, S.S., Salomons, T.V., Prkachin, K.M., Frey-Law, L.A., Lee, J.E., et al., 2016. Preserved emotional awareness of pain in a patient with extensive bilateral damage to the insula, anterior cingulate, and amygdala. Brain Struct. Funct. 221 (3), 1499–1511.
- Ferentzi, E., Bogdány, T., Szabolcs, Z., Csala, B., 2018. Multichannel investigation of interoception: sensitivity is not a generalizable feature. Front. Hum. Neurosci. 12, 223.
- Ferentzi, E., Drew, R., Tihanyi, B.T., Köteles, F., 2018. Interoceptive accuracy and body awareness – temporal and longitudinal associations in a non-clinical sample. Physiol. Behav. 184 (August 2017), 100–107.
- Ferguson, M.L., Katkin, E.S., 1996. Visceral perception, anhedonia, and emotion. Biol. Psychol. 42 (1–2), 131–145.
- Ferguson, E., Bibby, P.A., Rosamond, S., O'Grady, C., Parcell, A., Amos, C., et al., 2009. Alexithymia, cumulative feedback, and differential response patterns on the iowa gambling task. J. Pers. 77 (3), 883–902.
- Fernandez, E., Turk, D.C., 1992. Sensory and affective components of pain: separation and synthesis. Psychol. Bull. 112 (2), 205–217.
- Ferrè, E.R., Bottini, G., Haggard, P., 2012. Vestibular inputs modulate somatosensory cortical processing. Brain Struct. Funct. 217 (4), 859–864.
- Ferroni, F., Ardizzi, M., Sestito, M., Lucarini, V., Daniel, B.D., Paraboschi, F., et al., 2019. Shared multisensory experience affects Others' boundary: the enfacement illusion in schizophrenia. Schizophr. Res. 206, 225–235.
- Fiene, L., Ireland, M., Brownlow, C., 2018. The interoception sensory questionnaire (ISQ): a scale to measure interoceptive challenges in adults. J. Autism Dev. Disord. 48, 3354–3366.
- Foland-Ross, L.C., Altshuler, L.L., Bookheimer, S.Y., Lieberman, M.D., Townsend, J., Penfold, C., et al., 2010. Amygdala reactivity in healthy adults is correlated with prefrontal cortical thickness. J. Neurosci. 30 (49), 16673–16678.
- Fond, G., Loundou, A., Hamdani, N., Boukouaci, W., Dargel, A., Oliveira, J., et al., 2014. Anxiety and depression comorbidities in irritable bowel syndrome (IBS): a systematic review and meta-analysis. Eur. Arch. Psychiatry Clin. Neurosci. 264, 651–660.
- Forkmann, T., Scherer, A., Meessen, J., Michal, M., Schächinger, H., Vögele, C., Schulz, A., 2016. Making sense of what you sense: disentangling interoceptive awareness, sensibility and accuracy. Int. J. Psychophysiol. 109, 71–80.
- Fortune, E.E., Goodie, A.S., 2012. Cognitive distortions as a component and treatment focus of pathological gambling: a review. Psychol. Addict. Behav. 26 (2), 298–310.
- Foss-Feig, J.H., Heacock, J.L., Cascio, C.J., 2012. Tactile responsiveness patterns and their association with core features in autism spectrum disorders. Res. Autism Spectr. Disord. 6 (1), 337–344.

- Francis, S., Rolls, E.T., Bowtell, R., McGlone, F., O'Doherty, J., Browning, A., et al., 1999. The representation of pleasant touch in the brain and its relationship with taste and olfactory areas. Neuro. Rep. 10 (3), 453–459.
- Frank, G.K.W., 2015. Advances from neuroimaging studies in eating disorders. CNS Spectr. 20 (04), 391–400.
- Freeman, D., Sheaves, B., Goodwin, G.M., Yu, L.-M., Nickless, A., Harrison, P.J., et al., 2017. The effects of improving sleep on mental health (OASIS): a randomised controlled trial with mediation analysis. Lancet Psychiatry 4 (10), 749–758.
- Frewen, P.A., Pain, C., Dozois, D.J.A., Lanius, R.A., 2006. Alexithymia in PTSD: psychometric and fMRI studies. Ann. N. Y. Acad. Sci. 1071, 397–400.
- Frewen, P.A., Dozois, D.J.A., Neufeld, R.W.J., Lanius, R.A., 2008. Meta-analysis of alexithymia in posttraumatic stress disorder. J. Trauma. Stress 21 (2), 243–246.
- Friedman, S.R., Rapport, L.J., Lumley, M., Tzelepis, A., Vanvoorhis, A., Stettner, L., Kakaati, L., 2003. Aspects of social and emotional competence in adult attentiondeficit / hyperactivity disorder. Neuropsychology 17 (1), 50–58.
- Friston, K.J., Lawson, R., Frith, C.D., 2013. On hyperpriors and hypopriors: comment on Pellicano and Burr. Trends Cogn. Sci. 17 (1), 1.
- Frith, C., 2012. Explaining delusions of control: the comparator model 20years on. Conscious. Cogn. 21 (1), 52–54.
- Frith, U., Morton, J., Leslie, A.M., 1991. The cognitive basis of a biological disorder: autism. Trends Neurosci. 14 (10), 433–438.
- Fukushima, H., Terasawa, Y., Umeda, S., 2011. Association between interoception and empathy: evidence from heartbeat-evoked brain potential. Int. J. Psychophysiol. 79 (2), 259–265.
- Furman, D.J., Waugh, C.E., Bhattacharjee, K., Thompson, R.J., Gotlib, I.H., 2013. Interoceptive awareness, positive affect, and decision making in Major Depressive Disorder. J. Affect. Disord. 151 (2), 780–785.
- Füstös, J., Gramann, K., Herbert, B.M., Pollatos, O., 2012. On the embodiment of emotion regulation: interoceptive awareness facilitates reappraisal. Soc. Cogn. Affect. Neurosci. 8 (8), 911–917.
- Gabriele, E., Spooner, R., Brewer, R., Murphy, J., 2020. Dissociations between interoceptive accuracy and attention: evidence from the interoceptive attention scale. https://doi.org/10.31234/osf.io/vjph6. November 6.
- Gagliese, L., 2009. Pain and aging: the emergence of a new subfield of pain research. J. Pain 10 (4), 343–353.
- Gaigg, S.B., Cornell, A.S., Bird, G., 2016. The psychophysiological mechanisms of alexithymia in autism Spectrum disorder. J. Abnorm. Psychol.
- Gallese, V., 2001. The "Shared manifold" hypothesis: from mirror neurons to empathy. J. Conscious. Stud. 8 (5-7), 33-50.
- Gallese, V., 2003. The roots of empathy: the shared manifold hypothesis and the neural basis of intersubjectivity. Psychopathology 36 (4), 171–180.
- Ganos, C., Garrido, A., 2015. Premonitory urge to tic in Tourette's is associated with interoceptive awareness. Mov. Disord. 30 (9), 1198–1202.
- García-Cordero, I., Sedeño, L., De La Fuente, L., Slachevsky, A., Forno, G., Klein, F., et al., 2016. Feeling, learning from, and being aware of inner states: interoceptive dimensions in neurodegeneration and stroke. Philos. Trans. R. Soc. Lond., B, Biol. Sci. 371 (1708), 20160006.
- Garfinkel, S.N., Critchley, H.D., 2013. Interoception, emotion and brain: new insights link internal physiology to social behaviour. Commentary on: "Anterior insular cortex mediates bodily sensibility and social anxiety" by Terasawa et al. (2012). Soc. Cogn. Affect. Neurosci. 8 (3), 231–234.
- Garfinkel, Sarah N., Critchley, H.D., 2016. Threat and the body: how the heart supports fear processing. Trends Cogn. Sci. 20 (1), 34–46.
- Garfinkel, Sarah N., Liberzon, I., 2009. Neurobiology of PTSD: a review of neuroimaging findings. Psychiatr. Ann. 39 (6), 370–381.
- Garfinkel, Sarah N., Barrett, A.B., Minati, L., Dolan, R.J., Seth, A.K., Critchley, H.D., 2013. What the heart forgets: cardiac timing influences memory for words and is modulated by metacognition and interoceptive sensitivity. Psychophysiology 50 (6), 505–512.
- Garfinkel, Sarah N., Minati, L., Gray, M.A., Seth, A.K., Dolan, R.J., Critchley, H.D., 2014. Fear from the heart: sensitivity to fear stimuli depends on individual heartbeats. J. Neurosci. 34 (19), 6573–6582.
- Garfinkel, Sarah N., Seth, A.K., Barrett, A.B., Suzuki, K., Critchley, H.D., 2015. Knowing your own heart: distinguishing interoceptive accuracy from interoceptive awareness. Biol. Psychol. 104, 65–74.
- Garfinkel, Sarah N., Manassei, M.F., Hamilton-Fletcher, G., In Den Bosch, Y., Critchley, H.D., Engels, M., 2016. Interoceptive dimensions across cardiac and respiratory axes. Philos. Trans. Biol. Sci. 371, 20160014.
- Garfinkel, Sarah N., Tiley, C., O'Keeffe, S., Harrison, N.A., Seth, A.K., Critchley, H.D., 2016a. Discrepancies between dimensions of interoception in autism: implications for emotion and anxiety. Biol. Psychol. 114 (September), 117–126.
- Garfinkel, Sarah N., Tiley, C., O'Keeffe, S., Harrison, N.A., Seth, A.K., Critchley, H.D., 2016b. Discrepancies between dimensions of interoception in autism: implications for emotion and anxiety. Biol. Psychol. 114, 117–126.
- Garland, E.L., Gaylord, S.A., Palsson, O., Faurot, K., Mann, J.D., Whitehead, W.E., 2012. Therapeutic mechanisms of a mindfulness-based treatment for IBS: Effects on visceral sensitivity, catastrophizing, and affective processing of pain sensations. J. Behav. Med. 35 (6), 591–602.
- Garner, D.M., Olmsted, M.P., Polivy, J., 1983. The Eating Disorder Inventory: a measure of cognitive-behavioral dimensions of anorexia nervosa and bulimia. Anorexia Nervosa: Recent Dev. Res. 3 (October), 173–184.
- Gatta, M., Facca, I., Colombo, E., Svanellini, L., Montagnese, S., Schiff, S., 2014. Alexithymia, psychopathology and alcohol misuse in adolescence: a population based study on 3556 teenagers. Neurosci. Med. 5, 60–71.

Gaznick, N., Tranel, D., McNutt, a., Bechara, a., 2014. Basal ganglia plus insula damage yields stronger disruption of smoking addiction than basal ganglia damage alone. Nicotine Tob. Res. 16 (4), 445–453.

Georgiou, E., Matthias, E., Kobel, S., Kettner, S., Deyhaupt, J., Steinacker, J.M., Pollatos, O., 2015. Interaction of physical activity and interoception in children. Front. Psychol. 6, 502.

Georgiou, E., Mai, S., Fernandez, K.C., Pollatos, O., 2018. I see neither your Fear, nor your Sadness – interoception in adolescents. Conscious. Cogn. 60 (February), 52–61.

Gergely, G., Watson, J.S., 1996. The social biofeedback theory of parental affectmirroring: the development of emotional self-awareness and self-control in infancy.

Int. J. Psychoanal. 77 (6), 1181–1212. Giedd, J.N., Keshavan, M., Paus, T., 2008. Why do many psychiatric disorders emerge

during adolescence? Nat. Rev. Neurosci. 9 (12), 947–957. Glaros, A.G., Lumley, M.A., 2005. Alexithymia and pain in temporomandibular disorder. J. Psychosom. Res. 59, 85–88.

Gleichgerrcht, E., Tomashitis, B., Sinay, V., 2015. The relationship between alexithymia, empathy and moral judgment in patients with multiple sclerosis. Eur. J. Neurol. 22 (9), 1295–1303.

Goerlich, K.S., Votinov, M., Lammertz, S.E., Winkler, L., Spreckelmeyer, K.N., Habel, U., et al., 2016. Effects of alexithymia and empathy on the neural processing of social and monetary rewards. Brain Struct. Funct.

Goerlich-Dobre, K.S., Bruce, L., Martens, S., Aleman, A., Hooker, C.I., 2014. Distinct associations of insula and cingulate volume with the cognitive and affective dimensions of alexithymia. Neuropsychologia 53, 284–292.

Goerlich-Dobre, K.S., Votinov, M., Habel, U., Pripfl, J., Lamm, C., 2015. Neuroanatomical profiles of alexithymia dimensions and subtypes. Hum. Brain Mapp. 36 (10), 3805–3818.

Goldman, A.I., Sripada, C.S., 2005. Simulationist models of face-based emotion recognition. Cognition 94 (3), 193–213.

Goldstein, R.Z., Craig, A.D.(Bud), Bechara, A., Garavan, H., Childress, A.R., Paulus, M.P., Volkow, N.D., 2009. The neurocircuitry of impaired insight in drug addiction. Trends Cogn. Sci. 13 (9), 372–380.

Goodkind, M., Eickhoff, S.B., Oathes, D.J., Jiang, Y., Chang, A., Jones-Hagata, L.B., et al., 2015. Identification of a common neurobiological substrate for mental illness. JAMA Psychiatry 72 (4), 305–315.

Gordon, I., Voos, A.C., Bennett, R.H., Bolling, D.Z., Pelphrey, K.A., Kaiser, M.D., 2013. Brain mechanisms for processing affective touch. Hum. Brain Mapp. 34 (4), 914–922.

Gorka, S.M., Nelson, B.D., Phan, K.L., Shankman, S.A., 2016. Intolerance of uncertainty and insula activation during uncertain reward. Cogn. Affect. Behav. Neurosci. 16 (5), 929–939.

Gowin, J.L., Stewart, J.L., May, A.C., Ball, T.M., Wittmann, M., Tapert, S.F., Paulus, M.P., 2014. Altered cingulate and insular cortex activation during risk-taking in methamphetamine dependence: losses lose impact. Addiction 109 (2), 237–247.

Grabe, H.J., Wittfeld, K., Hegenscheid, K., Hosten, N., Lotze, M., Janowitz, D., et al., 2014. Alexithymia and brain gray matter volumes in a general population sample. Hum. Brain Mapp. 35 (12), 5932–5945.

Gray, M.A., Critchley, H.D., 2007. Interoceptive basis to craving. Neuron 54 (2), 183–186.

Gray, M.A., Taggart, P., Sutton, P.M., Groves, D., Holdright, D.R., Bradbury, D., et al., 2007. A cortical potential reflecting cardiac function. PNAS 104 (16), 6818–6823.

Gray, M., Beacher, F., Minati, L., Nagai, Y., Kemp, A.H., Harrison, N., Critchley, H., 2012. Emotional appraisal is influenced by cardiac afferent information. Emotion 12 (1), 180–191.

Green, D., Beaton, L., Moore, D., Warren, L., Wick, V., Sanford, J.E., Santosh, P., 2003. Clinical incidence of sensory integration difficulties in adults with learning disabilities and illustration of management. Br. J. Occup. Ther. 66 (10), 454–463.

Green, M.F., Horan, W.P., Lee, J., 2015. Social cognition in schizophrenia. Nat. Rev. Neurosci. 16 (10), 620–631.

Green, D., Chandler, S., Charman, T., Simonoff, E., Baird, G., 2016. Brief report: DSM-5 sensory behaviours in children with and without an autism Spectrum disorder. J. Autism Dev. Disord. 46 (11), 3597–3606.

Greenspan, J.D., Lee, R.R., Lenz, F.A., 1999. Pain sensitivity alterations as a function of lesion location in the parasylvian cortex. Pain 81 (3), 273–282.

Griffiths, T.D., Warren, J.D., Dean, J.L., Howard, D., 2004. "When the feeling's gone": a selective loss of musical emotion. J. Neurol. Neurosurg. Psychiatr. 75 (2), 344–345.

Gross, J.J., John, O.P., 2003. Individual differences in two emotion regulation processes: implications for affect, relationships, and well-being. J. Pers. Soc. Psychol. 85 (2), 348–362.

Grossi, D., Trojano, L., Postiglione, A., Soricell, A., Mansi, L., Salvatore, M., 1991. Mixed transcortical aphasia. Neuroimaging study on two cases. Eur. Neurol. 31, 204–211.

Grossi, D., Marcone, R., Cinquegrana, T., Gallucci, M., 2013. On the differential nature of induced and incidental echolalia in autism. J. Intellect. Disabil. Res. 57 (10), 903–912.

Grossman, P., 2015. Mindfulness: awareness informed by an embodied ethic. Mindfulness 6 (1), 17–22.

Grynberg, D., Chang, B., Corneille, O., Maurage, P., Vermeulen, N., Berthoz, S., Luminet, O., 2012. Alexithymia and the processing of emotional facial expressions (EFEs): systematic review, unanswered questions and further perspectives. PLoS One 7 (8), e42429.

Gu, X., Gao, Z., Wang, X., Liu, X., Knight, R.T., Hof, P.R., Fan, J., 2012. Anterior insular cortex is necessary for empathetic pain perception. Brain 135 (9), 2726–2735.

Guest, S., Grabenhorst, F., Essick, G., Chen, Y., Young, M., McGlone, F., et al., 2007.
 Human cortical representation of oral temperature. Physiol. Behav. 92 (5), 975–984.
 Gunzelmann, T., Kupfer, J., Brähler, E., 2002. Alexithymia in the elderly general

population. Compr. Psychiatry 43 (1), 74–80.

Neuroscience and Biobehavioral Reviews 130 (2021) 470-508

Guttman, H., Laporte, L., 2002. Alexithymia, empathy, and psychological symptoms in a family context. Compr. Psychiatry 43 (6), 448–455.

Habib, M., Daquin, G., Milandre, L., Royere, M.L., Rey, M., Lanteri, A., et al., 1995. Mutism and auditory agnosia due to bilateral insular damage-Role of the insula in human communication. Neuropsychologia 33 (3), 327–339.

Hajcak, G., Donald, N.M.C., Simons, R.F., 2003. To err is autonomic: error-related brain potentials, ANS activity, and post-error compensatory behavior. Psychophysiology 40, 895–903.

Hamilton, A.F.D.C., 2013. Reflecting on the mirror neuron system in autism: a systematic review of current theories. Dev. Cogn. Neurosci. 3 (1), 91–105.

Han, Z.S., Zhang, E.T., Craig, A.D., 1998. Nociceptive and thermoreceptive lamina I neurons are anatomically distinct. Nat. Neurosci. 1 (3), 218–225.

Harms, M.B., Martin, A., Wallace, G.L., 2010. Facial emotion recognition in autism spectrum disorders: a review of behavioral and neuroimaging studies. Neuropsychol. Rev. 20 (3), 290–322.

Harmsen, I.E., 2019. Empathy in autism Spectrum disorder. J. Autism Dev. Disord. 49, 3939–3955.

Harper, R.M., Bandler, R., Spriggs, D., Alger, J.R., 2000. Lateralized and widespread brain activation during transient blood pressure elevation revealed by magnetic resonance imaging. J. Comp. Neurol. 417 (2), 195–204.

Harrison, N.A., Brydon, L., Walker, C., Gray, M.A., Steptoe, A., Dolan, R.J., Critchley, H. D., 2009. Neural origins of human sickness in interoceptive responses to inflammation. Biol. Psychiatry 66 (5), 415–422.

Harrison, A., Tchanturia, K., Treasure, J., 2010. Attentional bias, emotion recognition,

and emotion regulation in anorexia: State or trait? Biol. Psychiatry 68 (8), 755–761. Harrison, N.A., Gray, M.A., Gianaros, P.J., Critchley, H.D., 2010. The embodiment of emotional feelings in the brain. J. Neurosci. 30 (38), 12878–12884.

Harshaw, C., 2008. Alimentary epigenetics: a developmental psychobiological systems

view of the perception of hunger, thirst and satiety. Dev. Rev. 28 (4), 541–569. Harshaw, C., 2015. Interoceptive dysfunction: toward and integrated frameword for understanding somatic and affective disturbance in depression. Psychol. Bull. 141 (2), 311–363.

Hart, N., Mcgowan, J., Minati, L., Critchley, H.D., 2013. Emotional regulation and bodily sensation: interoceptive awareness is intact in borderline personality disorder. J. Pers. Disord. 27 (4), 506–518.

Harver, A., Katkin, E.S., Bloch, E., 1993. Signal-detection outcomes on heartbeat and respiratory resistance detection tasks in male and female subjects. Psychophysiology 30, 223–230.

Hasin, D., Kilcoyne, B., 2012. Comorbidity of psychiatric and substance use disorders in the United States: current issues and findings from the NESARC. Curr. Opin. Psychiatry 25 (3), 165–171.

Hassanpour, M.S., Simmons, W.K., Feinstein, J.S., Luo, Q., Lapidus, R.C., Bodurka, J., et al., 2018. The insular cortex dynamically maps changes in cardiorespiratory interoception. Neuropsychopharmacology 43 (2), 426–434.

Hatfield, T.R., Brown, R.F., Giummarra, M.J., Lenggenhager, B., 2017. Autism spectrum disorder and interoception: Abnormalities in global integration? Autism.

Heaton, P., Reichenbacher, L., Sauter, D., Allen, R., Scott, S., Hill, E., 2012. Measuring the effects of alexithymia on perception of emotional vocalizations in autistic spectrum disorder and typical development. Psychol. Med. 42 (11), 2453–2459.

Heinzel, A., Schäfer, R., Müller, H.W., Schieffer, A., Ingenhag, A., Eickhoff, S.B., et al., 2010. Increased activation of the supragenual anterior cingulate cortex during visual emotional processing in male subjects with high degrees of alexithymia: an eventrelated fMRI study. Psychother. Psychosom. 79 (6), 363–370.

Henry, J.D., Phillips, L.H., Crawford, J.R., Theodorou, G., Summers, F., 2006. Cognitive and psychosocial correlates of alexithymia following traumatic brain injury. Neuropsychologia 44 (1), 62–72.

Henry, J.D., Bailey, P.E., von Hippel, C., Rendell, P.G., Lane, A., 2010. Alexithymia in schizophrenia. J. Clin. Exp. Neuropsychol. 32 (8), 890–897.

Herbert, B.M., Pollatos, O., 2014. Attenuated interoceptive sensitivity in overweight and obese individuals. Eat. Behav. 15 (3), 445–448.

Herbert, B.M., Ulbrich, P., Schandry, R., 2007. Interoceptive sensitivity and physical effort: implications for the self-control of physical load in everyday life. Psychophysiology 44 (2), 194–202.

Herbert, B.M., Herbert, C., Pollatos, O., 2011. On the relationship between interoceptive awareness and alexithymia: Is interoceptive awareness related to emotional awareness? J. Pers. 79 (5), 1149–1175.

Herbert, B.M., Herbert, C., Pollatos, O., Weimer, K., Enck, P., Sauer, H., Zipfel, S., 2012. Effects of short-term food deprivation on interoceptive awareness, feelings and autonomic cardiac activity. Biol. Psychol. 89 (1), 71–79.

Herbert, B.M., Muth, E.R., Pollatos, O., Herbert, C., 2012. Interoception across modalities: on the relationship between cardiac awareness and the sensitivity for gastric functions. PLoS One 7 (5), e36646.

Herbert, B.M., Blechert, J., Hautzinger, M., Matthias, E., Herbert, C., 2013. Intuitive eating is associated with interoceptive sensitivity. Effects on body mass index. Appetite 70, 22–30.

Herman, A.M., Critchley, H.D., Duka, T., 2018. Risk-taking and impulsivity: the role of mood states and interoception. Front. Psychol. 9, 1625.

Hermesdorf, M., Berger, K., Baune, B.T., Wellmann, J., Ruscheweyh, R., Wersching, H., 2016. Pain sensitivity in patients with major depression: differential effect of pain sensitivity measures, somatic cofactors, and disease characteristics. J. Pain 17 (5), 606–616.

Hernández, S.E., Suero, J., Barros, A., González-Mora, J.L., Rubia, K., 2016. Increased grey matter associated with long-Term Sahaja yoga meditation: a voxel-based morphometry study. PLoS One 11 (3), 1–16. Heshmati, R., Jafari, E., Hoseinifar, J., Ahmadi, M., 2010. Comparative study of alexithymia in patients with schizophrenia spectrum disorders, non-psychotic disorders and normal people. Procedia: Soc. Behav.Sci. 5, 1084–1089.

- Heyes, C., Bird, G., 2007. Mirroring, association, and the correspondence problem. In: Haggard, P., Rossetti, Y., Kawato, M. (Eds.), Sensorimotor Foundations of Higher Cognition, Attention and Performance. Oxford University Press, Oxford, England, pp. 461–479.
- Hezel, D.M., Riemann, B.C., McNally, R.J., 2012. Emotional distress and pain tolerance in obsessive-compulsive disorder. J. Behav. Ther. Exp. Psychiatry 43 (4), 981–987.
 Hickman, L., Seyedsalehi, A., Cook, J.L., Bird, G., Murphy, J., 2020. The Relationship
- between Heartbeat Counting and Heartbeat Discrimination: a Meta-analysis. Hill, E., Berthoz, S., Frith, U., 2004. Brief report: cognitive processing of own emotions in
- Hili, E., Bertnöz, S., Frith, U., 2004. Brief report: cognitive processing of own emotions in individuals with autistic spectrum disorder and in their relatives. J. Autism Dev. Disord. 34 (2), 229–235.
- Hintistan, S., Cilingir, D., Birinci, N., 2013. Alexithymia among elderly patients with diabetes. Pak. J. Med. Sci. 29 (6), 1344–1348.
- Hobday, D.I., Aziz, Q., Thacker, N., Hollander, I., Jackson, A., Thompson, D.G., 2001. A study of the cortical processing of ano-rectal sensation using functional MRI. Brain 124 (2), 361–368.
- Hobson, H., Hogeveen, J., Brewer, R., Catmur, C., Gordon, B., Krueger, F., et al., 2018. Language and alexithymia: evidence for the role of the inferior frontal gyrus in acquired alexithymia. Neuropsychologia 111, 229–240.
- Hobson, H., Brewer, R., Catmur, C., Bird, G., 2019. The role of language in alexithymia: moving towards a multiroute model of alexithymia. Emot. Rev. 11 (3), 247–261.

Hoehn-Saric, R., Mcleod, D.R., 2000. Anxiety and arousal: physiological changes and their perception. J. Affect. Disord. 61, 217–224.

- Hoffman, W.F., Schwartz, D.L., Meiri, G., Stevens, A.A., Mitchell, S.H., 2008. Cortical activation during delay discounting in abstinent methamphetamine dependent individuals. Psychopharmacology 201, 183–193.
- Hogeveen, J., Bird, G., Chau, A., Krueger, F., Grafman, J., 2016. Acquired alexithymia following damage to the anterior insula. Neuropsychologia 82, 142–148.
- Hölzl, R., Erasmus, L.P., Möltner, A., 1996. Detection, discrimination and sensation of visceral stimuli. Biol. Psychol. 42 (1–2), 199–214.
- Honkalampi, K., Hintikka, J., Laukkanen, E., Lehtonen, J., Viinamäki, H., 2001. Alexithymia and depression: a prospective study of patients with major depressive disorder. Psychosomatics 42 (3), 229–234.
- Honkalampi, K., Koivumaa-Honkanen, H., Hintikka, J., Antikainen, R., Haatainen, K., Tanskanen, A., Viinamäki, H., 2004. Do stressful life-events or sociodemographic variables associate with depression and alexithymia among a general population? - A 3-year follow-up study. Compr. Psychiatry 45 (4), 254–260.
- Honkalampi, K., Tolmunen, T., Hintikka, J., Rissanen, M.-L., Kylmä, J., Laukkanen, E., 2009. The prevalence of alexithymia and its relationship with Youth Self-Report problem scales among Finnish adolescents. Compr. Psychiatry 50 (3), 263–268.
- Hoorn, J., Van, Crone, E.A., Leijenhorst, L.Van., 2016. Hanging out with the right crowd: peer influence on risk-taking behavior in adolescence. J. Res. Adolesc. 27 (1), 189–200.
- Horing, B., Kugel, H., Brenner, V., Zipfel, S., 2013. Perception and pain thresholds for cutaneous heat and cold, and rectal distension: associations and disassociations. Neurogastroenterol. Motil. 25, e791–e802.
 Housiaux, M., Luminet, O., Broeck, N.V., Dorchy, H., 2010. Alexithymia is associated
- Housiaux, M., Luminet, O., Broeck, N.V., Dorchy, H., 2010. Alexithymia is associated with glycaemic control of children with type 1 diabetes. Diabetes Metab. 6 (1), 455–462.
- Ibañez, A., Gleichgerrcht, E., Manes, F., 2010. Clinical effects of insular damage in humans. Brain Struct. Funct. 214 (5–6), 397–410.
- Ihme, K., Dannlowski, U., Lichev, V., Stuhrmann, A., Grotegerd, D., Rosenberg, N., et al., 2013. Alexithymia is related to differences in gray matter volume: a voxel-based morphometry study. Brain Res. 1491, 60–67.
- Ikoma, A., Steinhoff, M., Ständer, S., Yosipovitch, G., Schmelz, M., 2006. The neurobiology of itch. Nat. Rev. Neurosci. 7 (7), 535–547.
- Iodice, P., Porciello, G., Bufalari, I., Barca, L., Pezzulo, G., 2019. An interoceptive illusion of effort induced by false heart-rate feedback. Proc. Natl. Acad. Sci. U.S.A. 116 (28), 13897–13902.
- Ishii, H., Ohara, S., Tobler, P.N., Tsutsui, K.-I., Iijima, T., 2012. Inactivating anterior insular cortex reduces risk taking. J. Neurosci. 32 (45), 16031–16039.
- Jakubczyk, A., Skrzeszewski, J., Trucco, E.M., Suszek, H., Zaorska, J., Nowakowska, M., et al., 2019. Interoceptive accuracy and interoceptive sensibility in individuals with alcohol use disorder – Different phenomena with different clinical correlations? Drug Alcohol Depend. 198, 34–38.
- James, W., 1894. The physical basis of emotion. Psychol. Rev. 1, 516-529.
- Jang, E.H., Park, B.J., Park, M.S., Kim, S.H., Sohn, J.H., 2015. Analysis of physiological signals for recognition of boredom, pain, and surprise emotions. J. Physiol. Anthropol. 34 (1), 25.
- Jarrahi, B., Mantini, D., Balsters, J.H., Michels, L., Kessler, T.M., Mehnert, U., Kollias, S. S., 2015. Differential functional brain network connectivity during visceral interoception as revealed by independent component analysis of fMRI TIME-Series. Hum. Brain Mapp. 36, 4438–4468.
- Jimerson, D.C., Wolfe, B.E., Franko, D.L., Covino, N.A., Sifneos, P.E., 1994. Alexithymia ratings in bulimia nervosa: clinical correlates. Psychosom. Med. 56 (2), 90–93.

Joel, D., Zohar, O., Afek, M., Hermesh, H., Lerner, L., Kuperman, R., et al., 2005. Impaired procedural learning in obsessive-compulsive disorder and Parkinson's disease, but not in major depressive disorder. Behav. Brain Res. 157 (2), 253–263.

Jørgensen, M.M., Zachariae, R., Skytthe, A., Kyvik, K., 2007. Genetic and environmental factors in alexithymia: a population-based study of 8,785 Danish twin pairs. Psychother. Psychosom. 76 (6), 369–375.

- John, O.P., Gross, J.J., 2004. Healthy and unhealthy emotion regulation: personality processes, individual differences, and life span development. J. Pers. 72 (6), 1301–1333.
- Johnson, S.A., Yechiam, E., Murphy, R.R., Queller, S., Stout, J.C., 2006. Motivational processes and autonomic responsivity in Asperger's disorder: evidence from the Iowa Gambling Task. J. Int. Neuropsychol. Soc. 12, 668–676.

Jones, G.E., 1994. Perception of visceral sensations: a review of recent findings, methodologies, and future directions. In: Jennings, J.R., Ackles, P.K. (Eds.), Advances in Psychophysiology, Vol. 5. Jessica Kingsley, London.

- Jones, Gary E., Hollandsworth, J.G., 1981. Heart Rate Discrimination Before and After Exercise-Induced Augmented Cardiac Activity. Psychophysiology 18 (3), 252–257.
- Jones, M.P., Roth, L.M., Crowell, M.D., 2005. Symptom reporting by functional dyspeptics during the water load test. Am. J. Gastroenterol. 100 (6), 1334.

Jones, C.L., Ward, J., Critchley, H.D., 2010. The neuropsychological impact of insular cortex lesions. J. Neurol. Neurosurg. Psychiatr. 81 (6), 611–618.

- Jongen, S., Axmacher, N., Kremers, N., a, W., Hoffmann, H., Limbrecht-Ecklundt, K., Traue, H.C., Kessler, H., 2014. An investigation of facial emotion recognition impairments in alexithymia and its neural correlates. Behav. Brain Res. 271, 129–139.
- Jordan, R., 1993. The nature of linguistic and communication difficulties of children with autism. In: Messer, D., Turner, G. (Eds.), Critical Influences on Child Language Acquisition and Development. St. Martin's Press, New York.

Joukamaa, M., Saarijärvi, S., Muuriaisniemi, M.L., Salokangas, R.K., 1996. Alexithymia in a normal elderly population. Compr. Psychiatry 37 (2), 144–147.

- Joukamaa, Matti, Taanila, A., Miettunen, J., Karvonen, J.T., Koskinen, M., Veijola, J., 2007. Epidemiology of alexithymia among adolescents. J. Psychosom. Res. 63, 373–376.
- Jovanovic, T., Kazama, A., Bachevalier, J., Davis, M., 2012. Impaired safety signal learning may be a biomarker of PTSD. Neuropharmacology 62 (2), 695–704.
- Joybari, M.T., 2014. Depression and interpersonal problems in adolescents: their relationship with alexithymia and coping styles. Iran. J. Psychiatry Behav. Sci. 8 (4), 38–45.
- Junghanns, K., Tietz, U., Dibbelt, L., Kuether, M., Jurth, R., Ehrenthal, D., et al., 2005. Attenuated salivary cortisol secretion under cue exposure is associated with early relapse. Alcohol Alcohol. 40 (1), 80–85.
- Kandasamy, N., Sarah, N.G., Page, L., Hardy, B., Critchley, H.D., Gurnell, M., Coates, J. M., 2016. Interoceptive ability predicts survival on a London trading floor. Sci. Rep. 6, 32986.
- Kang, J.I., Namkoong, K., Yoo, S.W., Jhung, K., Kim, S.J., 2012. Abnormalities of emotional awareness and perception in patients with obsessive–compulsive disorder. J. Affect. Disord. 141 (2–3), 286–293.
- Kanner, L., 1943. Autistic disturbances of affective contact. Nerv. Child 2 (3), 217–250. Kanner, L., 1946. Irrelevant and metaphorical language in early infantile autism. Am. J. Psychiatry 103, 242–246.
- Kano, M., Fukudo, S., 2013. The alexithymic brain: the neural pathways linking alexithymia to physical disorders. Biopsychosoc. Med. 7 (1), 1.
- Kano, M., Fukado, S., Jiro, G., Kamachi, M., Tagawa, M., Mochizuki, H., et al., 2003. Specific brain processing of facial expressions in people with alexithymia: an H2150-PET study. Brain 126 (6), 1474–1484.
- Kano, M., Hamaguchi, T., Itoh, M., Yanai, K., Fukudo, S., 2007. Correlation between alexithymia and hypersensitivity to visceral stimulation in human. Pain 132 (3), 252–263.
- Kano, M., Masatochi, I., Fukudo, S., 2011. Neural substrates of decision making as measured with the Iowa Gambling Task in men with alexithymia. Psychosom. Med. 73, 588–597.
- Kaplow, J.B., Widom, C.S., 2007. Age of onset of child maltreatment predicts long-term mental health outcomes. J. Abnorm. Psychol. 116 (1), 176–187.
- Karnath, H.O., Baier, B., 2010. Right insula for our sense of limb ownership and selfawareness of actions. Brain Struct. Funct. 1–7.
- Karukivi, M., Pölönen, T., Vahlberg, T., Saikkonen, S., Saarijärvi, S., 2014. Stability of alexithymia in late adolescence: results of a 4-year follow-up study. Psychiatry Res. 219 (2), 386–390.
- Katkin, E.S., Wiens, S., Ohman, A., 2001. Nonconscious fear conditioning, visceral perception, and the development of gut feelings. Psychol. Sci. 12 (5), 366–370.
- Kazour, F., Richa, S., Desmidt, T., Lemaire, M., Atanasova, B., El Hage, W., 2017. Olfactory and gustatory functions in bipolar disorders: a systematic review. Neurosci. Biobehav. Rev. 80, 69–79.
- Kench, S., Irwin, H.J., 2000. Alexithymia and childhood family environment. J. Clin. Psychol. 56 (6), 737–745.
- Kenyon, L.W., Ketterer, M.W., Gheorghiade, M., Goldstein, S., 1991. Psychological factors related to prehospital delay during acute myocardial infarction. Circulation 84 (5), 1969–1976.
- Kerr-Gaffney, J., Harrison, A., Tchanturia, K., 2019. Cognitive and affective empathy in eating disorders: a systematic review and meta-analysis. Front. Psychiatry 10, 102.

Kessler, R.C., Berglund, P., Demler, O., Jin, R., Merikangas, K.R., Walters, E.E., 2005. Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the national comorbidity survey replication. Arch. Gen. Psychiatry 62, 593–768.

- Kessler, R.C., Chui, W.T., Demler, O., Walters, E.E., 2005. Prevalence, severity, and comorbidity of 12-Month DSM-IV disorders in the national comorbidity survey replication. Arch. Gen. Psychiatry 62, 617–709.
- Kever, A., Pollatos, O., Vermeulen, N., Grynberg, D., 2015. Interoceptive sensitivity facilitates both antecedent- and response-focused emotion regulation strategies. Pers. Individ. Dif. 87, 20–23.
- Khalsa, Sahib S., Lapidus, R.C., 2016. Can interoception improve the pragmatic search for biomarkers in psychiatry? Front. Psychiatry 7, 121.

Khalsa, S.S., Rudrauf, D., Damasio, A.R., Davidson, R.J., Lutz, A., Tranel, D., 2008. Interoceptive awareness in experinced meditators. Psychophysiology 45 (4), 671–677.

Khalsa, Sahib S., Rudrauf, D., Tranel, D., 2009. Interoceptive awareness declines with age. Psychophysiology 46 (6), 1130–1136.

Khalsa, Sahib S., Rudrauf, D., Feinstein, J.S., Tranel, D., 2009. The pathways of interoceptive awareness. Nat. Neurosci. 12 (12), 1494–1496.

Khalsa, S.S., Rudrauf, D., Sandesara, C., Olshansky, B., Tranel, D., 2009. Bolus isoproterenol infusions provide a reliable method for assessing interoceptive awareness. Int. J. Psychophysiol. 72 (1), 34–45.

- Khalsa, Sahib S., Adolphs, R., Cameron, O.G., Critchley, H.D., Davenport, P.W., Feinstein, J.S., et al., 2018. Interoception and mental health: a roadmap. Biol. Psychiatry 3, 501–513.
- Khalsa, Sahib, Rudrauf, D., Hassanpour, M.S., Davidson, R.J., Tranel, D., 2020. The practice of meditation is not associated with improved interoceptive awareness of the heartbeat. Psychophysiology 57 (2), e13479.
- Khosravani, V., Alvani, A., Sharifi Bastan, F., Jamaati Ardakani, R., Akbari, H., 2016. The alexithymia, cognitive emotion regulation, and physical symptoms in Iranian asthmatic patients. Pers. Individ. Dif. 101, 214–219.
- Kilts, C.D., Schweitzer, J.B., Quinn, C.K., Gross, R.E., Faber, T.L., Muhammad, F., et al., 2001. Neural activity related to drug craving in cocaine addiction. Arch. Gen. Psychiatry 58, 334–341.
- Kilts, C.D., Gross, R.E., Ely, T.D., Drexler, K.P.G., 2004. The neural correlates of cueinduced craving in cocaine-dependent women. Am. J. Psychiatry 161, 233–241.

Kimhy, D., Gill, K.E., Brucato, G., Vakhrusheva, J., Arndt, L., Gross, J.J., Girgis, R.R., 2016. The impact of emotion awareness and regulation on social functioning in individuals at clinical high risk for psychosis. Psychol. Med. 46, 2907–2918.

Kinnaird, E., Stewart, C., Tchanturia, K., 2018. Taste sensitivity in anorexia nervosa: a systematic review. Int. J. Eat. Disord. 51, 771–784.

Kinnaird, E., Stewart, C., Tchanturia, K., 2019. Investigating alexithymia in autism: a systematic review and meta-analysis. Eur. Psychiatry 55, 80–89.

Kinnaird, E., Stewart, C., Tchanturia, K., 2020. Interoception in Anorexia Nervosa: exploring associations with alexithymia and autistic traits. Front. Psychiatry 11.

Kinomura, S., Kawashima, R., Yamada, K., Ono, S., Itoh, M., Yoshioka, S., et al., 1994. Functional anatomy of taste perception in the human brain studied with positron emission tomography. Brain Res. 659, 263–266.

Klabunde, M., Acheson, D., Boutelle, K., Matthews, S., Kaye, W., 2013. Interoceptive sensitivity deficits in women recovered from bulimia nervosa. Eat. Behav. 14 (4), 488–492.

Klabunde, M., Juszczak, H., Jordan, T., Baker, J.M., Bruno, J., Carrion, V., Reiss, A.L., 2019. Functional neuroanatomy of interoceptive processing in children and adolescents: a pilot study. Sci. Rep. 9 (1), 16184.

Kleckner, I.R., Baumann Wormwood, J., Simmons, W.K., Barrett, L.F., Quigley, K., 2015. Methodological recommnedations for a heartbeat detection-based measure of interoceptive sensitivity. Psychophysiology 52 (11), 1432–1440.

Klin, A., Jones, W., Schultz, R., Volkmar, F., Cohen, D., 2002. Visual fixation patterns during viewing of naturalistic social situations as predictors of social competence in individuals with autism. Arch. Gen. Psychiatry 59 (9), 809–816.

Klukowski, M., Wasilewska, J., Lebensztejn, D., 2015. Sleep and gastrointestinal disturbances in autism spectrum disorder in children. Dev. Period Med. 19 (2), 157–161.

Knapp, K., Ring, C., Brener, J., 1997. Sensitivity to mechanical stimuli and the role of general sensory and perceptal processes in heartbeat detection. Psychophysiology 34, 467–473.

Knapp-Kline, K., Kline, J.P., 2005. Heart rate, heart rate variability, and heartbeat detection with the method of constant stimuli: slow and steady wins the race. Biol. Psychol. 69 (3), 387–396.

Knoll, J.F., Hodapp, V., 1992. A comparison between two methods for assessing heartbeat perception. Psychophysiology 29 (2), 218–222.

Kobayashi, I., Boarts, J.M., Delahanty, D.L., 2007. Polysomnographically measured sleep abnormalities in PTSD: a meta-analytic review. Psychophysiology 44, 660–669 Vech. A. Pollyton O. 2014. Untergraphic review. Psychophysiology 44, 660–669

Koch, A., Pollatos, O., 2014. Interoceptive sensitivity, body weight and eating behavior in children: a prospective study. Front. Psychol. 5 (September), 1–11.

Koch, K.L., Hong, S.P., Xu, L., 2000. Reproducibility of gastric myoelectrical activity and the Water Load Test in patients with dysmotility-like dyspepsia symptoms and in control subjects. J. Clin. Gastroenterol. 31 (2), 125–129.

Koegel, R.L., Rincover, A., Egel, A.L., 1982. Educating and Understanding Autistic Children. College Hill., San Diego.

Kollenbaum, V.E., Dahme, B., Kirchner, G., 1996. "Interoception" of heart rate, blood pressure, and myocardial metabolism during ergometric work load in healthy young subjects. Biol. Psychol. 42 (1–2), 183–197.

Kong, J., White, N.S., Kwong, K.K., Vangel, M.G., Rosman, I.S., Gracely, R.H., Gollub, R. L., 2006. Using fMRI to dissociate sensory encoding from cognitive evaluation of heat pain intensity. Hum. Brain Mapp. 27 (9), 715–721.

- Kotov, R., Chang, S., Fochtmann, L.J., Mojtabai, R., Carlson, G.A., Sedler, M.J., Bromet, E.J., 2011. Schizophrenia in the Internalizing-Externalizing Framework: A Third Dimension? Schizophr. Bull. 37 (6), 1168–1178.
- Kotov, R., Ruggero, C.J., Krueger, R.F., Watson, D., Yuan, Q., Zimmerman, 2011. New dimensions in the quantitative classification of mental illness. Arch. Gen. Psychiatry 68 (10), 1003–1011.
- Kotov, R., Krueger, R.F., Watson, D., Achenbach, T.M., Althoff, R.R., Bagby, R.M., et al., 2017. The hierarchical taxonomy of psychopathology (HiTOP): A dimensional alternative to traditional nosologies. J. Abnorm. Psychol. 126 (4), 454–477.

Krolak-Salmon, P., Hénaff, M.A., Isnard, J., Tallon-Baudry, C., Guénot, M., Vighetto, A., et al., 2003. An attention modulated response to disgust in human ventral anterior insula. Ann. Neurol. 53 (4), 446–453. Krout, K.E., Loewy, A.D., 2000. Periaqueductal gray matter projections to midline and intralaminar thalamic nuclei of the rat. J. Comp. Neurol. 424 (1), 111–141.

Krueger, R.F., Caspi, A., Moffitt, T.E., Silva, P.A., 1998. The structure and stability of common mental disorders (DSM-III-R): a longitudinal-epidemiological study. J. Abnorm. Psychol. 107 (2), 216–227.

- Krukoff, T.L., Harris, K.H., Jhamandas, J.H., 1993. Efferent projections from the parabrachial nucleus demonstrated with the anterograde tracer Phaseolus vulgaris leucoagglutinin. Brain Res. Bull. 30 (1–2), 163–172.
- Kruschwitz, J.D., Kausch, A., Brovkin, A., Keshmirian, A., Paulus, M.P., Goschke, T., Walter, H., 2019. Self-control is linked to interoceptive inference: craving regulation and the prediction of aversive interoceptive states induced with inspiratory breathing load. Cognition 193, 104028.
- Kuhnen, C.M., Knutson, B., 2005. The neural basis of financial risk taking. Neuron 47 (5), 763–770.

Kuhtz-Buschbeck, J.P., van der Horst, C., Pott, C., Wolff, S., Nabavi, A., Jansen, O., Jünemann, K.P., 2005. Cortical representation of the urge to void: a functional magnetic resonance imaging study. J. Urol. 174 (1), 1477–1481.

Kutscheidt, K., Dresler, T., Hudak, J., Barth, B., Blume, F., Ethofer, T., et al., 2019. Interoceptive awareness in patients with attention - deficit/hyperactivity disorder (ADHD). Adhd Atten. Deficit Hyperact. Disord. 11 (4), 395–401.

Lackner, R.J., Fresco, D.M., 2016. Interaction effect of brooding rumination and interoceptive awareness on depression and anxiety symptoms. Behav. Res. Ther. 85, 43–52.

Lahey, B.B., Applegate, B., Hakes, J.K., Zald, D.H., Hariri, A.R., Rathouz, P.J., 2012. Is there a general factor of prevalent psychopathology during adulthood? J. Abnorm. Psychol. 121 (4), 971–977.

Lane, A.E., Young, Æ.R.L., Baker, Æ.A.E.Z., Angley, M.T., 2010. Sensory processing subtypes in autism: association with adaptive behavior. J. Autism Dev. Disord. 40, 112–122.

Lane, S.J., Reynolds, S., Dumenci, L., 2012. Sensory overresponsivity and anxiety in typically developing children and children With autism and attention deficit hyperactivity disorder: Cause or coexistence? Am. J. Occup. Ther. 66, 595–603.

- Lane, A.E., Molloy, C.A., Bishop, S.L., 2014. Classification of Children with Autism Spectrum Disorder by Sensory Subtype: A Case for Sensory-Based Phenotypes. Autism Res. 7 (3), 322–333.
- Lange, C., 1885. The Emotions. Williams & Wilkins, Baltimore, Md.
- Lapidus, R.C., Puhl, M., Kuplicki, R., Stewart, J.L., Paulus, M.P., Rhudy, J.L., et al., 2020. Heightened affective response to perturbation of respiratory but not pain signals in eating, mood, and anxiety disorders. PLoS One 15, 1–21.

Lattimore, P., Malinowski, P., Mead, B., Irwin, L., Grice, L., Carson, R., 2017. 'I can't accept that feeling': relationships between interoceptive awareness, mindfulness and eating disorder symptoms in females with, and at-risk of an eating disorder. Psychiatry Res. 247, 163–171.

Lazarov, A., Dar, R., Oded, Y., Liberman, N., 2010. Are obsessive-compulsive tendencies related to reliance on external proxies for internal states? Evidence from biofeedback-aided relaxation studies. Behav. Res. Ther. 48 (6), 516–523.

Leekam, S.R., Nieto, C., Libby, S.J., Wing, L., Gould, J., 2007. Describing the sensory abnormalities of children and adults with autism. J. Autism Dev. Disord. 37 (5), 894–910.

Leichnetz, G.R., 1972. Relationship of spontaneous vagal activity to wakefulness and sleep in the cat. Exp. Neurol. 35, 194–210.

Lemche, E., Klann-Delius, G., Koch, R., Joraschky, P., 2004. Mentalizing language development in a longitudinal attachment sample: implications for alexithymia. Psychother. Psychosom. 73 (6), 366–374.

Lenggenhager, B., Azevedo, R.T., Mancini, A., Aglioti, S.M., 2013. Listening to your heart and feeling yourself: effects of exposure to interoceptive signals during the ultimatum game. Exp. Brain Res. 230 (2), 233–241.

Levinson, D.J., 1978. Eras: the anatomy of the life cycle. Psychiatr. Opin. 15 (9), 39–48.

Li, D., Zucker, N.L., Kragel, P.A., Covington, V.E., Labar, K.S., 2017. Adolescent development of insula-dependent interoceptive regulation. Dev. Sci. 20, e12438.

Lieberman, M.D., Eisenberger, N.I., Crockett, M.J., Tom, S.M., Pfeifer, J.H., Way, B.M., 2007. Putting feelings into words: affect labeling disrupts amygdala activity in response to affective stimuli. Psychol. Sci. 18 (5), 421–428.

Light, A.R., Willcockson, H.H., 1999. Spinal laminae I-II neurons in rat recorded in vivo in whole cell, tight seal configuration: properties and opioid responses. J. Neurophysiol. 82 (6), 3316–3326.

- Liotti, M., Brannan, S., Egan, G., Shade, R., Madden, L., Abplanalp, B., et al., 2001. Brain responses associated with consciousness of breathlessness (air hunger). Proc. Natl. Acad. Sci. 98 (4), 2035–2040.
- London, E.D., Berman, S.M., Voytek, B., Simon, S.L., Mandelkern, M.A., Monterosso, J., et al., 2005. Cerebral metabolic dysfunction and impaired vigilance in recently abstinent methamphetamine abusers. Biol. Psychiatry 58, 770–778.

Longarzo, M., D'Olimpio, F., Chiavazzo, A., Santangelo, G., Trojano, L., Grossi, D., 2015. The relationships between interoception and alexithymic trait. The Self-Awareness Questionnaire in healthy subjects. Front. Psychol. 6 (August), 1149.

Ludewig, S., Geyer, M.A., Ramseier, M., Vollenweider, F.X., Rechsteiner, E., Cattapan-Ludewig, K., 2005. Information-processing deficits and cognitive dysfunction in panic disorder. J. Psychiatry Neurosci. 30 (1), 37–43.

Ludwick-Rosenthal, R., Neufeld, R.W., 1985. Heart beat interoception: a study of individual differences. Int. J. Psychophysiol. 31 (1), 57–65.

Luijten, M., Machielsen, M.W.J., Veltman, D.J., Hester, R., Haan, L.De, Franken, I.H.A., 2014. Systematic review of ERP and fMRI studies investigating inhibitory control and error processing in people with substance dependence and behavioural addictions. J. Psychiatry Neurosci. 39 (3), 149–169.

Luke, L., Clare, I.C.H., Ring, H., Redley, M., Watson, P., 2011. Decision-making difficulties experienced by adults with autism spectrum conditions. Autism 16 (6), 612–621.

Łukowska, M., Sznajder, M., Wierzchoń, M., 2018. Error-related cardiac response as information for visibility judgements. Sci. Rep. 8, 1131.

Luminet, O., Bagby, R.M., Taylor, G.J., 2001. An evaluation of the absolute and relative stability of alexithymia in patients with major depression. Psychother. Psychosom. 70 (5), 254–260.

Luminet, Olivier, Vermeulen, N., Demaret, C., Taylor, G.J., Bagby, R.M., 2006. Alexithymia and levels of processing: evidence for an overall deficit in remembering emotion words. J. Res. Pers. 40 (5), 713–733.

Lumley, M.A., Smith, J.A., Longo, D.J., 2002. The relationship of alexithymia to pain severity and impairment among patients with chronic myofascial pain Comparisons with self-efficacy, catastrophizing, and depression. J. Psychosom. Res. 53, 823–830.

Lumley, M.A., Neely, L.C., Burger, A.J., 2007. The assessment of alexithymia in medical settings: implications for understanding and treating health problems. J. Pers. Assess. 89 (3), 230–246.

Lutz, A.P.C., Schulz, A., Voderholzer, U., Koch, S., van Dyck, Z., Vögele, C., 2019. Enhanced cortical processing of cardio-afferent signals in anorexia nervosa. Clin. Neurophysiol. 130 (9), 1620–1627.

Lyvers, M., Duric, N., Thorberg, F.A., 2014. Caffeine use and alexithymia in university students. J. Psychoactive Drugs 46 (4), 340–346.

Ma, N., Liu, Y., Fu, X., Li, N., Wang, C., Zhang, H., et al., 2011. Abnormal brain defaultmode network functional connectivity in drug addicts. PLoS One 6 (1), e16560.

Macmillan, H.L., Streiner, D.L., Ph, D., Lin, E., Ph, D., Boyle, M.H., et al., 2001. Childhood abuse and lifetime psychopathology in a community sample. Am. J. Psychiatry 158, 1878–1883.

Mai, S., Ki, C., Georgiou, E., Pollatos, O., 2018. Interoception is associated with

heartbeat-evoked brain potentials (HEPs) in adolescents. Biol. Psychol. 137, 24–33. Maister, L., Tang, T., Tsakiris, M., 2017. Neurobehavioral evidence of interoceptive sensitivity in early infancy. ELife 6, 1–12.

Makovac, E., Garfinkel, S.N., Bassi, A., Basile, B., Macaluso, E., Cercignani, M., et al., 2015. Effect of parasympathetic stimulation on brain activity during appraisal of fearful expressions. Neuropsychopharmacology 40 (7), 1649–1658.

Mann, L., Wise, T., Trinidad, A., Kohanski, R., 1995. Alexithymia, affect recognition, and five factors of personality in substance abusers. Percept. Mot. Skills 81 (1), 35–40. Manninen, M., Therman, S., Suvisaari, J., Ebeling, H., Moilanen, I., Huttunen, M.,

Maininen, M., Hierman, S., Suvisaari, J., Ebening, H., Mohanen, L., Huttmen, M., Joukamaa, M., 2011. Alexithymia is common among adolescents with severe disruptive behavior. J. Nerv. Ment. Dis. 199 (7), 506–509.

Marchesi, C., Fontò, S., Balista, C., Cimmino, C., Maggini, C., 2005. Relationship between alexithymia and panic disorder: a longitudinal study to answer an open question. Psychother. Psychosom. 74 (1), 56–60.

Marco, E.J., Hinkley, L.B.N., Hill, S.S., Nagarajan, S.S., 2011. Sensory processing in autism: a review of neurophysiologic findings. Pediatric Res. Re 69 (5), 48–54.

Marshall, A.C., Gentsch, A., Schröder, L., Schütz-Bobach, S., 2018. Cardiac interoceptive learning is modulated by emotional valence perceived from facial expressions. Soc. Cogn. Affect. Neurosci. 13 (7), 677–686.

Martin, J.B., 1985. The stress-alexithymia hypothesis: theoretical and empirical considerations. Psychother. Psychosom. 43, 169–176.

Martin, J.B., Pihl, R.O., 1986. Influence of alexithymie characteristics on physiological and subjective stress responses in normal individuals. Psychother. Psychosom. 45, 66–77.

Martínez-Sánchez, F., Ato-García, M., Ortiz-Soria, B., 2003. Alexithymia - State or trait? Span. J. Psychol. 6 (1), 51–59.

Martins, Y., Young, R.L., Robson, D.C., 2008. Feeding and eating behaviors in children with autism and typically developing children. J. Autism Dev. Disord. 38 (10), 1878–1887.

Marucci, S., Ragione, L.D., De Iaco, G., Mococci, T., Vicini, M., Guastamacchia, E., Triggiani, V., 2018. Anorexia nervosa and comorbid psychopathology. Endocrine, Metab. & Immune Dis. Drug Targets 18 (4), 316–324.

Mata, F., Verdejo-Roman, J., Soriano-Mas, C., Verdejo-Garcia, A., 2015. Insula tuning towards external eating versus interoceptive input in adolescents with overweight and obesity. Appetite 93, 24–30.

Mattila, A.K., Salminen, J.K., Nummi, T., Joukamaa, M., 2006. Age is strongly associated with alexithymia in the general population. J. Psychosom. Res. 61 (5), 629–635.

Maurage, P., Grynberg, D., Noël, X., Joassin, F., Philippot, P., Hanak, C., et al., 2011. Dissociation between affective and cognitive empathy in alcoholism: a specific deficit for the. Alcoholism: Clin. Eperimental Res. 35 (9), 1–7.

May, A.C., Stewart, J.L., Tapert, S.F., Paulus, M.P., 2014. The effect of age on neural processing of pleasant soft touch stimuli. Front. Behav. Neurosci. 8 (February), 52. Mazefsky, C.A., White, S.W., 2014. Emotion regulation: concepts & practice in autism

spectrum disorder. Child Adolesc. Psychiatr. Clin. N. Am. 23 (1), 15–24.

Mazzola, L., Faillenot, I., Barral, F.G., Mauguière, F., Peyron, R., 2012. Spatial segregation of somato-sensory and pain activations in the human operculo-insular cortex. NeuroImage 60 (1), 409–418.

McCubbin, J.A., Merritt, M.M., Sollers I.I.I., J.J., Evans, M.K., Zonderman, A.B., Lane, R. D., Thayer, J.F., 2011. Cardiovascular emotional dampening: the relationship between blood pressure and recognition of emotion. Psychosom. Med. 73 (9), 743–750.

McDonald, S., Rosenfeld, J., Henry, J.D., Togher, L., Tate, R., Bornhofen, C., 2011. Emotion perception and Alexithymia in people with severe traumatic brain injury: one disorder or two?: a preliminary investigation. Brain Impair. 12 (3), 165–178.

McIntosh, R.C., Ironson, G., Antoni, M., Kumar, M., Fletcher, M.A., Schneiderman, N., 2014. Alexithymia is linked to neurocognitive, psychological, neuroendocrine, and immune dysfunction in persons living with HIV. Brain Behav. Immun. 36, 165–175. Medford, N., Critchley, H.D., 2010. Conjoint activity of anterior insular and anterior cingulate cortex: awareness and response. Brain Struct. Funct. 214 (5–6), 535–549.

Meessen, J., Mainz, V., Gauggel, S., Volz-Sidiropoulou, E., Sütterlin, S., Forkmann, T., 2016. The relationship between interoception and metacognition: a pilot study. J. Psychophysiol. 30 (2), 76–86.

Mehling, W., 2016. Differentiating attention styles and regulatory aspects of self-reported interoceptive sensibility. Philos. Trans. Biol. Sci. 371, 20160013.

Mehling, W.E., Price, C., Daubenmier, J.J., Acree, M., Bartmess, E., Stewart, A., 2012. The multidimensional assessment of interoceptive awareness (MAIA). PLoS One 7 (11), e48230.

Melloni, M., Sedeño, L., Couto, B., Reynoso, M., Gelormini, C., Favaloro, R., et al., 2013. Preliminary evidence about the effects of meditation on interoceptive sensitivity and social cognition. Behav. Brain Funct. 9 (1), 47.

Michael, R.P., 1957. Treatment of a case of compulsive swearing. Br. Med. J. 1, 1506–1508.

Michal, M., Reuchlein, B., Adler, J., Reiner, I., Beutel, M.E., Vögele, C., et al., 2014. Striking discrepancy of anomalous body experiences with normal interoceptive accuracy in depersonalization-derealization disorder. PLoS One 9 (2), 15–21.

Michalczuk, R., Bowden-Jones, H., Verdejo-Garcia, A., Clark, L., 2011. Impulsivity and cognitive distortions in pathological gamblers attending the UK National Problem Gambling Clinic: a preliminary report. Psychol. Med. 41, 2625–2635.

Migliorini, R., Stewart, J.L., May, A.C., Tapert, S.F., Paulus, M.P., 2013. What do you feel? Adolescent drug and alcohol users show altered brain response to pleasant interoceptive stimuli. Drug Alcohol Depend. 133 (2), 661–668.

Mikkelsen, M.B., O'Toole, M.S., Lyby, M.S., Wallot, S., Mehlsen, M., 2019. Emotional reactivity and interoceptive sensitivity: exploring the role of age. Psychon. Bull. Rev. 26 (4), 1440–1448.

Mikolajczak, M., Luminet, O., 2006. Is alexithymia affected by situational stress or is it a stable trait related to emotion regulation? Pers. Individ. Dif. 40 (7), 1399–1408.

Miller, L.C., Murphy, R., Buss, A.H., 1981. Consciousness of body: private and public. J. Pers. Soc. Psychol. 41 (2), 397–406.

Mizuhiki, T., Richmond, B.J., Shidara, M., 2012. Encoding of reward expectation by monkey anterior insular neurons. J. Neurophysiol. 107 (11), 2996–3007.

Mnif, L., Damak, R., Mnif, F., Ouanes, S., Abid, M., Jaoua, A., Masmoudi, J., 2014. Alexithymia impact on type 1 and type 2 diabetes: a case-control study. Ann. Endocrinol. (Paris) 75 (4), 213–219.

Moberg, E., 1983. The role of cutaneous afferents in position sense, kinaesthesia, and motor function of the hand. Brain 106, 1–19.

Mochizuki, H., Sadato, N., Saito, D.N., Toyoda, H., Tashiro, M., Okamura, N., Yanai, K., 2007. Neural correlates of perceptual difference between itching and pain: a human fMRI study. NeuroImage 36 (3), 706–717.

Moisset, X., Bouhassira, D., Denis, D., Dominique, G., Benoit, C., Sabaté, J.M., 2010. Anatomical connections between brain areas activated during rectal distension in healthy volunteers: a visceral pain network. Eur. J. Pain 14 (2), 142–148.

Monteleone, A.M., Castellini, G., Volpe, U., Ricca, V., Lelli, L., Monteleone, P., Maj, M., 2018. Neuroendocrinology and brain imaging of reward in eating disorders: a possible key to the treatment of anorexia nervosa and bulimia nervosa. Prog. Neuropsychopharmacol. Biol. Psychiatry 80, 132–142.

Moormann, P.P., Bermond, B., Vorst, H.C.M., Bloemendaal, A.F.T., Teijn, S.M., Rood, L., 2008. New avenues in alexithymia research: the creation of alexithymia types. In: Vingerhoets, Ad, Nyklicek, I., Denollet, J. (Eds.), Emotion Regulation. Springer, New York, NY, pp. 27–42.

Morie, K.P., Yip, S.W., Nich, C., Hunkele, K., Carroll, K.M., Potenza, M.N., 2016. Alexithymia and addiction: a review and preliminary data suggesting neurobiological links to reward / loss processing. Curr. Addict. Rep. 3 (2), 239–248.

Moriguchi, Y., Decety, J., Ohnishi, T., Maeda, M., Mori, T., Nemoto, K., et al., 2007a. Empathy and judging other's pain: an fMRI study of alexithymia. Cereb. Cortex 17 (9), 2223–2234.

Moriguchi, Y., Decety, J., Ohnishi, T., Maeda, M., Mori, T., Nemoto, K., et al., 2007b. Empathy and judging other's pain: an fMRI study of alexithymia. Cereb. Cortex 17 (9), 2223–2234.

Motomura, K., Terasawa, Y., Natsume, A., Iijima, K., Chalise, L., Sugiura, J., et al., 2019. Anterior insular cortex stimulation and its effects on emotion recognition. Brain Struct. Funct. 224 (6), 2167–2181.

Muir, K., Madill, A., Brown, C., 2017. Individual differences in emotional processing and autobiographical memory: interoceptive awareness and alexithymia in the fading affect bias. Cogn. Emot. 31 (7), 1392–1404.

Mul, C.-L., Stagg, S.D., Herbelin, B., Aspell, J.E., 2018. The feeling of me feeling for you: interoception, alexithymia and empathy in autism. J. Autism Dev. Disord. 48 (9), 2953–2967.

Mulcahy, J.S., Davies, M., Quadt, L., Critchley, H.D., Garfinkel, S.N., 2019. Interoceptive awareness mitigates deficits in emotional prosody recognition in Autism. Biol. Psychol. 146 (April), 107711.

Murphy, Jennifer, Brewer, R., Catmur, C., Bird, G., 2017. Interoception and psychopathology: a developmental neuroscience perspective. Dev. Cogn. Neurosci. 23, 45–56.

Murphy, Jennifer, Brewer, R., Hobson, H., Catmur, C., 2018. Is alexithymia characterised by impaired interoception? Further evidence, the importance of control variables, and the problems with the Heartbeat counting Task. Biol. Psychol. 136, 189–197.

Murphy, J., Catmur, C., Bird, G., 2018. Alexithymia is associated with a multidomain, multidimensional failure of interoception: Evidence from novel tests. J. Exp. Psychol. Gen. 147 (3), 398.

Murphy, J., Brewer, R., Hobson, H., Catmur, C., Bird, G., 2018. Is alexithymia characterised by impaired interoception? Further evidence, the importance of control variables, and the problems with the Heartbeat Counting Task. Biological psychology 136, 189–197.

- Murphy, Jennifer, Geary, H., Millgate, E., Catmur, C., Bird, G., 2017. Direct and indirect effects of age on interoceptive accuracy and awareness across the adult lifespan. Psychon. Bull. Rev.
- Murphy, Jennifer, Millgate, E., Geary, H., Ichijo, E., Coll, M.P., Brewer, R., et al., 2018. Knowledge of resting heart rate mediates the relationship between intelligence and the heartbeat counting task. Biol. Psychol. 133 (January), 1–3.
- Murphy, Jennifer, Wulff, K., Catmur, C., Bird, G., 2018. Alexithymic traits, independent of depression and anxiety, are associated with reduced sleep quality. Pers. Individ. Dif. 129, 175–178.
- Murphy, Jennifer, Brewer, R., Coll, M., Plans, D., Hall, M., Shiu, S.S., et al., 2019. I feel it in my finger: measurement device affects cardiac interoceptive accuracy. Biol. Psychol. 148, 107765.

Murphy, Jennifer, Catmur, C., Bird, G., 2019. Classifying individual differences in interoception: implications for the measurement of interoceptive awareness. Psychon. Bull. Rev. 26 (5), 1467–1471.

- Murphy, Jennifer, Cheesman, R., Gregory, A.M., Lau, J., Ehlers, A., Catmur, C., Eley, T. C., 2019. Estimating the stability of heartbeat counting in middle childhood: A twin study. Biol. Psychol. 148, 107764.
- Murphy, Jennifer, Millgate, E., Geary, H., Catmur, C., Bird, G., 2019. No effect of age on emotion recognition after accounting for cognitive factors and depression. Q. J. Exp. Psychol. 72 (11), 2690–2704.
- Murphy, Jennifer, Viding, E., Bird, G., 2019. Does atypical interoception following physical change contribute to sex differences in mental illness? Psychol. Rev. 126 (5), 787–789.
- Murphy, J., Brewer, R., Plans, D., Khalsa, S.S., Catmur, C., Bird, G., 2020. Testing the independence of self-reported interoceptive accuracy and attention. Q. J. Exp. Psychol. 73 (1), 115–133.
- Mussgay, L., Klinkenberg, N., Rüddel, H., 1999. Heart beat perception in patients with depressive, somatoform, and personality disorders. J. Psychophysiol. 13, 27–36.
- Nakano, T., Tanaka, K., Endo, Y., Yamane, Y., Yamamoto, T., Nakano, Y., et al., 2010. Atypical gaze patterns in children and adults with autism spectrum disorders dissociated from developmental changes in gaze behaviour. Proc. R. Soc. B: Biol. Sci. 277 (1696), 2935–2943.
- Nandrino, J.-L., Berna, G., Hot, P., Dodin, V., Latrée, J., Decharles, S., Sequeira, H., 2012. Cognitive and physiological dissociations in response to emotional pictures in patients with anorexia. J. Psychosom. Res. 72, 58–64.
- Naqvi, Nasir H., Bechara, A., 2009. The hidden island of addiction: the insula. Trends Neurosci. 32 (1), 56–67.
- Naqvi, Nasir H., Bechara, A., 2010. The insula and drug addiction: an interoceptive view of pleasure, urges and decision-making. Brain Struct. Funct. 214 (5–6), 435–450.
- Naqvi, N.H., Rudrauf, D., Damasio, H., Bechara, a., 2007. Damage to the insula disrupts addiction to cigarette smoking. Science 315 (5811), 531–534.
- Näring, G.W.B., van der Staak, C.P.F., 1995. Perception of heart rate and blood pressure: the role of alexithymia and anxiety. Psychother. Psychosom. 63, 193–200.
- Nemiah, J.C., Freyberger, H.J., Sifneos, P.E., 1976. Alexithymia: a view of the psychosomatic process. In: Hill, O.W. (Ed.), Modern Trends in Psychosomatic Medicine. Butterworths, London, England, pp. 430–439.

Nentjes, L., Meijer, E., Bernstein, D., Arntz, A., Medendorp, W., 2013. Brief communication: investigating the relationship between psychopathy and interoceptive awareness. J. Pers. Disord. 27 (5), 6229.

- Nestor, L.J., Ghahremani, D.G., Monterosso, J., London, E.D., 2011. Neuroimaging Prefrontal hypoactivation during cognitive control in early abstinent methamphetamine-dependent subjects. Psychiatry Res. Neuroimaging 194 (3), 287–295.
- Neumann, D., Zupan, B., Malec, J.F., Hammond, F., 2014. Relationships between alexithymia, affect recognition, and empathy after traumatic brain injury. J. Head Trauma Rehabil. 29 (1), E18–E27.
- Newton, T.L., Contrada, R.J., 1994. Alexithymia and repression: contrasting emotionfocused coping styles. Psychosom. Med. 56 (5), 457–462.
- Nicholson, T.M., Williams, D.M., Grainger, C., Christensen, J.F., Calvo-merino, B., Gaigg, S.B., 2018. Interoceptive impairments do not lie at the heart of autism or alexithymia. J. Abnorm. Psychol. 127 (6), 612–622.
- Nicholson, T., Williams, D., Carpenter, K., Kallitsounaki, A., 2019. Interoception is impaired in children, but not adults, with autism Spectrum disorder. J. Autism Dev. Disord. 49 (9), 3625–3637.
- Niedenthal, P.M., Barsalou, L.W., Winkielman, P., Krauth-gruber, S., Ric, F., 2005. Embodiment in attitudes, social perception, and emotion. Personal. Soc. Psychol. Rev. 9 (3), 184–211.
- Nummenmaa, L., Glerean, E., Hari, R., Hietanen, J.K., 2014. Bodily maps of emotions. Proc. Natl. Acad. Sci. U.S.A. 111 (2), 646–651.
- Nummenmaa, L., Hari, R., Hietanen, J.K., Glerean, E., 2018. Maps of subjective feelings. Proc. Natl. Acad. Sci. 115, 37.
- O'Brien, W.H., Reid, G.J., Jones, K.R., 1998. Differences in heartbeat awareness among males with higher and lower levels of systolic blood pressure. Int. J. Psychophysiol. 29 (1), 53–63.
- O'Driscoll, C., Laing, J., Mason, O., 2014. Cognitive emotion regulation strategies, alexithymia and dissociation in schizophrenia, a review and meta-analysis. Clin. Psychol. Rev. 34 (6), 482–495.
- Oakley, B.F.M., Brewer, R., Bird, G., Catmur, C., 2016. 'Theory of Mind' is not Theory of Emotion: a cautionary note on the reading the Mind in the Eyes test. J. Abnorm. Psychol. 125 (6), 818.

- Neuroscience and Biobehavioral Reviews 130 (2021) 470-508
- Olausson, H., Charron, J., Marchand, S., Villemure, C., Strigo, I.A., Bushnell, M.C., 2005. Feelings of warmth correlate with neural activity in right anterior insular cortex. Neurosci. Lett. 389, 1–5.
- Ondobaka, S., Kilner, J., Friston, K., 2017. The role of interoceptive inference in theory of mind. Brain Cogn. 112, 64–68.
- Oppenheimer, S.M., Gelb, A., Girvin, J.P., Hachinski, V.C., 1992. Cardiovascular effects of human insular cortex stimulation. Neurology 42 (9), 1727–1727.

Orem, J., Barnes, C.D., 2012. Physiology in Sleep. Academic Press.

- Ortigue, S., Grafton, S.T., Bianchi-Demicheli, F., 2007. Correlation between insula activation and self-reported quality of orgasm in women. NeuroImage 37, 551–560. Pace-schott, E.F., Amole, M.C., Aue, T., Balconi, M., Bylsma, L.M., Critchley, H., et al.,
- 2019. Physiological feelings. Neurosci. Biobehav. Rev. 103, 267–304. Palgi, S., Klein, E., Shamay-Tsoory, S., 2017. The role of oxytocin in empathy in PTSD.
- Psychol. Trauma Theor. Res. Pract. Policy 9 (1), 70–75.
- Palser, E.R., Fotopoulou, A., Pellicano, E., Kilner, J.M., 2018. The link between interoceptive processing and anxiety in children diagnosed with autism spectrum disorder: extending adult findings into a developmental sample. Biol. Psychol. 136 (May), 13–21.
- Palser, E.R., Fotopoulou, A., Pellicano, E., Kilner, J.M., 2019. Dissociation in how core autism features relate to interoceptive dimensions: evidence from cardiac awareness in children. J. Autism Dev. Disord. (0123456789).
- Panneton, W.M., 1991. Primary afferent projections from the upper respiratory tract in the muskrat. J. Comp. Neurol. 308 (1), 51–65.
- Papadimitriou, G.N., Linkowski, P., 2005. Sleep disturbance in anxiety disorders. Int. Rev. Psychiatry 17 (4), 229–236.
- Papciak, a S., Feuerstein, M., Spiegel, Ja., 1985. Stress reactivity in alexithymia: decoupling of physiological and cognitive responses. J. Human Stress 11 (March 2014), 135–142.
- Papezová, H., Yamamotová, A., Uher, R., 2005. Elevated pain threshold in eating disorders: physiological and psychological factors. J. Psychiatr. Res. 39 (4), 431–438.
- Paradiso, S., Vaidaya Jatin, G., McCormick, L.M., Jones, A., Robinson, R.G., 2008. Aging and alexithymia association with reduced right rostral cingulate volume. Am. J. Geriatr. Psychiatry 16 (9), 760–769.
- Park, H.D., Blanke, O., 2019. Coupling inner and outer body for self-consciousness. Trends Cogn. Sci. 23 (5), 377–388.
- Park, H.D., Tallon-Baudry, C., 2014. The neural subjective frame: from bodily signals to perceptual consciousness. Philos. Trans. Biol. Sci. 369 (1641).
- Parkin, L., Morgan, R., Rosselli, A., Howard, M., Sheppard, A., Evans, D., et al., 2014. Exploring the relationship between mindfulness and cardiac perception. Mindfulness 5 (3), 298–313.
- Pattinson, K.T.S., Governo, R.J., MacIntosh, B.J., Russell, E.C., Corfield, D.R., Tracey, I., Wise, R.G., 2009. Opioids depress cortical centers responsible for the volitional control of respiration. J. Neurosci. 29 (25), 8177–8186.
- Pauli, P., Hartl, L., Marquardt, C., Stalmann, H., Srian, F., 1991. Heartbeat and arrhythmia perception in diabetic autonomic neuropathy. Psychol. Med. 21, 413–421.
- Paulus, M.P., 2007. Neural basis of reward and craving a homeostatic point of view. Dialogues Clin. Neurosci. 9 (4), 379–387.
- Paulus, M.P., Stein, M.B., 2006. An insular view of anxiety. Biol. Psychiatry 60 (4), 383–387.
- Paulus, M.P., Stein, M.B., 2010. Interoception in anxiety and depression. Brain Struct. Funct. 214, 451–463.
- Paulus, M.P., Stewart, J.L., 2014. Interoception and drug addiction. Neuropharmacology 76 (0), 342–350.
- Paulus, M.P., Hozack, N., Frank, L., Brown, G.G., Schuckit, M.A., 2003. Decision making by methamphetamine-dependent subjects is associated with error-rate-independent decrease in prefrontal and parietal activation. Biol. Psychiatry 53, 65–74.
- Paulus, M.P., Rogalsky, C., Simmons, A., Feinstein, J.S., Stein, M.B., 2003. Increased activation in the right insula during risk-taking decision making is related to harm avoidance and neuroticism. NeuroImage 19 (4), 1439–1448.
- Paulus, M.P., Tapert, S.F., Schulteis, G., 2009. The role of interoception and alliesthesia in addiction. Pharmacol. Biochem. Behav. 94 (1), 1–7.
- Peled, A., Ritsner, M., Hirschmann, S., Geva, A.B., Modai, I., 2000. Touch feel illusion in schizophrenic patients. Soc. Biol. Psychiatry 48 (11), 1105–1108.
- Peled, A., Pressman, A., Geva, A.B., Modai, I., 2003. Somatosensory evoked potentials during a rubber-hand illusion in schizophrenia. Schizophr. Res. 64, 157–163.
- Pellicano, E., Burr, D., 2012. When the world becomes "too real": a Bayesian explanation of autistic perception. Trends Cogn. Sci. 16 (10), 504–510.
- Peñaloza Rojas, J.H., 1964. Electroencephalographic synchronization resulting from direct current application to the vagus nerves. Exp. Neurol. 9, 367–371.
- Penfield, W., Faulk, M.E., 1955. The insula: further observations on its function. Brain 78 (4), 445–470.
- Persons, J.B., 2008. The Case Formulation Approach to Cognitive-behavior Therapy. Guildford Press, New York.
- Petit, L., Beauchamp, M.S., 2003. Neural basis of visually guided head movements studied with fMRI. J. Neurophysiol. 89 (5), 2516–2527.
- Petzschner, F.H., Garfinkel, S.N., Paulus, M.P., Koch, C., Khalsa, S.S., 2021. Computational models of interoception and body regulation. Trends Neurosci. 44 (1), 63–76.
- Peyron, R., Peyron, R., Laurent, B., Laurent, B., García-Larrea, L., García-Larrea, L., 2000. Functional imaging of brain responses to pain. A review and meta-analysis (2000). Neurophysiol. Clin. Neurophysiol. 30 (5), 263–288.
- Pfeifer, G., Garfinkel, S.N., Gould van Praag, C.D., Sahota, K., Betka, S., Critchley, H.D., 2017. Feedback from the heart: emotional learning and memory is controlled by cardiac cycle, interoceptive accuracy and personality. Biol. Psychol. 126, 19–29.

Phillips, G.C., Jones, G.E., Rieger, E.J., Snell, J.B., 1999. Effects of the presentation of false heart-rate feedback on the performance of two common heartbeat-detection tasks. Psychophysiology 36, 504–510.

Phillips, M.L., Drevets, W.C., Rauch, S.L., Lane, R., 2003. Neurobiology of Emotion Perception I: The Neural Basis of Normal Emotion Perception. Biol. Psychiatry 54, 504–514.

Phillips, M.L., Gregory, L.J., Cullen, S., Cohen, S., Ng, V., Andrew, C., et al., 2003. The effect of negative emotional context on neural and behavioural responses to oesophageal stimulation. Brain 126 (3), 669–684.

Picardi, A., Toni, A., Caroppo, E., 2005. Stability of alexithymia and its relationships with the "big five" factors, temperament, character, and attachment style. Psychother. Psychosom. 74 (6), 371–378.

Piech, R.M., Strelchuk, D., Knights, J., Hjälmheden, J.V., Olofsson, J.K., Aspell, J.E., 2017. People with higher interoceptive sensitivity are more altruistic, but improving interoception does not increase altruism. Sci. Rep. 7 (1), 1–5.

Pierce, R.C., Crawford, C.A., Nonneman, A.J., Mattingly, B.A., Bardo, M.T., 1990. Effect of forebrain dopamine depletion on novelty-induced place preference behavior in rats. Pharmacol. Biochem. Behav. 36 (2), 321–325.

Pittenger, S.T., Bevins, R.A., 2013. Interoceptive conditioning with a nicotine stimulus is susceptible to reinforcer devaluation. Behav. Neurosci. 127 (3), 465–473.

Plailly, J., Radnovich, A.J., Sabri, M., Royet, J.P., Kareken, D.A., 2007. Involvement of the left anterior insula and frontopolar gyrus in odor discrimination. Hum. Brain Mapp. 28 (5), 363–372.

Plana, I., Lavoie, M.-A., Battaglia, M., Achim, A.M., 2014. A meta-analysis and scoping review of social cognition performance in social phobia, posttraumatic stress disorder and other anxiety disorders. J. Anxiety Disord. 28, 169–177.

Plans, D., Ponzo, S., Morelli, D., Cairo, M., Ring, C., Keating, C.T., et al., 2020. Measuring interoception: the phase adjustment task. PsyArXiv Preprints 44 (0), 1–23.

Poletti, M., Cavedini, P., Bonuccelli, U., Poletti, M., Cavedini, P., Bonuccelli, U., 2011. Iowa gambling task in parkinson's disease. J. Clin. Exp. Neuropsychol. 33 (4), 395–409.

Pollatos, O., Schandry, R., 2008. Emotional processing and emotional memory are modulated by interoceptive awareness. Cogn. Emot. 22 (2), 272–287.

Pollatos, O., Gramann, K., Schandry, R., 2007. Neural systems connecting interoceptive awareness and feelings. Hum. Brain Mapp. 28 (1), 9–18.

Pollatos, O., Herbert, B.M., Matthias, E., Schandry, R., 2007. Heart rate response after emotional presentation is modulated by interoceptive awareness. Int. J. Psychophysiol. 63, 117–124.

Pollatos, O., Traut-Mattausch, E., Schroeder, H., Schandry, R., 2007. Interoceptive awareness mediates the relationship between anxiety and the intensity of unpleasant feelings. J. Anxiety Disord. 21, 931–943.

Pollatos, O., Kurz, A.L., Albrecht, J., Schreder, T., Kleemann, A.M., Schöpf, V., et al., 2008. Reduced perception of bodily signals in anorexia nervosa. Eat. Behav. 9 (4), 381–388.

Pollatos, O., Traut-Mattausch, E., Schandry, R., 2009. Differential effects of anxiety and depression on interoceptive accuracy. Depress. Anxiety 26 (2), 167–173.

Pollatos, O., Herbert, B.M., Wankner, S., Dietel, A., Wachsmuth, C., Henningsen, P., Sack, M., 2011. Autonomic imbalance is associated with reduced facial recognition in somatoform disorders. J. Psychosom. Res. 71 (4), 232–239.

Pollatos, O., Werner, N.S., Duschek, S., Schandry, R., Matthias, E., Traut-Mattausch, E., Herbert, B.M., 2011. Differential effects of alexithymia subscales on autonomic reactivity and anxiety during social stress. J. Psychosom. Res. 70 (6), 525–533.

Pollatos, O., Füstös, J., Critchley, H.D., 2012. On the generalised embodiment of pain: how interoceptive sensitivity modulates cutaneous pain perception. Pain 153 (8), 1680–1686.

Pollatos, O., Dietel, A., Gündel, H., Duschek, S., 2015. Alexithymic trait, painful heat stimulation, and everyday pain experience. Front. Psychiatry 6 (OCT), 1–9.

Pollatos, O., Matthias, E., Keller, J., 2015. When interoception helps to overcome negative feelings caused by social exclusion. Front. Psychol. 6, 786.Pollatos, O., Herbert, B.M., Mai, S., Kammer, T., 2016. Changes in interoceptive

processes following brain stimulation. Philos. Trans. R. Soc. Lond., B, Biol. Sci. Ponzo, S., Kirsch, L.P., Fotopoulou, A., Jenkinson, P.M., 2018. Balancing body

ownership: visual capture of proprioception and affectivity during vestibular stimulation. Neuropsychologia 117, 311–321.

Ponzo, S., Kirsch, L.P., Fotopoulou, A., Jenkinson, P.M., 2019. Vestibular modulation of multisensory integration during actual and vicarious tactile stimulation. Psychophysiology 56 (10), 1–12.

Porges, S.W., 1993. Body Perception Questionnaire. University of Maryland: Laboratory of Developmental Assessment.

Porter, R.J., Gallagher, P., Thompson, J.M., Young, A.H., 2003. Neurocognitive impairment in drug-free patients with major depressive disorder. Br. J. Psychiatry 182 (March), 214–220.

Preston, S.D., de Waal, F.B.M., 2001. Empathy: its ultimate and proximate bases. Behav. Brain Sci. 25 (01).

Preuschoff, K., Quartz, S.R., Bossaerts, P., 2008. Human insula activation reflects risk prediction errors as well As risk. J. Neurosci. 28 (11), 2745–2752.

Pritchard, T.C., Macaluso, D.A., Eslinger, P.J., 1999. Taste perception in patients with insular cortex lesions. Behav. Neurosci. 113 (4), 663–671.

Prochnow, D., Donell, J., Schäfer, R., Jörgens, S., Hartung, H.P., Franz, M., Seitz, R.J., 2011. Alexithymia and impaired facial affect recognition in multiple sclerosis. J. Neurol. 258 (9), 1683–1688.

Pury, C.L.S., McCubbin, J.A., Helfer, S.G., Golloway, C., McMullen, L.J., 2004. Elevated resting blood pressure and dampened emotional response. Psychosom. Med. 66, 583–587.

Quadt, L., Critchley, H.D., Garfinkel, S.N., 2018. The neurobiology of interoception in health and disease. Ann. N. Y. Acad. Sci. 1428, 112–128. Neuroscience and Biobehavioral Reviews 130 (2021) 470-508

Quattrocki, E., Friston, K., 2014. Autism, oxytocin and interoception. Neurosci. Biobehav. Rev. 47, 410–430.

Quigley, K.S., Kanoski, S., Grill, W.M., Barrett, L.F., Tsakiris, M., 2021. Functions of interoception: from energy regulation to experience of the self. Trends Neurosci. 44 (1), 29–38.

Rae, C.L., Larsson, D.E.O., Garfinkel, S.N., Critchley, H.D., 2019. Dimensions of interoception predict premonitory urges and tic severity in Tourette syndrome. Psyciatry Research 271, 469–475.

Ray, R.D., Wilhelm, F.H., Gross, J.J., 2008. All in the mind's eye? Anger rumination and reappraisal. J. Pers. Soc. Psychol. 94 (1), 133–145.

Rebollo, I., Devauchelle, A.-D., Béranger, B., Tallon-Baudry, C., 2018. Stomach-brain synchrony reveals a novel, delayed-connectivity resting-state network in humans. ELife 7, e33321.

Reid, M.A., Stoeckel, L.E., White, D.M., Avsar, K.B., Bolding, M.S., Akella, N.S., et al., 2010. Assessments of function and biochemistry of the anterior cingulate cortex in schizophrenia. Biol. Psychiatry 68 (7), 625–633.

Reker, M., Ohrmann, P., Rauch, A.V., Kugel, H., Bauer, J., Dannlowski, U., et al., 2010. Individual differences in alexithymia and brain response to masked emotion faces. Cortex 46 (5), 658–667.

Repetti, R.L., Taylor, S.E., Seeman, T.E., 2002. Risky families: family social environments and the mental and physical health of offspring. Psychol. Bull. 128 (2), 330–366.

Riby, D.M., Hancock, P.J.B., 2008. Viewing it differently: social scene perception in Williams syndrome and autism. Neuropsychologia 46 (11), 2855–2860.

Ricciardi, L., Ferrazzano, G., Demartini, B., Morgante, F., Erro, R., Ganos, C., et al., 2016. Know thyself: exploring interoceptive sensitivity in Parkinson's disease. J. Neurol. Sci. 364, 110–115.

Rieffe, C., Oosterveld, P., Terwogt, M.M., 2006. An alexithymia questionnaire for children: factorial and concurrent validation results. Pers. Individ. Dif. 40 (1), 123–133.

Rietveld, S., Karsdorp, P.A., Mulder, B.J.M., 2004. Heartbeat sensitivity in adults with congenital heart disease. Int. J. Behav. Med. 11 (4), 203–211.

Ring, Christopher, Brener, J., 1992. The temporal locations of heartbeat sensations. Psychophysiology 29 (5), 535–545.

Ring, C., Brener, J., 1996. Influence of beliefs about heart rate and actual heart rate on heartbeat counting. Psychophysiology.

Ring, Christopher, Brener, J., 2018. Heartbeat counting is unrelated to heartbeat detection: a comparison of methods to quantify interoception. Psychophysiology 55, e13084.

Ring, Christopher, Brener, J., Knapp, K., Mailloux, J., 2015. Effects of heartbeat feedback on beliefs about heart rate and heartbeat counting: a cautionary tale about interoceptive awareness. Biol. Psychol. 104, 193–198.

Riquelme, I., Hatem, S.M., Montoya, P., 2016. Abnormal pressure pain, touch sensitivity, proprioception, and manual dexterity in children with Autism Spectrum disorders. Neural Plast. 2016, 1723401.

Romera, I., Perez, V., Menchón, J.M., Delgado-cohen, H., Polavieja, P., Gilaberte, I., 2010. Social and occupational functioning impairment in patients in partial versus complete remission of a major depressive disorder episode. A six-month prospective epidemiological study. Eur. Psychiatry 25, 58–65.

Rose, J.E., Zinser, M.D., Tashkin, D.P., Newcomb, R., Ertle, A., 1984. Subjective response to cigarette smoking following air-way anesthetization. Addict. Behav. 9, 211–215.

Rose, J.E., Tashkin, D.P., Ertle, A., Zinser, M.C., Lafer, R., 1985. Sensory blockade of smoking satisfaction. Pharmacol. Biochem. Behav. 23 (2), 289–293.

Rouse, C.H., Jones, G.E., Jones, K.R., 1988. The effect of body composition and gender on cardiac awareness. Psychophysiology 25 (4), 400–407.

Rozenstein, M.H., Latzer, Y., Stein, D., Eviatar, Z., 2011. Perception of emotion and bilateral advantage in women with eating disorders, their healthy sisters, and nonrelated healthy controls. J. Affect. Disord. 134 (1–3), 386–395.

Rufer, M., Hand, I., Braatz, A., Alsleben, H., Fricke, S., Peter, H., 2004. A prospective study of alexithymia in obsessive-compulsive patients treated with multimodal cognitive-behavioral therapy. Psychother. Psychosom. 73 (2), 101–106.

Rufer, Michael, Albrecht, R., Zaum, J., Schnyder, U., Mueller-Pfeiffer, C., Hand, I., Schmidt, O., 2010. Impact of alexithymia on treatment outcome: a naturalistic study of short-term cognitive-behavioral group therapy for panic disorder. Psychopathology 43 (3), 170–179.

Ruffman, T., Henry, J.D., Livingstone, V., Phillips, L.H., 2008. A meta-analytic review of emotion recognition and aging: implications for neuropsychological models of aging. Neurosci. Biobehav. Rev. 32 (4), 863–881.

Saarijärvi, S., Salminen, J.K., Toikka, T., 2006. Temporal stability of alexithymia over a five-year period in outpatients with major depression. Psychother. Psychosom. 75 (2), 107–112.

Säkkinen, P., Kaltiala-Heino, R., Ranta, K., Haataja, R., Joukamaa, M., 2007. Psychometric properties of the 20-item Toronto alexithymia scale and prevalence of alexithymia in a Finnish adolescent population. Psychosomatics 48 (2), 154–161.

Saladin, M.E., Santa Ana, E.J., Larowe, S.D., Simpson, A.N., Tolliver, B.K., Price, K.L., et al., 2012. Does alexithymia explain variation in cue-elicited craving reported by methamphetamine-dependent individuals? Am. J. Addiction 21 (2), 130–135.

Salminen, J.K., Saarijärvi, S., Toikka, T., Kauhanen, J., Äärelä, E., 2006. Alexithymia behaves as a personality trait over a 5-year period in Finnish general population. J. Psychosom. Res. 61 (2), 275–278.

Salvato, G., Mercurio, M., Sberna, M., Paulesu, E., Bottini, G., 2018. A very light lunch: interoceptive deficits and food aversion at onset in a case of behavioral variant frontotemporal dementia. Alzheimer's & Dementia: Diagnosis, Assess. Dis. Monitoring 10, 750–754.

Salvato, G., De Maio, G., Bottini, G., 2019. Interoceptive sensibility tunes risk-taking behaviour when body-related stimuli come into play. Sci. Rep. 9, 2396.

Sandsten, K.E., Nordgaard, J., Kjaer, T.W., Gallese, V., Ardizzi, M., Ferroni, F., et al., 2020. Altered self-recognition in patients with schizophrenia. Schizophr. Res. 218, 116–123.

- Saper, C.B., 2002. The central autonomic nervous system: conscious visceral perception and autonomic pattern generation. Annu. Rev. Neurosci. 25, 433–469.
- Sayed, S., Iacoviello, B.M., Charney, D.S., 2015. Risk factors for the development of psychopathology following trauma. Curr. Psychiatry Rep. 17 (8), 70.
- Sayin, A., Derinöz, O., Bodur, Ş., Şenol, S., Şener, Ş., 2007. Psychiatric symptoms and alexithymia in children and adolescents with non-organic pain: a controlled study. Gazi Med. J. 18 (4), 170–176.
- Scarpazza, C., Sellitto, M., Pellegrino, G., 2017. Now or not-now? The influence of alexithymia on intertemporal decision-making. Brain Cogn. 114, 20–28.
- Schaan, L., Schulz, A., Nuraydin, S., Bergert, C., Hilger, A., Rach, H., Hechler, T., 2019a. Interoceptive accuracy, emotion recognition, and emotion regulation in preschool children. Int. J. Psychophysiol. 138, 47–56.

Schaan, V.K., Schulz, A., Rubel, J.A., Bernstein, M., Domes, G., Schächinger, H., Vögele, C., 2019b. Childhood trauma affects stress-related interoceptive accuracy. Front. Psychiatry 10 (October), 1–10.

- Schachter, S., Singer, J.E., 1962. Cognitive, social, and psychological determinants of emotional state. Psychol. Rev. 69 (5), 379.
- Schaefer, M., Egloff, B., Witthöft, M., 2012. Is interoceptive awareness really altered in somatoform disorders? Testing competing theories with two paradigms of heartbeat perception. J. Abnorm. Psychol. 121 (3), 719–724.
- Schaefer, M., Egloff, B., Gerlach, A.L., Witthöft, M., 2014. Improving heartbeat perception in patients with medically unexplained symptoms reduces symptom distress. Biol. Psychol. 101 (1), 69–76.

Schandry, Rainer, 1981. Heart beat perception and emotional experience. Psychophysiology 18 (4), 483–488.

- Schandry, R., Weitkunat, R., 1990. Enhancement of heartbeat-related brain potentials through cardiac awareness training. Int. J. Neurosci. 53 (2–4), 243–253.
- Schandry, Rainer, Sparrer, B., Weitkunat, R., 1986. From the heart to the brain: a study of heartbeat contingent scalp potentials. Int. J. Neurosci. 30, 261–275.

Schauder, K.B., Mash, L.E., Bryant, L.K., Cascio, C.J., 2014. Interoceptive ability and body awareness in autism spectrum disorder. J. Exp. Child Psychol. 131, 193–200.

Schecklmann, M., Schwenck, C., Taurines, R., Freitag, C., Warnke, A., Gerlack, M., Romanos, M., 2013. A systematic review on olfaction in child and adolescent psychiatric disorders. J. Neural Transm. 120, 121–130.

- Schmidt, A.F., Eulenbruch, T., Langer, C., Banger, M., 2013. Interoceptive awareness, tension reduction expectancies and self-reported drinking behavior. Alcohol Alcohol. 48 (4), 472–477.
- Schreck, K.A., Williams, K., Smith, A.F., 2004. A comparison of eating behaviors between children with and without autism. J. Autism Dev. Disord. 34 (4), 433–438.

Schreiter, S., Pijnenborg, G.H.M., aan het Rot, M., 2013. Empathy in adults with clinical or subclinical depressive symptoms. J. Affect. Disord. 150, 1–16.

Schultchen, D., Zaudig, M., Krauseneck, T., Pollatos, O., 2019. Interoceptive deficits in patients with obsessive-compulsive disorder in the time course of cognitivebehavioral therapy. PLoS One.

Schulz, S.M., 2016. Neural correlates of heart-focused interoception: a functional magnetic resonance imaging meta-analysis. Philos. Trans. Biol. Sci. 371 (1708), 20160018.

- Schulz, André, Lass-Hennemann, J., Sütterlin, S., Schächinger, H., Vögele, C., 2013. Cold pressor stress induces opposite effects on cardioceptive accuracy dependent on assessment paradigm. Biol. Psychol. 93 (1), 167–174.
- Schulz, A., Köster, S., Beutel, M.E., Schächinger, H., Vögele, C., Rost, S., Michal, M., 2015. Altered patterns of heartbeat-evoked potentials in depersonalization/ derealization disorder: neurophysiological evidence for impaired cortical representation of bodily signals. Psychosom. Med. 77 (5), 506–516.
- Schulz, A., Stammet, P., Dierolf, A.M., Vögele, C., Beyenburg, S., Werer, C., Devaux, Y., 2018. Late heartbeat-evoked potentials are associated with survival after cardiac arrest. Resuscitation 126, 7–13.

Schwartz, G.E., Weinberger, D.A., Singer, J.A., 1981. Cardiovascular differentiation of happiness, sadness, anger, and fear following imagery and exercise. Psychosom. Med. 43 (4), 343–364.

Schwartz, E.K., Docherty, N.M., Najolia, G.M., Cohen, A.S., 2019. Exploring the racial diagnostic bias of schizophrenia using behavioral and clinical-based measures. J. Abnorm. Psychol. 128 (3), 263–271.

Sedeño, L., Couto, B., Melloni, M., Canales-Johnson, A., Yoris, A., Baez, S., et al., 2014. How do you feel when you can't feel your body? Interoception, functional connectivity and emotional processing in depersonalization-derealization disorder. PLoS One 9 (6).

Selzam, S., Coleman, J.R.I., Caspi, A., Moffitt, T.E., Plomin, R., 2018. A polygenic p factor for major psychiatric disorders. Transl. Psychiatry 8, 205.

Seseke, S., Baudewig, J., Kallenberg, K., Ringert, R.H., Seseke, F., Dechent, P., 2006. Voluntary pelvic floor muscle control — an fMRI study. NeuroImage 31, 1399–1407.

Seth, A.K., 2013. Interoceptive inference, emotion, and the embodied self. Trends Cogn. Sci. (Regul. Ed.) 17 (11), 565–573.

Seth, A.K., Suzuki, K., Critchley, H.D., 2012. An interoceptive predictive coding model of conscious presence. Front. Psychol. 2, 395.

Sforza, A., Bufalari, I., Haggard, P., Aglioti, S.M., 2010. My face in yours: visuo-tactile facial stimulation influences sense of identity. Soc. Neurosci. 5 (2), 148–162.

Shah, P., Catmur, C., Bird, G., 2016. Emotional decision-making in autism spectrum disorder: the roles of interoception and alexithymia. Mol. Autism 7 (1), 43.

Shah, P., Hall, R., Catmur, C., Bird, G., 2016. Alexithymia, not autism, is associated with impaired interoception. Cortex 81, 215–220.

Shah, P., Catmur, C., Bird, G., 2017. From heart to mind: linking interoception, emotion, and theory of mind. Cortex 93, 220.

- Shanahan, L., Copeland, W.E., Angold, A., Bondy, C.L., Costello, E.J., 2014. Sleep problems predict and are predicted by generalised anxiety/depression and oppositional defiant disorder. J. Am. Acad. Child Adolesc. Psychiatry 53 (5), 550–558.
- Shao, R., Read, J., Behrens, T.E.J., Rogers, R.D., 2013. Shifts in reinforcement signalling while playing slot-machines as a function of prior experience and impulsivity. Transl. Psychiatry 3 (1), e213–9.

Sharp, W.G., Berry, R.C., Mccracken, C., Nuhu, N.N., Marvel, E., Saulnier, C.A., et al., 2013. Feeding problems and nutrient intake in children with autism Spectrum disorders: a meta-analysis and comprehensive review of the literature. J. Autism Dev. Disord. 43, 2159–2173.

Shepherd, A.M., Matheson, S.L., Laurens, K.R., Carr, V.J., Green, M.J., 2012. Systematic meta-analysis of insula volume in schizophrenia. Biol. Psychiatry 72 (9), 775–784.

Shields, S., Mallory, M., Simon, A., 1989. Body awareness questionnaire: reliability and validity. J. Pers. Assess. 53 (4), 802–815.

Sifneos, P.E., 1973. The prevalence of "alexithymic" characteristics in psychosomatic patients. Psychother. Psychosom. 22 (2), 255–262.

- Silani, G., Bird, G., Brindley, R., Singer, T., Frith, C., Frith, U., 2008. Levels of emotional awareness and autism: an fMRI study. Soc. Neurosci. 3 (2), 97–112.
- Silver, A.J., 1990. Aging and risks for dehydration. Cleveland Clin. J. Med. 57 (4), 341-344.

Simmons, W.K., Avery, Ja., Barcalow, J.C., Bodurka, J., Drevets, W.C., Bellgowan, P., 2013. Keeping the body in mind: insula functional organization and functional connectivity integrate interoceptive, exteroceptive, and emotional awareness. Hum. Brain Mapp. 34 (11), 2944–2958.

Simms, M.L., Kemper, T.L., Timbie, C.M., Bauman, M.L., Blatt, G.J., 2009. The anterior cingulate cortex in autism: heterogeneity of qualitative and quantitative cytoarchitectonic features suggests possible subgroups. Acta Neuropathol. 118, 673–684.

- Sloan, D.M., Sandt, A.R., 2010. Depressed mood and emotional responding. Biol. Psychol. 84 (2), 368–374.
- Small, D.M., 2010. Taste representation in the human insula. Brain Struct. Funct. 214, 551–561.
- Smith, D.G., Xiao, L., Bechara, A., Smith, D.G., Xiao, L., Bechara, A., 2011. Decision making in children and adolescents: impaired iowa gambling task performance in early adolescence. Dev. Psychol. 48 (4), 1180.

Sokol-Hessner, P., Hsu, M., Curley, N.G., Delgado, M.R., Camerer, C.F., Phelps, E.A., 2009. Thinking like a trader selectively reduces individuals' loss aversion. Proc. Natl. Acad. Sci. 106 (13).

Sokol-Hessner, P., Hartley, Ca., Hamilton, J.R., Phelps, Ea., 2014. Interoceptive ability predicts aversion to losses. Cogn. Emot. 29 (4), 695–701.

South, S.C., Krueger, R.F., 2011. Genetic and environmental influences on internalizing psychopathology vary as a function of economic status. Psychol. Med. 41, 107–117.

Sowden, S., Koehne, S., Catmur, C., Dziobek, I., 2015. Intact automatic imitation and typical spatial compatibility in autism spectrum disorder: challenging the broken mirror theory. Autism Res.

Sowden, S., Brewer, R., Catmur, C., Bird, G., 2016. The specificity of the link between alexithymia, interoception and imitation. J. Exp. Psychol. Hum. Percept. Perform. 42 (11), 1687–1692.

- Sparrow, E.P., Spaniol, J., 2016. Age-related changes in decision making. Curr. Behav. Neurosci. Rep. 3, 285–292.
- Spek, A.A., van Rijnsoever, W., van Laarhoven, L., Kiep, M., 2019. Eating problems in men and women with an autism Spectrum disorder. J. Autism Dev. Disord. 23.
- Speranza, M., Corcos, M., Ste, P., Halfon, O., Jeammet, P., 2004. Alexithymia, depressive experiences and dependency in addictive disorders. Subst. Use Misuse 39 (4), 567–595.
- Sridharan, D., Levitin, D.J., Menon, V., 2008. A critical role for the right fronto-insular cortex in switching between central-executive and default-mode networks. Proc. Natl. Acad. Sci. U.S.A. 105 (34), 12569–12574.

Starita, F., Làdavas, E., di Pellegrino, G., 2016. Reduced anticipation of negative emotional events in alexithymia. Sci. Rep. 6, 27664.

Stasiewicz, P.R., Bradizza, C.M., Gudleski, G.D., Coffey, S.F., Schlauch, R.C., Bailey, S.T., et al., 2012. The relationship of alexithymia to emotional dysregulation within an alcohol dependent treatment sample. Addict. Behav. 37 (4), 469–476.

Steinberg, L., 2007. Risk taking in adolescence new perspectives from brain and behavioral science. Curr. Dir. Psychol. Sci. 16 (2), 55–59.

Stephan, E., Pardo, J.V., Faris, P.L., Hartman, B.K., Kim, S.W., Ivanov, E.H., et al., 2003. Functional neuroimaging of gastric distention. J. Gastrointest. Surg. 7 (6), 740–749.

Stephani, C., Fernandez-Baca Vaca, G., MacIunas, R., Koubeissi, M., Lüders, H.O., 2011. Functional neuroanatomy of the insular lobe. Brain Struct. Funct. 216 (2), 137–149.

Steptoe, A., Noll, A., 1997. The perception of bodily sensations, with special reference to hypochondriasis. Behav. Res. Ther. 35 (10), 901–910.

Steptoe, A., Vögele, C., 1992. Individual differences in the perception of bodily sensations: the role of trait anxiety and coping style. Behav. Res. Ther. 30 (6), 597–607.

Stern, E.R., 2014. Neural circuitry of interoception: new insights into anxiety and obsessive-compulsive disorders. Curr. Treat. Options Psychiatry 1 (September 2014), 235–247.

Stevens, J.C., Cruz, L.A., Hoffman, J.M., Patterson, M.Q., 1995. Taste sensitivity and aging: High incidence of decline revealed by repeated threshold measures. Chem. Senses 20 (4), 451–459.

Stevens, S., Gerlach, A.L., Cludius, B., Silkens, A., Craske, M.G., Hermann, C., 2011. Heartbeat perception in social anxiety before and during speech anticipation. Behav. Res. Ther. 49 (2), 138–143. Stingl, M., Bausch, S., Walter, B., Kagerer, S., Leichsenring, F., Leweke, F., 2008. Effects of inpatient psychotherapy on the stability of alexithymia characteristics. J. Psychosom. Res. 65 (2), 173–180.

- Stone, L.A., Nielson, K.A., 2001. Intact physiological response to arousal with impaired emotional recognition in alexithymia. Psychother. Psychosom. 70 (2), 92–102.
- Stubbs, B., Thompson, T., Acaster, S., Vancampfort, D., Gaughran, F., Correll, C.U., 2015. Decreased pain sensitivity among people with schizophrenia: a meta- analysis of experimental pain induction studies. Pain 156 (11), 2121–2131.
- Sturm, V.E., Levenson, R.W., 2011. Alexithymia in neurodegenerative disease. Neurocase 17 (3), 242–250.
- Sueyoshi, T., Sugimoto, F., Katayama, J., Fukushima, H., 2014. Neural correlates of error processing reflect individual differences in interoceptive sensitivity. Int. J. Psychophysiol.: Off. J. Int. Organization of Psychophysiol. 94 (3), 278–286.
- Suner-Soler, R., Grau, a., Gras, M.E., Font-Mayolas, S., Silva, Y., Davalos, a., et al., 2012. Smoking cessation 1 year poststroke and damage to the insular cortex. Stroke 43 (1), 131–136.
- Suslow, T., Junghanns, K., 2002. Impairments of emotion situation priming in alexithymia. Pers. Individ. Dif. 32 (3), 541–550.
- Sutherland, M.T., Carroll, A.J., Salmeron, B.J., Ross, T.J., Stein, Ea., 2013. Insula's functional connectivity with ventromedial prefrontal cortex mediates the impact of trait alexithymia on state tobacco craving. Psychopharmacology 228 (1), 143–155.
- Suzuki, K., Garfinkel, S.N., Critchley, H.D., Seth, A.K., 2013. Multisensory integration across exteroceptive and interoceptive domains modulates self-experience in the rubber-hand illusion. Neuropsychologia 51 (13), 2909–2917.
- Sze, J.A., Gyurak, A., Yuan, J.W., Levenson, R.W., 2010. Coherence between emotional experience and physiology: Does body awareness training have an impact? Emotion 10 (6), 803–814.
- Tajadura-Jiménez, A., Tsakiris, M., 2014. Balancing the "inner" and the "outer" self: interoceptive sensitivity modulates self-other boundaries. J. Exp. Psychol. Gen. 143 (2), 736–744.
- Tanidir, C., Mukaddes, N.M., Gu, B., 2007. Tactile sensitivity of normal and autistic children. Somatosens. Mot. Res. 24 (June), 21–33.
- Tavassoli, T., Frith, U., Blakemore, S., Tavassoli, T., Calò, S., Thomas, R.M., et al., 2006. Tactile sensitivity in Asperser syndrome Tactile sensitivity in Asperger syndrome. Brain Cogn. 61 (1), 5–13.
- Taylor, G.J., Parker, J.D.A., Michael Bagby, R., 1990. A preliminary investigation of alexithymia in men with psychoactive substance dependence. Am. J. Psychiatry 147 (9), 1228–1230.
- Taylor, K.S., Seminowicz, D.A., Davis, K.D., 2009. Two systems of resting state connectivity between the insula and cingulate cortex. Hum. Brain Mapp. 30 (9), 2731–2745.
- Teicher, M.H., Anderson, C.M., Ohashi, K., Polcari, A., 2014. Childhood maltreatment: altered network centrality of cingulate, precuneus, temporal pole and insula. Biol. Psychiatry 76 (4), 297–305.
- Terasawa, Y., Shibata, M., Moriguchi, Y., Umeda, S., 2012. Anterior insular cortex mediates bodily sensibility and social anxiety. Soc. Cogn. Affect. Neurosci. 8 (3), 259–266.
- Terasawa, Y., Moriguchi, Y., Tochizawa, S., Umeda, S., 2014. Interoceptive sensitivity predicts sensitivity to the emotions of others. Cogn. Emot. 28 (8), 1435–1448. Thorberg, F.A., Young, R.M., Sullivan, Ka., Lyvers, M., 2009. Alexithymia and alcohol
- Thorberg, F.A., Young, R.M., Sullivan, Ka., Lyvers, M., 2009. Alexithymia and alcohol use disorders: a critical review. Addict. Behav. 34 (3), 237–245.
- Todd, J., Aspell, J.E., Barron, D., Swami, V., 2019. An exploration of the associations between facets of interoceptive awareness and body image in adolescents. Body Image 31, 171–180.
- Tomchek, S.D., Dunn, W., 2007. Sensory processing in children with and without autism: a comparative study using the short sensory profile. Am. J. Occup. Ther. 61 (2), 190–200.
- Topsever, P., Filiz, T., Salman, S., Sengul, A., Sarac, E., Topalli, R., et al., 2006. Alexithymia in diabetes mellitus. Scott. Med. J. 51 (3), 15–20.
- Torregrossa, L.J., Snodgress, M.A., Hong, S.J., Nichols, H.S., Glerean, E., Nummenmaa, L., Park, S., 2019. Anomalous bodily maps of emotions in schizophrenia. Schizophr. Bull. 45 (5), 1060–1067.
- Tosic-Golubovic, S., Miljkovic, S., Nagorni, A., Lazarevic, D., Nikolic, G., 2010. Irritable bowel syndrome, anxiety, depression and personality characteristics. Psychiatr. Danub. 22 (3), 418–424.
- Trevisan, D.A., Bowering, M., Birmingham, E., 2016. Alexithymia, but not autism spectrum disorder, may be related to the production of emotional facial expressions. Mol. Autism 7, 46.
- Trevisan, Dominic A., Roberts, N., Lin, C., Birmingham, E., 2017. How do adults and teens with self-declared Autism Spectrum disorder experience eye contact? A qualitative analysis of first-hand accounts. PLoS One 12 (11), e0188446.
- Trevisan, Dominic A., Altschuler, M.R., Bagdasarov, A., Carlos, C., Duan, S., Hamo, E., et al., 2019. A meta-analysis on the relationship between interoceptive awareness and alexithymia: distinguishing interoceptive accuracy and sensibility. J. Abnorm. Psychol. 128 (8), 765–776.
- Trinkler, I., Devigneville, S., Achaibou, A., Ligneul, R.V., Brugières, P., de Langavant, L. C., et al., 2017. Embodied emotion impairment in Huntington's disease. Cortex 92, 44–56.
- Trojano, L., Fragassi, N.A., Postiglione, A., Grossi, D., 1988. Mixed transcortical aphasia. On relative sparing of phonological short-term store in a case. Neuropsychologia 26 (4), 633–638.
- Tsakiris, M., Critchley, H., 2016. Interoception beyond homeostasis: affect, cognition and mental health. Philos. Trans. Biol. Sci. 371, 20160002.
- Tsakiris, M., Tajadura-Jiménez, A., Costantini, M., 2011. Just a heartbeat away from one's body: interoceptive sensitivity predicts malleability of body-representations. Proc. R. Soc. B 278, 2470–2476.

- Tsuno, N., Besset, A., Ritchie, K., 2005. Sleep and depression. J. Clin. Psychiatry 66 (10), 1255–1270.
- Tuerke, K.J., Limebeer, C.L., Fletcher, P.J., Parker, L.A., 2012. Double dissociation between regulation of conditioned disgust and taste avoidance by serotonin availability at the 5-HT3 receptor in the posterior and anterior insular cortex. J. Neurosci. 32 (40), 13709–13717.
- Uddin, L.Q., Menon, V., 2009. The anterior insula in autism: under-connected and underexamined. Neurosci. Biobehav. Rev. 33 (8), 1198–1203.
- Uher, R., Murphy, T., Brammer, M.J., Dalgleish, T., Phillips, M.L., Ng, V.W., et al., 2004. Medial prefrontal cortex activity associated with symptom provocation in eating disorders. Am. J. Psychiatry 161, 1238–1246.
- Umeda, S., Tochizawa, S., Shibata, M., Terasawa, Y., 2016. Prospective memory mediated by interoceptive accuracy: a psychophysiological approach. Philos. Trans. Biol. Sci. 371, 20160005.
- Uzun, O., 2003. Alexithymia in male alcoholics: study in a Turkish sample. Compr. Psychiatry 44 (4), 349–352.
- Vaitl, D., 1996. Interoception. Biol. Psychiatry 42, 1-27.
- Vallbo, Å.B., Olausson, H., Wessberg, J., 1999. Unmyelinated afferents constitute a second system coding tactile stimuli of the human hairy skin. J. Neurophysiol. 81 (6), 2753–2763.
- van de Putte, E.M., Engelbert, R.H.H., Kuis, W., Kimpen, J.L.L., Uiterwaal, C.S.P.M., 2007. Alexithymia in adolescents with chronic fatigue syndrome. J. Psychosom. Res. 63, 377–380.
- van den Bergh, O., Winters, W., Devriese, S., van Diest, I., Vos, G., de Peuter, S., 2004. Accuracy of respiratory symptom perception in persons with high and low negative affectivity. Psychol. Health 19 (2), 213–222.
- Van Der Does, A.J.W., Antony, M.M., Ehlers, A., Barsky, A.J., 2000. Heartbeat perception in panic disorder: a reanalysis. Behav. Res. Ther. 38 (1), 47–62.
- Van den Bergh, O., Bogaerts, K., Walentynowicz, M., Van Diest, I., 2012. The interoceptive awareness questionnaire: unraveling the distinction between awareness of neutral and negative bodily sensations. In Annual Meeting of the International Society for the Advancement of Respiratory Psychophysiology (ISARP), 2012/09/28-2012/09/30, Orlando, FL
- van der Weiden, A., Prikken, M., van Haren, N.E.M., 2015. Self-other integration and distinction in schizophrenia: a theoretical analysis and a review of the evidence. Neurosci. Biobehav. Rev. 57, 220–237.
- Van Dyck, Z., Vögele, C., Blechert, J., Lutz, A.P.C., Schulz, A., Herbert, B.M., 2016. The water load test as a measure of gastric interoception: development of a two-stage protocol and application to a healthy female population. PLoS One 11 (9), 1–14.
- van Maanen, L., Fontanesi, L., Hawkins, G.E., Forstmann, B.U., 2016. Striatal activation reflects urgency in perceptual decision making. NeuroImage 139, 294–303.
- Vandenbergh, J., Dupont, P., Fischler, B., Bormans, G., Persoons, P., Janssens, J., Tack, J., 2005. Regional brain activation during proximal stomach distention in humans: a positron emission tomography study. Gastroenterology 128 (3), 564–573.
- Verdejo-Garcia, A., Clark, L., Dunn, B.D., 2012. The role of interoception in addiction: a critical review. Neurosci. Biobehav. Rev. 36 (8), 1857–1869.
- Villalta, L., Smith, P., Hickin, N., Stringaris, A., 2018. Emotion regulation difficulties in traumatized youth: a meta - analysis and conceptual review. Eur. Child Adolesc. Psychiatry 27, 527–544.
- Vizueta, N., Rudie, J.D., Townsend, J.D., Torrisi, S., Moody, T.D., Bookheimer, S.Y., Altshuler, L.L., 2012. Regional fMRI hypoactivation and altered functional connectivity during emotion processing in nonmedicated depressed patients with bipolar II disorder. Am. J. Psychiatry 169 (8), 831–840.
- von Leupoldt, A., Sommer, T., Kegat, S., Baumann, H.J., Klose, H., Dahma, B., Büchel, 2008. The unpleasantness of perceived dyspnea is processed in the anterior insula and amygdala. Am. J. Respir. Crit. Care Med. 177, 1026–1032.
- Vorst, H.C.M., Bermond, B., 2001. Validity and reliability of the bermond–Vorst alexithymia questionnaire. Pers. Individ. Dif. 30 (3), 413–434.
- Wadsworth, M.E., Achenbach, T.M., 2005. Explaining the link between low socioeconomic status and psychopathology: testing two mechanisms of the social causation hypothesis. J. Consult. Clin. Psychol. 73 (6), 1146–1153.
- Wakefield, A.J., Anthony, A., Murch, S.H., Thomson, M., Montgomery, S.M., Davies, S., et al., 2000. Enterocolitis in children with developmental disorders. Am. J. Gastroenterol. 95 (9), 2285–2295.
- Waldstein, S.R., Kauhanen, J., Neumann, S.A., Katzel, L.I., 2002. Alexithymia and cardiovascular risk in older adults: Psychosocial, psychophysiological, and biomedical correlates. Psychol. Health 17 (5), 597–610.
- Waltz, J.A., Frank, M.J., Robinson, B.M., Gold, J.M., 2007. Selective reinforcement learning deficits in schizophrenia support predictions from computational models of striatal-cortical dysfunction. Biol. Psychiatry 62 (7), 756–764.
- Wang, G.-J., Volkow, N.D., Fowler, J.S., Cervany, P., Hitxemann, R.J., Pappas, N.R., et al., 1999. Regional brain metabolic activation during craving elicited by recall of previous drug experiences. Life Sci. 64 (9), 775–784.
- Wang, L., Hermens, D.F., Hickie, I.B., Lagopoulos, J., 2012. A systematic review of
- resting-state functional-MRI studies in major depression. J. Affect. Disord. 142, 6–12. Watling, R.L., Deitz, J., White, O., 2001. Comparison of sensory profile scores of young children with and without autism spectrum disorders. Am. J. Occup. Ther. 55 (4), 416–423.
- Way, I., Yelsma, P., van Meter, A.M., Black-Pond, C., 2007. Understanding alexithymia and language skills in children: implications for assessment and intervention. Lang. Speech Hear. Serv. Sch. 38, 128–139.
- Wei, Y., Ramautar, J.R., Colombo, M.A., Stoffers, D., Gómez-Herrero, G., 2016. I keep a close watch on this heart of mine: increased interoception in insomnia. Sleep 39, 2113–2124.

Weiss, J.A., Thomson, K., Chan, L., 2014. A systematic literature review of emotion regulation measurement in individuals with autism spectrum disorder. Autism Res. 7 (6), 629–648.

Weiss, S., Sack, M., Henningsen, P., Pollatos, O., 2014. On the interaction of selfregulation, interoception and pain perception. Psychopathology 47, 377–382.

Weissman, D.G., Nook, E.C., Devs, A.A., Miller, A.B., Lambert, H.K., Sasse, S.F., et al., 2020. Low emotional awareness as a transdiagnostic mechanism underlying psychopathology in adolescence. Clin. Psychol. Sci.

- Weisz, J., Balazs, L., Adam, G., 1988. The influence of self-focused attention on heartbeat perception. Psychophysiology 25 (2), 193–199.
- Weller, J.A., Levin, I.P., Shiv, B., Bechara, A., Weller, J.A., Levin, I.P., et al., 2009. The effects of insula damage on decision-making for risky gains and losses. Soc. Neurosci. 4 (4), 347–358.

Weng, H.Y., Feldman, J.L., Leggio, L., Napadow, V., Park, J., Price, C.J., 2021.

Interventions and manipulations of interoception. Trends Neurosci. 44 (1), 52–62. Werner, N.S., Duschek, S., Mattern, M., Schandry, R., 2009. The relationship between pain perception and interoception. J. Psychophysiol. 23 (1), 35–42.

Werner, N.S., Jung, K., Duschek, S., Schandry, R., 2009. Enhanced cardiac perception is associated with benefits in decision-making. Psychophysiology 46 (6), 1123–1129.Werner, N.S., Peres, I., Duschek, S., Schandry, R., 2010. Implicit memory for emotional

- words is modulated by cardiac perception. Biol. Psychol. 85 (3), 370–376. Werner, N.S., Schweitzer, N., Meindl, T., Duschek, S., Kambeitz, J., Schandry, R., 2013. Interoceptive awareness moderates neural activity during decision-making. Biol.
- Psychol. 94, 498–506.
 Wessberg, J., Olausson, H., Fernström, K.W., Vallbo, A.B., 2003. Receptive field properties of unmyelinated tactile afferents in the human skin. J. Neurophysiol. 89, 1567–1575.
- Whitbourne, S.K., 2012. The Aging Body: Physiological Changes and Psychological Consequences. Media, Springer Science & Business.

White, J.F., 2003. Intestinal pathophysiology in autism. Exp. Biol. Med. 228 (6), 639-649.

Whitehead, W.E., Drescher, V.M., 1980. Perception of gastric contractions and selfcontrol of gastric motility. Psychophysiology 17 (6), 552–558.

Whitehead, William E., Drescher, V.M., Heiman, P., Blackwell, B., 1977. Relation of heart rate control to heartbeat perception. Biofeedback Self. 2 (4), 371–392.

Wicker, B., Keysers, C., Plailly, J., Royet, J.P., Gallese, V., Rizzolatti, G., 2003. Both of us disgusted in my insula: the common neural basis of seeing and feeling disgust. Neuron 40 (3), 655–664.

Wiebking, C., Northoff, G., 2015. Neural activity during interoceptive awareness and its associations with alexithymia — an fMRI study in major depressive disorder and non-psychiatric controls. Front. Psychol. 6, 589.

Wiens, S., 2005. Interoception in emotional experience. Curr. Opin. Neurol. 18 (4), 442–447.

Wiens, S., Mezzacappa, E.S., Katkin, E.S., 2000. Heartbeat detection and the experience of emotions. Cogn. Emot. 14 (3), 417–427.

Wigham, S., Rodgers, J., South, M., 2015. The interplay between sensory processing abnormalities, intolerance of uncertainty, anxiety and restricted and repetitive behaviours in Autism Spectrum disorder. J. Autism Dev. Disord. 45, 943–952.

Wildman, H.E., Jones, G.E., 1982. Consistency of heartbeat discrimination scores on the Whitehead procedure in knowledge of results trained and untrained subjects. Psychophysiology 19, 592.

Willem, C., Gandolphe, M.-C., Roussel, M., Verkindt, H., Pattou, F., 2019. Difficulties in emotion regulation and deficits in interoceptive awareness in moderate and severe obesity. Eat. Weight, Disord. - Stud. Anorex. Bulim. Obes. 24 (4), 633–644.

Williams, B.L., Hornig, M., Buie, T., Bauman, M.L., Cho Paik, M., Wick, I., et al., 2011. Impaired carbohydrate digestion and transport and mucosal dysbiosis in the intestines of children with autism and gastrointestinal disturbances. PLoS One 6 (9), e24585

Windmann, S., Schonecke, O.W., Fröhlig, G., Maldener, G., 1999. Dissociating beliefs about heart rates and actual heart rates in patients with cardiac pacemakers. Psychophysiology 36 (3), 339–342.

Wittkamp, M.F., Bertsch, K., Vögele, C., Schulz, A., 2018. A latent state-trait analysis of interoceptive accuracy. Psychophysiology 55 (e13055).

Wolf, F., Brüne, M., Assion, H.-J., 2010. Theory of mind and neurocognitive functioning in patients with bipolar disorder. Bipolar Disord. 12, 657–666.Wölk, J., Sütterlin, S., Koch, S., Vögele, C., Schulz, S.M., 2014. Enhanced cardiac

Wölk, J., Sütterlin, S., Koch, S., Vögele, C., Schulz, S.M., 2014. Enhanced cardiac perception predicts impaired performance in the Iowa Gambling Task in patients with panic disorder. Brain Behav. 4 (2), 238–246. Wolpert, N., Rebollo, I., Tallon-Baudry, C., 2020. Electrogastrography for psychophysiological research: practical considerations, analysis pipeline, and normative data in a large sample. Psychophysiology 57 (9), 1–25.

Wood, R.L.I., Williams, C., 2007. Neuropsychological correlates of organic alexithymia. J. Int. Neuropsychol. Soc. 13 (03), 471–479.

Woolf, C.J., Fitzgerald, M., 1983. The properties of neurones recorded in the superficial dorsal horn of the rat spinal cord. J. Comp. Neurol. 221 (3), 313–328.

- Wylie, K.P., Tregellas, J.R., 2010. The role of the insula in schizophrenia. Schizophr. Res. 123 (2–3), 93–104.
- Xiang, T., Lohrenz, T., Montague, P.R., 2013. Computational substrates of norms and their violations during social exchange. J. Neurosci. 33 (3), 1099–1108a.

Xue, G., Lu, Z., Levin, I.P., Bechara, A., 2010. The impact of prior risk experiences on subsequent risk decision-making: the role of the insula. NeuroImage 50 (2), 709–716.

Yamamotova, A., Bulant, J., Bocek, V., Papezova, H., 2017. Dissatisfaction with own body makes patients with eating disorders more sensitive to pain. J. Pain Res. 10, 1667–1675.

Yap, K., Dowling, N., Moulding, R., Mogan, C., Blair-west, S., Moriarty, A., Gelgec, C., 2018. Emotion regulation difficulties in obsessive-compulsive disorder. J. Clin. Psychol. 74, 695–709.

- Yehuda, R., Steiner, A., Kahan, B., Binder-Brynes, K., Southwick, S.M., Zemelman, S., Giller, E.L., 1997. Alexithymia in holocaust survivors with and without PTSD. J. Trauma. Stress 10 (1), 93–100.
- Yoris, Adrián, Esteves, S., Couto, B., Melloni, M., Kichic, R., Cetkovich, M., et al., 2015. The roles of interoceptive sensitivity and metacognitive interoception in panic. Behav. Brain Funct. 11 (14), 1–6.

Yoris, A., García, A.M., Traiber, L., Santamaría-García, H., Martorell, M., Alifano, F., et al., 2017. The inner world of overactive monitoring: neural markers of

interoception in obsessive-compulsive disorder. Psychol. Med. 47 (11), 1957–1970. Young, H.A., Williams, C., Pink, A.E., Freegard, G., Owens, A., Benton, D., 2017. Getting to the heart of the matter: Does aberrant interoceptive processing contribute towards emotional eating? PLoS One 12 (10), e0186312.

- Yu, X.H., Zhang, E.T., Craig, A.D., Shigemoto, R., Ribeiro-da-Silva, A., De Koninck, Y., 1999. NK-1 receptor immunoreactivity in distinct morphological types of lamina I neurons of the primate spinal cord. J. Neurosci. 19 (9), 3545–3555.
- Zachor, D.A., Ben-Itzchak, E., 2014. The relationship between clinical presentation and unusual sensory interests in autism Spectrum disorders: a preliminary investigation. J. Autism Dev. Disord. 44, 229–235.

Zaki, J., Davis, J.I., Ochsner, K.N., 2012. Overlapping activity in anterior insula during interoception and emotional experience. NeuroImage 62 (1), 493–499.

- Zaman, J., Weltens, N., Ly, H.G., Struyf, D., Vlaeyen, J.W.S., Van den Bergh, O., et al., 2016. Influence of interoceptive fear learning on visceral perception. Psychosom. Med. 78, 248–258.
- Zamariola, G., Vlemincx, E., Corneille, O., Luminet, O., 2018. Relationship between interoceptive accuracy, interoceptive sensibility, and alexithymia. Pers. Individ. Dif. 125, 14–20.

Zamariola, G., Frost, N., Van Oost, A., Corneille, O., Luminet, O., 2019. Relationship between interoception and emotion regulation: new evidence from mixed methods. J. Affect. Disord. 246, 480–485.

Zamariola, G., Luminet, O., Mierop, A., Corneille, O., 2019. Does it help to feel your body? Evidence is inconclusive that interoceptive accuracy and sensibility help cope with negative experiences accuracy and sensibility help cope with negative experiences. Cogn. Emot. 33 (8), 1627–1638.

Zampini, M., Guest, S., Shore, D.I., Spence, C., 2005. Audio-visual simultaneity judgments. Percept. Psychophys. 67 (3), 531-544.

Zhang, X., Salmeron, B.J., Ross, T.J., Geng, X., Yang, Y., Stein, E.A., 2011. Factors underlying prefrontal and insula structural alterations in smokers. Neuroimage 54 (1), 42–48.

Zhou, Y., Lin, F.-C., Du, Y.-S., Qin, L.-D., Zhao, Z.-M., Xu, J.-R., Lei, H., 2011. Gray matter abnormalities in Internet addiction: a voxel - based morphometry study. Eur. J. Radiol. 79 (November), 92–95.

Zimmermann, Ä., 2006. Delinquency in male adolescents: the role of alexithymia and family structure. J. Adolesc. 29, 321–332.

Zonnevijlle-Bender, M.J.S., van Goozen, S.H.M., Cohen-Kettenis, P.T., van Elburg, A., van Engeland, H., 2002. Do adolescent anorexia nervosa patients have deficits in emotional functioning? Eur. Child Adolesc. Psychiatry 11 (1), 38–42.

Zu Eulenburg, P., Baumgärtner, U., Treede, R.D., Dieterich, M., 2013. Interoceptive and multimodal functions of the operculo-insular cortex: tactile, nociceptive and vestibular representations. NeuroImage 83, 75–86.