

Increased CD1c⁺ mDC1 with mature phenotype regulated by TNF α -p38 MAPK in autoimmune ocular inflammatory disease

Chen, Ping; Denniston, Alastair; Hannes, Susan; Tucker, William; Wei, Lai; Liu, Baoying; Xiao, Tiaojiang; Hirani, Sima; Li, Zhiyu; Jawad, Shayma; Si, Han; Lee, Richard W.j.; Nida Sen, H.; Nussenblatt, Robert B.

DOI:

[10.1016/j.clim.2015.03.002](https://doi.org/10.1016/j.clim.2015.03.002)

License:

Other (please specify with Rights Statement)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Chen, P, Denniston, A, Hannes, S, Tucker, W, Wei, L, Liu, B, Xiao, T, Hirani, S, Li, Z, Jawad, S, Si, H, Lee, RWJ, Nida Sen, H & Nussenblatt, RB 2015, 'Increased CD1c⁺ mDC1 with mature phenotype regulated by TNF α -p38 MAPK in autoimmune ocular inflammatory disease', *Clinical Immunology*, vol. 158, no. 1, pp. 35-46. <https://doi.org/10.1016/j.clim.2015.03.002>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

NOTICE: this is the author's version of a work that was accepted for publication in *Clinical Immunology*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Clinical Immunology*, Vol 158, Issue 1, DOI: 10.1016/j.clim.2015.03.002.

Eligibility for repository checked April 2015

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Accepted Manuscript

Increased CD1c⁺ mDC1 with mature phenotype regulated by TNF α -p38 MAPK in autoimmune ocular inflammatory disease

Ping Chen, Alastair Denniston, Susan Hannes, William Tucker, Lai Wei, Baoying Liu, Tiaojiang Xiao, Sima Hirani, Zhiyu Li, Shayma Jawad, Han Si, Richard W.J. Lee, H. Nida Sen, Robert B. Nussenblatt

PII: S1521-6616(15)00092-3
DOI: doi: [10.1016/j.clim.2015.03.002](https://doi.org/10.1016/j.clim.2015.03.002)
Reference: YCLIM 7424

To appear in: *Clinical Immunology*

Received date: 8 August 2014
Revised date: 4 February 2015
Accepted date: 3 March 2015



Please cite this article as: Ping Chen, Alastair Denniston, Susan Hannes, William Tucker, Lai Wei, Baoying Liu, Tiaojiang Xiao, Sima Hirani, Zhiyu Li, Shayma Jawad, Han Si, Richard W.J. Lee, H. Nida Sen, Robert B. Nussenblatt, Increased CD1c⁺ mDC1 with mature phenotype regulated by TNF α -p38 MAPK in autoimmune ocular inflammatory disease, *Clinical Immunology* (2015), doi: [10.1016/j.clim.2015.03.002](https://doi.org/10.1016/j.clim.2015.03.002)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Increased CD1c⁺ mDC1 with mature phenotype regulated by TNF α -p38 MAPK in
autoimmune ocular inflammatory disease**

Ping Chen¹, Alastair Denniston^{2,3}, Susan Hannes¹, William Tucker¹, Lai Wei¹, Baoying Liu¹, Tiaojiang Xiao⁴, Sima Hirani¹, Zhiyu Li¹, Shayma Jawad¹, Han Si¹, Richard W.J. Lee⁵, H. Nida Sen¹, Robert B. Nussenblatt¹

¹Laboratory of Immunology, National Eye Institute, National Institutes of Health, Bethesda, Maryland 20892, USA

²Ophthalmology Department, Queen Elizabeth Hospital Birmingham, University Hospitals Birmingham NHSFT, Edgbaston, Birmingham B15 2WB, UK

³Centre for Translational Inflammation Research, University of Birmingham, UK

⁴Laboratory of Molecular Biology, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, MD 20892, USA.

⁵ Department of Clinical Sciences, University of Bristol, Bristol, UK

Running title: Mature CD1c⁺ mDC1 mediated by TNF α -p38 MAPK in autoimmune uveitis

Key words: CD1c⁺ mDC1, CD1c^{hi} mDC1 subpopulation, antigen uptake, TNF α , p38 MAPK, autoimmune uveitis

Corresponding author:

Robert B. Nussenblatt, M.D., M.P.H.

National Institutes of Health, National Eye Institute

9000 Rockville Pike, Building 10/10N109, Rockville, MD 20892

Email: nussenblatrr@nei.nih.gov

Phone: 301-435-5139

Abbreviation

mDC1, myeloid dendritic cells 1; HCs, healthy controls; MoDCs, monocyte-derived dendritic cells; PBMCs, peripheral blood mononuclear cells; Th, T helper; DCs, dendritic cells; mDCs, myeloid dendritic cells; pDCs, plasmacytoid dendritic cells; Lin1, lineage 1; Alb, albumin; iNKT, invariant natural killer T.

Abstract

In this study we investigated the role of blood CD1c⁺ myeloid dendritic cells 1 (mDC1), a key mDC subtype, in patients with autoimmune uveitis. We observed a significant increase of blood CD1c⁺ mDC1 in uveitis patients. The increased CD1c⁺ mDC1 exhibited high HLADR expression and less antigen uptake. CD1c⁺ mDC1 were divided into two subpopulations. CD1c^{hi} mDC1 subpopulation showed less antigen uptake and higher HLADR expression compared to CD1c^{lo} mDC1 subpopulation. Importantly, the CD1c^{hi} mDC1 subpopulation was increased in uveitis patients. *In vitro*, mature monocyte-derived dendritic cells (MoDCs), characterized by lower levels of antigen uptake, induced more CD62L⁻CD4⁺ T helper cell proliferation. The mature phenotype and function of CD1c⁺ mDC1 were regulated by TNF α via a p38 MAPK-dependent pathway. These data show that alterations in the systemic immune response are involved in the pathogenesis of autoimmune uveitis and invite the therapeutic possibility of attenuating uveitis by manipulating blood CD1c⁺ mDC1.

1. Introduction

Human autoimmune ocular inflammatory diseases, particularly non-infectious uveitis (abbreviated as uveitis), are currently responsible for 10% of the visual handicap in the United States [1]. The most commonly recognized mechanism of uveitis is CD4⁺ T helper (Th) cell mediated immune dysregulation [2-4]. It has been suggested that the immune dysregulation is characterized by the imbalance between regulatory and pathogenic effector T cells, increased inflammatory and decreased anti-inflammatory cytokines, hyper-responsiveness to self-antigens, and dysfunctional dendritic cells (DCs), which are all suspected to contribute to the pathogenesis of autoimmune disease [5].

Dendritic cells (DCs) play a central role in immune regulation since they are involved in both innate and adaptive immune responses. An indispensable part of initiating the immune response in the eye is the permeability of the blood-ocular barrier to inflammatory cells. Infiltrating cells include monocytes/macrophages, lymphocytes, neutrophils, and myeloid DCs (mDCs). Moreover, an emerging body of literature supports a pivotal role for mDCs in uveitis in humans and mice [6-8]. However, the actual source of mDCs in eye tissue, along with the factors that trigger DC maturation, take up antigens and help T cell responses, remain unidentified.

DCs are derived from either progenitors in the bone marrow or monocytes in the peripheral blood. Blood DCs contain three subsets: CD1c⁺ (BDCA1) myeloid dendritic cell 1 (mDC1), CD141⁺ (BDCA3) myeloid dendritic cell 2 (mDC2), and CD303⁺ (BDCA2) plasmacytoid DCs (pDCs) [9], which are from Lineage 1⁻ (Lin1, including CD3, CD14, CD16, CD19, CD20 and CD56) HLADR⁺ cells. Myeloid DCs actively infiltrate into the eye and can be seen suspending in the aqueous humor of uveitis patients [10].

In this study, we observed an increase in blood CD1c⁺ mDC1 with enhanced HLADR expression and lower levels of antigen uptake indicating a more mature phenotype. Moreover, this was associated with an increase in a subpopulation of CD1c^{hi} mDC1, which had very low levels of antigen uptake. In an *in vitro* model we observed that MoDCs with a similar low-antigen-uptake phenotype promoted more CD4⁺ T cell proliferation, than those taking up high levels of antigen. Our work also supports the hypothesis that in uveitis the mature phenotype of CD1c⁺ mDC1 were regulated by TNF α via a p38 MAPK-dependent pathway.

2. Material and methods

2.1. Study population

Peripheral blood was obtained from patients with non-infectious uveitis (n=74) in the Clinical Center of the National Eye Institute (NEI), and from healthy controls (HCs, n=96) in the Blood Bank of the National Institutes of Health (NIH). All protocols were approved by the NIH Neuroscience Institutional Review Board (IRB). Informed consents were obtained from all subjects before the study commenced. Ocular disease activity, i.e. active and inactive uveitis, were recorded according to the Standardization of Uveitis Nomenclature (SUN) [11]. According to SUN criteria, less than one cell, or alternatively, no flare in the anterior chamber, constitutes a graded score of 0. SUN criteria go on to address “trace” cells in the vitreous, correlating this clinical impression with a score of 0.5. A grade of 0 is regarded as inactive, whereas, a grade ≥ 0.5 is considered slightly active, and a grade ≥ 1 or haze in vitreous is considered active. Here, we put slightly active and active uveitis into one group as “active”.

To investigate the phenotype and function of mDC1 in the peripheral blood, the effects of systemic immunosuppressive therapy on mDC1 were considered. The presence or absence of systemic immunosuppressive therapy was thus documented; this included systemic corticosteroids, secondary immunomodulatory therapy and biologic response modifiers (see the table).

2.2. Identification of mDCs and pDCs in human peripheral blood

Blood DCs were identified by 4-color staining performed on whole fresh peripheral blood using the following monoclonal antibodies: PE-CD1c (BDCA1, Miltenyi Biotec, Auburn, CA) or PE-CD141 (BDCA3, Miltenyi Biotec), PERCP-HLADR (Miltenyi Biotec), APC-CD303 (BDCA2, Miltenyi Biotec), and FITC-labeled mAbs against lineage markers CD3, CD14, CD16, CD19, CD20 and CD56 (BD Biosciences, San Jose, CA). Cells that were not labeled with these lineage markers were designated as Lin¹⁻. Myeloid DC1 were CD1c⁺, mDC2 were CD141⁺, and pDCs were CD303⁺. These DCs were gated on Lin¹⁻ HLADR⁺ total DC population. Cells were analyzed on FACSCalibur (BD Biosciences). DC number was calculated by multiplying the percentage of DCs by the total number of white blood cells (WBC) in 1 ml blood (DC number = % of DCs × number of WBC/ml).

2.3. Generation of monocyte-derived DCs

Monocyte-derived dendritic cells (MoDCs) were generated as previously described and used as an *in vitro* model for mDCs for additional functional studies [12, 13]. Briefly, monocytes were isolated from mononuclear cell fractions of the peripheral blood of healthy controls and seeded in the presence of GM-CSF (40 ng/ml; R&D systems, Minneapolis, MN) and IL-4 (20 ng/ml; R&D Systems) at a concentration of 1×10^6

cells/well in a 6-well plate. Six to seven days after culture, monocytes were completely differentiated into dendritic cells and CD1c expression was evaluated by flow cytometry.

To investigate the effect of the TNF α -p38 MAPK signaling pathway on CD1c expression, recombinant TNF α (10 ng/ml, R&D, Minneapolis, MN) was added for 24 hours before measuring the protein expression of CD1c, and for 30 minutes before measuring phosphorylated-p38 MAPK (BD Biosciences) expression on MoDCs by flow cytometry. For blocking experiments, the selective MAPK inhibitor SB203580 (10 μ mol/l, Cell Signaling, Danvers, MA) was added at day 0 of MoDCs differentiation *in vitro*. CD1c protein expression was measured by flow cytometry on days 1, 3, and 5.

2.4. Evaluation of cytokine production

Peripheral blood were stimulated with lipopolysaccharides (LPS, 1 μ g/ml, Sigma-Aldrich, St. Louis, MO) and GolgiPlugTM (2 μ l/ml, BD Pharmingen, San Jose, California) for 5 h at 37°C in a humidified 5% CO₂ atmosphere. Following stimulation, cells were stained with Lin1, HLADR and CD1c to identify mDC1. Then, cells were fixed and permeabilized (Fix & Perm Cell Permeabilization Kit; BD Biosciences) before staining for IL6, IL10 and IL12p40/70 (BD Biosciences). Cytokine production was analyzed by flow cytometry.

2.5. Antigen uptake assay

Quantitative analysis of DC endocytosis was performed as described previously [14], with minor modifications. To study mDC endocytosis, 1ml of fresh blood was incubated with FITC-albumin (10 μ g/ml, Sigma-Aldrich, St. Louis, Mo) at either 37 or 4°C (as negative control) for 30 minutes. Freshly taken blood was stained with CD1c, HLADR

and CD19, followed by flow cytometric analysis. The CD1c⁺HLADR⁺CD19⁻ population was classified as mDC1. Myeloid DC1 expressing FITC represented mDC1 taking up antigen (albumin). In the functional studies, 1 µl sera from HC or uveitis patients was added into MoDCs in the volume of 200 µl for 24 hours before the antigen uptake assay was conducted. To investigate the effect of TNF α on antigen uptake, MoDCs were treated with recombinant TNF α at different concentrations (0.1 ng/ml, 1 ng/ml and 10 ng/ml) for 24 hours. The antigen uptake assay was subsequently conducted.

2.6. DC-stimulated T cell responses

MoDCs-stimulated T cell responses were performed as described previously [14], with minor modifications. Briefly, MoDCs with high albumin uptake (Alb⁺) and low albumin uptake (Alb⁻) were sorted based on FITC-albumin expression on MoDCs. Magnetic bead separation (Miltenyi Biotec) was used to isolate CD4⁺ T cells in peripheral blood mononuclear cells (PBMCs) from healthy controls. Purity of isolated fractions consistently averaged to be >95% for CD4⁺ T cells. Isolated DCs were co-cultured with CFSE-labeled allogenic CD4⁺ T cells at a ratio of 1:10 (DC:T) for 5 days with the help of anti-CD3 stimulation (1 µg/ml). CD4⁺ T cell proliferation was characterized by flow cytometric analysis (percentage of cells exhibiting CFSE fluorescence dilution) and negative expression of CD62L was characterized with activated CD4⁺ T cells.

2.7. Statistical analysis

Data is presented as mean \pm SD. Results shown were representative of at least three independent experiments. Statistical comparisons were performed using the Student's *t*

test for two groups, or ANOVA tests for analyzing more than two groups. $P < 0.05$ was considered as significant. Analyses were accomplished using Prism 6 software (GraphPad Software, Inc., La Jolla, CA).

3. Results

3.1. Elevation of blood CD1c⁺ mDC1 in uveitis patients

The percentages and absolute numbers of blood DC subsets were determined in uveitis patients and HCs. DC subsets are shown in figure 1A. Lin1⁻HLADR⁺ cells were defined as total DCs in which cells were subsequently identified as CD1c⁺ mDC1, CD141⁺ mDC2 and CD303⁺ pDCs [9]. In uveitis patients, the frequency ($p=0.0025$) and number ($p=0.026$) of blood CD1c⁺ mDC1 were significantly increased (Fig.1B and C), and the frequency of blood pDCs was decreased as compared to the levels observed in HCs (data not shown). No differences of mDC2 were detected between uveitis patients and HCs (data not show). Furthermore, we categorized the CD1c⁺ mDC1 frequency based on therapy and disease status in uveitis patients. No difference was observed in untreated versus treated patients (Fig.1D). However, CD1c⁺ mDC1 were increased in active uveitis compared to quiescent uveitis ($p=0.03$, Fig.1E). In summary, increased CD1c⁺ mDC1 expression was independent of the direct effect from therapy but highly correlated with disease activity. Demographic data of the study subjects were listed in the table.

3.2. Elevation of HLADR expression on CD1c⁺ mDC1 in uveitis patients

To determine whether blood CD1c⁺ mDC1 were more mature in uveitis patients compared with HCs, HLADR and co-stimulatory molecules (CD80 and CD86) on blood

CD1c⁺ mDC1 were analyzed by flow cytometry. HLADR expression, a marker of maturation and antigen presentation, was significantly higher on CD1c⁺ mDC1 in uveitis patients compared to HCs ($p=0.002$, Fig. 2A). We sought to determine whether HLADR expression was affected by therapy or disease activity. Interestingly, HLADR expression was significantly higher in the untreated group compared to the treated group ($p=0.0075$) and HCs ($p<0.0001$, Fig. 2B), indicating that therapy suppressed HLADR expression. We further evaluated whether higher HLADR expression correlated to disease status. Unexpectedly, higher HLADR expression was observed in the quiescent uveitis patients ($p=0.0009$) not in the active uveitis patients, as compared to HCs (Fig. 2C). The higher level of HLADR expression in the quiescent patients was probably due to the lower rates of therapy in this group as higher HLADR expression levels were observed in quiescent patients when not on treatment ($p=0.029$, Fig.2D).

We hypothesized that CD1c⁺ mDC1 from patients might be hyperresponsive to additional stimulation. To test this we stimulated CD1c⁺ mDC1 with LPS overnight and then measured HLADR expression. The HLADR expression level was significantly higher in uveitis patients after additional LPS stimulation, as compared to that in patients without stimulation ($p=0.01$, Fig. 2E). The co-stimulatory markers CD80 and CD86 were similar on CD1c⁺ mDC1 between uveitis patients and HCs (data not shown).

3.3. Impaired cytokine production of uveitic CD1c⁺ mDC1 upon additional TLR4 stimulation

Previous studies have shown that DCs exhaust their capacity to produce cytokines after the induction of maturation [15]. Consequently, we examined whether mature CD1c⁺ mDC1 in uveitis patients also displayed a similar exhaustion pattern in cytokine production. To test this, IL6, IL10 and IL12p40/70 produced from CD1c⁺ mDC1 of uveitis

patients (n=12) and HCs (n=9) were detected by flow cytometry. Uveitis patients were randomly selected from the cohort along with appropriately age-matched and gender-matched healthy controls. IL6 (Fig.3A), IL10 (Fig.3B) and IL12p40/70 (Fig.3C) production were similar between uveitis patients and HCs *ex vivo*. After overnight stimulation with LPS, the production of IL6 ($p=0.0002$, Fig.3A), IL10 ($p=0.01$, Fig.3B) and IL12p40/70 ($p=0.01$, Fig.3C) from mDC1 was significantly increased compared with the unstimulated group from HCs. The cytokine production was not significantly increased after LPS stimulation in uveitis patients when compared with the unstimulated group from uveitis (Fig. 3). The lack of *in vitro* responses to additional LPS stimulation suggests that cytokine production abilities of mature CD1c⁺ mDC1 from uveitis patients are exhausted.

3.4. Lower antigen uptake by CD1c⁺ mDC1 and its subpopulation CD1c^{hi} mDC1 in uveitis patients

The capacity of mDCs to take up antigens is an indicator of their level of maturation, with higher levels of antigen take-up in the immature state moving to an antigen presentation role (supported in part by increasing levels of HLADR) as the mDC matures [16]. We observed a significant decrease of antigen uptake in CD1c⁺ mDC1 in uveitis patients compared to HCs ($p=0.04$, Fig. 4A, B). Flow cytometric analysis showed that CD1c⁺ mDC1 contained two subpopulations, CD1c^{hi} and CD1c^{lo} mDC1 (Fig.4C). Interestingly, only 1% of CD1c^{hi} mDC1 took up antigens while 43% of CD1c^{lo} mDC1 took up antigens (Fig. 4D, E). Importantly, we observed that CD1c^{hi} mDC1 were increased in uveitis patients compared to HCs ($p=0.001$, Fig. 4F), indicating that the lower antigen uptake in the whole CD1c⁺ mDC1 population was likely due to the increased numbers of CD1c^{hi} mDC1 in the uveitis group. Furthermore, high expression of CD1c was associated with

higher expression of HLADR in both HCs ($p=0.0001$, Fig. 4G) and uveitis patients ($p=0.002$, Fig. 4H).

3.5. *In vitro* model showed MoDCs with lower antigen uptake associated with higher T cell proliferation

Having shown that patients with uveitis have higher levels of CD1c⁺ mDC1 and of the CD1c^{hi} subgroup, and that these mDCs have a relatively 'mature' profile characterized by higher HLADR and lower antigen uptake, we wished to investigate the functional consequences of this. For our functional studies, we used MoDCs as a substitute for mDCs in an *in vitro* model [12, 13, 17]. Six to Seven days are usually needed for monocytes to fully differentiate into mDCs *in vitro*. This allowed us to observe the expression of CD1c on MoDCs during differentiation (Fig. 7C, D). MoDCs were loaded with FITC-albumin for 30 minutes and two subpopulations were sorted by flow cytometry based on their ability to take up antigens, which were FITC-albumin⁺ and FITC-albumin⁻ MoDCs (Fig. 5A). We cultured allogenic CD4⁺ T cells with either Albumin⁺ or Albumin⁻ MoDCs for 5 days. CD62L⁻CD4⁺ T cell (activated T cells) proliferation was significantly increased in the group of Albumin⁻ MoDCs, as compared to the group of Albumin⁺ MoDCs. This was shown by the percentage of CFSE dilution on CD62L⁻CD4⁺ T cells (Fig. 5B, C), indicating that MoDCs with lower antigen uptake could induce more CD4⁺ T cell proliferation and activation.

3.6. Uveitic sera and exogenous TNF α reduced antigen uptake in MoDCs

Having confirmed in our *in vitro* model that the MoDCs with lower antigen uptake (modeling the low-antigen-uptake mDC1 identified earlier) induced higher levels of T cell proliferation with a higher proportion of activated CD62L⁺ T cells, we wished to investigate this further, specifically whether there was a factor in uveitis serum that might be inducing this more mature DC phenotype. In order to investigate this, we performed overnight incubation with MoDCs treated with the addition of sera from patients with uveitis or HCs. We found that MoDCs had less antigen uptake after they were treated with sera from uveitis patients compared with MoDCs treated with sera from HCs (Fig. 6A, B and C).

We hypothesized that this might be related to higher levels of inflammatory cytokines such as TNF α and IL-17a in patient sera. TNF α is known to contribute to DC maturation, plays a key role in the pathogenesis of uveitis, is present at high concentration levels both in the aqueous humor and in the serum [18-22], and is a successful target in the treatment of uveitis [23]. Although less well-established IL17a has also shown to be increased in patients with uveitis, and an anti-IL17 agent, secukinumab, has been trialed for treatment of uveitis [24, 25]. We found that TNF α , but not IL17a (data not shown), significantly reduced MoDCs' antigen uptake in a dose-dependent manner (Fig.6D, E and F). Furthermore we found that treatment with TNF α increased CD1c expression compared to controls (Fig.7A).

One of the key pathways that TNF α exerts its effect is via phosphorylation of p38 MAPK [26]. We hypothesized that TNF-induced upregulation of CD1c expression was mediated via this pathway. Treatment of MoDCs with TNF α caused an increased phosphorylation of p38 MAPK (Fig.7B). Conversely, blockage of this pathway by addition of SB203580, an inhibitor of p38 MAPK (Fig.7C, D), inhibited CD1c expression during MoDC differentiation. Although monocytes don't express CD1c, CD1c expression was

induced in MoDCs on day 1 and increased dramatically every subsequent day of differentiation. The inhibition of CD1c expression by SB203580 was observed on day 1 and reached statistical significance on days 3 and 5 (Fig. 7C, D). Taken together, this data suggests that the known increased TNF α levels in patient sera induces a more mature DC phenotype characterized by high CD1c expression and less antigen uptake, and this is mediated (at least in part) by the p38 MAPK pathway.

4. Discussion

In this study, we have shown that blood CD1c⁺ mDC1 is increased in autoimmune non-infectious uveitis patients compared with HCs. Blood CD1c⁺ mDC1 from uveitis patients show a mature phenotype, characterized by less antigen uptake and high HLADR expression. CD1c^{hi} mDC1 is identified as the mDC1 subpopulation with lower antigen uptake, and associated with higher HLADR expression and a capability to induce higher levels of CD4⁺ T cell activation and proliferation. Additionally, we present data suggesting these effects are mediated by the increased level of TNF α in uveitis patient sera, which appears to promote CD1c⁺ mDC1 maturation through the p38 MAPK pathway. Our data provide support for the role of blood CD1c⁺ mDC1, especially CD1c^{hi} mDC1, in the pathogenesis of non-infectious uveitis in humans.

Previous studies have provided strong evidence that non-infectious uveitis is an autoimmune disease [27]. A key step in autoimmune disease centers on mDC taking up self-antigens thereby initiating antigen-specific T cell immune responses. Among blood mDCs, CD1c⁺ mDC1 represent a major mDC population in the peripheral blood, and have been shown to be increased in other autoimmune disease such as rheumatoid arthritis [28]. Our study shows that CD1c⁺ mDC1 is increased in uveitis patients

compared with HCs. It should be noted from our data that the increased CD1c⁺ mDC1 appear to be independent of therapy but positively correlated with disease activity [29].

The maturation state of DCs affects their ability to induce adaptive immune responses [30]. We noted that CD1c⁺ mDC1 from uveitis patients were more mature characterized by an increased expression of HLADR, although interestingly we observed no differences in the expression of co-stimulatory molecules (CD80 and CD86). Interestingly, in another study of mDCs from patients with uveitis, Kim et al did observe an increase in CD86. The differences may lie in the broader spectrum of DCs considered in their study since their isolation process selects out mDCs with subsequent co-staining of CD11c. This will lead to both mDC1 and mDC2 populations being included whereas our study was restricted to changes in the mDC1 population. Interestingly even in their study the changes on CD86 were relatively minor compared to the significant elevation in HLADR seen in both studies [8]. It remains of interest to understand why CD1c⁺ mDC1 only exhibits high HLADR but no other co-stimulatory molecules, particularly as alterations in relative contribution of signal 1 and signal 2 may alter function and differentiation of T cells [31].

In addition to changes in their surface phenotype, maturation is associated with a number of other important alterations in mDC function. Cytokine production is affected, with DCs displaying a different cytokine repertoire as they mature [32], and activated DCs becoming refractory to further stimulation due to exhaustion of their cytokine-producing capacity [15]. In keeping with these findings, our data show that mature CD1c⁺ mDC1 from uveitis patients fail to increase their cytokine production in response to additional LPS stimulation *in vitro*.

It is also well established that normal DC maturation is associated with their transition from a phenotype with high levels of antigen uptake, to one, which primarily presents antigen (and has low levels of uptake). We have confirmed that this behavior is

also seen in the context of uveitis, with CD1c⁺ mDC1 in uveitis patients showing lower levels of antigen uptake than healthy controls, a characteristic which could also be induced by co-culture of MoDC with sera from uveitis patients [33]. Interestingly, we observed significant numbers of a CD1c^{hi} subpopulation in the uveitis patients, associated with a more mature phenotype, as indicated by higher HLADR expression and lower levels of antigen uptake, which is also supported by the study from mouse DCs [34]. The function of CD1 molecules is to present lipid, both endogenous and pathogen-derived [35]. Similar to class I MHC they associate with β 2-microglobulin on the surface of various types of cells, particularly antigen-presenting cells (APCs), but differ in that (1) they are not polymorphic, and (2) have hydrophobic binding pockets that allow them to bind the hydrocarbon chains of lipids [36-38]. CD1c (along with CD1a and CD1b) is recognized by CD1-restricted T cells whereas CD1d is recognized by invariant natural killer T (iNKT) cells. CD1c may present a wide range of antigens and indeed its ability to present antigen from important human pathogens such as mycobacteria, leading to its proposal as a candidate for vaccine development [36]. CD1c⁺ mDC1 can be found not only in the peripheral blood but also in various tissues including the intestinal lamina propria [39] and in the synovial fluid in patients with rheumatoid arthritis [28]. In both these studies the CD1c⁺ mDC1 identified in the tissues express higher levels of activation markers than those in the peripheral blood. CD1c⁺ mDC1 levels can however be affected by treatment, as demonstrated in the study by Jolivel, et al., where CD1c⁺ mDC1 levels were reduced in laquinimod-treated multiple sclerosis patients in comparison to untreated patients and healthy donors [40]. However, our study supports that level of CD1c is not due to the treatment but is associated with disease activity.

A critical role of mature mDCs is to promote T cell proliferation. In our study we showed, as expected, that not only did uveitis patients have a higher proportion of mDC1 with a more mature phenotype, but that (1) these mDC1 induced higher levels of T cell

proliferation and (2) a higher proportion of these were CD62L⁻. CD62L is expressed on naïve T cells and central memory T cells, which help T cells to localize in secondary lymphoid organs. Effector memory T cells do not express CD62L. They circulate in the periphery and have immediate effector functions upon encountering antigens [41]. It is likely that this lies at the center of the generation of autoimmunity in non-infectious uveitis. It is known that IRBP and S-antigen can initiate innate and adaptive immune responses by attracting immature DCs [42]. In this context our finding of increased levels of CD1c⁺ mDC1 cells in uveitis patients, is of interest, particularly in its implications for the generation of auto-reactive uveitic T cell immune responses, which may underlie the pathogenesis of noninfectious uveitis.

Several studies have demonstrated that TNF α is increased in sera of uveitis patients [18-22]. We have found that TNF α -treated MoDCs show lower antigen uptake and increase CD1c expression. TNF α can promote DC maturation [43] and activate p38 MAPK [26]. Studies have shown that SB203580 (p38 inhibitor) inhibits the up-regulation of CD1a, CD40, CD80, CD86, HLA-DR, and DC maturation marker CD83, which is induced by LPS and TNF α [26]. In addition, SB203580 inhibits the enhancement of the allostimulatory capacity and partially prevents the down-regulation of FITC-dextran uptake induced by LPS and TNF α [26]. Consistently, we have found that TNF α decreases FITC-albumin antigen uptake and increases CD1c expression through regulating p38 MAPK. These data suggest that phosphorylation of p38 MAPK may be a key pathway in the maturation of DCs in uveitis.

In conclusion, a significant increased blood CD1c⁺ mDC1 with mature phenotype is identified in uveitis patients. A CD1c^{hi} mDC1 subpopulation is identified which is increased in patients and is characterized by a relatively mature phenotype including lower antigen uptake and increased induction of T cell responses. We also provide

evidence that these changes may be caused by the higher levels of TNF α present in the serum of patients with uveitis, and which is p38 MAPK-dependent. Further understanding of the role of blood CD1c⁺ mDC1 in non-infectious uveitis may answer key questions regarding the initial events in the pathogenesis of uveitis and provide novel therapeutic opportunities for treating patients with this potentially blinding autoimmune disease.

Acknowledgements

This project was supported by the Intramural Research Program of the NIH, National Institute of Eye (NEI). We thank Rafael Villasmil and Julie Laux for assisting on flow cytometry. We thank Cheng-Rong Yu and Fan Pan for critical reading the manuscript.

Conflict of Interest Disclosure

The authors have no financial conflicts of interest.

References

- [1] T. Larson, R.B. Nussenblatt, H.N. Sen, Emerging drugs for uveitis, *Expert opinion on emerging drugs*, 16 (2011) 309-322.
- [2] A. Amadi-Obi, C.R. Yu, X. Liu, R.M. Mahdi, G.L. Clarke, R.B. Nussenblatt, I. Gery, Y.S. Lee, C.E. Egwuagu, TH17 cells contribute to uveitis and scleritis and are expanded by IL-2 and inhibited by IL-27/STAT1, *Nature medicine*, 13 (2007) 711-718.
- [3] R. Zhou, R. Horai, P.B. Silver, M.J. Mattapallil, C.R. Zarate-Blades, W.P. Chong, J. Chen, R.C. Rigden, R. Villasmil, R.R. Caspi, The living eye "disarms" uncommitted autoreactive T cells by converting them to Foxp3(+) regulatory cells following local antigen recognition, *Journal of immunology (Baltimore, Md. : 1950)*, 188 (2012) 1742-1750.
- [4] R.S. Grajewski, A.M. Hansen, R.K. Agarwal, M. Kronenberg, S. Sidobre, S.B. Su, P.B. Silver, M. Tsuji, R.W. Franck, A.P. Lawton, C.C. Chan, R.R. Caspi, Activation of invariant NKT cells ameliorates experimental ocular autoimmunity by a mechanism involving innate IFN-gamma production and dampening of the adaptive Th1 and Th17 responses, *Journal of immunology (Baltimore, Md. : 1950)*, 181 (2008) 4791-4797.

- [5] V.K. Kuchroo, P.S. Ohashi, R.B. Sartor, C.G. Vinuesa, Dysregulation of immune homeostasis in autoimmune diseases, *Nature medicine*, 18 (2012) 42-47.
- [6] J. Tang, W. Zhu, P.B. Silver, S.B. Su, C.C. Chan, R.R. Caspi, Autoimmune uveitis elicited with antigen-pulsed dendritic cells has a distinct clinical signature and is driven by unique effector mechanisms: initial encounter with autoantigen defines disease phenotype, *Journal of immunology (Baltimore, Md. : 1950)*, 178 (2007) 5578-5587.
- [7] D. Liang, A. Zuo, H. Shao, W.K. Born, R.L. O'Brien, H.J. Kaplan, D. Sun, Role of CD25+ dendritic cells in the generation of Th17 autoreactive T cells in autoimmune experimental uveitis, *Journal of immunology (Baltimore, Md. : 1950)*, 188 (2012) 5785-5791.
- [8] T.W. Kim, J.S. Kang, J.M. Kong, S. Bae, Y. Yu, H. Chung, H.G. Yu, Maturation profiles of peripheral blood dendritic cells in patients with endogenous uveitis, *Immunology letters*, 142 (2012) 14-19.
- [9] A. Dzionek, A. Fuchs, P. Schmidt, S. Cremer, M. Zysk, S. Miltenyi, D.W. Buck, J. Schmitz, BDCA-2, BDCA-3, and BDCA-4: three markers for distinct subsets of dendritic cells in human peripheral blood, *Journal of immunology (Baltimore, Md. : 1950)*, 165 (2000) 6037-6046.
- [10] A.K. Denniston, P. Tomlins, G.P. Williams, S. Kottoor, I. Khan, K. Oswal, M. Salmon, G.R. Wallace, S. Rauz, P.I. Murray, S.J. Curnow, Aqueous humor suppression of dendritic cell function helps maintain immune regulation in the eye during human uveitis, *Investigative ophthalmology & visual science*, 53 (2012) 888-896.
- [11] D.A. Jabs, R.B. Nussenblatt, J.T. Rosenbaum, Standardization of uveitis nomenclature for reporting clinical data. Results of the First International Workshop, *American journal of ophthalmology*, 140 (2005) 509-516.
- [12] J. Carrion, E. Scisci, B. Miles, G.J. Sabino, A.E. Zeituni, Y. Gu, A. Bear, C.A. Genco, D.L. Brown, C.W. Cutler, Microbial carriage state of peripheral blood dendritic cells (DCs) in chronic periodontitis influences DC differentiation, atherogenic potential, *Journal of immunology (Baltimore, Md. : 1950)*, 189 (2012) 3178-3187.
- [13] P. Chen, Q. Sun, Y. Huang, M.G. Atta, S. Turban, D.L. Segev, K.A. Marr, F.F. Naqvi, N. Alachkar, E.S. Kraus, K.L. Womer, Blood dendritic cell levels associated with impaired IL-12 production and T-cell deficiency in patients with kidney disease: implications for post-transplant viral infections, *Transplant international : official journal of the European Society for Organ Transplantation*, 27 (2014) 1069-1076.
- [14] Y. Huang, P. Johnston, B. Zhang, A. Zakari, T. Chowdhry, R.R. Smith, E. Marban, H. Rabb, K.L. Womer, Kidney-derived stromal cells modulate dendritic and T cell responses, *Journal of the American Society of Nephrology : JASN*, 20 (2009) 831-841.
- [15] A. Langenkamp, M. Messi, A. Lanzavecchia, F. Sallusto, Kinetics of dendritic cell activation: impact on priming of TH1, TH2 and nonpolarized T cells, *Nature immunology*, 1 (2000) 311-316.
- [16] I. Mellman, R.M. Steinman, Dendritic cells: specialized and regulated antigen processing machines, *Cell*, 106 (2001) 255-258.
- [17] B. Leon, C. Ardavin, Monocyte-derived dendritic cells in innate and adaptive immunity, *Immunology and cell biology*, 86 (2008) 320-324.

- [18] M. Santos Lacomba, C. Marcos Martin, J.M. Gallardo Galera, M.A. Gomez Vidal, E. Collantes Estevez, R. Ramirez Chamond, M. Omar, Aqueous humor and serum tumor necrosis factor-alpha in clinical uveitis, *Ophthalmic research*, 33 (2001) 251-255.
- [19] J.K. Ahn, H.G. Yu, H. Chung, Y.G. Park, Intraocular cytokine environment in active Behcet uveitis, *American journal of ophthalmology*, 142 (2006) 429-434.
- [20] M. Kramer, N. Goldenberg-Cohen, R. Axer-Siegel, D. Weinberger, Y. Cohen, Y. Monselise, Inflammatory reaction in acute retinal artery occlusion: cytokine levels in aqueous humor and serum, *Ocular immunology and inflammation*, 13 (2005) 305-310.
- [21] V. Perez-Guijo, M. Santos-Lacomba, M. Sanchez-Hernandez, C. Castro-Villegas Mdel, J.M. Gallardo-Galera, E. Collantes-Estevez, Tumour necrosis factor-alpha levels in aqueous humour and serum from patients with uveitis: the involvement of HLA-B27, *Current medical research and opinion*, 20 (2004) 155-157.
- [22] Y. Xu, W. Chen, H. Lu, X. Hu, S. Li, J. Wang, L. Zhao, The expression of cytokines in the aqueous humor and serum during endotoxin-induced uveitis in C3H/HeN mice, *Molecular vision*, 16 (2010) 1689-1695.
- [23] P. Neri, M. Zucchi, P. Allegri, M. Lettieri, C. Mariotti, A. Giovannini, Adalimumab (Humira): a promising monoclonal anti-tumor necrosis factor alpha in ophthalmology, *International ophthalmology*, 31 (2011) 165-173.
- [24] S. Jawad, B. Liu, E. Agron, R.B. Nussenblatt, H.N. Sen, Elevated serum levels of interleukin-17A in uveitis patients, *Ocular immunology and inflammation*, 21 (2013) 434-439.
- [25] A.D. Dick, I. Tugal-Tutkun, S. Foster, M. Zierhut, S.H. Melissa Liew, V. Bezlyak, S. Androudi, Secukinumab in the treatment of noninfectious uveitis: results of three randomized, controlled clinical trials, *Ophthalmology*, 120 (2013) 777-787.
- [26] J.F. Arrighi, M. Rebsamen, F. Rousset, V. Kindler, C. Hauser, A critical role for p38 mitogen-activated protein kinase in the maturation of human blood-derived dendritic cells induced by lipopolysaccharide, TNF-alpha, and contact sensitizers, *Journal of immunology (Baltimore, Md. : 1950)*, 166 (2001) 3837-3845.
- [27] Z. Li, B. Liu, A. Maminishkis, S.P. Mahesh, S. Yeh, J. Lew, W.K. Lim, H.N. Sen, G. Clarke, R. Buggage, S.S. Miller, R.B. Nussenblatt, Gene expression profiling in autoimmune noninfectious uveitis disease, *Journal of immunology (Baltimore, Md. : 1950)*, 181 (2008) 5147-5157.
- [28] F.M. Moret, C.E. Hack, K.M. van der Wurff-Jacobs, W. de Jager, T.R. Radstake, F.P. Lafeber, J.A. van Roon, Intra-articular CD1c-expressing myeloid dendritic cells from rheumatoid arthritis patients express a unique set of T cell-attracting chemokines and spontaneously induce Th1, Th17 and Th2 cell activity, *Arthritis research & therapy*, 15 (2013) R155.
- [29] P. Chen, W. Tucker, S. Hannes, B. Liu, H. Si, A. Gupta, R.W. Lee, H.N. Sen, R.B. Nussenblatt, Levels of Blood CD1c+ mDC1 and CD1chi mDC1 Subpopulation Reflect Disease Activity in Noninfectious Uveitis, *Investigative ophthalmology & visual science*, 56 (2015) 346-352.
- [30] D. Dissanayake, H. Hall, N. Berg-Brown, A.R. Elford, S.R. Hamilton, K. Murakami, L.S. Deluca, J.L. Gommerman, P.S. Ohashi, Nuclear factor-kappaB1 controls the functional maturation of dendritic cells and prevents the activation of autoreactive T cells, *Nature medicine*, 17 (2011) 1663-1667.

- [31] A. Noble, Review article: molecular signals and genetic reprogramming in peripheral T-cell differentiation, *Immunology*, 101 (2000) 289-299.
- [32] B. de Saint-Vis, I. Fugier-Vivier, C. Massacrier, C. Gaillard, B. Vanbervliet, S. Ait-Yahia, J. Banchereau, Y.J. Liu, S. Lebecque, C. Caux, The cytokine profile expressed by human dendritic cells is dependent on cell subtype and mode of activation, *Journal of immunology (Baltimore, Md. : 1950)*, 160 (1998) 1666-1676.
- [33] F. Sallusto, M. Cella, C. Danieli, A. Lanzavecchia, Dendritic cells use macropinocytosis and the mannose receptor to concentrate macromolecules in the major histocompatibility complex class II compartment: downregulation by cytokines and bacterial products, *The Journal of experimental medicine*, 182 (1995) 389-400.
- [34] M.B. Lutz, N. Kukutsch, A.L. Ogilvie, S. Rossner, F. Koch, N. Romani, G. Schuler, An advanced culture method for generating large quantities of highly pure dendritic cells from mouse bone marrow, *Journal of immunological methods*, 223 (1999) 77-92.
- [35] S. Porcelli, M.B. Brenner, J.L. Greenstein, S.P. Balk, C. Terhorst, P.A. Bleicher, Recognition of cluster of differentiation 1 antigens by human CD4-CD8-cytolytic T lymphocytes, *Nature*, 341 (1989) 447-450.
- [36] E.J. Adams, Diverse antigen presentation by the Group 1 CD1 molecule, CD1c, *Molecular immunology*, 55 (2013) 182-185.
- [37] M. Brigl, M.B. Brenner, CD1: antigen presentation and T cell function, *Annual review of immunology*, 22 (2004) 817-890.
- [38] D.B. Moody, D.M. Zajonc, I.A. Wilson, Anatomy of CD1-lipid antigen complexes, *Nature reviews. Immunology*, 5 (2005) 387-399.
- [39] S.M. Dillon, L.M. Rogers, R. Howe, L.A. Hostetler, J. Buhrman, M.D. McCarter, C.C. Wilson, Human intestinal lamina propria CD1c+ dendritic cells display an activated phenotype at steady state and produce IL-23 in response to TLR7/8 stimulation, *Journal of immunology (Baltimore, Md. : 1950)*, 184 (2010) 6612-6621.
- [40] V. Jolivel, F. Luessi, J. Masri, S.H. Kraus, M. Hubo, L. Poisa-Beiro, S. Klebow, M. Paterka, N. Yogev, H. Tumani, R. Furlan, V. Siffrin, H. Jonuleit, F. Zipp, A. Waisman, Modulation of dendritic cell properties by laquinimod as a mechanism for modulating multiple sclerosis, *Brain : a journal of neurology*, 136 (2013) 1048-1066.
- [41] R.M. Steinman, Decisions about dendritic cells: past, present, and future, *Annual review of immunology*, 30 (2012) 1-22.
- [42] O.M. Howard, H.F. Dong, S.B. Su, R.R. Caspi, X. Chen, P. Plotz, J.J. Oppenheim, Autoantigens signal through chemokine receptors: uveitis antigens induce CXCR3- and CXCR5-expressing lymphocytes and immature dendritic cells to migrate, *Blood*, 105 (2005) 4207-4214.
- [43] S. Miwa, H. Nishida, Y. Tanzawa, M. Takata, A. Takeuchi, N. Yamamoto, T. Shirai, K. Hayashi, H. Kimura, K. Igarashi, E. Mizukoshi, Y. Nakamoto, S. Kaneko, H. Tsuchiya, TNF-alpha and tumor lysate promote the maturation of dendritic cells for immunotherapy for advanced malignant bone and soft tissue tumors, *PloS one*, 7 (2012) e52926.

Figure legends

Figure 1.

Increased frequency and number of blood CD1c⁺ mDC1 in uveitis patients (n=32) as compared to HCs (n=64). **(A)** Representative scattergrams from flow cytometry analysis of dendritic cell subtypes CD1c⁺ mDC1, CD141⁺ mDC2 and CD303⁺ pDCs gated on Lin1⁻HLADR⁺ total DCs. Percentages **(B)** and absolute numbers **(C)** of mDC1 were analyzed between uveitis patients and HCs. Comparison of CD1c⁺ mDC1 percentage in treated and untreated groups **(D)**, as well as in quiescent and active uveitis groups **(E)**. Student's t test was used in **(B)** and **(C)**. One-way ANOVA analysis and post ANOVA analysis (Tukey's multiple comparisons test) were used. ANOVA p <0.0001 in **(D)** and **(E)**.

Figure 2.

Higher HLADR expression on CD1c⁺ mDC1 in uveitis patients compared to HCs. **(A)** Mean fluorescence intensity (MFI) of HLADR was measured by flow cytometry between uveitis patients (n=37) and HCs (n=39). **(B)** HLADR expression on mDC1 was compared among untreated, treated and HC groups. **(C)** HLADR expression on mDC1 was compared in quiescent, active and HC groups. **(D)** HLADR expression was compared in untreated and treated groups of quiescent uveitis patients, and HCs. **(E)** HLADR expression was measured on mDC1 with or without LPS stimulation overnight *in vitro* in both HCs (unstimulated group; LPS group) and uveitis patients (unstimulated group; LPS group). Student's t test was used in **(A)**. One-way ANOVA analysis and post ANOVA analysis (Tukey's multiple comparisons test) were used in **(B)** through **(E)**. The ANOVA p values are 0.03 in **(B)**, 0.002 in **(C)**, 0.0003 in **(D)** and less than 0.0001 in **(E)**.

Figure 3.

Cytokine production of mDC1 was exhausted upon additional TLR4 stimulation in uveitis patients. Myeloid DC1 producing-IL6 **(A)**, IL10 **(B)** and IL12P40/70 **(C)** were measured in HCs (n=9) and uveitis patients (n=12) with or without LPS stimulation *in vitro*. The cytokine production was detected by intracellular staining and analyzed by flow cytometry. One-way ANOVA analysis and post ANOVA analysis (Sidak's multiple comparisons test) were used. ANOVA overall p values are 0.049 **(A)**, 0.008 **(B)**, and 0.039 **(C)**.

Figure 4.

Low antigen uptake in CD1c⁺ mDC1 in uveitis patients. **(A)** Fresh blood from either HCs (n=23) or uveitis patients (n=30) was loaded with FITC-albumin (antigen) for 30 minutes. Representative antigen uptake of mDC1 measured by flow cytometry is shown. The negative control was MoDCs cultured in 4°C. **(B)** Significant differences in antigen uptake of mDC1 from uveitis patients and HCs. **(C)** CD1c⁺ mDC1 were separated as CD1c^{hi} and CD1c^{lo} mDC1 by flow cytometry. CD1c^{hi} mDC1 **(D)** had less antigen uptake when compared with CD1c^{lo} mDC1 **(E)**. **(F)** CD1c^{hi} mDC1 were increased in uveitis patients (n=20) as compared with HCs (n=23). High CD1c expression on mDC1 correlated with high HLADR expression in both HCs **(G)** and uveitis patients **(H)**.

Figure 5.

MoDCs with lower antigen uptake enhanced CD62L⁻CD4⁺ T cell proliferation. **(A)** To address the function of mDC1 with decreased antigen uptake, monocyte-derived DC (MoDCs) were used *in vitro*. MoDCs were loaded with FITC-albumin for 30 minutes and sorted by flow cytometry based on high and low expression of FITC-albumin (Albumin⁺ and Albumin⁻). **(B, C)** Albumin⁺ or Albumin⁻ MoDCs co-cultured with allogenic CFSE

labeled CD4⁺ T cells for 5 days. T cell proliferation and activation were measured by diluted expression of CFSE and low expression of CD62L. The figures are representative of four independent experiments. One-way ANOVA analysis was used; $p=0.04$ in **(C)**.

Figure 6.

Treatment with uveitis sera or TNF α reduced the antigen uptake of MoDCs. **(A)** Antigen uptake assay was done in two groups: MoDCs were treated with sera from either HCs (HC group, n=6) or patients with uveitis (Uveitis group, n=6). The percentage of FITC-albumin⁺ **(B)** and MFI of FITC-albumin⁺ **(C)** on MoDCs were calculated. The experiment was repeated six times. **(D)** An antigen uptake assay was done in five groups: MoDCs were cultured at 4°C (negative control); MoDCs were cultured at 37°C without TNF α stimulation (untreated group), or with TNF α stimulation overnight with different concentrations (0.1 ng/ml, 1 ng/ml and 10 ng/ml). The percentage of FITC-albumin⁺ **(E)** and MFI of FITC-albumin⁺ **(F)** on MoDCs were calculated. The dose-dependent experiment was repeated two times. One-way ANOVA analysis was used and the total $p=0.0008$ in **(E)** and 0.04 in **(F)**.

Figure 7.

TNF α increased CD1c expression by p38 MAPK signaling. **(A)** CD1c expression was measured by flow cytometry on MoDCs with (TNF α group) or without (control group) TNF α stimulation *in vitro* for 24 h. The experiment was repeated two times. Dotted line is control group; solid line is TNF α group. **(B)** Phosphorylated p38 was measured on MoDCs treated with (TNF α group) or without (control group) TNF α for 30 minutes. The experiment was repeated four times. Dotted line is isotype control; Dash line is control

group; solid line is TNF α group. **(C)** MoDCs were induced from monocytes by GM-CSF and IL-4. P38 inhibitor SB203580 was added at a concentration of 10 μ mol/L into the culture at day 0. During MoDC differentiation, CD1c expression was measured by flow cytometry on days 1, 3 and 5. The filled grey area is the p38 inhibitor group; and the solid line is the untreated group. The experiment was repeated two separate times. Unpaired student's *t* test was used in **(A)** and **(B)**. One-way ANOVA analysis was used and ANOVA $p < 0.0001$ in **(D)**.




Fig. 1




Fig.2




Fig.3

ACCEPTED




Fig.4




Fig.5

ACCEPTED




Fig.6




Fig.7

ACCEPTED

Table. Characteristics of Uveitis Patients and HCs

Parameter	Uveitis (n=74)	HCs (n=96)
Age (y), median (range)	44 (17-74)	46 (19-76)
Gender [n (%)]		
Male	44 (59)	61 (64)
Female	30 (41)	35 (36)
Race [n (%)]		
Caucasian	28 (38)	80 (83)
African American	28 (38)	15 (16)
Others	18 (24)	1 (1)
Anatomic type of uveitis [n (%)]		
Anterior	10 (14)	
Intermediate	7 (9)	
Posterior	17 (23)	
Panuveitis	40 (54)	
Disease association [n (%)]		
Sarcoidosis	17 (23)	
Idiopathic	32 (43)	
Birdshot	8 (11)	
VKH	5 (7)	
Behcet's disease	8 (11)	
Serpiginous choroidopathy	4 (5)	
Systemic Immunosuppressive Therapy [n (%)]		
Active uveitis [n (%)]	18 (24)	N/A

Highlights

- CD1c⁺ mDC1 were increased in uveitis patients.
- CD1c⁺ mDC1 exhibited less antigen uptake in uveitis patients.
- CD1c^{hi} mDC1 subpopulation with less antigen uptake was increased in patients.
- MoDCs with less antigen uptake induced more CD4⁺CD62L⁻ T cells proliferation.
- TNF α contribute to less antigen uptake in MoDCs through p38 MAPK pathway.

ACCEPTED MANUSCRIPT