

## Female gender and atrial fibrillation

Kornej, Jelena; Bollmann, Andreas; Hindricks, Gerhard; Lip, Gregory Y.h.

DOI:

[10.1016/j.ijcard.2015.02.062](https://doi.org/10.1016/j.ijcard.2015.02.062)

License:

Other (please specify with Rights Statement)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Kornej, J, Bollmann, A, Hindricks, G & Lip, GYH 2015, 'Female gender and atrial fibrillation: An association with worse prognosis and outcomes', *International Journal of Cardiology*, vol. 184, pp. 321-322.  
<https://doi.org/10.1016/j.ijcard.2015.02.062>

[Link to publication on Research at Birmingham portal](#)

### **Publisher Rights Statement:**

NOTICE: this is the author's version of a work that was accepted for publication in International Journal of Cardiology. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in International Journal of Cardiology, Vol 184, April 2015, DOI: 10.1016/j.ijcard.2015.02.062

Eligibility for repository checked April 2015

### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

## Accepted Manuscript


Female gender and atrial fibrillation: An association with worse prognosis and outcomes

Jelena Kornej, Andreas Bollmann, Gerhard Hindricks, Gregory Y.H. Lip

PII: S0167-5273(15)00189-8  
DOI: doi: [10.1016/j.ijcard.2015.02.062](https://doi.org/10.1016/j.ijcard.2015.02.062)  
Reference: IJCA 19714

To appear in: *International Journal of Cardiology*

Received date: 8 January 2015  
Accepted date: 21 February 2015



Please cite this article as: Kornej Jelena, Bollmann Andreas, Hindricks Gerhard, Lip Gregory Y.H., Female gender and atrial fibrillation: An association with worse prognosis and outcomes, *International Journal of Cardiology* (2015), doi: [10.1016/j.ijcard.2015.02.062](https://doi.org/10.1016/j.ijcard.2015.02.062)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Letter to the Editor

### Female gender and atrial fibrillation:

### An association with worse prognosis and outcomes

regarding article “Gender-related differences and risk of cardiovascular morbidity and all-cause mortality in patients hospitalized with incident atrial fibrillation without concomitant diseases: A nationwide cohort study of 9519 patients” by Andersson L et al, International Journal of Cardiology; doi: 10.1016/j.ijcard.2014.09.092

Jelena Kornej,<sup>1</sup> MD, Andreas Bollmann,<sup>1</sup> MD, PhD,

Gerhard Hindricks,<sup>1</sup> MD, Gregory YH Lip,<sup>2</sup> MD

<sup>1</sup>Leipzig University, Heart Center, Department of Electrophysiology, Germany

<sup>2</sup>University of Birmingham, Centre for Cardiovascular Sciences, Birmingham, UK

**Words (text):** 557

Address for correspondence:

Jelena Kornej, MD

Leipzig University

Heart Center

Department of Electrophysiology

Strümpellstr. 39

04289 Leipzig, Germany

Phone: ++49 341 865 1431

Fax: ++49 341 865 1460

Email: [jelena.kornej@gmx.de](mailto:jelena.kornej@gmx.de)

*To the Editor:*

Females with atrial fibrillation (AF) are at increased risk for ischemic stroke but have been under-represented in AF cohorts. Without anticoagulation, women are at higher risk than men for AF-related thromboembolic events (TE) but results are not consistent (1, 2). Interestingly, vascular complication rates after AF ablation had been recently shown to be also higher in females (3). Female sex has been included in the CHA<sub>2</sub>DS<sub>2</sub>-VASc score for assessing thromboembolic risk, which doubles in women between the ages of 55 and 65, the menopausal period, when estradiol levels decrease by about 60% (4). Following the menopause, a decline in endogenous estrogen receptors contributes to an up-regulated production of inflammatory cytokines, with further links to the hypercoagulable state (5).

Thus, it was with great interest that we read the article by Andersson and co-workers (6) on the gender-related differences in risk of cardiovascular morbidity and all-cause mortality in patients with lone and idiopathic AF. Using by far the largest nationwide set of patients with incident AF and no other co-morbidity at the time of diagnosis, the authors demonstrated a two-fold risk of thromboembolic events and three-fold increased risk of heart failure in patients with AF irrespective of age and gender, as compared to controls. Furthermore, the authors found that women were at higher risk of stroke and transient ischemic attack than men. The annual rate of thromboembolic events in men and women between 55 and 64 years was 1.0-1.1% and in controls, 0.4% and 0.3%, respectively. This is in accordance with other observational and registry studies that demonstrated low annual rates of thromboembolism (stroke and systemic embolism, TE) in selected populations after AF catheter ablation, reaching event rates that are comparable to patients without AF (7).

The authors should be congratulated for their interesting findings on this topical and timely performed study that are in accordance with our research (8). Assessing gender-specific differences in thromboembolic risk post-ablation and the predictive risk factors for thromboembolism separately in females and males, we found that the incidence of thromboembolic events after AF catheter ablation is low in both genders. In contrast to the current study (6), there were no differences between thromboembolic events in males and females (1.2% versus 2.0%,  $p=0.128$ ). In females, low ejection fraction (EF) and higher CHA<sub>2</sub>DS<sub>2</sub>-VASc score were independent predictors for thromboembolic risk, while renal dysfunction and all three stroke risk stratification scores, i.e. CHADS<sub>2</sub>, CHA<sub>2</sub>DS<sub>2</sub>-VASc, R<sub>2</sub>CHADS<sub>2</sub>, were associated with thromboembolic events risk in males. Because men constitute the large part of ablation cohorts, it was perhaps unsurprising to find an association between renal dysfunction and thromboembolic risk scores, as previously demonstrated (8).

Heart failure has been associated with higher stroke rates in AF (9). Whilst impaired left ventricular ejection fraction is more commonly been used as an equivalent for the heart failure, AF patients with heart failure and *normal* (preserved) EF also have greater thromboembolic risk compared with only AF patients (10). Our finding that impaired EF is significant predictor for TE in females supports previous studies; however, the novel aspect is the importance of EF for risk assessment in women after catheter ablation. Demonstrating higher risk for of thromboembolic complications and development of heart failure in females with lone/idiopathic AF, the current study by Andersson et al (6) supports our observations and emphasizes the necessity of more careful gender-specific stroke risk assessment.

## References

1. Fang MC, Singer DE, Chang Y, Hylek EM, Henault LE, Jensvold NG, Go AS. Gender differences in the risk of ischemic stroke and peripheral embolism in atrial fibrillation: the AnTicoagulation and Risk factors In Atrial fibrillation (ATRIA) study. *Circulation* 2005; 112(12):1687-1691.
2. Overvad TF, Rasmussen LH, Skjøth F, Overvad K, Albertsen IE, Lane DA, Lip GY, Larsen TB. Female sex as a risk factor for thromboembolism and death in patients with incident atrial fibrillation. The prospective Danish Diet, Cancer and Health study. *Thromb Haemost* 2014; 112(4):789-795.
3. Zhang XD, Tan HW, Gu J, Jiang WF, Zhao L, Wang YL, Liu YG, Zhou L, Gu JN, Liu X. Efficacy and safety of catheter ablation for long-standing persistent atrial fibrillation in women. *Pacing Clin Electrophysiol* 2013; 36(10):1236-1244.
4. Vaccarino V, Badimon L, Corti R, de Wit C, Dorobantu M, Hall A, Koller A, Marzilli M, Pries A, Bugiardini R. Ischaemic heart disease in women: are there sex differences in pathophysiology and risk factors? Position paper from the working group on coronary pathophysiology and microcirculation of the European Society of Cardiology. *Cardiovasc Res* 2011; 90(1):9-17.
5. Novella S, Heras M, Hermenegildo C, Dantas AP. Effects of estrogen on vascular inflammation: a matter of timing. *Arterioscler Thromb Vasc Biol* 2012; 32(8):2035-2042.
6. Andersson T, Magnuson A, Bryngelsson IL, Frobert O, Henriksson KM, Edvardsson N, Poci D. Gender-related differences in risk of cardiovascular morbidity and all-cause mortality in patients hospitalized with incident atrial fibrillation without concomitant diseases: A nationwide cohort study of 9519 patients. *Int J Cardiol* 2014; 177(1):91-99.
7. Hunter RJ, McCreedy J, Diab I, Page SP, Finlay M, Richmond L, French A, Earley MJ, Sporton S, Jones M, Joseph JP, Bashir Y, Betts TR, Thomas G, Staniforth A, Lee G, Kistler P, Rajappan K, Chow A, Schilling RJ. Maintenance of sinus rhythm with an ablation strategy in patients with atrial fibrillation is associated with a lower risk of stroke and death. *Heart* 2010; 98(1):48-53.
8. Kornej J, Hindricks G, Kosiuk J, Arya A, Sommer P, Husser D, Rolf S, Richter S, Piorkowski C, Gaspar T, Lip GY, Bollmann A. Renal dysfunction, stroke risk scores (CHADS2, CHA2DS2-VASc, and R2CHADS2), and the risk of thromboembolic events after catheter ablation of atrial fibrillation: the Leipzig Heart Center AF Ablation Registry. *Circ Arrhythm Electrophysiol* 2013; 6(5):868-874.
9. Agarwal M, Apostolakis S, Lane DA, Lip GY. The impact of heart failure and left ventricular dysfunction in predicting stroke, thromboembolism, and mortality in atrial fibrillation patients: a systematic review. *Clin Ther* 2014; 36(9):1135-1144.
10. Jang SJ, Kim MS, Park HJ, Han S, Kang DH, Song JK, Park SW, Park SJ, Kim JJ. Impact of heart failure with normal ejection fraction on the occurrence of ischaemic stroke in patients with atrial fibrillation. *Heart* 2013; 99(1):17-21.