

## Length of hospital stay is shorter in South Asian patients with ischaemic stroke

Potluri, Rahul; Wasim, Mohammed; Markandey, Bharat; Kapour, Arouna; Khouw, Niece; Carter, Paul; Uppal, Hardeep; Chandran, Suresh

DOI:

[10.1016/j.ijcard.2015.03.290](https://doi.org/10.1016/j.ijcard.2015.03.290)

License:

Other (please specify with Rights Statement)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Potluri, R, Wasim, M, Markandey, B, Kapour, A, Khouw, N, Carter, P, Uppal, H & Chandran, S 2015, 'Length of hospital stay is shorter in South Asian patients with ischaemic stroke', *International Journal of Cardiology*, vol. 187, pp. 190-191. <https://doi.org/10.1016/j.ijcard.2015.03.290>

[Link to publication on Research at Birmingham portal](#)

### **Publisher Rights Statement:**

NOTICE: this is the author's version of a work that was accepted for publication in International Journal of Cardiology. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in International Journal of Cardiology, Vol 187, May 2015, DOI: 10.1016/j.ijcard.2015.03.290

Eligibility for repository checked April 2015

### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

## Accepted Manuscript


Length of hospital stay is shorter in South Asian patients with Ischaemic Stroke

Rahul Potluri, Mohammed Wasim, Bharat Markandey, Arouna Kapour,  
Niece Khouw, Paul Carter, Hardeep Uppal, Suresh Chandran

PII: S0167-5273(15)00530-6  
DOI: doi: [10.1016/j.ijcard.2015.03.290](https://doi.org/10.1016/j.ijcard.2015.03.290)  
Reference: IJCA 20055

To appear in: *International Journal of Cardiology*

Received date: 27 January 2015  
Accepted date: 7 March 2015



Please cite this article as: Potluri Rahul, Wasim Mohammed, Markandey Bharat, Kapour Arouna, Khouw Niece, Carter Paul, Uppal Hardeep, Chandran Suresh, Length of hospital stay is shorter in South Asian patients with Ischaemic Stroke, *International Journal of Cardiology* (2015), doi: [10.1016/j.ijcard.2015.03.290](https://doi.org/10.1016/j.ijcard.2015.03.290)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Length of hospital stay is shorter in South Asian patients with Ischaemic Stroke****Short Communication to Editor – International Journal of Cardiology**

Rahul Potluri<sup>1</sup> MRCP

Mohammed Wasim<sup>2</sup> MBChB

Bharat Markandey<sup>3</sup> MBChB

Arouna Kapour<sup>3</sup> MBChB

Niece Khouw<sup>2</sup> MRCP

Paul Carter<sup>4</sup> BMedSc

Hardeep Uppal<sup>1</sup> MRCPsych

Suresh Chandran<sup>5</sup> FRCP

<sup>1</sup> ACALM Study Unit in collaboration with Aston Medical School, Aston University, Birmingham, UK

<sup>2</sup> Pennine Acute Hospitals NHS Trust, Manchester, UK

<sup>3</sup> Queen Elizabeth Hospital, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK

<sup>4</sup> The Medical School, University of Birmingham, Birmingham, UK

<sup>5</sup> Department of Acute Medicine, North Western Deanery, UK

**Key word(s):** – Ischaemic Stroke, Length of Hospital Stay, Ethnic, South Asian

**Short title:** - Ethnicity and duration of hospital stay in Ischaemic Stroke

**Conflict of interest statement:** No conflicts of interest to declare. No funding was obtained for this research.

**Correspondence to:** Dr Rahul Potluri, Honorary Clinical Lecturer in Cardiology, Aston University, Aston Triangle, Birmingham, B4 7ET, UK. Email: rahulpotluri@outlook.com. Telephone: 00447877482778. Fax: 00447877482778

Dear Editor,

Ischaemic stroke is a global healthcare issue and in the UK one percent of all admissions to National Health Service (NHS) hospitals were due to stroke<sup>1</sup>. It is the leading cause of disability, with more than half of the survivors left dependent on others, of which 11% are newly admitted to care homes. The majority of survivors will require rehabilitation and additional care in the community<sup>2,3</sup>. These additional needs culminate with longer hospital admissions, with a mean length of hospital stay (LOS) of 20 days<sup>3</sup>. These studies have predominantly been conducted on Caucasian population groups and there is a dearth of research investigating LOS amongst ethnic minority groups. In the UK, NHS hospitals serve large multi-ethnic populations and further research in this field is essential. We aimed to explore the effect of ethnicity on LOS in patients presenting with ischaemic stroke from a large catchment area in the North West of England, UK.

Using completely anonymous data, we traced adult patients admitted with ischaemic stroke across seven hospitals in the North West of England using the ACALM (Algorithm for Comorbidities, Associations, Length of stay and Mortality) study protocol between 1st January 2000 and 31st March 2013. The ACALM study used the International Classification of Disease, Version 10 (ICD-10) diagnosis codes and Office of Population Censuses and Surveys Classification of Interventions and Procedures (OPCS-4) to trace patients. Study participants admitted with a diagnosis of any type of ischaemic stroke, as defined by ICD-10 codes (including subsets), were examined by LOS, ethnicity, age, gender, co-morbidities and type of admission. The Caucasian population was used as a reference group, in comparison to other ethnic groups classified as South Asian, Afro-Caribbean, Oriental, mixed ethnicity, other ethnicity, and unknown ethnicity. The presence of co-morbidity was defined by the presence of any of the top eight causes of mortality in the UK apart from cerebrovascular disease, namely Ischaemic Heart Disease, Heart Failure, Lung Cancer, Breast Cancer, Dementia, Chronic Kidney Disease, Chronic Obstructive Airways Disease.

Data analysis was performed using SPSS Version 20.0. Variation in the LOS between ethnic groups was analysed by means of a logistic regression model adjusted for age, sex and co-morbidity with the Caucasian ethnic group as the reference. Completely anonymous patient data was used and processed in accordance with local ethical research and development policy. The methodology has been previously described and used by our group and similar methodology has been used by other groups.<sup>4-14</sup>

Within the study period, there were 929465 patients admitted. Of these, 17 415 (2.7%) were coded for ischaemic stroke using the International Classification of Disease (ICD) codes 433-434, inclusive of all subsets of these codes. Our results showed nearly equal proportion of male and female patients with ischaemic stroke amongst all groups combined (49% Male, 51% Female%). However, ethnic variations were present; Ischaemic stroke being more common amongst males in the South-Asian and Afro-Caribbean populations and more common amongst females in the Oriental and Mixed ethnic groups as shown in Table 1. 95.5% of the patients were admitted as an emergency and the average length of stay was 19.6 days. This is in keeping with the national data of mean 20 days<sup>3</sup>. Admissions from the black and ethnic minority population comprised 7.8% of total admissions with ischaemic stroke and these patients were also younger than the Caucasian population. The LOS was strikingly shorter in the South Asian ethnic group with a mean hospital stay of 12 days which compared with the Caucasian population after adjusting for age, gender and co-morbidity ( $p<0.01$ ). LOS was significantly longer in Afro-Caribbean patients. Full results are shown in table 1.

Our main findings were 1) Significantly shorter LOS in the South Asian ethnic group compared to the Caucasian group, even after modeling for differences in age, gender and co-morbidities. 2) Significantly longer LOS in Afro-Caribbean ethnic group (Table 1). Such ethnic disparities in LOS with ischaemic stroke warrant further investigation. Our data sets did not allow us to examine for any association with the presence of community based specialist discharge facilities, a factor which was previously recognised to support discharge and improve outcome for patients<sup>15</sup>.

Although the reasons for the shorter LOS are unknown, there is evidence which suggests that less support is offered to the ethnic minority populations at the time of discharge and/or that they may be discharged prematurely and consequently have a greater chance of readmission<sup>16</sup>. Similar findings were revealed in a recent study from Birmingham, UK, looking at LOS in patients with Diabetes<sup>5</sup> and three other studies by our group in the North West of England looking at length of hospital stay for patients post Myocardial Infarction, Pulmonary Embolism and Aortic Valve Disease<sup>6,8</sup>. A further two studies by our group looking at the LOS post coronary artery bypass grafting and cardiomyopathy showed that South Asians, Far-East Asians and Afro-Caribbean's respectively, have a longer length of stay<sup>7</sup>. The reason for these disparities is not clear but there is evidence that a more supportive home environment amongst the South Asian and mixed ethnicity groups allows for early discharge<sup>5,6,8</sup>. However, the possibility of an inappropriate early discharge warrants further investigation.

To assess the potential ethnic inequities in hospital care in the National Health Service, further studies with defined outcomes such as discharge support, the availability of community discharge facilities and the rate of adverse events are required. We believe the results of the study, alongside the issues highlighted will pave the way to better inpatient care with safe and appropriate discharges in centers with large multi-ethnic populations.

Despite awareness of inequities in health care quality, little is known about strategies that could improve the quality of healthcare for ethnic minority populations. We believe that the results of this study and the issues it raises may help design and improve care of inpatients with ischaemic strokes in regions with large multi-ethnic populations.

#### **Acknowledgements**

We would like to acknowledge all of those who were indirectly involved in this work

#### **Funding**

There was no formal funding for this study.

#### **Conflicts of Interest/Disclosure**

All authors declare no conflicts of interest. All authors do not have any relevant disclosures.

## References

1. Townsend N, Wickramasinghe K, Bhatnagar P, Smolina K, Nichols M, Leal J et al. Coronary heart disease statistics 2012 Edition. London, UK: British Heart Foundation. 2012:95-6
2. Adamson J, Beswick A, Ebrahim S. Stroke and Disability. *Journal of Stroke and Cerebrovascular Diseases* 2004;13:171-177
3. Stroke Working Party. National Sentinel Stroke Clinical Audit 2010 Round 7 Public report for England, Wales and Northern Ireland. London, UK: Royal College of Physicians. 2011:43.
4. Carter P, Mann J, Sangha J, Uppal H, Natalwala A, Aziz A et al. The burden of cardiovascular disease amongst psychiatric patients. *Int J Cardiol.* 2013;169:e65-6
5. Kimani KN, Potluri R, Natalwala A, Ghosh S, Heun R, Narendran P. Length of hospital stay is shorter in black and ethnic minority patients with diabetes. *Diabet Med.* 2012;29:830-1.
6. Khouw N, Wasim M, Aziz A, Uppal H, Chandran S, Potluri R. Length of hospital stay is shorter in South Asian patients with myocardial infarction. *Int J Cardiol.* 2014;171:e54-5.
7. Ciputra R, Sembiring Y, Prawato O, Khouw N, Baig M, Uppal H et al. Length of stay in hospital is longer in ethnic minority patients after coronary artery bypass surgery. *Int J Cardiol.* 2014 Apr 1;172(3):e434-5.
8. Smith S, Gollop N, Uppal H, Chandran S, Potluri R. Length of hospital stay is shorter in South Asian patients with acute pulmonary embolism. *Heart Asia.* 2014;6;1-2.
9. Carter P, Rai G, Aziz A, Mann J, Chandran S, Uppal H, Potluri R. Trends of cardiovascular disease amongst psychiatric patients between 2001 and 2012 in Greater Manchester, UK. *Int J Cardiol.* 2014 May 15;173(3):573-4.

10. Teo R, Gollop ND, Baig M, Uppal H, Chandran S, Potluri R. The burden and trends of psychiatric co-morbidities amongst patients with cardiomyopathy. *Int J Cardiol.* 2014 Jun 15;174(2):398-9.
11. Lavu D, Uppal H, Katragadda A, Ahmed A, Potluri R. Effect of ethnicity on live birth rates after in vitro fertilisation or intracytoplasmic sperm injection treatment: possible explanations and further observations. *BJOG.* 2014 Nov;121(12):1576-7.
12. Potluri R, Baig M, Mavi JS, Ali N, Aziz A, Uppal H, Chandran S. The role of angioplasty in patients with acute coronary syndrome and previous coronary artery bypass grafting. *Int J Cardiol.* 2014 Oct 20;176(3):760-3.
13. Thakker C, Gollop ND, Carter P, Lavu D, Uppal H, Chandran S, Potluri R. Prevalence and length of hospital stay in patients with aortic valve disease is lower amongst South Asians. *Int J Cardiol.* 2014 Nov 15;177(1):34-6.
14. Green B, Gollop ND, Baig M, Uppal H, Chandran S, Potluri R. Prolonged length of hospital stay in Far-East Asian and Afro-Caribbean patients with cardiomyopathy. *Int J Cardiol.* 2014 Sep;176(1):230-1.
15. Department of Health National Audit Office. Progress in improving stroke care. London, UK: The Stationary Office. 2010:8
16. 'You Can Take Him Home Now': Carer's Experience of Hospital Discharge. Carers National Association website. <http://www.carersuk.org/professionals/resources/research-library/item/517-you-can-take-him-home-now-carers-experiences-of-hospitaldischarge>. Accessed 8 Jan 2014



**Table 1 - Characteristics of admissions for patients with Ischaemic Stroke**

<b>Ethnicity</b>	<b>n(%)</b>	<b>Mean Age (yrs)</b>	<b>F:M Ratio</b>	<b>%admitted as emergency</b>	<b>%with co-morbidity</b>	<b>Mean LOS (days)</b>	<b>ODDs ratio for length of stay*</b>
<b>All Groups</b>	17415	73.8	1:1	95.5	57.2	19.6	-
<b>Caucasian</b>	14586 (83.8)	74.2	1:1	95.7	58.5	19.3	1
<b>South Asian</b>	764 (4.4)	65.1	0.8:1	92.8	53.8	12.8	0.993 (0.989– 0.996) #
<b>Afro-Caribbean</b>	275 (1.6)	66.9	0.7:1	83.6	46.2	25.6	1.003 (1.001-1.005) #
<b>Oriental</b>	59 (0.3)	73.5	1.3:1	88.1	37.3	24.1	1.002 (0.998-1.007)
<b>Mixed</b>	31(0.2)	58.6	1.2:1	87.1	54.8	12.3	0.993 (0.978-1.009)
<b>Other</b>	234(1.3)	70.9	1:1	94.9	44.0	26.9	1.004 (1.001-1.006) #
<b>Unknown</b>	1466 (8.4)	75.7	1.1:1	97.5	51.2	24.5	1.003 (1.001-1.006) #

LOS = Length of hospital stay \*adjusted for age, sex, co-morbidity, # statistically significant, p<0.05