

A well-preserved 'placoderm' (stem-group Gnathostomata) upper jaw from the Early Devonian of Mongolia clarifies jaw evolution

Brazeau, Martin D; Yuan, Haobo; Giles, Sam; Jerve, Anna L; Zorig, E; Ariunchimeg, Ya; Sansom, Robert S; Atwood, Robert C

DOI:
[10.1098/rsos.221452](https://doi.org/10.1098/rsos.221452)

License:
Creative Commons: Attribution (CC BY)

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (Harvard):
Brazeau, MD, Yuan, H, Giles, S, Jerve, AL, Zorig, E, Ariunchimeg, Y, Sansom, RS & Atwood, RC 2023, 'A well-preserved 'placoderm' (stem-group Gnathostomata) upper jaw from the Early Devonian of Mongolia clarifies jaw evolution', *Royal Society Open Science*, vol. 10, no. 2, 221452. <https://doi.org/10.1098/rsos.221452>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Research



Cite this article: Brazeau MD, Yuan H, Giles S, Jerve AL, Zorig E, Ariunchimeg Y, Sansom RS, Atwood RC. 2023 A well-preserved ‘placoderm’ (stem-group Gnathostomata) upper jaw from the Early Devonian of Mongolia clarifies jaw evolution. *R. Soc. Open Sci.* **10**: 221452. <https://doi.org/10.1098/rsos.221452>

Received: 9 November 2022

Accepted: 27 January 2023

Subject Category:

Organismal and Evolutionary Biology

Subject Areas:

palaeontology

Keywords:

acanthothoracid, placoderm, Devonian, jaws, Mongolia, evolution

Author for correspondence:

Martin D. Brazeau

e-mail: m.brazeau@imperial.ac.uk

[†]Present address: Department of Zoology, University of Cambridge, Downing Street, Cambridge CB2 3EJ, UK.

[‡]Present address: Department of Zoology, Naturhistoriska Riksmuseet, Frescativägen 40, 114 18 Stockholm, Sweden.

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.6414027>.

A well-preserved ‘placoderm’ (stem-group Gnathostomata) upper jaw from the Early Devonian of Mongolia clarifies jaw evolution

Martin D. Brazeau^{1,2}, Haobo Yuan^{1,†}, Sam Giles^{2,3}, Anna L. Jerve^{1,‡}, E. Zorig⁴, Ya. Ariunchimeg⁵, Robert S. Sansom⁶ and Robert C. Atwood⁷

¹Department of Life Sciences, Imperial College London, Silwood Park Campus, Ascot SL5 7PY, UK

²The Natural History Museum, London SW7 5BD, UK


³School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham B15 2TT, UK

⁴Institute of Paleontology, Mongolian Academy of Sciences, Ulaanbaatar 15160, Mongolia

⁵The Natural History Museum, Ulaanbaatar 1420, Mongolia

⁶Department of Earth and Environmental Sciences, University of Manchester, Manchester M13 9PT, UK

⁷Diamond Light Source, Didcot OX11 0DE, UK

 MDB, 0000-0002-0650-1282; SG, 0000-0001-9267-4392; RSS, 0000-0003-1926-2556

The origin of jaws and teeth remains contentious in vertebrate evolution. ‘Placoderms’ (Silurian-Devonian armoured jawed fishes) are central to debates on the origins of these anatomical structures. ‘Acanthothoracids’ are generally considered the most primitive ‘placoderms’. However, they are so far known mainly from disarticulated skeletal elements that are typically incomplete. The structure of the jaws—particularly the jaw hinge—is poorly known, leaving open questions about their jaw function and comparison with other placoderms and modern gnathostomes. Here we describe a near-complete ‘acanthothoracid’ upper jaw, allowing us to reconstruct the likely orientation and angle of the bite and compare its morphology with that of other known ‘placoderm’ groups. We clarify that the bite position is located on the upper jaw cartilage rather than on the dermal cheek and thus show that there is a highly conserved bite morphology among most groups of ‘placoderms’, regardless of their overall cranial geometry. Incorporation of the dermal skeleton appears to provide a sound

biomechanical basis for jaw origins. It appears that ‘acanthothoracid’ dentitions were fundamentally similar in location to that of arthrodire ‘placoderms’, rather than resembling bony fishes. Irrespective of current phylogenetic uncertainty, the new data here resolve the likely general condition for ‘placoderms’ as a whole, and as such, ancestral morphology of known jawed vertebrates.

1. Introduction

‘Placoderms’ are extinct fishes that ranged from the Silurian to the end of the Devonian period (444 to 359 Ma). Although their monophyly is highly debated [1–5], it is widely agreed that they are the only known jaw-bearing stem-group gnathostomes. They are thus key to reconstructing character transitions leading to modern (crown-group) gnathostomes. ‘Placoderms’ occupy a central role in debates on the origin of both jaws and teeth—major gnathostome innovations that led to diverse feeding ecologies. Recent discoveries from the Silurian and Early Devonian of south China [4,6,7] and Czechia [8] have delivered significant advances that diminish the morphological gap between ‘placoderm’ and crown-group vertebrate jaw morphology. There is now a growing consensus that the last common ancestor of crown gnathostomes possessed dermal jaw bones, a trait potentially extending to the earliest appearance of jaws [9]. This has transformed perspectives of both jaw and tooth evolution, which have long been modelled on shark-like conditions [10].

Owing to a lack of fossilized mandibular arch material in jawless fishes, the primitive structure of the gnathostome jaw has been difficult to reconstruct using palaeontological evidence. ‘Placoderms’ exhibit diverse jaw morphologies and presumed feeding ecologies [11–13]. These range from diverse biting modes, suspension feeding and possible grazing in arthrodires; durophagy in ptyctodonts; ambush predation in rhenanids; and benthic (detritus?) feeding in antiarchs. Most ‘placoderm’ dentitions were relatively simple: tubercles added in rows (or files) or centripetally to growing jaw plates [14,15]. Despite this simplicity, ‘placoderm’ dentitions are diverse, and detailed anatomical investigations in recent decades have highlighted similarities between some ‘placoderm’ groups (such as arthrodires) and crown-group gnathostomes [14,16]. More recently, it has been proposed [8] that some ‘acanthothoracids’ bore marginal teeth aligned in rows along the jaw edge, as in most crown-group gnathostomes. Understanding the relationship of ‘placoderm’ dentitions as well as overall jaw morphology and functional diversity are thus key to understanding the origin of modern jaws and teeth.

Here we describe a nearly complete, fully three-dimensional upper jaw (palatoquadrate and suborbital plate) of an ‘acanthothoracid’ from the Early Devonian (Pragian) of Mongolia. The specimen reveals the most complete quadrate (articular condyle connecting the lower jaw) of any ‘acanthothoracid’ known to date. This allows us to reconstruct and compare the orientation and angle of the bite in these ‘placoderms’, and we show that a highly similar jaw articulation and bite occurs between three groups of ‘placoderms’ with disparate presumed ecomorphologies: ‘acanthothoracids’, rhenanids and arthrodires, suggesting conservation of a shared ancestral morphology among these groups. Furthermore, we clarify the evolution of gnathostome dentitions by revising the interpretation of unusual dental morphology in a previously described ‘acanthothoracid’.

2. Material and methods

2.1. Specimens

The upper jaw described here occurs within a bedrock sample MPC-Fh200/10.4 from the Yamaat Gol locality in western Mongolia [17–19]. The absence of endochondral bone and the tight spacing of the tubercles indicate that this palatoquadrate does not belong to *Minjinia*, the only ‘placoderm’ taxon so far named from that locality. The Yamaat Gol fauna contains at least three distinct ‘acanthothoracid’ taxa, but all are based on isolated fossils, making taxonomic assignments fraught. This paper makes no further taxonomic attribution for the specimen as it lacks other diagnostic traits.

2.2. Synchrotron tomography

We performed synchrotron X-ray micro-computed tomography at the I12 beamline of the Diamond Light Source, United Kingdom [20]. The X-ray beam was set to a monochromatic energy of 90 keV (double bent Laue Si 111 monochromator). The regions of interest were scanned in I12 Experimental Hutch One using

the beamline's modular imaging system. This detector consists of a PCO.edge 5.5 sCMOS camera and four user-selectable optical modules, each comprising a scintillator, 90-degree turning mirrors and a visible light lens. We used module 2 with a magnification of 0.820, corresponding to a recorded pixel size of 7.91 μm . Each acquisition consisted of 1800 projections, of 15 ms exposure time each, over a 180° rotation of the sample. Additionally, 50 flatfield images (sample out of the beam) were recorded before and after the series of acquisition as well as 10 dark images (X-ray beam off to record the noise of the camera).

The tomographs were reconstructed using the SAVU tomographic processing software [21,22] developed at Diamond Light Source. In the reconstruction process, ring artefact removal [23] and auto-centring [24] were applied, as well as distortion correction [25]. The low-pass filter approach [26] was applied. Filtered back-projection reconstructions were performed using the Astra library [27,28], and the whole process was applied on an HPC cluster system using the SAVU tomography pipeline.

2.3. Data segmentation and visualization

We segmented the data using Materialise Mimics version 23.0 (<https://www.materialise.com/en/healthcare/mimics-innovation-suite/mimics>; Materialise, Leuven, Belgium). Three-dimensional models were exported and rendered in Blender v.3.2.2 (Blender Foundation; <https://www.blender.org>).

2.4. Phylogenetic analysis

We reproduced the analysis of Vaškaninová *et al.* [8] with modified scorings for *Radotina* to reflect uncertainties and interpretations presented here. A full list of score changes is provided in electronic supplementary material, table S1. We restricted our data re-codings to eight characters relating to the identification of teeth and sensory canals in *Radotina*. We analysed the matrix using TNT (v. 1.5) [29] using a 'new technology search' (xmult) set to level 10 including the parsimony ratchet with 50 replicates. We then applied branch-swapping (bbreak) to the trees in memory. To analyse the distribution of character states, we generated a fully resolved topology removing 'wildcard' taxa by generating an agreement subtree in PAUP* (v. 4.0 alpha test version build 168) [30]. We visually explored ancestral states conditions for jaw characters using parsimony with the 'Trace Character History' function in Mesquite (v. 3.70) [31].

3. Results

3.1. Description

MPC-Fh200/10.4 is a nearly complete upper jaw unit consisting of the suborbital plate and palatoquadrate ossification (figure 1). The palatoquadrate is preserved as a thin shell of perichondral bone; endochondral bone is absent. The specimen is mostly undeformed, retaining its original three-dimensional shape. The suborbital plate is covered in densely placed stellate tubercles. In lateral view, the suborbital plate has a geometry similar to *Radotina* (figure 2b): the outline is roughly trapezoidal; the posterior margin is deep (most of the depth of the plate) and slightly convex. There is no evidence of an orbital process on the dorsal surface, but we cannot rule out the possibility that it was worn or broken off. As in *Radotina*, the ventral margin of the suborbital plate is gently convex. This differs from *Romundina*, where the ventral profile has a marked angle between the autopalatine (anterior) portion and the adductor fossa area, contributing to a short posterior margin [32,33]. The most prominent feature of the external face of the suborbital is the infraorbital canal, which follows a course from the posterodorsal corner to the anterior margin of the plate. The upper margin of the plate forms a pronounced overhang of the infraorbital sensory canal. As in *Radotina*, the dorsal tubercles curl over the dorsal margin of this canal. The anterior end of the infraorbital canal curves downward to meet the supra-oral sensory canal. The latter traces the preserved ventral margin of the suborbital. This canal terminates between the autopalatine process and the quadrate.

In medial view, the palatoquadrate fills the perimeter of the suborbital plate. The surface of the perichondral bone in the virtual models has a nodular texture. However, these appear to be ferrous crystals formed on the surface. The palatoquadrate has the classic 'placoderm' omega shape, forming a thick arch around an almost circular adductor muscle chamber. The geometry and proportions of the palatoquadrate and adductor fossa again closely resemble *Radotina* [8]. The adductor fossa




Figure 1. MPC-Fh200/10.4 upper jaw complex in virtual three-dimensional rendering from synchrotron tomography. (a) Lateral view; (b) lateral view with in-fill of lateral line canals (red); (c) lateral view with suborbital plate rendered semi-transparent to reveal internal cavities and canals; (d) mesial view; (e) ventral (labial) view; (f) dorsal view; (g) posterior view. Blue-grey indicates suborbital (dermal) plate; mauve indicates perichondral sheath of palatoquadrate. Exact boundaries between these units are approximate. Dashed leader line indicates uncertain attribution.




Figure 2. (a) Lateral line canal pattern on the cheek of 'acanthothoracid' placoderms showing the position of the canals relative to the palatoquadrate (dotted line). (b) Drawing based on data from [8]; (c) interpretive drawing of specimen from [32].

occupies about half of the internal face. The autopalatine is commensurately short and turns medially at its anterior end. The ossification is well preserved, showing that it ended anteriorly with a single, open articulation facet as in rhenanid 'placoderms' [34]. On the dorsal side of the autopalatine is a low, blunt process, which we interpret as an anterodorsal articulation (figure 1*d*).

Exceptionally among 'acanthothoracid' fossils, the quadrate area is almost entirely preserved (figure 1*d–f*). For the first time, it allows a complete understanding of the jaw articulation in these

'placoderms'. The quadrate is mediolaterally broad and possibly double-headed as in osteichthyans. Slightly dorsal and anterior to the quadrate, a large-diameter canal pierces the element between the dermal bone and the palatoquadrate. This canal corresponds to the mandibularis canal (nerve V) described in *Holonema* [35]; a similar canal is seen but undescribed in *Radotina* [8]. Further anterior along the dorsal side of the palatoquadrate, the top of the palatoquadrate is pierced by an oblique, anteriorly directed canal similar to one seen in *Radotina* (figure 1f, 'mandibularis nerve canal (anterior branch)'). These canals branch into a multitude of ramules in the autopalatine cartilage (figure 1c).

The internal structure of the bone is relatively well preserved. Three rami of the mandibularis branch are preserved. These extend to the labial margin of the suborbital plate. Two extend in a nearly parallel course from the trunk canal in the posterior quarter of the plate all the way to the antero-labial corner, where fine ramules underly the junction of the supra-oral and infraorbital sensory canal. The suborbital plate of MPC-Fh200/10.4 carries short, buried dorsal and ventral cavities that parallel the sensory canals. The chambers do not have any apparent opening to the outside. We are not aware of any such structures in any other 'placoderms', but suggest they are homologues of the 'cutaneous sensory openings' seen in various 'placoderms' [36]. Such canals have not been observed in the suborbital plates of any 'acanthothoracids' so far, but no similarly complete examples have been studied by computed tomography scanning. Nevertheless, a cutaneous sensory opening extends parallel to the infraorbital canal in *Romundina* immediately posterior to the orbital margin (see Ørvig [32]; Dupret *et al.*, [37], termed 'sensory pits' by the latter). Examination of the tomograms by Dupret *et al.* confirms that the sensory pits are connected to a tubular cavity.

3.2. Phylogenetic analysis

The phylogenetic analysis resulted in 16 800 trees with a score of 947 steps (see electronic supplementary material, information). The strict consensus matches the topology recovered by Vaškaninová *et al.*, placing the 'acanthothoracids' as an unresolved assemblage joining at the base of mandibulate stem-group gnathostomes alongside antiarchs. However, the agreement subtree shows that the lack of resolution in this part of the tree was attributable to the highly incomplete, unnamed acanthothoracid snout from Drake Bay, referred to only by its specimen number CPW.9, and *Brindabellaspis*. Neither of these taxa is represented by mandibular arch material. The removal of these two taxa reveals an acanthothoracid clade immediately crownward of antiarchs. To view synapomorphies, we used the `describetrees` function in PAUP*. This clade is supported by two unambiguous synapomorphies according to this dataset (character numbers derive from the dataset of Qiao *et al.* [38]): character 97: the presence of a ventral notch between the parachordals (consistency index: 0.125); and character 103: the absence of complete dermal encirclement of the pectoral fin base (consistency index: 0.167).

4. Discussion

4.1. Comparative anatomy and function of upper jaws in 'placoderms'

MPC-Fh200/10.4 allows us to accurately reconstruct the axis of jaw closing and the position of the bite in an 'acanthothoracid'. To date, this has generally been inferred from fragmented and incompletely preserved fossils. Our material indicates that the lower jaw closed against the internal surface of the autopalatine, likely against a ridge that traverses this face (figure 3). This contrasts with the recent reconstruction of *Radotina*, which implied either that the bite was positioned on the labial margin of the suborbital plate or that an overbite of the submarginal was involved in prey capture or processing. Such a bite, where the suborbital carries the upper dentition, is anomalous compared with disparate other 'placoderms' such as rhenanids and arthrodires. In those taxa, the bite closes against the inner surface of the autopalatine and not the suborbital plate [34,39,41] (figure 3), regardless of their dramatically different craniofacial geometries (figure 4). These taxa are known from articulated specimens, or specimens in which the dental plates are intact. However, an alternative geometry is seen in *Bothriolepis*, where articulated specimens suggest the suborbital plate may have been involved in biting [40] superficially resembling the implied condition in *Radotina* [8]. Assessing the position of the bite in 'acanthothoracids' could help address which of these two 'bite models' is more appropriate.

The well-preserved quadrate of MPC-Fh200/10.4 allows us to reconstruct the orientation of the jaw hinge and mandible (electronic supplementary material, figure S2). As the quadrate is mediolaterally wide and nearly co-planar with the inner surface of the autopalatine, the lower jaw almost certainly




Figure 3. Upper jaws of placoderm fishes in internal view. (a) *Jagorina* (based on Museum für Naturkunde, Berlin, specimen MB.f.510.5); (b) *Buchanosteid* based on data published in [39]; (c) *Radotina* based on surface models from [8]; (d) MPC-Fh200/10.4 (original); (e) *Bothriolepis* (after [40]). Arrow indicates anterior. Horizontally hatched areas indicate articulation facet. Dashed label lines indicate uncertainty concerning facet as part of quadrate or possible hyoid attachment. Dashed leader line indicates uncertain attribution.

closed against the latter surface. This is supported by taking a line perpendicular to the quadrate hinge axis, corresponding to a hypothetical lower jaw which intersects the centre of the adductor chamber and traverses the autopalatine surface. Thus, we consider the bite in ‘acanthothoracids’ inclusive of *Radotina* to have been located on the autopalatine, not the suborbital. This agrees with the general morphological similarities in overall structure between MPC-Fh200/10.4, *Radotina*, *Jagorina* and arthrodires (figures 3 and 4). The role of the submarginal plate in biting, ‘cleaver-shaped’ palatoquadrate, and small adductor fossa of *Bothriolepis* represent a clear anatomical outlier (figures 3 and 4) and has been attributed recently to ‘scraping’ feeding [42].

4.2. Implications for ‘placoderm’ dentitions

The new data from MPC-Fh200/10.4 show that the aligned tubercles interpreted as teeth in *Radotina* are not associated with the oral margin. The evidence for this is twofold: firstly, as noted above, these tubercles lie outside of a functional bite (figures 2–4). Second, we interpret the groove that these tubercles lie in as being for the supra-oral sensory line canal rather than the oral margin (figures 2 and 3). This feature was described by Vaškaninová *et al.* as a ‘trough with vascular foramina and grooves’ and interpreted as the oral margin. MPC-Fh200/10.4 shows unambiguously that, instead, this is a sensory line canal connected to the infraorbital canal and facial innervation network. This is topologically identical to the same canal in *Romundina* (figure 2). The tubercles previously identified as teeth in *Radotina* lie dorsal to this supra-oral sensory line canal, and we propose that they are petal-shaped dermal tubercles. Petal-shaped tubercles with identical external morphology occur along the ventral margin of the infraorbital sensory canal in *Radotina* (see fig. S6 of the supplementary material of [8]), as well as elsewhere on the cheek. The comparative and functional anatomy above implies that the structures interpreted as teeth in *Radotina* could not have had a primary dental function and therefore have no direct bearing on the condition of teeth and jaw evolution in early gnathostomes.

4.3. Phylogenetic and evolutionary implications

A palatoquadrate fused to a dermal suborbital plate is a general condition of ‘placoderms’, as shown by numerous phylogenetically disparate examples (e.g. figures 3 and 4). Thus, this is very probably a primitive condition for jawed vertebrates. This is supported by our phylogenetic analysis, with this character (character 23) mapping to the ancestor of all jaw-bearing taxa. Although *Radotina* does not provide evidence of osteichthyan-like sutured dental plates in the mouths of ‘acanthothoracids’, Vaškaninová *et al.* convincingly demonstrate that tooth-bearing jaw bones were also more phylogenetically widespread in placoderms than previously understood. Indeed, our revised phylogenetic analysis supports an ancestral state of dermal bones borne on the jaws in jawed vertebrates (character 43). Coupled with the discoveries of Silurian ‘placoderms’ from China with osteichthyan-like jaw bones, there is an ever-growing case that heavily armoured jaws have an early phylogenetic origin. Although Vaškaninová *et al.* homologized the gnathal elements of placoderms with the dermal jaw bones of crown-group gnathostomes (e.g. premaxilla, maxilla and dentary), they did not re-code other placoderms for this new homology scheme. Nevertheless, this proposal was tested and further corroborated recently using a ‘dynamic homology’ approach [47]. If this is




Figure 4. Cheek morphology and bite pattern in placoderm fishes with suborbital plates. (a–d) Placoderms displaying pronounced overbite with lower jaw contact on internal surface of upper jaw; (e) highly ventralized jaw mechanics of antiarchs for inferred ‘scraping’ feeding [42]. Illustration in (a) is original composited from [39,43,44]. Illustrations in (b) and (c) adapted from [45]; (e) adapted from [42,46]. Red lines indicate paths of the sensory line canals.

extrapolated to the origin of jaws themselves, it raises important evolutionary implications. Dermal-associated jaws provide a functionally informative model for the origins of jaws themselves. Up to now, debates on the origin of jaws have strongly emphasized the endoskeleton and questions of whether the palatoquadrate and hyoid arch represent derived gill arches of a branchiomic ancestor. The absence of mineralized mandibular arches in fossil jawless fishes makes this scenario difficult to evaluate. This fact points to an overlooked problem: unmineralized endoskeletal cartilages are flexible and slightly elastic. Without support from the dermal skeleton, they would provide a structurally weak basis for the origin of jaws. We suggest that the phylogenetically deep association between exoskeletal and endoskeletal jaw presents a functionally plausible model.

5. Conclusion

The hypothesis that the ancestors of all jawed vertebrates possessed dermal jaw bones integrated with other facial jaw bones [9] is an important advance in our understanding of the origin of jaws. Dermal bone provides greater mechanical reinforcement than perichondrally ossified cartilage; it is therefore plausible that dermal bones provided a structural basis for jaws. As a purportedly primitive ‘placoderm’ assemblage, ‘acanthothoracids’ reveal jaw conditions with an important bearing on this hypothesis. Here we have shown that the morphology and function of ‘acanthoracid’ jaws resemble generalized ‘placoderm’ conditions seen also in arthrodires and rhenanids, consistent with past hypotheses [34]. Given the remote relationships between these assemblages, this indicates that jaw morphology was phylogenetically conserved across most ‘placoderms’. At present, there remains no evidence of an osteichthyan-like integration of tooth-bearing bones and dermal cheek plates deep in ‘placoderm’ evolution. A better understanding of jaw morphology in other ‘placoderms’, in conjunction with more stable phylogenetic hypotheses [5,6,19,38,48], will help to clarify these conditions further.

Data accessibility. The complete tomographic image data used in this analysis along with three-dimensional surface models can be accessed at: <https://doi.org/10.6084/m9.figshare.20581665>.

The data are provided in the electronic supplementary material [49].

Authors’ contributions. M.D.B.: conceptualization, data curation, formal analysis, funding acquisition, investigation, supervision, visualization, writing—original draft and writing—review and editing; H.Y.: visualization; S.G.: methodology, validation, visualization and writing—original draft; A.L.J.: funding acquisition and investigation; E.Z.: project administration and resources; Y.A.: project administration and resources; R.S.S.: funding acquisition and investigation; R.C.A.: methodology and visualization.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein. Conflict of interest declaration. The authors declare no competing interests.

Funding. The field expedition that recovered this fossil in 2012 was supported by the National Geographic Society's Global Exploration Fund (GEFNE35-12) to M.D.B. A portion of this work was supported by the European Research Council (ERC) under the European Union's Seventh Framework Programme (FP/20072013)/ERC Grant Agreement number 311092 to M.D.B. S.G. was supported by a Royal Society Dorothy Hodgkin Research Fellowship (grant no. DH160098). A.L.J. was supported in the field by the Anna Maria Lundin's stipend from Smålands Nation, Uppsala University. Y.H. was supported by an Undergraduate Research Opportunities Programme (UROP) bursary from Imperial College London. R.S.S. was supported by a Royal Society Research Grant and the University of Manchester. Synchrotron scanning was performed at Diamond Light Source, UK, supported by visit NT29710-1 in August 2021.

Acknowledgements. We thank P. Ahlberg, V. Vaškaninová and Z. Johanson for discussions regarding the dentition of *Radotina*, though they do not necessarily agree with our interpretations. TNT was provided free of charge by the Willi Hennig Society. We thank two anonymous reviewers for comments that helped improve the manuscript.

References

- Brazeau MD. 2009 The braincase and jaws of a Devonian 'acanthodian' and modern gnathostome origins. *Nature* **457**, 305–308. (doi:10.1038/nature07436)
- Young GC. 2010 Placoderms (armored fish): dominant vertebrates of the Devonian Period. *Annu. Rev. Earth Planet. Sci.* **38**, 523–550. (doi:10.1146/annurev-earth-040809-152507)
- Dupret V, Sanchez S, Goujet D, Tafforeau P, Ahlberg PE. 2014 A primitive placoderm sheds light on the origin of the jawed vertebrate face. *Nature* **507**, 500–503. (doi:10.1038/nature12980)
- Zhu M *et al.* 2013 A Silurian placoderm with osteichthyan-like marginal jaw bones. *Nature* **502**, 188–193. (doi:10.1038/nature12617)
- King B, Qiao T, Lee MSY, Zhu M, Long JA. 2016 Bayesian morphological dock methods resurrect placoderm monophyly and reveal rapid early evolution in jawed vertebrates. *Syst. Biol.* **66**, 499–516. (doi:10.1093/sysbio/syw107)
- Zhu M, Ahlberg PE, Pan Z, Zhu Y, Qiao T, Zhao W, Jia L, Lu J. 2016 A Silurian maxillate placoderm illuminates jaw evolution. *Science* **354**, 334–336. (doi:10.1126/science.aah3764)
- Li Q, Zhu Y, Lu J, Chen Y, Wang J, Peng L, Wei G, Zhu M. 2021 A new Silurian fish close to the common ancestor of modern gnathostomes. *Curr. Biol.* **31**, 3613–3620.e2. (doi:10.1016/j.cub.2021.05.053)
- Vaškaninová V, Chen D, Tafforeau P, Johanson Z, Ekrt B, Blom H, Ahlberg PE. 2020 Marginal dentition and multiple dermal jawbones as the ancestral condition of jawed vertebrates. *Science* **369**, 211–216. (doi:10.1126/science.aaz9431)
- Zhu Y-a, Ahlberg PE, Zhu M. 2019 The evolution of vertebrate dermal jaw bones in the light of maxillate placoderms. In *Evolution and development of fishes* (eds Z Johanson, C Underwood, M Richter), pp. 71–86. Cambridge, UK: Cambridge University Press.
- Reif WE. 1982 Evolution of dermal skeleton and dentition in vertebrates. In *Evolutionary biology* (eds M Hecht, B Wallace, G Prance), pp. 287–368. Boston, MA: Springer US.
- Miles RS. 1969 Features of placoderm diversification and the evolution of the arthrodire feeding mechanism. *Trans. R. Soc. Edinburgh* **68**, 123–170. (doi:10.1017/S0080456800014629)
- Anderson PSL, Friedman M, Brazeau MD, Rayfield EJ. 2011 Initial radiation of jaws demonstrated stability despite faunal and environmental change. *Nature* **476**, 206–209. (doi:10.1038/nature10207)
- Deakin WJ, Anderson PSL, Boer W den, Smith TJ, Hill JJ, Rüdcklin M, Donoghue PCJ, Rayfield EJ. 2022 Increasing morphological disparity and decreasing optimality for jaw speed and strength during the radiation of jawed vertebrates. *Sci. Adv.* **8**, eabl3644. (doi:10.1126/sciadv.abl3644)
- Rüdcklin M, Donoghue PCJ, Johanson Z, Trinajstić K, Marone F, Stamparoni M. 2012 Development of teeth and jaws in the earliest jawed vertebrates. *Nature* **491**, 748–751. (doi:10.1038/nature11555)
- Hu YZ, Young GC, Burrow C, Zhu Y, Lu J. 2019 High resolution XCT scanning reveals complex morphology of gnathal elements in an Early Devonian arthrodire. *Palaeoworld* **28**, 525–534. (doi:10.1016/j.palwor.2018.12.003)
- Smith M, Johanson Z. 2003 Separate evolutionary origins of teeth from evidence in fossil jawed vertebrates. *Science* **299**, 1235–1236. (doi:10.1126/science.1079623)
- Alekseeva RE, Mendbayar B, Erlanger OA. 1981 *Brachiopods and biostratigraphy of the lower devonian of Mongolia*. Moscow, Russia: Nauka.
- Alekseeva RE. 1993 *Devonian biostratigraphy of Mongolia*. Moscow, Russia: Nauka.
- Brazeau MD, Giles S, Dearden RP, Jerve A, Ariunchimeg Y, Zorig E, Sansom R, Guillaume T, Castiello M. 2020 Endochondral bone in an Early Devonian 'placoderm' from Mongolia. *Nat. Ecol. Evol.* **4**, 1477–1484. (doi:10.1038/s41559-020-01290-2)
- Drakopoulos M *et al.* 2015 112: the Joint Engineering, Environment and Processing (JEEP) beamline at Diamond Light Source. *J. Synchrotron Radiat.* **22**, 828–838. (doi:10.1107/S1600577515003513)
- Atwood RC, Bodey AJ, Price SWT, Basham M, Drakopoulos M. 2015 A high-throughput system for high-quality tomographic reconstruction of large datasets at Diamond Light Source. *Phil. Trans. R. Soc. A* **373**, 20140398. (doi:10.1098/rsta.2014.0398)
- Wadson N, Basham M. 2016 Savu: a Python-based, MPI framework for simultaneous processing of multiple, N-dimensional, large tomography datasets. (<https://arxiv.org/abs/1610.08015>)
- Vo NT, Atwood RC, Drakopoulos M. 2018 Superior techniques for eliminating ring artifacts in X-ray micro-tomography. *Opt. Express* **26**, 28 396–28 412. (doi:10.1364/OE.26.028396)
- Vo NT, Drakopoulos M, Atwood RC, Reinhard C. 2014 Reliable method for calculating the center of rotation in parallel-beam tomography. *Opt. Express* **22**, 19 078–19 086. (doi:10.1364/OE.22.019078)
- Vo NT, Atwood RC, Drakopoulos M. 2015 Radial lens distortion correction with sub-pixel accuracy for X-ray micro-tomography. *Opt. Express* **23**, 32 859–32 868. (doi:10.1364/OE.23.032859)
- Paganin D, Mayo SC, Gureyev TE, Miller PR, Wilkins SW. 2002 Simultaneous phase and amplitude extraction from a single defocused image of a homogeneous object. *J. Microsc.* **206**, 33–40. (doi:10.1046/j.1365-2818.2002.01010.x)
- Ramachandran GN, Lakshminarayanan AV. 1971 Three-dimensional reconstruction from radiographs and electron micrographs: application of convolutions instead of Fourier transforms. *Proc. Natl Acad. Sci. USA* **68**, 2236–2240. (doi:10.1073/pnas.68.9.2236)
- Aarle W van, Palenstijn WJ, Cant J, Janssens E, Bleichrodt F, Dabrovolski A, Beenhouwer JD, Batenburg KJ, Sijbers J. 2016 Fast and flexible X-ray tomography using the ASTRA toolbox. *Opt. Express* **24**, 25 129–25 147. (doi:10.1364/OE.24.025129)
- Goloboff PA, Catalano SA. 2016 TNT version 1.5, including a full implementation of phylogenetic morphometrics. *Cladistics* **32**, 221–238. (doi:10.1111/cla.12160)
- Swofford DL. 2019 PAUP* version 4.0a (build 168). See <http://phylosolutions.com/paup-test/>.
- Maddison DR, Maddison WP. 2021 Mesquite: a modular system for evolutionary analysis.

Version 3.70. See <http://www.mesquiteproject.org>.

32. Ørvig T. 1975 Description, with special reference to the dermal skeleton, of a new radotiid arthrodire from the Geddinnian of Arctic Canada. *Colloque International CNRS* **218**, 43–71.
33. Burrow CJ. 2006 Placoderm fauna from the Connemarra Formation (?late Lochkovian, Early Devonian), central New South Wales. *Alcheringa: Australasian J. Palaeontol.* **30**, 59–88. (doi:10.1080/03115510609506856)
34. Young G. 1986 The relationships of placoderm fishes. *Zool. J. Linn. Soc.* **88**, 1–57. (doi:10.1111/j.1096-3642.1986.tb00876.x)
35. Miles R. 1971 The Holonematidae (placoderm fishes), a review based on new specimens of *Holonema* from the Upper Devonian of Western Australia. *Phil. Trans. R. Soc. Lond. B* **263**, 101–234. (doi:10.1098/rstb.1971.0111)
36. King B, Hu Y, Long JA. 2017 Electroreception in early vertebrates: survey, evidence and new information. *Palaeontology* **61**, 325–358. (doi:10.1111/pala.12346)
37. Dupret V, Sanchez S, Goujet D, Ahlberg PE. 2017 The internal cranial anatomy of *Romundina stellina* Ørvig, 1975 (Vertebrata, Placodermi, Acanthothoraci) and the origin of jawed vertebrates—anatomical atlas of a primitive gnathostome. *PLoS ONE* **12**, e0171241. (doi:10.1371/journal.pone.0171241)
38. Qiao T, King B, Long JA, Ahlberg PE, Zhu M. 2016 Early gnathostome phylogeny revisited: multiple method consensus. *PLoS ONE* **11**, e0163157. (doi:10.1371/journal.pone.0163157)
39. Hu Y, Lu J, Young GC. 2017 New findings in a 400 million-year-old Devonian placoderm shed light on jaw structure and function in basal gnathostomes. *Sci. Rep.* **7**, 7813. (doi:10.1038/s41598-017-07674-y)
40. Young G. 1984 Reconstruction of the jaws and braincase in the Devonian placoderm fish *Bothriolepis*. *Palaeontology* **27**, 635–661.
41. Goujet D. 1984 *Les poissons placodermes du Spitsberg*. Paris, France: Cahiers de Paléontologie, Editions du CNRS.
42. Lebedev OA, Johanson Z, Kuznetsov AN, Tsessarsky A, Trinajstić K, Isakhodzayev FB. 2022 Feeding in the Devonian antiarch placoderm fishes: a study based upon morphofunctional analysis of jaws. *J. Paleontol.* **96**, 1413–1430. (doi:10.1017/jpa.2022.54)
43. White EI, Toombs HA. 1972 The buchanosteoid arthrodires of Australia. *Bull. Br. Mus. (Natural History) Geol.* **22**, 377–419.
44. Long JA, Mark-Kurik E, Young GC. 2014 Taxonomic revision of buchanosteoid placoderms (Arthrodira) from the Early Devonian of south-eastern Australia and Arctic Russia. *Aust. J. Zool.* **62**, 26. (doi:10.1071/ZO13081)
45. Goujet D, Young GC. 2004 Placoderm anatomy and phylogeny: new insights. In *Recent advances in the origin and early radiation of vertebrates* (eds G Arratia, MVH Wilson, R Cloutier), pp. 109–126. München, Germany: Verlag Dr Friedrich Pfeil.
46. Béchard I, Arsenault F, Cloutier R, Kerr J. 2014 The Devonian placoderm fish *Bothriolepis canadensis* revisited with three-dimensional digital imagery. *Palaeontol. Electron.* **17**, 17.1.2A. (doi:10.26879/417)
47. King B, Rücklin M. 2020 A Bayesian approach to dynamic homology of morphological characters and the ancestral phenotype of jawed vertebrates. *eLife* **9**, e62374. (doi:10.7554/eLife.62374)
48. Zhu YA, Giles S, Young GC, Hu Y, Bazzi M, Ahlberg PE, Zhu M, Lu J. 2021 Endocast and bony labyrinth of a Devonian ‘placoderm’ challenges stem gnathostome phylogeny. *Curr. Biol.* **88**, 1112–1118.e4. (doi:10.1016/j.cub.2020.12.046)
49. Brazeau MD, Yuan H, Giles S, Jerve AL, Zorig E, Ariunchimeg Y, Sansom RS, Atwood RC. 2023 Data from: A well-preserved ‘placoderm’ (stem-group Gnathostomata) upper jaw from the Early Devonian of Mongolia clarifies jaw evolution. Figshare. (doi:10.6084/m9.figshare.c.6414027)