

Long distance signalling and epigenetic changes in crop grafting

Jeynes-Cupper, Katie; Catoni, Marco

DOI:

[10.3389/fpls.2023.1121704](https://doi.org/10.3389/fpls.2023.1121704)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Jeynes-Cupper, K & Catoni, M 2023, 'Long distance signalling and epigenetic changes in crop grafting', *Frontiers in Plant Science*, vol. 14, 1121704. <https://doi.org/10.3389/fpls.2023.1121704>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



OPEN ACCESS

EDITED BY

Guillaume Moissiard,
UMR5096 Laboratoire Génome et
Développement des Plantes, France

REVIEWED BY

Matthew R. Willmann,
Pairwise, United States
Vicente Pallas,
Polytechnic University of Valencia, Spain

*CORRESPONDENCE

Katie Jeynes-Copper

✉ KEJ031@student.bham.ac.uk
Marco Catoni

✉ m.catoni@bham.ac.uk

SPECIALTY SECTION

This article was submitted to
Plant Cell Biology,
a section of the journal
Frontiers in Plant Science

RECEIVED 12 December 2022

ACCEPTED 10 March 2023

PUBLISHED 20 March 2023

CITATION

Jeynes-Copper K and Catoni M (2023)
Long distance signalling and epigenetic
changes in crop grafting.
Front. Plant Sci. 14:1121704.
doi: 10.3389/fpls.2023.1121704

COPYRIGHT

© 2023 Jeynes-Copper and Catoni. This is
an open-access article distributed under the
terms of the [Creative Commons Attribution
License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or
reproduction in other forums is permitted,
provided the original author(s) and the
copyright owner(s) are credited and that
the original publication in this journal is
cited, in accordance with accepted
academic practice. No use, distribution or
reproduction is permitted which does not
comply with these terms.

Long distance signalling and epigenetic changes in crop grafting

Katie Jeynes-Copper^{1*} and Marco Catoni^{1,2*}

¹School of Biosciences, University of Birmingham, Birmingham, United Kingdom, ²Institute for Sustainable Plant Protection, National Research Council of Italy, Torino, Italy

Humans have used grafting for more than 4000 years to improve plant production, through physically joining two different plants, which can continue to grow as a single organism. Today, grafting is becoming increasingly more popular as a technique to increase the production of herbaceous horticultural crops, where rootstocks can introduce traits such as resistance to several pathogens and/or improving the plant vigour. Research in model plants have documented how long-distance signalling mechanisms across the graft junction, together with epigenetic regulation, can produce molecular and phenotypic changes in grafted plants. Yet, most of the studied examples rely on proof-of-concept experiments or on limited specific cases. This review explores the link between research findings in model plants and crop species. We analyse studies investigating the movement of signalling molecules across the graft junction and their implications on epigenetic regulation. The improvement of genomics analyses and the increased availability of genetic resources has allowed to collect more information on potential benefits of grafting in horticultural crop models. Ultimately, further research into this topic will enhance our ability to use the grafting technique to exploit genetic and epigenetic variation in crops, as an alternative to traditional breeding.

KEYWORDS

solanaceous, cucurbitaceous, epigenetic, DNA methylation, grafting, mobile siRNA, signaling molecules, graft junction

1 Introduction

The global population is anticipated to grow to around 9.9 billion by 2050, and with that, food production will need to at least double to meet the increasing demands (World population data sheet 2020, <https://interactives.prb.org/2020-wpds/>). This is of particular concern considering that arable land dropped by more than half from 1961 to 2016 as a consequence of housing and industrial needs, as well as excessive agrochemical usage. One of the most obvious solutions to this problem is to increase crop productivity, which can be obtained by breeding more favourable traits or using more productive agronomical techniques. Plant grafting is an ancient horticultural technique traditionally used on cultivated trees which has more recently (since the late 1920s) been exploited as a

system to introduce resistance and improve productivity in horticultural herbaceous crops (Tateishi, 1927; Goldschmidt, 2014; Gaion et al., 2018). This technique involves joining two plants together, where one contributes to the upper part (scion) and the other contributes to the lower parts, including the roots (rootstock). In compatible combinations, the scion and the rootstock will grow together as a single plant (Mudge et al., 2009). Due to its simple application, this method is agronomically sustainable and has had many positive effects on food security. The most memorable use of grafting in agriculture involved the introduction of resistance to the soil-borne insect phylloxera (*Phylloxera vastatrix*) in the European wine industry in the 19th century (Pouget, 1990).

Until the 20th century, grafting was limited to woody species but has since been extended to herbaceous high-value crops (Lee and Oda, 2002; King et al., 2010). Research on herbaceous plant grafting was first described in a study in 1927, which found that watermelon (*Citrullus lanatus*) plants grafted onto pumpkin (*Cucurbita* spp.) were more resistant to pathogens and had a higher fruit yield (Tateishi, 1927). By the mid 1990s, grafting of *Arabidopsis thaliana* had been extensively exploited to understand various phenomena, including the movement of nutrients between the roots and shoots and the study of specific gene functions (Tsutsui and Notaguchi, 2017). The most grafted herbaceous crops with commercial relevance belong to the *Solanaceae* or *Cucurbitaceae* family which are grafted to increase their production and to transfer favourable traits from unproductive species or genotypes.

From 2000 to 2020, three regularly grafted species, tomatoes (*Solanum lycopersicum*), eggplant (*Solanum melongena*) and cucumber (*Cucumis sativus*), accounted for 42-45% of the total increase of the global vegetable production (FAO, 2022). This shows that the global cultivation of these crops is of high economic importance, and the use of grafting could contribute to improving their production. In this context, many seed companies have developed programs to breed rootstocks to improve the yield of these crops (Goldschmidt, 2014). These rootstocks are selected for specific traits, such as the ability to provide drought tolerance, soil-borne pathogen resistance, or scion vigour. Some effects of grafting can be explained simply by the intrinsic characteristics of the genotypes involved. For example, a rootstock can provide resistance to soil-borne pathogens due to their natural immunity, preventing the pathogen from accessing the aerial plant portion. However, it is unclear why a rootstock (or scion) can introduce a trait to a particular grafting partner, depending on the combination of genotypes used. This effect mainly involves scion vigour and changes in plant architecture, making it challenging to predict the outcome of grafting between the combination of two genotypes. It is commonly accepted that substances such as hormones, proteins and signalling molecules can be translocated between the scion and rootstock, and can impose a large biological effect on the recipient tissue (Haroldsen et al., 2012). Additional studies have proposed that such signalling molecules can introduce epigenetic changes that drive grafting-induced phenotypes (Molnar et al., 2010; Lewsey et al., 2016). Molecular alterations in the scion can ultimately change the transcriptome and plant phenotypes, thereby controlling critical aspects of plant development (as reviewed by

Feng et al., 2010). Although these effects have been observed in proof-of-concept studies mostly in *Arabidopsis thaliana*, for relevant agronomical traits we often lack direct evidence of the translation of such observations into crops. In this review, we summarise our understanding of the movement of signalling molecules that can trigger epigenetic effects in plant grafting and discuss the role of epigenetics in producing favourable horticultural traits in grafted crops and their progeny.

2 Phenotypic traits and transcriptomic alteration in crop grafting

Over the past decade, researchers have discovered a link between the introduction of grafting-induced phenotypic traits and changes at the transcriptional level. For example, cold-tolerance in tomatoes can be transferred to a susceptible scion by grafting it onto a cold-tolerant tomato rootstock, and this tolerance is associated with changes in the expression of defence-related genes (Ntatsi et al., 2017). Similarly, in watermelon, scions grafted onto bottle gourd (*Lagenaria siceraria*) rootstocks have been shown to trigger an increase in scion fruit size and rind thickness (Garcia-Lozano et al., 2020). Additionally, the improved scion phenotype has been linked to the differential expression of genes related to ripening, softening, cell wall strengthening, stress response and disease resistance (Garcia-Lozano et al., 2020). Interestingly, in the reverse grafting combination (using melon as a rootstock and bottle gourd as the scion), the produced fruits had a higher soluble solid content and thinner rinds (Garcia-Lozano et al., 2020). A previous study on watermelon grafted plants showed that productivity is strongly linked to the efficiency of water uptake (Rouphael et al., 2008). However, it is unclear whether this improved use of water is a consequence of the intrinsic propriety of the rootstock (e.g. extended root system) or the result of induced transcriptional changes.

In an experiment where tomatoes were grafted onto potatoes (*Solanum tuberosum*) to introduce late blight resistance, differences in the transcriptome were observed in both the scion and rootstock of hetero-grafted plants compared to self-grafted controls; however, it was unclear whether such alterations were simply due to changes in photosynthates production (Li and Zhao, 2021). Transcriptional alterations observed in grafted plants have not always been associated with phenotypical changes. In a study where watermelon was grafted onto bottle gourds or squash (*Cucurbita* spp.), the authors identified changes in the expression of many genes in the scion, but they did not observe phenotypic alterations between the heterografted and self-grafted plants (Liu et al., 2016). Similar results were obtained in a study on tomatoes, where heterografted scions displayed altered transcriptional profiles for genes related to oxidative stress, but in absence of phenotypic differences (Wang et al., 2019).

The study of the link between the phenotype and transcriptome in grafted plants is complicated by the fact that the grafting method itself can cause transcriptional changes in scions, as observed in experiments performed with *Arabidopsis* and tomato self-grafted plants (Kumari et al., 2015; Spanò et al., 2015). Studies performed in combination with pathogen viral infections have also suggested that

resistant or tolerant phenotypes could be due to a combination of the grafting method and the interactions between rootstock and scion. For example, a study examining the recovery from infection by *Potato virus Y* (PVY) in tomatoes demonstrated that the grafting technique itself improved recovery from the infection regardless of the use of either a susceptible or resistant rootstock (Spanò et al., 2020). Similarly, a report on the transmission of *Cucumber mosaic virus* (CMV) resistance by grafting in tomatoes showed that grafting a susceptible variety onto a resistant variety would enable the same disease recovery observed in the resistant lines, but recovery was also observed in the self-grafts of the susceptible variety (Spanò et al., 2017). Viruses move through the plant vascular tissue and must eventually pass through a graft junction to establish a systemic infection; thus, the alteration of vascular connections induced by grafting can directly affect the spread of the virus. Nonetheless, a transgenic tomato rootstock producing viral sRNAs and resistant to the *Tomato spotted wilt virus* (TSWV) was found to delay symptoms and strongly reduce virus accumulation in grafted scions, compared to scions grafted on susceptible tomato lines (Catoni et al., 2013). This indicates that a certain degree of tolerance to viral infection mediated by viral-derived sRNAs can, at least in certain instances, be transferred through grafting. Collectively, we have good evidence that grafting of different genotypes induces changes in gene expression, although this is not always associated with a clear transfer of a phenotype.

3 Movement of RNAs across the graft junction

Plants have evolved complex communication networks and transport systems, where roots and shoots constantly communicate to coordinate growth and development (Liu and Chen, 2018). Such communication is facilitated by the vascular system, which comprises of the xylem and phloem. The xylem is responsible for the transportation of inorganic salts and water, while the phloem is responsible for the transportation of plant hormones, photosynthates, amino acids and notably mobile RNA molecules such as messenger RNA (mRNA), small interfering RNA (siRNA) and micro RNA (miRNA) (Hu et al., 2016). When plants are grafted, the vascular systems of the scion and rootstock fuse in a compatible match. The phloem is the first to connect, followed by the xylem, along with the formation of plasmodesmata between adhered cells, which establishes cell-to-cell communication and facilitates long-distance signalling (Jeffree and Yeoman, 1983; Kollmann and Glockmann, 1985; Ferreres et al., 2011; Melnyk et al., 2015).

Mobile RNAs are well known to be transported inside plants and act as signalling macromolecules to coordinate an array of physiological processes (Lucas et al., 2001; Lough and Lucas, 2006; Luo et al., 2018). Grafting in *Arabidopsis thaliana* has been used extensively to prove that RNAs can travel bidirectionally between rootstock and scion tissues *via* the grafting junction (Harada, 2010; Spiegelman et al., 2013; TurnbullLopez-Cobollo, 2013). Some mobile RNAs can introduce epigenetic changes to their destination tissues and trigger alterations in plant physiology and

architecture (Kehr and Kragler, 2018; Liu and Chen, 2018). Numerous studies have demonstrated that also non-coding RNAs can regulate the expression of specific genes *via* transcriptional repression playing an important role in the emergence of specific phenotypes or traits in response to development or stress. Such transcriptional repression is achieved *via* changes in the epigenome, specifically DNA methylation (Song et al., 2003; Zhang et al., 2018a; Zhang et al., 2018b).

3.1 Movement of small non-coding RNA molecules

Small non-coding RNAs represent a cohort of different molecules in plants, with a size of 20 – 24 nucleotides. They can be broadly classified into two groups: i) Micro RNAs (miRNA), or ii) Small interfering RNAs (siRNA or sRNA). These two classes differ by the way siRNAs are synthesised and how they operate (Borges and Martienssen, 2015; Kamthan et al., 2015; Singh et al., 2018). In plants, miRNAs are encoded by endogenous genes that are transcribed into long primary miRNAs (pri-miRNAs) that are characteristically folded into a hairpin-like structures. These are then processed by RNase III-like Dicer 1 (DCL1), first into stem-loop structures, known as precursor miRNAs (pre-miRNAs), then into mature miRNA duplexes. The resulting miRNAs can prevent mRNA translation through Post Transcriptional Gene Silencing (PTGS), which ultimately causes gene repression (Borges and Martienssen, 2015) (Figure 1). In comparison, siRNAs are derived from long double-stranded RNA precursors and are grouped based on their function. The first group contains siRNAs with a nucleotide length of 21 - 22 and are derived from the activity of Dicer-like 2 and Dicer-like 4 (DCL2/DCL4) from trans-acting small interfering RNA (TAS) genes, transcripts or other aberrant RNA molecules (Berger et al., 2018). The 21/22-nt siRNAs are known to be involved in short-distance signalling and in the degradation of mRNAs (Tan et al., 2020), as well as in targeted Transcriptional Gene Silencing (TGS) (Figure 1) (Borges and Martienssen, 2015). The second group is known as 24-nt siRNAs which originate from heterochromatic regions by the continuous action of three enzymes: RNA Polymerase IV, RNA Dependent RNA polymerase 2 (RDR2) and Dicer-like 3 (DCL3). The 24-nt siRNAs are renowned for their involvement in the RNA-directed DNA methylation (RdDM) pathway by guiding the Domain rearranged DNA methyltransferases (DRMs) in TGS, but they are also known for their involvement in long-distance signalling (Berger et al., 2018) (Figure 1).

In the model species *Arabidopsis thaliana*, it has been demonstrated that the systemic movement of siRNAs can direct *de novo* DNA methylation in tissues distant from their origin (Borges and Martienssen, 2015; Xie and Yu, 2015) (Figure 2). This involves the recruitment of DNA methyltransferases in recipient cells or tissues, and the loading of the siRNA into ARGONAUTES (AGOs) proteins, which are then recruited into a complex with transcribed scaffold RNA by the action of RNA Polymerase V (Huang et al., 2021). More specifically, AGO1 protein in *Arabidopsis* has been proposed to remove sRNAs from

Four Classes of Mobile RNAs in Plants




FIGURE 1 Four classes of mobile RNA molecules in plants. Potential mobile RNA molecules include messenger RNA (mRNA) molecules which encode proteins; transfer RNA (tRNA) -like structures which are associated to movement of RNA molecules; micro RNA (miRNA) which regulate gene expression by guiding gene silencing mechanisms such as Post Transcriptional Gene Silencing (PTGS) and small interfering RNAs (siRNA) which mediate the silencing of gene expression through PTGS and/or through Transcriptional Gene Silencing (TGS) via the RNA-directed DNA methylation pathway (RdDM). Figure created with [BioRender.com](https://www.biorender.com).

traversed tissue, allowing their accumulation in recipient cells, a phenomenon defined as “consumption” of the mobile sRNAs (Denvers et al., 2020). Further studies identified in Squash pumpkin (*Cucurbita maxima*) phloem exudates a set of specific proteins able to bind sRNAs and mediate non-cell autonomous movement of both 21/22- and 24- nt classes (reviewed in Yan and

Ham, 2022). Among these, the PHLOEM SMALL RNA-BINDING PROTEIN1 (PSRP1) has being found able to form a ribonucleoprotein complex in the phloem sieve tube system (Ham et al., 2014), while the SMALL RNA-BINDING PROTEIN 1 (SRBP1) homologues in Arabidopsis has been found capable of cell-to-cell movement and has a role in sRNA trafficking into the




FIGURE 2 The movement of small RNA (siRNA) between root and shoot drive epigenetic modifications and lead to phenotypical changes in the plant. The siRNAs can move both from root-to-shoot and shoot-to-roots directions, via the phloem through the sieve tubes. Once at the destination tissue, the siRNAs can alter DNA methylation at specific loci via the RNA-directed DNA Methylation (RdDM) pathway, or prevent the translation of a gene product via the Post-Transcriptional Gene Silencing (PTGS) pathway. Messenger RNAs (mRNA) can move across a systemically (and across a graft junction) into cells where they are translated into transcription factors (TFs) to control gene expression. Although not yet demonstrated in grafting experiment, siRNAs can introduce histone modifications. Alterations to these epigenetic marks can result in changes to the plant phenotype including resistance to abiotic and biotic stress, as well as alteration in the plant architecture and vigour. Figure created with [BioRender.com](https://www.biorender.com).

apoplast (Yang et al., 2019; Karimi et al., 2022). These results suggest the existence of a specific mechanism for the systemic mobilization of sRNA molecules in plants, which is currently subject of research and discussion (Voinnet, 2022; Yan and Ham, 2022).

The translocation of siRNAs across the grafting junction has also been explored in several crops, including eggplant, pumpkin (*Cucurbita pepo*), cucumber, potato, *Nicotiana benthamiana*, grapevine (*Vitis vinifera*) and several bean (*Fabaceae*) species (Tournier et al., 2006; Bai et al., 2011; Zhang et al., 2014; Cerruti et al., 2021; Li et al., 2021; Davoudi et al., 2022; Rubio et al., 2022; Zhang et al., 2022). In addition, several miRNAs in potato have been demonstrated to be graft-transmissible, including *miR156*, *miR339*, *miR395* and *miR172* (Pant et al., 2008; Buhtz et al., 2010; Kasai et al., 2010). *MiR156* has been found to influence plant architecture and tuberization by targeting the regulation of specific genes (Bhogale et al., 2013). The movement of siRNAs was also studied by Kasai et al., who demonstrated that artificial siRNAs in *N. benthamiana* scions could travel into the potato rootstock to induce TGS via RdDM on both a transgene and an endogenous gene (Kasai et al., 2016). In a recent report, grafting was used to induce editing in scions using clustered regularly interspaced short palindromic repeats (CRISPR)-associated protein 9 (*Cas9*) and guide RNA transcripts moved from a transgenic rootstock (Yang et al., 2023). In addition to demonstrating the potential use of grafting in combination with genome editing, this study also showed that mobilised transcripts in the scion can accumulate at a sufficient abundance to introduce a gene editing effect.

In plants, the PTGS pathway falls under RNA interference (RNAi) (Ashfaq et al., 2020). This mechanism involves the suppression of gene expression through degradation of mRNA transcripts via the production of siRNAs (Ashfaq et al., 2020) (Figure 2) (Hamilton and Baulcombe, 1999; Jones et al., 1999; Mette et al., 1999; Dalmay et al., 2000; Marjori et al., 2001). Therefore, it is accepted that both PTGS and DNA methylation function cooperatively or in a dependent manner as initially suggested by Morel et al., 2000 (Morel et al., 2000), and subsequently proved in multiple experimental conditions (Palauqui et al., 1997; Sonoda and Nishiguchi, 2000; Voinnet et al., 2000; Crété et al., 2001; Mallory et al., 2001; Mallory et al., 2003; García-Pérez et al., 2004; Hewezi et al., 2005; Brosnan et al., 2007; Brumin et al., 2009; Kasai et al., 2011; Taochy et al., 2017; Chen et al., 2018). However, current literature exploring this aspect in grafted plants is limited, and phenotypic changes induced by grafting have only been indirectly linked to DNA methylation alterations (Cerruti et al., 2021).

3.2 Movement of mRNA transcripts

Unlike siRNAs, mobile mRNAs can directly contribute to the synthesis of proteins into recipient tissues. They have been found to be particularly relevant as transcription factors (TFs), which play a key role in controlling the expression of several genes in a specific pathway. The long-distance mobilisation of mRNAs has been

documented in several models using grafting as a tool. For example, in tomato it has been observed that the mRNA transcript *PPF-LeT6*, composed by a PYROPHOSPHATE-DEPENDENT PHOSPHOFRUCTOKINASE (PPF) and tomato KNOTTED-1-like homeobox (KNOX) domain (*LeT6*), is able to travel from roots to shoots and induce a change in the leaf morphology (Kim et al., 2001). Similarly, both RT-PCR and *in situ* hybridization revealed that mRNA transcripts of two partner transcription factors, the *potato homeobox1* (*POTH1*) gene and BEL1-type *StBEL5*, can move across the graft junction and regulate hormone accumulation, alter leaf architecture, and boost tuber formation (Banerjee et al., 2006; Mahajan et al., 2012; Sharma et al., 2014). Additional studies on pumpkin grafts have also identified other mobile mRNA transcripts that are likely to drive molecular and phenotypic changes, including those encoding transcriptional regulators of auxin signalling (Omid et al., 2007). This was followed by studies on *Arabidopsis*, which characterised specific auxin mRNA transcripts produced in leaf tissue and translocated to the root system, where they negatively regulate lateral root formation (Omid et al., 2007; Notaguchi et al., 2012). Within the same timeline, researchers have also identified similar hormone-related mRNA transcripts in tomato, cucumber, and pumpkin (Kim et al., 2001; Haywood et al., 2005; Banerjee et al., 2006; Notaguchi et al., 2012; Davoudi et al., 2022). In *Arabidopsis*, the dwarf phenotype observed in gibberellin (GA)-deficient mutants can be restored if plants are grafted onto WT (Magome et al., 2004), whereas tomato GA-deficient rootstocks can introduce a dwarf-like phenotype in the scion, which is mediated by the movement of transgenic mRNA transcripts up the graft junction (Wang et al., 2012). In tomato has been also found that the mRNA encoding for the GIBBERELLIC ACID INSENSITIVE (GAI) factor produced in a rootstock can transcriptionally regulate gibberellic acid response genes within the scion tissue (Haywood et al., 2005). In addition, there has been evidence that a GAI homologous can be transported in pumpkins by an RNA-protein complex (Ham et al., 2009). In a study investigating drought tolerance in cucumber and pumpkin (*Cucumis moschata*) grafts, authors identified that the movement of mobile mRNAs correlated with significant accumulation of abscisic acid (ABA) from the pumpkin rootstock in the cucumber scion tissue (Davoudi et al., 2022).

Collectively, it has been well-documented, both in model and crop species, that at least some mRNAs can move across the graft junction and induce phenotypic and developmental changes. However, it remains unclear whether these transcripts are primarily responsible for the observed traits or instead trigger more general epigenetic or post-transcriptional regulation mechanisms responsible for the scion/rootstock physiological response. Hence, further work could be directed to mechanistically associate mobile signalling molecules to an observed epigenetic and/or phenotypic change in grafted plants. In this context, the use of experimental grafting associated with genome-wide sequencing approaches appears to be a powerful tool for correlating phenotypes to mobile mRNA molecules. In experiments where scions and rootstock were from different sequenced genomes, it became possible to compare mRNA sequencing experiments with DNA sequence variants in the two genotypes and identify mobile mRNAs with greater confidence. The

methods used to analyse transcriptomic data are well established in plants, but only few analyses or pipelines have been specifically designed for the detection of mobile transcripts (Tomkins et al., 2022). Further development of such tools should be considered to support a better understanding of the molecular mechanisms underlying grafting-induced traits.

3.3 Transport mechanisms

The transport mechanism of mobile RNAs is still not well understood and has been predominantly investigated in the model species *Arabidopsis thaliana*. One of the first studies to identify an association between mRNA abundance and mobility in grafted plants proposed a hypothesis for RNA transport based on passive movement (Thieme et al., 2015). This was supported by a more recent study that found that long-distance movement of mRNA is linked to their local abundance (Calderwood et al., 2016). However, this work could only explain the movement of some mRNAs, and further hypotheses suggest the simultaneous passive and active movement of mobile RNA species (Kim et al., 2014; Notaguchi et al., 2015). The active transportation of RNA molecules has been then proposed in multiple subsequent studies (Zhang et al., 2009; Thieme et al., 2015; Zhang et al., 2016; Garcia-Lozano et al., 2020). Such active movement appears to be associated with the presence of transfer RNA-like structure (TLS) motifs, which can mediate the mobility of some mRNA transcripts to distant locations, as observed in both *Arabidopsis* and pumpkin (Zhang et al., 2009; Zhang et al., 2016). Experimental work has shown that the addition of a TLS motif, or a specific sequence derived from a TLS motif, is sufficient to introduce transcript mobility, while their removal from an RNA molecule prevents long-distance transportation (Garcia-Lozano et al., 2020; Yang et al., 2023). Collectively, these results suggest that TLSs or related structures may control the translocation of mobile transcripts in graft systems.

In addition to sequences associated with mobility in RNA molecules, also cytosine methylation can play a role in RNA mobility (Yang et al., 2019). Both plant phloem transfer RNAs (tRNAs) and graft-mobile mRNAs contain TLS motifs that have a high probability of being targeted by RNA methyltransferases (Yang et al., 2019). Hence, mobile RNA transcripts are enriched in cytosine methylation, and if such methylation is removed, mobility can be negatively affected (Yang et al., 2019). Much of what is known about the transport of RNA species in plants is credited to research conducted on viroids, which are circular non-coding RNA species often considered plant pathogens, mostly because of their recognition as an ideal model system for molecular transport studies (Capistrán-Barradas et al., 2006; Ding and Itaya, 2007; Wang et al., 2021). For example, research performed on viroids has found that their long-distance mobility is facilitated by selective trafficking based on the presence of RNA motifs or structures (Qi et al., 2004; Zhong et al., 2007; Zhong et al., 2008; Takeda et al., 2018) and a similar concept has been supported by several studies related to plant-derived mobile RNAs (Zhang et al., 2009; Thieme et al., 2015; Zhang et al., 2016; Garcia-Lozano

et al., 2020). In addition, experiments performed with grafting in cucumber demonstrated that viroid movement is mediated by specific host proteins (Gómez and Pallás, 2004), suggesting that a similar mechanism exists for endogenous RNA molecules. Interestingly, it was found that a specific plant protein assists the mobility of at least one endogenous mRNA in pear (*Pyrus betulaefolia*) (Duan et al., 2016), and recent studies have suggested that organelle-assisted movement is also used for the translocation of endogenous sRNAs in herbaceous plants (Luo et al., 2022). This hypothesis suggests that mRNAs move by interacting with organelles that are transported out of the phloem into the destination tissue (Luo et al., 2022). An example of this mechanism was observed with the mRNA of the mobile factor FLOWERING LOCUS T, which has been found to move through the plasmodesmata via microtubules in the endosomes (Luo et al., 2022). A similar finding was observed in maize, where mobilisation of the mRNA of the KNOTTED1 (KN1) TF, a highly conserved protein involved in stem cell production, moves in shoot meristems. This is of particular interest because, although the movement of TFs is widely known, for KN1 functionality it requires trafficking of both its protein and mRNA (Kitagawa et al., 2022).

Collectively, the data indicates the existence of active mechanism for the systemic transport of at least one group of mobile RNA molecules. Considering that distant mobile molecules can trigger effects through a graft junction, these studies are important for clarifying the mechanisms underlying how a rootstock can induce physiological changes in scions or vice versa.

4 Changes to the epigenome associate to grafting

Epigenetic marks are constituted by molecular alterations of chromatin which does not alter the sequence of the DNA itself, and can play critical roles in controlling the expression of genes, with consequences on the morphology, physiology, and ecology of many organisms (Rapp and Wendel, 2005; Fossey, 2009). In plants, there are two main categories of epigenetic marks: DNA methylation and histone modifications (Rapp and Wendel, 2005; Fossey, 2009). Cytosine is the only DNA nucleotide that is methylated, and there are a multitude of different histone modifications, including acetylation, methylation, phosphorylation, ribosylation, and ubiquitination (Pfluger and Wagner, 2007; Liu et al., 2010). These epigenetic changes play a role in the regulation of chromatin structure and control the accessibility of DNA (Robertson, 2005; Lang et al., 2017). Epigenetic variation can generate epigenetic alleles (or epialleles) with different degrees of stability, which are transmitted during DNA replication and often mitotically passed to progeny in a Mendelian fashion (Weigel and Colot, 2012; Catoni and Cortijo, 2018). In the previous chapter, we discussed how mobile RNA molecules travelling across the graft junction can trigger epigenetic changes at specific target loci in recipient tissues. Therefore, in this section, we discuss the evidence of epigenetic alterations associated with plant grafting.

4.1 Alterations of DNA methylation in grafted plants

Changes in DNA methylation associated with grafting have been observed in many plant families. The first studies were conducted on *Brassicaceae*, *Cucurbitaceae*, and *Solanaceae* species (Molnar et al., 2010; Avramidou et al., 2015; Lewsey et al., 2016). In *Arabidopsis*, a siRNA molecule originating from a transgene was found to move across the graft junction with prevalent translocation in the shoot-to-root direction. At specific loci, these molecules have been found to direct DNA methylation, resulting in a silencing signal (Molnar et al., 2010). The use of a transgenic construct has also been employed in tobacco to show that siRNA produced in the rootstock can travel across the graft junction and induce silencing of complementary DNA sequences in the scion by increasing DNA methylation (Bai et al., 2011). It was later revealed that the RdDM pathway has a critical role in triggering silencing by introducing DNA methylation at specific genomic loci (Lewsey et al., 2016). A general pattern of grafting-induced hypermethylation has been observed in *Cucurbitaceae* species when cucumber and melon were used a scion, grafted onto pumpkin rootstocks (Avramidou et al., 2015). In eggplant, however, hypomethylation in the scions was found to be associated with heterografting and linked to enhanced vigour (Cerruti et al., 2021). Similar studies across *Brassicaceae* and *Cucurbitaceae* identified methylation changes, mostly related to the non-CG context, associated with specific grafting combinations, but with different trends of hyper- and hypo-methylation (Avramidou et al., 2015; Lewsey et al., 2016; Cerruti et al., 2021). In these studies, the DNA methyltransferases and DNA demethylase genes in the scions were not found to be substantially differentially expressed, indicating that the epigenetic changes observed cannot be simply explained by altered expression levels of these epigenetic factors. Therefore, most of the DNA methylation changes associated with grafting could indeed be a consequence of a change in the siRNA targeting system or, alternatively, the result of other indirect effects of grafting (e.g. passive de-methylation or carry-over effects from the grafting procedure) (Wu et al., 2013; Cerruti et al., 2021).

Notably, hypomethylation and hypermethylation are associated with opposite effects on gene expression, and different combinations of scion and rootstock genotypes can lead to different phenotypic effects. Therefore, epigenetic changes could potentially explain part of the phenotypic alteration (e.g. vigour) observed in scions grafted onto different rootstock genotypes (Yang et al., 2018). In this context, it would be interesting to test the role of epigenetics in other grafting-induced phenotypes, such as disease resistance, which has been directly associated with epigenetic variation (Wada et al., 2004).

4.2 Grafting-induced histone modifications

While direct evidence of grafting-induced histone modifications has not yet been observed, there are studies suggesting that epigenetic modification induced by grafting is not limited to the alteration of the DNA methylation landscape (Figure 2). It is well established that there is a crosstalk between DNA methylation,

siRNAs accumulation and histone modifications, so that epigenetic changes induced by distant signaling are expected to occur also at level of chromatin structure (Saze et al., 2012).

In a study performed on grapevine heterografts, the vigorous phenotype observed with specific grafting combinations was associated with the overexpression of the core histone genes as well as two nucleosome-associated proteins (Cookson and Ollat, 2013), while on apple tree varieties has been found that the expression of genes encoding histone modification factors was central to control flowering (Fan et al., 2018).

Therefore, histone modifications might play a role in the physiological effects induced by grafting and the extension of specific approaches to study chromatin dynamic (e.g. ATAC-seq or 3C) to grafting experiments could reveal more details on how histone marks can be associated with crop traits.

4.3 Transgenerational inheritance of grafting-induced epigenetic changes and implication to breeding

Epigenetic modifications were originally thought to only affect an individual during their lifespan and were not perceived to impact the offspring. In 1956, these beliefs were challenged and ideas that traits induced by the environment could be inherited began to be discussed (Waddington, 1956). Recently, epigenetic inheritance has shown that environmentally controlled epialleles can be transmitted to progeny, bridging the gap between inheritance by genetic determinants and environmental influences (Van Otterdijk and Michels, 2016; Pang et al., 2017). Hence, determining whether grafting-induced epigenetic alterations are transgenerationally inherited is pivotal to better understand plant developmental mechanisms and to exploit their potential use in breeding schemes to consistently increase crop production. In *Brassicaceae* models, DNA methylation alterations induced by grafting in the scion was shown to be indeed heritable in the offspring of grafted plants (Cao et al., 2016). The first generation of self-fertilised progenies from tuber mustard (*Brassica juncea*) grafted onto red cabbage (*Brassica oleracea*) displayed a genome-wide change in DNA methylation levels by 5.29% to 6.59% (for hypermethylation and hypomethylation, respectively), compared to the parental lines. By the 5th generation, 31.58% of these changes were still present, while the rest had reverted to the original methylation levels (Cao et al., 2016). Similarly, in tomato and *Arabidopsis*, the vigorous phenotype induced by a mutation in *MutS HOMOLOGUE 1* (*MSH1*) can be passed through grafting via siRNAs, and is inherited in the offspring of wild-type scions (Kundariya et al., 2020).

Over the years, studies have proven that epialleles can arise naturally and can influence agronomically important traits, such as fruit ripening in tomato (Manning et al., 2006) or the seed size and leaf angle of rice (Zhang et al., 2015). Genome-wide and epigenome-wide association studies performed in maize and wheat have highlighted strong correlations between differentially methylated regions (DMRs) and the expression of many genes responsible for crop traits (Gardiner et al., 2018; Xu et al., 2019). These and other studies indicate the great potential of using epialleles in crop breeding for trait improvement. The

identification of epialleles linked to agronomically important traits in the progeny of crop grafts could give rise to novel epi-molecular markers (Noshay and Springer, 2021). The prospects of epigenetic breeding have been illustrated in soybean, where improved yields have been observed in the third generation by suppressing MSH1 (Raju et al., 2018). Similar results were observed when replicating MSH1 suppression in tomatoes and *Arabidopsis* (Kundariya et al., 2020). A clear benefit to producing heritable traits through grafting is the possibility of efficiently introducing new genetic and epigenetic diversity from wild species into crops, without the necessity to introgress the trait with breeding or by genetic transformation. Introducing stably transmissible epigenetic traits could ultimately represent a viable and unique technique for developing new crop varieties.

For crop species that are popularly grafted commercially, such as cucumber, tomato, and pepper, future endeavours should identify and preserve the biodiversity of natural varieties and species in light of their potential use as rootstock or scions in grafting combinations. As techniques continue to advance (e.g. micrografting), they will allow us to investigate the genetic and epigenetic variations in the genomes of plants improved by grafting. This will enable the development of new resources and drive comparative studies to fully connect the many facets of epigenetic regulation with the final goal of improving modern crop breeding.

Author contributions

KJ-C undertook the initial research and wrote the draft of the manuscript, as well as preparing Figures 1 and 2. MC provided constructive feedback and novel perspectives, edited the manuscript, introduced new written elements, and finalised the manuscript. All authors contributed to the article and approved the submitted version.

References

- Ashfaq, M. A., Dinesh Kumar, V., Soma Sekhar Reddy, P., Anil Kumar, C., Sai Kumar, K., Narasimha Rao, N., et al. (2020). Post-transcriptional gene silencing: Basic concepts and applications. *J. Biosci.* 45, 128. doi: 10.1007/s12038-020-00098-3
- Avramidou, E., Kapazoglou, A., Aravanopoulos, F. A., Xanthopoulou, A., Ganopoulos, I., Tsaballa, A., et al. (2015). Global DNA methylation changes in cucurbitaceae inter-species grafting. *Crop Breed. Appl. Biotechnol.* 15, 112–116. doi: 10.1590/1984-70332015v15n2n20
- Bai, S., Kasai, A., Yamada, K., Li, T., and Harada, T. (2011). A mobile signal transported over a long distance induces systemic transcriptional gene silencing in a grafted partner. *J. Exp. Bot.* 62, 4561–4570. doi: 10.1093/jxb/err163
- Banerjee, A. K., Chatterjee, M., Yu, Y., Suh, S.-G., Miller, W. A., and Hannapel, D. J. (2006). Dynamics of a mobile RNA of potato involved in a long-distance signaling pathway. *Plant Cell* 18, 3443–3457. doi: 10.1105/tpc.106.042473
- Berger, M. M., Gallusci, P., and Teyssier, E. (2018). Roles of epigenetic mechanisms in grafting and possible applications. *Adv. Botanical Res. Elsevier* 88, 203–246. doi: 10.1016/bs.abr.2018.10.003
- Bhogale, S., Mahajan, A. S., Natarajan, B., Rajabhoj, M., Thulasiram, H. V., and Banerjee, A. K. (2013). MicroRNA156: A potential graft-transmissible MicroRNA that modulates plant architecture and tuberization in solanum tuberosum ssp. andigena. *Plant Physiol.* 164, 1011–1027. doi: 10.1104/pp.113.230714
- Borges, F., and Martienssen, R. A. (2015). The expanding world of small RNAs in plants. *Nat. Rev. Mol. Cell Biol.* 16, 727–741. doi: 10.1038/nrm4085
- Brosnan, C., Mitter, N., Christie, M., Smith, N., Waterhouse, P., and Carroll, B. (2007). Nuclear gene silencing directs reception of long-distance mRNA silencing in *Arabidopsis*. *Proc. Natl. Acad. Sci.* 104, 14741–14746. doi: 10.1073/pnas.0706701104
- Brumin, M., Stukalov, S., Haviv, S., Muruganatham, M., Moskovitz, Y., Batuman, O., et al. (2009). Post-transcriptional gene silencing and virus resistance in *Nicotiana benthamiana* expressing a grapevine virus a minireplicon. *Transgenic Res.* 18, 331–345. doi: 10.1007/s11248-008-9222-3
- Buhtz, A., Pieritz, J., Springer, F., and Kehr, J. (2010). Research article phloem small RNAs, nutrient stress responses, and systemic mobility. *BMC Plant Biol.* 10, 64. doi: 10.1186/1471-2229-10-64
- Calderwood, A., Kopriva, S., and Morris, R. J. (2016). Transcript abundance explains mRNA mobility data in *Arabidopsis thaliana*. *Plant Cell* 28, 610–615. doi: 10.1105/tpc.15.00956
- Cao, L., Yu, N., Li, J., Qi, Z., Wang, D., and Chen, L. (2016). Heritability and reversibility of DNA methylation induced by *in vitro* grafting between brassica juncea and *b. oleracea*. *Sci. Rep.* 6, 27233. doi: 10.1038/srep27233
- Capistrán-Barradas, A., Moreno-Casasola, P., and Defeo, O. (2006). Postdispersal fruit and seed removal by the crab *Gecarcinus lateralis* in a coastal forest in Veracruz, Mexico 1. *Biotropica: J. Biol. Conserv.* 38, 203–209. doi: 10.1111/j.1744-7429.2006.00116.x
- Catoni, M., and Cortijo, S. (2018). EpiRILs: lessons from *Arabidopsis*. *Adv. Botanical Res. Elsevier* 88, 87–116. doi: 10.1016/bs.abr.2018.08.002

Funding

Work by MC is funded by the Royal Society Research Grant [RGS\R1\201297] (<https://royalsociety.org>) and the UK Research through the Biotechnology and Biological Sciences Research Council (BBSRC) grant [BB/W008866/1] and the Natural Environment Research Council grant [NE/V021346/1] (<https://www.ukri.org>). KJ-C is supported by the University of Birmingham Midlands Integrative Biosciences Training Partnership (MIBTP), funded by BBSRC with the grant [BB/T00746X/1], and partially funded by the company Enza Zaden (<https://www.enzazaden.com/>).

Conflict of interest

Author KJ-C PhD fellowship is partially funded by the vegetable-breeding company Enza Zaden (<https://www.enzazaden.com/>), which develops rootstocks for commercial grafting of horticultural species.

The remaining author declare that the research was conducted in the absence of any other commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

- Catoni, M., Lucioi, A., Doblaz-Ibáñez, P., Accotto, G. P., and Vaira, A. M. (2013). From immunity to susceptibility: virus resistance induced in tomato by a silenced transgene is lost as TGS overcomes PTGS. *Plant J.* 75, 941–953. doi: 10.1111/tpj.12253
- Cerruti, E., Gisbert, C., Drost, H.-G., Valentino, D., Portis, E., Barchi, L., et al. (2021). Grafting vigour is associated with DNA de-methylation in eggplant. *Horticulture Res.* 8, 241. doi: 10.1038/s41438-021-00660-6
- Chen, W., Zhang, X., Fan, Y., Li, B., Ryabov, E., Shi, N., et al. (2018). A genetic network for systemic RNA silencing in plants. *Plant Physiol.* 176, 2700–2719. doi: 10.1104/pp.17.01828
- Cookson, S. J., and Ollat, N. (2013). Grafting with rootstocks induces extensive transcriptional re-programming in the shoot apical meristem of grapevine. *BMC Plant Biol.* 13, 147. doi: 10.1186/1471-2229-13-147
- Crété, P., Leuenberger, S., Iglesias, V. A., Suarez, V., Schöb, H., Holtorf, H., et al. (2001). Graft transmission of induced and spontaneous post-transcriptional silencing of chitinase genes. *Plant J.* 28, 493–501. doi: 10.1046/j.1365-313X.2001.01171.x
- Dalmay, T., Hamilton, A., Mueller, E., and Baulcombe, D. C. (2000). Potato virus X amplicons in arabidopsis mediate genetic and epigenetic gene silencing. *Plant Cell* 12, 369–379. doi: 10.1105/tpc.12.3.369
- Davoudi, M., Song, M., Zhang, M., Chen, J., and Lou, Q. (2022). Long-distance control of pumpkin rootstock over cucumber scion under drought stress as revealed by transcriptome sequencing and mobile mRNAs identifications. *Horticulture Res.* 9, uhab033. doi: 10.1093/hr/uhab033
- Denvers, E. A., Brosman, C. A., Sarazim, A., Albertini, D., Amsler, A. C., Brioudes, F., et al. (2020). Movement and differential consumption of short interfering RNA duplexes underlie mobile RNA interference. *Nat. Plants* 6, 789–799. doi: 10.1038/s41477-020-0687-2
- Ding, B., and Itaya, A. (2007). Viroid: a useful model for studying the basic principles of infection and RNA biology. *Mol. Plant-Microbe Interact.* 20, 7–20. doi: 10.1094/MPMI-20-0007
- Duan, X., Zhang, W., Huang, J., Hao, L., Wang, S., Wang, A., et al. (2016). PbWoxT1 mRNA from pear (*Pyrus betulaefolia*) undergoes long-distance transport assisted by a polypyrimidine tract binding protein. *New Phytol.* 210, 511–524. doi: 10.1111/nph.13793
- Fan, S., Wang, J., Lei, C., Gao, C., Yang, Y., Li, Y., et al. (2018). Identification and characterization of histone modification gene family reveal their critical responses to flower induction in apple. *BMC Plant Biol.* 18, 1–21. doi: 10.1186/s12870-018-1388-0
- FAO (2022). Agricultural production statistics 2000–2020, *FAOSTAT Analytical, Brief 41*. Rome.
- Feng, S., Jacobsen, S. E., and Reik, W. (2010). Epigenetic reprogramming in plant and animal development. *Science* 330, 622–627. doi: 10.1126/science.1190614
- Ferreeres, F., Figueiredo, R., Bettencourt, S., Carqueijeiro, I., Oliveira, J., Gil-Izquierdo, A., et al. (2011). Identification of phenolic compounds in isolated vacuoles of the medicinal plant *catharanthus roseus* and their interaction with vacuolar class III peroxidase: an H₂O₂ affair? *J. Exp. Bot.* 62, 2841–2854. doi: 10.1093/jxb/erq458
- Fossey, A. (2009). Epigenetics: beyond genes. *South. Forests: J. For. Sci.* 71, 121–124. doi: 10.2989/SF.2009.71.2.6.821
- Gaion, L. A., Braz, L. T., and Carvalho, R. F. (2018). Grafting in vegetable crops: A great technique for agriculture. *Int. J. Vegetable Sci.* 24, 85–102. doi: 10.1080/19315260.2017.1357062
- García-Lozano, M., Dutta, S. K., Natarajan, P., Tomason, Y. R., Lopez, C., Katam, R., et al. (2020). Transcriptome changes in reciprocal grafts involving watermelon and bottle gourd reveal molecular mechanisms involved in increase of the fruit size, rind toughness and soluble solids. *Plant Mol. Biol.* 102, 213–223. doi: 10.1007/s11103-019-00942-7
- García-Pérez, R. D., Van Houdt, H., and Depicker, A. (2004). Spreading of post-transcriptional gene silencing along the target gene promotes systemic silencing. *Plant J.* 38, 594–602. doi: 10.1111/j.1365-313X.2004.02067.x
- Gardiner, L.-J., Joynson, R., Omony, J., Rusholme-Pilcher, R., Olohan, L., Lang, D., et al. (2018). Hidden variation in polyploid wheat drives local adaptation. *Genome Res.* 28, 1319–1332. doi: 10.1101/gr.233551.117
- Goldschmidt, E. E. (2014). Plant grafting: new mechanisms, evolutionary implications. *Front. Plant Sci.* 5, 727. doi: 10.3389/fpls.2014.00727
- Gómez, G., and Pallás, V. (2004). A long-distance translocatable phloem protein from cucumber forms a ribonucleoprotein complex *In vivo* with hop stunt viroid RNA. *J. Virol.* 78, 10104–10110. doi: 10.1128/JVI.78.18.10104-10110.2004
- Ham, B.-K., Bandom, J. L., Ringgold, V., Lough, T. J., and Lucas, W. J. (2009). A polypyrimidine tract binding protein, pumpkin RBP50, forms the basis of a phloem-mobile ribonucleoprotein complex. *Plant Cell* 21, 197–215. doi: 10.1105/tpc.108.061317
- Ham, B.-K., Li, G., Jia, W., Leary, J. A., and Lucas, W. J. (2014). Systemic delivery of siRNA in pumpkin by a plant PHLOEM SMALL RNA-BINDING PROTEIN 1-ribonucleoprotein complex. *Plant J.* 80, 683–694. doi: 10.1111/tpj.12662
- Hamilton, A. J., and Baulcombe, D. C. (1999). A species of small antisense RNA in posttranscriptional gene silencing in plants. *Science* 286, 950–952. doi: 10.1126/science.286.5441.950
- Harada, T. (2010). Grafting and RNA transport *via* phloem tissue in horticultural plants. *Scientia Hort.* 125, 545–550. doi: 10.1016/j.scienta.2010.05.013
- Haroldsen, V., Szczerba, M. W., Aktas, H., Lopez, J., Odias, M. J., Chi-Ham, C. L., et al. (2012). Mobility of transgenic nucleic acids and proteins within grafted rootstocks for agricultural improvement. *Front. Plant Sci.* 3, 39. doi: 10.3389/fpls.2012.00039
- Haywood, V., Yu, T. S., Huang, N. C., and Lucas, W. J. (2005). Phloem long-distance trafficking of GIBBERELLIC ACID-INSENSITIVE RNA regulates leaf development. *Plant J.* 42, 49–68. doi: 10.1111/j.1365-313X.2005.02351.x
- Hewezi, T., Alibert, G., and Kallerhoff, J. (2005). Local infiltration of high- and low-molecular-weight RNA from silenced sunflower (*Helianthus annuus* L.) plants triggers post-transcriptional gene silencing in non-silenced plants. *Plant Biotechnol. J.* 3, 81–89. doi: 10.1111/j.1467-7652.2004.00103.x
- Hu, C., Ham, B. K., El-Shabrawi, H. M., Alexander, D., Zhang, D., Ryals, J., et al. (2016). Proteomics and metabolomics analyses reveal the cucurbit sieve tube system as a complex metabolic space. *Plant J.* 87, 442–454. doi: 10.1111/tpj.13209
- Huang, P., Huang, H., Lin, X., Liu, P., Zhao, L., Nie, W.-F., et al. (2021). MSI4/FVE is required for accumulation of 24-nt siRNAs and DNA methylation at a subset of target regions of RNA-directed DNA methylation. *Plant J.* 108, 347–357. doi: 10.1111/tpj.15441
- Jeffrey, C., and Yeoman, M. (1983). Development of intercellular connections between opposing cells in a graft union. *New Phytol.* 93, 491–509. doi: 10.1111/j.1469-8137.1983.tb02701.x
- Jones, L., Hamilton, A. J., Voinnet, O., Thomas, C. L., Maule, A. J., and Baulcombe, D. C. (1999). RNA-DNA Interactions and DNA methylation in post-transcriptional gene silencing. *Plant Cell* 11, 2291–2301. doi: 10.1105/tpc.11.12.2291
- Kamthan, A., Chaudhuri, A., Kamthan, M., and Datta, A. (2015). Small RNAs in plants: recent development and application for crop improvement. *Front. Plant Sci.* 6, 208. doi: 10.3389/fpls.2015.00208
- Karimi, H., Baldrich, P., Rutter, B. D., Borniego, L., Zajt, K. K., and Meyers, (2022). Arabidopsis apoplastic fluid contains sRNA- and circular RNA-protein complexes that are located outside extracellular vesicles. *Plant Cell* 34, 1863–1881. doi: 10.1093/plcell/koac043
- Kasai, A., Bai, S., Hojo, H., and Harada, T. (2016). Epigenome editing of potato by grafting using transgenic tobacco as siRNA donor. *PLoS One* 11, e0161729. doi: 10.1371/journal.pone.0161729
- Kasai, A., Bai, S., Li, T., and Harada, T. (2011). Graft-transmitted siRNA signal from the root induces visual manifestation of endogenous post-transcriptional gene silencing in the scion. *PLoS One* 6 (2), e16895. doi: 10.1371/journal.pone.0016895
- Kasai, A., Kanehira, A., and Harada, T. (2010). miR172 can move long distances in *nicotiana benthamiana*. *Open Plant Sci. J.* 4, 1–6. doi: 10.2174/1874294701004010001
- Kehr, J., and Kragler, F. (2018). Long distance RNA movement. *New Phytol.* 218, 29–40. doi: 10.1111/nph.15025
- Kim, M., Canio, W., Kessler, S., and Sinha, N. (2001). Developmental changes due to long-distance movement of a homeobox fusion transcript in tomato. *Science* 293, 287–289. doi: 10.1126/science.1059805
- Kim, G., Leblanc, M. L., Wafula, E. K., Depamphilis, C. W., and Westwood, J. H. (2014). Genomic-scale exchange of mRNA between a parasitic plant and its hosts. *Science* 345, 808–811. doi: 10.1126/science.1253122
- King, S. R., Davis, A. R., Zhang, X., and Crosby, K. (2010). Genetics, breeding and selection of rootstocks for solanaceae and cucurbitaceae. *Scientia Hort.* 127, 106–111. doi: 10.1016/j.scienta.2010.08.001
- Kitagawa, M., Xu, X., and Jackson, D. (2022). Trafficking and localization of KNOTTED1 related mRNAs in shoot meristems. *Communicative Integr. Biol.* 15, 158–163. doi: 10.1080/19420889.2022.2095125
- Kollmann, R., and Glockmann, C. (1985). Studies on graft unions. i. plasmodesmata between cells of plants belonging to different unrelated taxa. *Protoplasma* 124, 224–235. doi: 10.1007/BF01290774
- Kumari, A., Kumar, J., Kumar, A., Chaudhury, A., and Singh, S. P. (2015). Grafting triggers differential responses between scion and rootstock. *PLoS One* 10, e0124438. doi: 10.1371/journal.pone.0124438
- Kundariya, H., Yang, X., Morton, K., Sanchez, R., Axtell, M. J., Hutton, S. F., et al. (2020). MSH1-induced heritable enhanced growth vigor through grafting is associated with the RdDM pathway in plants. *Nat. Commun.* 11, 1–14. doi: 10.1038/s41467-020-19140-x
- Lang, Z., Wang, Y., Tang, K., Tang, D., Datsenko, T., Cheng, J., et al. (2017). Critical roles of DNA demethylation in the activation of ripening-induced genes and inhibition of ripening-repressed genes in tomato fruit. *Proc. Natl. Acad. Sci.* 114, E4511–E4519. doi: 10.1073/pnas.1705233114
- Lee, J.-M., and Oda, M. (2002). Grafting of herbaceous vegetable and ornamental crops. *In Horticultural Reviews (John Wiley and Sons, Ltd)* 28, 61–124. doi: 10.1002/9780470650851.ch2
- Lewsey, M. G., Hardcastle, T. J., Melnyk, C. W., Molnar, A., Valli, A., Urich, M. A., et al. (2016). Mobile small RNAs regulate genome-wide DNA methylation. *Proc. Natl. Acad. Sci.* 113, E801–E810. doi: 10.1073/pnas.1515072113
- Li, S., Wang, X., Xu, W., Liu, T., Cai, C., Chen, L., et al. (2021). Unidirectional movement of small RNAs from shoots to roots in interspecific heterografts. *Nat. Plants* 7, 50–59. doi: 10.1038/s41477-020-00829-2
- Li, Y., and Zhao, D. (2021). Transcriptome analysis of scions grafted to potato rootstock for improving late blight resistance. *BMC Plant Biol.* 21, 1–13. doi: 10.1186/s12870-021-03039-w
- Liu, L., and Chen, X. (2018). Intercellular and systemic trafficking of RNAs in plants. *Nat. Plants* 4, 869–878. doi: 10.1038/s41477-018-0288-5

- Liu, C., Lu, F., Cui, X., and Cao, X. (2010). Histone methylation in higher plants. *Annu. Rev. Plant Biol.* 61, 395–420. doi: 10.1146/annurev.arplant.043008.091939
- Liu, N., Yang, J., Fu, X., Zhang, L., Tang, K., Guy, K. M., et al. (2016). Genome-wide identification and comparative analysis of grafting-responsive mRNA in watermelon grafted onto bottle gourd and squash rootstocks by high-throughput sequencing. *Mol. Genet. Genomics* 291, 621–633. doi: 10.1007/s00438-015-1132-5
- Lough, T. J., and Lucas, W. J. (2006). Integrative plant biology: role of phloem long-distance macromolecular trafficking. *Annu. Rev. Plant Biol.* 57, 203–232. doi: 10.1146/annurev.arplant.56.032604.144145
- Lucas, W. J., Yoo, B.-C., and Kragler, F. (2001). RNA As a long-distance information macromolecule in plants. *Nat. Rev. Mol. Cell Biol.* 2, 849–857. doi: 10.1038/35099096
- Luo, K.-R., Huang, N.-C., Chang, Y.-H., and Yu, T.-S. (2022). *Arabidopsis cyclophilins direct plasmodesmata-targeting of mobile mRNA via organelle hitchhiking*. Available at: 10.21203/rs.3.rs-1088339/v1.
- Luo, K.-R., Huang, N.-C., and Yu, T.-S. (2018). Selective targeting of mobile mRNAs to plasmodesmata for cell-to-cell movement. *Plant Physiol.* 177, 604–614. doi: 10.1104/pp.18.00107
- Magome, H., Yamaguchi, S., Hanada, A., Kamiya, Y., and Oda, K. (2004). Dwarf and delayed-flowering 1, a novel arabidopsis mutant deficient in gibberellin biosynthesis because of overexpression of a putative AP2 transcription factor. *Plant J.* 37, 720–729. doi: 10.1111/j.1365-313X.2003.01998.x
- Mahajan, A., Bhogale, S., Kang, I. H., Hannapel, D. J., and Banerjee, A. K. (2012). The mRNA of a Knotted1-like transcription factor of potato is phloem mobile. *Plant Mol. Biol.* 79, 595–608. doi: 10.1007/s11103-012-9931-0
- Mallory, A. C., Ely, L., Smith, T. H., Marathe, R., Anandalakshmi, R., Fagard, M., et al. (2001). Hc-pro suppression of transgene silencing eliminates the small RNAs but not transgene methylation or the mobile signal. *Plant Cell* 13, 571–583. doi: 10.1105/tpc.13.3.571
- Mallory, A. C., Mlotshwa, S., Bowman, L. H., and Vance, V. B. (2003). The capacity of transgenic tobacco to send a systemic RNA silencing signal depends on the nature of the inducing transgene locus. *Plant J.* 35, 82–92. doi: 10.1046/j.1365-313X.2003.01785.x
- Manning, K., Tör, M., Poole, M., Hong, Y., Thompson, A. J., King, G. J., et al. (2006). A naturally occurring epigenetic mutation in a gene encoding an SBP-box transcription factor inhibits tomato fruit ripening. *Nat. Genet.* 38, 948–952. doi: 10.1038/ng1841
- Marjori, M., Antonius, J., and Jan, M. K. (2001). RNA: guiding gene silencing. *Science* 293, 1080–1083. doi: 10.1126/science.1063051
- Melnyk, C. W., Schuster, C., Leyser, O., and Meyerowitz, E. M. (2015). A developmental framework for graft formation and vascular reconnection in arabidopsis thaliana. *Curr. Biol.* 25, 1306–1318. doi: 10.1016/j.cub.2015.03.032
- Mette, M. F., van der Winden, J., Matzke, M. A., and Matzke, A. J. M. (1999). Production of aberrant promoter transcripts contributes to methylation and silencing of unlinked homologous promoters in trans. *EMBO J.* 18, 241–248. doi: 10.1093/emboj/18.1.241
- Molnar, A., Melnyk, C. W., Bassett, A., Hardcastle, T. J., Dunn, R., and Baulcombe, D. C. (2010). Small silencing RNAs in plants are mobile and direct epigenetic modification in recipient cells. *Science* 328, 872–875. doi: 10.1126/science.1187959
- Morel, J.-B., Mourrain, P., Béclin, C., and Vaucheret, H. (2000). DNA Methylation and chromatin structure affect transcriptional and post-transcriptional transgene silencing in arabidopsis. *Struct. Biol.* 10, 1591–1594. doi: 10.1016/S0960-9822(00)00862-9
- Mudge, K., Janick, J., Scofield, S., and Goldschmidt, E. E. (2009). A history of grafting. In *Horticultural Reviews (John Wiley and Sons, Ltd)*, 473–493. doi: 10.1002/9780470593776.ch9
- Noshay, J. M., and Springer, N. M. (2021). Stories that can't be told by SNPs: DNA methylation variation in plant populations. *Curr. Opin. Plant Biol.* 61, 101989. doi: 10.1016/j.pbi.2020.101989
- Notaguchi, M., Higashiyama, T., and Suzuki, T. (2015). Identification of mRNAs that move over long distances using an RNA-seq analysis of Arabidopsis/Nicotiana benthamiana heterografts. *Plant Cell Physiol.* 56, 311–321. doi: 10.1093/pcp/pcu210
- Notaguchi, M., Wolf, S., and Lucas, W. J. (2012). Phloem-mobile Aux/IAA transcripts target to the root tip and modify root architecture. *J. Integr. Plant Biol.* 54, 760–772. doi: 10.1111/j.1744-7909.2012.01155.x
- Ntatsi, G., Savvas, D., Papisotiropoulos, V., Katsileros, A., Zrenner, R. M., Hinch, D. K., et al. (2017). Rootstock sub-optimal temperature tolerance determines transcriptomic responses after long-term root cooling in rootstocks and scions of grafted tomato plants. *Front. Plant Sci.* 8, 911. doi: 10.3389/fpls.2017.00911
- Omid, A., Keilin, T., Glass, A., Leshkowitz, D., and Wolf, S. (2007). Characterization of phloem-sap transcription profile in melon plants. *J. Exp. Bot.* 58, 3645–3656. doi: 10.1093/jxb/erm214
- Palauqui, J.-C., Elmayan, T., Pollien, J.-M., and Vaucheret, H. (1997). Systemic acquired silencing: transgene-specific post-transcriptional silencing is transmitted by grafting from silenced stocks to non-silenced scions. *EMBO J.* 16, 4738–4745. doi: 10.1093/emboj/16.15.4738
- Pang, T. Y., Short, A. K., Bredy, T. W., and Hannan, A. J. (2017). Transgenerational paternal transmission of acquired traits: stress-induced modification of the sperm regulatory transcriptome and offspring phenotypes. *Curr. Opin. Behav. Sci.* 14, 140–147. doi: 10.1016/j.cobeha.2017.02.007
- Pant, B. D., Buhtz, A., Kehr, J., and Scheible, W. R. (2008). MicroRNA399 is a long-distance signal for the regulation of plant phosphate homeostasis. *Plant J.* 53, 731–738. doi: 10.1111/j.1365-313X.2007.03363.x
- Pfluger, J., and Wagner, D. (2007). Histone modifications and dynamic regulation of genome accessibility in plants. *Curr. Opin. Plant Biol.* 10, 645–652. doi: 10.1016/j.pbi.2007.07.013
- Pouget, R. (1990). Histoire de la lutte contre le phylloxéra de la vigne en France. *Institut Natl. la Recherche Agronomique (in French)*, 1–168. Available at: <http://digital.casalini.it/9782759205875>.
- Qi, Y., Peëliissier, T., Itaya, A., Hunt, E., Wassenegger, M., and Ding, B. (2004). Direct role of a viroid RNA motif in mediating directional RNA trafficking across a specific cellular boundary. *Plant Cell* 16, 1741–1752. doi: 10.1105/tpc.021980
- Raju, S. K. K., Shao, M. R., Sanchez, R., Xu, Y. Z., Sandhu, A., Graef, G., et al. (2018). An epigenetic breeding system in soybean for increased yield and stability. *Plant Biotechnol. J.* 16, 1836–1847. doi: 10.1111/pbi.12919
- Rapp, R. A., and Wendel, J. F. (2005). Epigenetics and plant evolution. *New Phytol.* 168, 81–91. doi: 10.1111/j.1469-8137.2005.01491.x
- Robertson, K. D. (2005). DNA Methylation and human disease. *Nat. Rev. Genet.* 6, 597–610. doi: 10.1038/nrg1655
- Rouphael, Y., Cardarelli, M., Colla, G., and Rea, E. (2008). Yield, mineral composition, water relations, and water use efficiency of grafted mini-watermelon plants under deficit irrigation. *HortScience* 43, 730–736. doi: 10.21273/HORTSCI.43.3.730
- Rubio, B., Stammitti, L., Cookson, S. J., Teyssier, E., and Gallucci, P. (2022). Small RNA populations reflect the complex dialogue established between heterograft partners in grapevine. *Horticulture Res.* 9, uhab067. doi: 10.1093/hr/uhab067
- Saze, H., Tsugane, K., Kanno, T., and Nishimura, T. (2012). DNA Methylation in plants: relationship to small RNAs and histone modifications, and functions in transposon inactivation. *Plant Cell Physiol.* 53 (5), 766–784. doi: 10.1093/pcp/pcs008
- Sharma, P., Lin, T., Grandellis, C., Yu, M., and Hannapel, D. J. (2014). The BEL1-like family of transcription factors in potato. *J. Exp. Bot.* 65, 709–723. doi: 10.1093/jxb/ert432
- Singh, A., Gautam, V., Singh, S., Sarkar Das, S., Verma, S., Mishra, V., et al. (2018). Plant small RNAs: advancement in the understanding of biogenesis and role in plant development. *Planta* 248, 545–558. doi: 10.1007/s00425-018-2927-5
- Song, J.-J., Liu, J., Tolia, N. H., Schneidman, J., Smith, S. K., Martienssen, R. A., et al. (2003). The crystal structure of the Argonaute2 PAZ domain reveals an RNA binding motif in RNAi effector complexes. *Nat. Struct. Mol. Biol.* 10, 1026–1032. doi: 10.1038/nsb1016
- Sonoda, S., and Nishiguchi, M. (2000). Graft transmission of post-transcriptional gene silencing: Target specificity for RNA degradation is transmissible between silenced and non-silenced plants, but not between silenced plants. *Plant J.* 21, 1–8. doi: 10.1046/j.1365-313x.2000.00645.x
- Spanò, R., Ferrara, M., Montemurro, C., Mulè, G., Gallitelli, D., and Mascia, T. (2020). Grafting alters tomato transcriptome and enhances tolerance to an airborne virus infection. *Sci. Rep.* 10, 1–13. doi: 10.1038/s41598-020-59421-5
- Spanò, R., Gallitelli, D., and Mascia, T. (2017). Grafting to manage infections of top stunting and necrogenic strains of cucumber mosaic virus in tomato. *Ann. Appl. Biol.* 171, 393–404. doi: 10.1111/aab.12382
- Spanò, R., Mascia, T., Kormelink, R., and Gallitelli, D. (2015). Grafting on a non-transgenic tolerant tomato variety confers resistance to the infection of a Sw5-breaking strain of tomato spotted wilt virus via RNA silencing. *PLoS One* 10, e0141319. doi: 10.1371/journal.pone.0141319
- Spiegelman, Z., Golan, G., and Wolf, S. (2013). Don't kill the messenger: long-distance trafficking of mRNA molecules. *Plant Sci.* 213, 1–8. doi: 10.1016/j.plantsci.2013.08.011
- Takeda, R., Zirbel, C. L., Leontis, N. B., Wang, Y., and Ding, B. (2018). Allelic RNA motifs in regulating systemic trafficking of potato spindle tuber viroid. *Viruses* 10, 160. doi: 10.3390/v10040160
- Tan, H., Li, B., and Guo, H. (2020). The diversity of post-transcriptional gene silencing mediated by small silencing RNAs in plants. *Essays Biochem.* 64, 919–930. doi: 10.1042/EBC20200006
- Taochy, C., Gursansky, N. R., Cao, J., Fletcher, S. J., Dressel, U., Mitter, N., et al. (2017). A genetic screen for impaired systemic RNAi highlights the crucial role of DICER-LIKE 2. *Plant Physiol.* 175, 1424–1437. doi: 10.1104/pp.17.01181
- Tateishi, K. (1927). Grafting watermelon onto pumpkin. *J. Jpn. Hortic.* 39, 5–8.
- Thieme, C., Rojas-Triana, M., Stecyk, E., Schudoma, C., Zhang, W., Yang, L., et al. (2015). Endogenous arabidopsis messenger RNAs transported to distant tissues. *Nat. Plants* 1, 15025. doi: 10.1038/nplants.2015.25
- Tomkins, M., Hoerbst, F., Gupta, S., Apelt, F., Kehr, J., Kragler, F., et al. (2022). Exact Bayesian inference for the detection of graft-mobile transcripts from sequencing data. *J. R. Soc. Interface* 19, 20220644. doi: 10.1098/rsif.2022.0644
- Tournier, B., Tabler, M., and Kalantidis, K. (2006). Phloem flow strongly influences the systemic spread of silencing in GFP nicotiana benthamiana plants. *Plant J.* 47, 383–394. doi: 10.1111/j.1365-313X.2006.02796.x
- Tsutsui, H., and Notaguchi, M. (2017). The use of grafting to study systemic signaling in plants. *Plant Cell Physiol.* 58, 1291–1301. doi: 10.1093/pcp/pcx098
- Turnbull, C. G., and Lopez-Cobollo, R. M. (2013). Heavy traffic in the fast lane: long-distance signalling by macromolecules. *New Phytol.* 198, 33–51. doi: 10.1111/nph.12167
- Van Otterdijk, S. D., and Michels, K. B. (2016). Transgenerational epigenetic inheritance in mammals: how good is the evidence? *FASEB J.* 30, 2457–2465. doi: 10.1096/fj.201500083

- Voinnet, O. (2022). Revisiting small RNA movement in plants. *Nat. Rev. Mol. Cell Biol.* 23, 163–164. doi: 10.1038/s41580-022-00455-0
- Voinnet, O., Lederer, C., and Baulcombe, D. C. (2000). A viral movement protein prevents spread of the gene silencing signal in *Nicotiana benthamiana*. *Cell* 103, 157–167. doi: 10.1016/S0092-8674(00)00095-7
- Wada, Y., Miyamoto, K., Kusano, T., and Sano, H. (2004). Association between up-regulation of stress-responsive genes and hypomethylation of genomic DNA in tobacco plants. *Mol. Genet. Genomics* 271, 658–666. doi: 10.1007/s00438-004-1018-4
- Waddington, C. H. (1956). Genetic assimilation of the bithorax phenotype. *Evolution* 10, 1–13. doi: 10.2307/2406091
- Wang, T., Li, X., Zhang, X., Wang, Q., Liu, W., Lu, X., et al. (2021). RNA Motifs and modification involve in RNA long-distance transport in plants. *Front. Cell Dev. Biol.* 9, 651278. doi: 10.3389/fcell.2021.651278
- Wang, S.-S., Liu, Z.-Z., Sun, C., Shi, Q.-H., Yao, Y.-X., You, C.-X., et al. (2012). Functional characterization of the apple MhGAI1 gene through ectopic expression and grafting experiments in tomatoes. *J. Plant Physiol.* 169, 303–310. doi: 10.1016/j.jplph.2011.09.012
- Wang, H., Zhou, P., Zhu, W., and Wang, F. (2019). *De novo* comparative transcriptome analysis of genes differentially expressed in the scion of homografted and heterografted tomato seedlings. *Sci. Rep.* 9, 1–12. doi: 10.1038/s41598-019-56563-z
- Weigel, D., and Colot, V. (2012). Epialleles in plant evolution. *Genome Biol.* 13, 1–6. doi: 10.1186/gb-2012-13-10-249
- Wu, R., Wang, X., Lin, Y., Ma, Y., Liu, G., Yu, X., et al. (2013). Inter-species grafting caused extensive and heritable alterations of DNA methylation in solanaceae plants. *PLoS One* 8, e61995. doi: 10.1371/journal.pone.0061995
- Xie, M., and Yu, B. (2015). siRNA-directed DNA methylation in plants. *Curr. Genomics* 16, 23–31. doi: 10.2174/1389202915666141128002211
- Xu, J., Chen, G., Hermanson, P. J., Xu, Q., Sun, C., Chen, W., et al. (2019). Population-level analysis reveals the widespread occurrence and phenotypic consequence of DNA methylation variation not tagged by genetic variation in maize. *Genome Biol.* 20, 243. doi: 10.1186/s13059-019-1859-0
- Yan, Y., and Ham, B.-K. (2022). The mobile small RNAs: Important messengers for long-distance communication in plants. *Frontiers in Plant Science* 13. doi: 10.3389/fpls.2022.928729
- Yang, D., Huang, Z., Jin, W., Xia, P., Jia, Q., Yang, Z., et al. (2018). DNA Methylation: A new regulator of phenolic acids biosynthesis in *Salvia miltiorrhiza*. *Ind. Crops Products* 124, 402–411. doi: 10.1016/j.indcrop.2018.07.046
- Yang, L., Machin, F., Wang, S., Saploura, E., and Kragler, F. (2023). Heritable transgene-free genome editing in plants by grafting of wild-type shoots to transgenic donor rootstocks. *Nat. Biotechnol.* 1–10. doi: 10.1038/s41587-022-01585-8
- Yang, L., Perrera, V., Saploura, E., Apelt, F., Bahin, M., Kramdi, A., et al. (2019). m5C methylation guides systemic transport of messenger RNA over graft junctions in plants. *Curr. Biol.* 29, 2465–2476.e5. doi: 10.1016/j.cub.2019.06.042
- Zhang, W., Kollwig, G., Stecyk, E., Apelt, F., Dirks, R., and Kragler, F. (2014). Graft-transmissible movement of inverted-repeat-induced si RNA signals into flowers. *Plant J.* 80, 106–121. doi: 10.1111/tj.12622
- Zhang, X., Sun, J., Cao, X., and Song, X. (2015). Epigenetic mutation of RAV6 affects leaf angle and seed size in rice. *Plant Physiol.* 169, 2118–2128. doi: 10.1104/pp.15.00836
- Zhang, S., Sun, L., and Kragler, F. (2009). The phloem-delivered RNA pool contains small noncoding RNAs and interferes with translation. *Plant Physiol.* 150, 378–387. doi: 10.1104/pp.108.134767
- Zhang, W., Thieme, C. J., Kollwig, G., Apelt, F., Yang, L., Winter, N., et al. (2016). tRNA-related sequences trigger systemic mRNA transport in plants. *Plant Cell* 28, 1237–1249. doi: 10.1105/tpc.15.01056
- Zhang, H., Yu, P., Zhao, J., Jiang, H., Wang, H., Zhu, Y., et al. (2018a). Expression of tomato prosystemin gene in Arabidopsis reveals systemic translocation of its mRNA and confers necrotrophic fungal resistance. *New Phytol.* 217, 799–812. doi: 10.1111/nph.14858
- Zhang, J., Zhang, H., Srivastava, A. K., Pan, Y., Bai, J., Fang, J., et al. (2018b). Knockdown of rice MicroRNA166 confers drought resistance by causing leaf rolling and altering stem xylem development. *Plant Physiol.* 176, 2082–2094. doi: 10.1104/pp.17.01432
- Zhang, G., Zhou, J., Song, J., Guo, X., Nie, X., and Guo, H. (2022). Grafting-induced transcriptome changes and long-distance mRNA movement in the potato/*Datura stramonium* heterograft system. *Horticulture Environment Biotechnol.* 63, 229–238. doi: 10.1007/s13580-021-00387-2
- Zhong, X., Archual, A. J., Amin, A. A., and Ding, B. (2008). A genomic map of viroid RNA motifs critical for replication and systemic trafficking. *Plant Cell* 20, 35–47. doi: 10.1105/tpc.107.056606
- Zhong, X., Tao, X., Stombaugh, J., Leontis, N., and Ding, B. (2007). Tertiary structure and function of an RNA motif required for plant vascular entry to initiate systemic trafficking. *EMBO J.* 26, 3836–3846. doi: 10.1038/sj.emboj.7601812