# **UNIVERSITY** OF BIRMINGHAM University of Birmingham Research at Birmingham

# SOPTI Meeting Abstracts 2021

Recchioni, Alberto

DOI: 10.5384/sjovs.v14i2.149

License: Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

**Document Version** Publisher's PDF, also known as Version of record

Citation for published version (Harvard): Recchioni, A 2021, 'SOPTI Meeting Abstracts 2021', Scandinavian Journal of Optometry and Visual Science, vol. 14, no. 2. https://doi.org/10.5384/sjovs.v14i2.149

Link to publication on Research at Birmingham portal

#### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research. •User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)

•Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

# **SOPTI Meeting 2021: Abstracts**

After more than a year of blockade due to the Covid-19 pandemic, it was finally possible to return to the events in the presence. The 26th National Conference of the Italian Optometric Association (SOPTI) was held in Bologna on October 10–11, 2021.

The theme of the conference was "Good practice in Optometry and Contact Lenses", with the accent on two topics: the optometric management of the patient in old age and the progression of myopia.

Four keynote speakers were invited during the conference: Prof. Rigmor C. Baraas from the University of South-Eastern Norway in Kongsberg, Prof. David B. Elliot from the University of Bradford, Dr. Fabrizio Zeri from the University of Milano Bicocca and the IACLE President, Prof. Phil Morgan, from the University of Manchester.

The abstracts from accepted posters and free papers are presented here.

Received September 21, 2021 . Accepted November 8, 2021.

© Copyright abstract authors. This article is distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use and redistribution provided that the original author and source are credited.

# Corneal densitometry by sublayers: an alternative method for analyzing Scheimpflug images of normal or keratoconic eyes

Piermattia Gazzera, Francesca Carpi, Mauro Frisani, Michela Greco

Physics department, University of Optics and Optometry, Turin, Italy

Corresponding author: piergazzera@gmail.com

## Abstract

Loss of corneal transparency is a key element to monitor in ophthalmic practice, as it may be symptomatic of a wide range of conditions including corneal dystrophies. The most common is keratoconus, a non-inflammatory corneal ectasia characterized by a progressive thinning of corneal tissue and formation of corneal protrusion. Densitometric analysis has recently been introduced. It is an objective and non-invasive technique capable of quantifying corneal transparency with the aid of various imaging technologies including Scheimpflug photography. Pentacam Scheimpflug, the most used instrument, calculates density of corneal layers respecting fixed depth values. This work proposes an alternative method based on segmentation of the epithelium and stroma starting from an image obtained with Sirius Tomograph (CSO, Italy).

180 images of the anterior segment of 30 subjects  $(24.1 \pm 4.4 \text{ years})$  with healthy eyes and 50 images of eyes belonging to 22 subjects  $(28.8 \pm 8.4 \text{ years})$  with unilateral keratoconus were analyzed. These were acquired with the Sirius tomograph (CSO, Italy). The stroma and the epithelium associated with Bowman's membrane were segmented from each tomography corresponding to the horizontal meridian. Only the central 3-mm-diameter zone was considered. The computing platform used was MATLAB. In the process, the identification of the corneal apex is obtained by the apex.m function. It requires the coordinates of two points respectively to the left (L) and to the right (R) of the apex. On the other hand, the function used for the segmentation in the different layers of the corneal thickness was region3mm.m which exploits the region-growing technique.

Student's t-test showed that the subjective choice of points L and R does not determine significant changes in the identification of the apical coordinates. The thresholds used to segment are expressed in a grayscale from 1 (white) to 256 (black). They were 18 and 41 for the stroma, 41 and 63 for the epithelium. The segmentations obtained, introducing variations of  $\pm$  5 units to the thresholds, were classified into *adequate*, *almost adequate* and *inadequate* both for healthy and keratoconus eyes. Subsequently, the densitometric values or the stroma and epithelium were calculated by adopting two different criteria: the first referring to the extremes of the peaks identified by the intensity distribution of the pixels, the second referring to the thresholds.

The segmentations are adequate in healthy corneas with a success rate above 80%. The application of the method to corneas affected by keratoconus shows, as expected, a lower success rate around 50% (figure 1). The reliability of the apex.m algorithm is demonstrated. The region3mm.m function is more suitable for healthy corneas than corneas with keratoconus.

The densitometric results were converted to GSU (figure 1) and compared with the values reported in the studies by Dhubhghaill et al. (Ní Dhubhghaill et al., 2014) and Tekin et al. (Tekin et al., 2017) obtained from Pentacam Scheimpflug. The agreement with Tekin's study is the best among those considered, but not statistically significant. The comparison is weak because, with the Pentacam tomograph, corneal segments are evaluated for depth without recognition of different substrates.

Normal eyes (N=180)		stroma		epithelium	
		extreme s	threshold s	extremes	thresholds
adequate	mean value	11,3	11,2	18,9	18,7
	SD	0,9	0,5	1,4	0,7
	%	92%		79%	
almost adequate	mean value	12,1	11,7	20,5	19,4
	SD	0,5	0,3	1,2	0,6
	%	3%		4%	
inadequate	mean value	11,9	11,6	20,0	18,9
	SD	1,3	0,6	2,4	1,0
	%	5	5%	16%	
Normal eyes of monolateral KC (N=25)		stroma		epithelium	
		extreme s	threshold s	extremes	thresholds
	mean value	11,8	11,4	19,3	18,8
adequate	SD	1,0	0,5	1,3	0,7
	%	64%		64%	
almost adequate	mean value	11,9	11,7		
	SD	0,3	0,3		
	%	8%			
inadequate	mean value	10	10,3	21,2	19,3
	SD	2,8	1,5	6,2	0,5
	%	28%		36%	
Keratoconus eyes (N=25)		stroma		epithelium	
		extreme s	threshold s	extremes (N=22)	thresholds (N=25)
adequate	mean value	11,4	11,2	19,6	19,2
	SD	0,6	0,3	1,6	0,8
	%	52%		45%	40%
almost adequate	mean value	10,2	10,4		
	SD	1,3	0,9		
	%	20%			
inadequate	mean value	13,0	12,0	20,9	19,6
	SD	1,0	0,3	3,2	0,3
	%	28%		55%	60%

Figure 1: all densitometric values are reported in GSU and divided into the analyzed samples. The percentages obtained from the classification of segmentations in adequate, almost adequate and inadequate are also reported.

### References

Ní Dhubhghaill, S., Rozema, J. J., Jongenelen, S., Ruiz Hidalgo, I., Zakaria, N., & Tassignon, M.-J. (2014). Normative Values for Corneal Densitometry Analysis by Scheimpflug Optical Assessment. *Investigative Ophthalmology & Visual Science*, *55*(1), 162–168. https://doi.org/10.1167/iovs.13-13236

Tekin, K., Sekeroglu, M. A., Kiziltoprak, H., & & Yilmazbas, P. (2017). Corneal densitometry in healthy corneas and its correlation with endothelial morphometry. *Cornea*, *36*(11), 1336–1342. https://doi.org/10.1097/ICO.000000000001363

# Subjective measurement of fusional vergences, dissociated heterophorias and associated heterophorias: comparison between monocular and binocular prismatic apposition

Martina Gelain, Claudia Colandrea, Silvio Maffioletti, Marina Serio

Physics department, University of Optics and Optometry, Turin, Italy

Corresponding author: mg.martinagelain@gmail.com

## Abstract

The aim of the study is the measure of fusional vergences, dissociated heterophorias and associated heterophorias both in the phoropter and in the open field, comparing the data obtained when the prisms are placed in front of only one eye or on both eyes. Most of the studies in the literature, in fact, compare the various measurement methods but do not make a direct comparison between monocular and binocular prismatic apposition.

The study sample included 40 subjects (age  $22.5\pm3.1$ ) were examined in which the lenses of the subjective refraction performed at the phoropter were worn during the tests, then balanced with Humpriss in the open field. For each subject, the following were measured from afar (6 metres) and near (40 centimeters): the smooth fusional vergences (phoropter) and jumps (open field), the dissociated heterophorias by means of the von Graefe test (phoropter) and the of Maddox modified Thoringhton (open field) and finally, the heterophoria associated for distance, through the cross and needle test both to the phoropter and in open field and the associated phrases for near through the Wesson card (phoropter) and the Mallet Unit (open field). Wilconxon's non-parametric test was applied to study the comparison between the samples, considering a significance level of 0.05 (Figure 1).

		LEGEND
COMPARISONS	p-value	V: Varganaag
Vs FNEBMO – Vs FNEBBI	0,023	J: Jumps (open field)
Vs NNEREMO – Vs NNEREBI	0,003	S: Smooth (phoropter) F: Far
Vs NPOBMO – Vs NPOBBI	0,002	N: Near
Vs NPOREMO – Vs NPOREBI	0,041	NE: Negative
VJ FNEBMO – VJ FNEBBI	0,001	MO: Monocular prismatic apposition BI: Binocular prismatic apposition
VJ FNEREMO – VJ FNEREBI	0,012	B: Break
VJ NNEREMO – VJ NNEREBI	0,004	RE: Recovery OF: Open field
V <sub>S</sub> FPOREMO- V <sub>J</sub> FPOREMO	4*10^-4	CF: Closed field
Vs NPOBMO- V1 NPOBMO	0,001	
V5 FPOREBI- VJ FPOREBI	0,002	
V <sub>S</sub> NNEREBI- V <sub>J</sub> NPOREBI	2*10^-4	
MADDOX <sub>MOF</sub> VS MADDOX <sub>BIF</sub>	0,0455	
MADDOX MON VS MADDOXBIN	0,0062	
MALLET <sub>MON</sub> in OF VS MALLET <sub>BIN</sub> in OF	0,001	
CROSS TEST <sub>MOF</sub> in CF VS CROSS TEST <sub>MOF</sub> in OF	0,021	
NEEDLE TEST <sub>MOF</sub> in CF VS CROSS TEST <sub>MOF</sub> IN OF	0,036	
WESSONMON in CF VS MALLETMON in CA	0,002	
CROSS TEST BIF in CF $VS$ CROSS TEST BIF in OF	0,003	
NEEDLE TESTBIF in CF VS NEEDLE TESTBIF in OF	0,019	

Figure 1: Comparisons that showed significant differences with Wilconxon's nonparametric test.

In the study, the difference in fusional vergences was greater especially on breakages and recoveries for the most part negative and for the least part positive. In dissociated heterophorias, significant differences appear only on the data in the open field, so there may be a different adaptation to the monocular prism compared to the binocular when the data is not taken at the phoropter. In the associated heterophoria, the comparison histograms showed a greater presence of exophoric subjects in a monocular condition and a greater presence of exophoric subjects in a binocular condition.

The only test that did not show significant difference in open and closed field was that of the cross, probably due to the absence of a central fusional recall. The study results show a new perspective on binocular vision investigation methods. In this regard, in order to have a more specific and broader understanding of the commonly used procedures, it would be useful to deepen the study with a larger sample both in terms of number and variety.

# Contrast sensitivity measures: comparison between professional and casual drivers

Alessandro Guidani, Claudia Colandrea, Michela Greco Physics department, University of Optics and Optometry, Turin, Italy

Corresponding author: alexsoccer@live.it

### Abstract

Given the lack of evidence in the procedures defining visual standards for driving licenses and the lack of understanding of how visual decay impacts on safety and driving performance, in the study we aim to have contrast sensitivity measurements, paying particular attention to sample definition. Therefore the aim of the study is to analyze the differences between contrast sensitivity (CS) measurements obtained on a population of professional drivers (GP) and one of occasional drivers (GNP) weighting the role of binocular vision.

The subjects considered for this study are 60, of which 30 GP and 30 GNP. For the CS measurements, the Pelli-Robson subjective test in digital format was used. Before proceeding with the measures, all subjects were given a questionnaire with an indication of age, gender, profession, visual experiences driving in low-contrast conditions, number of accidents in the last year and reported data on the state of health. During the measurements all subjects wore their usual correction. First of all, visual acuity was measured, then subjective measurement of contrast sensitivity was carried out with Pelli-Robson test; the evaluation ended when the subject did not recognize at least 2 letters of the triplet presented.

The statistical analysis is presented, which shows on average better values of AV and CS for the sample of professional drivers. The statistically significant differences between the two categories and the role of binocularity for the measurement of contrast sensitivity are highlighted.

# Use of comfort questionnaires in a sample of irregular cornea contact lens wearers

# Mario Panzieri

Institute for Research and Study in Optics and Optometry (IRSOO), Vinci, Italy Corresponding author: mariopanzieri75@gmail.com

#### Abstract

To evaluate the possibility of administering short item questionnaires created for regular cornea subjects and daily disposable contact lens wearers to irregular cornea subjects wearing any kind of contact lens (CL). To compare two questionnaires among them.

In this study, 67 subjects were involved, wearing any kind of contact lens (Rigid, Soft, Hybrid contact lenses): 32 of them presented regular corneas (RC) and 37 irregular corneas (IC). Monocular visual acuity (VA) with spectacles (when available) and monocular VA with CL were taken.

Two questionnaires were used: the Contact Lens Dry Eye Questionnaire (CLDEQ-8), made of 8 questions, and a new questionnaire from IRSOO (Istituto di Ricerca e di Studi in Ottica e Optometria), made of 7 questions and related to evaluating vision comfort in CL wearers. Subjects were habitual CL wearers and independently filled the questionnaires, referring to the previous two weeks' CL wearing experience.

The statistical analysis used for comparing answers to both questionnaires was the Pearson's correlation coefficient (P). Comparison among VA with the different types of CL was assessed, assuming a p-value  $\leq 0.05$  as statistically significant.

Results and conclusions: Both questionnaires show a high correlation (P=0.73 for CLDEQ-8 and P=0.74 for IRSOO-7), for the two subgroups also: myopes (My) and keratoconus (Kc). There is no significant statistical difference (*p-value*>0.05) between VA with Soft and Rigid CL and with Soft and Hybrid CL, while the difference of VA between Hybrid and Rigid CL is statistically significant (*p-value*<0.05). Also, VA in the main Groups (RC, IC) and subgroups (My, Kc) was better while wearing CL rather than spectacles (RC 0.05 LogMAR vs 0.06 LogMAR; IC 0.09 LogMAR vs 0.27 LogMAR; My 0.04 LogMAR vs 0.05; Kc 0.09 LogMAR vs 0.29).

Both questionnaires therefore can be used in a routinary clinical practice to evaluate the use of a CL fitting quality over time or to compare the performance among different types of CL.