

## A Sticky Situation

Sindhi, Rithik Mohan Singh; Win, Kyaw Z.; Zafar, Athesham; Steeds, Richard P.

DOI:

[10.1016/j.jaccas.2023.101886](https://doi.org/10.1016/j.jaccas.2023.101886)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Sindhi, RMS, Win, KZ, Zafar, A & Steeds, RP 2023, 'A Sticky Situation: An Uncommon Cardiac Complication of Cyanoacrylate Glue Injection', *JACC: Case Reports*, vol. 17, 101886.  
<https://doi.org/10.1016/j.jaccas.2023.101886>

[Link to publication on Research at Birmingham portal](#)

### General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

CASE REPORT

ADVANCED

CLINICAL CASE

# A Sticky Situation

## An Uncommon Cardiac Complication of Cyanoacrylate Glue Injection



Rithik Mohan Singh Sindhi, MD,<sup>a</sup> Kyaw Z. Win, MBBS,<sup>a,b</sup> Athesham Zafar, MBBS,<sup>c</sup> Richard P. Steeds, MA, MD<sup>a,b</sup>

### ABSTRACT

This case report describes an uncommon cardiac complication related to the endoscopic injection of cyanoacrylate glue to treat gastric variceal bleeding. The presentation and management options are reviewed and compared with our step-by-step approach in a patient requiring a liver transplant. (**Level of Difficulty: Advanced.**) (J Am Coll Cardiol Case Rep 2023;17:101886) © 2023 The Authors. Published by Elsevier on behalf of the American College of Cardiology Foundation. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**A** 28-year-old man with iron deficiency anemia of unknown cause was referred to the local hospital for elective esophagogastroduodenoscopy (EGD). Before this episode, the patient's general practitioner had investigated him for iron deficiency anemia with intermittent diarrhea and rectal bleeding in the community. The computed

tomography (CT) scan revealed a cirrhotic appearance of the liver, with splenomegaly and gastric varices. During EGD, the endoscopist found lesions consistent with gastric varices at the gastric fundus and proceeded to treat them with glue injection consisting of 1.5 mL Histoacryl mixed with lipiodol. This unexpectedly resulted in an exsanguinating hemorrhage, requiring admission to the intensive care unit after resuscitation to achieve hemodynamic stability.

### LEARNING OBJECTIVES

- To recognize that TIPS can facilitate Histoacryl glue extension and embolization, complicating further management.
- To anticipate embolic complication of glue injection because the presentation can be acute, subacute, or delayed.
- To consider a multidisciplinary approach with multimodality imaging as early as possible to deliver the safest treatment option to the patient with embolic complications.

### MEDICAL HISTORY

The patient had a 2-year history of loose stools with occasional blood. He was not a regular alcohol drinker, with 1 episode of heavy drinking for 6 months a few years before this admission. No further personal or family history to indicate liver diseases was reported.

From the <sup>a</sup>Department of Cardiology, Queen Elizabeth Hospital Birmingham, University Hospitals Birmingham NHS Foundation Trust, United Kingdom; <sup>b</sup>Institute of Cardiovascular Sciences, University of Birmingham, United Kingdom; and the <sup>c</sup>Department of Gastroenterology, Queen Elizabeth Hospital Birmingham, University Hospitals Birmingham NHS Foundation Trust, United Kingdom.

The authors attest they are in compliance with human studies committees and animal welfare regulations of the authors' institutions and Food and Drug Administration guidelines, including patient consent where appropriate. For more information, visit the [Author Center](#).

Manuscript received March 8, 2023; revised manuscript received April 17, 2023, accepted April 20, 2023.

**ABBREVIATIONS  
AND ACRONYMS****EGD** =  
esophagogastroduodenoscopy**IR** = interventional radiology**IVC** = inferior vena cava**RA** = right atrium**TIPS** = transjugular  
intrahepatic portosystemic  
shunt**DIFFERENTIAL DIAGNOSES**

The differential diagnoses at this point were iatrogenic upper gastrointestinal hemorrhage and variceal bleeding with portal hypertension resulting from cirrhosis of uncertain etiology.

**INVESTIGATIONS**

On the following day, the patient underwent a repeated EGD under general anesthesia. Owing to the large volume of fresh blood obscuring the view, a Sengstaken-Blakemore tube was inserted as a rescue and bridge therapy to definitive treatment. The team investigated further with a CT angiogram, which did not suggest active bleeding.

**MANAGEMENT**


With a provisional diagnosis of portal hypertension and cirrhosis, a rescue transjugular intrahepatic portosystemic shunt (TIPS) was performed to lower portal venous pressure, with removal of the Sengstaken-Blakemore tube after a few days (**Figure 1**). TIPS was indicated to treat uncontrollable variceal bleeding secondary to portal hypertension.

Postoperatively, the patient experienced episodes of melena, requiring a further admission to the intensive care unit. A third EGD revealed oozing gastric varices requiring multiple bandings and glue injections. Because of further attacks of melaena, TIPS patency was investigated with Doppler ultrasonography and CT angiography, which showed TIPS patency to be inconclusive. A provisional diagnosis of cirrhosis of the liver with uncertain etiology, and a high United Kingdom Model for End-Stage Liver Disease score, resulted in the patient's being transferred to the regional liver transplantation unit.

In our center, an interventional radiologist then performed a TIPS venogram to investigate the patency and the need for potential embolization of varices. This surprisingly revealed the migration of glue in both the afferent and efferent systems of TIPS, further extending into the inferior vena cava (IVC) and the right atrium (RA), posing a severe risk factor for pulmonary embolization (**Figure 2, Video 1**). As this glue migration had not been reported in the previous CT scan, a repeat CT scan was performed to assess the risk of further bleeding from varices, the extent of glue migration, and feasibility to retrieve the migrated glue cast (**Figures 3 and 4**). The findings were: 1) good patency of the TIPS; 2) distribution of glue cast without imminent risk of further bleeding; and 3) mobile glue cast in the RA with embolized pieces in branches of the pulmonary arteries. Subsequent transthoracic echocardiography (**Figures 5 to 7**) found mobile echogenic, standlike structures within the RA, originating from the IVC, with normal biventricular size and function.

The cardiology team was then consulted for potential retrieval options. Our opinion was that snaring the glue cast was unlikely to be successful, with a further risk of breaking it into pieces flowing down to the pulmonary arteries. A similar view was provided by the interventional radiology (IR) team. Inasmuch as the risk of using a cardiopulmonary bypass machine for the patient was higher because of liver failure at that moment, the cardiothoracic

**FIGURE 1** Fluoroscopy Demonstrating Successful and Un-  
eventful Transjugular Intrahepatic Portosystemic Shunt  
Procedure



team also did not support open heart removal. The risk of forming a blood clot on the glue cast was managed by careful anticoagulation with therapeutic low-molecular-weight heparin subcutaneous injection. Then, comprehensive work-ups by the liver team, including magnetic resonance cholangiopancreatography and sigmoidoscopy, reached a final diagnosis for the liver transplantation candidate of cirrhosis of the liver due to primary sclerosing cholangitis with overlapping ulcerative colitis complicated by migration of glue used to treat gastric varices.


The multidisciplinary (interventional cardiology, cardiothoracic surgery, IR, and hepatology) discussion was then arranged to seek consensus on glue retrieval options. The teams agreed that the IR approach would carry the lowest risk, with the standby rescue cardiothoracic team.

During fluoroscopy by the IR team, the mobile glue cast in the RA was no longer visible (Video 2). The subsequent echocardiogram confirmed the presence of an echogenic glue cast (Videos 3 to 8). We concluded that lipiodol (radiopaque) was already dissolved; however, radiolucent Histoacryl glue remained in the RA. Therefore, the IR option was no longer feasible. Then, another multidisciplinary meeting proposed direct atrial extraction by the cardiothoracic team at the time of liver transplantation surgery before the liver could be removed, with cardiopulmonary bypass machine support.


## DISCUSSION



Histoacryl glue (a monomer, n-butyl-2-cyanoacrylate) polymerizes into a solid form once exposed to ionic materials such as water or blood. It is usually mixed with lipiodol (0.5 mL glue and 1 mL lipiodol at the authors' center; however, the proportion varies across units both nationally and internationally). Mixing with radiopaque lipiodol not only enables postprocedural radiological assessment but also slows the polymerization process, allowing the endoscopists more time for injection. It is licensed to treat emergency gastric variceal bleeding as first-line therapy. The potential complications include rebleeding, erosion, ulceration, extrusion of glue cast, portal and splenic vein thrombosis, sepsis, and embolization.

A recent retrospective study reported that the risk of distant embolization after glue injection was 0.5% (1 pulmonary and 1 cerebral).<sup>1</sup> The presentation can



vary from within hours to days after injection.<sup>2</sup> A catastrophic outcome with mortality from embolization was also reported.<sup>1</sup> In terms of management, a recent case report followed up the patient with conservative management with anticoagulation and





showed no change in the size of the glue in the right ventricle.<sup>3</sup> Few recent case reports of retrieval by a minimally invasive approach or open heart surgery were identified in the literature.

TIPS is a well-established procedure involving inserting a stent to connect the portal veins to adjacent hepatic veins under imaging guidance to relieve portal hypertension, thereby reducing the risk of

variceal hemorrhage. However, this therapeutic procedure unexpectedly served as a conduit that allowed glue extension and embolization in our patient. Additionally, liver failure requiring transplantation further limits the management options.

To the authors' knowledge, this case report is the first to describe the extension of glue cast through TIPS to the IVC down to the RA.

