

Fatores Sociodemográficos e de Saúde Associados aos Sintomas Depressivos Durante a Pandemia de COVID-19

Trettel, Ana Cláudia Pereira Terças; Muraro, Ana Paula; Nascimento, Vagner Ferreira do; Andrade, Amanda Cristina de Souza; Espinosa, Mariano Martinez; Oliveira, Elaine Cristina de; Wagstaff, Christopher; Pillon, Sandra C.

DOI:

[10.51338/rppsm.439](https://doi.org/10.51338/rppsm.439)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Trettel, ACPT, Muraro, AP, Nascimento, VFD, Andrade, ACDS, Espinosa, MM, Oliveira, ECD, Wagstaff, C & Pillon, SC 2023, 'Fatores Sociodemográficos e de Saúde Associados aos Sintomas Depressivos Durante a Pandemia de COVID-19: Um Estudo Transversal', *Revista Portuguesa de Psiquiatria e Saúde Mental*, vol. 9, no. 2, pp. 36-48. <https://doi.org/10.51338/rppsm.439>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

ORIGINAL ARTICLE/ ARTIGO ORIGINAL

The Health and Socio-Demographic Factors Associated with Depressive Symptoms During the COVID-19 Pandemic: A Cross-Sectional Study Fatores Sociodemográficos e de Saúde Associados aos Sintomas Depressivos Durante a Pandemia de COVID-19: Um Estudo Transversal

ANA CLÁUDIA PEREIRA TERÇAS TRETTEL^{*1,2,3}, ANA PAULA MURARO², VAGNER FERREIRA DO NASCIMENTO¹, AMANDA CRISTINA DE SOUZA ANDRADE², MARIANO MARTINEZ ESPINOSA², ELAINE CRISTINA DE OLIVEIRA⁴, CHRISTOPHER WAGSTAFF⁵, SANDRA CRISTINA PILLON³

1. Department of Nursing, Campus Tangará da Serra, State University of Mato Grosso (UNEMAT), Tangará da Serra – MT, Brazil.
2. Health Institute collective, university Federal of Mato Grosso (UFMT). Cuiabá - MT, Brazil.
3. School of Nursing of Ribeirão Preto, University from Sao Paulo (USP). Ribeirão Preto – SP, Brazil.
4. Secretary of State for Mato Grosso Health. Cuiabá - MT Brazil.
5. University of Birmingham. Birmingham, England.

Abstract

Introduction: Our objective was to analyze depressive symptoms during the COVID-19 pandemic and their association with health and sociodemographic factors.

Methods: A survey of 4203 adults from ten municipalities was conducted in Mato Grosso state, Amazon region of Brazil. Data collection was done at home, with the collection of sociodemographic characteristics, and aspects of both physical and mental health. The DSM-5 level 1 cross-sectional symptom scale was used to identify symptoms of depression, substance use disorders, sleep disturbances, and anxiety. Chemiluminescence was also used to detect IgG anti-SARS-CoV-2 antibodies.

Results: The estimated prevalence of anti-SARS-CoV-2 antibodies was 12.5% and the prevalence of depressive symptoms was 23.2%. In the multivariate analysis, depression was associated with being male, having a low income, the level of physical activity before the pandemic, chronic illness, substance use disorder, sleep and anxiety disorders.

Conclusion: Evidence on depression and its associated factors is critical to understanding the extent of the population situation from which recommendations can be made to guide public policy.

Resumo

Introdução: O nosso objetivo foi analisar a associação entre sintomas depressivos e fatores sociodemográficos e de saúde durante a pandemia de COVID-19.

Métodos: Foi realizado um estudo com 4203 adultos de dez municípios do estado de Mato Grosso, região amazônica do Brasil. A recolha de dados foi realizada na residência, com levantamento de informações sobre as características sociodemográficas, aspectos de saúde física e mental. A Escala de Sintomas Transversais de Nível 1 do DSM-5 foi utilizada para identificar sintomas de depressão, transtorno por uso de substâncias, distúrbios do sono e ansiedade. A quimiluminescência foi utilizada para detectar anticorpos IgG anti-SARS-CoV-2.

Recebido/Received: 2022-12-29

Aceite/Accepted: 2023-05-25

Publicado Online/Published Online: 2023-06-05

Publicado/Published:

* Autor Correspondente/Corresponding Author: Ana Cláudia Pereira Terças Trettel | ana.claudia@unemat.br | Departamento de Enfermagem, Campus Tangará da Serra, Universidade do Estado de Mato Grosso. MT 358, Jardim Industrial. 78300-000 Tangará da Serra MT Brasil.

© Author(s) (or their employer(s)) and SPPSM Journal 2023. Re-use permitted under CC BY 4.0.

© Autor (es) (ou seu (s) empregador (es)) e Revista SPPSM 2023. Reutilização permitida de acordo com CC BY 4.0.

Resultados: A prevalência estimada de anticorpos anti-SARS-CoV-2 foi de 12,5% e a prevalência de sintomas depressivos foi de 23,2%. Na análise multivariada, a depressão esteve associada a ser do sexo masculino, ter baixa renda, nível de atividade física antes da pandemia, ter doença crônica, apresentar transtorno por uso de substâncias, transtornos do sono e de ansiedade.

Conclusão: Os achados sobre a depressão e seus fatores associados são essenciais para a compreensão sobre a magnitude do quadro desses sintomas na população, a partir de onde podem ser feitas recomendações que norteiem as políticas públicas.

Keywords: Brazil; Depression; Mental Health; SARS-CoV-2; Surveys and Questionnaires

Palavras-chave: Brasil; Depressão; Inquéritos e Questionários; Saúde Mental; SARS-CoV-2

INTRODUCTION

After the emergence of COVID-19 and the changes arising from the development of the ‘new normal’ in societies, the world is now experiencing multidimensional challenges. In the health sector, in addition to the constant search for better therapies for the treatment of COVID-19 and the intensification of prevention activities, such as immunization, there is great concern about the impact of the pandemic on the mental health of the population.¹⁻³ There is evidence indicating that globally there has been a reduction in physical and psychological well-being since the start of the pandemic.⁴⁻⁶

A negative impact on people’s mental health has already been reported after previous epidemics, such as Ebola,⁷ however, the impact is greater with the current global pandemic because of the rapid worldwide spread, both viral and the infodemic and fake news.⁸ A meta-analysis study, comparing pandemic and pre-pandemic data, found a significant increase in depression rates.⁹ As well as a 34.3% prevalence of depression in major international studies since the identification of SARS-CoV-2.¹⁰ The COVID-19 Mental Disorders Collaborators estimated that since the beginning of the pandemic there was a 27.6% increase in global cases of depressive disorder in 2020 and major depressive disorder caused 49.4 million (33.6 to 68.7) DALYs.¹¹ However, the mechanisms of depression in COVID-19 cases are still unclear, even though it is the most common disorder of neurological symptoms post infection.¹²

Variables such as being married, having a lower level of education, having a chronic illness and having a previous history of depression in pregnant women were associated with depression in women during the pandemic.¹³ In students, severe symptoms of depression were observed, higher in males and among those with a lower level of social support.¹⁴ Among elderly Brazilians, women had a higher prevalence of depressive symptoms than men, which was again associated with low income.¹⁵ For individuals who had a previous history of major depressive disorders, the pandemic intensified symptoms and impacted psychopharmacological treatment.¹⁶ The complexity of the compounding factors of depression has already been described in international studies, however, studies must

be expanded worldwide to enable mental health services to confront the mental health consequences of the COVID-19 pandemic. Thus, the present study aims to analyze the health and socio-demographic factors associated with depressive symptoms during the COVID-19 pandemic in the population of three main Brazilian biomes.

METHODS

A cross-sectional household survey was carried out in the state of Mato Grosso, a state that, due to its wide territorial extension, is home to the three main biomes of the country, namely the Amazon, Cerrado and Pantanal. Ten cities were listed, whose populations ranged from 30 000 to 600 000 inhabitants, and which are located in 3 of the main Brazilian biomes (Cerrado [5 cities], Pantanal [1 city] and Amazônia [3 cities]). The main connections of regional integration are also considered, and it was possibly through these that the hierarchical diffusion process of COVID-19 in this region took place. This study is part of the matrix project “*Endemic and epidemic diseases in Mato Grosso*”.¹⁷

The sample size calculation was performed using the 2019 population estimate for the 10 selected municipalities (sample space: 1 650 643 inhabitants in the 10 municipalities of the study), considering a confidence level of 95%, design effect equal to 1.5, the prevalence of 3% and precision of 0.65%.¹⁸ The sample size was calculated using OpenEpi21. A 13% recomposition was added to the sample size, considering the anticipated losses resulting from refusals and the existence of closed households during attempted recruitment. The sample weight of each selected unit (census sector, household and individual) was calculated separately for each municipality, considering the inverse of the selection probability according to the proposed sampling plan, including calibrations for adjustments to the known population totals. Medium-sized municipalities, census sectors in urban areas and residents over 18 years of age were included, and municipalities with less than 25 000 inhabitants, census sectors that encompassed rural areas and indigenous reserves and unconscious bedridden residents were excluded.¹⁹

Data collection took place between September and October 2020 by professionals and undergraduate health

students. Prior training was carried out in August 2020 to standardize interviews, blood collection and pilot testing. The pilot test was carried out in a census sector not selected for sampling and its data were not included in the analysis. During data collection, the selected census sector was covered following a systematic approach of selecting households by “jumps” determined for each sector. If the selected household was empty at the time or the selected resident did not agree to participate in the research, the next residence on the left was taken as a substitute.

When the participants’ home was visited for recruitment, a resident over 18 years of age was randomly selected to answer the questionnaires, which were applied through Epi InfoTM²⁰ on Smartphones, which was also used to capture geographic coordinates. Fieldwork was carried out by a coordinator in each selected municipality. The data that was collected during the interview with the individual included (i) sociodemographic data, (ii) physical health data and (iii) mental health symptoms.

The socio-demographic data included simple questions about gender (female and male), race/skin colour (non-white and white), education (illiterate, elementary, secondary and higher education and more) and household income before the pandemic (< 1 minimum wage (MW), 1-3 and >3MW) (family income was assessed using R\$ 1045.00 or US\$ 183.91), reduced income during the pandemic (yes or no) and food insecurity (yes or no) analyzed by the question: “*Since the beginning of the COVID-19 Pandemic, has there been a time when food ran out in your house and you didn’t have the money to buy more?*”.

ii) Health – which involved questions about body weight change since the beginning of the pandemic; physical exercise before the pandemic; physical exercise during the pandemic; the presence of antibodies against SARS-CoV-2 (confirmed by chemoimmunofluorescence), self-reported symptoms of COVID-19; hospitalization for COVID-19; and the interaction between having a chronic disease and taking medication.

Mental Health Symptoms – Mental health variables were obtained using the DSM-5 Level 1 Symptoms Scale, except for alcohol use, which used the question about weekly alcohol use (yes/ no). Thus, the dependent variable depression (yes/ no), and the independent variables substance use disorder (yes/ no), sleep disorder (yes/ no) and anxiety disorder (yes/ no) were investigated using the DSM-5.

The DSM-5 is an instrument developed by the American Psychiatric Association [APA],²¹ which assesses mental health symptoms. This scale has also shown promise as a screening tool for research. The adult version contains 23 items with a response on a five-point scale (0=none; 1=rare, less than one or two days; 2=mild or several days; 3=moderate or more than half of the days; and 4=severe or nearly every day) to assess 13 psychiatric domains. The depression, medication use, sleep disorder and anxiety domains were selected for this study. For the substance use domain, scores are considered positive: very mild (1), mild (2), moderate (3) or severe (4) scores and only the score nothing (0) is considered negative. For depression, sleep disorders and

anxiety symptoms, the results of the DSM-5 domains when the scores are positive then the symptoms are considered mild, moderate or severe and when the scores are negative then there are no or very mild symptoms.

The presence of antibodies against SARS-CoV-2 was confirmed by chemoimmunofluorescence through biological samples that were collected at home and later cryopreserved at -20° to be transported to the Central Laboratory of Public Health of Mato Grosso (LACEN-MT), where laboratory tests were performed.

All analyzes were performed using Stata software version 16, mainly the “svy” module, which allows the addition of weighting factors and takes into account the complex design of the sample.

The depression variable was considered a dependent variable and all sociodemographic, health and mental health variables were considered independent variables. Initially, a descriptive analysis of the data was performed. Variables with *p* values lower than 0.20 were tested in the Poisson multiple regression model with robust variance in the bivariate analysis between the dependent and independent variables, considering the prevalence ratio adjusted by the method. The prevalence ratios (PR) and their respective 95% confidence intervals (95% CI) were calculated using the model mentioned above. Variables with *p*-values lower than 0.20 in the bivariate analysis were included in the multiple model and variables with *p*-values lower than 0.05 by the stepwise forward method remained in the final model.

RESULTS

The sample was characterized by women of working age (18 to 49 years), who were ethnically non-white, with a low level of education, with a low family income (1 and 3 minimum wages), and a reduced income after the start of the pandemic but had not been experiencing food shortages at home in September/October 2020. The prevalence of anti-SARS-CoV-2 antibodies was estimated at 12.5% (95% CI: 10.5; 14.7), ranging from 7.4% to 24.3% among the municipalities, and the prevalence of depressive symptoms (DSM-5) was 23.2%.

In the bivariate analysis, depression was associated with being ethnically non-white, having a low-income pre-pandemic (< 1 minimum wage and between 1-3 minimum wages), and a reduction in family income during the pandemic (Table 1). As regards physical health, there was a positive association between increased body weight after the start of the pandemic, physical activity before and after the start of the pandemic, chronic disease, weekly use of alcohol, as well as with symptoms of substance abuse disorder, sleep disorder and anxiety disorders (DSM-5) (Table 2).

In Table 3, the associations with depression continued to be being male (PR=1.04, CI=1.01;1.07, *p*=0.017), with the family income before the pandemic being less than one minimum wage (PR=0, 92, CI=10.86;0.99, *p*=0.020), not taking physical exercise after the start of the pandemic (PR=1.16, CI=1.07;1.27, *p*<0.001), the interaction between

chronic disease/medication use (PR=1.08, CI=1.01;1.15, $p=0.017$) was inversely associated with depression, with a 15% lower prevalence of depressive symptoms among those classified as having this disorder (Table 3).
 $p=0.036$), sleep disorders (PR=1.41, CI=1.32; 1.50, $p<0.001$) and anxiety disorders (PR=3.07, CI=2.57;3.66, $p<0.001$). Substance use disorder (PR=0.95, CI=0.91;0.99,

Table 1. Depression and socioeconomic factors, physical and mental health during the COVID-19 pandemic in the population of the three main Brazilian biomes, according to sociodemographic and economic variables. Mato Grosso, Brazil, 2020 (N=4,203).

Variables	Depression		RP _b **	IC (95%)**	p value
	No	Yes			
Gender					
Male	1146	330	1.03	(0.99; 1.08)	0.155
Female	2084	643	1.00	-	-
Race/skin colour					
No white	2233	601	1.07	(1.02; 1.12)	0.007*
White	997	372	1.00	-	-
Schooling					
Illiterate	149	46	0.99	(0.88; 1.10)	0.782
Elementary school	1150	388	0.98	(0.93; 1.04)	0.509
High school	1101	312	1.02	(0.93; 1.03)	0.416
Undergraduate and more	830	227	1.00	-	-
Family income before the pandemic					
< 1 minimum wage (MW)	318	281	0.62	(0.56; 0.68)	<0.001*
≤ 1 WM <3 WM	1747	455	0.94	(0.90; 0.97)	0.001*
>3 WM	1165	237	1.00	-	-
Reduction in income during the pandemic					
Yes	1814	477	1.08	(1.03; 1.12)	0.002*
No	1416	496	1.00	-	-
Food insecurity					
Yes	271	76	0.99	(0.91;1.07)	0.7451
No	2959	897	1.00	-	-

Notes: RPa: prevalence ratio estimated by the robust simple Poisson regression model. 95% CI: 95% confidence interval. **: Prevalence and 95% confidence intervals (95% CI) estimated under sample weights. p : p value. *: significant at the 5% level.

Table 2. Depression and socioeconomic aspects, health and mental health during the COVID-19 pandemic in the population of the three main Brazilian biomes, according to variable health aspects. Mato Grosso, Brazil, 2020 (N=4,203).

Variables	Depression		RP _b **	IC (95%)**	p value
	No	Yes			
HEALTH					
Body weight after the beginning of the pandemic					
Yes, increased	1120	286	1.07	(1.02; 1.12)	0.004*
Yes, decreased	463	127	1.06	(0.99; 1.13)	0.054
No	1647	560	1.00	-	-
Practice of physical exercises before the pandemic					
No	2972	582	2.08	(1.85; 2.3454)	<0.001*
Yes	258	391	1.00	-	-
Practice of physical exercises during the pandemic					
No	2628	495	1.51	(0.62; 0.71)	<0.001*
Yes	602	478	1.00	-	-
Presence of antibodies against SARS-CoV-2					
Positive	447	132	1.00	(0.94; 1.06)	0.99
Negative	2783	841	1.00	-	-
COVID-19 self-reported symptoms					
Yes	245	68	1.01	(0.93; 1.09)	0.793
No	2985	905	1.00	-	-
Hospitalization for COVID-19					
Yes	18	6	1.05	(0.84; 1.32)	0.667
No	3212	967	1.00	-	-
Interaction between having chronic illness and medication use					
Yes	1316	337	1.07	(1.03; 1.12)	0.001*
No	1914	636	1.00	-	-
MENTAL HEALTH SYMPTOMS					
Weekly alcohol use					
Yes	688	152	1.10	(1.06; 1.15)	<0.001*
No	2542	821	1.00	-	-
Substance use disorder					
Yes	2384	840	0.86	(0.82; 0.89)	<0.001*
No	846	133	1.00	-	-
Sleep disorder					
Yes	3057	334	4.19	(3.56; 4.94)	<0.001*
No	173	639			-
Anxiety disorder					
Yes	2608	185	2.12	(1.97; 2.28)	<0.001*
No	622	788	1.00	-	-

Notes: RPa: prevalence ratio estimated by the robust simple Poisson regression model. 95% CI: 95% confidence interval. **: Prevalence and 95% confidence intervals (95% CI) estimated under sample weights. p: p value. *: significant at the 5% level.

Table 3 - Final model of the multiple analysis between depression and socioeconomic factors, physical and mental health during the COVID-19 pandemic in the population of the three main Brazilian biomes, Mato Grosso, Brazil, 2020 (N= 4203).

Variables	Depression		RP _b **	IC (95%)**	p value
	No	Yes			
SOCIODEMOGRAPHIC					
Gender					
Male	1146	330	1.04	(1.01; 1.07)	0.017*
Female	2084	643	1.00	-	-
Family income before the pandemic					
< 1 minimum wage (MW)	318	281	0.92	(0.86; 0.99)	0.020*
≤ 1 WM <3 WM	1747	455	0.98	(0.95; 1.02)	0.150
>3 WM	1165	237	1.00	-	-
HEALTH					
Participating in physical exercises before the pandemic					
No	2972	582	1.16	(1.07; 1.27)	<0.001*
Yes	258	391	1.00	-	-
Interaction between having a chronic illness and medication use					
Yes	989	276	1.08	(1.01; 1.15)	0.036*
No	2241	697	1.00	-	-
MENTAL HEALTH SYMPTOMS					
Substance use disorder					
Yes	2384	840	0.95	(0.91; 0.99)	0.017*
No	846	133	1.00	-	-
Sleep disorder					
Yes	2608	185	1.41	(1.32; 1.50)	<0.001*
No	622	788	1.00	-	-
Anxiety disorder					
Yes	3057	334	3.07	(2.57; 3.66)	<0.001*
No	173	639	-	-	-

Notes: RPa: prevalence ratio estimated by the robust simple Poisson regression model. 95% CI: 95% confidence interval. **: Prevalence and 95% confidence intervals (95% CI) estimated under sample weights. *p*: *p* value. *: significant at the 5% level.

DISCUSSION

The main limitation of the present study is that its design does not allow causal conclusions between predictors and depression; however, it explores the relationship between the variables, with results consistent with those found in longitudinal studies. In addition, there was still the risk of interviewer bias and to minimize it, previous training of interviewers and field supervision were carried out. Memory bias and desirability bias related to mental health issues (responses to socially accepted standards) were potential factors for participants. To mitigate against these the interview was held in a discreet place, guaranteeing all the

necessary time, in addition to emphasizing the anonymity of their answers.

In the Brazilian biomes studied here, 23.2% of participants had depressive symptoms, similar to the average observed in other countries of 34.3%, ranging from 6.8% to 83.7%.²²⁻²⁷ Among the associated factors were, being male, low income, not taking physical exercise before the pandemic and other components of mental health such as anxiety, sleep and substance use.

Importantly, depression was not associated with having anti-SARS-CoV-2 antibodies (evidence of past infection), even when there was a COVID-19 prevalence of 12.5%

in the population studied. This finding is based on the fact that the limitations imposed by social distancing increased loneliness, stress in relationships and disconnection in communities, regardless of viral infection.²⁸ In this light, the findings of Tsang and Avery²⁹ remain, which describe that the relationship between depression and COVID-19 was importantly mediated by fear of the disease.

Regarding gender, studies point to a higher risk among women,^{27,30} with results highlighting a higher prevalence of depression.^{31,32} However, contradicting these studies, in this study there was an association between depression and being male. Possibly, this is related to the fact that men have a worse clinical course and higher mortality from COVID-19,^{33,34} or due to the behavior of men in maintaining healthy habits and lifestyles, thus tending to engage less in the search for care and prescribed treatments.³⁵ Another possibility lies in the sociogeographical aspect of this investigated region, which in the last pre-pandemic decade sustained high suicide rates among males, among all causes of death of this nature (ICD X60 to X84),³⁶ a circumstance that reflects among other aspects, the symptom and repercussions of depression, often not verbalized and/or treated by men, but which during the pandemic became familiar in this group, as manifesting depression in this period did not refer to the fragility so avoided by men at other times. Furthermore, it is important to note that in China, no association was found between gender and depression during the pandemic.³⁷

Another relevant point is the complex relationship between race/ethnicity, mental disorders and depression. Previous and contemporaneous research carried out in the United States^{38,39} and Brazil^{40,41} showed an association between people of colour and depressive symptoms. It is known that there is no biological relationship that associates ethnicity and mental health,⁴² however, belonging to socially and economically disadvantaged ethnic groups can trigger emotional overloads and reduce positive states of mood and well-being.⁴³

The negative economic impact caused by the pandemic was exorbitant, yet the financial weaknesses had an even more aggressive impact on people's mental health.⁴⁴ The current research contributes to the knowledge about this relationship in a developing country such as Brazil. In the population studied, representing 10 municipalities in the state of Mato Grosso, an association was detected between depressive symptoms and having a low income and reduced family income during the pandemic. Similar to results in the North American countries, such as a Canadian study in which the loss of income occurred in 10.3% of those evaluated, with a 12% increase in the cases of depression⁴⁵ and American adults who lost their jobs after the start of the pandemic.⁴⁶ This financial strain and income disruption have also been reported among the risk factors for depression in Asia.²⁷

Staying at home as a strategy to control viral spread in the first year of the pandemic was necessary. However, this strategy led to less physical activity, predisposing people to an increased risk of depression.⁴⁷ In the three Brazilian biomes studied, it was shown that not practising physical

activity before and after the beginning of the pandemic was statistically associated with depression. Dun *et al*⁴⁸ describe that prior physical fitness may be inversely associated with depression in adults during the pandemic.

An Australian study at the beginning of the pandemic described that the reduction in physical exercise and increased sedentary lifestyle had a direct impact on the mental health of the population,⁴⁹ with it being reported that reconciling healthy eating with physical activity is important to enhance the relief of depressive symptoms⁵⁰ and that improvement in these two behaviours were associated with low levels of depression in adolescents.⁵¹ Thus, the increase in body weight after the pandemic is related to the risk for depression in our study, noting that people's attitudes towards food are important when maintaining healthy lifestyle habits. Weight gain in uninfected people may occur due to decreased physical exercise, changes in food purchase and consumption habits, greater stress when eating, economic stress and anxiety.⁵²

The change in behaviour resulting from the challenges posed by the pandemic affected the global community, but for those with chronic diseases, there were additional aggravating factors, such as the suspension of routine monitoring health services, limiting access and weakening clinical management.² In our study, there was an association between having a chronic condition and depression during the pandemic, as also described in dialysis patients,⁵³ cancer patients,⁵⁴ diabetics, patients with hypertension, COPD and asthma,^{55,56} as well as Parkinson's.⁵⁷

The increase in negative emotions during the pandemic has been described as a motivating factor for alcohol consumption, including among those who were long-term abstainers.^{58,59} The phenomenon has been managed differently across countries since in some there was a loosening of access^{60,61} and in others, there was a temporary ban,^{62,63} a fact recorded unevenly in Brazilian municipalities. A systematic review presented data on the change in alcohol consumption, highlighting the concern about the increase in alcohol consumption after the onset of the pandemic, and its association with depression and anxiety in most of the studies investigated.⁶⁴

The consumption of alcohol during COVID-19 has been described differently in the world, with different findings concerning gender, age, skin colour/ethnicity and income, however, high rates of depression associated with increased use of alcohol stand out in Poland,⁶⁵ Canada,⁶⁶ the United States⁶⁷ and Australia.⁶⁸

Substance use disorder is defined as a pattern of substance use of any kind, leading to clinically significant distress or impairment.²¹ A study carried out in the United States showed that 13% of the population started or increased substance use after the beginning of the pandemic,⁶⁷ with a similar increase being detected in Canada,⁶⁹ while in Poland there was an increase in the consumption of recreational drugs, but in a lower percentage (1.4%).⁷⁰ In the present study, after adjusting for the other variables in the multiple models, there was a 15% lower prevalence of depression among those classified as having a substance use disorder. This result can be explained by the fact that this disorder is

related to depression, mainly in moments of abstinence and in chronic use, since initially use can minimize the effects of depression.⁷¹

Another relevant aspect associated with depression is sleep disorder; with pre-pandemic evidence, in both an international meta-analysis⁷² and a Brazilian study,⁷³ showing the correlation between sleep disorders and depression. With the changes in lifestyle and the restrictions associated with the pandemic, sleep disorders have increased over time with rates reaching 48% of insomnia in the population.⁷⁴

This complex relationship between sleep and depression is described in studies with the general population in Bangladesh,⁷⁵ Italy⁷⁶ and China,⁷⁷ as well as specific populations such as pregnant women⁷⁸ and health professionals.⁷⁹ It is also worth noting the negative emotional impact of the exacerbated use of online tools during the pandemic, due to the loneliness of the home office, online games, social media and internet shopping or the uninterrupted attack of *fake news*⁸⁰ may have contributed to the sleep disturbance.

The relationship between depression, sleep disorder and not practising physical activity is mostly permeated in a transversal way by anxiety. An alert issued by the World Health Organization in March 2022, suggested that countries pay attention to the impact of the pandemic on mental health and include more effective actions in their COVID-19 response plans, given the 25% increase in the prevalence of depression and anxiety.⁸¹ The COVID-19 Mental Disorders Collaborators, however, consider that at the beginning of the pandemic there was a significant increase in depression and anxiety rates in response to the acute and emerging crisis, however, over time it softened, approaching pre-pandemic values.

Depression and anxiety were prominent before the pandemic, and constituted the most prevalent mental health disorders, demanding a great global effort to address them. It should be noted that the disorganization of mental health services and the fragility of resources pre-pandemic⁸² were aggravated by the growing demand at the beginning of the pandemic, interruption of care in mental health services and personal limitations.⁸³

Data points to an average anxiety prevalence rate of 38% in several international studies with Chinese,⁸⁴ Italian,²⁵ Indian⁸⁵ and Turkish⁸⁶ populations. Despite the high prevalence

of depressive and anxious symptoms, both are not necessarily recorded together, but a complex relationship can exist so that one triggers the mechanisms of the other and vice versa, as they result from the interaction of the same social, psychological and biological factors.⁸⁷ Thus, this association between depression and anxiety is complex and must be understood considering several factors.

CONCLUSION

The high prevalence of depressive symptoms during the first year of the pandemic identified here refers highlights the reality for a significant proportion of the population during this difficult time, as well as socioeconomic conditions already present in everyday life before SARS-CoV-2. Multivariate analysis in this study demonstrates that depression is associated with being male, having a low income, having a sedentary lifestyle before the pandemic, chronic illness, substance use disorder, sleep and anxiety disorder (DSM-5).

The pandemic has created a greater urgency to strengthen mental health systems in most countries, so investigating the impact on the mental health of populations, especially depression and its associated factors, is fundamental in providing the acquisition of knowledge that directs assertive actions of interventions that should be implemented to cushion the negative effects of SARS-CoV-2 on mental health.

Research on depression must be continuous and comprehensive regarding associated factors to support the understanding of the duration, magnitude and severity of the pandemic's impact on the mental health of populations. Mitigation strategies must be recommended to incorporate ways to promote mental well-being, reduce the determinants of mental health and target appropriate interventions to treat those who develop a mental disorder.

ACKNOWLEDGMENTS

The authors are extremely grateful to everyone who agreed to participate in the research and to those who did the home visits, collection of biological samples, conduction of diagnostic tests, advised the population and delivered the results to the population visited.

Responsabilidades Éticas

Conflitos de Interesse: Os autores declaram a inexistência de conflitos de interesse na realização do presente trabalho.

Fontes de Financiamento: Este estudo foi realizado com o apoio do Governo do Estado de Mato Grosso, que disponibilizou os exames diagnósticos e estrutura para análises laboratoriais, bem como da UFMT e da UNEMAT, que disponibilizaram pessoal para auxiliar na coordenação, logística e divulgação de resultados.

Confidencialidade dos Dados: Os autores declaram ter seguido os protocolos da sua instituição acerca da publicação dos dados de doentes.

Proteção de Pessoas e Animais: Os autores declaram que os procedimentos seguidos estavam de acordo com os regulamentos estabelecidos pela Comissão de Ética responsável e de acordo com a Declaração de Helsínquia revista em 2013 e da Associação Médica Mundial.

Proveniência e Revisão por Pares: Não comissionado; revisão externa por pares.

Ethical Disclosures

Conflicts of Interest: The authors have no conflicts of interest to declare.

Financing Support: This study was carried out with the support of the Government of the State of Mato Grosso, which provided the diagnostic tests and structure for laboratory analysis, as well as UFMT and UNEMAT, which provided members of staff to assist with coordination, logistics and dissemination of results.

Confidentiality of Data: The authors declare that they have followed the protocols of their work center on the publication of data from patients.

Protection of Human and Animal Subjects: The authors declare that the procedures followed were in accordance with the regulations of the relevant clinical research ethics committee and with those of the Code of Ethics of the World Medical Association (Declaration of Helsinki as revised in 2013).

Provenance and Peer Review: Not commissioned; externally peer reviewed.

Declaração de Contribuição

ACPTT: Concepção, delineamento, coleta de dados, interpretação dos dados, redação, revisão crítica e aprovação da versão.

APM e ACSA: Concepção, projeto, coleta de dados, interpretação dos dados, redação, revisão crítica e aprovação da versão.

ECO: Coleta de dados, redação, revisão crítica e aprovação da versão.

VFN e MME: Interpretação dos dados, redação, revisão crítica e aprovação da versão.

CW: Tradução e a edição final do artigo.

SCP: Concepção, projeto, coleta de dados, interpretação dos dados, redação, revisão crítica e aprovação da versão.

Contributorship Statement

ACPTT: Contributed to the conception, design, data collection, data interpretation, writing, critical review and approval of the final version.

APM e ACSA: Contributed to the conception, design, data collection, data interpretation, writing, critical review and approval of the final version.

ECO: Contributed to data collection, writing, critical review and approval of the final version.

VFN e MME: Contributed to data interpretation, writing, critical review and approval of the final version.

CW: Contributed with the translation and final editing of the article.

SCP: Contributed to the design, design, data collection, data interpretation, writing, critical review and approval of the final version.

References

1. Pierce M, Hope H, Ford T, Hatch S, Hotopf M, John A, et al. Mental health before and during the COVID-19 pandemic: a longitudinal probability sample survey of the UK population. *Lancet Psychiatry*. 2020;7:883–92. doi: 10.1016/S2215-0366(20)30308-4.
2. Brooks SK, Webster RK, Smith LE, Woodland L, Wessely S, Greenberg N, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet*. 2020;395:912–20. doi: 10.1016/S0140-6736(20)30460-8.
3. Holmes EA, O'Connor RC, Perry VH, Tracey I, Wessely S, Arseneault L, et al. Multidisciplinary research priorities for the COVID-19 pandemic: a call for action for mental health science. *Lancet Psychiatry*. 2020;7:547–60. doi: 10.1016/S2215-0366(20)30168-1.
4. Vindegaard N, Benros ME. COVID-19 pandemic and mental health consequences: systematic review of the current evidence. *Brain Behav Immunity*. 2020;20:30954-5. doi: 10.1016/j.bbi.2020.05.048.
5. Xiong J, Lipsitz O, Nasri F, Lui LMW, Gill H, Phan L, et al. Impact of COVID-19 pandemic on mental health in the general population: a systematic review. *J Affective Disord*. 2020;277:55–64. doi: 10.1016/j.jad.2020.08.001.
6. Krishnamoorthy Y, Nagarajan R, Saya GK, Menon V. Prevalence of psychological morbidities among general population, healthcare workers and COVID-19 patients amidst the COVID-19 pandemic: a systematic review and meta-analysis. *Psychiatry Res*. 2020;293:113382. doi: 10.1016/j.psychres.2020.113382.
7. Cénat JM, Felix N, Blais-Rochette C, Rousseau C, Bukala J, Derivois D, et al. Prevalence of mental health problems in populations affected by the Ebola virus disease: a systematic review and meta-analysis. *Psychiatry Res*. 2020;289: 113033. doi: 10.1016/j.psychres.2020.113033.
8. Zielinski C. Infodemics and infodemiology: a short history, a long future. *Rev Panamericana Salud Publica*. 2021;5:e40. doi: 10.26633/RPSP.2021.40.

9. Kunzler AM, Röthke N, Günthner L, Stoffers-Winterling J, Tüscher O, Coenen M, et al. Mental burden and its risk and protective factors during the early phase of the SARS-CoV-2 pandemic: review and meta-analyses. *Global Health*. 2021;17:34. doi: 10.1186/s12992-021-00670-y.
10. Necho M, Tsehay M, Birkie M, Biset G, Tadesse E. Prevalence of anxiety, depression, and psychological distress among the general population during the COVID-19 pandemic: A systematic review and meta-analysis. *Int J Soc Psychiatry*. 2021;67:892-906. doi: 10.1177/00207640211003121.
11. COVID-19 Mental Disorders Collaborators. Global prevalence and burden of depressive and anxiety disorders in 204 countries and territories in 2020 due to the COVID-19 pandemic. *Lancet*. 2021;6,398:1700-12. doi: 10.1016/S0140-6736(21)02143-7.
12. Mohammadkhanizadeh A, Nikbakht F. Investigating the potential mechanisms of depression induced-by COVID-19 infection in patients. *J Clin Neurosci*. 201;91:283-7. doi: 10.1016/j.jocn.2021.07.023
13. Keliyo ET, Jibril MK, Wodajo GT. Prevalence of Antenatal Depression and Associated Factors among Pregnant Women Attending Antenatal Care at Health Institutions of Faafan Zone, Somali Region, Eastern Ethiopia. *Depress Res Treat*. 2021;27:2523789. doi: 10.1155/2021/2523789.
14. Guo K, Zhang X, Bai S, Minhat HS, Nazan AINM, Feng J, et al. Assessing social support impact on depression, anxiety, and stress among undergraduate students in Shaanxi province during the COVID-19 pandemic of China. *PLoS One*. 2021;23,16:e0253891. doi: 10.1371/journal.pone.0253891.
15. Silva Pereira-Ávila FMV, Lam SC, Goulart MCL, Góes FGB, Pereira-Caldeira NMV, Gir E. Fatores associados aos sintomas de depressão entre idosos durante a pandemia da COVID-19. *Texto Contexto Enferm*. 2021;30:e20200380.
16. De Filippis S, Lombardozzi G, Matrone M, Amici E, Trovini G, Perrini F, Di Giovanni A, Giovanetti V, Kotzalidis GD. Differential response to three antidepressants in patients with major depressive episode who suffered Covid-19-related trauma. *Curr Neuropharmacol*. 2022;20(2):393-407. doi: 10.2174/1570159X20666220310122849.
17. Trettel ACPT, Muraro AP, Oliveira EC, Nascimento VF, Andrade AC, Santos ESD, et al. Factors associated with suicidal ideation during the COVID-19 pandemic in a population in the Brazilian Legal Amazon. *Ciência Saúde Coletiva*. 2022;27: 3157-70. doi: 10.1590/1413-81232022278.04972022
18. Hallal PC, Horta BL, Barros AJD, Dellagostin OA, Hartwig FP, Pellanda LC, et al. Evolução da prevalência de infecção por COVID-19 no Rio Grande do Sul, Brasil: inquéritos sorológicos seriados. *Ciência Saúde Coletiva*. 2020;25:2395-401. doi: 10.1590/1413-81232020256.1.09632020.
19. Oliveira EC, Terças-Trettel AC, Andrade AC, Muraro AP, Santos ES, Espinosa MM, et al. Prevalência de anticorpos contra SARS-CoV-2 em Mato Grosso, Brasil: pesquisa de base populacional. *Cad Saúde Pública*. 2022;38:e00093021. doi: 10.1590/0102-311XPT093021.
20. Dean AG, Sullivan KM, Soe MM. *OpenEpi: Open Source Epidemiologic Statistics for Public Health*; 2013.
21. American Psychiatric Association. *Manual Diagnóstico e Estatístico de Transtornos Mentais-DSM-V*. Chicago: American Psychiatric Publishing; 2013.
22. Gao J, Zheng P, Jia Y, Chen H, Mao Y, Chen S, et al. Mental health problems and social media exposure during COVID-19 outbreak. *PLoS One*. 2020;15:e0231924. doi: 10.1371/journal.pone.0231924.
23. Lei L, Huang X, Zhang S, Yang J, Yang L, Xu M. Comparison of Prevalence and Associated Factors of Anxiety and Depression Among People Affected by versus People Unaffected by Quarantine During the COVID-19 Epidemic in Southwestern China. *Med Sci Monit*. 2020;26:e924609. doi: 10.12659/MSM.924609.
24. Li JB, Yang A, Dou K, Cheung RYM. Self-Control Moderates the Association Between Perceived Severity of Coronavirus Disease 2019 (COVID-19) and Mental Health Problems Among the Chinese Public. *Int J Environ Res Public Health*. 2020;4,17:4820. doi: 10.3390/ijerph17134820.
25. Mazza C, Ricci E, Biondi S, Colasanti M, Ferracuti S, Napoli C, et al. A Nationwide Survey of Psychological Distress among Italian People during the COVID-19 Pandemic: Immediate Psychological Responses and Associated Factors. *Int J Environ Res Public Health*. 2020;2,17: 3165. doi: 10.3390/ijerph17093165.
26. Zhang J, Lu H, Zeng H, Zhang S, Du Q, Jiang T, et al. The differential psychological distress of populations affected by the COVID-19 pandemic. *Brain Behav Immunity*. 2020;87:49-50. doi: 10.1016/j.bbi.2020.04.031.
27. Balakrishnan V, Ng KS, Kaur W, Govaichelvan K, Lee ZL. COVID-19 depression and its risk factors in Asia Pacific - A systematic review and meta-analysis. *J Affect Disord*. 2022;298:47-56. doi: 10.1016/j.jad.2021.11.048 See More.
28. Pfefferbaum B, North CS. Mental health and the Covid-19 pandemic. *New Engl J Med*. 2020;383:510-2. doi: 10.1056/NEJMp2008017.
29. Tsang S, Avery AR, Duncan GE. Fear and depression linked to COVID-19 exposure A study of adult twins during the COVID-19 pandemic. *Psychiatry Res*. 2021;296:113699. doi: 10.1016/j.psychres.2020.113699.
30. Rossell SL, Neill E, Phillipou A, Tan EJ, Toh WL, Van Rheenen TE, et al. An overview of current mental health in the general population of Australia during the COVID-19 pandemic: Results from the COL-LATE project. *Psychiatry Res*. 2021;296:113660. doi: 10.1016/j.psychres.2020.113660.
31. Flentje A, Obedin-Maliver J, Lubensky ME, Dastur Z, Neilands T, Lunn MR. Depression and anxiety changes

- among sexual and gender minority people coinciding with onset of COVID-19 pandemic. *J Gen Intern Med.* 2020;35:2788–90. doi: 10.1007/s11606-020-05970-4.
32. Moore SE, Wierenga KL, Prince DM, Gillani B, Mintz LJ. Disproportionate Impact of the COVID-19 Pandemic on Perceived Social Support, Mental Health and Somatic Symptoms in Sexual and Gender Minority Populations. *J Homosex.* 2021;68:577-91. doi: 10.1080/00918369.2020.1868184.
 33. Peckham H, de Gruijter NM, Raine C, Radziszewska A, Ciurtin C, Wedderburn LR, et al. Male sex identified by global COVID-19 meta-analysis as a risk factor for death and ICU admission. *Nat Commun.* 2020;11:6317. doi: 10.1038/s41467-020-19741-6
 34. Galvão MHR, Roncalli AG. Fatores associados a maior risco de ocorrência de óbito por COVID-19: análise de sobrevivência com base em casos confirmados. *Rev Brasil Epidemiol.* 2020;23:e200106. doi: 10.1590/1980-549720200106.
 35. Tibubos AN, Otten D, Ernst M, Beutel ME. A Systematic Review on Sex- and Gender-Sensitive Research in Public Mental Health During the First Wave of the COVID-19 Crisis. *Front Psychiatry.* 2021;12:712492. doi: 10.3389/fpsy.2021.712492.
 36. Brito VP, Carrijo AMM, Machado LMG, Nascimento VF, Oliveira SV. Perfil epidemiológico dos casos de suicídio no estado de Mato Grosso de 2008 a 2018: uma análise descritiva e quantitativa. In: Oliveira SV. *Pesquisas em saúde: Relatos, Fundamentos e Práticas.* Uberlândia: Editora Colab; 2021.
 37. Liu J, Zhu Q, Fan W, Makamure J, Zheng C, Wang J. Online mental health survey in a medical college in china during the COVID-19 outbreak. *Front Psychiatry.* 2020;11:459. doi: 10.3389/fpsy.2020.00459.
 38. Britt-Spells AM, Sledobnik M, Sands LP, Rollock D. Effects of perceived discrimination on depressive symptoms among Black men residing in the United States: A meta-analysis. *Am J Men Health.* 2018;12:52–63. doi: 10.1177/1557988315624509.
 39. Miller JA. Examining the relationship between discrimination and depressive symptomatology in African American men: A literature review utilizing a metanarrative approach. *J Am Psychiatric Nurs Assoc.* 2021;27:111–22. doi: 10.1177/1078390320907680.
 40. Smolen JR, Araújo E. Raça/cor da pele e transtornos mentais no Brasil: uma revisão sistemática. *Ciência & Saúde Coletiva.* 2017;22(12):4021-4030. doi: 10.1590/1413-812320172212.19782016.
 41. Almeida-Filho N, Lessa I, Magalhães L, Araújo MJ, Aquino E, James AS. Et al. Social inequality and depressive disorders in Bahia, Brazil: interactions of gender, ethnicity, and social class. *Soc Sci Med.* 2004;59:1339-53. doi: 10.1016/j.socscimed.2003.11.037.
 42. Williams DR, Yu Y, Jackson JS, Anderson NB. Racial differences in physical and mental health: socioeconomic status, stress and discrimination. *J Health Psychol.* 1997;2:335-351.
 43. Yelton B, Friedman DB, Noblet S, Lohman MC, Arant MA, Macaуда MM, et al. Social Determinants of Health and Depression among African American Adults: A Scoping Review of Current Research. *Int J Environ Res Public Health.* 2022;19:1498. doi: 10.3390/ijerph19031498.
 44. Frankham C, Richardson T, Maguire N. Psychological factors associated with financial hardship and mental health: A systematic review. *Clin Psychol Rev.* 2020;77:101832. doi: 10.1016/j.cpr.2020.101832.
 45. Robillard R, Daros AR, Phillips JL, Porteous M, Saad M, Pennestri MH, et al. Emerging New Psychiatric Symptoms and the Worsening of Pre-existing Mental Disorders during the COVID-19 Pandemic: a Canadian Multisite Study. *Can J Psychiatry.* 2021;66:815-26. doi: 10.1177/0706743720986786.
 46. Ganson KT, Tsai AC, Weiser SD, Benabou SE, Nagata JM. Job Insecurity and Symptoms of Anxiety and Depression Among U.S. Young Adults During COVID-19. *J Adolesc Health.* 2021;68:53-6. doi: 10.1016/j.jadohealth.2020.10.008.
 47. López-Bueno R, Calatayud J, Ezzatvar Y, Casajús JA, Smith L, Andersen LL et al. Association Between Current Physical Activity and Current Perceived Anxiety and Mood in the Initial Phase of COVID-19 Confinement. *Front Psychiatry.* 2020;11:729. doi: 10.3389/fpsy.2020.00729.
 48. Dun Y, Ripley-Gonzalez JW, Zhou N, Li Q, Chen M, Hu Z, et al. The association between prior physical fitness and depression in young adults during the COVID-19 pandemic-a cross-sectional, retrospective study. *PeerJ.* 2021;11:e11091. doi: 10.7717/peerj.11091.
 49. Stanton R, To QG, Khalesi S, Williams SL, Alley SJ, Thwaite TL, et al. Depression, anxiety and stress during COVID-19: associations with changes in physical activity, sleep, smoking and alcohol use in Australian adults. *Int J Environ Res Public Health.* 2020;17:4065. doi: 10.3390/ijerph17114065.
 50. Fullana M.A. , D. Hidalgo-Mazzei, E. Vieta, J. Radua. Coping behaviors associated with decreased anxiety and depressive symptoms during the COVID-19 pandemic and lockdown. *J Affect Disord.* 2020;275:80-1. doi: 10.1016/j.jad.2020.06.027.
 51. Chi X, Liang K, Chen S, Huang Q, Huang L, Yu Q, et al. Mental health problems among Chinese adolescents during the COVID-19: The importance of nutrition and physical activity. *Int J Clin Health Psychol.* 2021;21. doi: 10.1016/j.ijchp.2020.100218.
 52. Stefan N, Birkenfeld AL, Schulze MB. Global pandemics interconnected - obesity, impaired metabolic health and COVID-19. *Nat Rev Endocrinol.* 2021;17:135-49. doi: 10.1038/s41574-020-00462-1.
 53. Hao W, Tang Q, Huang X, Ao L, Wang J, Xie D. Analysis of the prevalence and influencing factors of depression and anxiety among maintenance dialysis patients during the COVID-19 pandemic. *Int Urol Nephrol.* 2021;53:1453-61. doi: 10.1007/s11255-021-02791-0.
 54. Faro JM, Mattocks KM, Nagawa CS, Lemon SC, Wang B, Cutrona SL, et al. Physical Activity, Mental Health, and Technology Preferences to Support Cancer Survivors During the COVID-19 Pandemic:

- Cross-sectional Study. *JMIR Cancer*. 2021;3;7:e25317. doi: 10.2196/25317.
55. Yadav R, Yadav P, Kumar SS, Kumar R. Assessment of Depression, Anxiety, and Sleep Disturbance in COVID-19 Patients at Tertiary Care Center of North India. *J Neurosci Rural Pract*. 2021;12:316-22. doi: 10.1055/s-0040-1722811.
 56. Hazarika M, Das S, Bhandari SS, Sharma P. The psychological impact of the COVID-19 pandemic and associated risk factors during the initial stage among the general population in India. *Open J Psychiatry Allied Sci*. 2021;12:31-5. doi: 10.5958/2394-2061.2021.00009.4.
 57. Suzuki K, Numao A, Komagamine T, Haruyama Y, Kawasaki A, Funakoshi K, et al. Impact of the COVID-19 Pandemic on the Quality of Life of Patients with Parkinson's Disease and Their Caregivers: A Single-Center Survey in Tochigi Prefecture. *J Parkinsons Dis*. 2021;11:1047-56. doi: 10.3233/JPD-212560.
 58. Ornell F, Moura HF, Scherer JN, Pechansky F, Kessler FHP, von Diemen L. The COVID-19 pandemic and its impact on substance use: Implications for prevention and treatment. *Psychiatry Res*. 2020;289:113096. doi: 10.1016/j.psychres.2020.113096.
 59. Da BL, Im GY, Schiano TD. Coronavirus Disease 2019 Hangover: A Rising Tide of Alcohol Use Disorder and Alcohol-Associated Liver Disease. *Hepatology*. 2020;72: 1102-1108. doi: 10.1002/hep.31307.
 60. Finlay I, Gilmore I. Covid-19 and alcohol—a dangerous cocktail. *BMJ*. 2020;20: m1987. doi: 10.1136/bmj.m1987.
 61. Rehm J, Kilian C, Ferreira-Borges C, Jernigan D, Monteiro M, Parry CDH, et al. Alcohol use in times of the COVID 19: Implications for monitoring and policy. *Drug Alcohol Rev*. 2020;39:301-4. doi: 10.1111/dar.13074
 62. Chu KM, Marco JL, Owolabi EO, Duvenage R, Londani M, Lombard C, et al. Trauma trends during COVID-19 alcohol prohibition at a South African regional hospital. *Drug Alcohol Rev*. 2022;41:13-9. doi: 10.1111/dar.13310.
 63. Mahadevan J, Shukla L, Benegal V. Alcohol controls in the aftermath of the COVID-19 pandemic in India: commentary on Stockwell. *Drug Alcohol Rev*. 2021;40:10-2. doi: 10.1111/dar.13158.
 64. Schmidt RA, Genois R, Jin J, Vigo D, Rehm J, Rush B. The early impact of COVID-19 on the incidence, prevalence, and severity of alcohol use and other drugs: A systematic review. *Drug Alcohol Depend*. 2021;1:109065. doi: 10.1016/j.drugaldep.2021.109065.
 65. Bartoszek A, Walkowiak D, Bartoszek A, Kardas G. Mental well-being (depression, loneliness, insomnia, daily life fatigue) during COVID-19 related home-confinement—a study from Poland. *Int J Environ Res Public Health*. 2020;17:7417. doi: 10.3390/ijerph17207417.
 66. Dozois DJA. Anxiety and depression in Canada during the COVID-19 pandemic: a national survey. *Can Psychol*. 2021;62:136-42. doi: 10.1037/cap0000251.
 67. Czeisler MÉ, Howard ME, Robbins R, Barger ER, Facer Childs SMW, Rajaratnam CA. COVID-19: public compliance with and public support for stay-at-home mitigation strategies. *medRxiv*. 2020.2004.2022.20076141
 68. Neill ED, Meyer WL, Toh TE, van Rheenen A, Philipou EJ, Tan SL. Alcohol use in Australia during the early days of the COVID-19 pandemic: initial results from the COLLATE project. *Psychiatry Clin Neurosci*. 2020;74:542-9. doi: 10.1111/pcn.13099.
 69. Taylor S, Paluszek MM, Rachor GS, McKay D, Asmundson GJG. Substance use and abuse, COVID-19-related distress, and disregard for social distancing: A network analysis. *Addict Behav*. 2021;114:106754. doi: 10.1016/j.addbeh.2020.106754
 70. Chodkiewicz J, Talarowska M, Miniszewska J, Nawrocka N, Bilinski P. Alcohol Consumption Reported during the COVID-19 Pandemic: The Initial Stage. *Int J Environ Res Public Health*. 2020;17:4677. doi: 10.3390/ijerph17134677.
 71. Barbosa LNF, Asfora GCA, Moura M. Ansiedade e depressão e uso de substâncias psicoativas em jovens universitários. *SMAD. Rev Eletrônica Saúde Mental Álcool Drogas*. 2020;16:1-8. doi: 10.11606/issn.1806-6976.smad.2020.155334
 72. Baglioni C, Battagliese G, Feige B, Spiegelhalder K, Nissen C, Voderholzer U. et al. Insomnia as a predictor of depression: a meta-analytic evaluation of longitudinal epidemiological studies. *J Affect Disord*. 2011;135:10-9. doi: 10.1016/j.jad.2011.01.011.
 73. Barros MBA, Lima MG, Ceolim M.F, Zancanella E, Cardoso TAMO. Quality of sleep, health status and subjective well-being in a population-based study. *Rev Saúde Pública*. 2019;53:82. doi: 10.11606/s1518-8787.2019053001067.
 74. Barros MBA, Lima MG, Malta DC, Szwarcwald C, Landmann ARCS, Romero D. et al. Relato de tristeza/depressão, nervosismo/ansiedade e problemas de sono na população adulta brasileira durante a pandemia de COVID-19. *Epidemiol Serv Saúde*. 2020;29:e2020427. doi: 10.1590/S1679-49742020000400018
 75. Barua L, Zaman MS, Omi FR, Faruque M. Psychological burden of the COVID-19 pandemic and its associated factors among frontline doctors of Bangladesh: a cross-sectional study. *F1000Research*. 2020;6:1304. doi: 10.12688/f1000research.27189.3.
 76. Di Renzo L, Gualtieri P, Pivari F, Soldati L, Attina A, Cinelli G, et al. Eating habits and lifestyle changes during COVID-19 lockdown: an Italian survey. *J Transl Med*. 2020;18:229. doi: 10.1186/s12967-020-02399-5
 77. Wang Y, Shi L, Que J, et al. The impact of quarantine on mental health status among general population in China during the COVID-19 pandemic. *Mol Psychiatry*. 2021;26:4813–22. doi: 10.1038/s41380-021-01019-y.
 78. Khoury JE, Atkinson L, Bennett T, Jack SM, Gonzalez A. COVID-19 and mental health during pregnancy: The importance of cognitive appraisal and social support. *J Affect Disord*. 2021;1:1161-9. doi: 10.1016/j.jad.2021.01.027.

79. Liu S, Yang L, Zhang C, Xu Y, Cai L, Ma S, et al. Gender differences in mental health problems of healthcare workers during the coronavirus disease 2019 outbreak. *J Psychiatry Res.* 2021;137:393-400. doi: 10.1016/j.jpsychires.2021.03.014
80. Garcia LP, Duarte E. Infodemia: excesso de quantidade em detrimento da qualidade das informações sobre a COVID-19. *Epidemiol Serv Saúde.* 2020;29:e2020186. doi: 10.1590/s1679-49742020000400019.
81. Organização Mundial de Saúde. Pandemia de COVID-19 desencadeia aumento de 25% na prevalência de ansiedade e depressão em todo o mundo: Chamado de alerta a todos os países para intensificar os serviços e apoio de saúde mental. Genebra: OMS; 2022. [consultado Jan 2022] Disponível em: <https://www.who.int/news/item/02-03-2022-covid-19-pandemic-triggers-25-increase-in-prevalence-of-anxiety-and-depression-worldwide>
82. Thornicroft G. Most people with mental illness are not treated. *Lancet.* 207;370:807-8. doi: 10.1016/S0140-6736(07)61392-0.
83. Kola L, Kohrt BA, Hanlon C. et al. COVID-19 mental health impact and responses in low-income and middle-income countries: reimagining global mental health. *Lancet Psychiatry.* 2021;8:535-50. doi: 10.1016/S2215-0366(21)00025-0.
84. Huang Y, Zhao N. Generalized anxiety disorder, depressive symptoms and sleep quality during the COVID-19 outbreak in China: a cross-sectional web-based survey. *Psychiatry Res.* 2020;288:112954. doi: 10.1016/j.psychres.2020.112954.
85. Roy D, Tripathy S, Kar SK, Sharma N, Verma SK, Kaushal V. Study of knowledge, attitude, anxiety and perceived need for mental health in the Indian population during the COVID-19 pandemic. *Asian J Psychiatry.* 2020;51:102083. doi: 10.1016/j.ajp.2020.102083
86. Özdin S, Bayrak Özdin Ş. Levels and predictors of anxiety, depression and health anxiety during COVID-19 pandemic in Turkish society: The importance of gender. *International J Soc Psychiatry.* 2020;66:504-11. doi: 10.1177/0020764020927051.
87. Gusmão EES, Plutarco LW, Moura DP, Silva CL, Meneses GO. Esquemas desadaptativos, ansiedade e depressão: proposta de um modelo explicativo. *Rev Bras Terap Cognit.* 2017;13:29-38. doi: 10.5935/1808-5687.20170006.