

Human guanylate-binding proteins in intracellular pathogen detection, destruction, and host cell death induction

Rivera-Cuevas, Yolanda; Clough, Barbara; Frickel, Eva-Maria

DOI:

[10.1016/j.coi.2023.102373](https://doi.org/10.1016/j.coi.2023.102373)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Rivera-Cuevas, Y, Clough, B & Frickel, E-M 2023, 'Human guanylate-binding proteins in intracellular pathogen detection, destruction, and host cell death induction', *Current Opinion in Immunology*, vol. 84, 102373. <https://doi.org/10.1016/j.coi.2023.102373>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Review

Human guanylate-binding proteins in intracellular pathogen detection, destruction, and host cell death induction

Yolanda Rivera-Cuevas, Barbara Clough and Eva-Maria Frickel



Cell-intrinsic defense is an essential part of the immune response against intracellular pathogens regulated by cytokine-induced proteins and pathways. One of the most upregulated families of proteins in this defense system are the guanylate-binding proteins (GBPs), large GTPases of the dynamin family, induced in response to interferon gamma. Human GBPs (hGBPs) exert their antimicrobial activity through detection of pathogen-associated molecular patterns and/or damage-associated molecular patterns to execute control mechanisms directed at the pathogen itself as well as the vacuolar compartments in which it resides. Consequently, hGBPs are also inducers of canonical and noncanonical inflammasome responses leading to host cell death. The mechanisms are both cell-type and pathogen-dependent with hGBP1 acting as a pioneer sensor for intracellular invaders. This review focuses on the most recent functional roles of hGBPs in pathways of pathogen detection, destruction, and host cell death induction.

Address

Institute of Microbiology and Infection, School of Biosciences, University of Birmingham, Edgbaston B15 2TT, United Kingdom

Corresponding author: Frickel, E.-M. (e.frickel@bham.ac.uk)**Current Opinion in Immunology** 2023, **84**:102373This review comes from a themed issue on **Host pathogens**Edited by **Liz Hartland** and **Sunny Shin**

Available online xxxx

<https://doi.org/10.1016/j.coi.2023.102373>0952-7915/© 2023 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

enhances the expression of innate immunity-related proteins necessary to *detect* and *destroy* intracellular pathogens, a pathway that often results in cellular *death*. An important set of IFN γ -stimulated genes (ISG) encodes a group of guanylate-binding proteins (GBPs) that play a crucial role in the control of intracellular pathogens not only in mammals, but across the eukaryotic kingdom [1]. GBPs execute antimicrobial functions against viruses, numerous bacteria, including *Salmonella enterica*, *Francisella novicida*, *Neisseria meningitidis*, *Shigella flexneri*, *Chlamydia trachomatis*, *Yersinia enterocolitica*, *Legionella pneumophila*, *Burkholderia thailandensis*, *Moraxella catarrhalis*, *Listeria monocytogenes*, group-A *Streptococcus*, and *Mycobacterium bovis* and against the protozoan parasites *Toxoplasma gondii* and *Leishmania donovani* [2–7].

Humans encode 7 GBP isoforms (hGBPs) that are composed of three main domains: a large guanosine triphosphatase (GTPase) (LG) domain located at the N-terminus, followed by a middle domain, and finally a GTPase effector domain at the C-terminus [8,9]. Three of these isoforms, hGBP1, hGBP2, and hGBP5, contain a CaaX-prenylation site at their C-terminus that facilitates membrane association [10,11]. GBPs' ability to physically interact with microbial surfaces or their replication compartments depend on this post-translational modification. Furthermore, heterodimerization enables non-prenylated GBP isoforms to be recruited to the site of prenylated GBP in a hierarchical manner, initiated by hGBP1 [10]. The biochemical properties of hGBP function during infection are expertly reviewed in [2].

This review compiles the recent literature of hGBPs in bacterial and protozoan microbial control and induction of host cell death. We partially include studies of murine Gbps (mGbps) to contrast and better understand the function of these proteins in pathogen-targeting destruction and induction of host cell death.


Introduction

Cells have intrinsic mechanisms for the *detection* of microbial infections through pattern recognition receptors (PRRs) that recognize pathogen-associated molecular patterns (PAMPs) resulting in the production of cytokines necessary to control the infection. Secretion of the inflammatory cytokine interferon gamma (IFN γ)

Cytosolic pathogen detection and destruction by guanylate-binding proteins

At early stages of infection, intracellular pathogens enter the host cell and reside in a membrane-bound compartment, known as the pathogen-containing vacuole (PCV). As part of the host's intrinsic defenses, this compartment should be targeted to the lysosome for

Figure 1



Mechanisms for hGBP-driven pathogen detection, destruction, and cellular death. **Detect:** hGBP1 targets invading microbes present in the host cytosol, in broken PCV, or intact PCVs. hGBP1 can detect cytosolic bacteria via its ability to bind LPS and upon membrane damage, demarked by Galectin-3 recruitment. Exactly how hGBP1 recognizes intact PCVs remains to be determined; however, ubiquitination has been proposed to play a role in this process. **Destroy:** hGBP1 binding to bacteria LPS affects membrane fluidity, making the pathogen more susceptible to antimicrobial peptides and/or direct killing by the APOL3. hGBP1 can also actively lyse intact PCV via an unknown mechanism. **Death:** hGBP1 microbial targeting results in PAMP release or it can initiate the recruitment of other hGBPs to cytosolic bacteria to initiate a signaling platform for inflammasome activation. hGBP1 function can result in noncanonical inflammasome (Caspase-4 activation) or canonical inflammasome activation (Caspase-1 activation), which primarily leads to pyroptotic cell death. However, in the case of *T. gondii* infection, hGBP1 releases parasite DNA that is detected by Absent In Melanoma 2 resulting in Caspase-8-dependent apoptotic cell death.

pathogen clearance. Pathogens often evade lysosome-mediated clearance by escaping from PCVs and establishing infection in the host cell cytosol. Emerging evidence uncovered a role for GBPs as cytosolic sensors to detect intracellular pathogens. During bacterial infection, hGBP1 binds to lipopolysaccharides (LPS) of Gram-negative bacteria exposed to the cell cytosol (Figure 1) [12–16]. The proposed model describes a detergent-like function of hGBP1 polymers that bind to LPS and subsequently diffuse from the polymer to the bacterial surface. This results in an increased outer membrane fluidity rendering the bacteria more susceptible to destruction by other immune-related effectors and/or antimicrobial peptides that otherwise would not have had access to the bacterial surface [2,14]. A recent study elucidated the contribution of hGBP1 in

apolipoprotein L3 (APOL3)-dependent killing of *S. enterica* serovar Typhimurium (*S. Typhimurium*) [17]. hGBP1 is shown to promote outer membrane permeability when interacting with the bacterial LPS, thus facilitating APOL3 access to the bacterial inner membrane for its dissolution [17] (Figure 1). The same surfactant-like activity of hGBP1 comes into play when binding *S. flexneri* with the consequence of inhibiting its actin tail formation [18,19]. In addition to its role in membrane fluidity, recent studies demonstrate a novel role for hGBP1 in bacterial LPS shedding into the host cytosol [20,21]. Furthermore, mGbp1 itself encodes a functional antimicrobial peptide-like sequence (mGbp1^{28–67}) in its globular domain, capable of targeting *F. novicida* and *N. meningitidis* but no other Gram-negative bacteria, including *S. Typhimurium* [4]. These data

suggest selectivity in Gbp1-mediated killing. Although the GBP1^{28–67} peptide is conserved across the murine and hGBPs [4], whether it functions similarly in hGBPs remains to be determined.

Recognition of broken and intact pathogen-containing vacuoles by guanylate-binding proteins

While recognition and clearance of cytosolic bacteria by GBP1 have undergone extensive study, less is known about the mechanisms for detection and control of intracellular pathogens encapsulated within a PCV. Currently, reports suggest sugar-binding galectins and/or ubiquitin as important contributors for GBP recruitment to the PCV [22–24] (Figure 1).

Galectins recognize sugar moieties present at the inner membrane leaflet that become exposed to the cell cytosol upon membrane damage, and PCV damage is often caused by the function of bacterial secretion systems [22,25]. Supporting this, it has been observed that recruitment of hGBP1 to *Y. enterocolitica*'s and group-A *Streptococcus*'s PCV occurs following membrane damage [7,26]. The damaged PCV, elicited by the bacterium itself through the function of its type-III secretion system (T3SS) or *Streptococcus* pore-forming protein, was recognized by the membrane damage sensor Galectin-3 and enabled hGBP1's access to the bacterial LPS [7,26]. Whether GBP1 also promotes membrane breakage in this scenario remains to be determined, however, hGBP1 was shown not to promote escape of *S. Typhimurium* from its PCV in both HeLa cells and macrophages [13,15]. Additionally, hGBP1 has been shown to detect sterile-damaged endosomes and this activity required the protein's GTPase activity, its farnesylation, and a triple-arginine polybasic motif at its C-terminus [19,24]. A potential role for Galectin-9 as an interactor of and contributor to mGbp2 recruitment to the *T. gondii* PCV has been proposed, but the precise state of the vacuole integrity at the time of recruitment of this complex is unclear [27].

GBPs also target pathogens that reside within intact PCVs, including *C. trachomatis* and *T. gondii* [23]. Whereas galectins seem to be contributing to the recruitment of GBPs toward damaged PCVs as a result of bacterial secretion systems [24], ubiquitination of the *C. trachomatis* and *T. gondii* PCV is an important marker for targeting the pathogens for clearance by cell-intrinsic defenses [23]. Interestingly, ubiquitination of *C. trachomatis* and *T. gondii* PCV has recently been demonstrated to be dependent on the E3 ubiquitin ligase RNF213 [28–30]. This provides a tool to assess whether ubiquitination plays a role upstream of hGBP1-mediated PCV detection and destruction (Figure 1). It is worthy to note that in HeLa cells, overexpressed hGBP1 targets latex

bead-containing phagosomes without IFN γ stimulation [31], while in THP-1 macrophages, IFN γ stimulation is needed to drive overexpressed hGBP1 to the *T. gondii* PCV [32]. An important gap in our understanding of GBP1-dependent pathogen control is how it differentiates PCVs from host organelles to avoid destroying self-lipid membranes. Lipid composition and phosphoinositide signatures of the PCV in interferon-stimulated cells probably play a role in the recognition mechanism. Murine immunity-related GTPases (Irgs) bind to certain phosphoinositides, a property that is reported to aid in their recognition of pathogen-containing compartments [33–35]. Another attractive hypothesis is that PCVs lack a self-molecule that enables GBP recognition akin to the 'missing self' hypothesis of murine Irg recruitment, where self-membranes are protected by murine Irgm proteins [36]. Active detection of pathogen proteins localized to the PCV is another possibility to drive GBP recruitment, or inversely, the presence of a negative regulator or a guard system that controls the cellular localization of GBP1.

How does guanylate-binding protein 1 drive membrane damage of intact vacuoles?

GBPs are part of the dynamin-like superfamily proteins, a group of GTPases that function in membrane-remodeling events such as fission or fusion through GTP hydrolysis. So, unsurprisingly, the hGBP1 isoform and its ortholog mGbp2 are capable of mediating tethering of giant unilamellar vesicles (GUV); however, the biological relevance of this observation in the context of their antimicrobial function remains unknown [11,31,37]. Although lipid composition was shown to be important for mGbp2 membrane binding and hinge motion, it was not sufficient to cause membrane curvature or damage [37]. Equally, hGBP1 recruitment to GUVs did not cause membrane damage to those compartments [11]. Therefore, it is possible that GBPs by themselves are not capable of causing membrane damage and that other proteins might be necessary. This is consistent with the observation that hGBP1-triggered membrane damage of *T. gondii* PCV is dependent on IFN γ stimulation, suggesting that there could be an undiscovered cofactor that is important [13]. We know that hGBP1 lyses PCVs of the bacteria *L. pneumophila* and the protozoan parasite *T. gondii*, but it remains unclear exactly how GBP1 disrupts these membranes (Figure 1) [13,38]. During *T. gondii* infection, hGBP1 isoprenylation and GTPase activity are important for PCV targeting in macrophages [39]. As a result, GTPase- and lipidation-deficient hGBP1 fail to control *T. gondii* infection [39]. Furthermore, a constitutive dimeric mutant was able to control *T. gondii* infection as effectively as wild-type hGBP1 [39]. This is consistent with structural analysis showing that hGBP1 lipid binding is facilitated by the conformational change employed upon GTPase activity and the stabilization of

the open state through dimerization [11]. Even though these data highlight important functional features in hGBP1 necessary for vacuolar pathogen immune control, the mechanism and coplayers driving GBP-mediated PCV membrane disruption remain elusive.

Guanylate-binding protein-dependent growth control independent of pathogen targeting

GBP-dependent pathogen control can also be exerted without active recruitment to the PCV. This is exemplified during infection with another protozoan parasite *L. donovani* where hGBP1 is necessary for parasite control, independent of PCV targeting [40]. A similar discovery was reported for hGBP1 control of *T. gondii* in A549 epithelial cells, and hGBP2 and hGBP5 *T. gondii* control in macrophages [39,41]. In the context of *L. donovani* infection, it appears that hGBP1 functions in pathogen restriction by promoting lysosome fusion with the PCV [42]. Although this could suggest a novel role for GBP1-dependent pathogen control, further elucidation of the molecular mechanisms involved in this process is required and how this might resemble the role of hGBP2 and hGBP5 in distal growth restriction of *T. gondii*.

Guanylate-binding protein-triggered host cell death

GBPs engage in canonical and noncanonical inflammasome activation as part of their role in cell-autonomous innate immune response (Figure 1) [43]. Inflammasomes are supramolecular complexes made of PRRs and inflammatory chemicals that, when activated, trigger pyroptotic cell death [44]. Most of our understanding of the role of GBPs in inflammasome activation comes from mouse GBPs (see reviews [2,43,45,46]). Here, multiple studies have focused on assessing the contribution of different mGbp and hGBP isoforms in pathogen control through host cell death [4,5,12,15,18,47].

hGBP1 contributes to noncanonical inflammasome activation through its role as a cytosolic LPS sensor, which results in Caspase-4 activation [12–15,21,48]. During *S. flexneri* and *S. Typhimurium* infection, it was observed that hGBP2, hGBP3, and hGBP4 form a complex that could target cytosolic bacteria but, unlike mGbps, localization of the different hGBP isoforms to the bacteria followed the initial recruitment of hGBP1 [12,15,18]. Thus, at least in human epithelial cells, there appears to be a hierarchy for the assembly and function of these proteins onto the bacterial surface. hGBP1 is the pioneering protein, hGBP4 and possibly hGBP2 control recruitment of Caspase-4, and hGBP3 binds LPS and drives Caspase-4 activation [12,15]. Cryo-electron tomography revealed insertion of hGBP1 molecules into the bacterial OM mediated by hGBP1 farnesylation and a C-terminal polybasic motif triggered release of LPS leading to activation of Caspase-4 [20]. The hGBP1

orthologs, mGbp2, mGbp1, Gbp3, and Gbp5, are also recruited to the cytosolic Gram-negative bacteria *M. catarrhalis* and *F. novicida* and are important for inflammasome activation [4,5]. Interestingly, a recent study employing *S. flexneri* mutants defective in antagonizing Caspase-4 and GBP1 bacterial binding or bacterial LPS release still observed pyroptosis [21,48]. Overexpressing hGBP2 or an hGBP1 defective in bacterial targeting restored pyroptotic cell death upon *S. flexneri* infection in cells deficient for hGBP1 [48]. hGBP1 and hGBP2 are capable of binding free LPS and acting as surfactants, clustering LPS molecules, and stimulating Caspase-4 activation [48].

hGBP5 was the first Gbp family member (murine or human) found to contribute to inflammasome activation by directly interacting with the intracellular PRR NOD-like receptor 3 (Nucleotide-binding and oligomerization domain–Leucine-rich repeat and pyrin domain-containing protein 3) [49]. The role of hGBP5 needs further attention to elicit its exact inflammasome-inducing mechanism, also in light of the recent finding that GBP5 deficiency leads to decreased proinflammatory mediator transcripts and secreted chemokines and cytokines [50]. Furthermore, the inflammasome can be activated through a novel pathway during *C. trachomatis* infection that is dependent on the GTPase activity of hGBP1 [51]. This study showed that whereas GTP hydrolysis is important for pathogen growth restriction, GMP produced by hGBP1 is metabolized to uric acid that can activate the inflammasome [51]. Thus, hGBP1's sequential GTP hydrolysis can employ different functions, including pathogen restriction and induction of cell death. Depending on the pathogen, hGBP1 can also drive host cell death to apoptosis. hGBP1-dependent disruption of the PCV and parasite membrane of *T. gondii* releases parasite DNA that is recognized by Absent In Melanoma 2 resulting in apoptosis-associated speck-like protein containing a caspase-recruitment domain Caspase-8-dependent death of human macrophages [13,20,32,52]. *S. Typhimurium*, however, elicits the expected hGBP1- and Caspase-4-dependent pyroptosis in human macrophages, but this pathogen also activates Caspase-1. In this case, hGBP1 is cleaved by Caspase-1, thereby reducing its recruitment to *S. Typhimurium* resulting in less Caspase-4 activation and reduced pyroptosis.

Conclusions/outlook

GBPs are important innate immune components that contribute to microbial control through pathogen detection and destruction and host cellular death induction. Current work has shed light on these functionalities, yet to fully understand GBPs, we have to address microbial counterdefense to GBPs, their cellular regulation, and targeting specificity. For GBP microbial counterdefense, the *Shigella* E3 IpaH9.8 is the only

identified microbial hGBP antagonist, ubiquitinating hGBP1 for proteasomal degradation [18,19,53], but there must be others. For cellular GBP regulation, what other post-translational mechanisms besides farnesylation control GBP function? Is there a broader role for caspase-driven regulation of GBP function that might extend into the extracellular space [13,54]? What additional mechanisms of ubiquitination impact GBP function? Does phosphorylation play a role in GBP regulation? For GBP targeting, can other PAMPs besides LPS be recognized and different GBPs distinguish LPS structural types [16]? How does hGBP1 target pathogens encapsulated within an intact vacuole and what entities are recognized on this non-self-membrane? What are the mechanisms of hGBP1-driven membrane disruption? GBP5 is robustly localized to the Golgi, yet we only know of its function to inhibit furin-mediated processing of viral glycoprotein at that location [55]. Employing GBP2/5 chimeras, bacterial targeting versus intracellular targeting to the Golgi are independent features in hGBPs [16]. How GBP targeting to self-membranes is regulated and what other functions specifically GBP5 performs at the Golgi are intriguing questions. Additionally, we await discovery of more GBP-interacting proteins that direct their function for the apparent selectivity in the context of different microbial infections and host cell death induction.

Beyond these immediate questions, it is important to understand that hGBPs vary dramatically in expression level depending on tissue and cell type and inflammatory condition [6]. Thus, some hGBPs will have cell-type-specific functions even when probing with the same pathogen infection. It is noteworthy, for example, that even after IFN γ stimulation, GBP3 and 4 show minimal expression in different human macrophages [32]. It is intriguing to note that GBP2, 3, and 4 generally are highly expressed at homeostasis, with GBP1 being almost absent and GBP5 demonstrating intermediate expression levels. Upon infection and inflammation, GBP1 is most highly upregulated. Whether this broadly will mean GBP1 (and possibly GBP5) are responsible for infection control, while other GBPs also serve additional homeostatic functions important for infection outcome, still needs to be established.

Data Availability

No data were used for the research described in the article.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could

have appeared to influence the work reported in this paper.

Acknowledgements

EMF received funding by Wellcome Trust Senior Research Fellowship 217202/Z/19/Z, an MRC Research and Innovation Grant MR/V030930/1, and Medical Research Council UK and Japan Agency for Medical Research and Development, UK-Japan (MRC-AMED) grant MR/T029323/1. This research was funded, in whole or in part, by The Wellcome Trust. A CC BY license is applied to the AAM arising from this submission, in accordance with the grant's open-access conditions.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Schelle L, Côte-Real JV, Esteves PJ, Abrantes J, Baldauf HM: **Functional cross-species conservation of guanylate-binding proteins in innate immunity.** *Med Microbiol Immunol* 2022, **212**:141-152, <https://doi.org/10.1007/s00430-022-00736-7>
2. Kutsch M, Coers J: **Human guanylate binding proteins: nanomachines orchestrating host defense.** *FEBS J* 2021, **288**:5826-5849, <https://doi.org/10.1111/febs.15662>
3. Zhang R, Li Z, Tang Y-D, Su C, Zheng C: **When human guanylate-binding proteins meet viral infections.** *J Biomed Sci* 2021, **28**:17, <https://doi.org/10.1186/s12929-021-00716-8>
4. Feng S, Enosi Tuipulotu D, Pandey A, Jing W, Shen C, Ngo C, *et al.*: **Pathogen-selective killing by guanylate-binding proteins as a molecular mechanism leading to inflammasome signaling.** *Nat Commun* (1) 2022, **13**:4395, <https://doi.org/10.1038/s41467-022-32127-0>.
This article describes pathogen-specific killing and inflammasome activation in mouse macrophages mediated by mGBPs. The molecular basis of this selectivity is determined by a peptide sequence at the N-terminus of the protein conserved in hGBPs and might be relevant to infection in human cells.
5. Enosi Tuipulotu D, Feng S, Pandey A, Zhao A, Ngo C, Mathur A, *et al.*: **Immunity against *Moraxella catarrhalis* requires guanylate-binding proteins and caspase-11-NLRP3 inflammasomes.** *EMBO J* 2023, **42**:e112558, <https://doi.org/10.15252/embj.2022112558>
6. Tretina K, Park E-S, Maminska A, MacMicking JD: **Interferon-induced guanylate-binding proteins: guardians of host defense in health and disease.** *J Exp Med* 2019, **216**:482-500, <https://doi.org/10.1084/jem.20182031>
7. Hikichi M, Toh H, Minowa-Nozawa A, Nozawa T, Nakagawa I: **Guanylate-binding protein 1 regulates infection-induced autophagy through TBK1 phosphorylation.** *Cell Microbiol* 2022, **2022**:1-18, <https://doi.org/10.1155/2022/8612113>.
This study demonstrates a role for hGBP1 in promoting autophagy of Group A *Streptococcus* through its interaction with TRAF Family Member Associated NFKB Activator binding kinase 1 (TBK1) and regulation of TBK1 phosphorylation. hGBP1 specifically localised to bacteria-induced membrane damage marked by galectin 3 recruitment.
8. Olszewski MA, Gray J, Vestal DJ: **In silico genomic analysis of the human and murine guanylate-binding protein (GBP) gene clusters.** *J Interferon Cytokine Res J Int Soc Interferon Cytokine Res* 2006, **26**:328-352, <https://doi.org/10.1089/jir.2006.26.328>
9. Prakash B, Praefcke GJK, Renault L, Wittinghofer A, Herrmann C: **Structure of human guanylate-binding protein 1 representing a unique class of GTP-binding proteins.** *Nature* 2000, **403**:567-571, <https://doi.org/10.1038/35000617>
10. Britzen-Laurent N, Bauer M, Berton V, Fischer N, Syguda A, Reipschläger S, *et al.*: **Intracellular trafficking of guanylate-binding proteins is regulated by heterodimerization in a hierarchical manner.** *PLoS One* 2010, **5**:e14246, <https://doi.org/10.1371/journal.pone.0014246>

6 Host pathogens

11. Sistemich L, Dimitrov Stanchev L, Kutsch M, Roux A, Günther Pomorski T, Herrmann C: **Structural requirements for membrane binding of human guanylate-binding protein 1**. *FEBS J* 2021, **288**:4098-4114, <https://doi.org/10.1111/febs.15703>.
This study demonstrates that GTP hydrolysis in hGBP1 drives dimerization and an open conformation of the protein, freeing the farnesyl tail of GBP1 to attach and assemble on artificial membranes in an upright configuration.
12. Wandel MP, Kim B-H, Park E-S, Boyle KB, Nayak K, Lagrange B, et al.: **Guanylate-binding proteins convert cytosolic bacteria into caspase-4 signaling platforms**. *Nat Immunol* 2020, **21**:880-891, <https://doi.org/10.1038/s41590-020-0697-2>
13. Fisch D, Clough B, Domart M-C, Encheva V, Bando H, Snijders AP, et al.: **Human GBP1 differentially targets Salmonella and Toxoplasma to license recognition of microbial ligands and caspase-mediated death**. *Cell Rep* 2020, **32**:108008, <https://doi.org/10.1016/j.celrep.2020.108008>
14. Kutsch M, Sistemich L, Lesser CF, Goldberg MB, Herrmann C, Coers J: **Direct binding of polymeric GBP1 to LPS disrupts bacterial cell envelope functions**. *EMBO J* 2020, **39**:e104926, <https://doi.org/10.15252/embj.2020104926>
15. Santos JC, Boucher D, Schneider LK, Demarco B, Dilucca M, Shkarina K, et al.: **Human GBP1 binds LPS to initiate assembly of a caspase-4 activating platform on cytosolic bacteria**. *Nat Commun* 2020, **11**:3276, <https://doi.org/10.1038/s41467-020-16889-z>
16. Valeva SV, Michal F, Degabriel M, Rohde JR, Randow F, Ernst RK: **Bacterial factors drive the differential targeting of Guanylate Binding Proteins to Francisella and Shigella**. *Pathog Dis.* 2023, **81**:ftad005, <https://doi.org/10.1093/femspd/ftad005>
17. Gaudet RG, Zhu S, Halder A, Kim B-H, Bradfield CJ, Huang S, et al.: **A human apolipoprotein L with detergent-like activity kills intracellular pathogens**. *Science* 2021, **373**:eabf8113, <https://doi.org/10.1126/science.abf8113>.
This study is the first to report the synergistic targeting and destruction of bacteria by ISGs hGBP1 and APOL3, through a sequential disruption of bacterial membranes in cytosolic *Salmonella*.
18. Wandel MP, Pathe C, Werner EI, Ellison CJ, Boyle KB, von der Malsburg A, et al.: **GBPs inhibit motility of Shigella flexneri but are targeted for degradation by the bacterial ubiquitin ligase IpaH9.8**. *Cell Host Microbe* 2017, **22**:507-518.e5, <https://doi.org/10.1016/j.chom.2017.09.007>
19. Piro AS, Hernandez D, Luoma S, Feeley EM, Finethy R, Yirga A, et al.: **Detection of cytosolic Shigella flexneri via a C-terminal triple-arginine motif of GBP1 inhibits actin-based motility**. *MBio* 2017, **8**:e01979-17, <https://doi.org/10.1128/mBio.01979-17>
20. Zhu S, Bradfield CJ, Mamińska A, Park E-S, Kim B-H, Kumar P, et al.: **Cryo-ET of a human GBP coatomer governing cell-autonomous innate immunity to infection**. 2021:2021.08.26.457804. (<https://doi.org/10.1101/2021.08.26.457804>).
21. Goers L, Kim K, Stedman TC, Canning PJ, Mou X, Ernst NH, et al.: **Shigella IpaH9.8 limits GBP1-dependent LPS release from intracytosolic bacteria to suppress caspase-4 activation**. *Proc Natl Acad Sci USA* 2023, **120**:e2218469120, <https://doi.org/10.1073/pnas.2218469120>
22. Coers J: **Sweet host revenge: galectins and GBPs join forces at broken membranes**. *Cell Microbiol* (12) 2017, **19**, <https://doi.org/10.1111/cmi.12793>
23. Haldar AK, Foltz C, Finethy R, Piro AS, Feeley EM, Pilla-Moffett DM, et al.: **Ubiquitin systems mark pathogen-containing vacuoles as targets for host defense by guanylate binding proteins**. *Proc Natl Acad Sci* 2015, **112**:E5628-E5637, <https://doi.org/10.1073/pnas.1515966112>
24. Feeley EM, Pilla-Moffett DM, Zwack EE, Piro AS, Finethy R, Kolb JP, et al.: **Galectin-3 directs antimicrobial guanylate binding proteins to vacuoles furnished with bacterial secretion systems**. *Proc Natl Acad Sci USA* 2017, **114**:E1698-E1706, <https://doi.org/10.1073/pnas.1615771114>
25. Bänfer S, Jacob R: **Galectins in intra- and extracellular vesicles**. *Biomolecules* 2020, **10**:1232, <https://doi.org/10.3390/biom10091232>
26. Rudolph M, Carsten A, Kulnik S, Aepfelbacher M, Wolters M: **Live imaging of Yersinia translocon formation and immune recognition in host cells**. *PLoS Pathog* 2022, **18**:e1010251, <https://doi.org/10.1371/journal.ppat.1010251>.
The authors use a high-resolution imaging technique to dissect the dynamic function of *Yersinia* T3SS pore complex during infection of human epithelial cells. The subsequent T3SS-induced disruption of phagosomal membranes led to sequential recruitment of galectin-3 and hGBP1.
27. Kravets E, Poschmann G, Hänsch S, Weidtkamp-Peters S, Degrandi D, Stühler K, et al.: **mGBP2 engages Galectin-9 and cytoskeleton-associated protein 4 for immunity against Toxoplasma gondii**. 2021:2021.12.28.474342. (<https://doi.org/10.1101/2021.12.28.474342>).
28. Hernandez D, Walsh S, Saavedra Sanchez L, Dickinson MS, Coers J: **Interferon-inducible E3 ligase RNF213 facilitates host-protective linear and K63-linked ubiquitylation of Toxoplasma gondii parasitophorous vacuoles**. *MBio* 2022, **13**:e0188822, <https://doi.org/10.1128/mbio.01888-22>
29. Walsh SC, Reitano JR, Dickinson MS, Kutsch M, Hernandez D, Barnes AB, et al.: **The bacterial effector GarD shields Chlamydia trachomatis inclusions from RNF213-mediated ubiquitylation and destruction**. *Cell Host Microbe* 2022, **30**:1671-1684.e9, <https://doi.org/10.1016/j.chom.2022.08.008>
30. Matta SK, Kohio HP, Chandra P, Brown A, Doench JG, Phillips JA, et al.: **Molecular basis for interferon-mediated pathogen restriction in human cells**. *Microbiology* 2022, <https://doi.org/10.1101/2022.10.21.513197>
31. Shydlovskiy S, Zienert AY, Ince S, Dovengerds C, Hohendahl A, Dargazanli JM, et al.: **Nucleotide-dependent farnesyl switch orchestrates polymerization and membrane binding of human guanylate-binding protein 1**. *Proc Natl Acad Sci* 2017, **114**:E5559-E5568, <https://doi.org/10.1073/pnas.1620959114>
32. Fisch D, Bando H, Clough B, Hornung V, Yamamoto M, Shenoy AR, et al.: **Human GBP1 is a microbe-specific gatekeeper of macrophage apoptosis and pyroptosis**. *EMBO J* 2019, **38**:e100926, <https://doi.org/10.15252/embj.2018100926>
33. Lee Y, Yamada H, Pradipta A, Ma JS, Okamoto M, Nagaoka H, et al.: **Initial phospholipid-dependent Irgb6 targeting to Toxoplasma gondii vacuoles mediates host defense**. *Life Sci Alliance* 2020, **3**:e201900549, <https://doi.org/10.26508/lsa.201900549>
34. Tiwari S, Choi H-P, Matsuzawa T, Pypaert M, MacMicking JD: **Targeting of the GTPase Irgm1 to the phagosomal membrane via PtdIns(3,4)P(2) and PtdIns(3,4,5)P(3) promotes immunity to mycobacteria**. *Nat Immunol* 2009, **10**:907-917, <https://doi.org/10.1038/ni.1759>
35. Saijo-Hamano Y, Sherif AA, Pradipta A, Sasai M, Sakai N, Sakihama Y, et al.: **Structural basis of membrane recognition of Toxoplasma gondii vacuole by Irgb6**. *Life Sci Alliance* 2022, **5**:e202101149, <https://doi.org/10.26508/lsa.202101149>
36. Dockterman J, Coers J: **How did we get here? Insights into mechanisms of immunity-related GTPase targeting to intracellular pathogens**. *Curr Opin Microbiol* 2022, **69**:102189, <https://doi.org/10.1016/j.mib.2022.102189>
37. Loschwitz J, Steffens N, Wang X, Schäffler M, Pfeffer K, Degrandi D, et al.: **Domain motions, dimerization, and membrane interactions of the murine guanylate binding protein 2**. *Sci Rep* 2023, **13**:679, <https://doi.org/10.1038/s41598-023-27520-8>.
The authors give insight into the specificity of mGBP membrane binding through membrane lipid composition. A preference of mGBP2 for cardiolipin led to dynamic changes in the protein structure which could influence membrane integrity. Selectivity for different lipids could determine membrane targeting for other prenylated GBPs and different pathogen membranes.
38. Bass AR, Shin S: **Human GBP1 promotes pathogen vacuole rupture and inflammasome activation during Legionella pneumophila infection**. 2020:2020.05.27.120477. (<https://doi.org/10.1101/2020.05.27.120477>).

39. Fisch D, Clough B, Khan R, Healy L, Frickel E-M: **Toxoplasma-proximal and distal control by GBPs in human macrophages.** *Pathog Dis* 2022, **79**:ftab058, <https://doi.org/10.1093/femspd/ftab058>.
- A key paper revealing that pathogen growth restriction in *Toxoplasma* can be mediated from a pathogen-distal position by hGBPs 2 and 5. In contrast, pathogen restriction and killing by hGBP1 required recruitment to PCVs accompanied by GTP hydrolysis.
40. Haldar AK, Nigam U, Yamamoto M, Coers J, Goyal N: **Guanylate binding proteins restrict *Leishmania donovani* growth in nonphagocytic cells independent of parasitophorous vacuolar targeting.** *MBio* 2020, **11**:e01464-20, <https://doi.org/10.1128/mBio.01464-20>
41. Johnston AC, Piro A, Clough B, Siew M, Virreira Winter S, Coers J, et al.: **Human GBP1 does not localize to pathogen vacuoles but restricts *Toxoplasma gondii*.** *Cell Microbiol* 2016, **18**:1056-1064, <https://doi.org/10.1111/cmi.12579>
42. Haldar AK: **Guanylate binding proteins promote anti-*Leishmania* host cell defense.** *J Cell Immunol* 2021, **3**:e01464-20, <https://doi.org/10.33696/immunology.3.113>
43. Kim B-H, Chee JD, Bradfield CJ, Park E-S, Kumar P, MacMicking JD: **IFN-induced guanylate binding proteins in inflammasome activation and host defense.** *Nat Immunol* 2016, **17**:481-489, <https://doi.org/10.1038/ni.3440>
44. Yi Y-S: **Functional crosstalk between non-canonical caspase-11 and canonical NLRP3 inflammasomes during infection-mediated inflammation.** *Immunology* 2020, **159**:142-155, <https://doi.org/10.1111/imm.13134>
45. Rafeld HL, Kolanus W, van Driel IR, Hartland EL: **Interferon-induced GTPases orchestrate host cell-autonomous defence against bacterial pathogens.** *Biochem Soc Trans* 2021, **49**:1287-1297, <https://doi.org/10.1042/BST20200900>
46. Johns CE, Galam L: **Guanylate binding protein 1 (GBP1): a key protein in inflammatory pyroptosis.** *Cell Biochem Biophys* 2022, **80**:295-299, <https://doi.org/10.1007/s12013-021-01056-y>
47. Park E-S, Kim B-H, Kumar P, Tretina K, Maminska A, Philbrick WM, et al.: **A hierarchical GBP network promotes cytosolic LPS recognition and sepsis.** 2021:2021.08.25.457662. (<https://doi.org/10.1101/2021.08.25.457662>).
48. Dickinson MS, Kutsch M, Sistemich L, Hernandez D, Piro AS, Needham D, et al.: **LPS-aggregating proteins GBP1 and GBP2 are each sufficient to enhance caspase-4 activation both in cellulo and in vitro.** *Proc Natl Acad Sci USA* 2023, **120**:e2216028120, <https://doi.org/10.1073/pnas.2216028120>.
- This study shows that both GBP1 and GBP2 can directly bind to and aggregate free LPS to enhance Caspase-4 activation and pyroptosis. This function of GBP1 and GBP2 is independent of GBP1's ability to form microcapsules on cytosolic bacteria, with GBP2 needing GBP1 to target to bacteria.
49. Shenoy AR, Wellington DA, Kumar P, Kassa H, Booth CJ, Cresswell P, et al.: **GBP5 promotes NLRP3 inflammasome assembly and immunity in mammals.** *Science* 2012, **336**:481-485, <https://doi.org/10.1126/science.1217141>
50. Li Y, Lin X, Wang W, Wang W, Cheng S, Huang Y, et al.: **The proinflammatory role of guanylate-binding protein 5 in inflammatory bowel diseases.** *Front Microbiol* 2022, **13**:926915, <https://doi.org/10.3389/fmicb.2022.926915>
51. Xavier A, Al-Zeer MA, Meyer TF, Daumke O: **hGBP1 coordinates *Chlamydia* restriction and inflammasome activation through sequential GTP hydrolysis.** *Cell Rep* 2020, **31**:107667, <https://doi.org/10.1016/j.celrep.2020.107667>
52. Dickinson MS, Kutsch M, Sistemich L, Hernandez D, Piro AS, Needham D: **GBP2 aggregates LPS and activates the caspase-4 inflammasome independent of the bacterial encapsulation factor GBP1.** *Proc Natl Acad Sci U S A* (15) 2023, **120**:e2216028120, <https://doi.org/10.1073/pnas.2216028120>
53. Li P, Jiang W, Yu Q, Liu W, Zhou P, Li J, et al.: **Ubiquitination and degradation of GBPs by a *Shigella* effector to suppress host defence.** *Nature* 2017, **551**:378-383, <https://doi.org/10.1038/nature24467>
54. Naschberger E, Geißdörfer W, Bogdan C, Tripal P, Kremmer E, Stürzl M, et al.: **Processing and secretion of guanylate binding protein-1 depend on inflammatory caspase activity.** *J Cell Mol Med* 2017, **21**:1954-1966, <https://doi.org/10.1111/jcmm.13116>
55. Braun E, Hotter D, Koepke L, Zech F, Groß R, Sparrer KMJ, et al.: **Guanylate-binding proteins 2 and 5 exert broad antiviral activity by inhibiting furin-mediated processing of viral envelope proteins.** *Cell Rep* 2019, **27**:2092-2104.e10, <https://doi.org/10.1016/j.celrep.2019.04.063>