

Dopamine dysregulation in Parkinson's disease flattens the pleasurable urge to move to musical rhythms

Pando-Naude, Victor; Matthews, Tomas Edward; Højlund, Andreas; Jakobsen, Sebastian; Østergaard, Karen; Johnsen, Erik; Garza-Villarreal, Eduardo A.; Witek, Maria A. G.; Penhune, Virginia; Vuust, Peter

DOI:
[10.1111/ejn.16128](https://doi.org/10.1111/ejn.16128)

License:
Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (Harvard):
Pando-Naude, V, Matthews, TE, Højlund, A, Jakobsen, S, Østergaard, K, Johnsen, E, Garza-Villarreal, EA, Witek, MAG, Penhune, V & Vuust, P 2023, 'Dopamine dysregulation in Parkinson's disease flattens the pleasurable urge to move to musical rhythms', *European Journal of Neuroscience*.
<https://doi.org/10.1111/ejn.16128>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.








When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Dopamine dysregulation in Parkinson's disease flattens the pleasurable urge to move to musical rhythms

Victor Pando-Naude¹  | Tomas Edward Matthews¹  | Andreas Højlund^{2,3}  |
 Sebastian Jakobsen^{2,3}  | Karen Østergaard^{2,4,5}  | Erik Johnsen^{2,4} |
 Eduardo A. Garza-Villarreal⁶ | Maria A. G. Witek⁷ | Virginia Penhune⁸  |
 Peter Vuust¹ 

¹Center for Music in the Brain, Department of Clinical Medicine, Aarhus University & The Royal Academy of Music Aarhus/Aalborg, Aarhus, Denmark

²Center of Functionally Integrative Neuroscience, Department of Clinical Medicine, Aarhus University Hospital, Aarhus, Denmark

³Department of Linguistics, Cognitive Science and Semiotics, School of Communication and Culture, Aarhus University, Aarhus, Denmark

⁴Department of Neurology, Aarhus University Hospital, Aarhus, Denmark

⁵Sano, Private Hospital, Aarhus, Denmark

⁶Instituto de Neurobiología, Universidad Nacional Autónoma de México (UNAM), Juriquilla, Querétaro, Mexico

⁷Department of Music School of Languages, Cultures, Art History and Music, University of Birmingham, Birmingham, UK

⁸Department of Psychology, Concordia University, Montreal, Quebec, Canada

Correspondence

Victor Pando-Naude, Center for Music in the Brain (MIB), Department of Clinical Medicine, Aarhus University, Universitetsbyen, 3-0-17, 8000 Aarhus, Denmark.
 Email: pandonaude@clin.au.dk

Funding information

Danish National Research Foundation, Grant/Award Number: DNRF 117

Edited by: Yoland Smith

Abstract

The pleasurable urge to move to music (PLUMM) activates motor and reward areas of the brain and is thought to be driven by predictive processes. Dopamine in motor and limbic networks is implicated in beat-based timing and music-induced pleasure, suggesting a central role of basal ganglia (BG) dopaminergic systems in PLUMM. This study tested this hypothesis by comparing PLUMM in participants with Parkinson's disease (PD), age-matched controls, and young controls. Participants listened to musical sequences with varying rhythmic and harmonic complexity (low, medium and high), and rated their experienced pleasure and urge to move to the rhythm. In line with previous results, healthy younger participants showed an inverted U-shaped relationship between rhythmic complexity and ratings, with preference for medium complexity rhythms, while age-matched controls showed a similar, but weaker, inverted U-shaped response. Conversely, PD showed a significantly flattened response for both the urge to move and pleasure. Crucially, this flattened response could not be attributed to differences in rhythm discrimination and did not reflect an overall decrease in ratings. For harmonic

Victor Pando-Naude and Tomas Edward Matthews shared first authorship.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2023 The Authors. *European Journal of Neuroscience* published by Federation of European Neuroscience Societies and John Wiley & Sons Ltd.

complexity, PD showed a negative linear pattern for both the urge to move and pleasure while healthy age-matched controls showed the same pattern for pleasure and an inverted U for the urge to move. This contrasts with the pattern observed in young healthy controls in previous studies, suggesting that both healthy aging and PD also influence affective responses to harmonic complexity. Together, these results support the role of dopamine within cortico-striatal circuits in the predictive processes that form the link between the perceptual processing of rhythmic patterns and the affective and motor responses to rhythmic music.

KEYWORDS

basal ganglia, dopamine, harmony, music-induced pleasure, Parkinson's disease, predictive processes, rhythm

1 | INTRODUCTION

When listening to rhythmic music, we often feel an automatic and pleasurable urge to tap our feet or bob our heads along with the underlying beat or pulse of the rhythm. This pleasurable urge to move to music (PLUMM) is a key component of groove (Câmara & Danielsen, 2018; Duman et al., 2022; Janata et al., 2012; Madison, 2006; Senn et al., 2019; Witek et al., 2014) and highlights the deep connection between music, movement and pleasure. In a recent neuroimaging study, we showed that PLUMM is associated with greater activity in dorsal and ventral striatal regions within the basal ganglia (BG), along with cortical regions associated with the motor and limbic cortico-striato-thalamo-cortical circuits (Alexander et al., 1986; Matthews et al., 2020) (henceforth, cortico-striatal circuits). This aligns with previous work implicating these loops in the predictive processes thought to drive both rhythm perception and music-induced pleasure, along with motor timing and reward processes more generally (Pando-Naude et al., 2021). For example, the motor cortico-striatal circuit is consistently engaged during rhythm and beat perception and production (Bengtsson et al., 2009; Chen et al., 2008; Grahn & Rowe, 2009, 2013; Kasdan et al., 2022; Schubotz et al., 2000), whereas the limbic cortico-striatal circuit is implicated in music-induced pleasure (Blood et al., 1999; Blood & Zatorre, 2001; Cheung et al., 2019; Gold et al., 2019; Salimpoor et al., 2011, 2013; Shany et al., 2019). Within cortico-striatal circuits, dopamine seems to have a functional role in both beat-based (Biswas et al., 2016; Cameron et al., 2016; Geiser & Kaelin-Lang, 2011; Grahn & Brett, 2009; Hsu et al., 2022) and motor timing (Jahanshahi et al., 2010), as well as music-induced pleasure (Ferreri et al., 2019; Gebauer et al., 2012; Salimpoor

et al., 2011). This confluence of musically relevant timing, motor and reward processes suggests a key role of dopamine in PLUMM (Matthews et al., 2020). To directly test this, we compared ratings of pleasure and the urge to move in two groups of participants with Parkinson's disease (PD), as well as age-matched and younger healthy controls.

Dopamine is thought to be necessary for the predictive processes that are fundamental to motor timing and reward mechanisms (Meck, 2006; Schultz et al., 1997; Tomassini et al., 2016), likely via separate pathways within motor and limbic cortico-striatal circuits. The motor cortico-striatal circuit, including the putamen, premotor cortex and supplementary motor area, encompasses the nigrostriatal dopaminergic pathway, wherein dopaminergic cells in the substantia nigra pars compacta project to the dorsal striatum, including the putamen and caudate. Nigrostriatal dopamine is implicated in timing processes in the hundreds of milliseconds range that are necessary for sensory anticipation and motor preparation (Behroozmand & Johari, 2019; Coull et al., 2012; Jones & Jahanshahi, 2014; Meck, 2006). The limbic cortico-striatal circuit, including the nucleus accumbens and medial prefrontal cortices, encompasses the mesolimbic dopaminergic pathway, wherein cells in the ventral tegmental area project to the ventral striatum including the nucleus accumbens. Longstanding models suggest that mesolimbic dopamine encodes reward prediction errors, indicating whether a reward was better or worse than expected (Schultz et al., 1997). Recent empirical and theoretical work suggest an expansion of the role of striatal dopamine to include sensory prediction errors (when sensory outcomes deviate from predictions), as well as the certainty of sensorimotor, state or value representations (Friston et al., 2014, 2012; Gershman & Uchida, 2019).

Predictive processes are crucial to the perception and production of music, as well as affective responses including PLUMM. Music is often highly structured both in time and tonal space. Embellishments on or deviations from this structure can elicit prediction confirmations or errors and, in turn, affective and motor responses within listeners (Huron, 2006; Koelsch et al., 2019; Meyer, 1956; Salimpoor et al., 2015; Vuust et al., 2022; Zald & Zatorre, 2011). In terms of PLUMM, there is an inverted U-shaped relation between rhythmic complexity (predictability) and ratings of both the urge to move and pleasure, as the highest ratings are elicited by rhythms that are neither too simple (predictable) nor too complex (Matthews et al., 2019, 2022; Sioros et al., 2014; Spiech et al., 2022; Stupacher, Wrede, & Vuust, 2022; Witek et al., 2014). A predictive processing account suggests that this pattern is driven by both the number of rhythm-based temporal prediction errors and the certainty of the antecedent predictions (Koelsch et al., 2019; Vuust et al., 2018, 2022; Vuust & Witek, 2014). According to this framework, predictions arise from an internal model based on prior experience. In the case of rhythmic music, the model is the meter, which, in western musical traditions, is the pattern of strong beats (when a note is very likely to occur) and weak beats (when a note is less likely). Therefore, moderately complex, high-PLUMM rhythms are regular enough to enable relatively precise metrical predictions and complex enough to challenge them. Recently, harmonic complexity was also shown to influence PLUMM via its effect on pleasure, in contrast to rhythmic complexity which directly affects both the urge to move and pleasure (Matthews et al., 2019). This suggests that the urge to move and pleasure can be at least partially disentangled via the differential effects of rhythm and harmony.

The role of dopamine in temporal and reward-related predictive processes is well established (Hollerman & Schultz, 1998; Nomoto et al., 2010); however, it has yet to be linked to PLUMM. A recent theoretical model suggests that nigrostriatal dopamine plays a key role in probabilistic representations of the musical beat and thus supports beat-based motor and perceptual timing (Cannon & Patel, 2020). This is consistent with studies involving participants with PD, which is characterized by cell loss in the substantia nigra leading to dysregulated dopamine in the nigrostriatal pathway and thus problems with motor timing (e.g., dysfunctional gait and bradykinesia). Specifically, PD show poorer performance and altered brain activity during perceptual and motor tasks that rely on beat-based timing (Bellinger et al., 2017; Benoit et al., 2014; Biswas et al., 2016; Breska & Ivry, 2018; Cameron et al., 2016; Elsinger et al., 2003; Geiser & Kaelin-Lang, 2011; Grahn & Brett, 2009; Hsu et al., 2022;

Jones et al., 2008; Merchant et al., 2008; O'Boyle et al., 1996; Vikene et al., 2019a, 2019b). There is evidence that neural oscillations at the beat frequency and the beta frequency range (13–30 Hz) are involved in beat and meter perception and synchronization and thus may provide a neural mechanism underlying beat-based predictive processes (Large & Snyder, 2009). Intriguingly, participants with PD show weaker entrainment of beta-range neural oscillations to rhythmic stimuli (te Woerd et al., 2017), suggesting that beta entrainment may mediate the link between dopamine dysregulation and beat-based timing. More generally, one study showed that participants with PD had a reduced urge to engage with music via dancing or singing compared with healthy controls (Morris et al., 2019), suggesting a weakened link between music and movement. Finally, rhythmic auditory stimulation, for example, cueing participants' gait with a metronome, improves motor deficits in participants with PD (Bella et al., 2017; Lei et al., 2019; Pau et al., 2016; Thaut et al., 1996), further supporting the link between dopamine and predictive motor timing.

The BG and dopamine are also implicated in music-induced pleasure (Ferreri et al., 2019; Gebauer et al., 2012; Salimpoor et al., 2011). Activity in the limbic cortico-striatal circuit, as well as functional and structural connectivity within this loop, and between the striatum and auditory cortex, are associated with music liking (Blood et al., 1999; Blood & Zatorre, 2001; Salimpoor et al., 2013), music wanting (i.e., the money one is willing to spend on a given piece; Salimpoor et al., 2013) and music reward sensitivity (Martinez-Molina et al., 2019; Martínez-Molina et al., 2016). Activity in the nucleus accumbens has also been linked to the predictive processes thought to underlie music-derived pleasure (Cheung et al., 2019; Gold et al., 2019; Seger et al., 2013; Shany et al., 2019). Evidence of a causal role for striatal dopamine in music-induced pleasure more generally comes from two parallel lines of research showing that modulations in BG activity or dopamine availability results in corresponding modulations in liking and wanting of music (Ferreri et al., 2019; Mas-Herrero et al., 2018). However, many of these studies have focused on harmonic predictions rather than the temporal predictions that dominate in PLUMM (Matthews et al., 2019).

Given the role of dopamine within cortico-striatal circuits in both rhythm-based predictive timing and music-derived pleasure, we investigated the link between dopamine and PLUMM by comparing the urge to move and pleasure in PD along with healthy older and younger controls. We hypothesized that the inverted U-shaped relation between rhythmic complexity, urge to move and pleasure would differ in PD compared with controls but

that the relationship between harmonic complexity and PLUMM would be unaffected. Aging has also been linked to changes in neural and behavioural correlates of rhythm (Al Jaja et al., 2020; Henry et al., 2017; Krampe et al., 2001, 2002, 2005; Sauvé et al., 2019; Turgeon & Wing, 2012) and reward processing (Chowdhury et al., 2013; Eppinger et al., 2013; Schott et al., 2007; Vink et al., 2015). Therefore, the inclusion of both healthy older and younger controls also allowed us to examine any age-related changes in overall PLUMM response.

2 | MATERIAL AND METHODS

2.1 | Participants

The study was conducted at the Center for Music in the Brain, Aarhus University, Denmark. A total of 112 participants performed the experiment. Data from 101 of these participants were analysed after 11 PD were excluded from the analysis due to incomplete data. Table 1 shows a summary of demographic data.

PD participants (age 46–80) were diagnosed with idiopathic PD, in early- to mid-stages based on the Hoehn and Yahr scale (Di Biase et al., 2017; Hoehn & Yahr, 1967). They were diagnosed by a certified neurologist according to the Movement Disorder Society (MDS) diagnostic criteria (Postuma et al., 2015). PD whose symptoms exceeded stage III on the Hoehn and Yahr

scale (Hoehn & Yahr, 1967) were excluded from the analysis to avoid confounding effects of cognitive impairments associated with late-stage PD. Additionally, PD were excluded if they were in treatment with DBS or if they suffered from severe walking and balance problems, had important uncontrolled metabolic problems (e.g., diabetes and hypertension), hearing loss or any other neurological or psychiatric disorders. PD were instructed to maintain their regular medication regimen which included levodopa for all PD except for three who were only taking a dopamine agonist. Dopamine agonist medication is often taken in combination with levodopa and the combination can lead to disruptions in motivation and reward processes (Cools & D'Esposito, 2011). To test for these effects, the PD were divided into two groups, those who were taking dopamine agonist medications (PD_{DA}; $n = 23$) and those who were not (PD_{nonDA}; $n = 24$). There were two groups of healthy controls: age-matched older adults (HC_{old}; age 47–77; $n = 27$) and younger adults (HC_{young}; age 21–33; $n = 27$). Both PD and HC were excluded if they had important uncontrolled metabolic problems (e.g., diabetes and hypertension), hearing loss or any other neurological or psychiatric disorders. The metabolic syndrome has been linked to cognitive dysfunction and brain abnormalities, with insulin resistance-induced impairment in cerebrovascular reactivity being a key mechanism (Yates et al., 2012). All participants in the HC_{old} group and in both PD groups reported their nationality as Danish.

TABLE 1 Demographics and behavioural questionnaires.

	HC _{young}	HC _{old}	PD _{DA}	PD _{nonDA}
Participants (n)	27	26	23	24
Gender (females)	14	21	4	10
Age (years)	24.9 ± 3.2	65.7 ± 8.2	68.2 ± 4.8	66.8 ± 9
Education (years)	18.2 ± 2.1***	14.7 ± 2.3	14.9 ± 2.9	14.9 ± 3.5
Musical training (years)	4.5 ± 6.4**	2.6 ± 5.9	.3 ± .9	2 ± 5.4
Music time (h/week)	22 ± 14	19 ± 15	18 ± 18	16 ± 15
MDI	3.1 ± 3**	5 ± 6*	8.4 ± 4.6	8.1 ± 5.7
MoCA	29 ± 1.1	29 ± 1.2	28 ± 1.4	28 ± 1.5
PD diagnosis (years)	-	-	6.6 ± 4.7	5.6 ± 4.9
MDS-UPDRS-III	-	-	29.7 ± 11	25.2 ± 12.3
Hoehn and Yahr ^a	-	-	2 (2–2)	2 (1–2)
MET: Melody	36.7 ± 5.8	34 ± 5.6	31.5 ± 4	32.4 ± 5.4
MET: Rhythm	38 ± 5.5	34 ± 5.7	32.7 ± 3.9	32.3 ± 3.9

Abbreviations: MDI, major depression inventory; MDS-UPDRS-III, motor part of the Movement Disorder Society Unified Parkinson's Disease Rating Scale; MET, Musical Ear Test; MoCA, Montreal cognitive assessment.

^aMedian (IQR).

*Significantly different than PD_{DA}.

**Significantly different than both PD groups.

***Significantly different than all other groups.

Nineteen participants in the HC_{young} group reported their nationality as Danish while the remaining participants reported their country of nationality as Poland, German, Egypt, the Philippines, the Netherlands, the United Kingdom, Turkey and Brazil (all were international exchange students at Aarhus University).

All participants gave their consent verbally and in written form before the experiment. The study was conducted in accordance with the Declaration of Helsinki and approved by the Internal Review Board at the DNC (*Dansk Neuroforskningssenter*), Aarhus University Hospital, Aarhus, Denmark. PD received no compensation for participating in the study, but the costs of transportation were covered. Healthy controls received 200 DKK, in compensation for their participation in the 2-h experiment.

2.2 | Behavioural questionnaires and Musical Ear Test (MET)

Participants filled out an initial background questionnaire which included variables such as age, gender, years of education, years of musical training, preferred musical genre, and number of hours spent per week interacting with music. Such musical interaction included active music listening or playing, or passively using music as background. Specifically for PD, we asked for a list of current medication prescribed by their neurologist. All participants filled out the Major Depression Inventory (MDI) to exclude depression (Olsen et al., 2003) and the Montreal Cognitive Assessment (MoCA) to exclude dementia (Nasreddine et al., 2005). PD filled out the MDS-Unified Parkinson's Disease Rating Scale (MDS-UPDRS-III) to ensure PD diagnosis (Goetz et al., 2008). Authorization to use materials owned by the International Parkinson and MDS was granted to author V.P.-N., for research purposes only. Additionally, rhythm and melody discrimination ability was assessed in all participants via the MET (Wallentin et al., 2010). Table 1 shows a summary of questionnaire and MET data.

2.3 | PLUMM rating paradigm

Participants listened to and rated their experienced pleasure and urge to move for rhythmic stimuli (see Figure S1 for musical notation of an example stimuli). The stimuli were selected from previous studies conducted in our lab (Matthews et al., 2019, 2020, 2022; Pando-Naude et al., 2023; Stupacher, Wrede, & Vuust, 2022; Stupacher, Matthews, et al., 2022) and varied in their degree of rhythmic complexity, which was

measured using the syncopation index (Fitch & Rosenfeld, 2007). The index quantifies the extent to which a pattern deviates from the standard 4/4 m. There were three levels of rhythmic complexity (low, medium and high) with three different rhythms in each level. The medium complexity rhythms consisted of the son and rumba claves along with an experimenter-created rhythm in a clave style. These claves are five-onset patterns that originate in Afro-Cuban music but are commonly used in many musical genres. The low complexity rhythms followed similar patterns but with nearly all onsets falling on strong beats, thus removing nearly all syncopations. The high complexity rhythms also followed similar patterns, but only the first of the five onsets fell on a strong beat while the remaining four onsets were shifted to off-beat positions. In this way, all rhythmic patterns had the same number of onsets and followed a similar general pattern, differing only in terms of syncopation.

The rhythmic patterns were made up of piano chords (six notes spanning four octaves in D major) with a single chord repeating to generate the rhythm. These chords varied across three levels of harmonic complexity, with three different chords per level. Harmonic complexity corresponded to the degree to which the chords deviated from the D major key and was quantified as roughness using the MIR Toolbox in Matlab (Lartillot et al., 2007; Plomp & Levelt, 1965; Sethares, 2004). The rhythmic patterns and chords were combined into 54 stimuli, which were a subset of the 81 possible rhythm-chord combinations. Each stimulus also included an isochronous hi-hat (corresponding to eighth notes) which provided a metrical context. The rhythmic patterns were repeated four times, were presented at 96 bpm and lasted 10 s. See Figure S2 for a schematic of all rhythmic patterns used and Figure S3 for musical notation of all chords used.

At the start of the task, participants listened to two musical sequences that were not part of the experiment, to adjust the volume to a comfortable level. Participants listened to each sequence and rated pleasure/urge to move on a 5-point Likert scale (1 = *not at all/none*, 5 = *very much/a lot*), by using numbers on the computer keyboard to select their rating. The ratings were made according to the following questions: (a) To what extent does this rhythm make you want to move? and (b) How much pleasure do you experience listening to this rhythm? Following stimulus presentation, participants were given 5 s to make their rating. Ratings were made for all 54 stimuli in two consecutive sessions, urge to move then pleasure, to avoid influences of one rating on the other. Within each session, each musical sequence was presented in a fully randomized order. Participants listened via Beyerdynamic DT 770 Pro headphones.

2.4 | Statistical analysis

Only data from participants who completed all questionnaires and all trials of the PLUMM rating paradigm were analysed. Therefore, the analyses described below are implemented on a complete data set with no missing values ($n = 101$).

2.4.1 | Behavioural questionnaires and MET

Descriptive and inferential statistics of data were performed using R (version 4.1.1) in RStudio (R Core Team, 2020). Non-parametric (Kruskal–Wallis rank sum) tests were used to check for differences in demographic, cognitive, motor and depression questionnaire responses between groups. Post hoc analyses were conducted using the Dunn test of multiple comparisons corrected with the Bonferroni method.

The MET results showed a normal distribution; therefore, a one-way ANOVA was conducted to compare the groups. Post hoc analyses were conducted using the Tukey HSD multiple comparisons of means test with a 95% family-wise confidence level.

2.4.2 | Urge to move and pleasure ratings

The correlation between pleasure and urge to move ratings was high but not so high as to be considered colinear (Spearman $r(99) = .68$, 95% CI [.56, .77]); therefore, the rating types were analysed separately. Effects of group, rhythmic and harmonic complexity, and their interactions, on the urge to move and pleasure ratings, were investigated using two separate regression models, one for each type of rating. Data from all four groups (HC_{old}, HC_{young}, PD_{DA} and PD_{nonDA}) were included in these models. As there were between-group differences in years of education, years of formal music training and MDI score, these variables were included as covariates. To account for inter-individual variability, as well as variability between versions of stimuli (items) within the same complexity levels, the regression models were estimated using linear mixed effects models on the raw pleasure and urge to move ratings using lme4 (Bates et al., 2015) and afex (Singmann et al., 2013) packages. Random effects structures were determined following the recommendations of Bates et al. (2015) and Matuschek et al. (2017). For both models, this resulted in by-participant random intercepts and random slopes for both rhythmic and harmonic complexity, as well as by-item random intercepts. Correlations between by-participant random slopes and intercepts were included

for the analysis of urge to move ratings but not pleasure ratings. Diagnostic plots of residuals indicated no violations of assumptions in either model.

The two models were submitted to separate type III sum of squares analysis of variance (ANOVA) to test for main effects and interactions. Since the interactions were our primary interest, we report these first, and only mention significant main effects if relevant. If the ANOVA showed a significant three-way interaction, F tests were used to test for a rhythm by harmony interaction simultaneously in all groups. Second-order polynomial and pairwise contrasts were used to test quadratic and linear effects of rhythmic and harmonic complexity and to compare these effects between groups, using the emmeans package (Lenth et al., 2021). These follow-up comparisons focused on the effects between PD groups and HC_{old}, and between HC_{old} and HC_{young}. Effects in HC_{young} were not compared to those in the PD groups. For the ANOVAs, the degrees of freedom were estimated using the Satterthwaite method. For the follow-up contrasts, degrees of freedom were estimated using the Kenward–Roger method, and the multivariate t method was used to correct for multiple comparisons.

Finally, we tested correlations between the ratings and PD motor symptom severity in PD_{DA} and PD_{nonDA} and age and MET scores in PD_{DA}, PD_{nonDA} and HC_{old}. A summary measure capturing the pattern of ratings (Medium – (Low + High)/2), calculated on estimated ratings from the models were used in the correlations. False discovery rate was used to correct for multiple correlations.

3 | RESULTS

3.1 | Demographic and behavioural questionnaires

Demographic and questionnaire data are presented in Table 1. Kruskal–Wallis tests indicated that PD_{DA}, PD_{nonDA} and HC_{old} were not significantly different in terms of age ($\chi^2(2) = 1.21$, $p = .55$) and MoCA scores ($\chi^2(3) = 7.03$, $p = .07$). Dunn tests showed no significant differences between PD_{nonDA} and PD_{DA} in years since PD diagnosis ($p = .25$), UPDRS ($p = .23$) or Hoehn and Yahr scale ($p = .24$). The groups differed in terms of gender ratio, with the HC_{old} identifying predominantly as female and the PD groups as predominantly male. The groups also differed in years of education ($\chi^2(3) = 25.66$, $p < .001$), with HC_{young} having more education than all other groups (all $p < .001$). A significant effect of group on musical training ($\chi^2(3) = 12.99$, $p = .004$) was driven by HC_{young} having more years of musical training than both

PD groups (both $p < .05$) but not HC_{old}. There was no significant effect of group on hours spent interacting with music ($\chi^2(3) = 3.98$, $p = .26$). There was a significant effect of group on MDI scores ($\chi^2(3) = 25.05$, $p < .001$) driven by significantly lower scores in HC_{old} and HC_{young} compared with PD_{DA} (both $p < .05$) and HC_{young} compared with PD_{nonDA} ($p = .001$). As there were between-group differences in years of education, years of formal music training and MDI score, these variables were included as covariates in the regression models.

A significant effect of group on the rhythm version of the MET ($\chi^2(3) = 17.37$, $p < .001$) was driven by significantly better performance in HC_{young} compared with all other groups (all $p < .05$). A significant effect of group on the melody version of the MET ($\chi^2(3) = 12.48$, $p = .006$) was driven by better performance in HC_{young} compared with PD_{DA} ($p = .006$), with a near-significant difference between HC_{young} and PD_{nonDA} ($p = .05$) but not HC_{old} ($p = .65$).

3.2 | Urge to move ratings

3.2.1 | Rhythmic complexity by group interaction

The ANOVA showed a significant group by rhythmic complexity interaction (Table S1). Follow-up tests showed a significant quadratic effect in both HC_{young} and HC_{old}, with HC_{young} showing a stronger effect compared with HC_{old} ($b(97) = -1.216$, 95% CI $[-1.840, -.589]$; Table S2 and Figure 1a). The quadratic effect was significantly stronger in HC_{old} compared with both PD_{nonDA} ($b(97) = -1.066$, 95% CI $[-1.821, -.312]$) and PD_{DA} ($b(97) = -1.113$, 95% CI $[-1.876, -.349]$). PD_{nonDA} showed a significant negative linear effect which was driven by significantly higher ratings for low compared with high complexity rhythms ($b(85.3) = .468$, 95% CI $[.091, .845]$). PD_{DA} showed a relatively flat pattern of ratings as reflected in no significant linear or quadratic effects. The difference in linear effects between PD_{nonDA} and PD_{DA} was not significant.

3.2.2 | Harmonic complexity by group interaction

The ANOVA also showed a group by harmonic complexity interaction (Table S1). Follow-up tests showed that both HC_{old} and HC_{young} showed a significant quadratic effect and that this effect was stronger in HC_{old} compared with HC_{young} ($b(97) = 1.043$, 95% CI $[.593, 1.490]$; Table S3 and Figure 1b). HC_{old} also showed a stronger




FIGURE 1 The effects of (a) rhythmic and (b) harmonic complexity on urge to move ratings across groups. Values are estimated ratings from regression model. Centre line, median; box limits, upper and lower quartiles; whiskers, 1.5 \times interquartile range; white points, means; black points, outliers.

quadratic effect compared with both PD_{nonDA} ($b(97) = -1.261$, 95% CI $[-1.801, -.720]$) and PD_{DA} ($b(97) = -1.289$, 95% CI $[-1.836, -.742]$). As can be seen in Figure 1b, both PD_{nonDA} and PD_{DA} showed significant negative linear effects. This was reflected by significant low–medium (PD_{nonDA}: $b(89.2) = .403$, 95% CI $[.153, .653]$; PD_{DA}: $b(90.1) = .333$, 95% CI $[.078, .589]$) and medium–high (PD_{nonDA}: $b(89.0) = .593$, 95% CI $[.343, .842]$; PD_{DA}: $b(90.0) = .495$, 95% CI $[.241, .749]$) contrasts. There were no significant differences in these effects between PD_{nonDA} and PD_{DA}.

3.2.3 | Rhythmic complexity by harmonic complexity interaction

The ANOVA also showed a significant rhythmic by harmonic complexity interaction and a near-significant three-way interaction (Table S1). Given that the three-way interaction was near significant, and each group showed such different patterns of ratings, F tests on the rhythmic by harmonic complexity interaction were performed per group. These revealed a significant rhythmic complexity by harmonic complexity interaction in HC_{old} ($F(4, 4908.05) = 4.38$, $p = .002$) and a near-significant interaction for HC_{young} ($F(4, 4908.05) = 2.11$, $p = .077$) but not for PD_{DA} or PD_{nonDA} (both $p > .2$). Follow-up contrasts in HC_{old} showed that the quadratic effect of rhythmic complexity was significantly stronger, here reflecting a more peaked inverted U, for medium

complexity chords compared with both low ($b(4908) = -.611$, 95% CI $[-1.177, -.045]$) and high ($b(4908) = -.833$, 95% CI $[-1.400, -.267]$) complexity chords (see Figure S6). Unlike HC_{old} , HC_{young} showed no significant differences in the quadratic effect of rhythmic complexity between medium and low or medium and high complexity chords.

3.2.4 | Main effects

The ANOVA showed that the main effect of group was not significant ($F(3, 94.1) = .78$, $p = .508$). Therefore, although the groups showed different patterns in their ratings, there was no overall difference when averaging over rhythmic and harmonic complexity. There were no significant main effects of the control variables (years of education, years of musical training, and MDI).

3.2.5 | Correlations between Urge to Move ratings, UPDRS and MET scores

In PD, correlations between UPDRS and the contrast estimates for both rhythmic ($r(47) = -.24$, 95% CI $[-.52, .09]$) and harmonic complexity were not significant ($r(47) = -.24$, 95% CI $[-.52, .09]$). Correlations with MET scores were also not significant (all $p > .09$).

3.3 | Pleasure ratings

3.3.1 | Rhythmic complexity by group interaction

The ANOVA showed a significant group by rhythmic complexity interaction (Table S4). Follow-up tests showed a stronger quadratic effect in HC_{young} compared with HC_{old} ($b(98.5) = -1.128$, 95% CI $[-1.87, -.389]$) and no significant quadratic effects in HC_{old} , PD_{nonDA} and PD_{DA} (Table S5 and Figure 2a). PD_{nonDA} showed a significant negative linear effect ($b(119.6) = -.602$, 95% CI $[-1.077, -.127]$) driven by higher ratings in medium compared with high complexity rhythms ($b(117.5) = .565$, 95% CI $[.091, 1.039]$). The same contrast was also significant in HC_{old} ($b(113.3) = .512$, 95% CI $[.060, .964]$). There were no significant differences in the linear effect between groups. As can be seen in Figure 2a, both HC_{old} and PD_{nonDA} show similar ratings for low and medium complexity rhythms, then a drop-off for high complexity rhythms. Conversely, PD_{DA} 's ratings are relatively flat across all levels.




FIGURE 2 The effects of (a) rhythmic and (b) harmonic complexity on pleasure ratings across groups. Values are estimated ratings from regression model. Centre line, median; box limits, upper and lower quartiles; whiskers, $1.5 \times$ interquartile range; white points, means; black points, outliers.

3.3.2 | Harmonic complexity by group interaction

The ANOVA also showed a significant group by harmonic complexity interaction (Table S4). Follow-up tests indicated that HC_{old} showed greater linear ($b(103) = -.93$, 95% CI $[-1.320, -.541]$) and quadratic ($b(103) = -.44$, 95% CI $[-.850, -.030]$) effects compared with HC_{young} , for whom neither effects were significant. As can be seen in Figure 2b, HC_{old} , PD_{nonDA} and PD_{DA} all show a negative linear pattern as supported by a significant linear contrast in these groups (Table S6). These three groups showed significant differences between low and medium and medium and high complexity chords. However, HC_{old} showed a smaller difference between low and medium chords leading to the significant quadratic effect. Follow-up test showed a significant quadratic effect in HC_{old} and significant negative linear effects for both PD_{nonDA} and PD_{DA} . There were no significant differences between HC_{old} , PD_{nonDA} , and PD_{DA} groups in either the quadratic or linear contrasts.

3.3.3 | Rhythmic complexity by harmonic complexity interaction

The ANOVA also showed a significant three-way interaction (Table S4, Figure S7). Comparing HC_{young} and HC_{old} , F-tests showed that this interaction was significant in HC_{young} ($F(4, 4914.92) = 4.464$, $p = .001$) but not in

HC_{old} ($F(4, 4914.92) = .422, p = .793$). Follow-up contrasts showed that the quadratic effect of rhythm in HC_{young} was stronger for low compared to high complexity chords ($b(4915) = -.735, 95\% \text{ CI } [-1.300, -.170]$). There were no differences in the quadratic effect of rhythm between low versus medium and medium versus high complexity chords.

PD_{DA} and HC_{old} did not show a significant or near-significant interaction (both $p > .16$), while PD_{nonDA} showed a near-significant rhythmic complexity by harmonic complexity interaction ($F(4, 4914.92) = 2.245, p = .061$). There was no significant difference between medium and low complexity rhythms at any level of harmonic complexity. Together, these results suggest that the three-way interaction was driven by differences between HC_{young} and the other groups.

3.3.4 | Main effects

The ANOVA also showed a main effect of group (Table S4). HC_{old} showed higher ratings overall compared to HC_{young} ($b(94) = .453, 95\% \text{ CI } [.141, .765]$). There were no significant differences between PD_{nonDA} , PD_{DA} and HC_{old} . There were no significant main effects of the control variables (years of education, years of musical training, and MDI).

3.3.5 | Correlations between pleasure ratings, UPDRS and MET scores

In PD, there was a significant negative correlation between UPDRS and rhythmic complexity contrast estimates ($r(47) = -.37, 95\% \text{ CI } [-.62, -.05]$; Figure 3) but not harmonic complexity contrast estimates ($r(47) = -.03, 95\% \text{ CI } [-.35, .30]$). This indicates that as PD symptoms worsen, the quadratic effect of rhythmic complexity flattens out and/or becomes more linear. No other correlations were significant.

4 | DISCUSSION

Our results support the hypothesis that dopamine within cortico-striatal circuits play a crucial role in the PLUMM (Matthews et al., 2020; Stupacher, Matthews, et al., 2022). Both PD groups showed a flattening of the U-shaped relation between rhythmic complexity and the urge to move compared with healthy age-matched controls. In addition, the severity of motor symptoms in PD was associated with a flatter effect of rhythmic complexity on pleasure. The inverted U-shaped relation was




FIGURE 3 Correlation between severity of PD-related motor symptoms (UPDRS) and contrast estimates (medium – (low + high)/2) from the effect of rhythmic complexity on pleasure.

significantly weaker in healthy older compared with younger adults, suggesting that aging also influences the effect of rhythmic complexity on PLUMM. In addition, both PD groups showed a strong negative linear effect of harmonic complexity for both pleasure and the urge to move whereas healthy older adults showed a negative linear effect for pleasure and a quadratic effect for the urge to move. Importantly, PD did not differ from age-matched controls on a test of rhythm discrimination and did not show overall lower ratings for the urge to move and pleasure but rather a difference in the pattern of ratings. These results indicate that the flattening of the PLUMM response cannot easily be attributed to deficits in rhythm perception or a general inability to experience PLUMM. Finally, PD taking a dopamine agonist in addition to the standard levodopa regime generally showed a similar flattening effect to the PD_{nonDA} group. Taken together, these findings indicate that the chronic dysregulation of dopamine associated with PD flattens the typical U-shaped relationship between rhythmic complexity and PLUMM. We propose that this points to the crucial involvement of dopamine in the beat- and meter-based predictive processes thought to drive PLUMM.

4.1 | Rhythmic complexity

The flattened response to rhythmic complexity in PD supports a role of dopamine in the effects of rhythm on both the urge to move and pleasure. The motor cortico-striatal circuit is part of a core timing network, along with prefrontal and cerebellar regions (Coull et al., 2011; Matell &

Meck, 2004; Teghil et al., 2019) and is implicated in relative or beat-based timing in particular (Bengtsson et al., 2009; Breska & Ivry, 2018; Chen et al., 2008; Cheng et al., 2022; Cope et al., 2014; Grahn & Rowe, 2009, 2013; Kasdan et al., 2022; Kung et al., 2013; Merchant et al., 2015; Schubotz et al., 2000; Teki et al., 2012; Thaut et al., 2008). This is consistent with a role of the motor system in driving temporal predictions in the context of auditory perception and attention more generally (Arnal, 2012; Morillon et al., 2015; Nobre & van Ede, 2018; Rimmele et al., 2018; Schubotz, 2007). Involvement of the motor cortico-striatal circuit implies a role of nigrostriatal dopamine. However, studies on beat perception and perceived rhythmic complexity in PD show mixed results (Benoit et al., 2014; Biswas et al., 2016; Cameron et al., 2016; Geiser & Kaelin-Lang, 2011; Grahn & Brett, 2009; Hsu et al., 2022; Vikene et al., 2019c). Conversely, many of these same studies show deficits in rhythm discrimination in PD (Benoit et al., 2014; Biswas et al., 2016; Cameron et al., 2016; Geiser & Kaelin-Lang, 2011; Grahn & Brett, 2009; Hsu et al., 2022), which could suggest that the flatter pattern of ratings seen here may be due to reduced ability to distinguish the stimuli. However, HC_{old}, PD_{nonDA} and PD_{DA} significantly differed in their patterns of ratings but not in their rhythm discrimination ability, as measured by the MET. Further, there were no significant associations between MET scores and the patterns of ratings or between MET scores and the severity of PD-related motor symptoms (UPDRS). Conversely, higher UPDRS was associated with a flatter effect of rhythmic complexity on pleasure. Therefore, our results suggest that PD disrupts the beat and meter-based predictive processes that form the link between perceptual processing of rhythmic patterns and the affective responses to rhythmic music.

According to predictive processing accounts of PLUMM, strongly weighted prediction errors, in the form of syncopations, drive the inverted U-shaped relation between rhythmic complexity and both the urge to move and pleasure (Koelsch et al., 2019; Vuust et al., 2018, 2022; Vuust & Witek, 2014). In this context, strongly weighted syncopations drive the urge to move, either directly to reinforce the metrical model or indirectly via covert adjustments to that model. In terms of pleasure, by challenging our metric models, these syncopations potentiate model improvement (Stupacher, Matthews, et al., 2022), which is itself intrinsically rewarding (Oudeyer et al., 2016; Schmidhuber, 2010). The weight of prediction errors due to syncopations depends on both the rhythmic context (Lumaca et al., 2019) and the strength of the listener's metrical model (Vuust et al., 2005). According to a recent theoretical model (Cannon & Patel, 2020), nigrostriatal dopamine underlies

both the timing and certainty of beat-based predictions. Therefore, one interpretation of our results is that disrupted dopamine in PD results in a weaker metrical model and more uncertain predictions, reducing pleasure and the urge to move for moderately complex rhythms. Similarly, high complexity rhythms are less disruptive to an already weak model and are thus less aversive, leading to a flatter pattern of ratings overall. Support for the role of metric model strength in PLUMM comes from studies comparing musicians and non-musicians. Behavioural and neural evidence suggests that musicians show stronger and/or more refined metrical models compared with non-musicians (Drake et al., 2000; Geiser et al., 2010; Jongsma et al., 2004; Palmer & Krumhansl, 1990; Perna et al., 2018; Vuust et al., 2005, 2009) and that musicians have a sharper U-shaped pattern of PLUMM ratings and greater preference for higher levels of complexity (Matthews et al., 2019, 2022). In the current study, healthy aging was also linked to a flatter effect of rhythmic complexity, particularly for pleasure ratings. Like PD, healthy aging is associated with differences in the neural correlates of predictive rhythmic pattern perception (Henry et al., 2017), an effect that is also linked to dopamine (Al Jaja et al., 2020). Therefore, metrical model strength, supported by dopaminergic signalling, may provide a common mechanism underlying the effects of PD, aging and musicianship on the shape and location of the inverted U relation between PLUMM and rhythmic complexity.

Neuroimaging evidence suggests that neural oscillations in the beta range (13–30 Hz) may encode the metrical model within the motor cortico-striatal circuit and thus may provide a potential neural mechanism underlying the altered link between rhythm perception and affective responses in PD. Beta dysfunction within the BG is a hallmark of PD (Hammond et al., 2007) suggesting a functional link between dopamine and beta activity. One proposal is that, by encoding the certainty (or salience) of state, sensorimotor or value representations (e.g., rhythmic context; Friston et al., 2014), dopamine signals the need for motor preparation or resource allocation which are governed by beta modulations (Jenkinson & Brown, 2011). Crucially, cortico-striatal beta activity is linked to beat and meter representations (Bauer et al., 2015; Fujioka et al., 2012, 2015; Graber & Fujioka, 2019; Iversen et al., 2009) and is sensitive to musical training (Doelling & Poeppel, 2015; Fujioka & Ross, 2017). Further, PD show weaker beta entrainment in motor regions during rhythmic auditory and multimodal perceptual tasks (te Woerd et al., 2017, 2018), an effect that is reversed following treatment with levodopa and DBS (Gulberti et al., 2015). Finally, the degree of beta dysfunction in cortico-striatal circuits is correlated with

the severity of motor symptoms in PD (Kühn et al., 2009) which aligns with the association between severity of motor symptoms and the flattening of the effect of rhythmic complexity on pleasure ratings seen here. Together, these results point to a functional link between dopamine, beta activity, metrical model strength and PLUMM; however, further work is necessary to solidify and clarify this link.

An alternative interpretation for the flattened effect of rhythmic complexity on PLUMM ratings in PD is that, rather than weakening the metrical model, dopamine dysregulation disrupts the signalling of syncopation-related prediction errors, thus reducing the urge to move and pleasure thought to result from these prediction errors. However, (reward) prediction errors are associated with mesolimbic dopamine (Schultz et al., 1997) while PD primarily affects nigrostriatal dopamine, at least in early and mid-stages of the disease (Dauer & Przedborski, 2003). One challenge is that the effects on metrical model strength would also be expected to affect the prediction errors, as the certainty of the meter-based prediction is thought to directly relate to the weight of the prediction error. Future work should use behavioural (Palmer & Krumhansl, 1990; Perna et al., 2018) and/or computational methods to explicitly test whether metrical representations do in fact differ between PD and healthy controls matched in age and musical training.

Interestingly, both PD groups showed a very similar pattern of ratings, aside from a drop in the urge to move and pleasure in PD_{nonDA} for high complexity rhythms. This constitutes an internal replication and suggests that dopamine agonist medication does not have a strong effect on PLUMM over-and-above the general dopamine dysregulation and counteracting effects of levodopa present for all PD. To further delineate the role of dopamine, future work should compare PLUMM in both PD and healthy controls on and off levodopa medication.

4.2 | Harmonic complexity

Contrary to our hypothesis, and in contrast to the pattern of ratings in young healthy adults shown here and elsewhere (Matthews et al., 2019, 2020), PD showed greater pleasure and urge to move for low complexity chords, while healthy older adults showed the same pattern for pleasure and an inverted U for the urge to move. Recent work suggests that musicianship is associated with an altered pattern liking responses as a function of harmonic complexity (Matthews et al., 2019; Witek et al., 2023). Specifically, non-musicians preferred low complexity chords to medium, high or octave chords, while

musicians preferred medium complexity chords (Witek et al., 2023). Therefore, as with rhythmic complexity, PD, aging and (lack of) musical training seem to a similar influence on chord preferences. Further, previous studies using the same type of stimuli showed that medium harmonic complexity enhances the effect of moderately complex rhythms and that this may depend on musical training (Matthews et al., 2019; Matthews et al., 2020). In the current study, this effect was only seen for the urge to move in healthy older adults. Therefore, PD may also dampen the influence of harmonic complexity on rhythmic complexity. Younger healthy controls also showed an interaction between harmonic and rhythmic complexity, but only for pleasure ratings and this was not the expected pattern as low harmonic complexity enhanced the quadratic effect of rhythm. This group had more years of musical training compared with both PD groups. However, years of musical training was included as a control nuisance variable in all analyses, which, in addition to controlling for group differences, may have also reduced its influence on the effects of harmonic complexity.

PD and healthy older adults may also have an altered association between the complexity of a single chord and the emotion evoked or referenced by this chord. For example, PD show differences in the ability to recognize emotions expressed in music (Gothwal et al., 2022), which is relevant as chords of varying complexity are often used to convey different emotions. However, the specific nature of this effect is so far inconclusive, as some studies show deficits in recognizing happiness and peacefulness in PD (Lima et al., 2013) and others show deficits in recognizing fear and anger (van Tricht et al., 2010). Other factors including changes to listening habits and genre preferences that come with aging may also have influenced the pattern of ratings seen here; however, research into these effects is lacking. Therefore, further work is necessary to disentangle the relative contributions of musical training, age and PD to the effects of harmonic complexity and its interaction with rhythmic complexity.

5 | CONCLUSION

Overall, our results are consistent with models implicating cortico-striatal circuits and dopamine in PLUMM (Matthews et al., 2020; Stupacher, Matthews, et al., 2022) and provide insight into the function of dopamine within these circuits more generally. We propose that dopamine depletion within the motor cortico-striatal circuit disrupts beat- and meter-based predictive processes that are supported by beta modulations and are crucial for the

affective response to rhythmic music. Beyond music, this proposal aligns with and complements theoretical and empirical work suggesting that premotor beta modulations transmit top-down predictive signals that temporally structure auditory perception, for example, in the context of speech (Abbasi & Gross, 2020; Arnal, 2012; Arnal & Giraud, 2012; Morillon et al., 2015; Rimmele et al., 2018). Importantly, the current results better our understanding of dopamine's role in linking temporal predictions and reward processing which can be leveraged to improve treatments for PD and other dopamine-related disorders.

AUTHOR CONTRIBUTIONS

Victor Pando-Naude: Conceptualization; data curation; formal analysis; investigation; methodology; project administration; visualization; writing—original draft; writing—review and editing. **Tomas Edward Matthews:** Conceptualization; formal analysis; writing—original draft; writing—review and editing. **Andreas Højlund:** Conceptualization; methodology; supervision. **Sebastian Jakobsen:** Data curation; investigation. **Karen Østergaard:** Conceptualization; supervision. **Erik Johnsen:** Investigation; supervision. **Eduardo A. Garza-Villarreal:** Conceptualization; supervision. **Maria A. G. Witek:** Conceptualization; supervision. **Virginia Penhune:** Conceptualization; supervision; writing—review and editing. **Peter Vuust:** Conceptualization, supervision, writing—review and editing, supervision, funding acquisition.

ACKNOWLEDGEMENTS

The Center for Music in the Brain (MIB) is supported by the Danish National Research Foundation (Grant DNR 117). The authors would like to thank the students who assisted in the acquisition of data: Victor Poulsen, Anders Weile Larsen, Caroline Steen Sundberg and Agata Patyczek. Special thanks to all PD participants and healthy controls for their participation in this study.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest for this work.

PEER REVIEW

The peer review history for this article is available at <https://www.webofscience.com/api/gateway/wos/peer-review/10.1111/ejn.16128>.

DATA AVAILABILITY STATEMENT

The data supporting the findings of this study is freely available at the Open Science Framework (OSF) website: https://osf.io/fszqe/?view_only=cffadc85e4004c8789ca5cf604f172c7.

ORCID

Victor Pando-Naude  <https://orcid.org/0000-0003-1963-5068>

Tomas Edward Matthews  <https://orcid.org/0000-0003-4538-0898>

Andreas Højlund  <https://orcid.org/0000-0001-8460-8635>

Sebastian Jakobsen  <https://orcid.org/0009-0002-9033-0179>

Karen Østergaard  <https://orcid.org/0000-0003-4829-9761>

Virginia Penhune  <https://orcid.org/0000-0002-9475-5724>

Peter Vuust  <https://orcid.org/0000-0002-4908-735X>

REFERENCES

- Abbasi, O., & Gross, J. (2020). Beta-band oscillations play an essential role in motor–auditory interactions. *Human Brain Mapping, 41*(3), 656–665. <https://doi.org/10.1002/hbm.24830>
- Al Jaja, A., Grahn, J. A., Herrmann, B., & MacDonald, P. A. (2020). The effect of aging, Parkinson's disease, and exogenous dopamine on the neural response associated with auditory regularity processing. *Neurobiology of Aging, 89*, 71–82. <https://doi.org/10.1016/j.neurobiolaging.2020.01.002>
- Alexander, G., DeLong, M. R., & Strick, P. L. (1986). Parallel organization of functionally segregated circuits linking basal ganglia and cortex. *Annual Review of Neuroscience, 9*(1), 357–381. <https://doi.org/10.1146/annurev.neuro.9.1.357>
- Arnal, L. H. (2012). Predicting “when” using the motor system's beta-band oscillations. *Frontiers in Human Neuroscience, 6*, 225. <https://doi.org/10.3389/fnhum.2012.00225>
- Arnal, L. H., & Giraud, A.-L. (2012). Cortical oscillations and sensory predictions. *Trends in Cognitive Sciences, 16*(7), 390–398. <https://doi.org/10.1016/j.tics.2012.05.003>
- Bates, D., Mächler, M., Bolker, B. M., & Walker, S. C. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software, 67*(1), 1–48. <https://doi.org/10.18637/jss.v067.i01>
- Bauer, A. K. R., Kreutz, G., & Herrmann, C. S. (2015). Individual musical tempo preference correlates with EEG beta rhythm. *Psychophysiology, 52*(4), 600–604. <https://doi.org/10.1111/psyp.12375>
- Behroozmand, R., & Johari, K. (2019). Sensorimotor impairment of speech and hand movement timing processing in Parkinson's disease. *Journal of Motor Behavior, 51*(5), 561–571. <https://doi.org/10.1080/00222895.2018.1528204>
- Bella, S. D., Benoit, C.-E., Farrugia, N., Keller, P. E., Obrig, H., Mainka, S., & Kotz, S. A. (2017). Gait improvement via rhythmic stimulation in Parkinson's disease is linked to rhythmic skills. *Scientific Reports, 7*, 42005. <https://doi.org/10.1038/srep42005>
- Bellinger, D., Altenmüller, E., & Volkman, J. (2017). Perception of time in music in patients with Parkinson's disease—The processing of musical syntax compensates for rhythmic deficits. *Frontiers in Neuroscience, 11*(February), 68. <https://doi.org/10.3389/fnins.2017.00068>
- Bengtsson, S. L., Ullen, F., Henrik Ehrsson, H., Hashimoto, T., Kito, T., Naito, E., Forssberg, H., & Sadato, N. (2009). Listening to rhythms activates motor and premotor cortices. *Cortex, 45*(1), 62–71. <https://doi.org/10.1016/j.cortex.2008.07.002>

- Benoit, C.-E., Dalla Bella, S., Farrugia, N., Obrig, H., Mainka, S., & Kotz, S. A. (2014). Musically cued gait-training improves both perceptual and motor timing in Parkinson's disease. *Frontiers in Human Neuroscience*, 8(July), 494. <https://doi.org/10.3389/fnhum.2014.00494>
- Biswas, A., Hegde, S., Jhunjhunwala, K., & Pal, P. K. (2016). Two sides of the same coin: Impairment in perception of temporal components of rhythm and cognitive functions in Parkinson's disease. *Basal Ganglia*, 6, 63–70. <https://doi.org/10.1016/j.baga.2015.12.001>
- Blood, A. J., & Zatorre, R. J. (2001). Intensely pleasurable responses to music correlate with activity in brain regions implicated in reward and emotion. *Proceedings of the National Academy of Sciences of the United States of America*, 98(20), 11818–11823. <https://doi.org/10.1073/pnas.191355898>
- Blood, A. J., Zatorre, R. J., Bermudez, P., & Evans, A. C. (1999). Emotional responses to pleasant and unpleasant music correlate with activity in paralimbic brain regions. *Nature Neuroscience*, 2(4), 382–387. <https://doi.org/10.1038/7299>
- Breska, A., & Ivry, R. B. (2018). Double dissociation of single-interval and rhythmic temporal prediction in cerebellar degeneration and Parkinson's disease. *PNAS*, 115(48), 12283–12288. <https://doi.org/10.1073/pnas.1810596115>
- Câmara, G. S., & Danielsen, A. (2018). Groove. In A. Rehding & S. Rings (Eds.), *The Oxford handbook of critical concepts in music theory* (pp. 271–294). Oxford University Press. <https://doi.org/10.1093/oxfordhb/9780190454746.013.17>
- Cameron, D. J., Pickett, K. A., Earhart, G. M., & Grahn, J. A. (2016). The effect of dopaminergic medication on beat-based auditory timing in Parkinson's disease. *Frontiers in Neurology*, 7(19), 1–8. <https://doi.org/10.3389/fpls.2015.00830>
- Cannon, J. J., & Patel, A. D. (2020). How beat perception co-opts motor neurophysiology. *Trends in Cognitive Sciences*, 25(2), 137–150. <https://doi.org/10.1016/j.tics.2020.11.002>
- Chen, J. L., Penhune, V. B., & Zatorre, R. J. (2008). Listening to musical rhythms recruits motor regions of the brain. *Cerebral Cortex*, 18(12), 2844–2854. <https://doi.org/10.1093/cercor/bhn042>
- Cheng, T.-H. Z., Creel, S. C., & Iversen, J. R. (2022). How do you feel the rhythm: Dynamic motor-auditory interactions are involved in the imagination of hierarchical timing. *The Journal of Neuroscience*, 42(3), 500–512. <https://doi.org/10.1523/jneurosci.1121-21.2021>
- Cheung, V. K. M., Harrison, P. M. C., Meyer, L., Pearce, M. T., Haynes, J.-D., & Koelsch, S. (2019). Uncertainty and surprise jointly predict musical pleasure and amygdala, hippocampus, and auditory cortex activity. *Current Biology*, 29(23), 4084–4092.e4. <https://doi.org/10.1016/j.cub.2019.09.067>
- Chowdhury, R., Guitart-Masip, M., Lambert, C., Dayan, P., Huys, Q., Düzel, E., & Dolan, R. J. (2013). Dopamine restores reward prediction errors in old age. *Nature Neuroscience*, 16(5), 648–653. <https://doi.org/10.1038/nn.3364>
- Cools, R., & D'Esposito, M. (2011). Inverted-U-shaped dopamine actions on human working memory and cognitive control. *Biological Psychiatry*, 69(12), e113–e125. <https://doi.org/10.1016/j.biopsych.2011.03.028>
- Cope, T. E., Grube, M., Singh, B., Burn, D. J., & Griffiths, T. D. (2014). The basal ganglia in perceptual timing: Timing performance in Multiple System Atrophy and Huntington's disease. *Neuropsychologia*, 52, 73–81. <https://doi.org/10.1016/j.neuropsychologia.2013.09.039>
- Coull, J. T., Cheng, R.-K., & Meck, W. H. (2011). Neuroanatomical and neurochemical substrates of timing. *Neuropsychopharmacology*, 36(1), 3–25. <https://doi.org/10.1038/npp.2010.113>
- Coull, J. T., Hwang, H. J., Leyton, M., & Dagher, A. (2012). Dopamine precursor depletion impairs timing in healthy volunteers by attenuating activity in putamen and supplementary motor area. *Journal of Neuroscience*, 32(47), 16704–16715. <https://doi.org/10.1523/JNEUROSCI.1258-12.2012>
- Dauer, W., & Przedborski, S. (2003). Parkinson's disease: Mechanisms and models. *Neuron*, 39, 889–909. <https://doi.org/10.1017/CCOL9780521851282.008>
- Di Biase, L., Brittain, J. S., Shah, S. A., Pedrosa, D. J., Cagnan, H., Mathy, A., Chen, C. C., Martín-Rodríguez, J. F., Mir, P., Timmerman, L., Schwingenschuh, P., Bhatia, K., Di Lazzaro, V., & Brown, P. (2017). Tremor stability index: A new tool for differential diagnosis in tremor syndromes. *Brain*, 140(7), 1977–1986. <https://doi.org/10.1093/brain/awx104>
- Doelling, K. B., & Poeppel, D. (2015). Cortical entrainment to music and its modulation by expertise. *Proceedings of the National Academy of Sciences*, 112(45), E6233–E6242. <https://doi.org/10.1073/pnas.1508431112>
- Drake, C., Penel, A., & Bigand, E. (2000). Tapping in time with mechanically and expressively performed music. *Music Perception*, 18(1), 1–23. <https://doi.org/10.2307/40285899>
- Duman, D., Snape, N., Toiviainen, P., & Luck, G. (2022). Redefining groove. In *Proceedings of Conference on Interdisciplinary Musicology*. <https://doi.org/10.2218/cim22.1a17>
- Elsinger, C. L., Rao, S. M., Zimelman, J. L., Reynolds, N. C., Blindauer, K. A., & Hoffmann, R. G. (2003). Neural basis for impaired time reproduction in Parkinson's disease: An fMRI study. *Journal of the International Neuropsychological Society*, 9(7), 1088–1098. <https://doi.org/10.1017/S1355617703970123>
- Eppinger, B., Schuck, N. W., Nystrom, L. E., & Cohen, J. D. (2013). Reduced striatal responses to reward prediction errors in older compared with younger adults. *Journal of Neuroscience*, 33(24), 9905–9912. <https://doi.org/10.1523/JNEUROSCI.2942-12.2013>
- Ferreri, L., Mas-herrero, E., Zatorre, R. J., Ripollés, P., Gomez-andres, A., & Aicart, H. (2019). Dopamine modulates the reward experiences elicited by music. *PNAS*, 116(9), 3793–3798. <https://doi.org/10.1073/pnas.1811878116>
- Fitch, W. T., & Rosenfeld, A. J. (2007). Perception and production of syncopated rhythms. *Music Perception: An Interdisciplinary Journal*, 25(1), 43–58. <https://doi.org/10.1525/mp.2007.25.1.43>
- Friston, K. J., Schwartenbeck, P., FitzGerald, T., Moutoussis, M., Behrens, T., & Dolan, R. J. (2014). The anatomy of choice: Dopamine and decision-making. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 369(1655), 20130481. <https://doi.org/10.1098/rstb.2013.0481>
- Friston, K. J., Shiner, T., FitzGerald, T., Galea, J. M., Adams, R., Brown, H., Dolan, R. J., Moran, R., Stephan, K. E., & Bestmann, S. (2012). Dopamine, affordance and active inference. *PLoS Computational Biology*, 8(1), e1002327. <https://doi.org/10.1371/journal.pcbi.1002327>
- Fujioka, T., & Ross, B. (2017). Beta-band oscillations during passive listening to metronome sounds reflect improved timing representation after short-term musical training in healthy older

- adults. *European Journal of Neuroscience*, 46, 2339–2354. <https://doi.org/10.1111/ejn.13693>
- Fujioka, T., Ross, B., & Trainor, L. J. (2015). Beta-band oscillations represent auditory beat and its metrical hierarchy in perception and imagery. *Journal of Neuroscience*, 35(45), 15187–15198. <https://doi.org/10.1523/JNEUROSCI.2397-15.2015>
- Fujioka, T., Trainor, L. J., Large, E. W., & Ross, B. (2012). Internalized timing of isochronous sounds is represented in neuromagnetic β oscillations. *The Journal of Neuroscience*, 32(5), 1791–1802. <https://doi.org/10.1523/JNEUROSCI.4107-11.2012>
- Gebauer, L., Kringelbach, M. L., & Vuust, P. (2012). Ever-changing cycles of musical pleasure: The role of dopamine and anticipation. *Psychomusicology: Music, Mind, and Brain*, 22(2), 152–167. <https://doi.org/10.1037/a0031126>
- Geiser, E., & Kaelin-Lang, A. (2011). The function of dopaminergic neural signal transmission in auditory pulse perception: Evidence from dopaminergic treatment in Parkinson's patients. *Behavioural Brain Research*, 225(1), 270–275. <https://doi.org/10.1016/j.bbr.2011.07.019>
- Geiser, E., Sandmann, P., Jancke, L., & Meyer, M. (2010). Refinement of metre perception—Training increases hierarchical metre processing. *European Journal of Neuroscience*, 32, 1979–1985. <https://doi.org/10.1111/j.1460-9568.2010.07462.x>
- Gershman, S. J., & Uchida, N. (2019). Believing in dopamine. *Nature Reviews Neuroscience*, 20(11), 703–714. <https://doi.org/10.1038/s41583-019-0220-7>
- Goetz, C. G., Tilley, B. C., Shaftman, S. R., Stebbins, G. T., Fahn, S., Martinez-Martin, P., Poewe, W., Sampaio, C., Stern, M. B., Dodel, R., Dubois, B., Holloway, R., Jankovic, J., Kulisevsky, J., Lang, A. E., Rascol, O., Schrag, A., Teresi, J. A., Van Hilten, J. J., & Lapelle, N. (2008). Movement Disorder Society-Sponsored Revision of the Unified Parkinson's Disease Rating Scale (MDS-UPDRS): Scale presentation and clinimetric testing results. *Movement Disorders*, 23, 2129–2170. <https://doi.org/10.1002/mds.22340>
- Gold, B. P., Mas-Herrero, E., Dagher, A., & Zatorre, R. J. (2019). Toward a fuller understanding of reward prediction errors and their role in musical pleasure. *Proceedings of the National Academy of Sciences of the United States of America*, 116(42), 20815–20816. <https://doi.org/10.1073/pnas.1913835116>
- Gothwal, M., Arumugham, S., Yadav, R., Pal, P., & Hegde, S. (2022). Deficits in emotion perception and cognition in patients with Parkinson's disease: A systematic review. *Annals of Indian Academy of Neurology*, 25(3), 367–375. https://doi.org/10.4103/aian.aian_573_21
- Graber, E., & Fujioka, T. (2019). Endogenous expectations for sequence continuation after auditory beat accelerations and decelerations revealed by P3a and induced beta-band responses. *Neuroscience*, 413, 11–21. <https://doi.org/10.1016/j.neuroscience.2019.06.010>
- Grahn, J. A., & Brett, M. (2009). Impairment of beat-based rhythm discrimination in Parkinson's disease. *Cortex*, 45(1), 54–61. <https://doi.org/10.1016/j.cortex.2008.01.005>
- Grahn, J. A., & Rowe, J. B. (2009). Feeling the beat: Premotor and striatal interactions in musicians and nonmusicians during beat perception. *The Journal of Neuroscience*, 29(23), 7540–7548. <https://doi.org/10.1523/JNEUROSCI.2018-08.2009>
- Grahn, J. A., & Rowe, J. B. (2013). Finding and feeling the musical beat: Striatal dissociations between detection and prediction of regularity. *Cerebral Cortex*, 23(4), 913–921. <https://doi.org/10.1093/cercor/bhs083>
- Gulberti, A., Moll, C. K. E., Hamel, W., Buhmann, C., Koeppen, J. A., Boelmans, K., Zittel, S., Gerloff, C., Westphal, M., Schneider, T. R., & Engel, A. K. (2015). Predictive timing functions of cortical beta oscillations are impaired in Parkinson's disease and influenced by L-DOPA and deep brain stimulation of the subthalamic nucleus. *NeuroImage: Clinical*, 9, 436–449. <https://doi.org/10.1016/j.nicl.2015.09.013>
- Hammond, C., Bergman, H., & Brown, P. (2007). Pathological synchronization in Parkinson's disease: Networks, models and treatments. *Trends in Neurosciences*, 30(7), 357–364. <https://doi.org/10.1016/j.tins.2007.05.004>
- Henry, M. J., Herrmann, B., Kunke, D., & Obleser, J. (2017). Aging affects the balance of neural entrainment and top-down neural modulation in the listening brain. *Nature Communications*, 8(May), 15801. <https://doi.org/10.1038/ncomms15801>
- Hoehn, M. M., & Yahr, M. D. (1967). Parkinsonism: Onset, progression, and mortality. *Neurology*, 17(5), 427–442. <https://doi.org/10.1212/wnl.50.2.318>
- Hollerman, J. R., & Schultz, W. (1998). Dopamine neurons report an error in the temporal prediction of reward during learning. *Nature Neuroscience*, 1(4), 304–309. <https://doi.org/10.1038/1124>
- Hsu, P., Ready, E. A., & Grahn, J. A. (2022). The effects of Parkinson's disease, music training, and dance training on beat perception and production abilities. *PLoS ONE*, 17(3), e0264587. <https://doi.org/10.1371/journal.pone.0264587>
- Huron, D. (2006). *Sweet anticipation: Music and the psychology of expectation*. MIT Press. <https://doi.org/10.1525/mp.2007.24.5.511>
- Iversen, J. R., Repp, B. H., & Patel, A. D. (2009). Top-down control of rhythm perception modulates early auditory responses. *Annals of the New York Academy of Sciences*, 1169, 58–73. <https://doi.org/10.1111/j.1749-6632.2009.04579.x>
- Jahanshahi, M., Jones, C. R. G., Zijlmans, J., Katzenschlager, R., Lee, L., Quinn, N., Frith, C. D., & Lees, A. J. (2010). Dopaminergic modulation of striato-frontal connectivity during motor timing in Parkinson's disease. *Brain*, 133(3), 727–745. <https://doi.org/10.1093/brain/awq012>
- Janata, P., Tomic, S. T., & Haberman, J. M. (2012). Sensorimotor coupling in music and the psychology of the groove. *Journal of Experimental Psychology: General*, 141(1), 54–75. <https://doi.org/10.1037/a0024208>
- Jenkinson, N., & Brown, P. (2011). New insights into the relationship between dopamine, beta oscillations and motor function. *Trends in Neurosciences*, 34(12), 611–618. <https://doi.org/10.1016/j.tins.2011.09.003>
- Jones, C. R. G., & Jahanshahi, M. (2014). Contributions of the basal ganglia to temporal processing: Evidence from Parkinson's disease. *Timing and Time Perception*, 2(1), 87–127. <https://doi.org/10.1163/22134468-00002009>
- Jones, C. R. G., Malone, T. J. L., Dirnberger, G., Edwards, M., & Jahanshahi, M. (2008). Basal ganglia, dopamine and temporal processing: Performance on three timing tasks on and off

- medication in Parkinson's disease. *Brain and Cognition*, 68(1), 30–41. <https://doi.org/10.1016/j.bandc.2008.02.121>
- Jongsma, M. L. A., Desain, P., & Honing, H. (2004). Rhythmic context influences the auditory evoked potentials of musicians and nonmusicians. *Biological Psychology*, 66(2), 129–152. <https://doi.org/10.1016/j.biopsycho.2003.10.002>
- Kasdan, A. V., Burgess, A. N., Pizzagalli, F., Scartozzi, A., Chern, A., Kotz, S. A., Wilson, S. M., & Gordon, R. L. (2022). Identifying a brain network for musical rhythm: A functional neuroimaging meta-analysis and systematic review. *Neuroscience & Biobehavioral Reviews*, 136, 104588. <https://doi.org/10.1016/j.neubiorev.2022.104588>
- Koelsch, S., Vuust, P., & Friston, K. (2019). Predictive processes and the peculiar case of music. *Trends in Cognitive Sciences*, 23(1), 63–77. <https://doi.org/10.1016/j.tics.2018.10.006>
- Krampe, R. T., Engbert, R., & Kliegl, R. (2001). Age-specific problems in rhythmic timing. *Psychology and Aging*, 16(1), 12–30. <https://doi.org/10.1037/0882-7974.16.1.12>
- Krampe, R. T., Engbert, R., & Kliegl, R. (2002). The effects of expertise and age on rhythm production: Adaptations to timing and sequencing constraints. *Brain and Cognition*, 48(1), 179–194. <https://doi.org/10.1006/brcg.2001.1312>
- Krampe, R. T., Kliegl, R., & Mayr, U. (2005). Timing, sequencing, and executive control in repetitive movement production. *Journal of Experimental Psychology: Human Perception and Performance*, 31(3), 379–397. <https://doi.org/10.1037/0096-1523.31.3.379>
- Kühn, A. A., Tsui, A., Aziz, T., Ray, N., Brücke, C., Kupsch, A., Schneider, G. H., & Brown, P. (2009). Pathological synchronisation in the subthalamic nucleus of patients with Parkinson's disease relates to both bradykinesia and rigidity. *Experimental Neurology*, 215(2), 380–387. <https://doi.org/10.1016/j.expneurol.2008.11.008>
- Kung, S.-J., Chen, J. L., Zatorre, R. J., & Penhune, V. B. (2013). Interacting cortical and basal ganglia networks underlying finding and tapping to the musical beat. *Journal of Cognitive Neuroscience*, 25(3), 401–420. https://doi.org/10.1162/jocn_a_00325
- Large, E. W., & Snyder, J. S. (2009). Pulse and meter as neural resonance. *Annals of the New York Academy of Sciences*, 1169, 46–57. <https://doi.org/10.1111/j.1749-6632.2009.04550.x>
- Lartillot, O., Toivainen, P., & Eerola, T. (2007). A Matlab toolbox for music information retrieval. In *Data analysis, machine learning and applications* (pp. 261–268). Springer. https://doi.org/10.1007/978-3-540-78246-9_31
- Lei, J., Conradi, N., Abel, C., Frisch, S., Brodski-Guerniero, A., Hildner, M., Kell, C. A., Kaiser, J., & Schmidt-Kassow, M. (2019). Cognitive effects of rhythmic auditory stimulation in Parkinson's disease: A P300 study. *Brain Research*, 1716, 70–79. <https://doi.org/10.1016/j.brainres.2018.05.016>
- Lenth, R. V., Buerkner, P., Herve, M., Love, J., Riebl, H., & Singmann, H. (2021). “emmeans”—R package.
- Lima, C. F., Garrett, C., & Castro, S. L. (2013). Not all sounds sound the same: Parkinson's disease affects differently emotion processing in music and in speech prosody. *Journal of Clinical and Experimental Neuropsychology*, 35(4), 373–392. <https://doi.org/10.1080/13803395.2013.776518>
- Lumaca, M., Haumann, N. T., Brattico, E., Grube, M., & Vuust, P. (2019). Weighting of neural prediction error by rhythmic complexity: A predictive coding account using mismatch negativity. *European Journal of Neuroscience*, 49, 1597–1609. <https://doi.org/10.1111/ejn.14329>
- Madison, G. (2006). Experiencing groove induced by music: Consistency and phenomenology. *Music Perception*, 24(2), 201–208. <https://doi.org/10.1525/mp.2006.24.2.201>
- Martínez-Molina, N., Mas-Herrero, E., Rodríguez-Fornells, A., Zatorre, R. J., & Marco-Pallarés, J. (2016). Neural correlates of specific musical anhedonia. *Proceedings of the National Academy of Sciences*, 113, E7337–E7345. <https://doi.org/10.1073/PNAS.1611211113>
- Martínez-Molina, N., Mas-Herrero, E., Rodríguez-Fornells, A., Zatorre, R. J., & Marco-Pallarés, J. (2019). White matter microstructure reflects individual differences in music reward sensitivity. *The Journal of Neuroscience*, 39(25), 5018–5027. <https://doi.org/10.1523/JNEUROSCI.2020-18.2019>
- Mas-Herrero, E., Dagher, A., & Zatorre, R. J. (2018). Modulating musical reward sensitivity up and down with transcranial magnetic stimulation. *Nature Human Behaviour*, 2, 27–32. <https://doi.org/10.1038/s41562-017-0241-z>
- Matell, M. S., & Meck, W. H. (2004). Cortico-striatal circuits and interval timing: Coincidence detection of oscillatory processes. *Cognitive Brain Research*, 21(2), 139–170. <https://doi.org/10.1016/j.cogbrainres.2004.06.012>
- Matthews, T. E., Witek, M. A. G., Heggli, O. A., Penhune, V. B., & Vuust, P. (2019). The sensation of groove is affected by the interaction of rhythmic and harmonic complexity. *PLoS ONE*, 14(1), e0204539. <https://doi.org/10.1371/journal.pone.0204539>
- Matthews, T. E., Witek, M. A. G., Lund, T., Vuust, P., & Penhune, V. B. (2020). The sensation of groove engages motor and reward networks. *NeuroImage*, 214, 1–12. <https://doi.org/10.1016/j.neuroimage.2020.116768>
- Matthews, T. E., Witek, M. A. G., Thibodeau, J. L. N., Vuust, P., & Penhune, V. B. (2022). Perceived motor synchrony with the beat is more strongly related to groove than measured synchrony. *Music Perception*, 39(5), 423–442. <https://doi.org/10.1525/mp.2022.39.5.423>
- Matuschek, H., Kliegl, R., Vasishth, S., Baayen, H., & Bates, D. (2017). Balancing Type I error and power in linear mixed models. *Journal of Memory and Language*, 94, 305–315. <https://doi.org/10.1016/j.jml.2017.01.001>
- Meck, W. H. (2006). Neuroanatomical localization of an internal clock: A functional link between mesolimbic, nigrostriatal, and mesocortical dopaminergic systems. *Brain Research*, 1109(1), 93–107. <https://doi.org/10.1016/j.brainres.2006.06.031>
- Merchant, H., Grahn, J., Trainor, L., Rohrmeier, M., Fitch, W. T., & Merchant, H. (2015). Finding the beat: A neural perspective across humans and non-human primates. *Philosophical Transactions of the Royal Society*, 370, 1–16. <https://doi.org/10.1098/rstb.2014.0093>
- Merchant, H., Luciana, M., Hooper, C., Majestic, S., & Tuite, P. (2008). Interval timing and Parkinson's disease: Heterogeneity in temporal performance. *Experimental Brain Research*, 184(2), 233–248. <https://doi.org/10.1007/s00221-007-1097-7>
- Meyer, L. B. (1956). *Emotion and meaning in music*. University of Chicago Press.
- Morillon, B., Hackett, T. A., Kajikawa, Y., & Schroeder, C. E. (2015). Predictive motor control of sensory dynamics in

- auditory active sensing. *Current Opinion in Neurobiology*, 31C, 230–238. <https://doi.org/10.1016/j.conb.2014.12.005>
- Morris, I. B., Vasudevan, E., Schedel, M., Weymouth, D., Loomis, J., Pinkhasov, T., & Muratori, L. M. (2019). Music to one's ears: Familiarity and music engagement in people with Parkinson's disease. *Frontiers in Neuroscience*, 13, 661. <https://doi.org/10.3389/fnins.2019.00661>
- Nasreddine, Z. S., Phillips, N. A., Bédirian, V., Charbonneau, S., Whitehead, V., Collin, I., Cummings, J. L., & Chertkow, H. (2005). The Montreal Cognitive Assessment, MoCA: A brief screening tool for mild cognitive impairment. *Journal of the American Geriatrics Society*, 53(4), 695–699. <https://doi.org/10.1111/j.1532-5415.2005.53221.x>
- Nobre, A. C., & van Ede, F. (2018). Anticipated moments: Temporal structure in attention. *Nature Reviews Neuroscience*, 19(1), 34–48. <https://doi.org/10.1038/nrn.2017.141>
- Nomoto, K., Schultz, W., Watanabe, T., & Sakagami, M. (2010). Temporally extended dopamine responses to perceptually demanding reward-predictive stimuli. *The Journal of Neuroscience*, 30(32), 10692–10702. <https://doi.org/10.1523/JNEUROSCI.4828-09.2010>
- O'Boyle, D. J., Freeman, J. S., & Cody, F. W. J. (1996). The accuracy and precision of timing of self-paced, repetitive movements in subjects with Parkinson's disease. *Brain*, 119(1), 51–70. <https://doi.org/10.1093/brain/119.1.51>
- Olsen, L. R., Jensen, D. V., Noerholm, V., Martiny, K., & Bech, P. (2003). The internal and external validity of the Major Depression Inventory in measuring severity of depressive states. *Psychological Medicine*, 33(2), 351–356. <https://doi.org/10.1017/S0033291702006724>
- Oudeyer, P. Y., Gottlieb, J., & Lopes, M. (2016). Intrinsic motivation, curiosity, and learning: Theory and applications in educational technologies. In *Progress in brain research* (1st ed., Vol. 229). Elsevier B.V. <https://doi.org/10.1016/bs.pbr.2016.05.005>
- Palmer, C., & Krumhansl, C. L. (1990). Mental representations for musical meter. *Journal of Experimental Psychology: Human Perception and Performance*, 16(4), 728–741. <https://doi.org/10.1037/0096-1523.16.4.728>
- Pando-Naude, V., Matthews, T. E., Højlund, A., Jakobsen, S., Østergaard, K., Johnsen, E., Garza-Villarreal, E. A., Witek, M. A. G., Penhune, V., & Vuust, P. (2023). Dopamine dysregulation in Parkinson's disease flattens the pleasurable urge to move to musical rhythms. *bioRxiv*. <https://doi.org/10.1101/2023.02.27.530174>
- Pando-Naude, V., Patyczek, A., Bonetti, L., & Vuust, P. (2021). An ALE meta-analytic review of top-down and bottom-up processing of music in the brain. *Scientific Reports*, 11(1), 20813. <https://doi.org/10.1038/s41598-021-00139-3>
- Pau, M., Corona, F., Pili, R., Casula, C., Sors, F., Agostini, T., Cossu, G., Guicciardi, M., & Murgia, M. (2016). Effects of physical rehabilitation integrated with rhythmic auditory stimulation on spatio-temporal and kinematic parameters of gait in Parkinson's disease. *Frontiers in Neurology*, 7(AUG), 126. <https://doi.org/10.3389/fneur.2016.00126>
- Perna, F., Pavani, F., Zampini, M., & Mazza, V. (2018). Behavioral dynamics of rhythm and meter perception: The effect of musical expertise in deviance detection. *Timing and Time Perception*, 6(1), 32–53. <https://doi.org/10.1163/22134468-00002100>
- Plomp, R., & Levelt, J. M. (1965). Tonal consonance and critical bandwidth. *The Journal of the Acoustical Society of America*, 38(4), 548–560. <https://doi.org/10.1063/1.3033202>
- Postuma, R. B., Berg, D., Stern, M., Poewe, W., Olanow, C. W., Oertel, W., Obeso, J., Marek, K., Litvan, I., Lang, A. E., Halliday, G., Goetz, C. G., Gasser, T., Dubois, B., Chan, P., Bloem, B. R., Adler, C. H., & Deuschl, G. (2015). MDS clinical diagnostic criteria for Parkinson's disease. *Movement Disorders*, 30(12), 1591–1601. <https://doi.org/10.1002/mds.26424>
- R Core Team. (2020). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing.
- Rimmele, J. M., Morillon, B., Poeppel, D., & Arnal, L. H. (2018). Proactive sensing of periodic and aperiodic auditory patterns. *Trends in Cognitive Sciences*, 22(10), 870–882. <https://doi.org/10.1016/j.tics.2018.08.003>
- Salimpoor, V. N., Benovoy, M., Larcher, K., Dagher, A., & Zatorre, R. J. (2011). Anatomically distinct dopamine release during anticipation and experience of peak emotion to music. *Nature Neuroscience*, 14(2), 257–262. <https://doi.org/10.1038/nn.2726>
- Salimpoor, V. N., van den Bosch, I., Kovacevic, N., McIntosh, A. R., Dagher, A., & Zatorre, R. J. (2013). Interactions between the nucleus accumbens and auditory cortices predict music reward value. *Science*, 340(6129), 216–219. <https://doi.org/10.1126/science.1231059>
- Salimpoor, V. N., Zald, D. H., Zatorre, R. J., Dagher, A., & McIntosh, A. R. (2015). Predictions and the brain: How musical sounds become rewarding. *Trends in Cognitive Sciences*, 19(2), 86–91. <https://doi.org/10.1016/j.tics.2014.12.001>
- Sauvé, S. A., Bolt, E. L. W., Fleming, D., & Zendel, B. R. (2019). The impact of aging on neurophysiological entrainment to a metronome. *Neuroreport*, 30(10), 730–734. <https://doi.org/10.1097/WNR.0000000000001267>
- Schmidhuber, J. (2010). Formal theory of creativity & intrinsic motivation (1990-2010). *IEEE Transactions on Autonomous Mental Development*, 2(3), 1–18. <https://doi.org/10.1109/TAMD.2010.2056368>
- Schott, B. H., Niehaus, L., Wittmann, B. C., Schütze, H., Seidenbecher, C. I., Heinze, H. J., & Düzel, E. (2007). Ageing and early-stage Parkinson's disease affect separable neural mechanisms of mesolimbic reward processing. *Brain*, 130(9), 2412–2424. <https://doi.org/10.1093/brain/awm147>
- Schubotz, R. I. (2007). Prediction of external events with our motor system: Towards a new framework. *Trends in Cognitive Sciences*, 11(5), 211–218. <https://doi.org/10.1016/j.tics.2007.02.006>
- Schubotz, R. I., Friederici, A. D., & von Cramon, D. Y. (2000). Time perception and motor timing: A common cortical and subcortical basis revealed by fMRI. *NeuroImage*, 11(1), 1–12. <https://doi.org/10.1006/nimg.1999.0514>
- Schultz, W., Dayan, P., & Montague, P. R. (1997). A neural substrate of prediction and reward. *Science*, 275(5306), 1593–1599. <https://doi.org/10.1126/science.275.5306.1593>
- Seger, C. A., Spiering, B. J., Sares, A. G., Quraini, S. I., Alpeter, C., David, J., & Thaut, M. H. (2013). Corticostriatal contributions to musical expectancy perception. *Journal of Cognitive Neuroscience*, 25(7), 1062–1077. https://doi.org/10.1162/jocn_a_00371
- Senn, O., Rose, D., Bechtold, T. A., Kilchenmann, L., Hoesl, F., Jerjen, R., Baldassarre, A., & Alessandri, E. (2019).

- Preliminaries to a psychological model of musical groove. *Frontiers in Psychology*, 10(June), 1228. <https://doi.org/10.3389/FPSYG.2019.01228>
- Sethares, W. A. (2004). *Tuning, timbre, spectrum, scale*. Springer Science and Business Media.
- Shany, O., Singer, N., Gold, B. P., Jacoby, N., Tarrasch, R., Hendler, T., & Granot, R. (2019). Surprise-related activation in the nucleus accumbens interacts with music-induced pleasantness. *Social Cognitive and Affective Neuroscience*, 14(4), 459–470. <https://doi.org/10.1093/scan/nsz019>
- Singmann, H., Bolker, B., Wetfall, J., & Aust, F. (2013). afex: Analysis of factorial experiments. R package.
- Sioros, G., Miron, M., Davies, M., Gouyon, F., & Madison, G. (2014). Syncopation creates the sensation of groove in synthesized music examples. *Frontiers in Psychology*, 5(SEP), 1–10. <https://doi.org/10.3389/fpsyg.2014.01036>
- Spiech, C., Sioros, G., Endestad, T., Danielsen, A., & Laeng, B. (2022). Pupil drift rate indexes groove ratings. *Scientific Reports*, 12, 11620. <https://doi.org/10.1038/s41598-022-15763-w>
- Stupacher, J., Matthews, T. E., Pando-Naude, V., Foster, O., Elst, V., & Vuust, P. (2022). The sweet spot between predictability and surprise: Musical groove in brain, body, and social interactions. *Frontiers in Psychology*, 13, 1–9. <https://doi.org/10.3389/fpsyg.2022.906190>
- Stupacher, J., Wrede, M., & Vuust, P. (2022). A brief and efficient stimulus set to create the inverted U-shaped relationship between rhythmic complexity and the sensation of groove. *PLoS ONE*, 17(5), e0266902. <https://doi.org/10.1371/journal.pone.0266902>
- te Woerd, E. S., Oostenfeld, R., de Lange, F. P., & Praamstra, P. (2017). Impaired auditory-to-motor entrainment in Parkinson's disease. *Journal of Neurophysiology*, 117, 1853–1864. <https://doi.org/10.1152/jn.00547.2016>
- te Woerd, E. S., Oostenfeld, R., de Lange, F. P., & Praamstra, P. (2018). Entrainment for attentional selection in Parkinson's disease. *Cortex*, 99, 166–178. <https://doi.org/10.1016/j.cortex.2017.11.011>
- Teghil, A., Boccia, M., Antonio, D., Di Vita, A., & de Lena, C. (2019). Neural substrates of internally-based and externally-cued timing: An activation likelihood estimation (ALE) meta-analysis of fMRI studies. *Neuroscience and Biobehavioral Reviews*, 96, 197–209. <https://doi.org/10.1016/j.neubiorev.2018.10.003>
- Teki, S., Grube, M., & Griffiths, T. D. (2012). A unified model of time perception accounts for duration-based and beat-based timing mechanisms. *Frontiers in Integrative Neuroscience*, 5-(January), 90. <https://doi.org/10.3389/fnint.2011.00090>
- Thaut, M. H., Demartin, M., & Sanes, J. N. (2008). Brain networks for integrative rhythm formation. *PLoS ONE*, 3(5), e2312. <https://doi.org/10.1371/journal.pone.0002312>
- Thaut, M. H., McIntosh, G. C., Rice, R. R., Miller, R. A., Rathbun, J., & Brault, J. M. (1996). Rhythmic auditory stimulation in gait training for Parkinson's disease patients. *Movement Disorders: Official Journal of the Movement Disorder Society*, 11(2), 193–200. <https://doi.org/10.1002/mds.870110213>
- Tomassini, A., Ruge, D., Galea, J. M., Penny, W., & Bestmann, S. (2016). The role of dopamine in temporal uncertainty. *Journal of Cognitive Neuroscience*, 28(1), 96–110. <https://doi.org/10.1162/jocn.1162/jocn>
- Turgeon, M., & Wing, A. M. (2012). Late onset of age-related difference in unpaced tapping with no age-related difference in phase-shift error detection and correction. *Psychology and Aging*, 27(4), 1152–1163. <https://doi.org/10.1037/a0029925>
- van Tricht, M. J., Smeding, H. M. M., Speelman, J. D., & Schmand, B. A. (2010). Impaired emotion recognition in music in Parkinson's disease. *Brain and Cognition*, 74(1), 58–65. <https://doi.org/10.1016/j.bandc.2010.06.005>
- Vikene, K., Skeie, G. O., & Specht, K. (2019a). Abnormal phasic activity in saliency network, motor areas, and basal ganglia in Parkinson's disease during rhythm perception. *Human Brain Mapping*, 40(3), 916–927. <https://doi.org/10.1002/hbm.24421>
- Vikene, K., Skeie, G. O., & Specht, K. (2019b). Compensatory task-specific hypersensitivity in bilateral planum temporale and right superior temporal gyrus during auditory rhythm and omission processing in Parkinson's disease. *Scientific Reports*, 9(1), 1–9. <https://doi.org/10.1038/s41598-019-48791-0>
- Vikene, K., Skeie, G. O., & Specht, K. (2019c). Subjective judgments of rhythmic complexity in Parkinson's disease: Higher baseline, preserved relative ability, and modulated by tempo. *PLoS ONE*, 14(9), e0221752. <https://doi.org/10.1371/journal.pone.0221752>
- Vink, M., Kleerekooper, I., van den Wildenberg, W. P. M., & Kahn, R. S. (2015). Impact of aging on frontostriatal reward processing. *Human Brain Mapping*, 36(6), 2305–2317. <https://doi.org/10.1002/hbm.22771>
- Vuust, P., Heggli, O. A., Friston, K. J., & Kringelbach, M. L. (2022). Music in the brain. *Nature Reviews Neuroscience*, 23(5), 287–305. <https://doi.org/10.1038/s41583-022-00578-5>
- Vuust, P., Ostergaard, L., Pallesen, K. J., Bailey, C., & Roepstorff, A. (2009). Predictive coding of music—Brain responses to rhythmic incongruity. *Cortex*, 45(1), 80–92. <https://doi.org/10.1016/j.cortex.2008.05.014>
- Vuust, P., Pallesen, K. J., Bailey, C., van Zuijen, T. L., Gjedde, A., Roepstorff, A., & Østergaard, L. (2005). To musicians, the message is in the meter: Pre-attentive neuronal responses to incongruent rhythm are left-lateralized in musicians. *NeuroImage*, 24(2), 560–564. <https://doi.org/10.1016/j.neuroimage.2004.08.039>
- Vuust, P., Witek, M., Dietz, M., & Kringelbach, M. L. (2018). Now you hear it: A predictive coding model for understanding rhythmic incongruity. *Annals of the New York Academy of Sciences*, 1423(1), 19–29. <https://doi.org/10.1111/nyas.13622>
- Vuust, P., & Witek, M. A. G. (2014). Rhythmic complexity and predictive coding: A novel approach to modeling rhythm and meter perception in music. *Frontiers in Psychology*, 5(1111), 1–14. <https://doi.org/10.3389/fpsyg.2014.01111>
- Wallentin, M., Højlund Nielsen, A., Friis-Olivarius, M., Vuust, C., & Vuust, P. (2010). The Musical Ear Test, a new reliable test for measuring musical competence. *Learning and Individual Differences*, 20(3), 188–196. <https://doi.org/10.1016/j.lindif.2010.02.004>

- Witek, M. A. G., Clarke, E. F., Wallentin, M., Kringelbach, M. L., & Vuust, P. (2014). Syncopation, body-movement and pleasure in groove music. *PLoS ONE*, 9(4), e94446. <https://doi.org/10.1371/journal.pone.0094446>
- Witek, M. A. G., Matthews, T., Bodak, R., Blausz, M. W., Penhune, V., & Vuust, P. (2023). Musicians and non-musicians show different preference profiles for single chords of varying harmonic complexity. *PLoS ONE*, 18(2), e0281057. <https://doi.org/10.1371/journal.pone.0281057>
- Yates, K. F., Sweat, V., Yau, P. L., Turchiano, M. M., & Convit, A. (2012). Impact of metabolic syndrome on cognition and brain: A selected review of the literature. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 32(9), 2060–2067. <https://doi.org/10.1161/ATVBAHA.112.252759>
- Zald, D. H., & Zatorre, R. J. (2011). On music and reward. In J. Gottfried (Ed.), *The neurobiology of sensation and reward*. Taylor & Francis.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Pando-Naude, V., Matthews, T. E., Højlund, A., Jakobsen, S., Østergaard, K., Johnsen, E., Garza-Villarreal, E. A., Witek, M. A. G., Penhune, V., & Vuust, P. (2023). Dopamine dysregulation in Parkinson's disease flattens the pleasurable urge to move to musical rhythms. *European Journal of Neuroscience*, 1–18. <https://doi.org/10.1111/ejn.16128>