UNIVERSITY^{OF} BIRMINGHAM University of Birmingham Research at Birmingham

Elbow defect coverage using a capillary perforator flap arising from the vasa nervosum of the ulnar nerve

Javaid, R.h.; Chamaidi, E.; Hardwicke, Joseph; Kannan, R.y.

DOI: 10.1016/j.bjps.2014.11.021

License: Other (please specify with Rights Statement)

Document Version Peer reviewed version

Citation for published version (Harvard):

Javaid, RH, Chamaidi, E, Hardwicke, J & Kannan, RY 2015, 'Elbow defect coverage using a capillary perforator flap arising from the vasa nervosum of the ulnar nerve', *Journal of Plastic, Reconstructive & Aesthetic Surgery*, vol. 68, no. 4, pp. 592-594. https://doi.org/10.1016/j.bjps.2014.11.021

Link to publication on Research at Birmingham portal

Publisher Rights Statement:

NOTICE: this is the author's version of a work that was accepted for publication in Journal of Plastic, Reconstructive & Aesthetic Surgery. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Journal of Plastic, Reconstructive & Aesthetic Surgery, Vol 68, Issue 4, April 2015, DOI: 10.1016/j.bjps.2014.11.021

Eligibility for repository checked

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

•Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.

•User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?) •Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Accepted Manuscript

Elbow defect coverage using a capillary perforator flap arising from the vasa nervosum of the ulnar nerve

R.H. Javaid , E. Chamaidi , J.T. Hardwicke , R.Y. Kannan

PII: S1748-6815(14)00686-X

DOI: 10.1016/j.bjps.2014.11.021

Reference: PRAS 4452

To appear in: Journal of Plastic, Reconstructive & Aesthetic Surgery

Received Date: 22 July 2014

Accepted Date: 28 November 2014

Please cite this article as: Javaid RH, Chamaidi E, Hardwicke JT, Kannan RY, Elbow defect coverage using a capillary perforator flap arising from the vasa nervosum of the ulnar nerve, *British Journal of Plastic Surgery* (2015), doi: 10.1016/j.bjps.2014.11.021.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



ELBOW DEFECT COVERAGE USING A CAPILLARY PERFORATOR FLAP ARISING FROM THE VASA NERVOSUM OF THE ULNAR NERVE

R.H. Javaid¹, E. Chamaidi¹, J.T. Hardwicke^{1,2}, R.Y. Kannan¹†

¹ Department of Plastic Surgery,

New Queen Elizabeth Hospital,

Mindelsohn Way, Edgbaston,

Birmingham B15 2WB, UK.

² University of Birmingham, Edgbaston, Birmingham,

B15 2TT, UK.

†corresponding author:
R.Y. Kannan,
Department of Plastic Surgery,
New Queen Elizabeth Hospital,
Mindelsohn Way,Edgbaston,
Birmingham B15 2WB, UK.
ruben.kannan@gmail.com

Keywords:

Freestyle flap; perforator flap; neurocutaneous; elbow; soft tissue; reconstruction. Dear Sir,

Traditionally the pedicled latissimus dorsi muscle flap has been the workhorse flap for large elbow defects. Other options include the pedicled radial forearm flap and the reverse lateral arm flap, which can also be raised as a perforator-based radial collateral artery perforator (RCAP) flap. More recently, freestyle perforator flaps have broadened the options for soft tissue reconstruction¹. We describe a case of a free-style perforator flap for elbow reconstruction based on the capillary perforators arising from the vasa nervorum of the ulnar nerve.

An 89-year-old female patient underwent a wide excision of a biopsy proven nodular malignant melanoma, of 6.4mm Breslow thickness, on the radial aspect of the left elbow area (Figure 1). Pre- operatively an RCAP over the mid-arm was identified using a hand-held Doppler device. The melanoma was then excised with a 3cms lateral margin up to the deep fascia. The excision resulted in a 15 x 10 cms defect around the elbow in full extension. The flap based on the RCAP was marked on the posterolateral portion of distal upper arm, to be raised as a free-style propeller flap.

During dissection however, a cluster of small perforators, 5 cms proximal to the elbow joint arising from the ulnar nerve vasa nervosum, was identified and preserved (Figure 2). One of the RCAPs, which was in the proximal third of the flap, was also identified and preserved, but it was not possible to transpose or pivot the flap based on the RCAP. Given this, the RCAP was clamped for ten minutes to determine the perfusion of the flap from the capillary perforators of

the ulnar nerve vasa nervorum, which was optimal. Next, the RCAP was divided and the flap rotated through 180 degrees before being inset to cover the defect. The donor site was closed primarily and a bulky elbow dressing was applied. Elbow movement was allowed from day one with no restriction. The flap survived completely and all the wounds healed uneventfully with good functional and aesthetic outcomes with six weeks' post-op pictures shown in figure 3.

The concept of neurocutaneous flaps is not a new one. Masquelet et al and Nakajima et al pioneered neurocutaneous flaps in the 1990s by demonstrating that vessels accompanying the superficial nerves enable skin flaps to survive^{2,3}. Bertelli and Khoury studied the vascularization of the radial and ulnar nerves on the dorsal aspect of the hand and were able to plan and raise cutaneous island flaps vascularized by the vessels around and inside the nerves⁴. These flaps survived on the dictum that their respective nerves and accompanying vasa nervorum were needed to be fully incorporated in these flaps. The question asked here is whether it is necessary to include the entire nerve complex in these flaps but base them instead upon vasa nervorum capillary perforators.

Koshima described the concept of capillary perforator flaps in the thoracodorsal vascular system⁵ as a viable option in 2010. This alongside the previously described superficial peroneal nerve accessory artery perforator (SPNAA) perforator flap; a flap supplied by vasa nervorum without sacrificing the donor artery and the nerve⁶ supports our finding of the capillary perforators around the ulnar nerve and its use as a free-styled perforator flap. There has also been another case of a perforator flap being raised on the vasa nervorum of the anterior femoral

cutaneous nerve⁷. This emphasises the fact that perforators do arise from the vasa nervorum in the vicinity of nerves and this can be used to our advantage when raising perforator flaps.

Using this concept, we were able to reconstruct a large defect around the elbow with a free-style perforator flap based on the capillary perforators arising from the vasa nervorum of the ulnar nerve, with preservation of the nerve itself. It is important to note here that the perforators described in this case were not individually distinct, but existed in the form of a 'vascular leash', akin to the pedicle of the Foucher flap. This cluster of perforators was able to sustain the flap because their distal vector was in the axis of the flap.

In the future, we expect more such flaps being described as surgeons' become increasingly aware of the presence of capillary perforators, even around indispensable nerves or structures. It also calls on those of us performing free-style perforator flaps to keep an open mind about source vessel(s) and to expect the unexpected.

Conflict of interest: None

Funding: N/A

Ethical Approval: N/A

REFERENCES

- 1. Lecours C, Saint-Cyr M, Wong C et al. Freestyle pedicle perforator flaps: clinical results and vascular anatomy. Plast Reconstr Surg. 2010;126:1589-1603
- Masquelet AC, Romana MC, Wolf G. Skin island flaps supplied by the vascular axis of the sensitive superficial nerves: anatomic study and clinical experience in the leg. Plast Reconstr Surg. 1992;89:1115-1121.
- Nakajima H, Imanishi N, Fukuzumi S et al. Accompanying arteries of the lesser saphenous vein and sural nerve: anatomic study and its clinical applications. Plast Reconstr Surg. 1999;103:104-120.
- Bertelli JA1, Khoury Z. Neurocutaneous island flaps in the hand: anatomical basis and preliminary results. Br J Plast Surg. 1992; 45:586-90.
- Koshima, I, Narushima, M, Mihara, M, Ida, T, Gonda, K, Uchida, G, Nakagawa, M, New Thoracodorsal Artery Perforator (TAPcp) Flap with Capillary Perforators for Reconstruction of Upper Limb, J Plast Reconstr Aesthet Surg, 2010;64:140-145.
- Kim JS, Son DG, Choi TH et al. An Anatomic Study of the Superficial Peroneal Nerve Accessory Artery and Its Perforators, and Clinical Application of Superficial Peroneal Nerve Accessory Artery Perforator Flaps. Ann Plast Surg. 2008;60:710-716.
- Manaf Khatib MK; Matthew Ives MI; Matthew Griffiths MG. Never fear, a perforator will always appear: the use of a perforator flap based on the vasa nervosum of the anterior branch of the femoral cutaneous nerve. J Plast Reconstr Aesthet Surg 2013;66:1607-9

FIGURE LEGENDS

- **Figure 1** Pre-operative appearance of a 3 x 3cms nodular melanoma on the radial aspect of the left elbow.
- Figure 2 Intra-operative view of the cluster of capillary perforators, 5 cms proximal to the elbow joint, arising from the ulnar nerve vasa nervosum. As can be seen, there is no accompanying superior ulnar collateral artery with the ulnar nerve and there are multiple capillary perforators arising from the vasa nervosum of the ulnar nerve. The forceps here points to the ulnar nerve
- Figure 3Post-operative photograph of the perforator flap used for soft tissue reconstruction
of the elbow, six weeks after surgery.

6







CHR CHR

