

# Kolmogorov-Smirnov type testing for structural breaks

Hong, Yongmiao; Linton, Oliver; McCabe, Brendan; Sun, Jiajing; Wang, Shouyang

DOI:

[10.1016/j.jeconom.2023.105603](https://doi.org/10.1016/j.jeconom.2023.105603)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Hong, Y, Linton, O, McCabe, B, Sun, J & Wang, S 2024, 'Kolmogorov-Smirnov type testing for structural breaks: A new adjusted-range based self-normalization approach', *Journal of Econometrics*, vol. 238, no. 2, 105603. <https://doi.org/10.1016/j.jeconom.2023.105603>

[Link to publication on Research at Birmingham portal](#)

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.



# Kolmogorov–Smirnov type testing for structural breaks: A new adjusted-range based self-normalization approach

Yongmiao Hong <sup>a,b</sup>, Oliver Linton <sup>c</sup>, Brendan McCabe <sup>d</sup>, Jiajing Sun <sup>b,e,\*</sup>,  
Shouyang Wang <sup>a,b</sup>

<sup>a</sup> Center for Forecasting Science, Chinese Academy of Sciences, No. 55 Zhongguancun East Road, Haidian District, Beijing, 100190, China

<sup>b</sup> School of Economics and Management, and MOE Social Science Laboratory of Digital Economic Forecasts and Policy Simulation, University of Chinese Academy of Sciences, No. 3, Zhongguancun South First Street, Haidian District, Beijing, 100190, China

<sup>c</sup> Faculty of Economics, University of Cambridge, Austin Robinson Building, Sidgwick Ave, Cambridge CB3 9DD, United Kingdom

<sup>d</sup> Management School, University of Liverpool, Chatham St, Liverpool L69 7ZH, United Kingdom

<sup>e</sup> School of Mathematics, University of Birmingham, Watson Building, Edgbaston, Birmingham, B15 2TT, United Kingdom

## ARTICLE INFO

### JEL classification:

C12  
C19

### Keywords:

Change-point testing  
CUSUM process  
Parameter constancy  
Studentization

## ABSTRACT

A popular self-normalization (SN) approach in time series analysis uses the variance of a partial sum as a self-normalizer. This is known to be sensitive to irregularities such as persistent autocorrelation, heteroskedasticity, unit roots and outliers. We propose a novel SN approach based on the adjusted-range of a partial sum, which is robust to these aforementioned irregularities. We develop an adjusted-range based Kolmogorov–Smirnov type test for structural breaks for both univariate and multivariate time series, and consider testing parameter constancy in a time series regression setting. Our approach can rectify the well-known power decrease issue associated with existing self-normalized KS tests without having to use backward and forward summations as in Shao and Zhang (2010), and can alleviate the “better size but less power” phenomenon when the existing SN approaches (Shao, 2010; Zhang et al., 2011; Wang and Shao, 2022) are used. Moreover, our proposed tests can cater for more general alternatives. Monte Carlo simulations and empirical studies demonstrate the merits of our approach.

## 1. Introduction

We propose a new approach to conducting valid statistical inference in time series settings in the presence of serial dependence and heteroskedasticity. The usual approach in the literature is based on consistent estimation of the long-run variance (LRV). Substantial efforts have been devoted to providing and improving estimators of the LRV that are valid under weak conditions on the dependence structure. Perhaps one of the most well-known methods is the so-called heteroskedasticity and autocorrelation consistent (HAC) LRV estimator, espoused by White (1980), Newey and West (1987, 1994), Andrews (1991) and Andrews and Monahan (1992), among many others. However, Andrews (1991) and Den Haan and Levin (1997), among others, find that the finite sample performance of these HAC LRV estimators is rather poor. Müller (2007) suggests that these asymptotically consistent HAC LRV estimators, despite being theoretically appealing and empirically simple to apply, often lead to tests with poor sizes in finite samples when realistic amounts of dependence are present.

\* Corresponding author at: School of Economics and Management, and MOE Social Science Laboratory of Digital Economic Forecasts and Policy Simulation, University of Chinese Academy of Sciences, No. 3, Zhongguancun South First Street, Haidian District, Beijing, 100190, China.

E-mail addresses: [yh20@cornell.edu](mailto:yh20@cornell.edu) (Y. Hong), [obl20@cam.ac.uk](mailto:obl20@cam.ac.uk) (O. Linton), [Brendan.McCabe@liverpool.ac.uk](mailto:Brendan.McCabe@liverpool.ac.uk) (B. McCabe), [jiajing.sun@gmail.com](mailto:jiajing.sun@gmail.com) (J. Sun), [sywang@amss.ac.cn](mailto:sywang@amss.ac.cn) (S. Wang).

<https://doi.org/10.1016/j.jeconom.2023.105603>

Received 8 February 2022; Received in revised form 20 September 2023; Accepted 4 October 2023

Available online 9 November 2023

0304-4076/© 2023 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

An alternative approach is to use the so called fixed- $b$  asymptotics approach, proposed by Kiefer and Vogelsang (2005). The HAC LRV is often estimated using the nonparametric kernel smoothing method, which involves the choice of a kernel function and a smoothing parameter called bandwidth. Instead of assuming that the bandwidth parameter  $b$  tends to zero as the sample size  $n$  approaches infinity, under the fixed- $b$  asymptotics,  $b \in (0, 1]$  is set to be a fixed number, in which case the HAC LRV estimator is asymptotically unbiased but inconsistent — it possesses a limiting distribution rather than concentrating at its target.<sup>1</sup> This approach, sometimes also called self-normalization (SN), leads to a pivotal (but non-Gaussian) limiting distribution that can be tabulated and used in inference. Kiefer and Vogelsang (2005) outline two advantages of the fixed- $b$  approach; specifically, it enables a more accurate first order approximation to the asymptotic distribution and informative local power analysis for HAC robust tests. The fixed- $b$  asymptotics can enhance the size performance of various tests in finite samples (Kiefer et al., 2000; Kiefer and Vogelsang, 2002, 2005). However, there is a trade-off between size distortion and power loss; for a larger  $b$  the size distortion rectifies itself but the power loss increases (Kiefer and Vogelsang, 2005; Shao, 2015). Furthermore, the distribution of the fixed- $b$  based HAC test statistics depends on the nuisance parameter  $b$ ; there is, however, no general guidance on choosing  $b$ , and it must be prespecified in some ad hoc fashion. Moreover, Shao (2015) summarizes the Monte Carlo simulation results from Kiefer and Vogelsang (2005), Shao (2010) and Shao and Zhang (2010) and finds that when the time series is a unit-root or near unit-root process, the size of the fixed- $b$  based HAC test statistics deteriorates.

The contribution of our paper is threefold. First, we introduce a novel generally applicable adjusted-range based SN method for time series analysis. For concreteness, we demonstrate its use in testing for structural breaks in the mean of a class of approximately linear statistics and in the correlation coefficients and matrices of a multivariate time series, as well as in testing the constancy of parameters in a time series regression setting. Second, we develop adjusted-range based tests for structural changes, which can cater for more general alternatives and hence are potentially powerful against a wide range of alternatives, including smooth structural changes. For abrupt structural changes, the number of break points does not need to be specified a priori in an ad hoc manner. This contrasts with the G test proposed by Shao and Zhang (2010), which must be formulated with a pre-specified number of change points. Third, we illustrate, through extensive simulation studies, that the adjusted-range based SN can ameliorate the poor finite sample performance of HAC based tests (Müller, 2007), and can help rectify the non-monotonic power issue, without having to use forward and backward summations as employed by Shao and Zhang (2010), and enable us to circumvent the specification of a contrast process as in Zhang and Lavitas (2018). Moreover, we find, through simulation studies, that for statistical quantities that vary slowly over time, such as the medians and correlation coefficients, instead of displaying the “better size but less power” phenomenon, identified by Shao (2010) and Zhang et al. (2011), Shao and Zhang’s (2010) G test suffers from an over-size problem. This is because the G test statistic relies on the SN approach of Lobato (2001) and Shao (2010), and for statistical quantities (e.g., median, correlation) that do not change much over time, and sometimes even almost remain constant as the estimation horizon increases under the null hypothesis, the variance of the partial sum process can be quite small, which would lead to over-rejection of the null hypothesis.

**Literature Review.** The concept of SN originates from Student (1908), whose widely applied and celebrated  $t$  statistic and distribution first utilized this idea. Despite the fact that the sample variance based on a small number of observations is an inadequate estimator of a population variance, it is stochastically proportional to the population variance and can be used as a normalizer in order to construct a test statistic. The concept of SN has become an important principle in conducting statistical inference (Shao, 2015).

The existing self-normalized approach to inference for time series is first introduced in Shao (2010) as a generalization of an idea devised and developed by Kiefer et al. (2000) and Lobato (2001). Since its introduction by Lobato (2001) and Shao (2010), SN has been deployed in various aspects of statistical inference, such as confidence interval construction (Shao, 2010), testing for autocorrelation (Lobato, 2001; Shao, 2010; Boubacar-Mainassara and Sausseureau, 2018), testing for structural breaks (Shao and Zhang, 2010; Zhang et al., 2011; Zhang and Lavitas, 2018), and has been applied to various types of data, such as functional time series (Zhang et al., 2011; Dette et al., 2020), spatial data (Zhang et al., 2014), censored dependent data (Huang et al., 2015) and alternating regime index datasets (Kim and Shin, 2020). SN has also been applied across many academic fields of study, including: economics (Lobato, 2001; Shao, 2010), finance (Choi and Shin, 2021, 2020), ecology (Zhang et al., 2014), climate studies (Dette et al., 2020) and epidemiology (Jiang et al., 2023).

The SN approach proposed by Lobato (2001) and Shao (2010) is based on the variance of the partial sum of a time series process, which is sensitive to irregularities such as persistent autocorrelation, heteroskedasticity, near-unit roots and outliers. To alleviate the adverse effects of these irregularities, in this paper we propose the use of the adjusted range of a partial sum instead of the sample variance. Similar to the work of Lobato (2001) and Shao (2010), the sample range of a partial sum is asymptotically proportional to the square root of the LRV up to a stochastic factor, and since its distribution is nuisance parameter free, it can be used as an alternative self-normalizer. As is well-known, the range has some appealing robustness properties, such as the ability to deal with persistent autocorrelation and to accommodate irregularities such as outliers, high levels of skewness/kurtosis and unit-root behavior in volatilities. For example, Mandelbrot and Wallis (1969) show by Monte Carlo simulation studies that a range statistic can effectively detect long-range dependence in highly non-Gaussian time series with large skewness and/or kurtosis. Mandelbrot (1972, 1975) shows the appealing almost-sure convergence property of the range statistic for stochastic processes with infinite variances, a distinct advantage over the methods based on autocorrelations and variance ratios which may not be well-defined for

<sup>1</sup> In the fixed- $b$  asymptotics,  $b$  is often chosen to be a fixed number in  $(0, 1]$ . In contrast, for standard asymptotics, the bandwidth  $b$  vanishes to 0 as the sample size  $n$  increases, and for this reason, Kiefer and Vogelsang (2005) refer to the standard asymptotics as small- $b$  asymptotics.

infinite variance processes. Moreover, the range as a statistical quantity has also been widely applied in financial volatility estimation (Parkinson, 1980; Alizadeh et al., 2002; Chou et al., 2010).

A cornerstone of time series analysis is the structural stability of the data or model under consideration and failure to incorporate structural breaks will inevitably lead to unreliable inferences and forecasts. The importance of testing for structural breaks is also reflected in the vast related literature in various contexts; see (e.g.) Hansen (2001) and Aue and Horváth (2013) for literature reviews on testing for structural breaks. Depending on the quantities of interest, structural break tests can be conducted for the mean, variance or covariance structure and for general model stability; see (e.g.) Stock and Watson (1996), Bai (1997), Altissimo and Corradi (2003) and Harchaoui and Lévy-Leduc (2010) for references on testing for structural breaks in the mean. Substantial efforts have been devoted to testing breaks in time series regression models. Noticeably, the constancy of parameter tests are shown to be equivalent to tests of the mean of certain residuals, such as the estimated one-step ahead prediction errors. In particular, since Brown et al. (1975) introduce the cumulative sum (CUSUM) test based on recursive residuals, a large number of tests for parameter constancy based on CUSUM processes have been developed. For instance, Krämer et al. (1988) consider the CUSUM test for structural breaks when lagged dependent variables are incorporated in the linear regression model, and show that the CUSUM tests retain their asymptotic significance levels in dynamic regression models. Ploberger and Krämer (1992) consider the use of a CUSUM test when testing for parameter constancy in a linear regression model. See Andrews (1993), Bai and Perron (1998) and Qu and Perron (2007) for more studies on testing for breaks in time series regression models.

Testing for a structural break in variance is particularly important in economics and finance, where the stability or otherwise of volatility is a crucial issue; see, for instance, Inclan and Tiao (1994), Chen and Gupta (1997) and Smith (2008). There are also efforts that combine tests for structural breaks in mean and variance, such as Wang and Zivot (2000), Pitarakis (2004) and Jin et al. (2018). Aue et al. (2009) test for structural breaks in the covariance structure. Moreover, given that many multivariate volatility models impose restrictions on the correlation structure to deal with the “curse of dimensionality”, such as the constant correlation model (Bollerslev, 1990), the diagonal model (Bollerslev et al., 1988), the dynamic conditional correlation model (Engle, 2002), and the orthogonal or principal component generalized autoregressive conditional heteroskedasticity (GARCH) model (Alexander, 1998), testing for the constancy of correlation coefficients/matrices is also an important aspect in the literature; see (e.g.) Wied et al. (2012), Choi and Shin (2021, 2020) and Wied (2017).

The remainder of this paper is organized as follows. Section 2 considers testing for changes in the mean of a time series and introduces a novel adjusted-range based SN in both the univariate and multivariate cases. Section 3 generalizes Section 2 and covers testing for changes in a class of approximately linear statistics, which includes the marginal mean, the marginal variance, the autocorrelation function, quantiles and the spectrum as special cases. Section 4 proposes adjusted-range based Kolmogorov–Smirnov-type tests for constancy of parameters. Sections 5 and 6 cover the simulation studies and empirical applications, respectively. Finally, Section 7 concludes.

## 2. Testing structural breaks in mean

### 2.1. The univariate case

Considering a univariate time series  $\{X_t\}$ , we wish to test the null hypothesis

$$\mathbb{H}_0^{(1)} : E(X_1) = \dots = E(X_n) = \mu, \tag{1}$$

versus the alternative hypothesis

$$\mathbb{H}_1^{(1)} : \mathbb{H}_0^{(1)} \text{ is false.} \tag{2}$$

Define the CUSUM process as

$$T_n(k) = n^{-1/2} \sum_{t=1}^k (X_t - \bar{X}_n), \quad k = \lfloor sn \rfloor, \quad s \in [0, 1],$$

where  $\bar{X}_n = n^{-1} \sum_{t=1}^n X_t$  and  $\lfloor x \rfloor = \max \{z \leq x : z \in \mathbb{Z}\}$ . Under appropriate moment and weak dependence conditions (e.g., Assumption 2.1 in Phillips (1987)),  $T_n(k) \Rightarrow \sigma B(s)$ , where  $B(s)$  is the one-dimensional Brownian motion (Wiener process),  $\sigma^2 = \lim_{n \rightarrow \infty} n \text{var}(\bar{X}_n) = \sum_{j \in \mathbb{Z}} \gamma(j)$  is the LRV,  $\gamma(j) = \text{cov}(X_0, X_j)$ , and “ $\Rightarrow$ ” denotes weak convergence. Let  $\mathbb{B}(s) = B(s) - sB(1)$  be a Brownian Bridge. Under the null hypothesis  $\mathbb{H}_0^{(1)}$ , we have  $T_n(k) = T_n(\lfloor sn \rfloor) \Rightarrow \sigma \mathbb{B}(s)$ . The CUSUM statistic, also known as the KS statistic, is defined as

$$\text{KS} = \sup_{s \in [0,1]} |T_n(\lfloor sn \rfloor) / \sigma| \xrightarrow{d} \sup_{s \in [0,1]} |\mathbb{B}(s)|,$$

where “ $\xrightarrow{d}$ ” denotes convergence in distribution. In practice, a consistent estimator for  $\sigma$  is needed, for example the HAC LRV estimator  $\hat{\sigma}_n^2 = \sum_{k=-n+1}^{n-1} \hat{\gamma}(k) K(k/b)$ , where  $K(\cdot)$  and  $b$  denote a kernel function and a bandwidth that depends on the sample size  $n$ , respectively. We use  $\text{KS}^0$  to denote the HAC standardized KS test statistic. It is known that HAC robust tests tend to have poor sizes in small and finite samples with a moderate degree of autocorrelation (Müller, 2007).

**Table 1**  
Simulated critical values for KS type test statistics.

Level	10.0%	5.0%	2.5%	1.0%	0.5%	0.1%
KS	1.2220	1.3640	1.4762	1.6175	1.7119	1.9111
KS <sup>V</sup>	2.8857	3.0585	3.2029	3.3896	3.4765	3.6735
KS <sup>R</sup>	0.8684	0.9117	0.9391	0.9634	0.9732	0.9869

Note: The number of Monte Carlo simulations is 10,000 and the Brownian motion is approximated by the normalized sum of 200,000 i.i.d.  $N(0, 1)$  realizations.

Let  $\bar{X}_t = t^{-1} \sum_{i=1}^t X_i$ . The self-normalizer of Shao (2010) then takes the form

$$V_n^2 = n^{-2} \sum_{t=1}^n t^2 \left( \bar{X}_t - \bar{X}_n \right)^2.$$

Shao and Zhang (2010) point out that a naive application of the SN idea of Shao (2010) fails. A well-known undesirable feature of the self-normalized KS test statistic

$$KS^V = \max_{1 \leq k \leq n} \left| V_n^{-1} T_n(k) \right|$$

is that it has decreasing power, as the level shift increases; see Figure 1 in Shao and Zhang (2010). Shao and Zhang (2010) attribute this result to the increase in  $V_n$  with respect to the break size, and address such issues by introducing a so-called G test statistic, whose self-normalizer accommodates both the forward partial sum before the break point  $k^*$ , and the backward partial sum after  $k^*$ , and so is invariant with respect to the structural shift  $\Delta_n := E(X_{k^*+1}) - E(X_{k^*})$ . As a result, Shao and Zhang’s (2010) G test can detect the change by formulating two piecewise stationary partitions. However, the G test, in its simplest form, can cater for one change-point alternative only.<sup>2</sup>

Here, we propose an alternative approach to improve the performance of  $KS^V$  by using a new type of SN, which is based on the adjusted-range of the partial sum,

$$R_n = \max_{1 \leq k \leq n} T_n(k) - \min_{1 \leq k \leq n} T_n(k). \tag{3}$$

To derive the asymptotic distribution of  $R_n$ , we first assume that  $X_t = \mu + \varepsilon_t$ , where  $\mu$  is a fixed, yet unknown finite parameter, and  $\varepsilon_t$  is a zero-mean time series process. Following Phillips (1987), we impose the following regularity conditions.

**Assumption 1.** (i)  $E(\varepsilon_t) = 0$  for all  $t$ ; (ii)  $\sup_t E(|\varepsilon_t|^{2\beta}) < \infty$  for some  $\beta > 2$ ; (iii)  $0 < \sigma^2 = \lim_{n \rightarrow \infty} E \left[ n^{-1} \left( \sum_{t=1}^n \varepsilon_t \right)^2 \right] < \infty$ ; (iv)  $\{\varepsilon_t\}$  is an  $\alpha$ -mixing process with mixing coefficients  $\alpha_k$  that satisfy  $\sum_{k=1}^{\infty} \alpha_k^{1-2/\beta} < \infty$ .

Assumption 1 provides regularity conditions on moments and serial dependence of  $(\varepsilon_t)$ ; see Phillips (1987) for discussions on the rationale behind these assumptions.

Under Assumption 1, we have

$$R_n \xrightarrow{d} \sigma \left( \sup_{s \in [0,1]} \mathbb{B}(s) - \inf_{s \in [0,1]} \mathbb{B}(s) \right).$$

Therefore, under the null hypothesis  $\mathbb{H}_0^{(1)}$ ,

$$KS^R = \max_{1 \leq k \leq n} \left| R_n^{-1} T_n(k) \right| \xrightarrow{d} U, \tag{4}$$

where the positive scalar random variable  $U$  is defined as

$$U = \frac{\sup_{s \in [0,1]} |\mathbb{B}(s)|}{\sup_{s \in [0,1]} \mathbb{B}(s) - \inf_{s \in [0,1]} \mathbb{B}(s)}. \tag{5}$$

The asymptotic distributions of  $KS^V$  and  $KS^R$  can be obtained through simulations. The simulated critical values for KS,  $KS^V$  and  $KS^R$  are summarized in Table 1.

The  $KS^R$  test statistic is valid under a broad range of alternatives, including multiple breaks, smooth changes, or a mixture of them. However, to discuss its consistency, we focus on the one break point alternative (6), which is the main alternative hypothesis considered by Shao and Zhang (2010):

$$\mathbb{H}_1^{(1)*} : E(X_1) = \dots = E(X_{k^*}) \neq E(X_{k^*+1}) = \dots = E(X_n), \tag{6}$$

<sup>2</sup> Shao and Zhang (2010) suggest that the number of change points can be estimated through treating change point estimation and testing as model selection or adopting a sequential testing procedure; Zhang and Lavitas (2018) propose to circumvent the estimation for the number of break points and the application of a sequential testing procedure through the construction of the contrast processes and formulate a so-called T test statistic, which detects change points by recursive scanning. However, both the forward and backward summations in Shao and Zhang (2010) and the construction of contrast processes in Zhang and Lavitas (2018) are computationally expensive. Despite covering the multivariate cases in their theoretical exposition, the simulation studies of Shao and Zhang (2010) and Zhang and Lavitas (2018) are restricted to univariate cases only. Zhang and Lavitas (2018) even introduce a grid approximation scheme to alleviate the computational burden in the univariate case.

where  $k^* = \lfloor s_0 n \rfloor$  is the actual break location and  $s_0 \in (0, 1)$ . Define  $\delta := E(X_{k^*+1}) - E(X_{k^*})$  as the level of structural shift, and denote  $c_\alpha$  as the critical value of  $KS^R$  at significance level  $\alpha\%$ .

**Theorem 1.** Suppose that Assumption 1 holds. Then,

- (i) if  $\delta \neq 0$  is fixed, then  $\Pr(KS^R > c_\alpha) = 1$  as  $n \rightarrow \infty$ ;
- (ii) if  $\delta = n^{-1/2}\eta \neq 0$ , then  $\lim_{n \rightarrow \infty} \Pr(KS^R > c_\alpha) > 0$ , and  $\lim_{|\eta| \rightarrow \infty} \lim_{n \rightarrow \infty} \Pr(KS^R > c_\alpha) = 1$ .

Theorem 1 shows that the proposed test  $KS^R$  has nontrivial power against the class of one change-point alternatives that approach zero at the parametric root- $n$  rate. The proof for Theorem 1 is summarized in Appendix A.1. From the proof of Theorem 1, we can see that any structural break(s) that renders either  $T_n(k^*) = \infty$  provided that  $\min_{1 \leq k \leq n} T_n(k)/T_n(k^*) = o_p(1)$ , or  $T_n(k^*) = -\infty$  provided that  $\max_{1 \leq k \leq n} T_n(k)/T_n(k^*) = o_p(1)$ , when  $n \rightarrow \infty$ , should suffice for the consistency of  $KS^R$ . This rules out some oscillating breaks that push both  $\max_{1 \leq k \leq n} |T_n(k)|$  and  $\min_{1 \leq k \leq n} T_n(k)$  to  $\pm\infty$  as  $n \rightarrow \infty$ . See the supplementary material for more detailed discussions.

### 2.2. The multivariate case

The generalization of the  $KS^R$  test statistic to the multivariate case is not straightforward, because the adjusted-range (3) is always non-negative, while the off-diagonal components for a valid covariance estimator can be negative, albeit the whole matrix being at least positive semi-definite (PSD). In this section, we focus on testing structural breaks in the mean of a multivariate series.<sup>3</sup> Let  $X_t = (X_{1,t}, \dots, X_{m,t}) \in \mathbb{R}^m$  for  $t = 1, \dots, n$ , where  $m \geq 2$  is a fixed finite integer. We assume that under the null hypothesis  $X_t$  is weakly stationary with  $E(X_t) = \mu$  and  $E((X_t - \mu)(X_t - \mu)^\top) = \Sigma_X$ .

#### 2.2.1. Triangular structure

We first suppose that there is a unit lower triangular matrix  $L$  (with 1's along the principal diagonal) such that the process  $v_t = LX_t$  has components that are pairwise uncorrelated at all leads and lags. Specifically, we suppose that  $\text{cov}(v_{it}, v_{js}) = 0$  for all  $i \neq j$  and all  $t, s = 1, 2, \dots$ , whereas  $\text{cov}(v_{it}, v_{i,t+k}) = \gamma_i(k)$  may not be zero for all  $k$ , although each such autocovariance satisfies a summability condition. Let  $\Sigma_X = CDC^\top$  be the LDL representation of the covariance matrix of  $X_t$ , where  $D$  is a diagonal matrix and  $C$  is the unique unit lower triangular matrix. Similar to independent component analysis (ICA), we refer to  $C$  as the “mixing matrix” and  $C^{-1}$  as the “demixing matrix” (Gouriéroux et al., 2017). Under our assumed structure,  $L = C^{-1}$  and the autocovariance function of the time series  $X_t$  satisfies  $\Gamma_X(k) = E[(X_t - \mu)(X_{t+k} - \mu)^\top] = CD(k)C^\top$  for  $k = 0, \pm 1, \pm 2$ , where  $D(k) = \text{diag}\{\gamma_1(k), \dots, \gamma_m(k)\}$ . The  $m^2$  functions in  $\Gamma_X(k)$  are driven by  $m$  freely chosen functions  $\gamma_1(k), \dots, \gamma_m(k)$  and the  $m(m-1)/2$  free parameters in  $C$ . This is a reasonable assumption in the case where level of serial dependence is small or have a simple structure, and is one of the main structures exploited in the structural vector autoregression (SVAR) literature. In practice, we form the LDL decomposition on the sample covariance matrix of  $X_t$ , denoted as  $\hat{\Sigma}_X$ , such that  $\hat{\Sigma}_X = \hat{C}\hat{D}\hat{C}^\top$ , where  $\hat{C}$  is a unique lower triangular matrix with 1's along the principal diagonal and  $\hat{D}$  is a unique diagonal matrix with positive entries along the principal diagonal. The original series  $X_t$  is then mapped into  $\hat{u}_t = (\hat{u}_{1,t}, \dots, \hat{u}_{m,t})^\top$ , using the linear transformation  $\hat{u}_t = \hat{C}^{-1}X_t$ . Since the series  $u_t = C^{-1}X_t$  has a negligible level of cross correlation, the same should approximately be true of  $\hat{u}_t$  — the estimation errors for  $\hat{C}$  have negligible effect, because  $C$  can be consistently estimated.<sup>4</sup> We assume that all  $\{u_{l,t}\}$ , for  $l = 1, \dots, m$ , satisfy Assumption 1. This is a kind of partial prewhitening transformation because its goal is to eliminate cross dependence but not own serial dependence. For  $X_t$  that suffers from persistent serial correlation, we recommend applying the Vector Autoregression (VAR) and conducting an LDL decomposition on the sample variance of the estimated errors of the VAR model; see Section 2.2.2 for more discussion, which allows for a richer dynamic structure.

The construction of the adjusted range-based extended KS (EKS) test statistic takes the following steps. First, generate a new multivariate CUSUM process,

$$T_n^*(k) = n^{-1/2} \sum_{t=1}^k (\hat{u}_t - \bar{u}_n) \Rightarrow \Delta_u \mathbf{B}(s), \tag{7}$$

where  $\bar{u}_n = (\bar{u}_{1,n}, \dots, \bar{u}_{m,n}) = n^{-1} \sum_{t=1}^n \hat{u}_t \xrightarrow{p} 0$ , and  $\Delta_u$  is a matrix constant, such that the LRV of  $\hat{u}_t$  obeys  $\Sigma_u = \Delta_u \Delta_u^\top$ . Note that (7) holds under Assumption 1.

Second, denote the  $l$ th component of  $T_n^*(k)$  as  $T_n^{(l)*}(k) = n^{-1/2} \sum_{t=1}^k (\hat{u}_{l,t} - \bar{u}_{l,n})$ , for  $l = 1, \dots, m$ . Generate the  $m \times 1$  vector

$$\tilde{R}_n = \begin{pmatrix} \max_{1 \leq k \leq n} (T_n^{(1)*}(k)) - \min_{1 \leq k \leq n} (T_n^{(1)*}(k)) \\ \vdots \\ \max_{1 \leq k \leq n} (T_n^{(m)*}(k)) - \min_{1 \leq k \leq n} (T_n^{(m)*}(k)) \end{pmatrix}. \tag{8}$$

The adjusted-range based covariance estimator for  $\hat{u}_t$  is, therefore,  $\text{diag}\{\tilde{R}_n^2\}$ , which is a matrix with diagonal elements equal to  $\tilde{R}_n^2(l)$ , for  $l = 1, \dots, m$ . The adjusted-range based covariance estimator for  $\{X_t\}$  is  $\hat{C}^\top \text{diag}\{\tilde{R}_n^2\} \hat{C}$ .

<sup>3</sup> Shao and Zhang (2010) also consider testing the structural breaks for the multivariate case, under a general framework of so-called approximately linear statistics, which include the mean as a special case. We discuss testing structural breaks for approximately linear statistics in Section 3.

<sup>4</sup> This follows from the fact the sample covariance is a consistent estimator for the population covariance, and the LDL decomposition is unique for any given sample covariance.

**Table 2**  
Simulated critical values for  $\mathbb{EKS}^R(m)$  for  $m = 2, 3, \dots, 10$ .

$m \backslash$ Level	10.0%	5.0%	2.5%	1.0%	0.5%	0.1%
$m = 2$	1.0339	1.1425	1.2518	1.3706	1.4530	1.5732
$m = 3$	1.2954	1.4216	1.5281	1.6720	1.7793	1.9893
$m = 4$	1.5456	1.6818	1.8003	1.9645	2.0482	2.2967
$m = 5$	1.7692	1.9149	2.0505	2.1939	2.3109	2.7045
$m = 6$	1.9829	2.1544	2.2870	2.4742	2.5882	2.9517
$m = 7$	2.1970	2.3614	2.5163	2.6841	2.8356	3.1257
$m = 8$	2.3971	2.5733	2.7497	2.9263	3.0924	3.3959
$m = 9$	2.6046	2.7860	2.9433	3.1438	3.2838	3.6567
$m = 10$	2.8039	2.9928	3.1787	3.3765	3.5211	3.7879

Note: The number of Monte Carlo replications is 10,000 and the Brownian motion is approximated using 5000 i.i.d.  $N(0, 1)$  realizations.

Third, the EKS test statistic is defined as

$$\begin{aligned} \mathbb{EKS}^R(m) &= \max_{1 \leq k \leq n-1} T_n^*(k)^\top \left[ \text{diag} \left\{ \tilde{R}_n^2 \right\} \right]^{-1} T_n^*(k) \\ &= \max_{1 \leq k \leq n-1} T_n(k)^\top \left\{ \hat{C}^\top \text{diag} \left\{ \tilde{R}_n^2 \right\} \hat{C} \right\}^{-1} T_n(k), \end{aligned} \tag{9}$$

where  $m \geq 2$ . Further, define the non-negative scalar random variable

$$W_m = \sup_{s \in [0, 1]} \left\{ \mathbb{B}_m(s)^\top \left[ \text{diag} \left\{ \sup_{s \in [0, 1]} \mathbb{B}_m(s) - \inf_{s \in [0, 1]} \mathbb{B}_m(s) \right\} \right]^{-2} \mathbb{B}_m(s) \right\}, \tag{10}$$

where  $\mathbb{B}_m(s)$  is the  $m$ -dimensional Brownian bridge. If all  $\{\hat{u}_{l,t}\}$ , for  $l = 1, \dots, m$ , satisfy Assumption 1, we have

$$\mathbb{EKS}^R(m) \xrightarrow{d} W_m \text{ as } n \rightarrow \infty. \tag{11}$$

This three-step procedure effectively resolves the problem of extending the univariate adjusted-range based self-normalized KS test to the multivariate case. The proof for (11) is omitted, as it is a direct result of the invariance principle and the continuous mapping theorem (CMT).

The simulated asymptotic critical values for the  $\mathbb{EKS}^R(m)$  test statistics are tabulated in Table 2.

To prove the consistency of the  $\mathbb{EKS}^R(m)$  test, we again focus on the one change-point alternative (6). Define  $\Delta_n := E(X_{k^*+1}) - E(X_{k^*})$  as the level of structural shift, and denote  $C_\alpha$  as the critical value of  $\mathbb{EKS}^R(m)$  at significance level  $\alpha\%$ .

**Theorem 2.** Suppose that (7) holds. Then under the alternative hypothesis,

- (i) if  $s_0 \in (0, 1)$  and  $\Delta_n \neq \mathbf{0}$  is fixed, then  $\Pr(\mathbb{EKS}^R(m) > C_\alpha) = 1$  as  $n \rightarrow \infty$ ;
- (ii) if  $\Delta_n = n^{-1/2}\eta \neq \mathbf{0}$ ,  $\eta = (\eta^{(1)}, \dots, \eta^{(m)})^\top \neq \mathbf{0}$ , then  $\lim_{n \rightarrow \infty} \Pr(\mathbb{EKS}^R(m) > C_\alpha) > 0$  and  $\lim_{\|\eta\| \rightarrow \infty} \lim_{n \rightarrow \infty} \Pr(\mathbb{EKS}^R(m) > C_\alpha) = 1$ .

### 2.2.2. Residual triangular structure

If the  $X_t$  suffer from a high level of heteroskedasticity and/or autocorrelation, we could apply the VAR approach and assume a triangular structure for the error process. The VAR(p) model is

$$X_t = \Psi_1 X_{t-1} + \dots + \Psi_p X_{t-p} + e_t,$$

where  $p$  can be selected using model selection criteria, such as the AIC (Akaike Information Criterion). We conduct the LDL decomposition on the sample variance for the error  $\hat{e}_t$  in the VAR model, denoted by  $\hat{\Sigma}_e$ . Intuitively, this procedure is similar to the VAR prewhitening approach of Andrews and Monahan (1992), where the LRV of  $X_t$ , denoted by  $\Sigma_X$ , is estimated by first estimating the LRV of  $\hat{e}_t$  using the HAC approach, and then using the estimated parameters  $\hat{\Psi}_i$ ,  $i = 1, \dots, p$ , to conduct the reverse transformation, such that  $\hat{\Sigma}_X = \left( I_m - \sum_{i=1}^p \hat{\Psi}_i \right)^{-1} \hat{\Sigma}_e \left( I_m - \sum_{i=1}^p \hat{\Psi}_i \right)^{-1}$ , where  $I_m$  denotes an  $m \times m$  identity matrix.

Similarly, we first obtain  $\hat{u}_t$  through the LDL decomposition on the sample variance of the VAR errors  $\hat{\Sigma}_e = \hat{A}_e \hat{D}_e \hat{A}_e^\top$ . Then, following (7) and (8), the adjusted-range based covariance estimator for  $\hat{e}_t$  is  $\hat{A}_e \text{diag} \left\{ \tilde{R}_n^2 \right\} \hat{A}_e^\top$ , and that for  $X_t$  is

$$\tilde{\Sigma}_X = \left( I_m - \sum_{i=1}^p \hat{\Psi}_i \right)^{-1} \hat{A}_e \text{diag} \left\{ \tilde{R}_n^2 \right\} \hat{A}_e^\top \left( I_m - \sum_{i=1}^p \hat{\Psi}_i \right)^{-1}.$$

Thus, the adjusted-range based EKS test statistic becomes

$$\mathbb{EKS}^R(m) = \max_{1 \leq k \leq n-1} T_n(k)^\top \tilde{\Sigma}_X^{-1} T_n(k),$$

which converges in distribution to  $W_m$  defined in (10). This is equivalent to setting  $\hat{u}_t = \hat{A}_e \left( I_m - \sum_{i=1}^p \hat{\Psi}_i \right)^{-1} X_t$ ; or equivalently  $\hat{C} = \left( I_m - \sum_{i=1}^p \hat{\Psi}_i \right) \hat{A}_e^{-1}$ . The validity of this VAR approach relies on consistent estimation for  $\hat{\Psi}_i$  for  $i = 1, \dots, p$ ; see Hamilton (1994 pp. 298–299) for the asymptotic properties of the maximum likelihood (ML) estimator of a VAR model.

For other statistical quantities, we recommend using the LDL decomposition on the sample variance of  $X_t$ . For instance, for testing constancy of correlation coefficients/matrices, the VAR approach would remove the correlation among  $X_{1,t}, \dots, X_{m,t}$ , which might render the EKS test statistic evaluated on  $\hat{u}_{1,t}, \dots, \hat{u}_{m,t}$  unable to detect any structural breaks in the correlations among  $X_t$  in finite samples. See the supplementary materials for tests for constancy of correlation coefficients/matrices.

A similar rationale applies to the ICA. We can apply the fast ICA algorithm developed by Hyvarinen (1999) and ICA via distance covariance by Matteson and Tsay (2017) to the sample variances of both  $X_t$  and  $\hat{e}_t$ , respectively. However, we find that they lead to lower powers; in the extreme case, when testing constancy of correlation coefficients/matrices, both the sizes and powers are diminished in finite samples.<sup>5</sup> For many statistical quantities other than the mean, all we need is a linear/affine transformation that can capture and summarize (off-diagonal) correlatedness into the diagonal components. In contrast, ICA algorithms that deliver independent  $(\hat{u}_{1,t}, \dots, \hat{u}_{m,t})^\top$  reduce the powers of the EKS test statistics.

In situations involving high-dimensional series, one may utilize techniques such as the singular value decomposition (SVD) prewhitening of the ICA, or the Karhunen–Loève expansion followed by a truncation to reduce the number of parameters to be estimated. However, these approaches are beyond the scope of this paper and will be pursued in subsequent research.

### 3. Testing for structural breaks in approximately linear statistics

In this section, we adopt the framework of Shao and Zhang (2010) and consider a general quantity of interest, known as approximately linear statistics (Kunsch, 1989).<sup>6</sup>

Let  $Y_t = (X_t, \dots, X_{t+m-1})^\top$ ,  $t = 1, \dots, n$ , and denote  $F_t^m$  as the distribution of  $Y_t$ . Define

$$\theta_t = \mathbf{T}(F_t^m) \in \mathbb{R}^q, \quad t = 1, \dots, n, \tag{12}$$

as the quantity of interest, where  $\mathbf{T}$  is a functional that takes values in  $\mathbb{R}^q$ . Examples of  $\theta$  include, but are not limited to, the marginal mean of  $X_t$ , the marginal variance of  $X_t$ , the autocovariance function of  $X_t$  and the quantiles of the distribution of  $\mathbf{F}^1$  (Shao and Zhang, 2010).

Here, we are interested in testing the null hypothesis

$$\mathbb{H}_0^{(2)} : \theta_1 = \dots = \theta_n,$$

versus the alternative hypothesis

$$\mathbb{H}_1^{(2)} : \mathbb{H}_0^{(2)} \text{ is false.}$$

We replace the partial sum process, in the case of mean, by a sequence of recursive estimators of the quantity of interest, which are functionals of the distribution function  $F_t^m$ . Because  $F_t^m$  is unknown, these recursive estimators are obtained using the empirical distribution function. Let  $\rho_{1,k}$  be the empirical distribution based on  $\{Y_t\}_{t=1}^k$ , namely

$$\rho_{1,k} = k^{-1} \sum_{t=1}^k \delta_{Y_t}, \tag{13}$$

for  $1 \leq k \leq n$ , where  $\delta_Y$  is the probability measure which puts mass 1 at point  $Y$ ; see Definition 1 in Hampel et al. (1986, p.84).

The approximately linear statistic  $\hat{\theta}_{1,k}$  satisfies the following expansion (Shao and Zhang, 2010),

$$\hat{\theta}_{1,k} = \mathbf{T}(\rho_{1,k}) = \mathbf{T}(F^m) + k^{-1} \sum_{t=1}^k \mathbf{IF}(Y_t; F^m) + \mathbf{R}_{1,k}, \tag{14}$$

for  $1 \leq k \leq n$ , where  $\mathbf{IF}(Y; F^m)$  is the influence function of  $\mathbf{T}$  at  $F^m$ , such that

$$\mathbf{IF}(Y; F^m) = \lim_{\epsilon \downarrow 0} \frac{\mathbf{T}[(1 - \epsilon)F^m + \epsilon\delta_Y] - \mathbf{T}(F^m)}{\epsilon},$$

and  $\mathbf{R}_{1,k}$  is the remainder term of the expansion.

We define a process based on  $\hat{\theta}_{1,k}$ , i.e.,

$$T_n(k) = kn^{-1/2} \left( \hat{\theta}_{1,k} - \hat{\theta}_{1,n} \right), \tag{15}$$

where  $\hat{\theta}_{1,k}$  is estimated using the subsample  $\{Y_j\}_{j=1}^k$ .

<sup>5</sup> The simulation studies are available from the authors upon request.

<sup>6</sup> We also consider testing structural breaks in correlation coefficients and matrices, reflecting their importance in volatility modeling. Since correlations are not approximately linear statistics, adjusted-range based self-normalized tests for correlations are introduced in a slightly different asymptotic setting; see the supplementary material for detailed theoretical exposition.



Following Shao and Zhang (2010), let  $D[0, 1]$  be the space of functions on  $[0, 1]$ , which are càdlàg functions endowed with the Skorokhod topology (Billingsley, 1968); furthermore, we denote “ $\Rightarrow$ ” as weak convergence in  $D[0, 1]$ , hereafter. We impose the following condition.

**Assumption 2.**  $E\{\mathbf{IF}(Y_t; \mathbf{F}^m)\} = \mathbf{0}$  and  $n^{-1/2} \sum_{t=1}^{\lfloor sn \rfloor} \mathbf{IF}(Y_t; \mathbf{F}^m) \Rightarrow \Delta \mathbf{B}_q(s)$ , where  $\Delta$  is a  $q \times q$  lower triangular matrix with nonnegative diagonal entries and  $\mathbf{B}_q(\cdot)$  is a  $q$ -dimensional vector of independent Brownian motions. The long-run variance covariance matrix  $\Sigma(\mathbf{F}^m) = \Delta \Delta^\top = \sum_{k=-\infty}^{\infty} \text{cov}\{\mathbf{IF}(Y_0; \mathbf{F}^m), \mathbf{IF}(Y_k; \mathbf{F}^m)\}$  is positive definite.

Assumption 2 is referred to as Assumption 1 in Shao (2010), or Assumption 3.1 in Shao and Zhang (2010). Shao (2010) notes that it is not primitive, and cite Assumption 2.1 in Phillips (1987) as a primitive condition, which is identical to Assumption 1 imposed earlier.

To ensure that the remainder term for the expansion in (14) vanishes to zero asymptotically, we impose the following condition.

**Assumption 3.**  $\sup_{1 \leq k \leq n} |k \mathbf{R}_{1,k}| = o_p(n^{1/2})$ .

Assumption 3 is similar to Assumption 3.2 in Shao and Zhang (2010). The only difference is that since we do not need the backward and forward summations as in the G test of Shao and Zhang (2010), there is no need to assume that  $\sup_{1 \leq k \leq n} |k \mathbf{R}_{n-k+1,n}| = o_p(n^{1/2})$ . Under the expansion in (14), if Assumptions 2 and 3 are satisfied, we have  $\sqrt{n}(\hat{\theta}_{1,n} - \theta) \xrightarrow{d} N(\mathbf{0}, \Sigma(\mathbf{F}_m))$  as  $n \rightarrow \infty$ , where  $\theta = E(\hat{\theta}_{1,n})$ .

The subtleness for applying the EKS test to the present context is the existence of cross dependence in  $\hat{\theta}_{1,k}$ , which follows from the cross dependence in  $Y_t$ .<sup>7</sup> Following a procedure similar to that in Section 2.2, we work with  $\hat{u}_t = \hat{C}^{-1} Y_t$ , where  $\hat{C}$  is obtained through an LDL decomposition on the sample variance of  $Y_t$ , such that  $\hat{\Sigma}_Y = \hat{C} \hat{D} \hat{C}^\top$ . Because  $u_t$  is a linear/affine transformation of  $Y_t$ , from the multivariate change of variables theorem, the Jacobian of such an inverse transformation is  $\det(\hat{C})$ , which, together with the approximately linear form of the quantity of interest as in (14), implies that detecting structural changes in the same approximately linear statistics of  $\hat{u}_t$ , denoted as  $\theta^*$ , would be identical to testing structural changes in  $\theta$ .<sup>8</sup>

The adjusted-range based EKS test statistic is obtained as follows. First, construct a multivariate CUSUM process

$$T_n^*(k) = kn^{-1/2} (\hat{\theta}_{1,k}^* - \hat{\theta}_{1,n}^*). \tag{16}$$

Second, put  $T_n^*(k) = (T_n^{(1)*}(k), \dots, T_n^{(q)*}(k))^\top$ . Third, generate a  $q \times 1$  vector

$$R_n = \begin{pmatrix} \max_{1 \leq k \leq n} (T_n^{(1)*}(k)) - \min_{1 \leq k \leq n} (T_n^{(1)*}(k)) \\ \vdots \\ \max_{1 \leq k \leq n} (T_n^{(q)*}(k)) - \min_{1 \leq k \leq n} (T_n^{(q)*}(k)) \end{pmatrix}. \tag{17}$$

The adjusted-range based covariance estimator for  $\theta^*$  is  $\text{diag}\{R_n^2\}$ . The EKS test statistic is

$$\mathbb{EKS}^R(q) = \max_{1 \leq k \leq n-1} T_n^*(k)^\top [\text{diag}\{R_n^2\}]^{-1} T_n^*(k). \tag{18}$$

**Theorem 3.** Suppose Assumptions 2 and 3 hold. Then under the null hypothesis  $\mathbb{H}_0^{(2)}$ , as  $n \rightarrow \infty$ ,

$$\mathbb{EKS}^R(q) \xrightarrow{d} W_q, \tag{19}$$

where  $W_q$  is defined in (10).

The proof for Theorem 3 is relegated to Appendix A.2. Table 2 provides the simulated critical values.

#### 4. Testing parameter constancy

We now demonstrate that the adjusted range-based KS/EKS statistics can be used to test for parameter constancy. The notations and assumptions largely follow those of Chan et al. (2021), who use the SN approach of Shao (2010) and Lobato (2001) for sequential change point monitoring. See Chan et al. (2021) for a list of references adopting similar asymptotic settings.

Suppose that  $\{X_t\}_{t=1}^n$  is a stationary ergodic time series sample, with the joint density  $f_\theta$ , where  $\theta \in \mathbb{R}^d$  lies in a compact space  $\Theta$ , where  $d \geq 1$  is a positive integer.  $\{f_\theta : \theta \in \Theta\}$  can be regarded as a class of parametric models indexed by parameter  $\theta$ . The parameter of interest  $\theta$  satisfies  $E[L(X_t, \theta)] = \mathbf{0}$ . As a result,  $\theta$  can be consistently estimated by solving the system of equations

<sup>7</sup> As it can be seen from Section 2.2, our transformation methods are “partial prewhitening” in the sense that we only remove cross dependence, but not own temporal dependence of a time series. However, for series that suffer from a high level of heteroskedasticity and/or autocorrelation, and when we are testing for structural breaks in the mean, prewhitening will help to reduce serial dependence.

<sup>8</sup> If we have explicit information or would like to impose the condition that different components of  $\theta$  are uncorrelated, as in the case when  $\theta$  comprises the mean and variance of normally distributed random variables, then  $C$  should be an identity matrix.

$\sum_{t=1}^n L(\mathbf{X}_t, \hat{\theta}) = \mathbf{0}$ . This framework includes classical estimators such as ML estimators, M-estimators, least-squares estimators, and generalized method of moments estimators (Chan et al., 2021). We are interested in testing the null hypothesis

$$\mathbb{H}_0^p : \theta = \theta_0,$$

against the alternative hypothesis

$$\mathbb{H}_1^p : \mathbb{H}_0^p \text{ is false.}$$

Let  $L'(\mathbf{X}_t, \theta)$  be the gradient matrix of  $L(\mathbf{X}_t, \theta)$  with respect to the parameter  $\theta$ . Denote  $\|c\|$  as the supremum norm of a vector  $c$ . Define the matrix norm of a matrix  $A$  as  $\|A\| = \sup_{x: \|x\|=1} \|Ax\|$ . We impose the following regularity conditions as in Chan et al. (2021).

**Assumption 4.** The true parameter value  $\theta_0$  is in the interior region of  $\Theta$ , where  $\Theta$  is a compact set of  $\mathbb{R}^d$ .

**Assumption 5.** The time series process  $\{\mathbf{X}_t\}$  is stationary and ergodic.

**Assumption 6.**  $E \left[ \sup_{\theta \in \Theta} \|L(\mathbf{X}_t, \theta)\| \right] < \infty$  and  $\theta_0$  is the unique zero solution of  $E[L(\mathbf{X}_t, \theta)]$ . That is, for any given constant  $\epsilon > 0$ , there exists a constant  $\kappa > 0$ , such that  $E[L(\mathbf{X}_t, \theta)] > \kappa$  for all  $\theta$ , with  $\|\theta - \theta_0\| > \epsilon$ .

**Assumption 7.**  $E \left[ \sup_{\theta \in \Theta} \|L(\mathbf{X}_t, \theta_0)\|^{2+\delta} \right] < \infty$ , for some  $\delta > 0$ , and  $\{\mathbf{X}_t\}$  is a strong mixing sequence with mixing coefficients  $\alpha_k$  satisfying  $\sum_{k=1}^{\infty} \alpha_k^{\delta/(2+\delta)} < \infty$ .

**Assumption 8.**  $L(\mathbf{X}_t, \theta)$  is continuously differentiable with respect to  $\theta$  in a neighborhood  $V_{\theta_0}$  of  $\theta_0$ . In addition,  $E[L'(\mathbf{X}_t, \theta_0)]$  is positive definite, and  $E \left( \sup_{\theta \in V_{\theta_0}} \|L'(\mathbf{X}_t, \theta)\| \right) < \infty$ .

We refer to  $\{L(\mathbf{X}_t, \hat{\theta})\}$  as “generalized residuals”. Under Assumptions 4–8, testing for parameter constancy is equivalent to testing structural breaks in  $\{L(\mathbf{X}_t, \hat{\theta})\}$ , which implies that structural break tests in mean, such as the G test proposed by Shao and Zhang (2010) and the KS type tests, are all asymptotically valid tests for parameter constancy under this framework. Here, we focus on developing the adjusted-range based KS and EKS tests.

First, we show the consistency of the parameter estimator  $\hat{\theta}$  and the invariance principle of the partial sum process  $S(k, \hat{\theta}) = n^{-1/2} \sum_{t=1}^k L(\mathbf{X}_t, \hat{\theta})$ , for  $k = 1, \dots, n$ , as stated in Lemma 1 below.

**Lemma 1.**

- (i) Under Assumptions 4–8,  $\hat{\theta} = \theta_0 + O_p(n^{-1/2})$ .
- (ii) Under Assumptions 5 and 7,  $S(k, \hat{\theta}) = n^{-1/2} \sum_{t=1}^{\lfloor sn \rfloor} L(\mathbf{X}_t, \hat{\theta}) \Rightarrow \Delta_M \mathbf{B}_d(s)$ , where  $\Delta_M \Delta_M^\top = \sum_{k=-\infty}^{\infty} E[L(\mathbf{X}_1, \theta_0) L(\mathbf{X}_{k+1}, \theta_0)^\top]$ .

Lemma 1 is similar to Lemma 1 of Chan et al. (2021), thus we refer its proof to those of Theorem 3 of Kirch and Kamgaing (2012) and Theorem 3.2.1 of Lin and Lu (1996) as cited in Chan et al. (2021).

With Lemma 1, we then apply the functional central limit theorem (FCLT) and obtain

$$R^M = \max_{1 \leq k \leq n} S(k, \hat{\theta}) - \min_{1 \leq k \leq n} S(k, \hat{\theta}) \xrightarrow{d} \Delta_M \left[ \sup_{s \in [0,1]} \mathbb{B}_d(s) - \inf_{s \in [0,1]} \mathbb{B}_d(s) \right].$$

The adjusted-range based tests for constancy of parameter follow from CMT. When the parameter dimension  $d = 1$ , the adjusted-range based test statistic for constancy of parameters boils down to

$$M^R = \max_{1 \leq k \leq n} \left| (R^M)^{-1} S(k, \hat{\theta}) \right|. \tag{20}$$

When  $d \geq 2$ , the construction of the test statistic depends on whether the “generalized residuals” are correlated or not, which in turn depends on the estimation method. Let  $L^*(X_t, \hat{\theta}) = \hat{C}^{-1} L(\mathbf{X}_t, \hat{\theta})$ , where  $\hat{C}$  is an identity matrix when  $\{L(\mathbf{X}_t, \hat{\theta})\}_{t=1}^n$  exhibits no cross dependence, e.g. when  $\hat{\theta}$  is obtained through estimating a correctly specified model by the ordinary least squares (OLS) method,  $\{L(\mathbf{X}_t, \hat{\theta})\}_{t=1}^n$  is the time series of residuals, which are uncorrelated. In general,  $\{L^*(X_t, \hat{\theta})\}_{t=1}^n$  can be correlated; for example, when we use the ML estimation,  $\{L(\mathbf{X}_t, \hat{\theta})\}_{t=1}^n$  correspond to the scores. It is well known that under a correctly specified parametric model, the variance covariance matrix of the scores is equal to the negative expected value of the Hessian matrix of the log-likelihood, according to the Fisher information equality.<sup>9</sup> In that case, following Section 2.2,  $\hat{C}$  is obtained through the LDL decomposition on the sample variance of  $\{L(\mathbf{X}_t, \hat{\theta})\}_{t=1}^n$ , which converges in probability to the Fisher information matrix.

<sup>9</sup> Noticeably, Chan et al. (2021) consider the scores when sequentially monitoring the changes in parameter in stochastic volatility (SV) models; Pape et al. (2021) detect structural shifts in the dynamic conditional correlation (DCC) model of Engle (2002) by testing breaks in the score of the quasi-likelihood.

When  $d \geq 2$ , define  $S^*(k, \hat{\theta}) = (S^{(1)*}(k, \hat{\theta}), \dots, S^{(d)*}(k, \hat{\theta}))^\top = n^{-1/2} \sum_{t=1}^k \hat{C}^{-1} L(\mathbf{X}_t, \hat{\theta})$ , and then we generate a  $d \times 1$  adjusted-range based self-normalizer

$$\tilde{R}_n^M = \begin{pmatrix} \max_{1 \leq k \leq n} (S^{(1)*}(k, \hat{\theta})) - \min_{1 \leq k \leq n} (S^{(1)*}(k, \hat{\theta})) \\ \vdots \\ \max_{1 \leq k \leq n} (S^{(d)*}(k, \hat{\theta})) - \min_{1 \leq k \leq n} (S^{(d)*}(k, \hat{\theta})) \end{pmatrix}.$$

Then, the adjusted-range based test statistic for parameter constancy is

$$\mathbb{M}^R = \max_{1 \leq k \leq n-1} S^*(k, \hat{\theta})^\top \left[ \text{diag} \left\{ \tilde{R}_n^M \right\} \right]^{-2} S^*(k, \hat{\theta}).$$

The asymptotic properties of the adjusted-range based test statistics for constancy of parameter are summarized in the following theorem.

**Theorem 4.** Suppose Assumptions 4–8 hold. Then under the null hypothesis  $\mathbb{H}_0^p$ , as  $n \rightarrow \infty$ ,

$$\mathbb{M}^R \xrightarrow{d} U \text{ for } d = 1, \quad \text{and } \mathbb{M}^R \xrightarrow{d} W_d, \text{ for } d \geq 2,$$

where  $U$  is defined in (5), and  $W_d$  is defined in (10).

The proof for Theorem 4 is omitted for brevity as it follows from CMT. The consistency of  $\mathbb{M}^R$  and  $\mathbb{M}^R$  follow from Theorems 1 and 2, respectively.

### 5. Simulation studies

In this section, we consider structural shifts in the mean of a multivariate series  $\{X_t\}$ . Simulation studies for the univariate test  $\text{KS}^R$ , the tests for structural breaks in the median, and the tests for constancy of correlations are all relegated to the supplementary material. Throughout this section, we set the number of Monte Carlo simulations to 1000, and the significance level to 5%.

#### 5.1. Data generating processes (DGPs)

**DGP1 [Simple homoskedastic errors].** We consider a level shift in a bivariate VAR(1) model. Put  $X_t = \Psi X_{t-1} + \varepsilon_t$ , where  $\Psi = \begin{pmatrix} 0.5 & 0.0 \\ 0.0 & 0.5 \end{pmatrix}$  and  $\{\varepsilon_t\}$  follows an i.i.d.  $MN(\mathbf{0}, I_2)$ , where “MN” stands for a “multivariate normal distribution” and  $I_2$  is a  $2 \times 2$  identity matrix. This case is identical to two independent series generated from a shift in level in a univariate linear AR process considered by Shao and Zhang (2010), being aligned together to form a bivariate series.

**DGP2 [VAR with homoskedastic errors].** Everything else remains the same as in DGP1, except that we now allow for some cross dependence in  $\{X_t\}$  and  $\{\varepsilon_t\}$ , i.e.  $\Psi = \begin{pmatrix} 0.5 & 0.1 \\ 0.1 & 0.5 \end{pmatrix}$  and  $\{\varepsilon_t\} \sim MN(\mathbf{0}, \Sigma_\varepsilon)$  and  $\Sigma_\varepsilon = \begin{pmatrix} 1.0 & 0.1 \\ 0.1 & 1.0 \end{pmatrix}$ .

**DGP3 [VAR with conditional heteroskedastic errors].** This case is similar to DGP2, except that now the error terms are conditionally heteroscedastic, such that  $\{\varepsilon_t\}$  follows a GARCH(1,1) process

$$\varepsilon_t = \Sigma_t^{1/2} e_t, \sigma_{i,t}^2 = (1 - \alpha_1 - \beta_1) + \alpha_1^2 \varepsilon_{i,t-1}^2 + \beta_1 \sigma_{i,t-1}^2, \quad i = 1, 2,$$

where  $\Sigma_t = \begin{pmatrix} \sigma_{1,t}^2 & 0 \\ 0 & \sigma_{2,t}^2 \end{pmatrix}$ ,  $(\alpha_1, \beta_1) = (0.1, 0.79)$ , and  $\{e_t\}$  is a vector of innovations following an i.i.d.  $MN(\mathbf{0}, \Sigma_\varepsilon)$ .

**DGP4 [VAR with unconditional heteroskedastic errors].** This case resembles DGP2 and DGP3, except that now there exists a structural break in volatilities:

$$\sigma_{i,t}^2 = \sigma_0 [1 + \delta I(t > n/2)], \quad \sigma_0 = \delta = 1, \quad i = 1, 2.$$

We consider the following structural breaks, and set  $\eta \in \{0.5, 1.0, \dots, 2.0\}$ .

#### (i) Level shift:

$$Y_t = \begin{cases} X_t, & 1 \leq t \leq \lfloor n/2 \rfloor, \\ \eta + X_t, & \lfloor n/2 \rfloor + 1 \leq t \leq n. \end{cases}$$

#### (ii) Smooth changes/multiple breaks:

$$Y_t = X_t + \eta(t/n).$$

**Table 3**  
 Sizes, power, and size-adjusted power of the EKS<sup>R</sup>(2) and G(2) tests for detecting structural change(s) in the mean.

		$\eta$	n = 250				n = 500			
			DGP1	DGP2	DGP3	DGP4	DGP1	DGP2	DGP3	DGP4
EKS	Size	0	0.062	0.068	0.085	0.079	0.057	0.056	0.068	0.065
G	Size	0	0.099	0.095	0.119	0.124	0.112	0.106	0.095	0.092
Break type (i)										
EKS	Power	0.5	0.437	0.246	0.283	0.217	0.745	0.457	0.496	0.349
		1.0	0.898	0.711	0.687	0.569	0.984	0.961	0.951	0.847
		1.5	0.868	0.861	0.836	0.770	0.935	0.977	0.950	0.953
		2.0	0.739	0.835	0.803	0.816	0.841	0.935	0.882	0.944
	Size-adjusted power	0.5	0.405	0.202	0.209	0.172	0.729	0.446	0.441	0.310
		1.0	0.882	0.625	0.573	0.478	0.981	0.960	0.923	0.818
		1.5	0.845	0.810	0.744	0.699	0.929	0.972	0.928	0.939
		2.0	0.708	0.776	0.709	0.750	0.831	0.930	0.852	0.928
G	Power	0.5	0.541	0.394	0.416	0.305	0.808	0.593	0.639	0.458
		1.0	0.957	0.857	0.872	0.728	0.996	0.982	0.973	0.932
		1.5	1.000	0.988	0.988	0.948	1.000	1.000	0.999	0.997
		2.0	1.000	0.999	1.000	0.995	1.000	1.000	1.000	1.000
	Size-adjusted power	0.5	0.417	0.260	0.242	0.177	0.695	0.435	0.504	0.337
		1.0	0.916	0.750	0.720	0.591	0.992	0.959	0.960	0.865
		1.5	0.996	0.969	0.954	0.861	1.000	0.999	0.998	0.993
		2.0	1.000	0.995	0.995	0.980	1.000	1.000	1.000	1.000
Break type (ii)										
EKS	Power	0.5	0.187	0.148	0.145	0.114	0.384	0.236	0.241	0.186
		1.0	0.589	0.373	0.364	0.290	0.898	0.699	0.659	0.498
		1.5	0.848	0.650	0.676	0.514	0.969	0.936	0.896	0.803
		2.0	0.856	0.785	0.781	0.675	0.927	0.963	0.943	0.932
	Size-adjusted power	0.5	0.164	0.111	0.087	0.082	0.365	0.227	0.197	0.157
		1.0	0.551	0.304	0.282	0.231	0.882	0.682	0.604	0.457
		1.5	0.812	0.558	0.547	0.439	0.965	0.930	0.867	0.765
		2.0	0.829	0.708	0.677	0.589	0.920	0.956	0.918	0.901
G	Power	0.5	0.295	0.216	0.207	0.195	0.427	0.326	0.323	0.245
		1.0	0.562	0.474	0.520	0.380	0.708	0.628	0.610	0.536
		1.5	0.733	0.624	0.669	0.599	0.761	0.761	0.754	0.695
		2.0	0.749	0.749	0.721	0.688	0.789	0.755	0.767	0.745
	Size-adjusted power	0.5	0.176	0.123	0.096	0.097	0.282	0.202	0.210	0.163
		1.0	0.414	0.340	0.321	0.238	0.550	0.464	0.470	0.385
		1.5	0.591	0.436	0.462	0.428	0.634	0.619	0.611	0.566
		2.0	0.624	0.577	0.504	0.498	0.640	0.598	0.639	0.598

5.2. Structural breaks in mean

Given the presence of conditional/unconditional heteroskedastic errors in DGP3 and DGP4, we consider the LDL decomposition of the sample variance VAR prewhitened errors  $\{\hat{e}_t\}$ . This is because, under DGP3 and DGP4, the LDL decomposition based on the sample variance of  $\{X_t\}$  leads to a slightly over-inflated size, when the sample size  $n$  is small.<sup>10</sup>

The results for size, power, and size-adjusted power for the proposed EKS test and Shao and Zhang’s (2010) G test are summarized in Table 3. The EKS test demonstrates reasonable size and power. The size performance of the EKS test is better than that of the G test. On the other hand, the G test has higher power under the single structural break in (i), a result that aligns with our expectation, since the G test is formulated under one structural break point. However, its power performance is worse than EKS under smooth changes/multiple structural breaks in (ii).

5.3. Further discussion

In the supplementary material, we compare the performances of the proposed univariate KS<sup>R</sup> test with Shao and Zhang’s (2010) G test, the KS tests using the SN approach of Shao (2010) and Lobato (2001) (KS<sup>V</sup>), and the KS<sup>0</sup> test based on standard asymptotics. We also compare the performance of our proposed KS<sup>R</sup> test with Zhang and Lavitas’s (2018) T test. Furthermore, we consider the structural breaks in the median for the same DGPs in Section 5.1. Finally, we apply the proposed ESK test to test the constancy of the correlation matrix. We report powers using both asymptotic and empirical critical values respectively, we place our focus on the size-adjusted power of various tests for a fair comparison. The main findings are summarized as follows.

<sup>10</sup> The results are available from the authors upon request.

First, unlike the  $KS^V$  test, our proposed  $KS^R$  test does not suffer from the notorious decreasing power problem when the break size increases, which is illustrated by Fig. 1 in Shao and Zhang (2010). Moreover, because there is no need to use forward and backward summations as in the G test of Shao and Zhang (2010),  $KS^R$  is much more computationally efficient.

Second,  $KS^R$  offers superior power and size-adjusted power compared to Shao and Zhang's (2010) G test when addressing gradual shifts in mean. On the other hand, the G test typically surpasses  $KS^R$  under the one change-point alternative, viz., the specific scenario for which it is designed. Notably, simulation studies indicate that the adjusted-range based  $KS^R$  test might outperform Shao and Zhang's (2010) G test even when there is only one change point. This is particularly the case when the signal-to-noise ratio is low, and when the error terms follow a highly skewed Gamma distribution. The first set of results further confirms the findings of Mandelbrot (1972, 1975) regarding the appealing property of almost-sure convergence of the range statistic for stochastic processes with infinite variance; the second set of results further highlights the robustness of the range.

Third, under the same DGPs as in Zhang and Lavitas (2018), we find that  $KS^R$  delivers more accurate sizes when the level of autocorrelation in  $\{X_t\}$  is small; while Shao and Zhang's (2010) G test is more powerful than  $KS^R$  under the one change-point alternative,  $KS^R$  has good power under exact self-canceling breaks; while the G test has low power (DPG2), and both  $KS^R$  and the G test have inadequate size-adjusted-powers under oscillating breaks (DGP3).

Fourth, for testing structural breaks in median, Shao and Zhang's (2010) G test may suffer from the "over-size" problem; whereas the  $EKS^R(q)$  test delivers better power and size-adjusted powers under DGP2, DGP3 and DGP4 — multivariate series with autocorrelation and/or conditional/unconditional heteroskedasticity.

Fifth, the adjusted-range based EKS test for constancy of the correlation matrix, denoted as  $\mathbb{H}^R(q)$ , demonstrates adequate sizes, powers and size-adjusted powers, which show its merit in volatility modeling. In contrast, Shao and Zhang's (2015) G test suffers from an "over-size" problem; its size-adjusted powers are also lower than those of the  $\mathbb{H}^R(q)$  test.

Sixth, the adjusted-range based KS and EKS tests pose a substantially smaller computational burden. While the theoretical frameworks of Shao and Zhang (2010) and Zhang and Lavitas (2018) consider the approximately linear statistics in a multivariate context, both of their simulation studies focus on univariate cases, apparently due to computational cost. The computational burden for Zhang and Lavitas's (2018) T test statistic is huge for multivariate cases. In fact, the computational burden for Zhang and Lavitas's (2018) T test statistic is so severe that Zhang and Lavitas (2018) introduce a grid approximation scheme.

Finally, we find, through simulation studies, that for statistical quantities which vary slowly over time, such as the median, or become "almost constants" as the estimation horizon increases, such as the correlations, Shao and Zhang's (2010) G test suffers from an "over-size" problem. This is exactly the opposite of the "better size and less power" phenomenon documented in the SN literature (Shao, 2010; Zhang et al., 2011; Wang and Shao, 2022). Arguably, in spite of formulating multiple piecewise stationary partitions according to the prespecified change points, in order to reduce the increases of the denominator/self-normalizer, Shao and Zhang's (2010) G test still relies on the SN approach of Lobato (2001) and Shao (2010), whose self-normalizers are the variances of partial sum processes. For a robust statistical quantity that does not change much (e.g. the median), and is sometimes even "almost constants" (e.g. the correlation), as the estimation horizon increases, the variance of such a partial sum can become quite small, which can lead to over-rejection of the null hypothesis.

## 6. Empirical application

Motivated by the fact that range has been widely applied in volatility estimation (Parkinson, 1980; Alizadeh et al., 2002; Chou et al., 2010), we consider structural changes in conditional heteroskedasticity in five of the world's major stock indices from the 1st January 2012 to the 31st December 2020. The stock indices considered are the Dow Jones Industrial Average (DJIA), the S&P Composite Index, the FTSE 100, the CAC 40 and the DAX, which cover the stock markets in the United States, Canada, the United Kingdom, France and Germany.<sup>11</sup>

Not all markets open on the same days, but too many observations would get lost if we were to remove all the days when there were no observations. Thus, we only remove weekend days, we use the R command "na.interp" in the "forecast" package to interpolate the indices, and compute the continuously compounded returns, such that  $r_{j,t} = 100(\ln P_{j,t} - \ln P_{j,t-1})$ , where  $P_{j,t}$  is the closing price of the stock index  $j$  at day  $t$ . There are  $n = 2385$  observations in each return series  $\{r_{j,t}\}$ ,  $1 \leq t \leq n$  and  $1 \leq j \leq 5$ . The rates of return for the DJIA are visualized in Fig. 1, which indicates significant volatility clustering and dependence; particularly, the volatility burst due to COVID-19.<sup>12</sup>

The descriptive statistics for daily returns for each stock index are summarized in Table 4. Because the return series are serially dependent, the normality test of Bai and Ng (2005) is applied here, instead of Bera and Jarque's (1981) JB test.

Following the standard practice in modeling return series by first considering temporal dependence in conditional mean and then conditional heteroskedasticity, we first consider the parameter constancy of the conditional mean equation for the ARMA(1,1)-GARCH(1,1) models, and then extend our approach to multivariate conditionally heteroskedastic scenarios, viz. checking the rationality of both the constant correlation (CC) model (Bollerslev, 1990) and the DCC model (Engle, 2002), both of which are widely used in modeling multivariate volatility in empirical finance.

<sup>11</sup> The reason why we only consider five stocks here is due to the "curse of dimensionality". For a  $p \times p$  correlation matrix, there are  $q = p(p-1)/2$  correlation coefficients to be tested. An increase of one dimension for a  $p \times p$  correlation matrix, will result in an increase of  $p$  degrees of freedom, i.e.  $(p+1)p/2 - p(p-1)/2 = p$ . This will greatly increase the computational burden, especially for Shao and Zhang's (2010) G test statistic.

<sup>12</sup> The patterns of rates of return for the other stock indices are similar, and are relegated to the supplementary material due to page limits.




Fig. 1. Daily continuously compounded rates of return for the DJIA.

Table 4  
Summary statistics of daily continuously compounded rates of return for five major stock indices.

	Min	Max	Mean	Std Dev	Skewness	Kurtosis	ADF test	Normality test
Dow Jones	-13.8418	10.7643	0.0398	1.0571	-1.1809	28.8425	-12.3617***	12.4313***
SP Composite	-13.1761	11.2945	0.0171	0.9273	-1.8542	48.2642	-12.4516***	28.9615***
FTSE 100	-11.5125	8.6667	0.0089	0.9820	-0.8562	14.2314	-13.7099***	6.6640**
CAC 40	-13.0983	8.0561	0.0188	1.2074	-0.8108	10.7613	-13.7702***	7.4173**
DAX	-13.0549	10.4143	0.0303	1.2123	-0.6684	10.5644	-13.5273***	6.3851**

Note: \*\*\*, \*\* and \* stand for significance at the 1%, 5% and 10% significance levels, respectively.

Table 5  
Average statistics and rejection rates for testing the constancy of parameters for the mean equation of the ARMA(1,1)-GARCH(1,1) model.

	Average statistics			Rejection rates		
	M <sup>R</sup>	KS <sup>0</sup>	G	M <sup>R</sup>	KS <sup>0</sup>	G
Dow Jones	0.7061	0.7926	10.2119	0.1002	0.0000	0.0212
SP Composite	0.7489	0.9672	13.2733	0.1119	0.0159	0.0366
FTSE 100	0.6651	0.6719	7.9438	0.0440	0.0053	0.0069
CAC 40	0.6825	0.7133	9.6142	0.0429	0.0021	0.0090
DAX	0.6838	0.7403	9.6745	0.0551	0.0011	0.0027

6.1. Parameter constancy of the conditional mean equation for the ARMA(1,1)-GARCH(1,1) models

Similar to Chan et al. (2021), we specify an ARMA(1,1)-GARCH(1,1) model for each stock index return series, viz.

$$r_t = \phi_0 + \phi_1 r_{t-1} + \xi_t + \psi \xi_{t-1} = \mu_t + \xi_t, \tag{21}$$

$$\xi_t = \sigma_t v_t, \sigma_t^2 = \omega + \alpha r_{t-1}^2 + \beta \sigma_{t-1}^2, \tag{22}$$

where (21) and (22) are mean and variance equations respectively,  $\theta = (\phi_0, \phi_1, \psi, \omega, \alpha, \beta)$  are parameters, and  $v_t$  represents an unobservable shock to  $\xi_t$ , which is usually assumed to be i.i.d. with zero mean and unit variance. The conditional mean of  $r_t$  based on  $F_{t-1}$ , the information set at  $t - 1$ , is  $\hat{\mu}_t = E(r_t | F_{t-1}) = \hat{\phi}_0 + \hat{\phi}_1 r_{t-1} + \hat{\psi} \hat{\xi}_{t-1}$ .

Thus, the parameter constancy test  $M^R$  for conditional mean equations is obtained by plugging  $S(k, \hat{\theta}) = n^{-1/2} \sum_{t=1}^k (r_t - \hat{\mu}_t)$  into (20). Additionally, we compute the KS test statistic,  $KS^0$ , based on standard asymptotics and Shao and Zhang's (2010) G test statistic.<sup>13</sup> To analyze the data, we employ the rolling window estimation, using a window length of 500, which roughly equals the number of trading days in two years. We set the step size to 1 to ensure that all data points are included in the analysis. There are 1886 steps/windows in total. The results are summarized in Table 5. As indicated in Table 4, all the ADF tests strongly reject the null hypothesis of a unit root, thus favoring stationarity; we would expect the filtered series  $\{r_t - \hat{\mu}_t\}$  to be stationary, with a constant unconditional mean, which is supported by the low rejection rates for all statistics in Table 5. The results of  $M^R$  appear to be more reasonable, as the rejection percentages are close to the 5% significance level.

The test statistic values are visualized in Fig. 2,  $KS^0$  is omitted from the visualization, because it rarely rejects. Throughout the empirical analysis, the blue dashed lines represent the 5% critical values, and the time index corresponds to the end of each window.

<sup>13</sup> When generating  $KS^0$ , we apply the default setting for the "getLongRunVar" function in R, where the bandwidth selection follows Andrews's (1991) method. Because the SN approach of Shao (2010) can be viewed as a special case of the fixed- $b$  asymptotics in Kiefer and Vogelsang (2005), when  $b = 1$  and the kernel is the Bartlett kernel, we would like to keep the use of the kernel the same for comparison purposes.




Fig. 2. Statistic values for the proposed  $M^R$  test and the G test.




Fig. 3. Values of the proposed  $H^R(10)$  test and the G(10) test for suitability of the CC model.

From Fig. 2, we can see that the rejections based on  $KS^R$  are more evenly distributed across the sample, which is consistent with the overall stationarity of the rates of return of the DJIA, as seen from Table 4.<sup>14</sup>

6.2. Suitability of the CC model

Next, we consider the suitability of the CC model. Following Andreou and Ghysels’s (2003) procedure, we test for breaks in conditional correlations for normalized return series. For each rolling window, we first obtain the  $\hat{\sigma}_{j,t}^2$  for each stock index return series, using ARMA(1,1)-GARCH(1,1) models, in order to generate the normalized return  $r_{j,t}^* = r_{j,t}/\hat{\sigma}_{j,t}$ , for  $j = 1, \dots, 5$ . We then stack the normalized rates of return to form a 5-variate normalized return vector  $r_t^* = (r_{1,t}^*, r_{2,t}^*, \dots, r_{5,t}^*)'$ , and compute the proposed EKS test statistic  $H^R(10)$  and Shao and Zhang’s (2010) G test statistic for constancy of correlation coefficients among the 5-variate normalized return series,

$$\hat{\rho}_{1,k}^{ij} = \frac{\sum_{t=1}^k (r_{i,t}^* - \bar{r}_{i,k}^*) (r_{j,t}^* - \bar{r}_{j,k}^*)}{\sqrt{\sum_{t=1}^k (r_{i,t}^* - \bar{r}_{i,k}^*)^2 \sum_{t=1}^k (r_{j,t}^* - \bar{r}_{j,k}^*)^2}}$$

for  $1 \leq i \leq j \leq p$ ,  $p = 5$ , where  $\bar{r}_{i,k}^* = \sum_{t=1}^k r_{i,t}^*/k$  and  $\bar{r}_{j,k}^* = \sum_{t=1}^k r_{j,t}^*/k$ . Note  $\hat{\rho}_{1,k}^{ij}$  is the  $(i, j)$ ’th component from the sample correlation matrix calculated from the sub-sample  $t = 1, 2, \dots, k$ . In the supplementary material, we demonstrate, both through theoretical derivation and simulation studies, the use of the adjusted-range based EKS test and Shao and Zhang’s (2010) G test on testing constancy of correlation matrices. We perform the proposed EKS test and Shao and Zhang’s (2010) G test for the 1886 rolling windows, their averaged statistic values are 2.879 and 207.099, and the rejection rates are 0.411 and 0.371, respectively. Both results clearly suggest that the CC model is inadequate; in other words, the correlation structures among the 5 stock indices change over time. The statistic values are visualized in Fig. 3, the rejection patterns of both the  $H^R(10)$  test and the G test are similar.<sup>15</sup>

<sup>14</sup> The plots based on results from other stock indices reveal the same finding, see the supplementary material for detail.

<sup>15</sup> The maximum value of the G test statistics is 19,606.0600, which happens during the window ending on the 12th of March 2020, the value was removed from visualization as it pushes the plot almost flat. The value of the proposed  $H^R(10)$  test on that date is 5.0564, which shows up as a peak as well.

**Table 6**  
Average  $\mathbb{M}^R(8)$  and  $G(8)$  test statistics and rejection rates for constancy of parameter tests for the bivariate DCC-GARCH(1,1) model.

Stock indices		Average statistics		Rejection rates	
		$\mathbb{M}^R(8)$	$G(8)$	$\mathbb{M}^R(8)$	$G(8)$
Dow Jones	SP Composite	2.0931	205.8848	0.1625	0.1292
Dow Jones	FTSE 100	2.2257	180.3565	0.2168	0.0641
Dow Jones	CAC 40	2.1329	195.0296	0.1919	0.0913
Dow Jones	DAX	2.1666	197.6643	0.1948	0.1086
SP Composite	FTSE 100	2.1357	185.5688	0.1629	0.0782
SP Composite	CAC 40	2.0673	183.8792	0.1316	0.0885
SP Composite	DAX	2.0864	184.7184	0.1414	0.0815
FTSE 100	CAC 40	2.1365	201.9721	0.1489	0.1287
FTSE 100	DAX	2.1712	186.8859	0.1489	0.0698
CAC 40	DAX	2.2103	193.8053	0.1774	0.0698

6.3. Parameter constancy of the DCC model

Like Pape et al. (2021), we consider the suitability of the DCC model. To avoid the “curse of dimensionality” problem, we consider the following bivariate DCC-GARCH(1,1) model:

$$\tilde{r}_t = H_t^{1/2} \epsilon_t, \quad H_t = D_t S_t D_t, \tag{23}$$

$$h_{11,t} = c_{11} + a_{11,1} \tilde{r}_{1,t-1}^2 + g_{11,1} h_{11,t-1}, \quad h_{22,t} = c_{11} + a_{22,1} \tilde{r}_{1,t-1}^2 + g_{22,1} h_{22,t-1}, \tag{24}$$

$$S_t = \begin{pmatrix} 1 & s_{12,t} \\ s_{12,t} & 1 \end{pmatrix}, \quad s_{12,t} = \frac{q_{12,t}}{\sqrt{q_{11,t} q_{22,t}}}, \tag{25}$$

$$q_{12,t} = (1 - \alpha - \beta) + \alpha \frac{\tilde{r}_{1,t-1}}{\sqrt{h_{11,t}}} \frac{\tilde{r}_{2,t-1}}{\sqrt{h_{22,t}}} + \beta q_{12,t-1}, \tag{26}$$

$$q_{11,t} = (1 - \alpha - \beta) + \alpha \frac{\tilde{r}_{1,t-1}^2}{h_{11,t}} + \beta q_{11,t-1}, \quad q_{22,t} = (1 - \alpha - \beta) + \alpha \frac{\tilde{r}_{2,t-1}^2}{h_{22,t}} + \beta q_{22,t-1}, \tag{27}$$

where  $\{\tilde{r}_t\}$  is the filtered rates of returns by removing temporal dependence using ARMA(1,1) models,  $H_t$  is the conditional variance matrix of  $\{\tilde{r}_t\}$ ,  $\epsilon_t$  is the innovation at time  $t$ ,  $D_t$  is the diagonal matrix with conditional standard deviations, and  $S_t$  is the time-varying conditional correlational matrix at time  $t$ . There are eight parameters ( $\alpha, \beta, c_{11}, c_{22}, a_{11,1}, a_{22,1}, g_{11,1}, g_{22,1}$ ) in total. The estimation is conducted by assuming that  $\{\epsilon_t\}$  follows an i.i.d.  $MN(0, I_2)$ . When the multivariate normality of  $\{\epsilon_t\}$  is violated, the ML estimation method becomes the quasi-ML method. As demonstrated by Pape et al. (2021) and Section 4, testing parameter constancy is equivalent to testing structural breaks in scores. Similar to the first and second cases, we consider rolling window estimation, we set the length of the window to be 250 and the step to be 1, and thus, there are 2136 rolling windows in total. The reduction of window length is due to the computational burden of generating Shao and Zhang’s (2010) G test statistic.

The average statistics and rejection rates are presented in Table 6. The rejection rates based on  $G(8)$  are close to the 5% significance level, while those from  $\mathbb{M}^R(8)$  are considerably higher. Given that self-normalized tests, based on Shao and Zhang’s (2010) SN approach, tend to suffer from a “better size but less power” phenomenon, these results suggest that there might be periods in these stock markets when the DCC-GARCH(1,1) model does not fully capture the underlying dynamics. This is potentially because we specified the model to be the DCC-GARCH(1,1), which is a presumption rather than a model specification informed by the actual volatility dynamics. Nevertheless, for both the  $\mathbb{M}^R(8)$  and  $G(8)$  test statistics, the rejection rates are substantially lower for the CC model. This suggests the importance of considering dynamic changes in correlation structures.<sup>16</sup>

7. Conclusion

In this paper, we propose using the adjusted range of the partial sum of a time series as a novel self-normalizer instead of its sample variance, thus developing an alternative SN approach to that of Lobato (2001) and Shao and Zhang (2010). Since the range has the well-known robustness properties, the proposed adjusted-range based SN approach has the appealing properties of being robust to different types of structural breaks under different DGPs. Three scenarios are considered: testing for structural change in the mean of a time series, testing for structural changes for approximately linear statistics, and testing parameter constancy in time series regression. Testing for constancy of correlation coefficients/matrices is relegated to the supplementary materials, due to page limits.

Like Shao and Zhang’s (2010) G test, our proposed adjusted-range based KS and EKS test statistics do not involve any user specified inputs or tuning parameters. And there is also no need to use forward and backward summations or pre-specification

<sup>16</sup> Visualizations related to these findings are provided in the supplementary material.



of structural break points, as in Shao and Zhang’s (2010) G test statistic, or the construction of a contrast process and a grid approximation to speed computation even for univariate series as in Zhang and Lavitas (2018). As a result, the adjusted-range based KS and EKS test statistics can greatly simplify and speed up the computation involved.

Monte Carlo simulations show that the use of the adjusted-range as a self-normalizer can rectify the nonmonotonic power problem when the break size increases, which are present under the SN approach of Shao (2010) and Lobato (2001). In general, adjusted-range based test statistics offer reasonable sizes and adequate power even under autocorrelation and conditional/unconditional heteroskedastic errors, whereas Shao and Zhang’s (2010) G test is optimal if the break points are correctly specified. Notably, simulation studies indicate that the adjusted-range based  $KS^R$  test might outperform Shao and Zhang’s (2010) G test even when there is only one change point. This is particularly the case when the signal-to-noise ratio is low. Such results further confirm the findings of Mandelbrot (1972, 1975) regarding the appealing property of almost-sure convergence of the range statistic for stochastic processes with infinite variance. Another notable instance is when the error terms follow a highly skewed Gamma distribution, the adjusted-range based  $KS^R$  can outperform Shao and Zhang’s (2010) G, which further underscores the robustness of the range.

Our simulation results also confirm the merits of the adjusted-range based KS type statistics. In particular, for statistical quantities that do not vary much over time, such as medians and correlation coefficients, Shao and Zhang’s (2010) G test statistic suffers from an “over-size problem”, which supplements the existing finding that the self-normalized tests usually suffer from a “better size but less power” problem (Shao, 2010; Zhang et al., 2011; Wang and Shao, 2022). Finally, the empirical studies demonstrate the merit of the adjusted-range based KS type statistics in examining the suitability of CC and DCC models.

As a generally applicable SN approach, we could extend the adjusted-range based SN approach to construct confidence intervals, detecting parameter changes sequentially, estimating the locations of break points, or extending the structural break tests to functional data possibly of infinite dimension. These topics will be pursued in subsequent research.

**Acknowledgments**

The research is supported by National Natural Science Foundation of China (NSFC) grants No. 71988101 and No. 72173120. We would like to thank Serena Ng (Editor), an associate editor, two anonymous referees, as well as the participants at Gregory Chow Seminar in Center of Forecasting Sciences, Chinese Academy of Sciences, and Paula and Gregory Chow Institute for Studies in Economics, Xiamen University (2023), Australian Meeting of the Econometric Society (2021), North American Summer Meeting of the Econometric Society (2021) and the 4th International Conference on Computational and Financial Econometrics, for their insightful comments and suggestions, which have led to significant improvements to our paper.

**Appendix A. Proof of the main results**

*A.1. Proof of Theorem 1*

As discussed in the supplementary material, the lower and upper bounds of  $KS^R$  are 0 and 1, respectively. Therefore, to establish the consistency of  $KS^R$ , it suffices to demonstrate that  $KS^R = 1$  almost surely (a.s.). Under the one change-point alternative hypothesis  $\mathbb{H}_1^{(1)*}$ , we have  $X_t = x_t$  for  $1 < t \leq k^*$  and  $X_t = y_t = x_t + \delta$  for  $k^* < t \leq n$ ; finally, set  $\bar{x} = n^{-1} \sum_{t=1}^n x_t \xrightarrow{P} \mu < \infty$  and  $s_0 = k^*/n$ .

Consider the mean of  $X_t$ ,

$$\begin{aligned} \bar{X}_n &= \frac{1}{n} \sum_{t=1}^n X_t = \frac{1}{n} \left[ \sum_{t=1}^{k^*} x_t + \sum_{t=k^*+1}^n (x_t + \delta) \right] = \frac{1}{n} \left[ \sum_{t=1}^n x_t + (n - k) \delta \right] \\ &= \bar{x} + (1 - s_0) \delta \xrightarrow{P} \mu + (1 - s_0) \delta < \infty \text{ if } \delta \text{ is fixed.} \end{aligned}$$

At  $k = k^*$ ,  $T_n(k) \rightarrow \infty$  when  $n \rightarrow \infty$ . To see how, first consider  $1 < k \leq k^*$ ,

$$T_n(k) = n^{-1/2} \sum_{t=1}^k (X_t - \bar{X}_n) = n^{-1/2} \sum_{t=1}^k [(x_t - \bar{x}) - (1 - s_0) \delta],$$

where  $\bar{x} = n^{-1} \sum_{t=1}^n x_t$ ; so for each increment  $(X_t - \bar{X}_n)$ , there is an “ $-n^{-1/2} (1 - s_0) \delta$ ” negative/positive shift, if  $\delta$  is positive/negative; consequently, the resulting CUSUM process  $T_n(k)$  has a deterministic downward/upward trend. However, when  $k^* < k \leq n$ ,

$$T_n(k) = \frac{1}{n^{1/2}} \sum_{t=1}^{k^*} (X_t - \bar{X}_n) = \frac{1}{n^{1/2}} \sum_{t=1}^{k^*} [(x_t - \bar{x}) - (1 - s_0) \delta] + \sum_{t=k^*+1}^n [(x_t - \bar{x}) + \delta s_0],$$

$T_n(k)$  begins to exhibit an upward/downward trend instead; in other words, the trend gets reversed.

Specifically,  $T_n(k)$  can be written as

$$T_n(k) = \frac{1}{n^{1/2}} \left( \sum_{t=1}^k X_t - \bar{X}_n \right) = \frac{1}{n^{1/2}} \left[ \frac{n-k}{n} \sum_{t=1}^k X_t - \frac{k}{n} \sum_{t=k+1}^n X_t \right]$$

$$= \frac{(n-k)}{n^{3/2}} \sum_{t=1}^k X_t - \frac{k}{n^{3/2}} \sum_{t=k+1}^n X_t = \frac{k(n-k)}{n^{3/2}} \left[ k^{-1} \sum_{t=1}^k X_t - (n-k)^{-1} \sum_{t=k+1}^n X_t \right].$$

As a result,

$$\begin{aligned} T_n(k^*) &= \frac{k^*(n-k^*)}{n^{3/2}} \left[ (k^*)^{-1} \sum_{t=1}^{k^*} (x_t) - (n-k^*)^{-1} \sum_{t=k^*+1}^n (x_t + \delta) \right] \\ &= \frac{k^*(n-k^*)}{n^{3/2}} \left\{ (k^*)^{-1} \sum_{t=1}^{k^*} (x_t - \mu) - (n-k^*)^{-1} \sum_{t=k^*+1}^n (x_t - \mu) - \delta \right\} \\ &= \frac{n-k^*}{n^{3/2}} \sum_{t=1}^{k^*} (x_t - \mu) - \frac{k^*}{n^{3/2}} \sum_{t=k^*+1}^n (x_t - \mu) - \frac{k^*(n-k^*)}{n^{3/2}} \delta; \end{aligned}$$

if  $s_0 \in (0, 1)$ , we have

$$T_n(k^*) \xrightarrow{d} s_0^{1/2} (1-s_0) \sigma_x B(1) - s_0 (1-s_0)^{1/2} \sigma_x \widehat{B}(1) - s_0 (1-s_0) (n^{1/2} \delta),$$

where  $B(1)$  and  $\widehat{B}(1)$  are two independent copies, and  $\sigma_x^2$  is the LRV of  $x$ . Thus, if  $\delta > 0$ ,  $T_n(k^*) \rightarrow -\infty$  and  $\max_{1 \leq k \leq n} T_n(k) / T_n(k^*) = o_p(1)$ ; and if  $\delta < 0$ ,  $T_n(k^*) \rightarrow \infty$  and  $\min_{1 \leq k \leq n} T_n(k) / T_n(k^*) = o_p(1)$ .

Consequently, when  $\delta > 0$ , and as  $n \rightarrow \infty$ ,

$$KS^R = \frac{\max_{1 \leq k \leq n} |T_n(k)|}{\max_{1 \leq k \leq n} T_n(k) - \min_{1 \leq k \leq n} T_n(k)} = \frac{|T_n(k^*)|}{\max_{1 \leq k \leq n} T_n(k) - T_n(k^*)} = 1 \text{ a.s.}$$

When  $\delta < 0$ , and as  $n \rightarrow \infty$ ,

$$KS^R = \frac{\max_{1 \leq k \leq n} |T_n(k)|}{\max_{1 \leq k \leq n} |T_n(k)| - \min_{1 \leq k \leq n} T_n(k)} = \frac{|T_n(k^*)|}{T_n(k^*) - \min_{1 \leq k \leq n} T_n(k)} = 1 \text{ a.s.}$$

If  $\delta = n^{-1/2} \eta$  and  $\eta \neq 0$ , we have

$$T_n(k^*) \xrightarrow{d} s_0^{1/2} (1-s_0) \sigma_x B(1) - s_0 (1-s_0)^{1/2} \sigma_x \widehat{B}(1) - s_0 (1-s_0) \eta.$$

As  $\eta \rightarrow \infty$ ,  $T_n(k^*) \rightarrow -\infty$ , and  $\max_{1 \leq k \leq n} T_n(k) / T_n(k^*) = o_p(1)$ ; while as  $\eta \rightarrow -\infty$ ,  $T_n(k^*) \rightarrow \infty$ , and  $\min_{1 \leq k \leq n} T_n(k) / T_n(k^*) = o_p(1)$ ; thus, as  $n \rightarrow \infty$  and  $|\eta| \rightarrow \infty$ , we have  $KS^R = 1$  a.s.  $\square$

### A.2. Proof of Theorem 2

The proof of Theorem 2 is similar to that of Theorem 1.  $T_n(k) = (T_n^{(1)}(k), \dots, T_n^{(m)}(k))^T$ , where  $T_n^{(i)}(k) = n^{-1/2} \sum_{t=1}^k (X_{i,t} - \bar{X}_{i,n})$  and  $i = 1, \dots, m$ . Consider the case when  $C$  is an identity matrix, or equivalently,  $\{X_{i,t}\}$  are independent of each other,  $\mathbb{E}KS^R(m)$  can be expanded such that

$$\begin{aligned} \mathbb{E}KS^R(m) &= \max_{1 \leq k \leq n} \left[ \left( \frac{T_n^{(1)}(k)}{\max T_n^{(1)}(k) - \min T_n^{(1)}(k)} \right)^2 + \dots + \left( \frac{T_n^{(m)}(k)}{\max T_n^{(m)}(k) - \min T_n^{(m)}(k)} \right)^2 \right] \\ &\leq (KS^R(k_1^*))^2 + \dots + (KS^R(k_m^*))^2, \end{aligned} \tag{A.1}$$

where

$$KS^R(k_i^*) = \left( \frac{\max_{1 \leq k_i \leq n} |T_n^{(i)}(k_i)|}{\max_{1 \leq k_i \leq n} T_n^{(i)}(k_i) - \min_{1 \leq k_i \leq n} T_n^{(i)}(k_i)} \right)^2.$$

$KS^R(k_i^*)$ ,  $i = 1, \dots, m$ , are independent copies of each other; and “=” in (A.1) holds when  $k = k_1^* = \dots = k_m^*$ , which is precisely the alternative hypothesis under consideration.

Following the proof of Theorem 1,  $KS^R(k_i^*)$  attains its maximum at  $k_i^*$  which converges to 1. Therefore,

$$\mathbb{E}KS^R(m) = m \text{ a.s.}$$

as  $n \rightarrow \infty$ , when  $s_0 \in (0, 1)$  and  $\Delta_n \neq 0$  is fixed, alternatively, if  $\Delta_n = n^{-1/2} \eta$ ,  $\eta = (\eta^{(1)}, \dots, \eta^{(m)})^T \neq \mathbf{0}$ , and  $|\eta| \rightarrow \infty$ . The conclusion follows.

When we consider the case when there is cross dependence in  $X_{i,t}$ , all we need is the finite variance for  $X_i$  to obtain an estimate for  $C$ , which is guaranteed by Assumptions 1 or 2. We know that  $\widehat{C} \xrightarrow{p} C$  as  $n \rightarrow \infty$ . The result of Theorem 2 follows from CMT.  $\square$

A.3. Proof of Theorem 3

From (14), for  $1 \leq t_1 \leq t_2 \leq n$ , we have

$$t_1^* (\hat{\theta}_{1,t_1}^* - \hat{\theta}_{1,t_2}^*) = \left( \sum_{t=1}^{t_1} \mathbf{IF}(u_t; \tilde{\mathbf{F}}^m) - \frac{t_1}{t_2} \mathbf{IF}(u_t; \tilde{\mathbf{F}}^m) \right) + \left( t_1 \tilde{\mathbf{R}}_{1,t_1} - \frac{t_1}{t_2} \tilde{\mathbf{R}}_{1,t_2} \right), \tag{A.2}$$

where  $\tilde{\mathbf{F}}^m$  is the  $m$ th marginal distribution of  $\hat{u}_t$ . From Assumption 2, we have

$$n^{-1/2} \sum_{t=1}^{\lfloor sn \rfloor} \mathbf{IF}(\hat{u}_t; \tilde{\mathbf{F}}^m) \Rightarrow \tilde{\Delta \mathbf{B}}_q(s). \tag{A.3}$$

Because  $\hat{u}_t$  is a linear/affine transformation of  $Y_t$ , from Assumption 3, we know that both  $\tilde{\mathbf{R}}_{1,t_1}$  and  $\tilde{\mathbf{R}}_{1,t_2}$  are negligible, which combined with (A.3), indicates the joint convergences of

$$\mathbf{T}_n^*(k) \Rightarrow \tilde{\Delta \mathbf{B}}(s)$$

and

$$\left[ \text{diag}(\tilde{\mathbf{R}}_n) \right]^2 \Rightarrow \tilde{\Delta} \text{diag} \left( \sup_{s \in (0,1]} \mathbb{B}_q(s) - \inf_{s \in (0,1]} \mathbb{B}_q(s) \right)^2 \tilde{\Delta}^T.$$

The result of Theorem 3 follows from CMT.  $\square$

Appendix B. Supplementary data

Supplementary material related to this article can be found online at <https://doi.org/10.1016/j.jeconom.2023.105603>.

References

Alexander, C., 1998. Volatility and correlation: Measurement, models and applications. In: Alexander, C. (Ed.), Risk Management and Analysis. John Wiley, New York, USA, pp. 125–171.

Alizadeh, S., Brandt, M.W., Diebold, F.X., 2002. Range-based estimation of stochastic volatility models. *J. Finance* 57 (3), 1047–1091.

Altissimo, F., Corradi, V., 2003. Strong rules for detecting the number of breaks in a time series. *J. Econometrics* 117 (2), 207–244.

Andreou, E., Ghysels, E., 2003. Tests for breaks in the conditional co-movements of asset returns. *Statist. Sinica* 13 (4), 1045–1073.

Andrews, D.W., 1991. Heteroskedasticity and autocorrelation consistent covariance matrix estimation. *Econometrica* 59 (3), 817–858.

Andrews, D.W., 1993. Tests for parameter instability and structural change with unknown change point. *Econometrica* 61 (4), 821–856.

Andrews, D.W., Monahan, J.C., 1992. An improved heteroskedasticity and autocorrelation consistent covariance matrix estimator. *Econometrica* 60 (4), 953–966.

Aue, A., Hörmann, S., Horváth, L., Reimherr, M., 2009. Break detection in the covariance structure of multivariate time series models. *Ann. Statist.* 37 (6B), 4046–4087.

Aue, A., Horváth, L., 2013. Structural breaks in time series. *J. Time Series Anal.* 34 (1), 1–16.

Bai, J., 1997. Estimating multiple breaks one at a time. *Econom. Theory* 13 (3), 315–352.

Bai, J., Ng, S., 2005. Tests for skewness, kurtosis, and normality for time series data. *J. Bus. Econom. Statist.* 23 (1), 49–60.

Bai, J., Perron, P., 1998. Estimating and testing linear models with multiple structural changes. *Econometrica* 66 (1), 47–78.

Bera, A.K., Jarque, C.M., 1981. Efficient tests for normality, homoscedasticity and serial independence of regression residuals: Monte Carlo evidence. *Econom. Lett.* 7 (4), 313–318.

Billingsley, P., 1968. *Convergence of Probability Measures*. John Wiley & Sons, New York, USA.

Bollerslev, T., 1990. Modelling the coherence in short-run nominal exchange rates: A multivariate generalized ARCH model. *Rev. Econ. Stat.* 72 (3), 498–505.

Bollerslev, T., Engle, R.F., Wooldridge, J.M., 1988. A capital asset pricing model with time-varying covariances. *J. Polit. Econ.* 96 (1), 116–131.

Boubacar-Mainassara, Y., Saussereau, B., 2018. Diagnostic checking in multivariate ARMA models with dependent errors using normalized residual autocorrelations. *J. Amer. Statist. Assoc.* 113 (524), 1813–1827.

Brown, R.L., Durbin, J., Evans, J.M., 1975. Techniques for testing the constancy of regression relationships over time. *J. R. Stat. Soc. Ser. B Stat. Methodol.* 37 (2), 149–163.

Chan, N.H., Ng, W.L., Yau, C.Y., 2021. A self-normalized approach to sequential change-point detection for time series. *Statist. Sinica* 31 (1), 491–517.

Chen, J., Gupta, A.K., 1997. Testing and locating variance change-points with application to stock prices. *J. Amer. Statist. Assoc.* 92 (438), 739–747.

Choi, J.-E., Shin, D.W., 2020. A self-normalization test for correlation change. *Econom. Lett.* 193, 108363.

Choi, J.-E., Shin, D.W., 2021. A self-normalization break test for correlation matrix. *Statist. Papers* 62 (5), 2333–2353.

Chou, R.Y., Chou, H., Liu, N., 2010. Range volatility models and their applications in finance. In: Lee, C.-F., Lee, A.C., Lee, J. (Eds.), *Handbook of Quantitative Finance and Risk Management*. Springer, Boston, USA, pp. 1273–1281.

Den Haan, W.J., Levin, A.T., 1997. A practitioner’s guide to robust covariance matrix estimation. *Handbook of Statist.* 15, 299–342.

Dette, H., Kokot, K., Volgushev, S., 2020. Testing relevant hypotheses in functional time series via self-normalization. *J. R. Stat. Soc. Ser. B Stat. Methodol.* 82 (3), 629–660.

Engle, R., 2002. Dynamic conditional correlation: A simple class of multivariate generalized autoregressive conditional heteroskedasticity models. *J. Bus. Econom. Statist.* 20 (3), 339–350.

Gouriéroux, C., Monfort, A., Renne, J.-P., 2017. Statistical inference for independent component analysis: Application to structural VAR models. *J. Econometrics* 196 (1), 111–126.

Hamilton, J.D., 1994. *Time Series Analysis*. Princeton University Press, Princeton, USA.

Hampel, F.R., Ronchetti, E.M., Rousseeuw, P.J., Stahel, W.A., 1986. *Robust statistics: The approach based on influence functions*. John Wiley & Sons, New York, USA.

Hansen, B.E., 2001. The new econometrics of structural change: Dating breaks in US labour productivity. *J. Econ. Perspect.* 15 (4), 117–128.

Harchaoui, Z., Lévy-Leduc, C., 2010. Multiple change-point estimation with a total variation penalty. *J. Amer. Statist. Assoc.* 105 (492), 1480–1493.

Huang, Y., Volgushev, S., Shao, X., 2015. On self-normalization for censored dependent data. *J. Time Series Anal.* 36 (1), 109–124.

- Hyvarinen, A., 1999. Fast and robust fixed-point algorithms for independent component analysis. *IEEE Trans. Neural Netw.* 10 (3), 626–634.
- Inclan, C., Tiao, G.C., 1994. Use of cumulative sums of squares for retrospective detection of changes of variance. *J. Amer. Statist. Assoc.* 89 (427), 913–923.
- Jiang, F., Zhao, Z., Shao, X., 2023. Time series analysis of COVID-19 infection curve: A change-point perspective. *J. Econometrics* 232 (1), 1–17.
- Jin, H., Zhang, S., Zhang, J., Hao, H., 2018. Modified tests for change points in variance in the possible presence of mean breaks. *J. Stat. Comput. Simul.* 88 (14), 2651–2667.
- Kiefer, N.M., Vogelsang, T.J., 2002. Heteroskedasticity-autocorrelation robust testing using bandwidth equal to sample size. *Econom. Theory* 18 (6), 1350–1366.
- Kiefer, N.M., Vogelsang, T.J., 2005. A new asymptotic theory for heteroskedasticity-autocorrelation robust tests. *Econom. Theory* 21 (6), 1130–1164.
- Kiefer, N.M., Vogelsang, T.J., Bunzel, H., 2000. Simple robust testing of regression hypotheses. *Econometrica* 68 (3), 695–714.
- Kim, B.G., Shin, D.W., 2020. A mean-difference test based on self-normalization for alternating regime index data sets. *Econom. Lett.* 193, 108334.
- Kirch, C., Kamgaing, J.T., 2012. Testing for parameter stability in nonlinear autoregressive models. *J. Time Series Anal.* 33 (3), 365–385.
- Krämer, W., Ploberger, W., Alt, R., 1988. Testing for structural change in dynamic models. *Econometrica* 56 (6), 1355–1369.
- Kunsch, H.R., 1989. The jackknife and the bootstrap for general stationary observations. *Ann. Statist.* 17 (3), 1217–1241.
- Lin, Z., Lu, C., 1996. *Limit Theorems of Mixing Dependent Random Variables*. Academic Publisher, USA, New York.
- Lobato, I.N., 2001. Testing that a dependent process is uncorrelated. *J. Amer. Statist. Assoc.* 96 (455), 1066–1076.
- Mandelbrot, B., 1972. Statistical methodology for nonperiodic cycles: From the covariance to R/S analysis. *Ann. Econ. Soc. Meas.* 1 (3), 259–290.
- Mandelbrot, B.B., 1975. Limit theorems on the self-normalized range for weakly and strongly dependent processes. *Z. Wahrscheinlichkeitstheorie Verwandte Geb.* 31 (4), 271–285.
- Mandelbrot, B.B., Wallis, J.R., 1969. Computer experiments with fractional Gaussian noises: Part 1, averages and variances. *Water Resour. Res.* 5 (1), 228–241.
- Matteson, D.S., Tsay, R.S., 2017. Independent component analysis via distance covariance. *J. Amer. Statist. Assoc.* 112 (518), 623–637.
- Müller, U.K., 2007. A theory of robust long-run variance estimation. *J. Econometrics* 141 (2), 1331–1352.
- Newey, W.K., West, K.D., 1987. A simple, positive semi-definite, heteroskedasticity and autocorrelation consistent covariance matrix. *Econometrica* 55 (3), 703–708.
- Newey, W.K., West, K.D., 1994. Automatic lag selection in covariance matrix estimation. *Rev. Econom. Stud.* 61 (4), 631–653.
- Pape, K., Galeano, P., Wied, D., 2021. Sequential detection of parameter changes in dynamic conditional correlation models. *Appl. Stoch. Models Bus. Ind.* 37 (3), 475–495.
- Parkinson, M., 1980. The extreme value method for estimating the variance of the rate of return. *J. Bus.* 61–65.
- Phillips, P.C., 1987. Time series regression with a unit root. *Econometrica* 55 (2), 277–301.
- Pitarakis, J.-Y., 2004. Least squares estimation and tests of breaks in mean and variance under misspecification. *Econom. J.* 7 (1), 32–54.
- Ploberger, W., Krämer, W., 1992. The CUSUM test with OLS residuals. *Econometrica* 60 (2), 271–285.
- Qu, Z., Perron, P., 2007. Estimating and testing structural changes in multivariate regressions. *Econometrica* 75 (2), 459–502.
- Shao, X., 2010. A self-normalized approach to confidence interval construction in time series. *J. R. Stat. Soc. Ser. B Stat. Methodol.* 72 (3), 343–366.
- Shao, X., 2015. Self-normalization for time series: A review of recent developments. *J. Amer. Statist. Assoc.* 110 (512), 1797–1817.
- Shao, X., Zhang, X., 2010. Testing for change points in time series. *J. Amer. Statist. Assoc.* 105 (491), 1228–1240.
- Smith, D.R., 2008. Testing for structural breaks in GARCH models. *Appl. Financial Econ.* 18 (10), 845–862.
- Stock, J.H., Watson, M.W., 1996. Evidence on structural instability in macroeconomic time series relations. *J. Bus. Econom. Statist.* 14 (1), 11–30.
- Student, 1908. The probable error of a mean. *Biometrika* 6 (1), 1–25.
- Wang, R., Shao, X., 2020. Hypothesis testing for high-dimensional time series via self-normalization. *Ann. Statist.* 48 (5), 2728–2758.
- Wang, R., Zhu, C., Volgushev, S., Shao, X., 2022. Inference for change points in high-dimensional data via self-normalization. *Ann. Statist.* 50 (2), 781–806.
- Wang, J., Zivot, E., 2000. A Bayesian time series model of multiple structural changes in level, trend, and variance. *J. Bus. Econom. Statist.* 18 (3), 374–386.
- White, H., 1980. A heteroskedasticity-consistent covariance matrix estimator and a direct test for heteroskedasticity. *Econometrica* 48 (4), 817–838.
- Wied, D., 2017. A nonparametric test for a constant correlation matrix. *Econometric Rev.* 36 (10), 1157–1172.
- Wied, D., Krämer, W., Dehling, H., 2012. Testing for a change in correlation at an unknown point in time using an extended functional delta method. *Econom. Theory* 28 (3), 570–589.
- Zhang, T., Lavitas, L., 2018. Unsupervised self-normalized change-point testing for time series. *J. Amer. Statist. Assoc.* 113 (522), 637–648.
- Zhang, X., Li, B., Shao, X., 2014. Self-normalization for spatial data. *Scand. J. Stat.* 41 (2), 311–324.
- Zhang, X., Shao, X., Hayhoe, K., Wuebbles, D.J., 2011. Testing the structural stability of temporally dependent functional observations and application to climate projections. *Electron. J. Stat.* 5, 1765–1796.