

The PM10 fraction of road dust in the UK and India

Pant, Pallavi; Baker, Stephen J.; Shukla, Anuradha; Maikawa, Caitlin; Godri Pollitt, Krystal J.; Harrison, Roy M.

DOI:

[10.1016/j.scitotenv.2015.05.084](https://doi.org/10.1016/j.scitotenv.2015.05.084)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Peer reviewed version

Citation for published version (Harvard):

Pant, P, Baker, SJ, Shukla, A, Maikawa, C, Godri Pollitt, KJ & Harrison, RM 2015, 'The PM10 fraction of road dust in the UK and India: Characterization, source profiles and oxidative potential', *Science of the Total Environment*, vol. 530-531, pp. 445-452. <https://doi.org/10.1016/j.scitotenv.2015.05.084>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

After an embargo period this document is subject to the terms of a Creative Commons Attribution Non-Commercial No Derivatives license

Checked September 2015

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

1

2 **The PM₁₀ Fraction of Road Dust in the UK and**
3 **India: Characterization, Source Profiles and**
4 **Oxidative Potential**

5

6 **Pallavi Pant^{1†}, Stephen J. Baker¹, Anuradha Shukla²,**
7 **Caitlin Maikawa³, Krystal J. Godri Pollitt^{3,4} and**
8 **Roy M. Harrison^{*1††}**

9

10

11 **¹Division of Environmental Health and Risk Management, School of Geography,**
12 **Earth and Environmental Sciences, University of Birmingham, Edgbaston,**
13 **Birmingham, B15 2TT, United Kingdom**

14

15 **²Environmental Sciences Division, Central Road Research Institute, Mathura**
16 **Road, New Delhi 110025, India**

17

18 **³ Southern Ontario Centre for Atmospheric Aerosol Research, Department of**
19 **Chemical Engineering & Applied Chemistry, University of Toronto, 200 College**
20 **St., Toronto, Ontario M5S 3E5, Canada**

21 **⁴Department of Environmental Health Sciences, School of Public Health and**
22 **Health Sciences, University of Massachusetts, Amherst, MA 01003, USA**

23

* To whom correspondence should be addressed.
Tele: +44 121 414 3494; Email: r.m.harrison@bham.ac.uk

† Now at Department of Environmental Health Sciences, School of Public Health and Health Sciences, University of Massachusetts, Amherst, MA 01003, USA

†† Also at: Department of Environmental Sciences / Center of Excellence in Environmental Studies, King Abdulaziz University, PO Box 80203, Jeddah, 21589, Saudi Arabia

24 **ABSTRACT**

25 Most studies of road dust composition have sampled a very wide range of particle sizes, but from
26 the perspective of respiratory exposure to resuspended dusts, it is the PM₁₀ fraction which is of most
27 importance. The PM₁₀ fraction of road dust samples was collected at two sites in Birmingham, UK
28 (major highway and road tunnel) and one site in New Delhi, India. Dust loadings were found to be
29 much higher for New Delhi compared to Birmingham, while concentrations of several species were
30 much higher in the case of Birmingham. Detailed chemical source profiles were prepared for both
31 cities and previously generated empirical factors for source attribution to brake wear, tyre wear and
32 crustal dust were successfully applied to the UK sites. However, 100% of the mass for the Indian
33 site could not be accounted for using these factors. This study highlights the need for generation of
34 local empirical estimation factors for non-exhaust vehicle emissions. A limited number of bulk
35 road dust and brake pad samples were also characterized. Oxidative Potential (OP) was also
36 determined for a limited number of PM₁₀ and bulk road dust samples, and Cu was found to be a
37 factor significantly associated with OP in PM₁₀ and bulk road dust.

38

39 **Keywords:** Non-exhaust emissions; source profile; source apportionment; road dust; Oxidative
40 Potential

41

42 1. INTRODUCTION

43 Emissions from road vehicles are of importance for human health as they are emitted in the
44 vicinity of human activity, and street canyon effects can enhance pollutant levels by limiting
45 dispersion (Colville et al., 2001). Particulate matter (PM) emissions from road vehicles include
46 both exhaust and non-exhaust (wear and tear of vehicle parts, resuspension of dust) constituents
47 (Pant and Harrison, 2013). Non-exhaust emissions are more abundant in the coarse size fraction
48 (particles with aerodynamic diameter of 2.5-10 μm) and can deposit on the roadside and be
49 resuspended subsequently due to vehicle activity. Road dust (RD) consists primarily of coarse-sized
50 PM derived from different sources such as wear of vehicle components (brakes, tyres and
51 clutches), the road surface, engine corrosion, tailpipe emissions, crustal dust and other emission
52 sources. However, similarity in composition between direct emission sources such as brake wear
53 and tyre wear and particles resuspended from the road surface renders it difficult to distinguish the
54 contribution of the different sources (Bukowiecki et al., 2010). RD can be a significant source of
55 trace metals, particle-bound polycyclic aromatic hydrocarbons (PAHs) and other chemical species
56 in the atmosphere (Amato et al., 2009; Duong and Lee, 2011; Kwon and Castaldi, 2012). Several
57 studies on RD have reported elemental and organic marker concentrations for RD in different size
58 fractions (Han et al., 2007; Morillo et al., 2007; Agarwal, 2009; Amato et al., 2009; Faiz et al.,
59 2009; Duong and Lee, 2011; Gunawardana et al., 2011; Martuzevicius et al., 2011). Crustal dust is
60 characterized by elements such as Si, Al, Ti, Mn and Fe while non-exhaust emissions are typically
61 characterized by trace metals such as Cu, Ba, Sb, Sn (brake wear) and Zn (tyre wear) (Wahlin et al.,
62 2006; Gietl et al., 2010; Hulskotte et al., 2014). However, several authors have also used organic
63 markers (e.g. PAHs, n-alkanes, benzothiazoles) as source tracers (Lough et al., 2005; Kwon and
64 Castaldi, 2012). Enrichment of both trace elements and organic species has been reported in RD
65 compared to background soils worldwide (Han et al., 2007; Liu et al., 2007; Agarwal, 2009; Duong
66 and Lee, 2011; Luo et al., 2011; Peltier et al., 2011). Elements such as Pt, Pd and Rh have also

67 been reported to be present in RD and are attributed to emissions from catalytic converters
68 (Prichard and Fisher, 2012).

69

70 It is important to note that the chemical source profile (i.e. percentage of species with respect to
71 total PM mass) for contaminated soil as well as non-exhaust emissions varies geographically
72 dependent upon parameters such as traffic (volume and pattern, fleet characteristics), and road
73 surface type, and the climate and geology of the region (Omstedt et al., 2005; Amato et al.,
74 2011; Pant and Harrison, 2013; Kwak et al., 2013). Most of the research, as well as policy action,
75 in the last few decades has focused largely on exhaust emissions, and with a decline in percentage
76 contribution of vehicle exhaust emissions to ambient PM, the contribution of non-exhaust PM is of
77 increasing importance (Thorpe and Harrison, 2008; Denier van der Gon et al., 2013). Detailed
78 accounts of sources and properties of non-exhaust emissions and ambient measurements of non-
79 exhaust components are provided elsewhere (Luhana et al., 2004; Thorpe and Harrison, 2008; Pant
80 and Harrison, 2013; Amato et al., 2014). Typical methods for characterization of RD and non-
81 exhaust emissions include use of a brake dynamometer (Garg et al., 2000), rolling resistance testing
82 machine (Rogge et al., 1993), rotating drum method (Camatani et al., 2001), sweep/vacuum
83 collection of particles followed by sieving or resuspension (Duong and Lee, 2011; Martuzevicius et
84 al., 2011) and use of a specific RD sampler (Amato et al., 2009). Several researchers have analysed
85 the chemical composition of brake components (Rogge et al., 1993; Garg et al., 2000; Kukutschova
86 et al., 2010, 2011; Hulskotte et al., 2014) and tyres (Camatani et al., 2001; Adachi and Tainosho,
87 2004; Councill et al., 2004; Milani et al., 2004; Aatmeeyata and Sharma, 2010).

88

89 Much of the reported research has been conducted on bulk or very coarse fractions of RD,
90 with a very few studies focused on the characterization and source apportionment of the PM_{10}
91 fraction of RD (Wahlin et al., 2006; Amato et al., 2009; Han et al., 2009; Amato et al., 2011). In
92 order to quantify RD as a source of airborne PM, it is important to prepare detailed chemical source

93 profiles. Both India and UK lack detailed chemical source profiles for the RD source. For New
94 Delhi, source profiles for PM₁₀ RD (paved, unpaved and soil) were published as a part of a
95 larger study but the molecular markers were not analysed in detail (Sethi and Patil, 2008).

96
97 The aims of this study were to characterize the chemical composition of the PM₁₀ fraction RD, to
98 prepare a PM₁₀ RD source profile and to calculate the contribution of non-exhaust emission
99 sources to samples of RD in Birmingham (UK) and New Delhi (India). In addition, Oxidative
100 Potential of the RD samples from the UK was assessed. To the best of our knowledge, this is the
101 first study to undertake detailed elemental as well as molecular marker and oxidative
102 characterization of the PM₁₀ fraction of RD either in India or the UK.

103

104 **2. METHODS**

105 **2.1 Sampling Sites**

106 Samples were collected at two high traffic locations in Birmingham: Bristol Road- a heavily
107 trafficked road (hereafter referred to as Site A) and Queensway A38 tunnel – a road traffic tunnel
108 (hereafter referred to as Site B) and one heavily trafficked site in New Delhi: Mathura Road-
109 NH2 (hereafter referred to as Site C). Site A (UK roadside) is one of the major arterial roads in
110 south-west Birmingham with a dual carriageway with 32,000 vehicles/day and includes both
111 gasoline and diesel vehicles. RD samples were collected on both sides of the road close to the
112 traffic junction. Site B (UK road tunnel) is one of the major road tunnels in Birmingham and
113 runs in the North/South direction. The tunnel is naturally ventilated and typical driving speed
114 through the tunnel is 30 miles/hour with an estimated 89,000 vehicles travelling each day. Several
115 studies have been conducted at both these sites in the past (Smith and Harrison, 1996; Birmili et
116 al., 2006). Site C (India roadside) is one of the major roads in Delhi and falls on the National
117 Highway 2. An estimated 174,200 vehicles per day run on this road including intra-state and interstate

118 traffic. Public transport (including buses and auto-rickshaws) runs on compressed natural gas (CNG)
119 while personal vehicles run on both gasoline and diesel. Figure S1 (SI) provides the modal split at
120 the various sites (Azzi, 2012; DoT, 2014; Pant et al., 2015).

121

122 **2.2 Sampling**

123 A custom-built PM₁₀ dust sampler (as described in Amato et al., 2009) was used to collect the RD
124 samples. Samples were collected onto 47 mm PTFE and quartz fibre filters to enable analysis of
125 elements and organic species. Both PTFE (n=10) and quartz filter (n=10) samples were collected at
126 Sites A and C (UK and India roadside) while only PTFE filter samples (n=10) were collected at
127 Site B (UK tunnel). All the samples were collected after five days of dry weather in September
128 2012 (Site B) and June-July 2013 (Sites A and C- UK and India roadside). Individual samples were
129 collected over an area of 1m² for a period of 15 minutes and include samples from the kerbside as
130 well as middle of the road.

131

132 In spring 2014, soil, bulk RD, brake pad and tyre samples were collected in Birmingham while soil
133 and brake pad samples were collected in India in June 2014. The soil samples were collected from
134 the lawns of University of Birmingham and the CRRI campus remote from traffic in Birmingham
135 and New Delhi respectively while the brake pads were sourced from local garages. Only one brake
136 pad per city was analysed. Care was taken to ensure that the soil samples were collected from an
137 uncontaminated area. An additional set of PM₁₀ RD samples were also collected at Site A (UK
138 roadside). The soil and RD samples were dried, ground using a pestle and mortar and sieved using a
139 2 mm sieve; 0.5 g of each sample was then extracted and analyzed using the methods detailed in the
140 next section. Brake pad (UK and India) and tyre (UK) samples were frozen using liquid nitrogen
141 (N₂) and subsequently ground. The samples were passed through a 2 mm sieve and extracted and
142 analyzed.

143

144 **2.3 Chemical Analysis**

145 Details of the analytical procedures are provided in the SI.

146

147 **3. RESULTS AND DISCUSSION**

148 **3.1 Mass Loadings of PM₁₀ RD Fraction**

149 Average PM₁₀ mass loading on the road surface was found to be much higher in New Delhi
150 compared to Birmingham: $9.34 \pm 5.56 \text{ mg/m}^2$, $12.1 \pm 9.3 \text{ mg/m}^2$ and $72.9 \pm 24.3 \text{ mg/m}^2$ for sites
151 A (UK roadside), B (UK road tunnel) and C (India roadside) respectively. Mass loadings varied
152 between $3.78\text{-}21.8 \text{ mg/m}^2$, $3.01\text{-}36.1 \text{ mg/m}^2$ and $44\text{-}106 \text{ mg/m}^2$ at Sites A, B and C respectively.
153 In comparison, an average mass loading of 9 mg/m^2 has been reported for the city centre in
154 Barcelona (Spain) while mass loadings of $2.4\text{-}21.6 \text{ mg/m}^2$ have been reported for different site
155 types across Spain (Amato et al., 2009; Amato et al., 2013). Higher mass loading in Delhi can
156 be attributed to several factors including local meteorology (lower annual rainfall resulting in dry
157 conditions), prevalence of unpaved areas, larger proportion of construction activities, and during
158 the summer season, intrusion of dust from the Thar Desert. The site also had a higher proportion of
159 heavy duty vehicles (HDVs) which could be contributing to the higher mass loading since previous
160 studies have reported higher dust loadings for areas with HDVs (Abu Allaban et al., 2003).
161 Between the road and tunnel sites in Birmingham, elemental concentrations were consistently
162 higher at Site B (UK road tunnel) compared to Site A (UK roadside) with the exception of Al and
163 Si. Elemental concentrations are presented in Figure 1 and average mass loadings for elements and
164 organic species are presented in Table S1, SI. Si had the highest concentration among the crustal
165 elements, and Zn, Ti, Ba and Cu were the most abundant traffic-related elements at Sites A and B
166 whereas Zn, Mn, Ti and Ba were the most abundant in the case of Site C. Mass loadings for all the
167 species excluding Sb were found to be higher in the case of Site C (New Delhi) compared to the
168 Birmingham sites (A and B) while the concentrations ($\mu\text{g/g}$) of elements associated with traffic
169 emissions (i.e. Cu, Zn, Ba, Sb) were found to be highest in the road tunnel environment in

170 Birmingham (Site B). Previous studies in Houston (USA), by Spada et al. (2012) reported similar
171 results with concentrations of Cu, Zn, Pb, Ba, Sn and Sb found to be higher for bulk RD samples
172 collected in a tunnel compared to roads.

173

174 Concentrations of brake wear-related elements such as Sb and Ba in PM₁₀ RD were lower in the
175 case of New Delhi compared to Birmingham as well as other European sites while the Ca
176 concentration in Delhi (Site C) was found to be higher than other reported studies. Ca has been
177 used as a marker for crustal dust as well as construction activity (Chen et al., 2012; Pant and
178 Harrison, 2013). In Delhi, previous studies of bulk RD have reported a higher abundance of Ca-
179 rich dust attributed to the dust from the Thar Desert. The concentration of Sn was found to be
180 higher in Birmingham compared to New Delhi. Chemical composition of brakes and tyres is
181 expected to be different in UK and India, and the brake wear mode for elements such as Cu and Sb
182 has not been observed in elemental particle size distributions in New Delhi (Pant et al., 2015 in
183 preparation). For the organic species, concentrations were found to be close to the higher end of
184 the range of the previously reported values. It is important to note that elemental concentrations can
185 vary based on the type of tyres, brakes and pavement used at the site type. For example, summer
186 tyres create a lower dust loading (tyre dust) compared to non-studded and studded winter tyres and
187 both Birmingham and New Delhi use summer tyres all year round (Pant and Harrison, 2013). In
188 addition, concentrations of various crustal elements can vary based on soil type. For example, Fe
189 was found to be 7.4%, 7.3% and 3.1% for Sites A, B and C respectively while Al was found to be
190 8.7%, 7.2% and 3.98%. Chen et al. (2012) reported the abundance of Fe and Al as 2% and 3.5% in
191 China while Amato et al. (2011) reported Fe abundance of 4.6%, 5% and 5.8% and Al
192 abundance of 2.8%, 8% and 13.6% for Zurich, Barcelona and Girona respectively in the
193 PM₁₀ RD.

194

195 Based on Pearson correlation analysis (significance at $p < 0.01$), two major groups of elements
196 were identified in the PM_{10} RD in samples from sites A and B (UK roadside and road tunnel),
197 each with statistically significant correlations among the elements in the group. These were
198 crustal elements including Al, Si and Fe (group 1) and traffic-related elements such as Cu, Zn,
199 Sb, Ba, Sn and Mn (group 2). In Beijing (China), Chen et al. (2012) reported four groups: crustal
200 elements (Al, Fe, Ca, Ti), salt (K, Mg, Na), sulphur and other trace elements. At site B, Ni was not
201 found to be correlated to either group. Ni is typically attributed to industrial and fuel oil
202 combustion emissions and has also been attributed to engine metal wear and tear (Hays et al.,
203 2011). Ca showed a high correlation with the traffic-associated elements at Site A. At site C (India
204 roadside), a high correlation was observed between Zn, Cu and Ba but was weaker for Sn, and
205 Sb was not found to be correlated to other traffic-associated elements. Interestingly, while Ti was
206 found to be correlated to traffic-related elements at Sites A and B, a stronger correlation was
207 observed for Ti with Al and Si at Site C. The traffic-related elements could not be subdivided into
208 tyre/brake wear categories based on correlation analysis.

209

210 High molecular weight PAHs showed very high correlation with one another, attributable to
211 common sources and deposition pathways.

212

213

214 **3.2 Bulk Samples**

215 **3.2.1 Soil and road dust**

216 In addition to the PM_{10} RD, bulk soil and road dust samples were analysed to understand the
217 similarities (using Pearson Correlation) in the chemical compositions across the fractions both in
218 UK and India (Figure S3, SI). In the UK, bulk RD (< 2 mm) composition was found to be
219 correlated with bulk soil (< 2 mm) as well as PM_{10} RD while in India, bulk RD (< 2 mm) was found
220 to be correlated with bulk soil (< 2 mm) but the correlation was not significant for PM_{10} RD.

221

222 To understand the contribution of traffic emissions to the RD composition (Figure S3, SI), a
223 roadside enrichment factor was calculated based on equation 1 (Amato et al., 2011).

$$\text{Roadside Enrichment} = \left(\frac{C_r - C_s}{C_r} \right) * 100 \quad \text{Eq (1)}$$

224

225 Where

226 C_r is concentration at roadside

227 C_s is the concentration in soil

228

229 Highest enrichment was observed for Ti, Sb and Cr in Birmingham with enrichment of more than
230 70% while Cu, Zn and Sn showed enrichment of more than 50%. V was not found to be enriched in
231 Birmingham while in Delhi, Sb, Ba and Ni were not found to be enriched. In Delhi, Zn was the only
232 element with an enrichment of > 70% although elements such as Ti, Cu and Sn showed enrichment
233 of more than 30%.

234

235 **3.2.2 Brake pad and tyre composition**

236 Individual samples of four-wheeler passenger vehicle brake pads from India and UK (one from each
237 country) were analysed to assess the chemical composition. This is particularly relevant in the case
238 of India since there is a need to identify an elemental tracer for brake wear estimation. Pant et al.
239 (2015 in preparation) have highlighted that particle size distributions measured in New Delhi do not
240 show the brake wear modes for elements such as Cu, Sb and Ba which are observed in European
241 samples (e.g. Gietl et al., 2010). However, it is important to note that the sample size is limited, and
242 cannot be used to draw significant conclusions. The differences highlighted using this sample need
243 to be investigated further, and further tests are planned for the next phase of the analysis.

244

245 The bulk brake pad dust was found to be rich in Ti (0.12%), Sb (1.12%) and Cu (0.49%) in India,
246 while in the UK, Ba (0.12%) and Sb (0.16%) were the most abundant elements among those
247 analyzed (Figure S4, SI). This is consistent with previous studies in which differences in the
248 composition of brake pads have been reported between different types (e.g. NAO vs low-metallic)
249 (Sanders et al., 2003) and between countries (Kukutschova et al., 2010; Hulskotte et al., 2014).
250 However, it is important that only one sample was analysed per city, and may not be representative
251 of the range of brake pads in use. Future work is underway to further evaluate varying brake pad
252 formulations across geographic regions to understand the chemical composition of brake pads as
253 well as brake wear particles.

254

255 In the case of the tyre sample, Zn was found to be the most abundant element (Figure S5, SI).

256

257 **3.3 Enrichment Factor Analysis for the PM₁₀ RD**

258 In order to further understand the sources of the elements (crustal vs. anthropogenic), enrichment
259 factors (EFs) were calculated for PM₁₀ RD based on continental crust concentrations using Al as the
260 reference element (Taylor and McLennan, 1995) (Figure).

261

$$Enrichment\ Factor\ (X) = \frac{\left\{ \frac{Concentration(X)}{Concentration(Reference)} \right\}_{sample}}{\left\{ \frac{Concentration(X)}{Concentration(Reference)} \right\}_{crustal}} \quad Eq\ (2)$$

262

263 Use of Al as the reference element was based on previous studies in Birmingham that identified a
264 minimal traffic-associated increment for Al (Birmili et al., 2006). Corresponding to the correlations
265 observed in the dataset, highest EFs were observed for Cu, Zn, Sb, Ni and Sn. Across the three
266 sites, Site A (UK roadside) was found to have the lowest enrichment factors. Other studies have

267 also reported high EFs for Ba, Cr, Cu and Sb in airborne particles (Birmili et al., 2006; Dongarra et
268 al., 2009; Oliveira et al., 2011). At the UK sites (Sites A and B), Si, Fe and Ca showed no
269 enrichment with respect to the crustal concentrations; however, Ca was observed to have a high EF
270 at Site C (India roadside). This could be linked with sources associated with construction activity
271 and desert dust which is rich in Ca.

272

273 **3.4 Elemental Ratios in the PM₁₀ RD**

274 Ratios between different elements can be used to distinguish between different sources. Various
275 authors have proposed characteristic Cu/Sb ratios for brake wear particles including 4.6 ± 2.3 in
276 USA (Sternbeck et al., 2002) and 7.0 ± 1.9 in Europe (Amato et al., 2009). In comparison, this ratio
277 is 125 for the upper continental crust indicating relatively much higher concentration of Cu in the
278 crust compared to Sb. It is important to note that the Cu/Sb ratio can vary across geographical
279 regions based on the type of brake or the elemental content of the brakes, which differs among
280 manufacturers (Pant and Harrison, 2013; Hulskotte et al., 2014). For example, non-asbestos
281 organic (NAO) brake lining material typically has a Cu/Sb ratio of 11.5 (Iijima et al., 2007) whereas
282 the ratio is 1.33 for semi-metallic brake pad material (Schauer et al., 2006). Recently, Hulskotte et
283 al. (2014) have reported an average ratio value of 4.6 for brake pad material in the Netherlands.

284

285 In the current study, the overall Cu/Sb ratio was observed to be 4.95 ± 0.50 for Site A, 7.50 ± 0.79 for
286 Site B and 11.5 ± 0.82 for Site C. The ratio values for the Birmingham sites fall within the
287 proposed range of characteristic ratios, consistent with a contribution of brake wear particles to
288 the PM₁₀ RD, and are similar to the Cu/Sb ratio of 7.5 reported for low-metallic brake pads in
289 the USA (Schauer et al., 2006). However, a higher ratio (11.5) was observed in the case of Site
290 C (India) which could be due to either other sources of Cu in the area, or due to differences in the

291 composition of brakes. For the bulk RD, the Cu/Sb ratio was 4.80 ± 0.40 for Delhi and 6.22 ± 3.16 for
292 Birmingham.

293
294 Further discussion on elemental ratios is presented in SI.

295

296 **3.5 Chemical Source Profiles of the PM₁₀ RD**

297 Source profiles (weight % of species with respect to the mass of PM) were generated for the
298 road traffic sites in Birmingham (Sites A and B) and New Delhi (Site C) and appear in Table 1
299 (see Table S2 in SI for other molecular markers).

300

301 Previously reported source profiles for RD and various non-exhaust emission sources were
302 compared against data obtained at Sites A-C in order to achieve a qualitative understanding of
303 source contributions (Figure 3).

304

305 The source profiles were extracted from the USEPA SPECIATE 4.3 database and other published
306 literature. In the case of elements, profiles included paved RD [highway (profile number 3565),
307 tunnel (profile number 4112410)] and RD (Mexico) (profile number 4106), from the SPECIATE
308 database, paved RD from India (Sethi and Patil, 2008), Hong Kong (Ho et al., 2003), Barcelona
309 (Amato et al., 2009) and Beijing (China) (Chen et al., 2012). Concentrations of various elements
310 were broadly similar to other compositional profiles of PM₁₀ RD, and in most cases, PM₁₀ RD
311 concentrations in the tunnel were higher than roadside samples. A previously reported
312 compositional profile of PM₁₀ RD for Delhi (Sethi and Patil, 2008) was found to be different from
313 the current profile. Concentrations of several elements including Si, Al, Ca and Ba were much
314 lower in the case of the profile from Sethi and Patil (2008), while concentrations were lower for the
315 current profile for Sn, OC and EC and comparable for Ti, Fe and Zn. The Si/Al ratio was found to
316 be lower than the UCC value of 3.9 for the profile from Sethi and Patil (2008) while the Cu/Sb ratio

317 at 21.5 was much higher compared to the profile generated in this study. The OC/EC ratios were,
318 however, comparable (3.6 for current study, 3.3 for Sethi and Patil, 2008).

319

320 **3.6 Source Apportionment**

321 Recently, a novel method was proposed for the estimation of source contributions of non-exhaust
322 sources using Ba, Zn and Si as source markers for brake dust, tyre dust and crustal dust respectively
323 (Harrison et al., 2012). The factors reported by Harrison et al. (2012) (91 for Ba, 50 for Zn and 3.6
324 for Si) were used to convert masses of these elements to mass contributions of brake dust, tyre dust
325 and crustal dust respectively to the total PM₁₀ mass (Table 2 and Figure S6, SI). It is important to
326 note that Zn is emitted from various sources which include engine emissions and brake dust, and it
327 is possible that using Zn as a tyre wear tracer results in an over-estimation of the tyre dust
328 emissions. In Asia, Zn has also been associated with 2-stroke gasoline engine emissions and
329 emissions from the galvanization industry (Begum et al., 2011). Further, preliminary analysis of
330 brake pad samples from New Delhi (Section 3.2) reveals that levels of Ba are quite low in the brake
331 pads in New Delhi. In the case of New Delhi, Fe was used as the source marker for crustal dust
332 due to its high abundance in crustal material. In addition, a factor of 1.35 was used for EC to
333 estimate the contribution of vehicular exhaust (Pio et al. 2011).

334

335 Application of these factors accounted for a high proportion of PM₁₀ RD mass in the UK samples
336 (99.3% for Site A and 105% for Site B), with the highest contribution from crustal dust. For site B
337 (road tunnel), the contribution of tyre wear was calculated to be 15.6% which might be an over-
338 estimation since Zn is emitted from other sources as well. The contribution of brake wear was
339 also the highest for Site B. In the case of Delhi, 79.2% of the total mass is accounted for by this
340 method but if Si is used as the source marker for crustal dust, only 51% of the total PM₁₀ mass is
341 estimated. This indicates that this estimation method cannot be used universally, and it is important

342 to adjust the markers/factors according to the local soil characteristics. Other sources which
343 could make important contributions in the case of New Delhi are construction activity and
344 deposition from other sources which are currently not included in the estimation. Vehicle exhaust
345 was found to contribute a very small percentage to the total PM₁₀ mass which is plausible since
346 most of the vehicle exhaust particles are in the smaller size fraction and are less likely to deposit.

347

348 Chen et al. (2012) estimated soil dust, construction-related particles, vehicle exhaust, particle
349 deposition and coal burning-associated particles as the key sources for RD in Beijing. Further
350 evaluation of brake wear, tyre wear, crustal dust and vehicle exhaust by Amato et al. (2013)
351 demonstrated these emission sources contributed 27%, 16%, 37% and 20%, respectively of the total
352 PM mass in Barcelona and 39%, 41%, 12%, 8%, respectively, in Utrecht, the Netherlands.

353

354 To compare the attribution of crustal dust using Si against an independent estimate, crustal dust
355 mass was reconstructed using the concentrations of Al, Si, Fe, Ca and Ti using the equation (3)
356 (Chan et al., 1997). Results indicated good correlation between measured and reconstructed mass in
357 all cases with $r^2 > 0.85$ in all cases. However, as with the source apportionment, the amount of mass
358 apportioned as crustal dust was observed to be low at Site C (Indian roadside). This may be related
359 to the varied soil composition at different locations. Hence, it can be concluded that it is important
360 to analyse uncontaminated soil samples (bulk and PM₁₀ fraction where possible) to determine the
361 local chemical composition which can be then be used to adjust the crustal dust factor.

362

$$\text{Crustal Dust} = 1.16 (1.90 \text{ Al} + 2.15 \text{ Si} + 1.41 \text{ Ca} + 1.67 \text{ Ti} + 2.09 \text{ Fe}) \quad \text{Eq (3)}$$

363

364

365 3.7 Oxidative Potential

366 While it is well known that PM exposure is a contributor to adverse cardiopulmonary health, the
367 hazards posed by PM emitted from different sources may vary dependent upon the characteristics of
368 the emitted particles. Particle toxicity is thought to vary based on chemical composition. Specific
369 constituents of PM such as redox active trace metals (e.g. Cu, Zn, Ni) and PAHs may be
370 particularly harmful as they can lead to the generation of reactive oxygen species (ROS) and
371 subsequent inflammation and oxidative stress (Kelly, 2003; Ayres et al., 2008). Multiple PM
372 constituents have been shown to exert Oxidative Potential, which may lead to oxidative stress in
373 biological systems, and Oxidative Potential has been suggested as a biologically informative
374 expression for response assessment (Weichenthal et al 2013). Recent studies of PM have
375 highlighted the high Oxidative Potential of non-exhaust road traffic emissions (Godri et al., 2011;
376 Kelly et al, 2011; Yanosky et al., 2012; Janssen et al 2014; Amato et al., 2014). The oxidative
377 potential (OP) of a subset of bulk and PM₁₀ RD samples was measured in the current study.

378

379 Both OP associated with ascorbate depletion per unit PM mass (OP^{AA}) and OP associated with
380 glutathione depletion per unit PM mass (OP^{GSH}) were found to be higher for the bulk RD compared
381 to the PM₁₀ RD (Figure 4). OP^{AA} and OP^{GSH} were not found to be correlated with each other,
382 suggesting that distinct chemical species were contributing to the depletion of each antioxidant.
383 This result supports previous studies that have shown that glutathione and ascorbate respond to
384 different chemical components of PM (Godri et al., 2010a; Kelly et al., 2011).

385

386 Published studies have shown significant associations between OP and transition metals in ambient
387 PM₁₀ samples, specifically the recurring correlation between OP^{GSH} and Fe (Godri et al., 2010a;
388 Kelly et al., 2011; Godri et al., 2011). Yang et al. (2014) and Janssen et al. (2014) also found
389 significant positive associations between Fe, Cu and OP^{AA} for PM_{2.5} while Godri et al. (2010b)

390 found positive association between Fe, Pb and OP^{AA} for PM_{10} . Fe and Cu are traffic related redox
391 active transition metals thought to drive antioxidant depletion (Kelly et al. 2011). Linear regression
392 analysis was performed to understand the contribution of individual elements to the oxidative
393 potential. For OP^{GSH} , no significant relationships were observed for either PM_{10} or bulk RD. On the
394 other hand, OP^{AA} showed some significant relationships for both fractions ($p < 0.05$). In terms of
395 individual elements, Cu was the only significant element ($r^2 = 0.995$, $p = 0.044$) for bulk RD (Figure
396 S7, SI) while for PM_{10} , Ti ($r^2 = 0.54$), Mn ($r^2 = 0.52$), S ($r^2 = 0.57$), As ($r^2 = 0.55$), Cu ($r^2 = 0.61$) and
397 V ($r^2 = 0.57$) were found to be significant for $p < 0.05$ (Figure S8, SI). Results of the regression
398 analysis are summarized in Table 3. Cu is typically associated with non-exhaust traffic emissions
399 (Amato et al., 2009; Pant and Harrison, 2013), and was found to be associated with OP^{AA} in both
400 fractions of RD. On the other hand, Mn, V and Ti can be associated with traffic as well as other
401 sources such as fuel oil combustion. It is useful to note that at Site A (UK roadside), high
402 correlations were observed between elements. Interestingly, Fe and Zn were not found to be a
403 significant in the regression models for either of the RD fractions.

404

405 **4. CONCLUSIONS**

406 Non-exhaust particles constitute an important source of PM emissions in urban areas, and the
407 chemical composition as well as contribution can vary from region to region. Since there are a
408 number of contributions to non-exhaust emissions, it is often difficult to estimate the contribution of
409 different sources to RD. In this study, an attempt was made to undertake detailed characterization
410 of the PM_{10} RD and the empirical method proposed for estimation of the contribution of non-
411 exhaust sources was applied with some success to the dataset.

412

413 Chemical source profiles generated for the PM_{10} RD at sites in Birmingham and New Delhi
414 correspond well with previously reported RD profiles, although differences were observed between

415 element concentrations in UK and India. This can be attributed to the difference in soil composition
416 as well as the chemical composition of tyre and brake pads. Comparison carried out between
417 elemental and PAH ratios for RD PM and airborne PM reveal that the ratios can be quite similar
418 between RD and ambient PM, particularly in the case of coarse PM.

419 The empirical factors generated by Harrison et al. (2012) relating tracer elements to particle mass
420 were found to be able to account well for source contributions in the area where they were
421 empirically determined (i.e. high traffic areas in UK), but these factors were not able to apportion
422 all the PM mass for the Indian site. Low mass closure in New Delhi is perhaps due to the different
423 sources that influence the dust concentrations in New Delhi, as well as due to differences in the
424 overall concentrations of different elements, and perhaps their sources as well. It is therefore
425 necessary to adjust the factors before using them for analysis in other regions. The estimated
426 contribution of brake wear was very similar between Sites A (UK roadside) and C (Indian
427 roadside), both of which experience heavy-trafficked roads with stop-and-go traffic flows.

428

429 The dataset generated in this study will be useful in receptor modelling studies. It is critical to note
430 that the contribution of non-exhaust emissions to ambient PM concentration varies based on site
431 characteristics, and data generated with pilot studies in specific areas/site types cannot be used to
432 generalize the role and quantitative contribution of non-exhaust emissions to ambient air quality.

433

434 A preliminary assessment of the Oxidative Potential of RD (bulk and PM₁₀) has been made in this
435 study. Cu, often used as a traffic emissions marker, was found to be significant in both cases.
436 However, it is important to note that multiple metals are redox active, and can lead to antioxidant
437 depletion (Godri et al., 2011), and it is possible that the complex interactions are not completely
438 expressed through this limited dataset. Further analysis is required to understand the drivers for the

439 differences across sample fractions (PM₁₀ vs. bulk) and to quantify the health risks associated with
440 exposure to non-exhaust emissions vis-à-vis other emission types.

441

442 **ACKNOWLEDGEMENTS**

443 The authors would like to thank Rees Jeffreys Road Fund and Simon Wolff Charitable Foundation
444 for financial support and Pallavi Pant gratefully acknowledges financial support from the University
445 of Birmingham. The authors would also like to thank Amey for access to the Queensway Tunnel in
446 Birmingham and the AMC facility at University of Birmingham for access to the XRF.

447

448 **REFERENCES**

- 449
- 450 Aatmeeyata, K. D. S. and Sharma, M., 2010. Polycyclic aromatic hydrocarbons, elemental and
451 organic carbon emissions from tire-wear. *Science of the Total Environment* 408, 4563-4568.
- 452
- 453 Abu-Allaban, M., Gillies, J.A., Gertler, A.W., Clayton, R. and Proffitt, D., 2003. Tailpipe,
454 resuspended road dust, and brake wear emission factors from on-road vehicles. *Atmospheric*
455 *Environment* 37, 5283-5293.
- 456
- 457 Adachi, K. and Tainosho, Y., 2004. Characterization of heavy metal particles embedded in tire dust.
458 *Environment International* 30, 1009-1017.
- 459
- 460 Agarwal, T. (2009). Concentration level, pattern and toxic potential of PAHs in traffic soil of Delhi,
461 India. *Journal of Hazardous Materials* 171, 894-900.
- 462
- 463 Amato, F., Pandolfi, M., Viana, M., Querol, X., Alastuey, A. and Moreno, T., 2009. Spatial and
464 chemical patterns of PM10 in road dust deposited in urban environment. *Atmospheric Environment*
465 43, 1650-1659.
- 466
- 467 Amato, F., Pandolfi, M., Moreno, T., Furger, M., Pey, J., Alastuey, A., Bukowiecki, N., Prevot,
468 A.S.H., Baltensperger, U. and Querol, X., 2011. Sources and variability of inhalable road dust
469 particles in three European cities. *Atmospheric Environment* 45, 6777-6787.
- 470
- 471 Amato, F., Pandolfi, M., Alastuey, A., Lozano, A., Gonzalez, J.C., Querol, X., 2013. Impact of
472 traffic intensity and pavement aggregate size on road dust particles loading. *Atmospheric*
473 *Environment* 77, 711-717.
- 474
- 475 Amato, F., Cassee, F.R., Denier van der Gon, H.A.C., Gehrig, R., Gustaffson, M., Hafner, W.,
476 Harrison, R.M., Jozwicka, M., Kelly, F.J., Moreno, T., Prevot, A.H., Schaap, M., Sunyer, J. and
477 Querol, X., 2014. Urban air quality: The challenge of traffic non-exhaust emissions. *Journal of*
478 *Hazardous Materials* 275, 31-36.
- 479
- 480 Ayres, J.G., Borm, P., Cassee, F.R., Castranova, V., Donaldson, K., Ghio, A., Harrison, R.M.,
481 Hider, R., Kelly, F., Ingeborg, M.K., Marano, F., Maynard, R.L., Mudway, I., Nel, A., Sioutas, C.,
482 Smith, S., Baeza-Squiban, A., Cho, A., Duggan, S. and Froines, J., 2008. Evaluating the toxicity of
483 airborne particulate matter and nanoparticles by measuring oxidatve stress potential- a workshop
484 report and consensus statement. *Inhalation Toxicology* 20, 75-99.
- 485
- 486 Azzi, S. (salim.azzi@amey.co.uk) (18 September 2012). Re: Air sampling in the Queensway road
487 tunnel, Birmingham. E-mail to P. Pant (pxp024@bham.ac.uk)
- 488
- 489 Begum, B.A., Biswas, S.K. and Hopke, P.K., 2011. Key issues in controlling air pollutants in
490 Dhaka, Bangladesh. *Atmospheric Environment* 45, 7705-7713.
- 491
- 492 Birmili, W., Allen, A. G., Bary, F. and Harrison, R.M., 2006. Trace Metal Concentrations and
493 Water Solubility in Size-Fractionated Atmospheric Particles and Influence of Road Traffic.
494 *Environmental Science and Technology* 40, 1144-1153.
- 495
- 496 Bukowiecki, N., Lienemann, P., Hill, M., Furger, M., Richard, A., Amato, F., Prevot, A.S.H.,
497 Baltensperger, U., Buchmann, B. and Gehrig, R., 2010. PM10 emission factors for non-exhaust
498 particles generated by road traffic in an urban street canyon and along a freeway in Switzerland.
499 *Atmospheric Environment* 44, 2330-2340.

500 Camatani, M., Crosta, G. F., Dolukhanyan, T., Sung, C., Giuliani, G., Corbetta, G. M., Cencetti, S.
501 and Regazzoni, C., 2001. Microcharacterization and identification of tire debris in heterogenous
502 laboratory and environmental specimens. *Materials Characterization* 46, 271-283.
503

504 Chan, Y.C., Simpson, R.W., Mctainsh, G.H., Vowles, P.D., Cohen, D.D. and Bailey, G.M., 1997.
505 Characterisation of chemical species in PM2.5 and PM10 aerosols in Brisbane, Australia.
506 *Atmospheric Environment* 31, 3773- 3785.
507

508 Chen, J., Wang, W., Liu, H., Ren, L., 2012. Determination of road dust loadings and chemical
509 characteristics using resuspension. *Environmental Monitoring and Assessment* 184, 1693-1709.
510

511 Colville, R.N., Hutchinson, E.J., Mindell, J.S. and Warren, R.F., 2001. The transport sector as a
512 source of air pollution. *Atmospheric Environment* 35, 1537-1565.

513 Councell, T. B., Duckenfield, K. U., Landa, E. R. and Callender, E., 2004. Tire wear particles as a
514 source of Zn to the environment. *Environmental Science and Technology*, 38, 4206-4214.
515

516 Denier van der Gon, H., Gerlofs-Nijland, M.E., Gehrig, R., Gustafsson, M., Janssen, N., Harrison,
517 R.M., Hulskotte, J., Johansson, C., Jozwicka, M., Keuken, M., Krijgsheld, K., Ntziachristos, L.,
518 Riediker, M. and Cassee, F.R., 2013. The Policy Relevance of Wear Emissions from Road
519 Transport, Now and in the Future- An International Workshop Report and Consensus Statement.
520 *Journal of the Air and Waste Management Association* 63, 136-149.
521

522 Department of Transport (2014) Traffic Counts Birmingham. Accessed at
523 <http://www.dft.gov.uk/traffic-counts/cp.php?la=Birmingham#81577> .
524

525 Dongarra, G., Manno, E. and Varrica, D., 2009. Possible markers of traffic-related emissions.
526 *Environmental Monitoring and Assessment* 154, 117-125.
527

528 Duong, T. T. and Lee, B. K., 2011. Determining contamination level of heavy metals in road dust
529 from busy traffic areas with different characteristics. *Journal of Environmental Management* 92(3),
530 554-562.
531

532 Faiz, Y., Tufail, M., Tayyeb Javed, M., Chaudry, M.M. and Naila-Siddique (2009). Road dust
533 pollution of Cd, Ni, Pb and Zn along Islamabad Expressway, Pakistan. *Microchemical Journal* 92,
534 186-192.
535

536 Garg, B.D., Cadle, S.H., Mulawa, P.A., Groblicki, P.J., Laroo, C. and Parr, G.A., 2000. Brake Wear
537 Particulate Matter Emissions. *Environmental Science and Technology* 34, 4463-4469.
538

539 Gietl, J.K., Lawrence, R., Thorpe, A.J., Harrison, R.M., 2010. Identification of brake wear particles
540 and derivation of a quantitative tracer for brake dust at a major road. *Atmospheric Environment* 44 ,
541 141-146.
542

543 Godri, K.J., Duggan, S.T., Fuller, G.W., Baker, T., Green, D., Kelly, F.J. and Mudway, I.S., 2010a.
544 Particulate oxidative burden associated with firework activity. *Environmental Science and*
545 *Technology* 44, 8295-8301.
546

547 Godri, K.J., Duggan, S.T., Fuller, G.W., Baker, T., Green, D., Kelly, F.J. and Mudway, I.S., 2010b.
548 Particulate matter oxidative potential from waste transfer station activity. *Environmental Health*
549 *Perspectives* 118, 493-498.
550

551
552 Godri, K.J., Harrison, R.M., Evans, T., Baker, T., Dunster, C., Mudway, I.S. and Kelly, F.J., 2011.
553 Increased oxidative burden associated with traffic component of ambient particulate matter at
554 roadside and urban background school sites in London. *PloS ONE* 6, e21961,
555 doi:10.1371/journal.pone.0021961.
556
557 Gunawardana, C., Goonetilleke, A., Egodawatta, P., Dawes, L. and Kokot, S., 2011. Source
558 characterisation of road dust based on chemical and mineralogical composition. *Chemosphere* 87,
559 163-170.
560
561 Han, L., Zhuang, G., Cheng, S., Wang, Y. and Li, J., 2007. Characteristics of re-suspended road
562 dust and its impact on the atmospheric environment in Beijing. *Atmospheric Environment* 41, 7485-
563 7499.
564
565 Han, B., Bai, Z., Guo, G., Wang, F., Li, F., Liu, Q., Ji, Y., Li, X. and Hu, Y., 2009. Characterization
566 of PM(10) fraction of road dust for polycyclic aromatic hydrocarbons (PAHs) from Anshan, China.
567 *Journal of Hazardous materials* 170, 934-940.
568
569 Harrison, R.M., Jones, A.M., Gietl, J., Yin, J. and Green, D.C., 2012. Estimation of the contributions
570 of brake dust, tire wear, and resuspension to nonexhaust traffic particles derived from atmospheric
571 measurements. *Environmental Science and Technology* 46, 6523-6529.
572
573 Hays, M.D., Cho, S.-H., Baldauf, R., Schauer, J.J. and Shafer, M., 2011. Particle size distributions
574 of metal and non-metal elements in an urban near-highway environment. *Atmospheric Environment*
575 45, 925-934.
576
577 Ho, K.F., Lee, S.C., Chow, J.C., Watson, J.G., 2003. Characterization of PM₁₀ and PM_{2.5} source
578 profiles for fugitive dust in Hong Kong. *Atmospheric Environment* 37, 1023-1032.
579
580 Hulskotte, J.H.J, Roskam, G.D. and Denier Van Der Gon, H.A.C., 2014. Elemental composition of
581 current automotive braking materials and derived air emission factors. *Atmospheric Environment*
582 99, 436-445.
583
584 Iijima, A., Sato, K., Yano, K., Tago, H., Kato, M., Kimura, H. and Furuta, N., 2007. Particle size
585 and composition distribution analysis of automotive brake abrasion dusts for the evaluation of
586 antimony sources of airborne particulate matter. *Atmospheric Environment* 41, 4908-4919.
587
588 Janssen, N.A.H., Yang, A., Strak, M., Steenhof, M., Hellack, B., Gerlofs-Nijland, M.E., Kuhlbusch,
589 T., Kelly, F., Harrison, R., Brunekreef, B., Hoek, G. and Cassee, F., 2014. Oxidative potential of
590 particulate matter collected at sites with different source characteristics. *Science of the Total*
591 *Environment* 472, 572-581.
592
593 Kelly, F.J., 2003. Oxidative stress: its role in air pollution and adverse health effects. *Occupational*
594 *and Environmental Medicine* 60, 612- 616.
595
596 Kelly, F., Anderson, H.R., Armstrong, B., Atkinson, R., Barratt, B., Beevers, S., Derwent, D.,
597 Green, D., Mudway, I. and Wilkinson, P. Part 2: Analysis of the oxidative potential of particulate
598 matter. In: *The Impact of the Congestion Charging Scheme on Air Quality in London*. Boston, MA:
599 Research Report 155. Health Effects Institute; 2011.
600

601 Kukutschova, J., Roubicek, V., Maslan, M., Jancik, D., Slovak, V., Malachova, K., Pavlickova, Z.
602 and Filip, P., 2010). Wear performance and wear debris of semimetallic automotive brake materials.
603 Wear 268, 86-93.
604

605 Kukutschova, J., Moravec, P., Tomasek, V., Matejka, V., Smolik, J., Schwarz, J., Seidlerova, J.,
606 Safarova, K. and Filip, P., 2011. On airborne nano/micro-sized wear particles released from low-
607 metallic automotive brakes. Environmental Pollution, 159, 998- 1006.
608

609 Kwak, J., Kim, H., Lee, J., Lee, S., 2013. Characterization of non-exhaust coarse and fine particles
610 from on-road driving and laboratory measurements. Science of the Total Environment 458-460,
611 273-282.
612

613 Kwon, E.E. and Castaldi, M.J., 2012. Mechanistic understanding of polycyclic aromatic
614 hydrocarbons (PAHs) from the thermal degradation of tires under various oxygen concentration
615 atmospheres. Environmental Science and Technology 46, 12921- 6.
616

617 Liu, M., Cheng, S.B., Ou, D.N., Hou, L.J., Gao, L., Wang, L.L., Xie, Y.S., Yang, Y., Xu, S.Y.,
618 2007. Characterization, identification of road dust PAHs in central Shanghai areas, China.
619 Atmospheric Environment 41, 8785-8795.
620

621 Lough, G.C., Schauer, J.J., Park, J., Shafer, M.M., Deminter, J.T., Weinstein, J.P., 2005. Emissions
622 of metals associated with motor vehicle roadways. Environmental Science and Technology 39, 826-
623 836.
624

625 Luhana, L., Sokhi, R., Warner, L., Mao, H., Boulter, P., McCrae, I., Wright, J., Osborn, D., 2004.
626 Characterisation of Exhaust Particulate Emissions from Road Vehicles (PARTICULATES),
627 Deliverable 8: Measurement of non-exhaust particulate matter. Contract Number 2000-RD.11091,
628 European Commission – DG TrEn, 5th Framework Programme Competitive and Sustainable
629 Growth Sustainable Mobility and Intermodality.
630

631 Luo, X.S., Yu, S., Zhu, Y. G. and Li, X. D., 2011. Trace metal contamination in urban soils of
632 China. Science of the Total Environment 421-422, 17-30.
633

634 Martuzevicius, D., Kliucininkas, L., Prasauskas, T., Krugly, E., Kauneliene, V. and Strandberg, B.,
635 2011. Resuspension of particulate matter and PAHs from street dust. Atmospheric Environment, 45,
636 310-317.
637

638 Milani, M., Pucillo, F.P., Ballerini, M., Camatani, M., Gualtieri, M. and Martino, S., 2004. First
639 evidence of tyre debris characterization at the nanoscale by focused ion beam. Materials
640 Characterization 52, 283-288.
641

642 Morillo, C., Romero, A.S., Maqueda, C., Madrid, L., Ajmone-Marsan, F., Grcman, H., Davidson,
643 C.M., Hursthouse, A.S. and Villaverde, J., 2007). Soil pollution by PAHs in urban soils: a
644 comparison of three European cities. Journal of Environmental Monitoring 9, 1001-1008.
645

646 Oliveira, C., Martins, N., Tavares, J., Pio, C., Cerqueira, M., Matos, M., Silva, H., Oliveira, C. and
647 Camoes, F., 2011. Size distribution of polycyclic aromatic hydrocarbons in a roadway tunnel in
648 Lisbon, Portugal. Chemosphere 83, 1588-1596.
649

650 Omstedt, G., Bringfelt, B., Johansson, C., 2005. A model for vehicle-induced non-tailpipe
651 emissions of particles along Swedish roads. Atmospheric Environment 39, 6088-6097.
652

653 Pant, P. and Harrison, R.M., 2013. Estimation of the contribution of road traffic emissions to
654 particulate matter concentrations from field measurements: A review. *Atmospheric Environment*
655 77, 78-97.
656
657 Pant, P., Shukla, A., Kohl, S.D., Chow, J.C., Watson, J.G. and Harrison, R.M., 2015a.
658 Characterization of ambient PM_{2.5} at a pollution hotspot in New Delhi, India and inference of
659 sources. *Atmospheric Environment* 109, 178.-189.
660
661 Pant, P., Baker, S.J., Guttikunda, S., Goel, A., Shukla, A. and Harrison, R.M., 2015b. Analysis of
662 size-segregated winter season aerosol data from New Delhi, India. Manuscript in preparation.
663
664 Peltier, R. E., Cromar, K. R., Ma, Y., Fan, Z.-H. and Lippmann, M., 2011. Spatial and seasonal
665 distribution of aerosol chemical components in New York City: (2) Road dust and other tracers of
666 traffic-generated air pollution. *Journal of Exposure Science and Environmental Epidemiology* 21,
667 484-494.
668
669 Pio, C., Cerqueira, M., Harrison, R.M., Nunes, T., Mirante, F., Alves, C., Oliveira, C., Sanchez de
670 la Campa, A., Artinano, B. and Matos, M., 2011. OC/EC ratio observations in Europe: Re-thinking
671 the approach for apportionment between primary and secondary organic carbon. *Atmospheric*
672 *Environment* 45, 6121-6132.
673
674 Prichard, H.M. and Fisher, P.C., 2012. Identification of Platinum and Palladium Particles Emitted
675 from Vehicles and Dispersed into the Surface Environment. *Environmental Science and*
676 *Technology* 46, 3149-3154.
677
678 Rogge, W.F., Hildemann, L.M., Mazurek, M.A., Cass, G.R. and Simoneit, B. R. T., 1993. Sources
679 of fine organic aerosol. 3. Road dust, tire debris, and organometallic brake lining dust: roads as
680 sources and sinks. *Environmental Science and Technology* 27(9), 1892-1904.
681
682 Sanders, P.G., Xu, N., Dalka, T.M. and Maricq, M.M., 2003. Airborne brake wear debris: Size
683 distributions, composition, and a comparison of dynamometer and vehicle tests. *Environmental*
684 *Science and Technology* 37(4), 4060-4069.
685
686 Schauer, J.J., Lough, G.C., Shafer, M.M., Christensen, W.C., Arndt, M.F., DeMinter, J.T. and Park,
687 J.-S., 2006. Characterization of Emissions of Metals Emitted from Motor Vehicles. HEI Research
688 Report 133. Health Effects Institute, Boston, MA.
689
690 Sethi, V. and Patil, R.S., 2008. Development of Air Pollution Source Profiles- Stationary Sources-
691 Volumes 1 and 2. Available at
692 http://www.cpcb.nic.in/Source_Emission_%20Profiles_NVS_Volume%20One.pdf [Accessed April
693 21, 2011].
694
695 Smith, D.J.T. and Harrison, R.M., 1996. Concentrations, trends and vehicle source profile of
696 polynuclear aromatic hydrocarbons in the U.K. atmosphere. *Atmospheric Environment* 30, 2513-
697 2525.
698
699 Spada, N., Bozlaker, A. and Chellam, S., 2012. Multi-elemental characterization of tunnel and road
700 dusts in Houston, Texas using dynamic reaction cell-quadrupole-inductively coupled plasma-mass
701 spectrometry: Evidence for the release of platinum group and anthropogenic metals from motor
702 vehicles. *Analytica Chimica Acta* 735, 1-8.
703

704 Sternbeck, J., Sjödin, Å. and Andréasson, K., 2002. Metal emissions from road traffic and the
705 influence of resuspension—results from two tunnel studies. *Atmospheric Environment* 36, 4735-
706 4744.
707
708 Taylor, S.R. and McLennan, S.M., 1995. The geochemical evolution of the continental crust.
709 *Reviews of Geophysics* 33, 241-265.
710
711 Thorpe, A. and Harrison, R.M., 2008. Sources and properties of non-exhaust particulate matter from
712 road traffic: a review. *Science of the Total Environment* 400, 270-282.
713
714 Wahlin, P., Berkowicz, R. and Palmgren, F., 2006. Characterisation of traffic-generated particulate
715 matter in Copenhagen. *Atmospheric Environment* 40, 2151-2159.
716
717 Weichenthal, S.A., Godri Pollitt, K. and Villeneuve, P.J., 2013. PM_{2.5}, oxidant defense and
718 cardiorespiratory health: a review. *Environmental Health* 12, 40.
719
720 Yang, A., Jedynska, A., Hellack, B., Kooter, I., Hoek, G., Brunekreef, B., Kuhlbusch, T.A.J.,
721 Cassee, F.R. and Janssen, N.A.H., 2014. Measurement of the oxidative potential of PM_{2.5} and its
722 constituents: The effect of extraction solvent and filter type. *Atmospheric Environment* 83, 35-42.
723
724 Yanosky, J.D., Tonne, C.D., Beevers, S.D., Wilkinson, P. and Kelly, F.J., 2012. Modeling
725 exposures to the oxidative potential of PM. *Environmental Science and Technology* 46, 7612-7620.
726
727
728

729 **TABLE LEGENDS**

730

731

732 **Table 1:** Chemical source profiles for heavy traffic sites in Birmingham (Sites A and B) and
733 New Delhi (Site C) (% w/w).

734

735 **Table 2:** Source contributions of various non-exhaust sources to the PM₁₀ fraction of road dust
736 in Birmingham and Delhi (in %).

737

738 **Table 3:** Parameters of the regression model to explain oxidative potential variance for road
739 dust (more significant outcomes only).

740

741

742 **FIGURE LEGENDS**

743

744 **Figure 1:** Concentrations of various elements in the PM₁₀ fraction of road dust at Sites A-C
745 ($\mu\text{g/g}$).

746

747 **Figure 2:** Enrichment factors for sites A-C.

748

749 **Figure 3:** Comparison of species concentration (weight %) observed in the current study with
750 the PM₁₀ fraction of road dust sampled elsewhere.

751

752 **Figure 4:** Oxidative Potential (OP) per μg dust for the PM₁₀ fraction and bulk road dust
753 samples for site A.

754

755

756

757 **Table 1:** Chemical source profiles for heavy traffic sites in Birmingham (Sites A and B) and New
 758 Delhi (Site C) (% w/w)

759

<i>Species</i>	<i>Site A</i>		<i>Site B</i>		<i>Site C</i>	
	Conc.	SD	Conc.	SD	Conc.	SD
OC	7.91	4.80			1.82	1.69
EC	0.13	0.41			0.51	0.29
Si	28.8	17.6	28.2	8.98	13.0	3.04
Al	8.66	4.81	7.22	2.34	3.98	1.04
Ca	3.24	1.62			5.00	0.64
Fe	7.39	8.92	7.32	2.00	3.11	0.49
Ti	0.032	0.025	0.078	0.029	0.047	0.005
Mn	0.044	0.037	0.077	0.025	0.044	0.005
Cu	0.035	0.029	0.108	0.032	0.016	0.002
Sb	0.007	0.006	0.014	0.003	0.001	0.000
Ba	0.033	0.027	0.087	0.027	0.042	0.005
Sn	0.005	0.004			0.001	0.000
Cr	0.007	0.006	0.017	0.007	0.006	0.001
V	0.002	0.002	0.005	0.002	0.003	0.000
Zn	0.071	0.058	0.366	0.142	0.068	0.009
Ni			0.012	0.009		
S	0.11	0.18	0.99	0.29		
Benzo(b)fluoranthene (BbF)	0.004	0.002			0.004	0.003
Benzo(k)fluoranthene (BkF)	0.004	0.002			0.003	0.002
Benzo(e)pyrene (BeP)	0.003	0.002			0.002	0.001
Benzo(a)pyrene (BaP)	0.002	0.001			0.002	0.002
Indeno(123-cd)pyrene (IcdP)	0.002	0.001			0.002	0.002
Benzo(ghi)perylene (BghiPe)	0.002	0.001			0.002	0.002
Coronene (Cor)	0.001	0.000			0.001	0.000

760

761

762 **Table 2:** Source contributions of various non-exhaust sources to the PM₁₀ fraction of road dust in
 763 Birmingham and Delhi (in %)

<i>Site ID</i>	<i>Site characteristics</i>	<i>Brake Wear (Ba)</i>	<i>Tyre Wear (Zn)</i>	<i>Crustal Dust (Si/Fe)</i>	<i>Vehicle Exhaust</i>	<i>Total Mass Estimated</i>
Site A	High traffic volume, stop-and-go traffic, mixed LDVs and HDVs	3.8	4.7	89.5	1.3	99.3
Site B	High traffic volume, smooth traffic flow with stop-and-go during congestion, mixed LDVs and HDVs	6.6	15.6	82.8	-	105
Site C	High traffic volume, stop-and-go traffic, mixed LDVs and HDVs	3.9	3.5	71.2	0.59	79.2
	Open unpaved area in vicinity					

764

765

766

767 **Table 3:** Parameters of the regression model to explain oxidative potential variance for road dust
 768 (significant outcomes only)


<i>Sample type</i>	<i>Metric</i>	<i>Metal ($\mu\text{g/g}$)</i>	<i>Explanatory Variable</i>		<i>Model</i>		
			Slope	Intercept	R²	SE	p-value
Bulk RD ($<2\text{mm}$)	OP ^{AA} μg^{-1}	Cu	6019 \pm 416.4	-668 \pm 52.7	0.995	4.40	0.044
PM ₁₀ RD ($<10\ \mu\text{m}$)	OP ^{AA} μg^{-1}	Cu	2363 \pm 768	4.28 \pm 45.6	0.61	32.2	0.022
		Ti	5460 \pm 2076	-57.4 \pm 123	0.54	86.9	0.039
		V	331 \pm 118	-4.04 \pm 7.03	0.57	4.96	0.031
		Mn	4214 \pm 1643	-19.7 \pm 97.5	0.52	68.8	0.043
		As	64.8 \pm 23.9	-0.93 \pm 1.42	0.55	1.00	0.035
		S	86326 \pm 30757	-2705 \pm 1825	0.57	1288	0.031

769

770

771

772




773

774

775 **Figure 1:** Concentrations of various elements in the PM₁₀ fraction of road dust at Sites A-C (µg/g)
776 (Site A- UK roadside, Site B- UK tunnel and Site C- India roadside).

777


778



780

781 **Figure 2:** Crustal enrichment factors for sites A-C.


782



783

784 **Figure 3:** Comparison of species concentration (weight %) observed in the current study with the
 785 PM_{10} fraction of road dust sampled elsewhere.

786



787

788 Figure 4: Oxidative Potential (OP) per μg dust for the PM_{10} fraction and bulk road dust
 789 samples for site A.

790