

Front propagation in cellular flows for fast reaction and small diffusivity

Tzella, Alexandra; Vanneste, Jacques

DOI:

[10.1103/PhysRevE.90.011001](https://doi.org/10.1103/PhysRevE.90.011001)

License:

None: All rights reserved

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Tzella, A & Vanneste, J 2014, 'Front propagation in cellular flows for fast reaction and small diffusivity', *Physical Review E*, vol. 90, 011001. <https://doi.org/10.1103/PhysRevE.90.011001>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Published as: Tzella, Alexandra, and Jacques Vanneste. "Front propagation in cellular flows for fast reaction and small diffusivity." *Physical Review E* 90.1 (2014): 011001. Available online at: <http://dx.doi.org/10.1103/PhysRevE.90.011001>
© 2014 American Physical Society

Eligibility for repository checked September 2015

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Front propagation in cellular flows for fast reaction and small diffusivity

Alexandra Tzella^{1,*} and Jacques Vanneste²

¹*School of Mathematics, University of Birmingham, Birmingham, United Kingdom*

²*School of Mathematics and Maxwell Institute for Mathematical Sciences, University of Edinburgh, Edinburgh, United Kingdom*

(Received 4 April 2014; published 11 July 2014)

We investigate the influence of fluid flows on the propagation of chemical fronts arising in Fisher-Kolmogorov-Petrovsky-Piskunov (FKPP) type models. We develop an asymptotic theory for the front speed in a cellular flow in the limit of small molecular diffusivity and fast reaction, i.e., large Péclet (Pe) and Damköhler (Da) numbers. The front speed is expressed in terms of a periodic path—an instanton—that minimizes a certain functional. This leads to an efficient procedure to calculate the front speed, and to closed-form expressions for $(\log \text{Pe})^{-1} \ll \text{Da} \ll \text{Pe}$ and for $\text{Da} \gg \text{Pe}$. Our theoretical predictions are compared with (i) numerical solutions of an eigenvalue problem and (ii) simulations of the advection-diffusion-reaction equation.

DOI: [10.1103/PhysRevE.90.011001](https://doi.org/10.1103/PhysRevE.90.011001)

PACS number(s): 47.70.Fw, 05.10.-a, 05.40.-a, 82.40.Ck

The spreading of chemical or biological populations in fluid flows is a fundamental problem in many areas of science and engineering with applications ranging from plankton blooms to combustion [1,2]. In the absence of flow, this spreading results from the competition between spatial diffusion, local growth, and saturation, and leads to the formation of wave fronts that travel undeformed at constant speed [3]. A number of theoretical results describe the influence that divergence-free, spatially smooth flows have on such fronts (see Refs. [4,5] for comprehensive reviews). These have stimulated experiments using a variety of reactions and flow configurations [6–8], with much effort devoted to steady spatially periodic cellular flows [9–12].

Using the classic Fisher-Kolmogorov-Petrovsky-Piskunov (FKPP) model [13,14] based on logistic type growth, Gärtner and Freidlin [15] showed that the speed of the pulsating front that arises in such periodic flows can be obtained by solving an eigenvalue problem (see below). In practice, this procedure requires rather involved numerical computations; there is therefore a need for simplified results that provide scaling predictions or closed-form expressions in asymptotic regimes. Results of this type have been derived in the limit of slow reactions and small diffusivity [16,17]. Here we consider the opposite limit of fast reaction (e.g., Ref. [18]) relevant, for instance, to premixed flame propagation [19]. In this limit, we obtain a compact expression for the speed in terms of a single periodic path that minimizes an action functional. This brings physical insight, in particular, into the role of the flow's stagnation points, and yields closed-form results valid for a remarkably large range of reaction rates. It also provides an efficient way to compute the speed in a regime where direct numerical computations are most challenging because of the widely disparate spatial scales (see, e.g., Fig. 1).

Model. The effect of a background flow \mathbf{u} is incorporated in the FKPP advection-diffusion-reaction equation

$$\partial_t \theta + \mathbf{u} \cdot \nabla \theta = \text{Pe}^{-1} \Delta \theta + \text{Da} r(\theta), \quad (1)$$

for the population concentration θ . Here, the reaction term is $r(\theta) = \theta(1 - \theta)$ or, more generally, any function $r(\theta)$

that satisfies $r(0) = r(1) = 0$ with $r(\theta) > 0$ for $\theta \in (0, 1)$, $r(\theta) < 0$ for $\theta \notin [0, 1]$, and $r'(0) = \sup_{0 < \theta < 1} r(\theta)/\theta = 1$. The nondimensional parameters are the Péclet and Damköhler numbers $\text{Pe} = U\ell/\kappa$ and $\text{Da} = \ell/(U\tau)$, where U and ℓ are the characteristic amplitude and length scale of the flow, κ the molecular diffusivity, and τ the reaction time. We consider the two-dimensional cellular flow $\mathbf{u} = (u_1, u_2) = (-\partial_y \psi, \partial_x \psi)$ with stream function

$$\psi(x, y) = -\sin x \sin y. \quad (2)$$

We take the domain to be the strip $D = (-\infty, \infty) \times [0, \pi]$ along which an infinite array of identical cells are arranged, each of which is composed of two half cells of opposite circulation with hyperbolic stagnation points at each corner (streamlines are shown for a single cell in Fig. 2). As the initial condition we take $\theta(x, y, 0) = \Theta(-x)$, where Θ is the Heaviside step function. The boundary conditions are $\theta(\infty, y, t) = 0$ and $\theta(-\infty, y, t) = 1$, so that the front advances rightwards, and no flux, $\partial_y \theta = 0$ at $y = 0, \pi$. The front characteristics change drastically with Da. When Da is small, the front's leading edge is confined near the cell boundaries (see left column in Fig. 1). As Da increases, the front sharpens and the imprint of the flow, the boundary-layer structure in particular, is less prominent (see middle and right columns in Fig. 1).

The long-time speed of propagation of the front is determined by the behavior of the solution near the front's leading edge, where $0 < \theta \ll 1$ and $r(\theta) \approx r'(0)\theta = \theta$. For steady periodic flows such as (2), this is given by

$$c = \inf_{q>0} \frac{f(q) + \text{Da}}{q}, \quad (3)$$

where $f(q)$ is the largest eigenvalue of

$$\mathcal{L}v = f(q)v, \quad (4a)$$

with the operator \mathcal{L} defined by

$$\mathcal{L} = \text{Pe}^{-1} \Delta - \mathbf{u} \cdot \nabla - 2\text{Pe}^{-1} q \partial_x + (u_1 q + \text{Pe}^{-1} q^2), \quad (4b)$$

and periodic and no-flux boundary conditions in x and y , respectively (see Refs. [15,20 (Chap. 7),21]). This result is intimately connected with the long-time large-deviation rate

*a.tzella@bham.ac.uk




FIG. 1. (Color online) Successive snapshots of the concentration θ for $Pe = 50$ and $Da = 0.5$ (left), $Da = 5$ (middle), and $Da = 20$ (right), with time increasing from the top to the bottom rows. The corresponding front speeds are $c \approx 0.5$, $c \approx 1$, and $c \approx 1.6$.

function associated with the concentration of the nonreacting passive scalar (i.e., $Da = 0$); specifically, $f(q)$ is the Legendre transform of this rate function [15,20,22,23].

In the absence of a background flow, $f(q) = q^2/Pe$, recovering the classical formula for the bare speed $c_0 = 2\sqrt{Da/Pe} = 2\sqrt{\kappa/\tau}$ (see Refs. [13,14,24,25] and references therein). For general $\mathbf{u} \neq \mathbf{0}$, the eigenvalue problem (4) cannot be solved explicitly.

Small diffusivity, fast reaction. Our purpose is to use asymptotic analysis to obtain the speed of the front in the large-Péclet limit with

$$\gamma = Da/Pe = O(1),$$

corresponding to the geometric optics regime defined by $\kappa, \tau \rightarrow 0$ with $\kappa/\tau = O(1)$ [26]. This can be achieved by analyzing the large-Pe limit of the eigenvalue problem (4), or by considering the large- t limit of the geometric optics theory treatment of [20 (Chap. 6)] (see also Refs. [18,27]). It is, however, convenient to start with the basic result exploited by Refs. [15,20], namely, that the front speed is controlled by the point at which the solution to the linearization of Eq. (1) neither grows nor decays.




FIG. 2. (Color online) Streamlines of the cellular flow with stream function (2) (thin solid black lines) and trajectories of the instantons minimizing (9) (thick red lines). The instantons are calculated numerically for $c = 0.5$ (corresponding to $\gamma \approx 0.002$), $c = 1$ ($\gamma \approx 0.09$), and $c = 5$ ($\gamma \approx 5.9$), and become closer to the straight line $y = \pi/2$ as c increases. The dashed lines show the small- c and large- c asymptotic approximations obtained using (10) for $c = 0.5$ and (15) for $c = 5$, respectively.

We seek a solution to this equation in the WKB form

$$\theta(x, y, t) \sim \exp\{-Pe[I(x, y, t) - \gamma t]\}, \quad \text{for } Pe \gg 1, \quad (5)$$

where $I(x, y, t)$ can be recognized as the small-noise large-deviation rate function [28]. At leading order, $\partial_t I + H(\nabla I, x, y) = 0$, where $H = \|\nabla I\|^2 + \mathbf{u}(x, y) \cdot \nabla I$ is the Hamiltonian and $\|\cdot\|$ the usual norm. Its solution is well known from Hamilton-Jacobi theory (e.g., Refs. [28,29]) and is given by

$$I(x, y, t) = \frac{1}{4} \inf_{\phi(\cdot)} \int_0^t \|\dot{\phi}(s) - \mathbf{u}(\phi(s))\|^2 ds, \quad (6)$$

subject to $\phi(t) = (x, y)$ and $\phi(0) = (0, \cdot)$. Thus, the behavior of (5) is controlled by a single path $\phi^*(s)$ that minimizes this integral. This optimal path is often called *instanton* [30]. Expression (5) then indicates that for $t \gg 1$, the front speed satisfies

$$\gamma = \lim_{t \rightarrow \infty} \frac{I(ct, y, t)}{t} \equiv \mathcal{G}(c), \quad (7)$$

where the limit $t \rightarrow \infty$ eliminates the dependence on y . This leads to

$$c = \mathcal{G}^{-1}(\gamma). \quad (8)$$

The front speed is therefore obtained by calculating $\mathcal{G}(c)$. This calculation is significantly simplified by observing that the limit in (7) is determined in terms of periodic trajectories, an observation justified rigorously in Ref. [31]. We take solutions $\phi(s)$ to be periodic in the sense that $\phi(\tau) = \phi(0) + (2\pi, 0)$, where the period is $\tau = 2\pi/c$. Letting $\sigma = cs$, we obtain the simplified expression

$$\mathcal{G}(c) = \frac{1}{8\pi} \inf_{\phi(\cdot)} \int_0^{2\pi} \|c\phi'(\sigma) - \mathbf{u}(\phi(\sigma))\|^2 d\sigma, \quad (9)$$

subject to $\phi(2\pi) = \phi(0) + (2\pi, 0)$. Expressions (8) and (9) are the main result of this Rapid Communication. They provide a direct way of computing the instantons and thus the front speed. Note that $\mathcal{G}(c)$ may be interpreted as the Legendre transform of \bar{H} , the effective Hamiltonian of the homogenized Hamilton-Jacobi equation $\partial_t I + \bar{H}(\nabla I) = 0$ [18,32]. In order to derive (8), we have formally assumed that $\gamma = O(1)$. However, the asymptotic results apply over a broad range of values of γ , specifically $\gamma \gg (Pe \log Pe)^{-1}$, as we show below.

The solution to the minimization problem (9) is easy to obtain numerically. We first discretize (9) and use MATLAB to minimize the action. We iterate over c , starting with large




FIG. 3. Front speed c for $Pe \gg 1$ as a function of γ . Prediction (8) derived from the numerical minimization of (9) (solid black line) is compared with its small- c approximation (12) (lower dashed line), its large- c approximation (16) (upper dashed line), and the bare speed $c_0 = 2\sqrt{\gamma}$ (dotted line). The inset focuses on smaller values of γ .

values for which the straight line $\phi^*(\sigma) = (\sigma, \pi/2)$ is a good initial guess. Characteristic examples of instantons obtained for different values of c are shown in Fig. 2 (the two cases with $c = 0.5$ and $c = 1$ correspond to the first two columns in Fig. 1). For large values of c , the instanton is close to a straight line. For small values of c , it follows closely a streamline near the cell boundaries. Figure 3 shows the behavior of c as a function of γ deduced from (8).

Asymptotic limits. We now obtain closed-form expressions for the speed c in two asymptotic limits. The first one corresponds to $\gamma \ll 1$ and hence $c \ll 1$. Numerical results (see Fig. 2 for $c = 0.5$) suggest that the instanton departs from the streamline only for $y \approx \pi/2$ when it crosses the separatrix between adjacent cells. It is also clear that (9) is minimized when $\phi^*(\sigma) = (x(\sigma), y(\sigma))$ satisfies $cy' \approx -\cos x \sin y$ (so that the instanton and flow speeds differ only in the x direction). Exploiting symmetry to consider $0 \leq \sigma \leq \pi/2$ only, with

$x(0) = 0$, $y(0) = x(\pi/2) = \pi/2$, and $y'(\pi/2) = 0$, we divide the instanton into two regions (see Fig. 2). In region 1, where $x \ll 1$, the integrand in (9) is approximately $(cx' - x \cos y)^2$, leading to the Euler-Lagrange equation $c^2 x'' = x$ (since $cy' \approx -\sin y$). In region 2, $y \ll 1$, $cx' = \sin x \cos y \approx -\sin x$, and $cy' = -\cos x \sin y \approx -y \cos x$. Matching the solutions for $x, y \ll 1$ (the cell corner) gives the approximation

$$\phi^*(\sigma) \sim \begin{cases} (C_1(\sigma), C_2(\sigma)) & \text{for } \sigma \ll \pi/2, \\ (C_2(\pi/2 - \sigma), C_3(\pi/2 - \sigma)) & \text{for } \sigma \gg c, \end{cases} \quad (10)$$

where

$$\begin{aligned} C_1(\sigma) &= 4 \exp\left(-\frac{\pi}{2c}\right) \sinh\left(\frac{\sigma}{c}\right), \\ C_2(\sigma) &= 2 \tan^{-1}\left[\exp\left(-\frac{\sigma}{c}\right)\right], \\ C_3(\sigma) &= 4 \exp\left(-\frac{\pi}{2c}\right) \cosh\left(\frac{\sigma}{c}\right). \end{aligned}$$

Expression (10) is in very good agreement with our numerical solution (see Fig. 2 for $c = 0.5$).

Using (10) gives the integrand in (9) as $(cx' - x \cos y)^2 \approx 16 \exp(-\pi/c) \cosh^{-2}(\sigma/c)$, leading to

$$\mathcal{G}(c) \sim 4 \times (2/\pi) c e^{-\pi/c}, \quad \text{where } c \ll 1, \quad (11)$$

and the factor 4 appears because, for $\sigma \in [0, 2\pi]$, there are four regions that are similar to region 1. Inverting (11) and using (8) finally gives

$$c \sim \frac{\pi}{W_p(8\gamma^{-1})}, \quad \text{for } \gamma \ll 1, \quad (12)$$

where W_p denotes the principal real branch of the Lambert W function [33]. This approximation holds for $(\log Pe)^{-1} \ll Da \ll Pe$: the lower bound follows from requiring that the argument of the exponential in (5) be large, $Pe I \sim Da \tau \gg 1$, where the period is roughly estimated as $\tau = 2\pi/c \sim \log Pe/2$ (see (13) below and Ref. [23] for a complete argument). Figure 3 shows that this approximation is excellent within its range of validity. Since $\gamma \ll 1$, it is consistent to approximate $W_p(8\gamma^{-1})$ to reduce (12) to

$$c \approx \frac{\pi}{\log Pe}. \quad (13)$$

A qualitatively similar expression was obtained in Refs. [34,35] using a heuristic approach based on the so-called G equation. Note that the logarithmic dependence of the speed on Pe and its slow growth with Da [captured by (12) but not (13)] is associated with the holdup of the instanton near the hyperbolic stagnation points at the cell corners.

The second limit leading to closed-form results corresponds to $\gamma \gg 1$, hence $c \gg 1$. In this case, we seek an instanton as a power series $\phi^*(\sigma) = (\sigma, y_0) + c^{-1}(x_1(\sigma), y_1(\sigma)) + \dots$, where x_1, y_1 are functions of period 2π that satisfy $x_1(0) = y_1(0) = 0$. Substituting into (9), we find that at $O(c^{-1})$,

$$\mathcal{G}(c) = \frac{c^2}{4} + \frac{1}{8\pi} \inf_{x_1, y_0, y_1} \int_0^{2\pi} (x_1'^2 + y_1'^2 + 4y_1 \sin \sigma \sin y_0) d\sigma, \quad (14)$$




FIG. 4. (Color online) Front speed c as a function of Da for different values of Pe . The large- Pe prediction (8) derived from the numerical minimization of (9) (dashed lines) is compared with the exact expression (3) estimated by solving the eigenvalue problem (4) numerically (solid lines) and with direct numerical simulations of (1) (symbols).

after some manipulations. Minimizing this integral leads to the instanton

$$\phi^*(\sigma) = \left(\sigma, \frac{\pi}{2}\right) + c^{-1}(0, -2 \sin \sigma) + \dots, \quad \text{for } c \gg 1, \quad (15)$$

in excellent agreement with our numerical solution (see Fig. 2 for $c = 5$). Combining (14) and (15) yields $\mathcal{G}(c) = c^2/4 - 3/8 + O(c^{-2})$. We now use (8) to find that

$$c \sim 2\sqrt{\gamma} \left(1 + \frac{3}{16\gamma} + \dots\right), \quad \text{for } \gamma \gg 1, \quad (16)$$

which corresponds to $Da \gg Pe$. The leading-order term in (16) is the bare speed c_0 . As Fig. 3 shows, the second term in the expansion is necessary for a good agreement between asymptotic and full results.

Comparison with numerical results. We now compare our predictions for c derived from (8)–(9) with values obtained from (i) a numerical evaluation of the principal eigenvalue in (4), and (ii) direct numerical simulations of (1) with $r(\theta) = \theta(1 - \theta)$. For (i) we use a standard second-order discretization to approximate (4) and choose the spatial resolution Δ to satisfy $\pi/\Delta = 750$. The resulting matrix eigenvalue problem is solved for a range of values of q using MATLAB. For (ii) we discretize (1) using a fractional-step method with a Godunov splitting which alternates between advection (using a first-order upwind method with a minmod limiter—see Ref. [36] for details), diffusion (using an alternating direction implicit method), and reaction (solved exactly). We choose the same spatial resolution Δ as for (4). The computational domain is made finite using artificial boundaries at $x = \pm N\pi$, with $N = 5$, so that boundary effects are negligible. The front is tracked for long times by modifying the computational domain: When the solution at $x = (N - 1)\pi$ exceeds $\delta = 10^{-6}$, we eliminate

the nodes with $-N\pi \leq x \leq (-N + 1)\pi$ and add new nodes with $N\pi \leq x \leq (N + 1)\pi$ where we set $\theta = 0$. We calculate the front speed using a linear fit of the right endpoint of the front, $x_\epsilon^+(t) = \max\{x : \theta(x, t) = \epsilon\}$ where $\epsilon = 10^{-3}$. Results are insensitive to the exact values of ϵ and δ .

The three sets of numerical results are shown in Fig. 4. The speeds derived from the eigenvalue equation (4) are in excellent agreement with the corresponding values obtained from the full numerical simulations of Eq. (1). With increasing values of Pe , the asymptotic expression (9) becomes increasingly accurate, with excellent agreement for $Pe = 250, 500$ and satisfactory agreement for the moderate values $Pe = 50, 125$. As expected, (8)–(9) is valid for a broad range of values of Da , restricted only by $Da \gg (\log Pe)^{-1}$. Note that the use of both the eigenvalue equation and the full numerical simulations is restricted: as Pe increases, the solutions to (1) and (4) become progressively localized, with $O(1/\sqrt{Da Pe})$ length scales that are challenging to resolve when $Da, Pe \gg 1$.

Conclusion. We have derived a compact expression for the front speed based on the minimization of the large-deviation action (9) over periodic instantons. This leads to the efficient computation of the speed for a large range of values of Da . For the particular case of cellular flows, this expression provides the closed-form results (12) and (16) valid for $(\log Pe)^{-1} \ll Da \ll Pe$ and $Da \gg Pe$. In the first regime, the passage of the front near the stagnation points at the cell corners is shown to control the front speed; as a result this is almost insensitive to the reaction rate and depends logarithmically on the Péclet number. For $Da = O(\log Pe)^{-1}$ and smaller, the front speed is not controlled by a single minimizing trajectory, and asymptotic solutions to the eigenvalue problem (4) must be sought by other means; this will be the subject of future work [23].

Acknowledgments. The authors thank P. H. Haynes, G. C. Papanicolaou, and A. Pocheau for helpful discussions. The work was supported by EPSRC (Grant No. EP/I028072/1).

-
- [1] T. Tel, A. de Moura, C. Grebogi, and G. Károlyi, *Phys. Rep.* **413**, 91 (2005).
- [2] Z. Neufeld and E. Hernández-García, *Chemical and Biological Processes in Fluid Flows: A Dynamical Systems Approach* (Imperial College Press, London, 2009).
- [3] J. Murray, *Mathematical Biology I: An Introduction*, 3rd ed. (Springer, Berlin, 2002).
- [4] J. Xin, *SIAM Rev.* **42**, 161 (2000).
- [5] J. Xin, *An Introduction to Fronts in Random Media* (Springer, Berlin, 2000).
- [6] M. Leconte, J. Martin, N. Rakotomalala, and D. Salin, *Phys. Rev. Lett.* **90**, 128302 (2003).
- [7] M. S. Paoletti, C. R. Nugent, and T. H. Solomon, *Phys. Rev. Lett.* **96**, 124101 (2006).
- [8] M. E. Schwartz and T. H. Solomon, *Phys. Rev. Lett.* **100**, 028302 (2008).
- [9] M. S. Paoletti and T. H. Solomon, *Phys. Rev. E* **72**, 046204 (2005).
- [10] A. Pocheau and F. Harambat, *Phys. Rev. E* **73**, 065304 (2006).
- [11] A. Pocheau and F. Harambat, *Phys. Rev. E* **77**, 036304 (2008).
- [12] B. W. Thompson, J. Novak, M. C. T. Wilson, M. M. Britton, and A. F. Taylor, *Phys. Rev. E* **81**, 047101 (2010).
- [13] R. Fisher, *Ann. Eugen.* **7**, 355 (1937).
- [14] A. N. Kolmogorov, I. G. Petrovsky, and N. S. Piskunov, *Bull. Univ. Moskov. Ser. Int. Sect.* **1**, 1 (1937).
- [15] J. Gärtner and M. I. Freidlin, *Sov. Math. Dokl.* **20**, 1282 (1979).
- [16] B. Audoly, H. Berestycki, and Y. Pomeau, *C. R. Acad. Sci., Ser. IIB: Mec.* **328**, 255 (2000).
- [17] A. Novikov and L. Ryzhik, *Arch. Ration. Mech. Anal.* **184**, 23 (2007).
- [18] A. J. Majda and O. E. Sogadinis, *Nonlinearity* **7**, 1 (1994).
- [19] N. Peters, *Turbulent Combustion* (Cambridge University Press, Cambridge, UK, 2000).
- [20] M. I. Freidlin, *Functional Integration and Partial Differential Equations*, Annals of Mathematics Studies Vol. 109 (Princeton University Press, Princeton, 1985).
- [21] L. Evans and P. Souganidis, *Ann. Inst. Poincaré Anal. Non Linéaire* **S6**, 229 (1989).

- [22] P. H. Haynes and J. Vanneste, *J. Fluid Mech.* **745**, 321 (2014).
- [23] A. Tzella and J. Vanneste (unpublished).
- [24] J. Leach and D. Needham, *Matched Asymptotic Expansions in Reaction-Diffusion Theory*, Springer Monographs in Mathematics (Springer, Berlin, 2003).
- [25] W. van Saarloos, *Phys. Rep.* **386**, 29 (2003).
- [26] F. A. Williams, in *The Mathematics of Combustion*, edited by J. D. Buckmaster (SIAM, Philadelphia, PA, 1985), pp. 97–131.
- [27] M. I. Freidlin and R. B. Sowers, *Stochastic Process. Appl.* **82**, 23 (1999).
- [28] M. I. Freidlin and A. D. Wentzell, *Random Perturbations of Dynamical Systems* (Springer-Verlag, New York, 1984).
- [29] C. L. Evans, *Partial Differential Equations*, Vol. 19 (American Mathematical Society, Providence, RI, 2010).
- [30] M. I. Dykman, E. Mori, J. Ross, and P. M. Hunt, *J. Chem. Phys.* **100**, 5735 (1994).
- [31] A. Piatnitski, *Commun. Math. Phys.* **197**, 527 (1998).
- [32] P. L. Lions, G. C. Papanicolaou, and S. Varadhan (unpublished).
- [33] “NIST digital library of mathematical functions”, <http://dlmf.nist.gov/>, Release 1.0.6 of 2013-05-06 .
- [34] M. Abel, M. Cencini, D. Vergni, and A. Vulpiani, *Chaos* **12**, 481 (2002).
- [35] M. Cencini, A. Torcini, D. Vergni, and A. Vulpiani, *Phys. Fluids* **15**, 679 (2003).
- [36] R. J. Leveque, *Finite Volume Methods for Hyperbolic Problems* (Cambridge University Press, Cambridge, UK, 2002).