

A generalization of de Vries duality to closed relations between compact Hausdorff spaces

Abbadini, Marco; Bezhanishvili, Guram; Carai, Luca

DOI:

[10.1016/j.topol.2023.108641](https://doi.org/10.1016/j.topol.2023.108641)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Abbadini, M, Bezhanishvili, G & Carai, L 2023, 'A generalization of de Vries duality to closed relations between compact Hausdorff spaces', *Topology and its Applications*, vol. 337, 108641.
<https://doi.org/10.1016/j.topol.2023.108641>

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.


Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.



A generalization of de Vries duality to closed relations between compact Hausdorff spaces



Marco Abbadini^a, Guram Bezhanishvili^b, Luca Carai^{a,c,*}

^a Department of Mathematics, University of Salerno, Piazza Renato Caccioppoli, 2, 84084, Fisciano (SA), Italy

^b Department of Mathematical Sciences, New Mexico State University, 88003, Las Cruces, NM, USA

^c Department of Philosophy, University of Barcelona, Carrer de Montalegre 6, 08001, Barcelona, Spain

ARTICLE INFO

Article history:

Received 12 June 2022

Received in revised form 22 June 2023

Accepted 7 July 2023

Available online 13 July 2023

MSC:

54D30

54G05

18F70

06E15

18B10

18E08

Keywords:

Compact Hausdorff space

Closed relation

Subordination

De Vries algebra

ABSTRACT

Stone duality generalizes to an equivalence between the categories Stone^R of Stone spaces and closed relations and BA^S of boolean algebras and subordination relations. Splitting equivalences in Stone^R yields a category that is equivalent to the category KHaus^R of compact Hausdorff spaces and closed relations. Similarly, splitting equivalences in BA^S yields a category that is equivalent to the category DeV^S of de Vries algebras and compatible subordination relations. Applying the machinery of allegories then gives that KHaus^R is equivalent to DeV^S , thus resolving a problem recently raised in the literature.

The equivalence between KHaus^R and DeV^S further restricts to an equivalence between the category KHaus of compact Hausdorff spaces and continuous functions and the wide subcategory DeV^F of DeV^S whose morphisms satisfy additional conditions. This yields an alternative to de Vries duality. One advantage of this approach is that composition of morphisms is usual relation composition.

© 2023 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

In [12], de Vries generalized Stone duality to a duality for the category KHaus of compact Hausdorff spaces and continuous maps. The objects of the dual category DeV are complete boolean algebras equipped with a proximity relation, known as de Vries algebras. The morphisms of DeV are functions satisfying certain conditions. One drawback of DeV is that composition of morphisms is not usual function composition. In this

* Corresponding author at: Department of Philosophy, University of Barcelona, Carrer de Montalegre 6, 08001, Barcelona, Spain.

E-mail addresses: marco.abbadini.uni@gmail.com (M. Abbadini), guram@nmsu.edu (G. Bezhanishvili), luca.carai.uni@gmail.com (L. Carai).

paper we propose an alternative approach to de Vries duality, where morphisms between de Vries algebras become certain relations and composition is usual relation composition.

For our purpose, it is more natural to start with the category \mathbf{KHaus}^R whose objects are compact Hausdorff spaces and morphisms are closed relations (i.e., relations $R: X \rightarrow Y$ such that R is a closed subset of $X \times Y$). This category was studied in [5], and earlier in [23] in the more general setting of stably compact spaces. The latter paper establishes a duality for \mathbf{KHaus}^R that generalizes Isbell duality [19] between \mathbf{KHaus} and the category of compact regular frames and frame homomorphisms. This is obtained by generalizing the notion of a frame homomorphism to that of a preframe homomorphism. However, a similar duality in the language of de Vries algebras remained problematic (see [5, Rem. 3.14]). We resolve this problem as follows.

By Stone duality, the category \mathbf{Stone} of Stone spaces (zero-dimensional compact Hausdorff spaces) and continuous maps is dually equivalent to the category \mathbf{BA} of boolean algebras and boolean homomorphisms. Halmos [15] generalized Stone duality by showing that continuous relations between Stone spaces dually correspond to functions $f: A \rightarrow B$ between boolean algebras that preserve finite meets. This was further generalized by Celani [11] to closed relations between Stone spaces and certain functions from a boolean algebra A to the ideal frame $\mathcal{J}(B)$ of a boolean algebra B . We show that such functions correspond to subordination relations between A and B studied in [4]. Consequently, we obtain that the category \mathbf{Stone}^R of Stone spaces and closed relations is equivalent to the category \mathbf{BA}^S of boolean algebras and subordination relations (identity is \leq and composition is usual composition of relations). This equivalence is in fact an equivalence of allegories, hence self-dual categories. Thus, the choice of direction of morphisms is ultimately a matter of taste and the equivalence can alternatively be stated as a dual equivalence.

We point out that the equivalence between \mathbf{Stone}^R and \mathbf{BA}^S is also a consequence of a more general result of Jung, Kurz, and Moshier [24, Thm. 5.9] who worked with order-enriched categories to show that Priestley duality extends to a dual equivalence between the category of bounded distributive lattices and subordination relations and the category of Priestley spaces and Priestley relations.

As we pointed out above, both \mathbf{Stone}^R and \mathbf{BA}^S are allegories, and the equivalence between \mathbf{Stone}^R and \mathbf{BA}^S is an equivalence of allegories. Therefore, we may utilize the machinery of allegories [14] to split equivalences in both \mathbf{Stone}^R and \mathbf{BA}^S . Splitting equivalences in \mathbf{Stone}^R yields the category \mathbf{StoneE}^R which is equivalent to \mathbf{KHaus}^R . On the other hand, splitting equivalences in \mathbf{BA}^S yields the category $\mathbf{SubS5}^S$ which we show is equivalent to its full subcategory \mathbf{DeV}^S consisting of de Vries algebras. By piggybacking the equivalence of \mathbf{Stone}^R and \mathbf{BA}^S , we obtain an equivalence between \mathbf{StoneE}^R and $\mathbf{SubS5}^S$, which then yields our desired equivalence between \mathbf{KHaus}^R and \mathbf{DeV}^S . This resolves an open problem raised in [5, Rem. 3.14] (see Remark 4.8).

The category \mathbf{StoneE}^R has a full subcategory \mathbf{Gle}^R of Gleason spaces [4,5] which is more directly related to \mathbf{DeV}^S . We thus obtain a series of equivalences of allegories (see the diagram after Corollary 4.7). We prove that, unlike \mathbf{StoneE}^R and $\mathbf{SubS5}^S$, isomorphisms in \mathbf{DeV}^S and \mathbf{Gle}^R correspond to structure-preserving bijections.

The equivalence between \mathbf{KHaus}^R and \mathbf{DeV}^S can also be obtained by directly generalizing the regular open functor of de Vries duality. Indeed, associate with each compact Hausdorff space X the de Vries algebra $(\mathbf{RO}(X), \prec)$, where $\mathbf{RO}(X)$ is the complete boolean algebra of regular open subsets of X and \prec is the de Vries proximity on $\mathbf{RO}(X)$ defined by $U \prec V$ iff $\text{cl}(U) \subseteq V$. The key is to associate with each closed relation $R: X \rightarrow Y$ the relation $S_R: \mathbf{RO}(X) \rightarrow \mathbf{RO}(Y)$ given by

$$U S_R V \iff R[\text{cl}(U)] \subseteq V$$

(here $R[-]$ denotes the direct image under R). This gives a more explicit description of the equivalence between \mathbf{KHaus}^R and \mathbf{DeV}^S , which further restricts to an equivalence between \mathbf{KHaus} and a wide subcategory \mathbf{DeV}^F of \mathbf{DeV}^S , thus providing an alternative to de Vries duality.

This paper is related to the line of research that was initiated by D. Scott [33], and further developed in [28,18,36,2,25,22,23,26,31,24], that uses certain relations as morphisms. We apply the insights of these works specifically to the category of compact Hausdorff spaces and closed relations. We do so in a way that is optimal to explain the connection with de Vries duality. The setting of compact Hausdorff spaces allows the use of simple entities such as closed equivalence relations on Stone spaces and their quotients, which are familiar to a wide range of mathematicians approaching topology from the algebraic or logical perspective.

2. Lifting Stone duality to closed relations

For two sets X and Y , we write $R: X \rightarrow Y$ to indicate that R is a relation from X to Y . As usual, for $F \subseteq X$, we write $R[F]$ for the R -image of F in Y ; and for $G \subseteq Y$, we write $R^{-1}[G]$ for the R -inverse image of G in X .

If X, Y are Stone spaces, then we call $R: X \rightarrow Y$ *closed* if R is a closed subset of $X \times Y$ (equivalently, $R[F]$ is closed for each closed $F \subseteq X$ and $R^{-1}[G]$ is closed for each closed $G \subseteq Y$).

Definition 2.1. Let Stone^R be the category of Stone spaces and closed relations between them. Identity morphisms in Stone^R are identity relations and composition in Stone^R is relation composition.

As we pointed out in the introduction, Celani [11] extended Stone duality to Stone^R by generalizing boolean homomorphisms to what he termed “quasi-semi-homomorphisms.” For a boolean algebra B , let $\mathcal{J}(B)$ be the complete lattice of ideals of B .

Definition 2.2. [11, Def. 1] Let A, B be boolean algebras. A *quasi-semi-homomorphism* is a function $\Delta: A \rightarrow \mathcal{J}(B)$ such that $\Delta(1) = B$ and $\Delta(a \wedge b) = \Delta(a) \cap \Delta(b)$ for all $a, b \in A$.

By [11, Lem. 2], boolean algebras and quasi-semi-homomorphisms between them form a category, which we denote by BA^Q . The identity quasi-semi-homomorphism on A is given by $I_A(a) = \downarrow a$ and the composition of two quasi-semi-homomorphisms $\Delta_1: A \rightarrow \mathcal{J}(B)$ and $\Delta_2: B \rightarrow \mathcal{J}(C)$ by

$$(\Delta_2 \circ \Delta_1)(a) = \bigcup \{ \Delta_2(b) \mid b \in \Delta_1(a) \}$$

(note that this union is an ideal because $\{ \Delta_2(b) \mid b \in \Delta_1(a) \}$ is directed).

Theorem 2.3. [11, Thm. 4] Stone^R is dually equivalent to BA^Q .

Quasi-semi-homomorphisms $A \rightarrow \mathcal{J}(A)$ were studied in [9] under the name of quasi-modal operators. It was pointed out in [4, Rem. 2.6] that quasi-modal operators on A are in one-to-one correspondence with subordinations on A . We show that this generalizes to a dual isomorphism between BA^Q and the category of boolean algebras and subordination relations between them.

Definition 2.4.

1. A *subordination relation* from a boolean algebra A to a boolean algebra B is a relation $S: A \rightarrow B$ satisfying, for $a, b \in A$ and $c, d \in B$:

- (S1) $0 S 0$ and $1 S 1$;
- (S2) $a, b S c$ implies $(a \vee b) S c$;
- (S3) $a S c, d$ implies $a S (c \wedge d)$;
- (S4) $a \leq b S c \leq d$ implies $a S d$.

2. Let \mathbf{BA}^S be the category of boolean algebras and subordination relations. The identity morphism on A is the order relation \leq on A , and composition in \mathbf{BA}^S is relation composition.

Theorem 2.5. \mathbf{BA}^S is dually isomorphic to \mathbf{BA}^Q .

Proof. For a subordination $S: A \rightarrow B$, define $\Delta_S: B \rightarrow \mathcal{J}(A)$ by $\Delta_S(b) = S^{-1}[b]$. It is straightforward to check that Δ_S is a well-defined quasi-semi-homomorphism. Moreover, $\Delta_{\leq_A}(a) = \downarrow a$, so $\Delta_{\leq_A} = I_A$. Furthermore, if $S_1: A \rightarrow B$ and $S_2: B \rightarrow C$ are subordinations, then

$$\Delta_{S_2 \circ S_1}(c) = (S_2 \circ S_1)^{-1}[c] = S_1^{-1}S_2^{-1}[c] = (\Delta_{S_1} \circ \Delta_{S_2})(c).$$

For a quasi-semi-homomorphism $\Delta: A \rightarrow \mathcal{J}(B)$, define $S_\Delta: B \rightarrow A$ by $b S_\Delta a$ iff $b \in \Delta(a)$. Again, it is straightforward to check that S_Δ is a well-defined subordination. Moreover, $b S_{I_A} a$ iff $b \in \downarrow a$ iff $b \leq a$, so $S_{I_A} = \leq_A$. Furthermore, if $\Delta_1: A \rightarrow \mathcal{J}(B)$ and $\Delta_2: B \rightarrow \mathcal{J}(C)$ are quasi-semi-homomorphisms, then

$$\begin{aligned} c S_{\Delta_2 \circ \Delta_1} a &\iff c \in (\Delta_2 \circ \Delta_1)(a) \iff c \in \bigcup \{\Delta_2(b) \mid b \in \Delta_1(a)\} \\ &\iff \exists b \in \Delta_1(a) : c \in \Delta_2(b) \iff c (S_{\Delta_1} \circ S_{\Delta_2}) a. \end{aligned}$$

In addition, for each subordination S , we have $b S_{\Delta_S} a$ iff $b \in \Delta_S(a)$ iff $b S a$, so $S_{\Delta_S} = S$. Also, for each quasi-semi-homomorphism Δ , we have $\Delta_{S_\Delta}(a) = S_\Delta^{-1}[a] = \Delta(a)$, and hence $\Delta_{S_\Delta} = \Delta$. Thus, \mathbf{BA}^S is dually isomorphic to \mathbf{BA}^Q . \square

Putting Theorems 2.3 and 2.5 together yields:

Corollary 2.6. \mathbf{Stone}^R is equivalent to \mathbf{BA}^S .

Remark 2.7. The functors establishing the equivalence of Corollary 2.6 generalize the well-known clopen and ultrafilter functors \mathbf{Clop} and \mathbf{Ult} of Stone duality. Indeed, if $R: X \rightarrow Y$ is a closed relation between Stone spaces, then composing the functors $\mathbf{Stone}^R \rightarrow \mathbf{BA}^Q \rightarrow \mathbf{BA}^S$, we obtain that the corresponding subordination relation $S_R: \mathbf{Clop}(X) \rightarrow \mathbf{Clop}(Y)$ is given by

$$U S_R V \iff R[U] \subseteq V.$$

Conversely, if $S: A \rightarrow B$ is a subordination relation, then composing the functors $\mathbf{BA}^S \rightarrow \mathbf{BA}^Q \rightarrow \mathbf{Stone}^R$ in the other direction yields that the corresponding closed relation $R_S: \mathbf{Ult}(A) \rightarrow \mathbf{Ult}(B)$ is given by

$$x R_S y \iff S[x] \subseteq y.$$

We will slightly abuse notation and use \mathbf{Clop} and \mathbf{Ult} also for the functors establishing the equivalence between \mathbf{Stone}^R and \mathbf{BA}^S .

Remark 2.8. That in Corollary 2.6 we have an equivalence instead of a dual equivalence is explained by the fact that the categories \mathbf{Stone}^R and \mathbf{BA}^S are self-dual (see Theorem 2.14), and hence equivalence implies dual equivalence.

Remark 2.9. Corollary 2.6 is also a consequence of [24, Thm. 5.9], where a more general order-enriched duality result is established between distributive lattices and Priestley spaces, with appropriate relations as morphisms.

From now on we will work with the additional structure of allegory on Stone^R and BA^S [14,21]. In fact, it is enough to have the structure of order-enriched category with involution [34,27]. However, we prefer to work with allegories because their well-developed machinery is readily available to us. One could instead work with bicategories (see [8,7]).

Definition 2.10.

1. [17, p. 105] A category C is *order enriched* if each hom-set of C is equipped with a partial order \leq such that $f \leq f'$ and $g \leq g'$ imply $gf \leq g'f'$ for all $f, f': C \rightarrow D$ and $g, g': D \rightarrow E$.
2. [16, p. 74] A *dagger* on a category C is a contravariant endofunctor $(-)^{\dagger}: C \rightarrow C$ that is the identity on objects and the composition of $(-)^{\dagger}$ with itself is the identity on C . A *dagger category* is a category equipped with a dagger.
3. [21, p. 136] An *allegory* A is an order-enriched dagger category such that:
 - (i) each partially ordered hom-set of A has binary meets;
 - (ii) $(-)^{\dagger}$ preserves the order on the hom-sets ($f \leq g \implies f^{\dagger} \leq g^{\dagger}$);
 - (iii) the modular law holds: $gf \wedge h \leq (g \wedge hf^{\dagger})f$ for all $f: C \rightarrow D, g: D \rightarrow E$, and $h: C \rightarrow E$.

We next extend the notion of equivalence of categories to allegories.

Definition 2.11.

1. [21, p. 142] If $F: A \rightarrow A'$ is a functor between two allegories, we say that F is a *morphism of allegories* if it preserves the binary meets of morphisms and commutes with $(-)^{\dagger}$.
2. A morphism of allegories is an *equivalence of allegories* if it is an equivalence of categories.

Remark 2.12. If $F: A \rightarrow A'$ is an equivalence of allegories with quasi-inverse $G: A' \rightarrow A$, then G is also an equivalence of allegories. To see this, if f is an isomorphism in A' , then $f^{-1} = f^{\dagger}$ (see [14, p. 199]). Therefore, since F is an essentially surjective, full, and faithful functor that preserves the dagger, the proof of [35, Lem. V.1] shows that G also preserves the dagger. Because F is a bijection on the hom-sets that preserves meets, it is an order-isomorphism on the hom-sets. Thus, G is also an order-isomorphism on the hom-sets, and hence an equivalence of allegories.

Lemma 2.13. Stone^R is an allegory.

Proof. Stone^R is an order-enriched category if we order its hom-sets by inclusion. For a relation $R: X \rightarrow Y$, let $R^{\smile}: Y \rightarrow X$ be its converse defined by $y R^{\smile} x$ iff $x R y$. If R is a closed relation, then R^{\smile} is also closed. The assignment $R \mapsto R^{\smile}$ defines a dagger on Stone^R . It is straightforward to check that all the conditions of Definition 2.10(3) hold in Stone^R . \square

Theorem 2.14. BA^S is an allegory and Clop and Ult yield an equivalence of Stone^R and BA^S as allegories.

Proof. If we order the hom-sets of BA^S by reverse inclusion, then BA^S becomes an order-enriched category. For a subordination $S: A \rightarrow B$, define $\widehat{S}: B \rightarrow A$ by $b \widehat{S} a$ iff $\neg a S \neg b$. It is straightforward to check that \widehat{S} is a subordination and that the assignment $S \mapsto \widehat{S}$ defines a dagger on BA^S . We show that the bijections between the hom-sets induced by the functors Clop and Ult preserve and reflect the orders, and commute with the daggers.

Suppose $R_1 \subseteq R_2$. Let $U \in \text{Clop}(X)$ and $V \in \text{Clop}(Y)$ such that $U S_{R_2} V$. Then $R_1[U] \subseteq R_2[U] \subseteq V$, and so $U S_{R_1} V$. Thus, $S_{R_2} \subseteq S_{R_1}$. Conversely, assume that $R_1 \not\subseteq R_2$. Let $x \in X$ and $y \in Y$ be such that

$x R_1 y$ but $x \not R_2 y$. Since R_2 is a closed relation, $R_2[x]$ is a closed subset of Y and $y \notin R_2[x]$. Because Y is a Stone space, there is clopen $V \subseteq Y$ such that $R_2[x] \subseteq V$ and $y \notin V$. Since R_2 is closed, $X \setminus R_2^{-1}[Y \setminus V]$ is an open subset of X containing x . Therefore, there is clopen $U \subseteq X$ such that $x \in U \subseteq X \setminus R_2^{-1}[Y \setminus V]$, so $R_2[U] \subseteq V$. Thus, $x \in U$, $y \notin V$, and $x R_1 y$, yielding $R_1[U] \not\subseteq V$. Consequently, $U S_{R_2} V$ but $U \not S_{R_1} V$, which implies that $S_{R_2} \not\subseteq S_{R_1}$.

Suppose $S_1 \subseteq S_2$. Let $x \in \text{Ult}(A)$ and $y \in \text{Ult}(B)$ such that $x R_{S_2} y$. Then $S_1[x] \subseteq S_2[x] \subseteq y$, and so $x R_{S_1} y$. Thus, $R_{S_2} \subseteq R_{S_1}$. Conversely, assume that $S_1 \not\subseteq S_2$. Let $a \in A$ and $b \in B$ be such that $a S_1 b$ but $a \not S_2 b$. Since S_2 is a subordination, $S_2^{-1}[b]$ is an ideal of A such that $a \notin S_2^{-1}[b]$. By the ultrafilter lemma, there is $x \in \text{Ult}(A)$ such that $a \in x$ and $x \cap S_2^{-1}[b] = \emptyset$. Therefore, $b \notin S_2[x]$ and $S_2[x]$ is a filter of B . Thus, there is $y \in \text{Ult}(B)$ such that $b \in y$ and $S_2[x] \subseteq y$. Since $a \in x$, $b \in y$, and $a S_1 b$, we have $S_1[x] \not\subseteq y$. Consequently, $x R_{S_2} y$ but $x \not R_{S_1} y$, which implies that $R_{S_2} \not\subseteq R_{S_1}$. This shows that the bijections between the hom-sets preserve and reflect the orders.

Let $R: X \rightarrow Y$ be a morphism in Stone^R , $U \in \text{Clop}(X)$, and $V \in \text{Clop}(Y)$. Then $R^\sim[V] \subseteq U$ iff $R[X \setminus U] \subseteq Y \setminus V$. Therefore, $S_{R^\sim} = \widehat{S}_R$. Also, let $S: A \rightarrow B$ be a morphism in BA^S , $x \in \text{Ult}(A)$, and $y \in \text{Ult}(B)$. Then $\widehat{S}[y] \subseteq x$ iff $S[x] \subseteq y$. Thus, $R_{\widehat{S}} = (R_S)^\sim$. Thus, the functors Clop and Ult commute with the daggers on Stone^R and BA^S .

Since Stone^R is an allegory and the functors Clop and Ult are quasi-inverses of each other that preserve and reflect the orders on the hom-sets and commute with the daggers, it follows that BA^S is also an allegory and Clop and Ult are morphisms of allegories. Thus, Stone^R and BA^S are equivalent as allegories. \square

Remark 2.15. Each hom-set $\text{hom}_{\text{Stone}^R}(X, Y)$ is a complete lattice because it is the set of closed subsets of $X \times Y$. Thus, Theorem 2.14 implies that each hom-set $\text{hom}_{\text{BA}^S}(A, B)$ is also a complete lattice. In [1] we give an explicit description of meets and joins of subordinations.

Remark 2.16. A closed relation $R: X \rightarrow Y$ is an isomorphism in Stone^R iff it is a homeomorphism. To see this, the inverse of R in Stone^R is a closed relation $Q: Y \rightarrow X$ such that $Q \circ R = \text{id}_X$ and $R \circ Q = \text{id}_Y$. Since id_X and id_Y are identities, R is a bijective function and Q is its inverse. Because a function that is closed as a relation is a continuous function, we conclude that R is a homeomorphism. Therefore, by Corollary 2.6, two boolean algebras are isomorphic in BA^S iff they are isomorphic as boolean algebras.

3. Further lifting to closed relations between compact Hausdorff spaces

The goal of this section is to lift the equivalence between Stone^R and BA^S to the category KHaus^R of compact Hausdorff spaces and closed relations. Guided by the fact that KHaus is the exact completion of Stone (see, e.g., [30, Thm. 8.1]), we think of compact Hausdorff spaces as the quotients of Stone spaces by closed equivalence relations. We will obtain a category equivalent to KHaus^R by splitting closed equivalence relations in Stone^R , a construction that we describe in the language of allegories.

Closed equivalence relations on Stone spaces and their corresponding subordinations on boolean algebras are instances of the notion of an equivalence in an allegory.

Definition 3.1. [14, p. 198] Let C be an object of an allegory. A morphism $f: C \rightarrow C$ is called an *equivalence* if $\text{id}_C \leq f$ (*reflexivity*), $f = f^\dagger$ (*symmetry*), and $ff \leq f$ (*transitivity*).

It is immediate to see that a closed relation R on a Stone space is reflexive, symmetric, and transitive as a morphism in Stone^R iff it is reflexive, symmetric, and transitive as a relation. Thus, R is an equivalence in Stone^R iff it is a closed equivalence relation. We have a similar result for subordinations.

Lemma 3.2. A subordination $S: B \rightarrow B$ is an equivalence in BA^S iff it satisfies the following conditions:

- (S5) $a S b$ implies $a \leq b$.
- (S6) $a S b$ implies $\neg b S \neg a$.
- (S7) $a S b$ implies there is $c \in B$ such that $a S c$ and $c S b$.

Proof. Since the hom-sets of BA^S are ordered by reverse inclusion, a subordination S on B is a reflexive morphism in BA^S iff S is contained in the partial order \leq on B , which holds iff S satisfies (S5). Since $S = \widehat{S}$ iff $S \subseteq \widehat{S}$, it follows that S is a symmetric morphism in BA^S iff S satisfies (S6). Finally, S is a transitive morphism in BA^S iff $S \subseteq S \circ S$, which means that (S7) holds. Thus, S is an equivalence in BA^S iff it satisfies conditions (S5)–(S7). \square

Definition 3.3. An *S5-subordination* is a subordination $S: B \rightarrow B$ satisfying conditions (S5)–(S7).

Remark 3.4. The terminology in the above definition is motivated by the connection to the modal logic S5, which is the logic of relational structures (X, R) such that R is an equivalence relation (see, e.g., [6, Table 4.1]).

Definition 3.5. Let KHaus^R be the category of compact Hausdorff spaces and closed relations between them. Identity morphisms in KHaus^R are identity relations and composition in KHaus^R is relation composition.

Clearly Stone^R is a full subcategory of KHaus^R . By arguing as we did in Theorem 2.14 for Stone^R , we obtain:

Lemma 3.6. KHaus^R is an allegory.

We will show that KHaus^R is equivalent to the allegory obtained by splitting the equivalences in Stone^R .

Definition 3.7. [14, p. 15] Let \mathcal{E} be a class of idempotent morphisms of a category \mathbf{C} , where $e: C \rightarrow C$ is *idempotent* if $ee = e$. The category $\text{Split}(\mathcal{E})$, obtained by *splitting* \mathcal{E} , is defined as follows. The objects of $\text{Split}(\mathcal{E})$ are pairs (C, e) where $C \in \mathbf{C}$ and $e \in \mathcal{E}$ with $e: C \rightarrow C$. A morphism $f: (C, e) \rightarrow (C', e')$ in $\text{Split}(\mathcal{E})$ is a morphism $f: C \rightarrow C'$ in \mathbf{C} such that $fe = f = e'f$.

$$\begin{array}{ccc}
 C & \xrightarrow{f} & C' \\
 e \downarrow & \searrow f & \downarrow e' \\
 C & \xrightarrow{f} & C'
 \end{array}$$

When \mathcal{E} is the class of all idempotents of \mathbf{C} , then $\text{Split}(\mathcal{E})$ is called the *Karoubi envelope* or *Cauchy completion* of \mathbf{C} .

Proposition 3.8. [14, p. 211] *Let \mathbf{A} be an allegory and \mathcal{E} a class of symmetric idempotent morphisms of \mathbf{A} . Then $\text{Split}(\mathcal{E})$ inherits the structure of an allegory from \mathbf{A} . In particular, if $\text{Eq}_{\mathbf{A}}$ is the class of all equivalences in \mathbf{A} , then $\text{Split}(\text{Eq}_{\mathbf{A}})$ is an allegory.*

Notation 3.9.

1. To simplify notation, we denote $\text{Split}(\text{Eq}_{\text{Stone}^R})$ by StoneE^R . Thus, an object of StoneE^R is a pair (X, E) where X is a Stone space and E is a closed equivalence relation on X . We call such pairs *S5-subordination spaces*. A morphism in StoneE^R between (X, E) and (X', E') is a closed relation $R: X \rightarrow X'$ satisfying $R \circ E = R = E' \circ R$. We call such relations *compatible closed relations*. The composition of compatible closed relations in StoneE^R is the standard relation composition and the identity on (X, E) is the relation E .

2. Similarly, we denote $\text{Split}(\text{Eq}_{\text{BA}^S})$ by SubS5^S , so an object of SubS5^S is a pair (B, S) where B is a boolean algebra and S is an $S5$ -subordination on B . We call such pairs $S5$ -subordination algebras.¹ A morphism in SubS5^S between (B, S) and (B', S') is a subordination relation $T: B \rightarrow B'$ satisfying $T \circ S = T = S' \circ T$. We call such subordinations *compatible subordinations*. The composition of compatible subordinations in SubS5^S is the standard relation composition and the identity on (B, S) is the subordination S .

By Proposition 3.8, StoneE^R is an allegory with the hom-sets ordered by inclusion and the dagger defined as in Stone^R . In addition, SubS5^S is an allegory with the hom-sets ordered by reverse inclusion and the dagger defined as in BA^S . The following is now immediate because Stone^R and BA^S are equivalent as allegories (see Theorem 2.14):

Theorem 3.10. StoneE^R and SubS5^S are equivalent allegories.

Remark 3.11. The functors establishing the equivalence of allegories are obtained by lifting the functors Clop and Ult yielding the equivalence between Stone^R and BA^S . An $S5$ -subordination space (X, E) is mapped to $(\text{Clop}(X), S_E)$ and a compatible closed relation R to S_R . An $S5$ -subordination algebra (B, S) is mapped to $(\text{Ult}(B), R_S)$ and a compatible subordination T to R_T .

We next show that KHaus^R and StoneE^R are equivalent as allegories. We first observe that each compact Hausdorff space is homeomorphic to a quotient of a Stone space. Indeed, any compact Hausdorff space is a quotient of its Gleason cover (see, e.g., [20, pp. 107–108]). The Gleason cover is not the only possible option. In fact, it follows from the universal mapping property of Stone-Čech compactifications (see, e.g., [13, Thm. 3.6.1]) that any compact Hausdorff space X is homeomorphic to a quotient of the Stone-Čech compactification of the discrete space with the same underlying set as X . There are many other options as well. Clearly, if a compact Hausdorff space X is homeomorphic to a quotient Y/E of a Stone space Y , then E is necessarily closed (see, e.g., [13, Thm. 3.2.11]). We will use this in the proof of the next theorem.

Theorem 3.12. KHaus^R and StoneE^R are equivalent allegories.

Proof. Define $\mathcal{Q}: \text{StoneE}^R \rightarrow \text{KHaus}^R$ as follows: for each $(X, E) \in \text{StoneE}^R$ let $\mathcal{Q}(X, E)$ be the quotient space X/E , which is well known to be compact Hausdorff (see, e.g., [13, Thm. 3.2.11]); and for each morphism $R: (X, E) \rightarrow (X', E')$ in StoneE^R let $\mathcal{Q}(R): \mathcal{Q}(X, E) \rightarrow \mathcal{Q}(X', E')$ be the morphism in KHaus^R given by $\mathcal{Q}(R) = \pi' \circ R \circ \pi^\smile$, where $\pi: X \rightarrow X/E$ and $\pi': X' \rightarrow X'/E'$ are the projections onto the quotients.

We show that \mathcal{Q} preserves identities and compositions. Note that if $\pi: X \rightarrow X/E$ is the projection onto the quotient, then $E = \pi^\smile \circ \pi$ and $\pi \circ \pi^\smile = \text{id}_{X/E}$. Therefore,

$$\mathcal{Q}(\text{id}_{(X,E)}) = \mathcal{Q}(E) = \pi \circ E \circ \pi^\smile = \pi \circ \pi^\smile \circ \pi \circ \pi^\smile = \text{id}_{X/E}.$$

Let $R_1: (X_1, E_1) \rightarrow (X_2, E_2)$ and $R_2: (X_2, E_2) \rightarrow (X_3, E_3)$ be compatible closed relations. Then

$$\begin{aligned} \mathcal{Q}(R_2) \circ \mathcal{Q}(R_1) &= (\pi_3 \circ R_2 \circ \pi_2^\smile) \circ (\pi_2 \circ R_1 \circ \pi_1^\smile) = \pi_3 \circ R_2 \circ E_2 \circ R_1 \circ \pi_1^\smile \\ &= \pi_3 \circ R_2 \circ R_1 \circ \pi_1^\smile = \mathcal{Q}(R_2 \circ R_1). \end{aligned}$$

We prove that \mathcal{Q} is full, faithful, and essentially surjective. Let $R': X'/E' \rightarrow X'/E'$ be a closed relation. Then $R = (\pi')^\smile \circ R' \circ \pi$ is a compatible closed relation such that $\mathcal{Q}(R) = R'$ because

$$\mathcal{Q}(R) = \mathcal{Q}((\pi')^\smile \circ R' \circ \pi) = \pi' \circ (\pi')^\smile \circ R' \circ \pi \circ \pi^\smile = \text{id}_{X'/E'} \circ R' \circ \text{id}_{X/E} = R'.$$

¹ Celani [10] calls these algebras *quasi-monadic algebras* because they generalize Halmos' monadic algebras [15].

Therefore, \mathcal{Q} is full. If $R: (X, E) \rightarrow (X', E')$ is a compatible closed relation, then $R = (\pi')^\smile \circ \mathcal{Q}(R) \circ \pi$ because

$$(\pi')^\smile \circ \mathcal{Q}(R) \circ \pi = (\pi')^\smile \circ \pi' \circ R \circ \pi^\smile \circ \pi = E' \circ R \circ E = R.$$

Thus, if $R, R': (X, E) \rightarrow (X', E')$ are two compatible closed relations such that $\mathcal{Q}(R) = \mathcal{Q}(R')$, then $R = (\pi')^\smile \circ \mathcal{Q}(R) \circ \pi = (\pi')^\smile \circ \mathcal{Q}(R') \circ \pi = R'$. This shows that \mathcal{Q} is faithful. As we pointed out before the theorem, if Y is a compact Hausdorff space, then there exists $(X, E) \in \text{StoneE}^{\mathbb{R}}$ such that Y is homeomorphic to X/E . It follows that \mathcal{Q} is essentially surjective, hence an equivalence [29, p. 93]. It remains to show that \mathcal{Q} is a morphism of allegories.

Let $R_1, R_2: (X, E) \rightarrow (X', E')$ be compatible closed relations. If $R_1 \subseteq R_2$, then

$$\mathcal{Q}(R_1) = \pi' \circ R_1 \circ \pi^\smile \subseteq \pi' \circ R_2 \circ \pi^\smile = \mathcal{Q}(R_2).$$

Conversely, if $\mathcal{Q}(R_1) \subseteq \mathcal{Q}(R_2)$, then

$$R_1 = (\pi')^\smile \circ \mathcal{Q}(R_1) \circ \pi \subseteq (\pi')^\smile \circ \mathcal{Q}(R_2) \circ \pi = R_2.$$

Therefore, \mathcal{Q} preserves and reflects the inclusion of relations, and so it yields an order-isomorphism on hom-sets. Finally, since $(-)^{\smile}$ is a contravariant endofunctor, $\mathcal{Q}(R^\smile) = \pi \circ R^\smile \circ (\pi')^\smile$ is the converse relation of $\pi' \circ R \circ \pi^\smile = \mathcal{Q}(R)$. Thus, \mathcal{Q} commutes with the dagger, and hence is a morphism of allegories. \square

Remark 3.13. The following is a more categorical approach to Theorem 3.12. If \mathbf{A} is an allegory, then a morphism $f: C \rightarrow D$ in \mathbf{A} is a *map* if $f^\dagger f \leq \text{id}_C$ and $\text{id}_D \leq f f^\dagger$ (see [14, p. 199]). The wide subcategory of \mathbf{A} whose morphisms are maps is denoted by $\text{Map}(\mathbf{A})$.

If \mathbf{C} is a regular category, then we can define the allegory $\text{Rel}(\mathbf{C})$ whose objects are the same as \mathbf{C} and whose morphisms are internal relations, where an *internal relation* $R: C \rightarrow D$ is a subobject of $C \times D$ (see [21, Sec. A3.1]). Both KHaus and Stone are regular categories and their internal relations correspond to closed relations. Therefore, $\text{KHaus}^{\mathbb{R}}$ and $\text{Rel}(\text{KHaus})$ are isomorphic allegories, and so are $\text{Stone}^{\mathbb{R}}$ and $\text{Rel}(\text{Stone})$. Thus, the allegories $\text{Split}(\text{Eq}_{\text{Rel}(\text{Stone})})$ and $\text{Split}(\text{Eq}_{\text{Stone}^{\mathbb{R}}})$ are isomorphic.

If \mathbf{C} is a regular category, then $\text{Map}(\text{Split}(\text{Eq}_{\text{Rel}(\mathbf{C})}))$ is the effective reflection of \mathbf{C} in the category of regular categories [14, p. 213]. This is also known as the exact completion or ex/reg completion.² Roughly speaking, the exact completion is obtained by closing under quotients. Since KHaus is the exact completion of Stone (see, e.g., [30, Thm. 8.1]), it follows that KHaus is equivalent to $\text{Map}(\text{Split}(\text{Eq}_{\text{Rel}(\text{Stone})}))$. Therefore, $\text{Rel}(\text{KHaus})$ and $\text{Rel}(\text{Map}(\text{Split}(\text{Eq}_{\text{Rel}(\text{Stone})})))$ are equivalent allegories (see, e.g., [14, p. 204]). By applying [21, A3.2.10, A3.3.4(i), A3.3.9(ii)], the allegories $\text{Rel}(\text{Map}(\text{Split}(\text{Eq}_{\text{Rel}(\text{Stone})})))$ and $\text{Split}(\text{Eq}_{\text{Rel}(\text{Stone})})$ are isomorphic. Thus, we have the following chain of equivalences of allegories:

$$\text{KHaus}^{\mathbb{R}} \cong \text{Rel}(\text{KHaus}) \simeq \text{Rel}(\text{Map}(\text{Split}(\text{Eq}_{\text{Rel}(\text{Stone})}))) \cong \text{Split}(\text{Eq}_{\text{Rel}(\text{Stone})}) \cong \text{Split}(\text{Eq}_{\text{Stone}^{\mathbb{R}}}) = \text{StoneE}^{\mathbb{R}},$$

where \cong stands for isomorphism and \simeq for equivalence of allegories.

As an immediate consequence of Theorems 3.10 and 3.12 we obtain:

Corollary 3.14. $\text{KHaus}^{\mathbb{R}}$, $\text{StoneE}^{\mathbb{R}}$, and $\text{SubS5}^{\mathbb{S}}$ are equivalent as allegories.

We conclude this section by characterizing isomorphisms in $\text{StoneE}^{\mathbb{R}}$ and $\text{SubS5}^{\mathbb{S}}$.

² This should not be confused with the effective reflection of \mathbf{C} in the category of lex categories, also known as ex/lex completion (see, e.g., <https://ncatlab.org/nlab/show/regular+and+exact+completions>).

Remark 3.15.

- (1) By Theorem 3.12, \mathcal{Q} is full and faithful. Therefore, a morphism $R: (X_1, E_1) \rightarrow (X_2, E_2)$ in StoneE^R is an isomorphism iff $\mathcal{Q}(R): X_1/E_1 \rightarrow X_2/E_2$ is a homeomorphism. Thus, (X_1, E_1) and (X_2, E_2) are isomorphic in StoneE^R iff X_1/E_1 and X_2/E_2 are homeomorphic.
- (2) Let $R: (X_1, E_1) \rightarrow (X_2, E_2)$ be an isomorphism in StoneE^R . Since inverses in an allegory are daggers (see [14, p. 199]) and $R^\dagger = R^\smile$ in StoneE^R (see Lemma 2.13 and the paragraph before Theorem 3.10), it follows that the inverse of R in StoneE^R is R^\smile .
- (3) Not every isomorphism in StoneE^R is a structure-preserving bijection; two objects (X_1, E_1) and (X_2, E_2) may be isomorphic even if X_1 and X_2 do not share the same cardinality (see (1)). However, the converse is true. If $f: X_1 \rightarrow X_2$ is a structure-preserving bijection, then f is a homeomorphism such that $x E_1 y$ iff $f(x) E_2 f(y)$. Therefore, f and $f^{-1} = f^\smile$ are compatible closed relations, and hence f is an isomorphism in StoneE^R .
- (4) A similar result is true for SubS5^S : two objects (B_1, S_1) and (B_2, S_2) may be isomorphic even if B_1 and B_2 do not share the same cardinality. However, every structure-preserving bijection gives rise to an isomorphism in SubS5^S . Indeed, let $(B_1, S_1), (B_2, S_2) \in \text{SubS5}^S$ and suppose there is a boolean isomorphism $f: B_1 \rightarrow B_2$ such that for all $a, b \in B_1$ we have $a S_1 b$ iff $f(a) S_2 f(b)$. Similarly to Remark 2.16, an isomorphism between (B_1, S_1) and (B_2, S_2) is given by the relation $T: B_1 \rightarrow B_2$ defined by

$$a T b \iff f(a) S_2 b \iff a S_1 f^{-1}(b),$$

and its inverse in SubS5^S is the relation $Q: B_2 \rightarrow B_1$ defined by

$$b Q a \iff b S_2 f(a) \iff f^{-1}(b) S_1 a.$$

By (S6),

$$b \widehat{T} a \iff \neg a T \neg b \iff f(\neg a) S_2 \neg b \iff b S_2 f(a) \iff b Q a.$$

Thus, every structure-preserving bijection $f: B_1 \rightarrow B_2$ gives rise to an isomorphism $T: B_1 \rightarrow B_2$ in SubS5^S whose inverse is $\widehat{T}: B_2 \rightarrow B_1$.

4. De Vries algebras and Gleason spaces

As we saw in Section 3, KHaus^R is equivalent to both StoneE^R and SubS5^S . In this section we introduce two important full subcategories Gle^R of StoneE^R and DeV^S of SubS5^S . The objects of Gle^R are Gleason spaces and those of DeV^S are de Vries algebras. We prove that Gle^R and DeV^S are also equivalent to KHaus^R . In Section 5 we will see that, unlike StoneE^R and SubS5^S , isomorphisms in Gle^R and DeV^S are structure-preserving bijections.

For a compact Hausdorff space X , let $g_X: \widehat{X} \rightarrow X$ be the Gleason cover of X (see, e.g., [20, pp. 107–108]). Let E be the equivalence relation on \widehat{X} given by

$$x E y \iff g_X(x) = g_X(y). \tag{*}$$

Since X is homeomorphic to the quotient space \widehat{X}/E , in a certain sense, we can identify X with the pair (\widehat{X}, E) . This was made precise in [5], where an equivalence is exhibited between KHaus^R and the full subcategory of StoneE^R that we next define.

Definition 4.1.

1. [13, p. 368] A topological space is *extremally disconnected* if the closure of each open set is open.
2. [4, Def. 6.6] A *Gleason space* is an object (X, E) of $\text{StoneE}^{\mathbf{R}}$ such that X is extremally disconnected and E is *irreducible* (i.e., if F is a proper closed subset of X , then so is $E[F]$).
3. [5, Def. 3.6] Let $\text{Gle}^{\mathbf{R}}$ be the full subcategory of $\text{StoneE}^{\mathbf{R}}$ whose objects are Gleason spaces.

We next define the category $\text{DeV}^{\mathbf{S}}$.

Definition 4.2.

1. [3, Def. 3.2] A *de Vries algebra* is an $\mathbf{S5}$ -subordination algebra (B, S) such that B is a complete boolean algebra and S satisfies the following axiom:

$$(S8) \text{ If } a \neq 0, \text{ then there is } b \neq 0 \text{ such that } b S a.$$

2. Let $\text{DeV}^{\mathbf{S}}$ be the full subcategory of $\text{SubS5}^{\mathbf{S}}$ consisting of de Vries algebras.

Remark 4.3. A relation S on a boolean algebra B satisfying (S1)–(S8) is called a *de Vries proximity* on B and is usually denoted by \prec .

Lemma 4.4. $\text{Gle}^{\mathbf{R}}$ and $\text{DeV}^{\mathbf{S}}$ are allegories.

Proof. By [21, Examples A.3.2.2(c)], every full subcategory of an allegory inherits the allegory structure. \square

Theorem 4.5. *The equivalence between $\text{StoneE}^{\mathbf{R}}$ and $\text{SubS5}^{\mathbf{S}}$ restricts to an equivalence of allegories between $\text{Gle}^{\mathbf{R}}$ and $\text{DeV}^{\mathbf{S}}$.*

Proof. Let $B \in \mathbf{BA}$. By Stone duality, B is complete iff $\text{Ult}(B)$ is extremally disconnected. Let S be an $\mathbf{S5}$ -subordination on B and R_S the corresponding closed equivalence relation on $\text{Ult}(B)$. By [4, Lem. 6.3], R_S is irreducible iff S satisfies (S8). Thus, the restriction of $\text{Clop}: \text{StoneE}^{\mathbf{R}} \rightarrow \text{SubS5}^{\mathbf{S}}$ and $\text{Ult}: \text{SubS5}^{\mathbf{S}} \rightarrow \text{StoneE}^{\mathbf{R}}$ gives the desired equivalence. \square


For a compact Hausdorff space X , let $\mathcal{G}(X) = (\widehat{X}, E)$, where \widehat{X} is the Gleason cover of X and E the corresponding equivalence relation defined in (*). For a closed relation $R: X \rightarrow X'$, let $\mathcal{G}(R): \mathcal{G}(X) \rightarrow \mathcal{G}(X')$ be given by $\mathcal{G}(R) = (g_{X'})^\vee \circ R \circ g_X$. This defines a functor $\mathcal{G}: \mathbf{KHaus}^{\mathbf{R}} \rightarrow \text{StoneE}^{\mathbf{R}}$, which is a quasi-inverse of \mathcal{Q} :

Theorem 4.6. *Each object (X, E) of $\text{StoneE}^{\mathbf{R}}$ is isomorphic in $\text{StoneE}^{\mathbf{R}}$ to the Gleason space $\mathcal{G}(X/E)$. Thus, \mathcal{G} is a quasi-inverse of \mathcal{Q} and the inclusion of $\text{Gle}^{\mathbf{R}}$ into $\text{StoneE}^{\mathbf{R}}$ is an equivalence of allegories.*

Proof. Let $(X, E) \in \text{StoneE}^{\mathbf{R}}$ and let $(X', E') = \mathcal{G}(X/E)$. Then $(X', E') \in \text{Gle}^{\mathbf{R}}$ and (X, E) is isomorphic to (X', E') by Remark 3.15(1). Thus, the inclusion of $\text{Gle}^{\mathbf{R}}$ into $\text{StoneE}^{\mathbf{R}}$ is full, faithful, and essentially surjective, hence an equivalence of allegories (because it is a morphism of allegories by [21, Examples A.3.2.2(c)]). Since any Gleason space is an object of $\text{StoneE}^{\mathbf{R}}$, what we observed above implies that any $(X, E) \in \text{Gle}^{\mathbf{R}}$ is isomorphic to $\mathcal{G}(X/E) = \mathcal{G}(\mathcal{Q}(X, E))$. Straightforward computations show that the isomorphisms $(X, E) \cong \mathcal{G}(\mathcal{Q}(X, E))$ for $(X, E) \in \text{Gle}^{\mathbf{R}}$ and $\mathcal{Q}(\mathcal{G}(X)) \cong X$ for $X \in \mathbf{KHaus}^{\mathbf{R}}$ yield natural isomorphisms $\text{id}_{\text{Gle}^{\mathbf{R}}} \cong \mathcal{G} \circ \mathcal{Q}$ and $\text{id}_{\mathbf{KHaus}^{\mathbf{R}}} \cong \mathcal{Q} \circ \mathcal{G}$. Thus, \mathcal{G} is a quasi-inverse of \mathcal{Q} . \square

Corollary 4.7. \mathbf{KHaus}^R , \mathbf{Gle}^R , and \mathbf{DeV}^S are equivalent allegories.

As a consequence, we arrive at the following diagram of equivalences of allegories that commutes up to natural isomorphisms.



Remark 4.8. That \mathbf{KHaus}^R and \mathbf{Gle}^R are equivalent categories was first established in [5, Thm. 3.13]. By Corollary 4.7, \mathbf{KHaus}^R is equivalent to \mathbf{DeV}^S . This resolves the problem raised in [5, Rem. 3.14] to find a generalization of the category of de Vries algebras that is (dually) equivalent to \mathbf{KHaus}^R . The key is to work with relations between de Vries algebras instead of functions.

We conclude this section by providing an explicit description of the functor from \mathbf{KHaus}^R to \mathbf{DeV}^S yielding the equivalence, which is a direct generalization of the regular open functor of de Vries duality.

For a compact Hausdorff space X , let $\mathbf{RO}(X)$ be the complete boolean algebra of regular open subsets of X . We recall that joins in $\mathbf{RO}(X)$ are computed by $\bigvee_i U_i = \text{int}(\text{cl}(\bigcup_i U_i))$ and the negation is computed by $\neg U = \text{int}(X \setminus U)$. Similarly, if $\mathbf{RC}(X)$ is the complete boolean algebra of regular closed subsets of X , then meets in $\mathbf{RC}(X)$ are computed by $\bigwedge_i F_i = \text{cl}(\text{int}(\bigcap_i F_i))$ and the negation is computed by $\neg F = \text{cl}(X \setminus F)$.

Parts of the next lemma are known, but it is convenient to collect all the relevant isomorphisms together.

Lemma 4.9. Let X be a compact Hausdorff space and $g_X: \widehat{X} \rightarrow X$ its Gleason cover. The boolean algebras $\mathbf{RO}(X)$, $\mathbf{RC}(X)$, and $\mathbf{Clop}(\widehat{X})$ are isomorphic, with the isomorphisms given by

$$\begin{array}{ccc}
 \mathbf{RO}(X) \longleftrightarrow \mathbf{RC}(X) & \mathbf{Clop}(\widehat{X}) \longleftrightarrow \mathbf{RC}(X) & \mathbf{Clop}(\widehat{X}) \longleftrightarrow \mathbf{RO}(X) \\
 U \mapsto \text{cl}(U) & V \mapsto g_X[V] & V \mapsto \text{int}(g_X[V]) \\
 \text{int}(F) \longleftarrow F, & \text{cl}(g_X^{-1}[\text{int}(F)]) \longleftarrow F, & \text{cl}(g_X^{-1}[U]) \longleftarrow U.
 \end{array}$$

Proof. (Sketch) That $\text{cl}: \mathbf{RO}(X) \rightarrow \mathbf{RC}(X)$ and $\text{int}: \mathbf{RC}(X) \rightarrow \mathbf{RO}(X)$ are inverses of each other is an immediate consequence of the definition of regular open and regular closed sets.

Since $g_X: \widehat{X} \rightarrow X$ is an onto irreducible map, the direct image function $g_X[-]: \mathbf{RC}(\widehat{X}) \rightarrow \mathbf{RC}(X)$ that maps $F \in \mathbf{RC}(\widehat{X})$ to $g_X[F]$ is a boolean isomorphism (see [32, p. 454]). We note that $\mathbf{RC}(\widehat{X}) = \mathbf{Clop}(\widehat{X})$ because \widehat{X} is extremally disconnected. It follows from the proof given in [32, p. 454] that the inverse of $g_X[-]$ maps each $F \in \mathbf{RC}(X)$ to $\text{cl}(g_X^{-1}[\text{int}(F)])$.

By composing the isomorphism between $\mathbf{RO}(X)$ and $\mathbf{RC}(X)$ with the isomorphism between $\mathbf{RC}(X)$ and $\mathbf{Clop}(\widehat{X})$, we obtain the isomorphism between $\mathbf{RO}(X)$ and $\mathbf{Clop}(\widehat{X})$ described in the statement. \square

Definition 4.10. Let $\mathcal{D}: \mathbf{KHaus}^R \rightarrow \mathbf{DeV}^S$ be defined by associating with each compact Hausdorff space X the de Vries algebra $(\mathbf{RO}(X), \prec)$ of regular open subsets of X (with the de Vries proximity defined by $U \prec V$ iff $\text{cl}(U) \subseteq V$) and with each closed relation $R: X \rightarrow X'$ the subordination $\mathcal{D}(R): \mathbf{RO}(X) \rightarrow \mathbf{RO}(X')$ given by

$$U \mathcal{D}(R) V \iff R[\text{cl}(U)] \subseteq V.$$

It is straightforward to see that \mathcal{D} is a well-defined functor. Recalling the functors $\mathcal{G}: \text{KHaus}^{\mathbb{R}} \rightarrow \text{Gle}^{\mathbb{R}}$ and $\text{Clop}: \text{Gle}^{\mathbb{R}} \rightarrow \text{DeV}^{\mathbb{S}}$, we have:

Theorem 4.11. *The functor \mathcal{D} is naturally isomorphic to $\text{Clop} \circ \mathcal{G}$.*

Proof. By Lemma 4.9, for every compact Hausdorff space X , the function $\eta_X: \text{RO}(X) \rightarrow \text{Clop}(\mathcal{G}X)$ that maps U to $\text{cl}(g_X^{-1}[U])$ is a boolean isomorphism.

If $R: X \rightarrow X'$ is a closed relation, then $\mathcal{G}R: \mathcal{G}X \rightarrow \mathcal{G}X'$ is given by $\mathcal{G}R = (g_{X'})^\smile \circ R \circ g_X$. Moreover, $\text{Clop}(\mathcal{G}R)$ is the $\text{DeV}^{\mathbb{S}}$ -morphism $S_{\mathcal{G}R}: \text{Clop}(\mathcal{G}X) \rightarrow \text{Clop}(\mathcal{G}X')$ given by $U S_{\mathcal{G}R} V$ iff $(\mathcal{G}R)[U] \subseteq V$. Consequently, for all $U \in \text{RO}(X)$ and $V \in \text{RO}(X')$ we have

$$\begin{aligned} \eta_X(U) (\text{Clop } \mathcal{G}R) \eta_{X'}(V) &\iff \mathcal{G}R[\eta_X(U)] \subseteq \eta_{X'}(V) \\ &\iff \mathcal{G}R[\eta_X(U)] \cap (\mathcal{G}X' \setminus \eta_{X'}(V)) = \emptyset \\ &\iff g_{X'}^{-1}[R[g_X[\eta_X(U)]]] \cap (\mathcal{G}X' \setminus \eta_{X'}(V)) = \emptyset \\ &\iff R[g_X[\eta_X(U)]] \cap g_{X'}[\mathcal{G}X' \setminus \eta_{X'}(V)] = \emptyset. \end{aligned}$$

Since $U \in \text{RO}(X)$ and $\text{cl}(U) \in \text{RC}(X)$, it follows from Lemma 4.9 that

$$R[g_X[\eta_X(U)]] = R[g_X[\text{cl}(g_X^{-1}[\text{int}(\text{cl}(U))]]] = R[\text{cl}(U)].$$

Because $\eta_{X'}$ is an isomorphism,

$$g_{X'}[\mathcal{G}X' \setminus \eta_{X'}(V)] = g_{X'}[\eta_{X'}(\text{int}(X' \setminus V))] = g_{X'}[\text{cl}(g_{X'}^{-1}[\text{int}(X' \setminus V)])] = X' \setminus V,$$

where the last equality follows from Lemma 4.9 since $X' \setminus V \in \text{RC}(X')$. Therefore,

$$\eta_X(U) (\text{Clop } \mathcal{G}R) \eta_{X'}(V) \iff R[\text{cl}(U)] \cap (X' \setminus V) = \emptyset \iff R[\text{cl}(U)] \subseteq V.$$

When $X = X'$, taking R to be the identity 1_X on X , we get

$$\eta_X(U) (\text{Clop } \mathcal{G}1_X) \eta_X(V) \iff U \prec V$$

for all $U, V \in \text{RO}(X)$. Thus, the isomorphism $\eta_X: \text{RO}(X) \rightarrow \text{Clop}(\mathcal{G}X)$ is a structure-preserving bijection. By Remark 3.15(4), $\text{RO}(X)$ and $\text{Clop}(\mathcal{G}X)$ are isomorphic in $\text{DeV}^{\mathbb{S}}$, and the isomorphism is given by the relation $S_X: \text{RO}(X) \rightarrow \text{Clop}(\mathcal{G}X)$ defined by

$$U S_X V \iff \eta_X(U) (\text{Clop } \mathcal{G}1_X) V \iff U \prec \eta_X^{-1}(V).$$

To prove naturality, we show that the following diagram commutes for every morphism $R: X \rightarrow X'$ in KHaus .

$$\begin{array}{ccc} \text{RO}(X) & \xrightarrow{S_X} & \text{Clop}(\mathcal{G}X) \\ \mathcal{D}(R) \downarrow & & \downarrow \text{Clop } \mathcal{G}R \\ \text{RO}(X') & \xrightarrow{S_{X'}} & \text{Clop}(\mathcal{G}X') \end{array}$$

Let $U \in \text{RO}(X)$ and $V \in \text{Clop}(\mathcal{G}X')$. Since $\text{Clop } \mathcal{G}R$ and $\mathcal{D}(R)$ are compatible subordinations, we have

$$\begin{aligned}
 U(S_{X'} \circ \mathcal{D}(R)) V &\iff \exists C \in \text{RO}(X') : U \mathcal{D}(R) C S_{X'} V \\
 &\iff \exists C \in \text{RO}(X') : U \mathcal{D}(R) C \prec \eta_{X'}^{-1}(V) \\
 &\iff U \mathcal{D}(R) \eta_{X'}^{-1}(V) \\
 &\iff \eta_X(U) (\text{Clop } \mathcal{G}R) V \\
 &\iff \exists D \in \text{Clop}(\mathcal{G}X) : \eta_X(U) (\text{Clop } \mathcal{G}1_X) D (\text{Clop } \mathcal{G}R) V \\
 &\iff \exists D \in \text{Clop}(\mathcal{G}X) : U S_X D (\text{Clop } \mathcal{G}R) V \\
 &\iff U ((\text{Clop } \mathcal{G}R) \circ S_X) V. \quad \square
 \end{aligned}$$

We thus obtain:

Corollary 4.12. *The functor $\mathcal{D}: \text{KHaus}^R \rightarrow \text{DeV}^S$ is an equivalence of allegories.*

Proof. By Theorem 4.11, \mathcal{D} is naturally isomorphic to the composition $\text{Clop} \circ \mathcal{G}$, each of which is an equivalence of allegories (see Remark 3.11 and Theorem 4.6). \square

5. Isomorphisms in DeV^S and Gle^R

It follows from Remark 3.15 that isomorphisms in SubS5^S and StoneE^R are not structure-preserving bijections. In this section we show that in DeV^S and Gle^R isomorphisms are given by structure-preserving bijections, thus making the latter categories more convenient to work with.

For a subset E of a boolean algebra B , we write $U(E)$ and $L(E)$ for the sets of upper and lower bounds of E , respectively. We will freely use the fact that in a de Vries algebra (A, S) we have $a = \bigwedge S[a] = \bigvee S^{-1}[a]$ for every $a \in A$.

Lemma 5.1. *Let (A, S_A) and (B, S_B) be isomorphic objects in SubS5^S with $T: A \rightarrow B$ and $Q: B \rightarrow A$ inverses of each other. Suppose that (A, S_A) is a de Vries algebra.*

(1) *For all $a_1, a_2 \in A$, we have*

$$a_1 \leq a_2 \iff T[a_1] \supseteq T[a_2] \iff Q^{-1}[a_1] \subseteq Q^{-1}[a_2].$$

(2) *For all $a \in A$, we have*

$$Q^{-1}[a] = S_B^{-1}[L(T[a])] \quad \text{and} \quad T[a] = S_B[U(Q^{-1}[a])].$$

Proof. (1). We only prove $a_1 \leq a_2 \iff T[a_1] \supseteq T[a_2]$; the equivalence $a_1 \leq a_2 \iff Q^{-1}[a_1] \subseteq Q^{-1}[a_2]$ is proved similarly. The left-to-right implication is immediate. For the right-to-left implication, suppose that $T[a_1] \supseteq T[a_2]$. Then $QT[a_1] \supseteq QT[a_2]$. Since T and Q are inverses of each other, $Q \circ T = S_A$, and hence $S_A[a_1] = QT[a_1] \supseteq QT[a_2] = S_A[a_2]$. Thus, $a_1 = \bigwedge S_A[a_1] \leq \bigwedge S_A[a_2] = a_2$ because (A, S_A) is a de Vries algebra.

(2). We only prove $Q^{-1}[a] = S_B^{-1}[L(T[a])]$; the second equality is proved similarly. For the left-to-right inclusion, let $b \in Q^{-1}[a]$, so $b Q a$. Since Q is a compatible subordination, there is $b' \in B$ such that $b S_B b' Q a$. For every $b'' \in T[a]$ we have $b' Q a T b''$, so $b' S_B b''$, and hence $b' \leq b''$ by (S5). Therefore, $b' \in L(T[a])$, and so $b \in S_B^{-1}[L(T[a])]$.

For the right-to-left inclusion, let $b \in S_B^{-1}[L(T[a])]$. Then there is $b' \in L(T[a])$ such that $b S_B b'$. Since Q and T are inverses of each other, there is $a' \in A$ such that $b Q a' T b'$. But then $T[a] \subseteq T[a']$ because for

every $c \in T[a]$, we have $a' T b' \leq c$, and so $a' T c$. Thus, $a' \leq a$ by (1). Consequently, $b Q a' \leq a$, so $b Q a$, and hence $b \in Q^{-1}[a]$. \square

Lemma 5.2. *Let (A, S_A) and (B, S_B) be isomorphic objects in DeV^S with $T: A \rightarrow B$ and $Q: B \rightarrow A$ inverses of each other. For all $a \in A$ and $b \in B$,*

- (1) $T[a] = S_B[\bigvee Q^{-1}[a]] = S_B[\bigwedge T[a]]$;
- (2) $Q[b] = S_A[\bigvee T^{-1}[b]] = S_A[\bigwedge Q[b]]$;
- (3) $T^{-1}[b] = S_A^{-1}[\bigwedge Q[b]] = S_A^{-1}[\bigvee T^{-1}[b]]$;
- (4) $Q^{-1}[a] = S_B^{-1}[\bigwedge T[a]] = S_B^{-1}[\bigvee Q^{-1}[a]]$.

Proof. We only prove the first equality of (4) and the second equality of (1). The rest are proved similarly. To see that $Q^{-1}[a] = S_B^{-1}[\bigwedge T[a]]$, by Lemma 5.1(2) it is sufficient to prove that $S_B^{-1}[L(T[a])] = S_B^{-1}[\bigwedge T[a]]$. But this is obvious because $\bigwedge T[a]$ is the greatest lower bound of $T[a]$.

To see that $S_B[\bigvee Q^{-1}[a]] = S_B[\bigwedge T[a]]$, since (B, S_B) is a de Vries algebra, for each $c \in B$, we have $c = \bigvee S_B^{-1}[c]$. Thus, by the first equality of (4),

$$S_B[\bigvee Q^{-1}[a]] = S_B[\bigvee S_B^{-1}[\bigwedge T[a]]] = S_B[\bigwedge T[a]]. \quad \square$$

Lemma 5.3. *Let (A, S_A) and (B, S_B) be isomorphic objects in DeV^S with $T: A \rightarrow B$ and $Q: B \rightarrow A$ inverses of each other. Define $f: A \rightarrow B$ and $g: B \rightarrow A$ by*

$$f(a) = \bigwedge T[a] \quad \text{and} \quad g(b) = \bigwedge Q[b].$$

Then f and g are structure-preserving bijections that are inverses of each other. Moreover, for each $a \in A$ and $b \in B$ we have

$$a T b \text{ iff } f(a) S_B b \quad \text{and} \quad b Q a \text{ iff } g(b) S_A a.$$

Proof. Let $a \in A$. Since Q is a compatible subordination and T is its inverse, by Lemma 5.2(1) we have

$$Q[f(a)] = Q[\bigwedge T[a]] = QS_B[\bigwedge T[a]] = QT[a] = S_A[a].$$

Thus, since S_A is a de Vries proximity,

$$gf(a) = \bigwedge Q[f(a)] = \bigwedge S_A[a] = a.$$

A similar proof yields that $fg(b) = b$ for each $b \in B$. It is an immediate consequence of Lemma 5.1 that f and g are order-preserving. Therefore, f, g are boolean isomorphisms that are inverses of each other.

We next show that f preserves and reflects S_A . That g preserves and reflects S_B is proved similarly. Let $a, a' \in A$. As we saw above, $Q[f(a)] = S_A[a]$. Also, by Lemma 5.2(4), $S_B^{-1}[f(a')] = S_B^{-1}[\bigwedge T[a']] = Q^{-1}[a']$. Therefore,

$$\begin{aligned} f(a) S_B f(a') &\iff f(a) \in S_B^{-1}[f(a')] \iff f(a) \in Q^{-1}[a'] \iff a' \in Q[f(a)] \\ &\iff a' \in S_A[a] \iff a S_A a'. \end{aligned}$$

Finally, let $a \in A$ and $b \in B$. To see that $a T b$ iff $f(a) S_B b$, it is sufficient to observe that Lemma 5.2(1) implies $S_B[f(a)] = S_B[\bigwedge T[a]] = T[a]$. A similar reasoning gives that $b Q a$ iff $g(b) S_A a$. \square

As an immediate consequence of Remark 3.15 and Lemma 5.3 we obtain:

Theorem 5.4. *Isomorphisms in DeV^S are given by structure-preserving bijections.*

Remark 5.5. The above theorem generalizes a similar result for the category DeV of de Vries algebras (see [12, Prop. 1.5.5]).

Remark 5.6. Since Gle^R is equivalent to DeV^S (see Theorem 4.5), an analogue of Theorem 5.4 is that isomorphisms in Gle^R are homeomorphisms that preserve and reflect the equivalence relation.

6. An alternative approach to de Vries duality

In this final section we show that the equivalence between KHaus^R and DeV^S restricts to an equivalence between KHaus and the wide subcategory DeV^F of DeV^S whose morphisms satisfy two additional conditions (the superscript F signifies that these morphisms correspond to those morphisms in KHaus^R that are functions). This provides an alternative of de Vries duality. We finish the paper by giving a direct proof that DeV is dually isomorphic to DeV^F . An advantage of DeV^F over DeV is that composition in DeV^F is usual relation composition.

Definition 6.1. We define the category StoneE^F as $\text{Map}(\text{StoneE}^R)$. Explicitly, this is the category whose objects are pairs (X, E) where E is a closed equivalence relation on a Stone space X , and whose morphisms from (X_1, E_1) to (X_2, E_2) are the compatible closed relations $R: X_1 \rightarrow X_2$ such that $E_1 \subseteq R^\smile \circ R$ and $R \circ R^\smile \subseteq E_2$ (see Remark 3.13).

Proposition 6.2. *The categories KHaus and StoneE^F are equivalent.*

Proof. By Theorem 3.12, KHaus^R and StoneE^R are equivalent allegories. Therefore, $\text{Map}(\text{KHaus}^R)$ and $\text{Map}(\text{StoneE}^R)$ are equivalent categories (see [14, p. 204]). The result follows since $\text{KHaus} = \text{Map}(\text{KHaus}^R)$ and $\text{StoneE}^F = \text{Map}(\text{StoneE}^R)$. \square

Remark 6.3. Since $\text{StoneE}^F = \text{Map}(\text{StoneE}^R) = \text{Map}(\text{Split}(\text{Eq}_{\text{Stone}^R}))$ and $\text{Map}(\text{Split}(\text{Eq}_{\text{Stone}^R}))$ is isomorphic to $\text{Map}(\text{Split}(\text{Eq}_{\text{Rel}(\text{Stone})}))$ (see Remark 3.13), we have that StoneE^F is the exact completion of Stone . Thus, as a consequence we obtain an alternative proof of the fact that KHaus is the exact completion of Stone .

Definition 6.4. We define the category SubS5^F as $\text{Map}(\text{SubS5}^S)$. Explicitly, this is the category whose objects are pairs (B, S) where S is a $S5$ -subordination on a Boolean algebra B , and whose morphisms from (B_1, S_1) to (B_2, S_2) are the morphisms $T: (B_1, S_1) \rightarrow (B_2, S_2)$ in SubS5^S satisfying $\widehat{T} \circ T \subseteq S_1$ and $S_2 \subseteq T \circ \widehat{T}$ (see Remark 3.13).

We next give a slightly more explicit description of morphisms in SubS5^F .

Lemma 6.5. *A morphism $T: (B_1, S_1) \rightarrow (B_2, S_2)$ in SubS5^S is a morphism in SubS5^F iff the following two conditions hold.*

- (1) $\forall a \in B_1 (a T 0 \Rightarrow a = 0)$.
- (2) $\forall b_1, b_2 \in B_2 (b_1 S_2 b_2 \Rightarrow \exists a \in B_1 : \neg a T \neg b_1 \text{ and } a T b_2)$.

Proof. It is immediate that (2) is equivalent to $S_2 \subseteq T \circ \widehat{T}$. Moreover, $\widehat{T} \circ T \subseteq S_1$ is equivalent to the following condition:

$$\forall a_1, a_2 \in B_1 \forall b \in B_2 ((a_1 T b \text{ and } \neg a_2 T \neg b) \Rightarrow a_1 S_1 a_2). \tag{**}$$

We show that (**) is equivalent to (1). Suppose that (**) holds, $a \in B_1$, and $a T 0$. Letting $a_1 = a$, $a_2 = 0$, and $b = 0$ in (**) yields $a S_1 0$. Therefore, $a = 0$ by (S5). Conversely, suppose that (1) holds. Let $a_1, a_2 \in B_1$ and $b \in B_2$ such that $a_1 T b$ and $\neg a_2 T \neg b$. Since T is a compatible subordination, from $a_1 T b$ it follows that there is $a \in B_1$ such that $a_1 S_1 a T b$. From $\neg a_2 T \neg b$ and $a T b$ it follows that $(\neg a_2 \wedge a) T (\neg b \wedge b) = 0$. Therefore, (1) implies $\neg a_2 \wedge a = 0$, so $a \leq a_2$. Thus, $a_1 S_1 a \leq a_2$, and hence $a_1 S_1 a_2$. \square

Proposition 6.6. *The categories StoneE^F and SubS5^F are equivalent.*

Proof. This follows from the fact that StoneE^R and SubS5^S are equivalent allegories (see Theorem 3.10), together with the facts that StoneE^F = Map(StoneE^R) and SubS5^F = Map(SubS5^S). \square


Putting Propositions 6.2 and 6.6 together yields:

Corollary 6.7. *The categories KHaus, StoneE^F, and SubS5^F are equivalent.*

Definition 6.8. Following [5, Def. 6.5], we let Gle denote the full subcategory of StoneE^F whose objects are Gleason spaces. We also let DeV^F denote the full subcategory of SubS5^F consisting of de Vries algebras.

We then have Gle = Map(Gle^R) and DeV^F = Map(DeV^S). Consequently, we arrive at the following.

Theorem 6.9. *The categories KHaus, StoneE^F, Gle, SubS5^F, and DeV^F are equivalent.*



Proof. By Corollaries 3.14 and 4.7, the allegories KHaus^R, StoneE^R, Gle^R, SubS5^S and DeV^S are equivalent. Therefore, the categories Map(KHaus^R), Map(StoneE^R), Map(Gle^R), Map(SubS5^S), and Map(DeV^S) are equivalent. Thus, KHaus, StoneE^F, Gle, SubS5^F, and DeV^F are equivalent. \square

In particular, the equivalence between KHaus and DeV^F provides an alternative of de Vries duality. In the rest of the paper we show how to derive de Vries duality from this result. We start by recalling the definition of a de Vries morphism. From now on, following Remark 4.3, we denote a de Vries proximity on a boolean algebra by \prec .

Definition 6.10. A function $f: A \rightarrow B$ between de Vries algebras (A, \prec) and (B, \prec) is a *de Vries morphism* if it satisfies the following conditions:

- (M1) $f(0) = 0$;
- (M2) $f(a \wedge b) = f(a) \wedge f(b)$;
- (M3) $a \prec b$ implies $\neg f(\neg a) \prec f(b)$;
- (M4) $f(a) = \bigvee \{f(b) \mid b \prec a\}$.

The composition of two de Vries morphisms $f: A \rightarrow B$ and $g: B \rightarrow C$ is the de Vries morphism $g * f: A \rightarrow C$ given by

$$(g * f)(a) = \bigvee \{gf(b) \mid b \prec a\}$$

for each $a \in A$. Let \mathbf{DeV} be the category of de Vries algebras and de Vries morphisms, where identity morphisms are identity functions and composition is defined as above.

Remark 6.11. Each de Vries morphism $f: A \rightarrow B$ satisfies that $a \prec b$ implies $f(a) \prec f(b)$ for each $a, b \in A$. Moreover, for a de Vries morphism $g: B \rightarrow C$ we have that $a \prec b$ implies $gf(a) \leq (g * f)(b) \leq gf(b)$. This will be used in what follows.

We recall from the introduction that de Vries duality is induced by the contravariant functor $\mathbf{RO}: \mathbf{KHaus} \rightarrow \mathbf{DeV}$ that associates to each $X \in \mathbf{KHaus}$ the de Vries algebra $(\mathbf{RO}(X), \prec)$ of regular opens of X where $U \prec V$ iff $\text{cl}(U) \subseteq V$. The functor \mathbf{RO} sends each continuous function $f: X \rightarrow Y$ between compact Hausdorff spaces to the de Vries morphism $f^*: \mathbf{RO}(Y) \rightarrow \mathbf{RO}(X)$ given by $f^*(V) = \text{int}(\text{cl}(f^{-1}[V]))$ for each $V \in \mathbf{RO}(Y)$.

We show that \mathbf{DeV} is dually isomorphic to \mathbf{DeV}^F . The definition of contravariant functors between \mathbf{DeV} and \mathbf{DeV}^F is motivated by the following result.

Proposition 6.12. *Let $f: X \rightarrow Y$ be a continuous function between compact Hausdorff spaces.*

(1) *For every $V \in \mathbf{RO}(Y)$,*

$$\text{int}(\text{cl}(f^{-1}[V])) = \bigvee \{U \in \mathbf{RO}(X) \mid f[\text{cl}(U)] \subseteq V\},$$

where the join is computed in $\mathbf{RO}(X)$.

(2) *For every $U \in \mathbf{RO}(X)$ and $V \in \mathbf{RO}(Y)$,*

$$f[\text{cl}(U)] \subseteq V \iff \exists V' \in \mathbf{RO}(Y) : \text{cl}(V') \subseteq V \text{ and } U \subseteq \text{int}(\text{cl}(f^{-1}[V'])).$$

Proof. (1). Since $f^{-1}[V]$ is open, we have

$$f^{-1}[V] = \bigcup \{U \in \mathbf{RO}(X) \mid \text{cl}(U) \subseteq f^{-1}[V]\}.$$

Therefore,

$$\text{int}(\text{cl}(f^{-1}[V])) = \bigvee \{U \in \mathbf{RO}(X) \mid \text{cl}(U) \subseteq f^{-1}[V]\} = \bigvee \{U \in \mathbf{RO}(X) \mid f[\text{cl}(U)] \subseteq V\}.$$

(2). To prove the left-to-right implication, suppose $f[\text{cl}(U)] \subseteq V$. Since $f[\text{cl}(U)]$ is closed and

$$V = \bigcup \{V' \in \mathbf{RO}(Y) \mid \text{cl}(V') \subseteq V\},$$

where the union is directed, there is $V' \in \mathbf{RO}(Y)$ such that $f[\text{cl}(U)] \subseteq V' \subseteq \text{cl}(V') \subseteq V$. Therefore, $U \subseteq \text{cl}(U) \subseteq f^{-1}[V'] \subseteq \text{int}(\text{cl}(f^{-1}[V']))$.

To prove the right-to-left implication, suppose that there is $V' \in \mathbf{RO}(Y)$ such that $\text{cl}(V') \subseteq V$ and $U \subseteq \text{int}(\text{cl}(f^{-1}[V']))$. Then

$$\text{cl}(U) \subseteq \text{cl}(\text{int}(\text{cl}(f^{-1}[V']))) = \text{cl}(f^{-1}[V']) \subseteq \text{cl}(f^{-1}[\text{cl}(V')]) = f^{-1}[\text{cl}(V')] \subseteq f^{-1}[V],$$

which implies $f[\text{cl}(U)] \subseteq V$. \square

Proposition 6.12 suggests the following definition of two contravariant functors providing a dual isomorphism between DeV and DeV^F .

Definition 6.13.

1. The contravariant functor from DeV^F to DeV is the identity on objects and maps a morphism $S: A \rightarrow B$ in DeV^F to the function $f_S: B \rightarrow A$ given by

$$f_S(b) = \bigvee \{a \in A \mid a S b\} = \bigvee S^{-1}[b].$$

2. The contravariant functor from DeV to DeV^F is the identity on objects and maps a de Vries morphism $f: A \rightarrow B$ to the relation $S_f: B \rightarrow A$ given by

$$b S_f a \iff \exists a' \in A : a' \prec a \text{ and } b \leq f(a').$$

To show that the above functors are well defined, we need the following lemma.

Lemma 6.14. *Let $S: A \rightarrow B$ be a morphism in DeV^F and let $b_1, b_2 \in B$ be such that $b_1 \prec b_2$. Then $f_S(b_1) S b_2$.*

Proof. By Lemma 6.5(2), from $b_1 \prec b_2$ it follows that there is $a_0 \in A$ such that $\neg a_0 S \neg b_1$ and $a_0 S b_2$. We show that $a S b_1$ implies $a \leq a_0$. From $a S b_1$ and $\neg a_0 S \neg b_1$ it follows that $(a \wedge \neg a_0) S (b_1 \wedge \neg b_1) = 0$. By Lemma 6.5(1), $a \wedge \neg a_0 = 0$, so $a \leq a_0$. Thus, $\bigvee \{a \in A \mid a S b_1\} \leq a_0$, and hence $\bigvee \{a \in A \mid a S b_1\} S b_2$, which gives $f_S(b_1) S b_2$. \square

Lemma 6.15. *The assignment in Definition 6.13(1) is a well-defined contravariant functor from DeV^F to DeV .*

Proof. Let $S: A \rightarrow B$ be a morphism in DeV^F . We show that f_S is a de Vries morphism. That f_S satisfies (M1) follows from Lemma 6.5(1). The proof of (M2) is straightforward and (M4) follows from the fact that S is a compatible subordination. We prove (M3). Suppose $b_1 \prec b_2$. We must show that $\neg f_S(\neg b_1) \prec f_S(b_2)$, i.e. that $\neg \bigvee S^{-1}[\neg b_1] \prec \bigvee S^{-1}[b_2]$, which is equivalent to $\bigwedge \{\neg c \mid c \in S^{-1}[\neg b_1]\} \prec \bigvee S^{-1}[b_2]$. By Lemma 6.5(2), there is $a \in A$ such that $\neg a S \neg b_1$ and $a S b_2$. Since S is compatible, $a S b_2$ implies that there is $a' \in A$ such that $a \prec a' S b_2$. Because $\neg a \in S^{-1}[\neg b_1]$ and $a' \in S^{-1}[b_2]$, we have

$$\bigwedge \{\neg c \mid c \in S^{-1}[\neg b_1]\} \leq a \prec a' \leq \bigvee S^{-1}[b_2].$$

This proves that f_S is a de Vries morphism.

Let $S_1: A \rightarrow B$ and $S_2: B \rightarrow C$ be morphisms in DeV^F . We prove that $f_{S_2 \circ S_1} = f_{S_1} * f_{S_2}$.

Claim 6.16. *For every $a \in A$ and $c \in C$,*

$$a (S_2 \circ S_1) c \iff \exists c' \in C : c' \prec c \text{ and } a \leq f_{S_1} f_{S_2}(c').$$

Proof of claim. For the left-to-right implication, suppose $a (S_2 \circ S_1) c$. Then there is $b \in B$ such that $a S_1 b S_2 c$. From $a S_1 b$ and the definition of f_{S_1} it follows that $a \leq f_{S_1}(b)$. Since S is compatible, from $b S_2 c$ it follows that there is $c' \in C$ such that $b S_2 c' \prec c$. Therefore, $b \leq f_{S_2}(c')$, and so $a \leq f_{S_1}(b) \leq f_{S_1} f_{S_2}(c')$.

For the right-to-left implication, suppose that there is $c' \in C$ such that $c' \prec c$ and $a \leq f_{S_1} f_{S_2}(c')$. By Lemma 6.14, $f_{S_2}(c') S_2 c$. Since S is compatible, there is $b \in B$ such that $f_{S_2}(c') \prec b S_2 c$. Applying Lemma 6.14 again, $a \leq f_{S_1} f_{S_2}(c') S_1 b S_2 c$, which implies that $a S_1 b S_2 c$. Thus, $a (S_2 \circ S_1) c$. \square

For every $c \in C$, we have

$$\begin{aligned} (f_{S_1} * f_{S_2})(c) &= \bigvee \{f_{S_1} f_{S_2}(c') \mid c' \prec c\} \\ &= \bigvee \{a \in A \mid \exists c' \in C : c' \prec c \text{ and } a \leq f_{S_1} f_{S_2}(c')\} \\ &= \bigvee \{a \in A \mid a (S_2 \circ S_1) c\} && \text{(by Claim 6.16)} \\ &= f_{S_2 \circ S_1}(c). \end{aligned}$$

This proves $f_{S_2 \circ S_1} = f_{S_1} * f_{S_2}$.

Let (A, \prec) be a de Vries algebra. Since for every $a \in A$, we have $a = \bigvee \{b \in A \mid b \prec a\}$, the identity on (A, \prec) in DeV^F is mapped to the identity on (A, \prec) in DeV . \square

Lemma 6.17. *The assignment in Definition 6.13(2) is a well-defined contravariant functor from DeV to DeV^F .*

Proof. Let $f: A \rightarrow B$ be a de Vries morphism. We show that $S_f: B \rightarrow A$ is a morphism in DeV^F . It is straightforward to see that S_f is a subordination. We only verify (S2). Suppose that $b_1, b_2 S_f a$. Then there exist $a_1, a_2 \prec a$ such that $b_1 \leq f(a_1)$ and $b_2 \leq f(a_2)$. Therefore, $(a_1 \vee a_2) \prec a$ and

$$b_1 \vee b_2 \leq f(a_1) \vee f(a_2) \leq f(a_1 \vee a_2)$$

because f is order-preserving. Thus, $(b_1 \vee b_2) S_f a$. We next show that S_f is compatible. For this we need to show that $S_f \circ \prec = S_f = \prec \circ S_f$. To see that $S_f \subseteq S_f \circ \prec$, let $b S_f a$. Then there is $a' \in A$ such that $a' \prec a$ and $b \leq f(a')$. By (S7), there is $a'' \in A$ such that $a' \prec a'' \prec a$. Therefore, $b \leq f(a') \prec f(a'')$, so $f(a'') S_f a$, and hence $S_f \subseteq S_f \circ \prec$. The other inclusions are proved similarly. Finally, we show that S_f satisfies the two conditions of Lemma 6.5. Condition (1) is immediate from the definition of S_f and the fact that $f(0) = 0$. For (2), let $a_1 \prec a_2$. By (S7), there is $a \in A$ such that $a_1 \prec a \prec a_2$. By (M3), $\neg f(\neg a_1) \prec f(a)$. Set $b = f(a)$. It is left to show that $\neg b S_f \neg a_1$ and $b S_f a_2$. The latter is obvious because $a \prec a_2$ and $b = f(a)$. We prove the former. By (S7), there is $c \in A$ such that $a_1 \prec c \prec a$. Then $\neg b = \neg f(a) \prec f(\neg c)$ by (M3) and (S6), and hence $\neg b \leq f(\neg c)$ by (S5). Moreover, $\neg c \prec \neg a_1$ by (S6). Thus, $\neg b S_f \neg a_1$. This proves that S_f is a morphism in DeV^F .

Let $f: A \rightarrow B$ and $g: B \rightarrow C$ be de Vries morphisms. We prove that $S_{g*f} = S_f \circ S_g$. To see that $S_f \circ S_g \subseteq S_{g*f}$, let $a \in A$ and $c \in C$ be such that $c (S_f \circ S_g) a$. Then there is $b \in B$ such that $c S_g b S_f a$. Since $c S_g b$, it follows from the definition of S_g that there is $b' \in B$ such that $b' \prec b$ and $c \leq g(b')$. Also, since $b S_f a$, there is $a' \in A$ such that $a' \prec a$ and $b \leq f(a')$. By (S7), there is $a'' \in A$ such that $a' \prec a'' \prec a$. Therefore, $c \leq g(b') \leq g(b) \leq g f(a') \leq (g * f)(a'')$. This proves $c S_{g*f} a$. To see that $S_{g*f} \subseteq S_f \circ S_g$, let $a \in A$ and $c \in C$ be such that $c S_{g*f} a$. Then there is $a' \in A$ such that $a' \prec a$ and $c \leq (g * f)(a') \leq g f(a')$. By (S7), there is $a'' \in A$ such that $a' \prec a'' \prec a$. Therefore, $f(a') \prec f(a'')$, and so $c S_g f(a'') S_f a$. Thus, $c (S_f \circ S_g) a$.

Finally, if f is the identity on (A, \prec) in DeV , then $S_f = \prec$, and hence it is the identity on (A, \prec) in DeV^F . \square

Theorem 6.18. *The contravariant functors described in Definition 6.13 establish a dual isomorphism between DeV^F and DeV .*

Proof. It is sufficient to show that for each morphism $S: A \rightarrow B$ in DeV^F we have $S_{S_f} = S$, and that for each morphism $f: A \rightarrow B$ in DeV we have $f_{S_f} = f$.

Let $S: A \rightarrow B$ be a morphism in DeV^F . Suppose $a \in A$, $b \in B$, and $a S_{f_S} b$. Then there is $b' \in B$ such that $b' \prec b$ and $a \leq f_S(b')$. By Lemma 6.14, $a \leq f_S(b') S b$, so $a S b$. This proves $S_{f_S} \subseteq S$. For the other inclusion, suppose $a \in A$, $b \in B$, and $a S b$. Since S is compatible, there is $b' \in B$ such that $a S b' \prec b$. We have $a \leq \bigvee \{c \in A \mid c S b'\} = f_S(b')$. Therefore, the element b' witnesses that we have $a S_{f_S} b$. Thus, $S \subseteq S_{f_S}$, and hence $S_{f_S} = S$.

Let $f: A \rightarrow B$ be a de Vries morphism. For $a \in A$ we have $f_{S_f}(a) = \bigvee \{b \in B \mid b S_f a\}$. Also, $f(a) = \bigvee \{f(a') \mid a' \prec a\}$ by (M4). If $a' \prec a$, then $f(a') \prec f(a)$, so $f(a') S_f a$. Therefore, $f(a')$ is one of the $b \in B$ such that $b S_f a$, and hence $f(a) \leq f_{S_f}(a)$. On the other hand, if $b S_f a$, then there is $a' \in A$ such that $a' \prec a$ and $b \leq f(a')$. Therefore, $b \leq f(a') \prec f(a)$, and so $f_{S_f}(a) \leq f(a)$. Thus, $f_{S_f}(a) = f(a)$, and hence $f_{S_f} = f$. \square

Remark 6.19. Combining Theorems 6.9 and 6.18 yields de Vries duality. Consequently, all the categories that appear in Theorem 6.9 are dually equivalent to DeV .

Acknowledgements

We thank Sergio Celani for drawing our attention to [11], which simplified our original proof of Corollary 2.6. We are also thankful to the referee for suggesting the machinery of order-enriched categories, which has considerably shortened our proofs. Remarks 3.13 and 6.3 were inspired by the referee.

Marco Abbadini and Luca Carai were supported by the Italian Ministry of University and Research through the PRIN project n. 20173WKCM5 *Theory and applications of resource sensitive logics*. Luca Carai acknowledges partial support from the Juan de la Cierva-Formación 2021 programme (FJC2021-046977-I) funded by MCIN/AEI/10.13039/501100011033 and by the European Union “NextGenerationEU”/PRTR.

References

- [1] M. Abbadini, G. Bezhanishvili, L. Carai, Ideal and MacNeille completions of subordination algebras, arXiv:2211.02974, 2022.
- [2] S. Abramsky, A. Jung, Domain Theory, Handbook of Logic in Computer Science, vol. 3, Oxford Univ. Press, New York, 1994, pp. 1–168.
- [3] G. Bezhanishvili, Stone duality and Gleason covers through de Vries duality, Topol. Appl. 157 (6) (2010) 1064–1080.
- [4] G. Bezhanishvili, N. Bezhanishvili, S. Sourabh, Y. Venema, Irreducible equivalence relations, Gleason spaces, and de Vries duality, Appl. Categ. Struct. 25 (3) (2017) 381–401.
- [5] G. Bezhanishvili, D. Gabelaia, J. Harding, M. Jibladze, Compact Hausdorff spaces with relations and Gleason spaces, Appl. Categ. Struct. 27 (6) (2019) 663–686.
- [6] P. Blackburn, M. de Rijke, Y. Venema, Modal Logic, Cambridge Tracts in Theoretical Computer Science, vol. 53, Cambridge University Press, Cambridge, 2001.
- [7] A. Carboni, E.M. Vitale, Regular and exact completions, J. Pure Appl. Algebra 125 (1–3) (1998) 79–116.
- [8] A. Carboni, R.F.C. Walters, Cartesian bicategories. I, J. Pure Appl. Algebra 49 (1–2) (1987) 11–32.
- [9] S.A. Celani, Quasi-modal algebras, Math. Bohem. 126 (4) (2001) 721–736.
- [10] S.A. Celani, Subdirectly irreducible quasi-modal algebras, Acta Math. Univ. Comen. (N.S.) 74 (2) (2005) 219–228.
- [11] S.A. Celani, Quasi-semi-homomorphisms and generalized proximity relations between Boolean algebras, Miskolc Math. Notes 19 (1) (2018) 171–189.
- [12] H. de Vries, Compact spaces and compactifications. An algebraic approach, Ph.D. thesis, University of Amsterdam, 1962.
- [13] R. Engelking, General Topology, second ed., Sigma Series in Pure Mathematics, vol. 6, Heldermann Verlag, Berlin, 1989.
- [14] P.J. Freyd, A. Scedrov, Categories, Allegories, North-Holland Mathematical Library, vol. 39, North-Holland Publishing Co., Amsterdam, 1990.
- [15] P.R. Halmos, Algebraic logic. I. Monadic Boolean algebras, Compos. Math. 12 (1956) 217–249.
- [16] C. Heunen, J. Vicary, Categories for Quantum Theory, Oxford Graduate Texts in Mathematics, vol. 28, Oxford University Press, Oxford, 2019.
- [17] D. Hofmann, G.J. Seal, W. Tholen (Eds.), Monoidal Topology. A Categorical Approach to Order, Metric, and Topology, Encycl. Math. Appl., vol. 153, Cambridge University Press, Cambridge, 2014.
- [18] R. Hoofman, Continuous information systems, Inf. Comput. 105 (1) (1993) 42–71.
- [19] J. Isbell, Atomless parts of spaces, Math. Scand. 31 (1972) 5–32.
- [20] P.T. Johnstone, Stone Spaces, Cambridge Studies in Advanced Mathematics, vol. 3, Cambridge University Press, Cambridge, 1982.

- [21] P.T. Johnstone, *Sketches of an Elephant: a Topos Theory Compendium*, vol. 1, Oxford Logic Guides, vol. 43, The Clarendon Press, Oxford University Press, New York, 2002.
- [22] A. Jung, M. Kegelmann, M.A. Moshier, Multilingual sequent calculus and coherent spaces, *Fundam. Inform.* 37 (4) (1999) 369–412.
- [23] A. Jung, M. Kegelmann, M.A. Moshier, Stably compact spaces and closed relations, *Electron. Notes Theor. Comput. Sci.* 45 (2001) 209–231.
- [24] A. Jung, A. Kurz, M.A. Moshier, Stone duality for relations, arXiv:1912.08418, 2019.
- [25] A. Jung, P. Sünderhauf, On the duality of compact vs. open, in: *Papers on General Topology and Applications*, Gorham, ME, 1995, in: *Ann. New York Acad. Sci.*, vol. 806, 1996, pp. 214–230, New York.
- [26] M. Kegelmann, Continuous domains in logical form, *Electron. Notes Theor. Comput. Sci.* 49 (2002).
- [27] J. Lambek, Diagram chasing in ordered categories with involution, *J. Pure Appl. Algebra* 143 (1–3) (1999) 293–307.
- [28] K.G. Larsen, G. Winskel, Using information systems to solve recursive domain equations effectively, in: *Semantics of Data Types*, Valbonne, 1984, in: *Lecture Notes in Comput. Sci.*, vol. 173, Springer, Berlin, 1984, pp. 109–129.
- [29] S. Mac Lane, *Categories for the Working Mathematician*, second ed., Graduate Texts in Mathematics, vol. 5, Springer-Verlag, New York, 1998.
- [30] V. Marra, L. Reggio, A characterisation of the category of compact Hausdorff spaces, *Theory Appl. Categ.* 35 (2020) 1871–1906.
- [31] M.A. Moshier, On the relationship between compact regularity and Gentzen’s cut rule, *Theor. Comput. Sci.* 316 (1–3) (2004) 113–136.
- [32] J.R. Porter, R.G. Woods, *Extensions and Absolutes of Hausdorff Spaces*, Springer-Verlag, New York, 1988.
- [33] D.S. Scott, Domains for denotational semantics, in: *Automata, Languages and Programming*, Aarhus, 1982, in: *Lecture Notes in Comput. Sci.*, vol. 140, Springer, Berlin-New York, 1982, pp. 577–613.
- [34] M.Sh. Tsalenko, V.B. Gisin, D.A. Rajkov, Ordered categories with involution, *Diss. Math.* 227 (1984).
- [35] J. Vicary, Completeness of \dagger -categories and the complex numbers, *J. Math. Phys.* 52 (8) (2011) 082104.
- [36] S. Vickers, Information systems for continuous posets, *Theor. Comput. Sci.* 114 (2) (1993) 201–229.