UNIVERSITY^{OF} BIRMINGHAM University of Birmingham Research at Birmingham

Health-related fitness in medical students

Paez, Diana C; Cortes Corrales, Sebastian; Jimenez-Mora, Mario A; Gutiérrez, Agustín; Arango-Paternina, Carlos M. ; Duperly, John

DOI: 10.1186/s12889-024-17748-y License: Creative Commons: Attribution (CC BY)

Document Version Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Paez, DC, Cortes Corrales, S, Jimenez-Mora, MA, Gutiérrez, A, Arango-Paternina, CM & Duperly, J 2024, 'Health-related fitness in medical students: a curricular intervention in Bogota, Colombia', *BMC Public Health*, vol. 24, no. 1, 320. https://doi.org/10.1186/s12889-024-17748-y

Link to publication on Research at Birmingham portal

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

•Users may freely distribute the URL that is used to identify this publication.

Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)

•Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

RESEARCH

Open Access

Health-related fitness in medical students: a curricular intervention in Bogota, Colombia



Diana C. Paez^{1*}, Sebastián Cortés-Corrales², Mario A. Jimenez-Mora¹, Agustín Gutiérrez³, Carlos M. Arango-Paternina⁴ and John Duperly^{1,5}

Abstract

Objective To evaluate the impact of a curricular intervention to promote health-related fitness (HRF) among medical students in Bogota, Colombia.

Method The study was conducted between May 2014 and December 2015 as part of the medical physiology course, in which 208 medical students were enrolled. The curricular intervention included two lectures on physical activity (PA) and student-led group presentations on the physiological effects of exercise on human physiology. An academic incentive (10% of final grade) was given to students who reported and documented regular PA practice during the semester. This study assessed students' HRF variables, perceptions of the curriculum intervention, and PA practices using quantitative and qualitative approaches.

Results 55% of the students were female, with a mean age of 19.5 years. Body fat, estimated maximum oxygen consumption (VO2max), handgrip, and sit-up strength showed statistically significant improvements at the end of the intervention. Students reported that PA practice was positively influenced by the curriculum intervention, particularly the academic incentive and the HRF tests. Students reported a wide variety of PA practices, which were mainly done with friends, classmates, or family members. Lack of time was the main reported barrier to PA practice.

Conclusion The curricular intervention was effective in improving HRF and promoting PA. It remains to be investigated whether these gains are sustained over time.

Keywords Curriculum, Intervention, Lifestyles, Health-related fitness, Physical activity

*Correspondence:

¹School of Medicine, Universidad de los Andes, Bogotá, Colombia
²Department of Economics, University of Birmingham, Birmingham, UK
³Department of Social Medicine, Universidad de los Andes, Bogotá, Colombia

© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence are on your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Dublic Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

The prevalence of chronic non-communicable diseases (NCDs) is on the rise globally, with a pronounced impact in developing nations like Colombia [1]. In Colombia, these persistent NCDs stand as the primary contributors to both illness and death recognized [2]. Global surveillance data show that the NCD pandemic could be prevented by reducing common risk factors, mainly tobacco use, physical inactivity, unhealthy diets, and excessive alcohol consumption [3–5]. Accordingly, international health organisations advocate the control of these key behavioural risk factors through the promotion of healthy lifestyles, including physical activity (PA) [6].

Diana C. Paez

paez.d.carolina@gmail.com

⁴Instituto Universitario de Educación Física y Deportes, Universidad de Antioquia, Medellín, Colombia

⁵Institute of Exercise Medicine and Rehabilitation, Fundación Santa Fe de Bogotá, Bogotá, Colombia

The medical community has a key role to play in promoting healthy lifestyles for the prevention and management of diseases [7]. However, few healthcare providers, such us physicians, nurses, medical assistants, psychologists, and physical therapists, have the appropriate environment, knowledge, self-efficacy and lifestyle medicine skills to counsel patients on prevention [8, 9]. Integration of skills into routine clinical practice remains a challenge [10–12]; partly due to infrequent and inadequate training during medical education. Medical schools reported including PA [13, 14] or exercise prescription [15] as part of the curriculum. However, the medical curriculum is insufficiently focused on prevention, is rarely evaluated, and is occasionally adopted [16].

Strategies to integrate PA into lifestyle medicine curricula are needed for current and future physicians to effectively promote behaviour change in patients [17, 18]. Furthermore, not many medical education models promote the adoption of active lifestyles among students. This is important because studies have shown that, in addition to knowledge, personal PA behaviours and physical fitness levels positively influence physicians' PA counselling practices [19–24]. Knowledge of exercise and fitness provides medical students with a solid foundation in exercise physiology, allowing them to better understand the physiological responses in terms of benefits and adaptations to exercise [25]. This knowledge would enhance physicians ability to provide PA recommendations, and write PA or exercise prescriptions to patients.

There is some encouraging evidence that PA curricular interventions in medical schools have been successful in increasing students' physical fitness [26], personal PA habits, knowledge, skills, and self-efficacy [27]. However, efforts are needed to adapt and evaluate feasible interventions that can be easily integrated in LMICs. Therefore, the aim of this study is to evaluate the impact of a curricular intervention focused on improving healthrelated fitness (HRF) and PA knowledge, attitudes, and behaviours among second-year medical students at the Universidad de los Andes in Bogota, Colombia.

Methods

Population, setting and study design

A mixed methods intervention design was conducted between May 2014 and December 2015 to assess the impact of an intervention to improve HRF in second year medical students. This design is useful for exploring in more detail the impact of an intervention and participants' perceptions of [28]. In this case, the integration of the quantitative and qualitative research components occurred after the intervention. The intervention was embedded in the compulsory medical physiology course across three cohorts of students. Data were available from 208 enrolled medical students (68 in the second semester of 2014, 88 in the first semester of 2015 and 52 in the second semester of 2015) at the Universidad de los Andes in Bogota, Colombia.

The outcomes assessed were students' HRF variables, their perceptions of the curriculum intervention and PA practices.

Medical Physiology course description: The course provided second year (i.e. fourth academic semester) medical students with a basic knowledge of how the human body functions to further understand disease processes and therapeutic approaches. The course covered molecular and cellular physiology and major physiological topics such as cardiopulmonary, neuromuscular, endocrine, gastrointestinal and renal systems. The course consisted of 60 h during the 20-week academic semester. A numerical scale from 0 to 5 was used for grading. Course assessment was based on the following: three exams (70% of total grade), quizzes (10% of total grade) and group presentations (10% of total grade). In addition, 10% of the total course mark was allocated to students' self-assessed personal commitment to PA. A total of 10% of the questions in the exams and guizzes were dedicated to exercise physiology.

Quantitative component

A quasi-experimental study design was conducted. In total, 12 of the 60 h (20%) of the medical physiology course were spent on the curriculum intervention. The curriculum intervention consisted of lectures on muscle contraction and exercise physiology for a total of 6 h during the academic semester. In addition, 3 h were allocated to student-led group presentations on the effects of healthy habits (i.e. exercise, diet, tobacco cessation) on specific pathophysiological processes. Basic HRF assessments were administered at the beginning (week 1) and at the end of the course (week 20), and took a total of 3 h to complete.

Group presentations Students were asked to form groups of 8. Each group was randomly assigned an academic topic on the prevention and/or treatment of disease through the promotion of PA, to be researched through a literature review. At the end of the course, students were asked to submit an abstract (500 words) and give a 10-minute academic presentation to classmates, the professor and experts in the field. The presentations were evaluated by the members of the evaluation panel according to a preestablished rubric. The pre-established rubric comprises five key criteria (1. Quality of research and content, 2. Structure and organization of the presentation, 3. Quality of the presentation, 4. Ability to comprehend the topic and respond to questions from the audience and the evaluation panel, and 5. Group collaboration and contribution). Academic incentive for PA practice An important component of the curriculum intervention was an academic incentive to encourage students to practise PA during the semester. Students received support from teaching assistants and resources available at the university. For example, students were encouraged to attend the sports centre free of charge (i.e. the sports centre serves as a free resource available to the student community) and those who attended were trained by sports professionals. The 10% of the final grade, as described above, was agreed between the professor and the student independently of the HRF test results. No student received a grade lower than 3 out of 5. Students were also told that their answers would not affect their final grade.

Outcome variables HRF variables were assessed before and after the curricular intervention. All measurements were carried out at the Universidad de los Andes Sports Centre according to a standardised protocol. A detailed manual of procedures was developed by the research group and was reviewed by the staff prior to the measurements. The staff consisted of research assistants and trained medical students. After basic screening for PA readiness (PAR-Q and YOU) [29], students received instructions (e.g., use of sports clothing, hydration) and recommendations for possible contraindications (i.e., acute illness, fever, chest pain). None of the students had any contraindications.

Weight and height were measured in lightly dressed students with shoes removed. Body weight was measured using calibrated electronic scales (Beurer Bg 21 Tanita Omron) and height using an acrylic stadiometer. Body mass index (BMI) was calculated from weight and height (kg/m2) and categories (i.e. underweight, normal weight, overweight and obesity) were calculated according to the World Health Organization standards (WHO, 2000) [30]. A handheld device was used to measure body fat percentage using a bioelectrical impedance device (Omron HBF-306 C).

Cardiorespiratory fitness was assessed using the 20-metre multistage shuttle run test (20-MST), previously validated for the study population [31]. Students were required to walk or run between two lines 20 m apart at a pace announced by recorded tones at standard intervals. For any given participant, the test was terminated if the participant failed to reach the line in time for the beep on two consecutive occasions. Heart rates were recorded continuously throughout the test (Polar heart rate monitor, model FT7). The last stage completed was recorded and then used to estimate maximum oxygen consumption (VO2max) according to the equation of Leger et al. [32].

Grip strength was assessed using the JAMAR[®] hand dynamometer (JAMAR, 5030J1), which was calibrated

prior to the measurements. Students were asked to assume a seated position and grasp the dynamometer with the dominant hand, with the shoulder adducted and neutrally rotated, the elbow flexed at 90°, the forearm in neutral position, and the wrist between 0° and 30° dorsiflexion and between 0° and 15° ulnar deviation. Students positioned the instrument so that it fit comfortably in their hands. Once the student was in the correct position, the staff gave a verbal instruction to 'squeeze as hard as possible' and apply maximum force. The test was repeated three times. Maximum grip force was recorded in kilograms [33].

Abdominal muscle endurance was measured as the maximal number of sit-ups in 30 seconds. Students were asked to lie on a mat with their knees bent at right angles, their feet flat on the floor and held down by a partner. Their fingers had to be interlocked behind their heads. On the command 'go' the participants had to touch their knees with their elbows and then return to the floor. The maximum number of repetitions in 30 s was recorded.

Statistical analysis Means and standard deviations were calculated for demographic, body composition and fitness variables for all participants, stratified by sex. T-tests assuming equal and unequal variance (according to F-test results) were performed to analyse baseline differences by sex. Two-tailed paired Student's t-tests were used to compare body composition and fitness variables before and after the intervention in men and women separately. The effect of the intervention was calculated using a potential outcome framework using the mean difference between pre and post scores [34]. This methodology considers the pre-test measure as the contra-factual (i.e., control) for each student. Five multilevel mixed-effects generalized linear regression models were run to estimate the effect of the intervention on body composition, cardiorespiratory fitness and muscular strength and endurance. An individual mixed effect was used in the regression analyses to control for the individual inherent time-invariant characteristics (i.e., confounds) that may affect the students' progress during the semester. Specifically, we use a series of linear regression models to estimate the relationship between our dependent variables and various factors, taking into account random effects at both the cohort and student levels. In this way, we control for all time invariant-confounds that may arise by the cohort and students' idiosyncratic characteristics that might affect our variables of interest. For each of our dependent variables, including BMI, body fat percentage, VO2max, handgrip strength and sit-up strength, we adopt the following specification.

$$y_{ict} = \beta_0 + \beta_1 Intervention_t + \beta_2 Gender_i + \beta_3 Intervention_t * Gender_i + \gamma_i + \delta_c + \epsilon_{ict}$$

where i refers to each student, while t denotes the period, i.e. before and after the intervention. The multivariate regression model includes the following regressors $Intervention_t$ a dichotomous variable that takes the value 0 before the intervention and 1 after the intervention, and $Gender_i$ another dichotomous variable based on the gender of each student. Here, female students are coded as 1, while male students are coded as 0. In addition, we account for the random effects of each student and cohort using the normal standard random variables γ_i and δ_c , respectively. For the regression with the dependent variable VO2max, we included peak heart rate as a control as it is a potential confounder. Statistical analyses were performed using Stata 12 and a 95% confidence interval for hypothesis testing (StataCorp LP, College Station, TX, USA).

Qualitative component

An interpretive descriptive qualitative design [35] was used to assess students' PA practices and their perceptions of the curriculum intervention. Data were collected through self-report assessment and semi-structured interviews over two academic semesters (II-2014 and I-2015).

At the end of the course, students were asked to submit self-reports detailing their level of PA adherence, perceptions of sport skills, and personal experiences of engaging in PA. These self-reports were submitted in written form. In addition, students were encouraged to provide supporting documentation to supplement their reports, including photographs and sports centre attendance records.

The interview process was conducted by a diverse team of four trained researchers specialising in qualitative data collection methods (i.e. an anthropologist, a psychologist,

Table 1	Demographic characteristics, body composition, and	
baseline	fitness tests in second year medical students (<i>n</i> = 166)	

Variable	Male (n=70)	Female (<i>n</i> = 96)	<i>p</i> - value	
	Mean (SD)	Mean (SD)	(Paired t test)	
Age (years)	19.7 (1.8)	19.4 (1.3)	0.337	
Weight (kg)	68.4 (10.5)	58.6 (9.9)	< 0.001	
Height (cm)	174.9 (6.0)	162.3 (6.3)	0.003	
Body mass index (kg·m⁻²)	22.3 (2.7)	22.0 (3.8)	0.558	
Body fat (%)	16.0 (6.4)	25.0 (5.7)	< 0.001	
Estimated VO2max (ml·kg ⁻¹ ·min ⁻¹)	40.5 (5.1)	31.1 (3.7)	< 0.001	
Peak heart rate (bpm)	192.2 (11.4)	179.1 (19.6)	< 0.001	
Handgrip strength (kg)	43.2 (8.3)	26.7 (4.3)	< 0.001	
Sit up strength (# sit ups in 30 s)	26.1 (5.3)	20.8 (3.9)	< 0.001	

an economist and a physiotherapist). Several themes were explored during the interviews, including the students' motivations and their support network (i.e. friends and family members). Students were also asked to provide insights into the benefits and barriers they encountered when engaging in PA, their preferred types of PA, the locations where they preferred to engage in PA, and the people with whom they usually engaged in PA. In addition, the interviews proved valuable in understanding students' perceptions of the curriculum intervention and in identifying the factors influencing behaviour change over the semester.

Participants: Purposive sampling was used to select participants. The selection criteria were medical students enrolled in the medical physiology course who participated in the physical fitness tests. During II-2014, interviews were conducted with students (n=6) as a pilot interview guide for clarity and applicability of questions. During the academic term I-2015, 36 students participated in the semi-structured interviews: 22 students were included in the pre-intervention and 14 students in the post-intervention.

Data analysis The interviews were audio-recorded, transcribed verbatim by the interviewers and analysed using ATLAS.ti 7. The data collected were coded into categories and then analysed according to suggested procedures [36]. Theoretical saturation was achieved at all interview stages [37]. Notes were translated into English by the bilingual research staff of the Universidad de los Andes.

Ethical aspects

This study was approved by the Ethical Review Committee of the University of Los Andes (Act No. 538/2015). Students' participation in the physical fitness tests, selfreports, and interviews was voluntary. Those who agreed to participate in the study signed an informed consent form.

Results

A total of 208 students were enrolled in the physiology course, 68 in II-2014, 88 in I-2015 and 52 in II-2015. One hundred and ninety-one students participated in the preintervention physical fitness tests and 172 students participated in the post-intervention physical fitness tests. One hundred and sixty-six students completed information on all dependent variables (i.e. BMI, body fat, estimated VO2max, handgrip strength and sit-up test) in both the pre- and post-tests. Of the participating students, 58% were female with a mean age of 19.5 years. As expected, significant differences in body composition variables and fitness test performance were observed between males and females at baseline (Table 1).

Table 2 Pre and post levels of the bo	y composition and fitness tests in second	year medical students (n = 166)
---------------------------------------	---	---------------------------------

Variable	Male (n=70)			Female (<i>n</i> = 96)		
	Pre	Post	<i>p</i> -value (Paired t test)	Pre	Post	<i>p</i> -value (Paired t test)
	Mean (SD)	Mean (SD)		Mean (SD)	Mean (SD)	
Body mass index (kg·m-2)	22.3 (2.7)	22.5 (2.5)	0.131	22.0 (3.8)	22.0 (2.9)	0.943
Body fat (%)	16.3 (6.3)	14.9 (5.6)	< 0.001	25.0 (5.7)	24.2 (5.2)	0.035
Estimated VO2max (ml·kg-1·min-1)	40.7 (5.1)	43.8 (5.3)	< 0.001	31.1 (3.7)	34.6 (3.8)	< 0.001
Peak heart rate (bpm)	192.0 (11.3)	193.8 (10.4)	0.197	178.7 (20.2)	181.9 (16.6)	0.284
Handgrip strength (kg)	43.0 (8.1)	45.7 (9.8)	0.001	26.7 (4.3)	28.5 (5.3)	< 0.001
Sit up strength (# sit ups in 30 s)	26.1 (5.3)	28.7 (4.4)	< 0.001	20.8 (3.9)	24.2 (3.6)	< 0.001

Table 3 Impact of curricular intervention on measures and physical fitness tests in medical students (n = 166)

	Body mass index (kg∙m-2)	Body fat (%)	Estimated VO2max (ml·kg-1·min-1)	Handgrip strength (kg)	Sit-up strength (#sit- up in 30 s)
Intervention's impact	0.16	-1.54***	3.22***	2.48***	2.84***
	[-0.33, 0.64]	[-2.42, -0.67]	[2.43, 4.01]	[1.03, 3.93]	[2.00, 3.68]
Sex (reference Male)	-0.39	8.77***	-6.31***	-16.11***	-5.30***
	[-1.24, 0.46]	[7.19, 10.34]	[-9.75, -2.88]	[-18.08, -14.14]	[-6.53, -4.07]
Sex*intervention	-0.11	0.54	0.38	-0.65	0.65
	[-0.76, 0.54]	[-0.63, 1.70]	[-0.67, 1.42]	[-2.57, 1.27]	[-0.47, 1.77]
Peak heart rate			-0.00		
			(-0.03, 0.02)		
Constant	22.44***	16.21***	39.96***	43.07***	25.96***
	[21.79, 23.08]	[15.02, 17.40]	[34.22, -45.71]	[41.44, 44.69]	[24.66, 27.26]

Confidence interval in square parentheses

**** *p*<0.01, *** *p*<0.05, * *p*<0.1

The sample was homogeneous within the three academic semesters, there were no significant differences for age, BMI, estimated VO2max or sit-up test. However, for the sit-up test, students from different semesters showed slight differences with a variation of 2 sit-ups.

Table 2 shows the means and standard deviations for the body composition and fitness variables for male and female participants. There were no statistically significant changes in BMI or peak heart rate on the Leger test between the two assessments. However, statistically significant differences were observed in body fat percentage, estimated VO2max, handgrip strength and sit-up strength over time.

In the regression adjusted models, women had higher body fat and lower estimated VO2max, handgrip strength and sit-up strength at baseline (Table 3). After the intervention, body fat decreased by 1.54% [CI: -2.42 to -0.67; *p*-value<0.01], estimated VO2max improved by 3.22 ml·kg-1·min-1 [CI: 2.43 to 4.01; *p*-value<0.01], handgrip strength increased by 2.48 kg [CI: 1.03 to 3.93; *p*-value<0.01], and the 30-second sit-up test increased by 2.84 sit-ups [CI: 2.00 to 3.68; *p*-value<0.01]. Conversely, no differences in BMI were observed after the intervention. The effect of the intervention on all dependent variables did not differ by sex.

Curriculum intervention

Fifty-seven per cent of the students surveyed (8 out of 14) said that the curriculum - lectures, case observation and discussions - had a positive impact on their PA practice.

"I think it is important for us to be motivated to be physically active (...). As the problems associated with a sedentary lifestyle have become more apparent during the lectures, we are more aware of the need to improve our fitness" (Male student, 20 years).

The academic incentive was a motivational strategy for most students to practice PA, start a routine or exercise during the semester, and maintain active behaviours.

"For a lot of us, the academic incentive is very important, the university looks at you for your grades, so it's important to try to get the highest grade possible. Anything that improves the average grade is really motivating for me (...), I would say it contributed to 50% of my decision to be more active" (Male student, 21 years old).

Similarly, students reported that the physical fitness tests encouraged them to set goals and adopt healthy behaviours during the semester. "The fitness test is used to evaluate yourself; you set goals at the beginning of the semester and are encouraged to achieve them. (Male student, 19 years old)

PA practice

Only 8 out of 22 students (36.4%) interviewed before the intervention reported practising PA (150 min of moderate intensity PA or 75 min of vigorous intensity PA per week), compared to 12 out of 14 students (85.7%) interviewed after the intervention.

Most students reported practising PA three times a week, with the duration of each session ranging from 45 min to 2 h. Selected types of activities were sports (volleyball, football, tennis, ultimate taekwondo and kickboxing), workouts (TRX, CrossFit, Zumba, Pilates and gym strengthening routines) and aerobic activities such as running and swimming. Walking and cycling were the modes of transport used by students to and within the campus.

Most students used university infrastructure facilities (e.g., sports centre, healthy circuits) for physical activity and participated in PA courses (1-credit electives) offered as part of the degree credits for all academic programmes at the university.

Student's external motivations and barriers for PA practice Motivators such as sharing healthy behaviours with friends and family were important motivators for students to adopt and maintain the behaviour change.

Among students who reported that family members and friends were physically active, 85.7% (12 out of 14) and 35.7% (5 out of 14), respectively, reported engaging in PA after the intervention.

"During the semester, I started to encourage my family to go outside and do some physical activity. We now play basketball on Sundays" (Female student, 21 years old).

Family members and friends played an important role in the adoption of students' PA habits.

Two elements were identified:

(1) previous PA habits of family members,

"I consider myself lucky to have my father as a role model, he is an excellent professional and an admirable athlete. (...) I grew up with a lifestyle where outdoor activities were plentiful and always involved some kind of physical effort. This made me love sport" (Female student, 21 years). (2) feedback students received from family members after adopting healthy habits

"I was able to influence some members of my family to exercise, first my mother started walking with me around the neighbourhood, then she was no longer able to walk, but my brother joined me" (Female student, 19 years).

Group activities were reported as motivational and social support elements to start and continue healthy habits.

Academic workload and lack of time were the main barriers to PA reported by students, consistent with the decrease in PA practices during the semester (i.e., final exams in November 2014 and May 2015).

"I don't exercise, I tried to exercise a month ago but I couldn't keep up the habit (...) Being at university is hard, I don't have time and also I've never had a routine" (Female student, 19 years old, 2015).

Healthy benefits

Improvements in fitness, appearance, stress, anxiety, sleep, smoking cessation, diet and weight control were the most commonly perceived benefits of PA by students.

"I like to exercise because it gives me energy and I do it to improve my fitness, (...) I think it is important to maintain healthy habits" (Female student, 18 years).

Students also reported that the health benefits of PA were associated with improved attitudes to preventive care for future medical practice.

"As a future doctor, I feel morally obliged to recommend physical activity because I now know its physical and mental health benefits (...) I want my patients to adopt and maintain a healthy lifestyle" (Male student, 19 years).

Discussion

We found that a one-semester curricular intervention was effective in improving the HRF variables studied in medical students. Cardiorespiratory fitness, body fat and muscular fitness improved significantly from baseline. These results are consistent with findings from intervention studies of similar duration. For example, Project Active, a randomised trial (n=235) comparing two interventions in a sedentary middle-aged adult population, showed an increase in estimated VO2max of 1.58 ml·kg-1·min-1 (95% CI: 0.93 to 2.22) for the lifestyle intervention group and 3.64 ml·kg-1·min-1 (95% CI: 2.98 to 4.29) for the structured exercise programme group after 24 weeks of intervention [38]. Compared to our study, we found a similar increase in estimated VO2max (3.11 ml·kg-1·min-1) after the 20th week of intervention. In addition, a systematic review [39] of health behaviour change programmes found a median net increase in estimated VO2max of 6.3% (interquartile range, 5.1–9.8%) in adult participants. The present study calculated a net increase in estimated VO2max of 8.9%, which is consistent with the data from the available studies.

Regarding changes in body composition, the Project Active study found a -1.4% (95% CI: -1.8 to -0.9) decrease in body fat percentage for the lifestyle intervention group and a -1.7% (95% CI: -2.1 to -1.2) decrease for the structured exercise programme group [40], similar to the decreases found in this study. Contrastingly, we did not find significant changes in BMI after the intervention. The unchanged BMI levels highlights the fact that the intervention did not lead to changes in body weight. However, this finding does not rule out beneficial changes in body composition, which are not considered by the BMI formula. The significant decrease in body fat percentage that we observed, along with the increase in muscle fitness and the stable BMI, indicate an increase in fat-free mass. Several studies have reported the same favorable changes in body composition (decreased fat mass and increased fat-free mass) with minimal or no change in BMI after controlled aerobic and resistance training programmes [41] usually lasting 15 to 20 weeks; these changes are usually small in magnitude [42]. Furthermore, weight loss was not expected in this population with healthy baseline BMI. Evidence suggests that high amounts of PA (≥250 min·week-1) plus dietary changes involving energy restriction are required to achieve weight loss [43]. According to the apparent doseresponse relationship between PA and weight loss, PA levels between 150 and 250 min-week-1 are associated with weight maintenance [43], which is consistent with our results.

In terms of muscular fitness, handgrip strength and abdominal muscle endurance increased significantly from baseline. On the one hand, resistance training programmes alone or in combination with aerobic exercise have been reported to increase handgrip strength in healthy medical students [44]. On the other hand, several studies lasting 7 to 24 weeks have reported improved sit-up performance after traditional sit-up/curl up programmes [45, 46], core stabilisation exercise programmes [46], integrated resistance training programmes with functional movement patterns [47], upper/total body resistance training programmes [48] and unstable resistance training programmes [49], but not after aerobic training alone [48]. Therefore, the students' reports of participation in both resistance and aerobic PA training are consistent with the results of the physical tests.

There is a paucity of studies on PA promotion among medical students. A systematic review [27] included 11 studies that evaluated the effectiveness of 10 PA counselling programmes for undergraduate medical students. There are several similarities between these programmes and our study. First, all studies in this area were guasiexperimental, reflecting trends and limitations in medical education research. Second, most interventions were incorporated into already established courses; as in this curricular intervention, the PA promotion strategy was incorporated into the medical physiology course. Thirdly, all studies included experiences or didactics of PA as a health behaviour. Fourthly, half of the studies asked students to report on their personal experience of PA. Three of the included studies asked students to change their own healthy habits, and only Frank et al. [50] required this habit to be PA. In addition, only the latter included pre- and post-intervention assessments.

Behavioural change theories and socio-ecological models are essential for effectively understanding and planning interventions to promote healthy lifestyles [51]. Bauman et al. propose individual, interpersonal and environmental dimensions as determinants of PA [52]. The qualitative methodology used in this study provided complementary information on the three dimensions described.

Our intervention resulted in the vast majority of students meeting the recommended amounts of PA, much more than the average medical student in Colombia [23] and higher than the PA levels of physicians and medical students in the United States [53–55] and the United Kingdom [56].

We found that students preferred to participate in group sports or activities, and that having an active family member or friend was an important motivator for engaging in PA. These findings add to the existing literature linking social support with increased PA in LMIC [57, 58], and specifically among medical students [15]. Promoting group sports and activities (such as aerobics classes, running clubs or football teams) may be an important element to include in future interventions for medical students.

Previous reviews have shown that lifestyle education for medical students needs to be an important part of the curriculum to have a positive impact on PA counselling [15, 27, 59]. In the present study, we showed that a 20-week intervention can be effective in promoting healthy lifestyles in relation to PA in second-year medical students. In a previous study, a lifestyle course delivered within 24 h was shown to be ineffective in improving health behaviours in medical students [60]. The frequency of PA among students in this study (i.e. an average of 3 days per week) is consistent with recent recommendations for health and fitness benefits. The position statement of the American College of Sports Medicine also states that healthy adults, such as our students, whose goal is to improve physical fitness and health, can also benefit from more intense advanced training techniques [61].

Our study found that the academic incentive was an effective motivator for compliance, similar findings have been documented [62]. However, it remains to be understood whether such an incentive produces lasting changes. In addition to improvements in HRF variables, our intervention provided perceived benefits such as a reduction in stress and an improvement in sleep patterns according to student self-reports. Previous research has shown that medical students have poorer mental health and perceive more stress than the general population [63, 64]. Research among undergraduates has found a negative relationship between PA levels, perceived stress and mental health [65, 66], which is consistent with our current findings. In addition, more active medical students were less likely to report symptoms of depression and anxiety [67]. Thus, these types of physical activity programmes appear to be an effective way to help students cope better with the stressors inherent in medical school, but more research is needed to better assess this claim [68].

Finally, one of the most commonly cited barriers to engagement in PA is lack of time, particularly among medical students [69–71]. Time constraints are particularly prevalent among medical students when considering stressful factors such as academic pressure, social issues and financial difficulties, which have been found to be higher compared to other undergraduate academic programmes and as academic careers progress [72, 73]. Nevertheless, time constraints for PA practice among medical students have been linked to a lack of adequate time management in similar studies [74, 75]. These findings highlight the importance of addressing time constraints by, for example, encouraging medical students to use more active transport, engage in more vigorous activities and make use of existing campus infrastructure.

Strengths and limitations Strengths of this study include the use of both quantitative and qualitative methods. Objective data such as body composition and fitness test performance were chosen as outcome variables for the quantitative analyses. Limitations include the lack of a control group for intervention comparison. Students' PA practices and perceptions of the curriculum were based on self-reported data. Findings on the benefits students perceived from PA (e.g. improved sleep patterns and better eating habits) were self-reported and there were no objective measures to confirm such benefits. Finally, the results may not be generalisable to other situations.

Conclusion

A curricular intervention was effective in improving second year medical students' HRF variables and promoting personal PA habits. Students were positively influenced by the curricular intervention to promote PA during the semester, which is expected to be maintained over time and to be reflected in favourable attitudes towards health promotion for clinical practice, prevention programmes and policy initiatives. These findings should encourage medical schools to incorporate curricular strategies to promote PA and related knowledge among medical students. Further evaluation is needed to assess long-term effects on HRF variables and personal PA behaviours, as well as counselling practices among medical students.

Abbreviations

PA	Physical activity
HRF	health-related fitness
NCDs	Non-communicable diseases
LMICs	low and middle-income countries
VO2max	maximum rate of oxygen consumption
BMI	Body Mass Index
20-MST	20-meter shuttle run test

Acknowledgements

We thank the students for their participation in this study. We are deeply indebted to Tatiana García-Betancourt for her contribution to the qualitative research component.

Author contributions

DCP, SCC and JD conceived the study and its design. DCP, SCC and MAJM contributed in acquisition of data and methodology. DCP, SCC, MAJM, AG, CMAP and JD participated in data analysis and interpretation. DCP, SCC, MAJM, AG and CMAP wrote the manuscript. All authors read and approved the final manuscript.

Funding

This work was supported by the Research Committee of the Faculty of Education and Faculty of Medicine from Universidad de los Andes in Bogotá, Colombia (Call for Interfaculty Research projects, 2015).

Data availability

The data used and/or analyzed in the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

The study was approved by the Ethics Committee of the Universidad de los Andes in Bogotá, Colombia (Act no. 538/2015). The study adhered to the Declaration of Helsinki. Written informed consent was obtained from all participants who agreed to participate in the study.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests. They have not received any honorarium, grant, or other form of payment for the preparation of this manuscript.

Received: 29 August 2023 / Accepted: 11 January 2024

Published online: 29 January 2024

References

- 1. Roth GA, Abate D, Abate KH, Abay SM, Abbafati C, Abbasi N, et al. Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980–2017: a systematic analysis for the global burden of Disease Study 2017. The Lancet. 2018;392(10159):1736–88.
- World Health Organization. Noncommunicable diseases country profiles 2018. Geneva: World Health Organization; 2018.
- Benziger CP, Roth GA, Moran AE. The global burden of Disease Study and the preventable burden of NCD. Glob Heart. 2016;11(4):393–7.
- Lee IM, Shiroma EJ, Lobelo F, Puska P, Blair SN, Katzmarzyk PT. Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. Lancet. 2012;380(9838):219–29.
- GBD 2017 Risk Factor Collaborators. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990–2017: a systematic analysis for the global burden of Disease Study 2017. Lancet. 2018;392(10159):1923–94.
- Kohl HW 3rd, Craig CL, Lambert EV, Inoue S, Alkandari JR, Leetongin G, et al. The pandemic of physical inactivity: global action for public health. Lancet. 2012;380(9838):294–305.
- World Health Organization. Tackling NCDs: 'best buys' and other recommended interventions for the prevention and control of noncommunicable diseases. Geneva: World Health Organization; 2017.
- Albert FA, Crowe MJ, Malau-Aduli AEO, Malau-Aduli BS. Physical activity promotion: a systematic review of the perceptions of healthcare professionals. Int J Environ Res Public Health. 2020;17(12).
- Huijg JM, Gebhardt WA, Verheijden MW, van der Zouwe N, de Vries JD, Middelkoop BJC, et al. Factors influencing primary health care professionals' physical activity promotion behaviors: a systematic review. Int J Behav Med. 2015;22(1):32–50.
- 10. Lobelo F, Stoutenberg M, Hutber A. The Exercise is Medicine Global Health Initiative: a 2014 update. Br J Sports Med. 2014;48(22):1627–33.
- Garrett S, Elley CR, Rose SB, O'Dea D, Lawton BA, Dowell AC. Are physical activity interventions in primary care and the community cost-effective? A systematic review of the evidence. Br J Gen Pract. 2011;61(584):e125–33.
- Sanchez A, Bully P, Martinez C, Grandes G. Effectiveness of physical activity promotion interventions in primary care: a review of reviews. Prev Med. 2015;76(Suppl):56–67.
- Garry JP, Diamond JJ, Whitley TW. Physical activity curricula in medical schools. Acad Med. 2002;77(8):818–20.
- Richard W, Stephen C, Ngaire C, Mark H, Emmanuel S. Physical activity education in the undergraduate curricula of all UK medical schools. Are tomorrow's doctors equipped to follow clinical guidelines? Br J Sports Med. 2012;46(14):1024.
- Wattanapisit A, Tuangratananon T, Thanamee S. Physical activity counseling in primary care and family medicine residency training: a systematic review. BMC Med Educ. 2018;18(1):159.
- Frenk J, Chen L, Bhutta ZA, Cohen J, Crisp N, Evans T, et al. Health professionals for a new century: transforming education to strengthen health systems in an interdependent world. Lancet. 2010;376(9756):1923–58.
- 17. Phillips E, Pojednic R, Polak R, Bush J, Trilk J. Including lifestyle medicine in undergraduate medical curricula. Med Educ Online. 2015;20:26150.
- Rea B, Worthman S, Shetty P, Alexander M, Trilk JL. Medical education transformation: Lifestyle medicine in undergraduate and graduate medical education, fellowship, and continuing medical education. Am J Lifestyle Med. 2021;15(5):514–25.
- Lobelo F, de Quevedo IG. The evidence in support of physicians and health care providers as physical activity role models. Am J Lifestyle Med. 2016;10(1):36–52.
- Selvaraj CS, Abdullah N. Physically active primary care doctors are more likely to offer exercise counselling to patients with cardiovascular diseases: a crosssectional study. BMC Prim Care. 2022;23(1):59.
- Binns HJ, Mueller MM, Ariza AJ. Healthy and fit for prevention: the influence of clinician health and fitness on promotion of healthy lifestyles during health supervision visits. Clin Pediatr. 2007;46(9):780–6.
- 22. Pasarica M, Harris DM, Franklin AL. Medical students teaching peer athletes: an innovative way of instructing the physiology of exercise, nutrition,

and sleep as fundamentals for lifestyle medicine. Adv Physiol Educ. 2017;41(3):432–5.

- Lobelo F, Duperly J, Frank E. Physical activity habits of doctors and medical students influence their counselling practices. Br J Sports Med. 2009;43(2):89–92.
- Morris A, Do D, Gottlieb-Smith R, Ng J, Jain A, Wright S, et al. Impact of a fitness intervention on medical students. South Med J. 2012;105(12):630–4.
- Gates A. Training tomorrow's doctors, in exercise medicine, for tomorrow's patients. Br J Sports Med. 2015;49(4):207–8.
- 26. Tovar G, López G, Ibáñez M, Alvarado R, Lobelo F, Duperly J. Institutionalized physical activity curriculum benefits of medical students in Colombia. Educ Health (Abingdon). 2016;29(3):203–9.
- Dacey ML, Kennedy MA, Polak R, Phillips EM. Physical activity counseling in medical school education: a systematic review. Med Educ Online. 2014;19:24325.
- 28. Creswell J, Plano V. Designing and conducting mixed methods research. Third edition. Thousand Oaks, California: Sage Publications; 2018.
- Warburton D, Jamnik V, Bredin S, Gledhill N. The physical activity readiness questionnaire for everyone (PAR-Q+) and electronic physical activity Readiness Medical Examination (ePARmed-X+). HFJC. 2011;4(2):3–17.
- World Health Organization. Obesity: Preventing and managing the global epidemic. Report of a WHO Consultation (WHO Technical Report Series 894). Geneva: World Health Organization; 2000.
- 31. Duperly J, Serrato M, Forero NI, Jimenez-Mora MA, Mendivil CO, Lobelo F. Validation of maximal, submaximal, and nonexercise indirect V $^{\circ}O_2$ max estimations at 2600 m altitude. High Alt Med Biol. 2020;21(2):135–43.
- Leger L, Mercier D, Gadoury C, Lambert J. The multistage 20 metre shuttle run test for aerobic fitness. J Sports Sci. 1988;6(2):93–101.
- Roberts HC, Denison HJ, Martin HJ, Patel HP, Syddall H, Cooper C, et al. A review of the measurement of grip strength in clinical and epidemiological studies: towards a standardised approach. Age Ageing. 2011;40(4):423–9.
- 34. Rubin DB. Estimating causal effects of treatments in randomized and nonrandomized studies. J Educ Psychol. 1974;66:688–701.
- Thorne S, Kirkham SR, O'Flynn-Magee K. The analytic challenge in interpretive description. Int J Qual Methods. 2004;3(1):1–11.
- Thorne S, Kirkham SR, MacDonald-Emes J. Interpretive description: a noncategorical qualitative alternative for developing nursing knowledge. Res Nurs Health. 1997;20(2):169–77.
- 37. Pitney W, Parker J. Qualitative research in physical activity and the health professions. Champaign, IL: Human Kinetics; 2009.
- Dunn AL, Garcia ME, Marcus BH, Kampert JB, Kohl HW, Blair SN. Six-month physical activity and fitness changes in project active, a randomized trial. Med Sci Sports Exerc. 1998;30(7):1076–83.
- Kahn EB, Ramsey LT, Brownson RC, Heath GW, Howze EH, Powell KE, et al. The effectiveness of interventions to increase physical activity. A systematic review. Am J Prev Med. 2002;22(4 Suppl):73–107.
- Dunn AL, Marcus BH, Kampert JB, Garcia ME, Kohl HW 3rd, Blair SN. Reduction in cardiovascular disease risk factors: 6-month results from project active. Prev Med. 1997;26(6):883–92.
- O'Donoghue G, Blake C, Cunningham C, Lennon O, Perrotta C. What exercise prescription is optimal to improve body composition and cardiorespiratory fitness in adults living with obesity? A network meta-analysis. Obes Rev. 2021;22(2):e13137.
- 42. Stiegler P, Cunliffe A. The role of diet and exercise for the maintenance of fat-free mass and resting metabolic rate during weight loss. Sports Med. 2006;36(3):239–62.
- Donnelly JE, Blair SN, Jakicic JM, Manore MM, Rankin JW, Smith BK. American College of Sports Medicine Position Stand. Appropriate physical activity intervention strategies for weight loss and prevention of weight regain for adults. Med Sci Sports Exerc. 2009;41(2):459–71.
- 44. Al-Asadi JN. Handgrip strength in medical students: correlation with body mass index and hand dimensions. Asian J Med Sci. 2018;9(1):21–6.
- 45. Learman K, Pintar J, Ellis A. The effect of abdominal strength or endurance exercises on abdominal peak torque and endurance field tests of healthy participants: a randomized controlled trial. Phys Ther Sport. 2015;16(2):140–7.
- Childs JD, Teyhen DS, Benedict TM, Morris JB, Fortenberry AD, McQueen RM, et al. Effects of sit-up training versus core stabilization exercises on sit-up performance. Med Sci Sports Exerc. 2009;41(11):2072–83.
- Distefano LJ, Distefano MJ, Frank BS, Clark MA, Padua DA. Comparison of integrated and isolated training on performance measures and neuromuscular control. J Strength Cond Res. 2013;27(4):1083–90.

- Kibele A, Behm DG. Seven weeks of instability and traditional resistance training effects on strength, balance and functional performance. J Strength Cond Res. 2009;23(9):2443–50.
- 50. Frank E, Elon L, Hertzberg V. A quantitative assessment of a 4-year intervention that improved patient counseling through improving medical student health. Medscape J Med. 2007;9(2):58.
- Sallis JF, Cervero RB, Ascher W, Henderson KA, Kraft MK, Kerr J. An ecological approach to creating active living communities. Annu Rev Public Health. 2006;27:297–322.
- Bauman A, Reis RS, Sallis JF, Wells JC, Loos RJ, Martin BW. Correlates of physical activity: why are some people physically active and others not? Lancet. 2012;380(9838):258–71.
- 53. Stanford FC, Durkin MW, Stallworth JR, Blair SN. Comparison of physical activity levels in physicians and medical students with the general adult population of the United States. Phys Sportsmed. 2013;41(4):86–92.
- Dyrbye LN, Satele D, Shanafelt TD. Healthy exercise habits are associated with lower risk of burnout and higher quality of life among U.S. medical students. Acad Med. 2017;92(7):1006–11.
- Frank E, Tong E, Lobelo F, Carrera J, Duperly J. Physical activity levels and counseling practices of U.S. medical students. Med Sci Sports Exerc. 2008;40(3):413–21.
- Blake H, Stanulewicz N, McGill F. Predictors of physical activity and barriers to exercise in nursing and medical students. J Adv Nurs. 2017;73(4):917–29.
- Del Díaz A, Sarmiento OL, Reis RS, Brownson RC. Translating evidence to policy: urban interventions and physical activity promotion in Bogotá, Colombia and Curitiba, Brazil. Transl Behav Med. 2011;1(2):350–60.
- Torres A, Sarmiento OL, Stauber C, Zarama R. The Ciclovia and Cicloruta programs: promising interventions to promote physical activity and social capital in Bogotá, Colombia. Am J Public Health. 2013;103(2):e23–30.
- Mastrocola MR, Roque SS, Benning LV, Stanford FC. Obesity education in medical schools, residencies, and fellowships throughout the world: a systematic review. Int J Obes. 2020;44(2):269–79.
- Malatskey L, Essa-Hadad J, Willis TA, Rudolf MCJ. Leading healthy lives: Lifestyle medicine for medical students. Am J Lifestyle Med. 2019;13(2):213–9.
- 61. Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee IM, et al. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: Guidance for prescribing exercise. Med Sci Sports Exerc. 2011;43(7):1334–59.
- DeVahl J, King R, Williamson JW. Academic incentives for students can increase participation in and effectiveness of a physical activity program. J Am Coll Health. 2005;53(6):295–8.
- Ludwig AB, Burton W, Weingarten J, Milan F, Myers DC, Kligler B. Depression and stress amongst undergraduate medical students. BMC Med Educ. 2015;15:141.

- Rosiek A, Rosiek-Kryszewska A, Leksowski Ł, Leksowski K. Chronic stress and suicidal thinking among medical students. Int J Environ Res Public Health. 2016;13(2):212.
- Fares J, Saadeddin Z, Al Tabosh H, Aridi H, El Mouhayyar C, Koleilat MK, et al. Extracurricular activities associated with stress and burnout in preclinical medical students. J Epidemiol Glob Health. 2016;6(3):177–85.
- Fares J, Al Tabosh H, Saadeddin Z, El Mouhayyar C, Aridi H. Stress, burnout and coping strategies in preclinical medical students. N Am J Med Sci. 2016;8(2):75–81.
- Baj-Korpak J, Zaworski K, Szymczuk E, Shpakou A. Physical activity and mental health of medical students from Poland and Belarus-countries with different restrictive approaches during the COVID-19 pandemic. Int J Environ Res Public Health. 2022;19(21).
- Leuchter RK, Stuber ML, McDonald AL, Croymans DM. Relationship between exercise intensity and stress levels among U.S. medical students. Med Educ Online. 2022;27(1):2027651.
- Awadalla NJ, Aboelyazed AE, Hassanein MA, Khalil SN, Aftab R, Gaballa II, et al. Assessment of physical inactivity and perceived barriers to physical activity among health college students, south-western Saudi Arabia. East Mediterr Health J. 2014;20(10):596–604.
- Williams AS, Williams CD, Cronk NJ, Kruse RL, Ringdahl EN, Koopman RJ. Understanding the exercise habits of residents and attending physicians: a mixed methodology study. Fam Med. 2015;47(2):118–23.
- Likus W, Milka D, Bajor G, Jachacz-Łopata M, Dorzak B. Dietary habits and physical activity in students from the Medical University of Silesia in Poland. Rocz Panstw Zakl Hig. 2013;64(4):317–24.
- Dyrbye LN, Thomas MR, Shanafelt TD. Systematic review of depression, anxiety, and other indicators of psychological distress among U.S. and Canadian medical students. Acad Med. 2006;81(4):354–73.
- Rotenstein LS, Ramos MA, Torre M, Segal JB, Peluso MJ, Guille C, et al. Prevalence of depression, depressive symptoms, and suicidal ideation among medical students: a systematic review and meta-analysis. JAMA. 2016;316(21):2214–36.
- Sajwani RA, Shoukat S, Raza R, Shiekh MM, Rashid Q, Siddique MS, et al. Knowledge and practice of healthy lifestyle and dietary habits in medical and non-medical students of Karachi, Pakistan. J Pak Med Assoc. 2009;59(9):650–5.
- Shaikh BT, Kahloon A, Kazmi M, Khalid H, Nawaz K, Khan N, et al. Students, stress and coping strategies: a case of Pakistani medical school. Educ Health. 2004;17(3):346–53.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.