

Search for vector-boson resonances decaying into a top quark and a bottom quark using pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector

ATLAS Collaboration

DOI:

[10.1007/JHEP12\(2023\)073](https://doi.org/10.1007/JHEP12(2023)073)

License:

Creative Commons: Attribution (CC BY)

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

ATLAS Collaboration 2023, 'Search for vector-boson resonances decaying into a top quark and a bottom quark using pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector', *JHEP*, vol. 2023, no. 12, 73.
[https://doi.org/10.1007/JHEP12\(2023\)073](https://doi.org/10.1007/JHEP12(2023)073)

[Link to publication on Research at Birmingham portal](#)

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Search for vector-boson resonances decaying into a top quark and a bottom quark using pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector



The ATLAS collaboration

E-mail: atlas.publications@cern.ch

ABSTRACT: A search for a new massive charged gauge boson, W' , is performed with the ATLAS detector at the LHC. The dataset used in this analysis was collected from proton-proton collisions at a centre-of-mass energy of $\sqrt{s} = 13$ TeV, and corresponds to an integrated luminosity of 139 fb^{-1} . The reconstructed tb invariant mass is used to search for a W' boson decaying into a top quark and a bottom quark. The result is interpreted in terms of a W' boson with purely right-handed or left-handed chirality in a mass range of 0.5–6 TeV. Different values for the coupling of the W' boson to the top and bottom quarks are considered, taking into account interference with single-top-quark production in the s -channel. No significant deviation from the background prediction is observed. The results are expressed as upper limits on the $W' \rightarrow tb$ production cross-section times branching ratio as a function of the W' -boson mass and in the plane of the coupling vs the W' -boson mass.

KEYWORDS: Beyond Standard Model, Hadron-Hadron Scattering

ARXIV EPRINT: [2308.08521](https://arxiv.org/abs/2308.08521)

Contents

1	Introduction	2
2	ATLAS detector	3
3	Data and simulated event samples	4
3.1	Signal and interference samples	5
3.2	Background samples for the 0-lepton channel	6
3.3	Background samples for the 1-lepton channel	6
4	Object reconstruction	7
5	Analysis strategy in the 0-lepton channel	10
5.1	Event selection	10
5.2	Event categorisation	11
5.3	Further categorisation	12
6	Analysis strategy in the 1-lepton channel	14
6.1	Event preselection	16
6.2	Event reconstruction	16
6.3	Event categorisation	17
7	Background estimation for the 0-lepton channel	21
8	Background estimation for the 1-lepton channel	22
9	Systematic uncertainties	24
9.1	Experimental uncertainties	25
9.2	Modelling uncertainties in background simulations	25
9.3	Uncertainties related to the data-driven background estimation	27
9.4	Uncertainty impact	27
10	Statistical analysis and results	29
11	Conclusions	36
	The ATLAS collaboration	44

1 Introduction

Multiple theories beyond the Standard Model (SM) involve enhanced symmetries that predict new gauge bosons, usually referred to as W' or Z' bosons. The W' boson is the mediator of a new charged vector current and can be massive enough to decay into a top quark and a bottom quark. Many models, such as those with extra dimensions [1], strong dynamics [2–5], or a composite Higgs boson [6], predict new vector charged-current interactions. Some models predict W' bosons that preferentially couple to third-generation particles [7–10] and are only observable in third-generation decay modes. Some of those models predict W' bosons that can only couple to quarks and are therefore not observable in leptonic decay modes [8, 9].

In the Sequential Standard Model (SSM) [11], an effective Lagrangian is used to capture the phenomenology of a W' boson decaying into a top quark and bottom quark ($W' \rightarrow tb$), which includes $W'^+ \rightarrow t\bar{b}$ and $W'^- \rightarrow \bar{t}b$ [12, 13]. This effective Lagrangian has a W' boson with the same coupling structure as the SM W boson. It has three free parameters: the mass of the W' boson, the chirality of the interaction, and an overall strength parameter that multiplies the fermion couplings of the new boson, which makes it possible to study W' bosons with different widths. This choice of Lagrangian has a wide applicability: in many beyond-the-SM theories predicting a W' boson, the top-quark phenomenology is independent of the light quarks because of its high mass.

Figure 1 shows the leading-order (LO) Feynman diagram for W' -boson production and its decay into tb . The top quark decays into a W boson and a bottom quark, with the W boson subsequently decaying either into quarks (figure 1(a), all-hadronic decay mode) or into a lepton and a neutrino (figure 1(b), lepton+jets decay mode). Two chirality scenarios are considered for the W' boson: right-handed chirality and left-handed chirality. A W' boson with right-handed chirality couples only to right-handed fermions. Its production and decay does not interfere with any SM processes. A W' boson with left-handed chirality couples only to left-handed fermions. Its production and decay interferes with the SM s -channel single-top-quark process, where the W boson replaces the W' boson in figure 1. In this paper, the reconstructed mass of the tb system is used to search for the W' -boson signal in both of these scenarios. In the absence of a signal, limits are set on the W' -boson production cross-section times branching ratio $W' \rightarrow tb$ as a function of the mass of the W' boson. For the scenario with right-handed chirality, the mass of the right-handed neutrino is assumed to be much higher than that of the right-handed W' boson, so the W' boson cannot decay leptonically [13]. As a result, the branching ratio for decay of a right-handed W' boson into tb is about 10% higher than that for a left-handed W' boson with the same mass, which can also decay into a lepton and a neutrino.

Searches for a W' boson decaying into tb have been performed at the Tevatron [14, 15] and the Large Hadron Collider (LHC) [16–23]. The most recent search by the CMS Collaboration, using $\sqrt{s} = 13$ TeV proton-proton (pp) collision data with an integrated luminosity of 137 fb^{-1} and targeting the all-hadronic decay mode, excluded a right-handed or left-handed W' boson with a mass below 3.4 TeV [23]. Previous searches by the ATLAS Collaboration for $W' \rightarrow tb$, using 36.1 fb^{-1} of pp collision data at $\sqrt{s} = 13$ TeV, excluded




Figure 1. Representative leading-order Feynman diagrams for s -channel W' -boson production with decay into tb , for (a) a top quark decaying into a W boson that decays hadronically (all-hadronic decay mode) and (b) a top quark decaying into a W boson that decays into a lepton and neutrino (lepton+jets decay mode).

a right-handed W' boson with a mass below 3.25 TeV by combining the all-hadronic and lepton+jets decay modes [21, 22] and assuming a W' -boson coupling equal to the SM W coupling.

This paper presents a search for W' bosons using the full Run 2 dataset collected by the ATLAS detector. Compared to the previous ATLAS analyses [21, 22], this search has improved top-quark and b -quark identification, better multi-jet background estimation and a refined selection strategy. The search is performed in both the all-hadronic (0-lepton) channel and the lepton+jets (1-lepton, either electron or muon) channel. Tau-leptons are not considered explicitly in either channel, and the electrons and muons are simply referred to as “leptons” in this paper. The 1-lepton analysis allows studies of the lower transverse momentum (p_T) region, which is out of reach for the 0-lepton channel’s trigger selection. In exchange, the 0-lepton channel provides optimal sensitivity to high W' masses.

The paper is organised as follows. The ATLAS detector at the LHC is described in section 2. Section 3 provides details of the data and simulated event samples. Object reconstruction is described in section 4. The analysis strategy, including event selection and categorisation, is described in sections 5 and 6 for the 0-lepton and 1-lepton channels, respectively. The background estimation for the two channels is described in sections 7 and 8. Systematic uncertainties considered in the statistical analysis are discussed in section 9. The results of the fit to data are presented in section 10, and section 11 provides the conclusions.

2 ATLAS detector

The ATLAS detector [24] at the LHC covers nearly the entire solid angle around the collision point.¹ It consists of an inner tracking detector surrounded by a thin superconducting

¹ATLAS uses a right-handed coordinate system with its origin at the nominal interaction point (IP) in the centre of the detector and the z -axis along the beam pipe. The x -axis points from the IP to the centre of the LHC ring, and the y -axis points upwards. Cylindrical coordinates (r, ϕ) are used in the transverse plane, ϕ being the azimuthal angle around the z -axis. The pseudorapidity is defined in terms of the polar angle θ as $\eta = -\ln \tan(\theta/2)$. Angular distance is measured in units of $\Delta R \equiv \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$.

solenoid, electromagnetic and hadron calorimeters, and a muon spectrometer incorporating three large superconducting air-core toroidal magnets.

The inner-detector system (ID) is immersed in a 2 T axial magnetic field and provides charged-particle tracking in the range $|\eta| < 2.5$. The high-granularity silicon pixel detector covers the vertex region and typically provides four measurements per track, the first hit normally being in the insertable B-layer (IBL) installed before Run 2 [25, 26]. It is followed by the silicon microstrip tracker, which usually provides eight measurements per track. These silicon detectors are complemented by the transition radiation tracker (TRT), which enables radially extended track reconstruction up to $|\eta| = 2.0$. The TRT also provides electron identification information based on the fraction of hits (typically 30 in total) above a higher energy-deposit threshold corresponding to transition radiation.

The calorimeter system covers the pseudorapidity range $|\eta| < 4.9$. Within the region $|\eta| < 3.2$, electromagnetic calorimetry is provided by barrel and endcap high-granularity lead/liquid-argon (LAr) calorimeters, with an additional thin LAr presampler covering $|\eta| < 1.8$ to correct for energy loss in material upstream of the calorimeters. Hadron calorimetry is provided by the steel/scintillator-tile calorimeter, segmented into three barrel structures within $|\eta| < 1.7$, and two copper/LAr hadron endcap calorimeters. The solid angle coverage is completed with forward copper/LAr and tungsten/LAr calorimeter modules optimised for electromagnetic and hadronic energy measurements respectively.

The muon spectrometer comprises separate trigger and high-precision tracking chambers measuring the deflection of muons in a magnetic field generated by the superconducting air-core toroidal magnets. The field integral of the toroids ranges between 2.0 and 6.0 T m across most of the detector. Three layers of precision chambers, each consisting of layers of monitored drift tubes, cover the region $|\eta| < 2.7$, complemented by cathode-strip chambers in the forward region, where the background is highest. The muon trigger system covers the range $|\eta| < 2.4$ with resistive-plate chambers in the barrel, and thin-gap chambers in the endcap regions.

Interesting events are selected by the first-level trigger system implemented in custom hardware, followed by selections made by algorithms implemented in software in the high-level trigger [27]. The first-level trigger accepts events from the 40 MHz bunch crossings at a rate below 100 kHz, which the high-level trigger reduces in order to record events to disk at about 1 kHz.

An extensive software suite [28] is used in data simulation, in the reconstruction and analysis of real and simulated data, in detector operations, and in the trigger and data acquisition systems of the experiment.

3 Data and simulated event samples

This analysis is performed using data from pp collisions at $\sqrt{s} = 13$ TeV collected with the ATLAS detector during Run 2, from 2015 to 2018. After applying a number of criteria to ensure that the detector was in good operating condition [29], the data used in this analysis have an integrated luminosity of 139 fb^{-1} .

Monte Carlo (MC) event generators were used to simulate signal and background events. The generation of all simulated event samples includes the effect of multiple pp interactions per bunch crossing, as well as changes in detector response due to interactions in bunch crossings before or after the one containing the hard interaction, modelled by overlaying simulated inelastic events on the physics event. These two effects are referred to as pile-up. The simulated event samples were processed with the GEANT4-based ATLAS detector simulation [30, 31].

All samples are weighted to match the pile-up distribution observed in data and are processed with the same reconstruction algorithms as data [32].

3.1 Signal and interference samples

Signal events were generated at LO in QCD with MADGRAPH5_AMC@NLO 2.6.7 [33], using a chiral W' -boson model that implements the effective Lagrangian described in section 1. In this model the coupling strength of the W' boson to right- or left-handed fermions (g') can be freely scaled relative to the SM coupling (g) by an arbitrary factor. Only purely right-handed or purely left-handed W' bosons are considered. The right-handed W' boson cannot decay to leptons because the right-handed neutrino is assumed to be more massive than the W' boson. MADGRAPH5_AMC@NLO was also used to decay the top quark and W boson, with spin correlations taken into account. PYTHIA 8.244 [34] was used for the modelling of the parton shower, fragmentation and underlying event. The PDF4LHC15 set of parton distribution functions (PDF) [35] and a set of tuned parameters called the A14 tune [36] were used for the event generation. Signal samples were normalised to the next-to-leading-order (NLO) cross-section computed by ZTOP [13]. Several contributions to the uncertainty in the NLO cross-section are considered for each coupling value. An uncertainty accounting for missing higher-order terms is estimated by doubling and halving both the renormalisation and factorisation scales independently. Uncertainties associated with the choice of PDF set and strong coupling constant value are obtained using the PDF4LHC15 PDF set. Finally, an uncertainty due to the choice of top-quark mass value (172.5 GeV) is obtained by raising and lowering the chosen value by 1 GeV. The NLO/LO cross-section normalisation ratios (K -factors) range from 1.3 to 1.4, depending on the mass of the W' boson. The width of the W' boson is about 3% of its mass for a coupling equal to the SM coupling ($g'/g = 1$) and scales with $(g'/g)^2$; it was calculated at NLO with ZTOP.

Signal samples were generated in 0.5 TeV steps for W' -boson masses between 0.5 and 6.0 TeV for lepton+jets top-quark decays and between 1.5 and 6.0 TeV for all-hadronic top-quark decays. Samples corresponding to right-handed and left-handed chiralities were produced separately. The coupling in the event generation was set to $g'/g = 2.0$. Weights were computed by MADGRAPH5_AMC@NLO during the parton-level event generation to reweight each sample to coupling values between $g'/g = 0.1$ and $g'/g = 0.5$ in steps of 0.1 and between $g'/g = 0.5$ and $g'/g = 5.0$ in steps of 0.5. Masses below 1.5 TeV were not generated in the all-hadronic case because the trigger selection utilised in the 0-lepton channel is completely inefficient in that region of phase space.

Interference between left-handed W' -boson production and SM single-top-quark production in the s -channel was modelled by reweighting the nominal signal samples using a parameterisation of the ratio of W' boson production to the interference contributions as a function of the parton-level invariant mass of the $t\bar{b}$ system [37]. The interference effects are destructive on the low-mass side of the W' mass peak and constructive on the high-mass side. Their size and relative importance depends strongly on the mass and coupling values considered, but their effect on the results shown in this paper is small.

3.2 Background samples for the 0-lepton channel

The dominant SM background process for all-hadronic events is QCD multi-jet production. This background is estimated with data-driven methods as described in section 7. The second most important background is top-quark-pair production ($t\bar{t}$), with an inclusive cross-section of 832 ± 51 pb for a top-quark mass of 172.5 GeV, as obtained from calculations at next-to-next-to-leading order (NNLO) in QCD including the resummation of next-to-next-to-leading logarithmic (NNLL) soft-gluon terms with TOP++ 2.0 [38–44]. Other small backgrounds, such as V +jets ($V = W$ or Z boson) or single-top production, are accounted for in the data-driven multi-jet estimate.

The production of $t\bar{t}$ events was modelled using the POWHEG BOX v2 [45–48] generator at NLO with the NNPDF3.0NLO [49] PDF set and the h_{damp} parameter² set to 1.5 times the mass of the top quark [50]. The events were passed to PYTHIA 8.230 to model the parton shower, hadronisation, and underlying event, with parameter values set according to the A14 tune and using the NNPDF2.3LO PDF set [51]. The decays of bottom and charm hadrons were performed by EVTGEN 1.6.0 [52].

Two $t\bar{t}$ background contributions are considered: all-hadronic $t\bar{t}$ events where both W bosons decay into quarks, resulting in a signature that is similar to the signal, and non-all-hadronic $t\bar{t}$ events where at least one of the W bosons decays leptonically. These events can contribute to the background in two ways: when none of the charged leptons from W -boson decays are identified or through hadronic tau decays in $W \rightarrow \tau\nu$ events.

The modelling of the $t\bar{t}$ background is improved by correcting the $t\bar{t}$ samples so that the top-quark p_{T} distribution matches that predicted at NNLO in QCD and NLO EW accuracy [53]. The corrections entail an implicit change in the PDF set and top-quark mass value considered with respect to those used in sample generation. The NNLO differential calculations are performed using the NNPDF3.0QED PDF set and a top-quark mass of 173.3 GeV.

3.3 Background samples for the 1-lepton channel

The largest background in the lepton+jets channel is $t\bar{t}$ production, already described for the 0-lepton channel in section 3.2. Other important backgrounds arise from V +jets production, especially W +jets in which the W boson decays leptonically. Other subdominant backgrounds such as single-top-quark and multi-boson production were also considered.

²The h_{damp} parameter is a resummation damping factor and one of the parameters that controls the matching of POWHEG matrix elements to the parton shower and thus effectively regulates the high- p_{T} radiation against which the $t\bar{t}$ system recoils.

Finally, a small multi-jet contribution was also taken into account and was estimated with data-driven methods as described in section 8.

The production of V +jets was simulated with the SHERPA 2.2.11 [54] generator using NLO matrix elements for up to two partons, and LO matrix elements for up to four partons, calculated with the Comix [55] and OPENLOOPS [56–58] libraries. They were matched with the SHERPA parton shower [59] using the MEPS@NLO prescription [60–63] with the set of tuned parameters developed by the SHERPA authors. The NNPDF3.0NNLO set of PDFs was used and the samples were normalised to the NNLO prediction [64].

The three single-top production modes (s -channel, t -channel, and tW -channel) were considered. They were modelled with the POWHEG BOX v2 [46–48, 65] generator at NLO in QCD, using the five-flavour scheme (four-flavour scheme for t -channel production) and the corresponding NNPDF3.0NLO set of PDFs. The events were interfaced with PYTHIA 8.230, which used the A14 tune and the NNPDF2.3LO set of PDFs. The diagram removal scheme [66] was used to remove interference and overlap between the tW -channel and $t\bar{t}$ production.

Samples of diboson final states (VV) were simulated with the SHERPA 2.2.1 generator, including off-shell effects and Higgs boson contributions where appropriate. Fully leptonic final states (where both bosons decay leptonically) and lepton+jets final states (where one decays leptonically and the other hadronically) were generated using matrix elements at NLO accuracy in QCD for up to one additional parton emission and at LO accuracy for up to three additional parton emissions. Samples for loop-induced $gg \rightarrow VV$ processes were generated using matrix elements calculated at LO accuracy for up to one additional parton emission. The matrix element calculations were matched and merged with the SHERPA parton shower based on Catani-Seymour dipole factorisation [55, 59] using the MEPS@NLO prescription. The virtual QCD corrections were provided by the OPENLOOPS library. The NNPDF3.0NNLO set of PDFs was used, along with the dedicated set of tuned parton-shower parameters developed by the SHERPA authors.

4 Object reconstruction

The signal process $W' \rightarrow tb$ targeted in this search results in a final state with a high- p_T top quark and a high- p_T b -quark. The b -quark is reconstructed as a small-radius (small- R) jet, while the reconstruction of the top quark depends on the decay mode of the W boson from the top-quark decay. In the 0-lepton channel, the W boson decays hadronically, and the top quark is reconstructed as a high- p_T large-radius (large- R) jet. In the 1-lepton channel, the W boson decays leptonically, and the top quark is reconstructed from the lepton (electron or muon), the missing transverse momentum, and a small- R jet.

For each event, collision vertices are reconstructed from inner-detector tracks with $p_T > 0.5$ GeV. The primary vertex in each event is chosen to be the one with the largest sum of the squared transverse momenta of all associated tracks.

Large- R jets are built from three-dimensional topological clusters of energy deposits in the calorimeter, which are calibrated to the hadronic energy scale with the local cluster weighting (LCW) procedure [67]. The anti- k_t [68, 69] algorithm with radius parameter

$R = 1.0$ is used to reconstruct large- R jets. These jets are trimmed [70] to reduce contributions from pile-up and soft interactions by reclustering the jet constituents into subjects using the k_t algorithm [71, 72] with a radius parameter $R = 0.2$ and discarding constituents belonging to subjects with p_T less than 5% of the p_T of the parent jet. The large- R jet four-momentum is then recomputed from the four-momenta of the remaining constituents and corrected using simulation and data [73]. Only large- R jets with $|\eta| < 2.0$ and $p_T > 500$ GeV are considered in this analysis.

Large- R jets are identified as containing a hadronically decaying top quark (henceforth called a top-tagged jet) using a multivariate classification algorithm implemented as a deep neural network (DNN) [74]. In most of the kinematic region of interest in the 0-lepton channel, a single large- R jet captures the top-quark decay products, resulting in a characteristic three-prong substructure within the jet, in contrast to a typical one-prong substructure associated with jets in multi-jet background processes. The DNN uses multiple features of the jet as inputs, e.g. calibrated jet p_T and mass, information about the dispersion of the jet constituents such as N-subjettiness [75], splitting scales [76], and energy correlation functions [77]. A DNN score between zero and one is obtained, with top-quark-initiated jets having values close to one and light-parton-initiated jets (including gluon jets) having values close to zero.

The top-tagging algorithm used in this analysis is optimised for top-quark-initiated jets that satisfy the “contained” criteria, where most of the top-quark decay products are contained inside the large- R jet [78]. The criteria are defined in the simulation as follows. First, trimmed large- R jets are built at particle level from all stable particles (with $c\tau > 10$ mm), excluding muons and neutrinos, and using a radius parameter $R = 1.0$. This trimmed particle-level jet must be matched to a generator-level top quark within $\Delta R < 0.75$ and have a mass larger than 140 GeV. At least one b -hadron must be associated with the jet [79]. Finally, a detector-level large- R jet is considered contained if it is within $\Delta R = 0.75$ of such a particle-level jet.

Two different efficiency working points, based on the DNN score, are used to define the signal regions in the 0-lepton channel: one in which the requirements correspond to a top-tagging efficiency of 80% (DNN score cut of ~ 0.6 – 0.7 , depending on p_T , to keep the efficiency constant), and a tighter one in which they correspond to an efficiency of 50% (DNN score cut of ~ 0.9 , depending on p_T , to keep the efficiency constant). Both efficiencies are calculated using simulated $t\bar{t}$ events. The corresponding light-jet rejection factors are between 10 and 40 (80% working point) and between 30 and 150 (50% working point) depending on p_T . Scale factors are used to correct for possible efficiency differences between simulated event samples and data [74]. Two additional efficiency working points are used to define control regions and to estimate the multi-jet background in the 0-lepton channel. A DNN score boundary of e^{-4} is used to divide the events in the control regions used for background estimation into two roughly equal-size samples, and a very loose DNN score cut of e^{-7} is used in the definition of the top-proxy jets that are used in the multi-jet background estimation (section 5.2). These working points have efficiencies higher than 95% and light-jet rejection factors ranging between 1.5 and 2.5 approximately.

Small- R jets are reconstructed by applying the anti- k_t algorithm with a radius parameter $R = 0.4$ to inner-detector tracks associated with the primary vertex and calorimeter clusters selected by a particle-flow reconstruction algorithm [80]. An energy calibration is applied to both the input calorimeter clusters [67] and the final reconstructed jets [81]. The latter takes into account both pile-up effects and flavour dependencies. Only small- R jets with $|\eta| < 2.5$ and $p_T > 25$ GeV are considered in this analysis. To reject jets arising from pile-up, a jet-vertex-tagging technique using a multivariate likelihood [82] is applied to jets with $p_T < 60$ GeV, ensuring that selected jets are matched to the primary vertex.

Small- R jets are identified as containing a b -hadron (henceforth called b -tagged) using the “DL1r” algorithm [83, 84]. This algorithm is based on a multivariate classification technique with a DNN combining information about the impact parameters of tracks and topological properties of secondary and tertiary decay vertices reconstructed from the tracks associated with the jet. In this analysis, the b -tagged jets are selected by using a working point corresponding to an efficiency of 85% for identifying true b -jets in simulated $t\bar{t}$ events. Light-jet rejection factors range between 20 and 50, depending on p_T [84]. Scale factors are used to correct for possible differences between the b -tagging efficiencies in simulated events and data events [83, 85, 86].

A third kind of jet is used in the 1-lepton channel to reject $t\bar{t}$ events containing hadronically decaying top quarks without using the previously defined traditional large- R jets. They are obtained by reclustering [87] small- R jets (passing the aforementioned selection) with a variable- R anti- k_t algorithm with a density parameter ρ of 350 GeV and a maximum radius of 1.0 [88]. Since the inputted constituent small- R jets are fully calibrated, their calibration and uncertainties can be propagated directly to the reclustered jet, and no further calibration step is necessary. In order to suppress contributions from pile-up and soft radiation, the reclustered variable- R jets are trimmed by removing all associated small- R jets that have p_T below 5% of the p_T of the reclustered jet. These jets, henceforth referred to as vRC-jets, are required to have $p_T > 100$ GeV and $|\eta| < 2.0$.

Electron candidates are reconstructed from energy deposits in the electromagnetic (EM) calorimeter that are matched to charged-particle tracks in the ID [89]. They are required to satisfy $p_T > 25$ GeV and $|\eta| < 2.47$, excluding the transition region between the barrel and endcap EM calorimeters ($1.37 < |\eta| < 1.52$). They are identified using the “tight” likelihood identification operating point [89]. The number of hits in the innermost pixel layer, the IBL, is used to discriminate between electrons and converted photons, and the longitudinal impact parameter z_0 relative to the primary vertex is required to satisfy $|z_0 \sin(\theta)| < 0.5$ mm. The significance of the transverse impact parameter d_0 must satisfy $|d_0/\sigma_{d_0}| < 5$. Electrons are also required to be isolated from other activity in the tracking and calorimeter systems, using the “FCTight” isolation working point [89]. The isolation criteria must be satisfied in a cone of size $\Delta R = 0.2$ around the electron in the calorimeter and a cone of p_T -dependent size in the ID. The latter choice improves the performance for electrons produced in the decay of high- p_T particles.

Muon candidates are reconstructed from matching tracks in the ID and the muon spectrometer, refined by a global fit which makes use of the hits in both subdetectors [90]. Muons must have $p_T > 25$ GeV and $|\eta| < 2.5$, and satisfy the “medium” identification

criteria [90]. Like the electrons, their longitudinal impact parameter is required to satisfy $|z_0 \sin(\theta)| < 0.5$ mm. The significance of the transverse impact parameter d_0 must satisfy $|d_0/\sigma(d_0)| < 3$. Muons are required to be isolated from other activity in the tracking system, using the “TightTrackOnly” isolation working point [90]. Similarly to the electrons, the isolation criterion must be satisfied in a cone of p_T -dependent size around the muon in the ID.

For both the electrons and muons, correction factors are applied to compensate for differences between data and simulation in trigger, reconstruction efficiency, particle identification, and isolation, usually as a function of relevant kinematic variables.

To resolve any reconstruction ambiguities between electrons, muons and jets, an overlap removal procedure is applied in a prioritised sequence as follows. First, if an electron shares the same ID track with another electron, the electron with lower p_T is discarded. Any electron sharing the same ID track with a muon is rejected. Next, jets are rejected if they lie within $\Delta R = 0.2$ of an electron. Similarly, jets within $\Delta R = 0.2$ of a muon are rejected if the jet has fewer than three associated tracks or if the muon is matched to the jet through ghost association [79]. Finally, electrons that are close to a remaining jet are discarded if their distance from the jet is $\Delta R < 0.4$, while for muons the distance is $\Delta R < \min(0.4, 0.04 + 10 \text{ GeV}/p_T)$.

The missing transverse momentum \vec{p}_T^{miss} , with magnitude E_T^{miss} , is calculated as the negative vectorial sum of the transverse momenta of all reconstructed physics objects (electrons, muons and jets) [91] and the soft term. The soft term includes all tracks associated with the primary vertex but not matched to any reconstructed physics object. Only tracks associated with the primary vertex are considered, improving the E_T^{miss} resolution by suppressing the effect of pile-up.

5 Analysis strategy in the 0-lepton channel

Events containing at least one high- p_T large- R jet and one high- p_T small- R jet that do not overlap are selected, according to the decay products in the all-hadronic decay mode of the tb final state. Events are separated into signal, control, validation and template regions based on the properties of the event’s large- R and small- R jets. Signal regions are signal-enriched regions, with a top-tagged large- R jet and a b -tagged small- R jet. Template and control regions are used to estimate the multi-jet background. Template regions are used to obtain the initial shape of the reconstructed tb mass distribution of the multi-jet background. Control regions are used to normalise those templates and obtain the final background distributions as well as to estimate their uncertainty. The validation region is used to validate the background estimation method.

5.1 Event selection

Events are first selected at the trigger level by requiring at least one large- R jet with p_T exceeding a threshold which depends on the data-taking year: 360 GeV for 2015, 420 GeV for 2016, and 460 GeV for 2017 and 2018. In order to perform the analysis in the regime where the trigger selection is fully efficient, events are required to have at least one reconstructed large- R jet with $p_T > 500$ GeV.




Figure 2. Flow chart of the event categorisation, starting from events containing one or more large- R jets that meet the preselection criteria. If that jet is top-tagged, using the 80% efficiency working point, it is a top-candidate jet. If no top-candidate jet is found in the event, each large- R jet becomes a top-proxy jet as long as it has a DNN score of at least e^{-7} . The b -candidate jet is the small- R jet that is back-to-back with a top-candidate jet or top-proxy jet. Top-proxy jets not back-to-back with a b -candidate jet are skipped.

Events with noise bursts or coherent noise in the calorimeters are removed, as are events containing large energy deposits from non-collision or cosmic sources of background. Events without a reconstructed primary vertex are rejected.

Events containing charged leptons (electron or muon) are removed to ensure orthogonality to the 1-lepton channel. The definition of lepton candidates described in section 4 is used for this veto, with the exception of the isolation requirement, which is dropped.

5.2 Event categorisation

Selected events are categorised into regions according to the procedure outlined in figure 2, separately for events where the large- R jet is b -tagged and for those where it is not, as described in section 5.3.

First, the number of top-candidate jets in an event is checked. A top-candidate jet is defined as a large- R jet with $p_T > 500$ GeV that is top-tagged using the 80% efficiency working point. Events with more than one such jet are vetoed, which reduces $t\bar{t}$

contamination. Events with exactly one top-candidate jet are kept and are considered for the signal regions, the validation region, and the template regions.

Events with no top-candidate jet are considered for the control regions. For this type of event, all large- R jets with $p_T > 500$ GeV and a top-tagging DNN score higher than e^{-7} are considered. This minimal top-tagging DNN score requirement removes less than 5% of events in data. The removed events show a flavour composition different from the rest when studied in simulation. Removing these events makes the flavour composition of jets more uniform in the control regions. The large- R jets thus defined are referred to as top-proxy jets in the following. They are a good representation of the light-flavour and gluon-initiated jets that form the multi-jet background in the signal region.

Next, an attempt to find a b -candidate jet is made. Pairs consisting of a large- R jet and a small- R jet are formed using all of the top-candidate and top-proxy jets selected in the previous step. For each top-candidate or top-proxy jet (J), the leading small- R jet (j) with $\Delta\phi(j, J) > 2.0$ and $p_T(j) > 500$ GeV is found. If an event has no pair formed in this way, it is rejected. The small- R jet thus selected is the b -candidate jet for that specific top-candidate or top-proxy jet.

Events are also rejected if the pseudorapidity difference, $|\Delta\eta|$, between the top-candidate (or top-proxy) jet and its associated b -candidate jet is greater than 2.0. This selection requirement reduces the background from t -channel multi-jet processes, which are important in the high- p_T regime.

Events with a top-candidate jet and a b -candidate jet that is b -tagged are assigned to the signal region SR1, SR2 or SR3, or the validation region VR. Which region they are assigned to depends on the DNN score of the large- R jet and the presence of additional b -tagged jets. This assignment is detailed in section 5.3 and figure 3.

Events with a top-candidate jet and a b -candidate jet that is not b -tagged but otherwise satisfies all criteria outlined above are assigned to the template regions TR1–TR4. These events are used to estimate the shape of the reconstructed m_{tb} distribution of the multi-jet background.

Finally, events with at least one top-proxy jet are separated similarly. Pairs consisting of a top-proxy jet and a b -candidate jet are assigned to different regions according to whether the b -candidate jet is b -tagged: if it is, then the pair is assigned to control regions 1 (CR1a or CR1b); if not, then the pair is assigned to control regions 2 (CR2a or CR2b). Each pair of top-proxy and b -candidate jets is categorised independently for events with more than one such configuration. By using all possible pairs, any bias that could arise by having to choose only one large- R jet as the top-proxy jet is avoided without reducing the number of pairs available in the CR. The statistical correlations introduced by this choice are negligible.

The variable of interest in this analysis is the reconstructed mass of the top-quark-bottom-quark system, m_{tb} , which is defined in all cases as the invariant mass of the top-candidate (or top-proxy) jet and its associated b -candidate jet.

5.3 Further categorisation

The regions defined in section 5.2 are further refined, taking advantage of the top-tagging properties of the large- R jet and the presence of b -tagged jets in the event. In particular, the




Figure 3. Events in the 0-lepton channel are categorised according to the top-tagging and b -tagging status of the large- R jets selected as top-candidate or top-proxy jets, and the b -tagging status of the small- R jets selected as b -candidate jets. Events are assigned to the right grid or the left grid, depending on whether the top-tagged or top-proxy jet is also b -tagged (1- b -tag-in-top category) or not (0- b -tag-in-top category). Events with exactly one top-candidate jet are assigned to one of the top two rows depending on whether the top-candidate jet also fulfils the 50% efficiency top-tagging working point. In the bottom two rows, each top-proxy jet from events without a top-candidate jet is considered. Events are further separated into columns based on the b -candidate jet: the right column if it passes the b -tag requirement, the left column if it does not.

top-quark decay leads to a bottom quark, which can be reconstructed and identified with the b -tagging algorithm. Events are therefore separated into those where the top-candidate jet is b -tagged (1- b -tag-in-top category), and those where it is not (0- b -tag-in-top category). A top-candidate (or top-proxy) jet J is considered b -tagged if at least one b -tagged small- R jet (j) is close to it ($\Delta R(j, J) < 1.0$). This corresponds to the expected signature of a $W' \rightarrow tb$ decay, characterised by the presence of one large- R jet that is top-tagged and b -tagged and is back-to-back with a small- R jet that is b -tagged. The remaining events are in the 0- b -tag-in-top category; they include some signal events where the b -quark from the top-quark decay is not identified. The background estimation procedure is performed separately for these two categories (see figure 3).

In both the 0- b -tag-in-top and 1- b -tag-in-top categories, events are further categorised according to the top-tagging DNN score of the top-candidate (or top-proxy) jet and the b -tagging score of the b -candidate jet. Signal-, validation- and template-region events in each category are assigned to one of the regions in the upper half of figure 3. Events with a top-candidate jet passing the 50% efficiency working point are assigned to the upper row, while events with a top-candidate jet failing the 50% efficiency working point but passing the 80% efficiency working point are assigned to the second row from the top. Events in the signal regions and the validation region, with a b -tagged b -candidate jet, populate the top-right quadrant of each category, while events in the template regions populate the top-left quadrant.

Every pair consisting of a top-proxy jet and a b -candidate jet from events without a top-candidate jet is assigned to one of the control regions in the lower half of each category. Pairs in which the DNN score of the top-proxy jet is above e^{-4} are assigned to the third row from the top, while the rest are assigned to the bottom row. Pairs with a b -tagged b -candidate jet are assigned to a region in the right column of each category, while events without one are assigned to a region in the left column.

The resulting regions are used in different ways:

- SR1, SR2 and SR3 are those where the signal-to-background ratio is the largest and the ones to be used in the statistical analysis described in section 10.
- VR is used to validate the data-driven multi-jet background estimation described in section 7.
- TR1, TR2, TR3 and TR4 provide the initial template for the multi-jet background in SR1, SR2, SR3 and VR respectively.
- CR1a and CR2a are used to obtain the multi-jet background in SR1 and SR2, while CR3a and CR4a are used to obtain the same background in SR3 and VR. They are also used to assess its uncertainty.
- CR1b and CR2b are used to assess the uncertainty in the multi-jet background in regions SR1 and SR2, while CR3b and CR4b are used for the same uncertainty in regions SR3 and VR.

The distribution of the reconstructed m_{tb} in each of the three signal regions is shown in figure 4 for selected simulated signal samples with a right-handed W' boson and a coupling value of $g'/g = 1$. The distribution peaks at the W' -boson mass, but exhibits a tail to lower m_{tb} that is more pronounced for higher W' -boson masses. The tail is due to the fact that, when the W' pole-mass is high, the PDF values for producing on-shell W' bosons are suppressed relative to the ones for producing low-mass off-shell W' bosons. The product of fiducial acceptance and selection efficiency is shown in figure 5 for the three signal regions and the right- and left-handed chirality scenarios with a coupling value of $g'/g = 1$. The fraction of W' -boson signal events in the template regions is small. Its impact on the background estimate is less than 3.5% of the background for a W' -boson mass of 4 TeV when the signal cross-section is normalised to the expected limit at 4 TeV. It is negligible compared to the systematic uncertainties of the signal and background. The signal contamination in the control regions is negligible.

6 Analysis strategy in the 1-lepton channel

Events containing exactly one isolated lepton, two or more jets and a certain amount of E_T^{miss} are selected, based on the expected decay products of the tb final state in the lepton+jets decay mode. Events passing these preselection requirements are categorised into different regions based on the number of jets, the number of b -tagged jets, and other kinematic variables.




Figure 4. Reconstructed m_{tb} distributions for the right-handed W' -boson signal with a coupling value of $g'/g = 1$ in (a) signal region 1, (b) signal region 2 and (c) signal region 3 of the 0-lepton channel. Distributions are normalised to unit area. The first and last bin in each distribution includes the underflow and overflow, respectively.




Figure 5. The product of fiducial acceptance and selection efficiency for the three signal regions of the 0-lepton channel as a function of the mass of the W' boson, for W' bosons with (a) right-handed chirality and (b) left-handed chirality. The W' boson's coupling strength is set to $g'/g = 1$.

6.1 Event preselection

Events are selected using a combination of single-lepton and E_T^{miss} triggers [27]. The E_T^{miss} triggers are only considered for events with a reconstructed $E_T^{\text{miss}} > 200$ GeV to ensure 100% efficiency of the trigger selection. The single-lepton triggers require the presence of a muon or an electron with p_T higher than a certain threshold and, in some cases, impose identification and lepton-isolation requirements. The lowest p_T threshold was 24 (20) GeV for electrons (muons) during the 2015 data-taking period and 26 GeV for both the electrons and muons in the data-taking periods from 2016 to 2018. A trigger-matching requirement is applied to the reconstructed lepton, which must be within $\Delta R = 0.1$ of the corresponding object at the trigger level [27]. The addition of E_T^{miss} triggers offsets a small loss of signal efficiency that occurs for the muon trigger.

Events with noise bursts or coherent noise in the calorimeters are removed, as are events containing large energy deposits from non-collision or cosmic sources of background. Events without a reconstructed primary vertex are rejected.

Events are required to contain one lepton with $p_T > 50$ GeV and $|\eta| < 2.47$, and no additional lepton with $p_T > 30$ GeV and $|\eta| < 2.47$. Electrons in the transition region between the barrel and endcap EM calorimeters ($1.37 < |\eta| < 1.52$) are not considered. Events are required to contain two or more jets with $p_T > 30$ GeV and $|\eta| < 2.5$. Finally, events are required to have $E_T^{\text{miss}} > 100$ GeV. These lepton selection criteria ensure that the trigger selection has a high efficiency for signal events, generally above 95%.

6.2 Event reconstruction

Selected events contain exactly one lepton, missing transverse momentum, and at least two jets. These objects are used to reconstruct the W' boson and the intermediate top quark and leptonically decaying W boson from its decay, as shown in the right diagram of figure 1. The neutrino is reconstructed starting from the missing transverse momentum in the event. Assuming that all of the E_T^{miss} in an event is carried by the neutrino, $p_{x,\nu}$ and $p_{y,\nu}$ are given by the x - and y -component of the \vec{p}_T^{miss} . The $p_{z,\nu}$ component is estimated by requiring that the squared sum of the lepton and neutrino four-momenta must yield the W -boson mass, which results in a quadratic equation. The possible solutions for $p_{z,\nu}$ are given by

$$p_{z,\nu}^{\pm} = \frac{\mu \cdot p_{z,\ell}}{p_{T,\ell}^2} \pm \sqrt{\frac{\mu^2 \cdot p_{z,\ell}^2}{p_{T,\ell}^4} - \frac{E_\ell^2 \cdot (E_T^{\text{miss}})^2 - \mu^2}{p_{T,\ell}^2}},$$

with

$$\mu = \frac{m_W^2}{2} + \cos \Delta\phi \cdot p_{T,\ell} \cdot p_{T,\nu}.$$

In these formulae, m_W is set to 80.4 GeV, $p_{T,\nu}$ is the transverse momentum of the neutrino and $\Delta\phi$ is the azimuthal angle between the charged lepton and the reconstructed \vec{p}_T^{miss} . The p_z , transverse momentum, and energy of the charged lepton are given by $p_{z,\ell}$, $p_{T,\ell}$ and E_ℓ , respectively.

If there are two real solutions for $p_{z,\nu}$, the one with the smaller absolute value is chosen. If the radicand is negative, the imaginary solution is avoided by multiplying \vec{p}_T^{miss} by a factor

chosen to make the radicand exactly zero. This adjustment satisfies $m_{\text{T}}^W = m_W$, where m_{T}^W is the transverse mass of the reconstructed W boson, and results in a single real solution.

The W boson is reconstructed as the sum of the four-vectors of the lepton and the neutrino. The top quark is then reconstructed by combining the W boson with one of the jets, without considering b -tagging. The jet j that provides the invariant mass of the Wj system closest to the top-quark mass ($m_{\text{top}} = 172.5$ GeV) is chosen and is referred to as b_{top} . Events with $p_{\text{T}}^{\text{top}} \leq 200$ GeV are rejected. Finally, the jet with the highest transverse momentum not selected as b_{top} is added to the top quark to obtain the reconstructed W' boson and its mass m_{tb} . This jet is referred to as $b_{W'}$ in the following. Events with $p_{\text{T}}^{b_{W'}} \leq 200$ GeV or $m_{tb} \leq 500$ GeV are rejected. This simple method to reconstruct and identify the tb candidate provides a W' -boson mass peak with good resolution without any efficiency reduction.

6.3 Event categorisation

Events selected and reconstructed as described in the previous subsections are categorised into regions based on the reconstructed objects. Both lepton flavours, electron and muon, are kept together in the same region. Signal, validation and control regions are defined by selecting events with two or three jets, one or two of which are required to be b -tagged, resulting in four possible combinations. These regions are referred to as 2j1b, 3j1b, 2j2b and 3j2b. Additional region-specific requirements are imposed to further suppress SM backgrounds:

- A requirement of $m_{\text{T}}^W > 20$ GeV on the transverse mass of the reconstructed W boson in regions with one b -tagged jet suppresses multi-jet events.
- In regions with three jets, events in which the third jet (neither the b_{top} nor the $b_{W'}$) is b -tagged are rejected. This requirement reduces the $t\bar{t}$ background where the third jet is more likely to originate from a bottom quark.
- In regions with three jets, events are rejected if they contain a reclustered jet with mass close to the top quark ($140 \text{ GeV} < m_{\text{vRC-jet}} < 200 \text{ GeV}$). This requirement removes both the W +jets and $t\bar{t}$ background where the third jet is less likely to be b -tagged.
- In regions with two jets and one b -tagged jet, only events in which the $b_{W'}$ is b -tagged are kept. This requirement reduces the $t\bar{t}$ background, which has a high fraction of events in which the b_{top} is b -tagged but the $b_{W'}$ is not.

Each of the four initial regions is further divided into signal, validation and control regions. The signal regions are referred to as SR 2j1b, SR 3j1b, SR 2j2b and SR 3j2b and are defined by two additional requirements:

- The separation between the top-quark decay products in the (η, ϕ) plane is required to be small, as expected for a high- p_{T} top quark. A requirement of $\Delta R(\ell, b_{\text{top}}) < 1.0$ is imposed.

Regions	SR	CR_{W+jets}	VR_{W+jets}	$VR_{t\bar{t}}$
	SR 2j1b, SR 2j2b, SR3j1b, SR3j2b	CR 2j1b, CR 3j1b	VR 2j1b, VR 3j1b	VR 2j2b, VR 3j2b
Trigger	E_T^{miss} OR one-lepton			
N_{jets}	2,3			
N_{b-jets}	1,2	1	1	2
p_T^{lepton}	> 50 GeV			
E_T^{miss}	> 100 GeV			
m_T^W (in 1-tag)	> 20 GeV			
$p_T^{b_{W'}}$	> 200 GeV			
p_T^{top}	> 200 GeV			
m_{tb}	> 500 GeV			
$ \Delta\eta(\text{top}, b_{W'}) $	< 2.0	n/a	n/a	n/a
$\Delta R(\ell, b_{top})$	< 1.0	> 1.5, ≤ 2.4	> 1.0, ≤ 1.5	> 1.0, ≤ 2.4
b -tagging (2-jet regions)	$b_{W'}$ is b -tagged			
b -tagging (3-jet regions)	third jet is not b -tagged			
vRC-jet (3-jet regions)	veto events with $140 \text{ GeV} < m_{\text{vRC-jet}} < 200 \text{ GeV}$			

Table 1. Definition of the signal, control and validation regions in the 1-lepton channel.

- The pseudorapidity difference between the top quark and the $b_{W'}$ is required to satisfy $|\Delta\eta(\text{top}, b_{W'})| < 2.0$. This requirement rejects $t\bar{t}$ and multi-jet backgrounds, where the top quark and the $b_{W'}$ are expected to be well separated.

Control regions for the W +jets background are defined using the same jet-multiplicity criteria as the signal regions (two or three jets) and requiring one of those jets to be b -tagged. Signal events are suppressed by requiring that the lepton and the b_{top} are separated in ΔR . As the distance $\Delta R(\ell, b_{top})$ increases, the amount of signal and $t\bar{t}$ background decreases, while the amount of W +jets background increases. Events are assigned to the control regions if they satisfy $1.5 < \Delta R(\ell, b_{top}) \leq 2.4$. These regions are referred to as CR 2j1b and CR 3j1b and are orthogonal to the signal regions.

Events with the same jet multiplicity and b -jet multiplicity as in CR 2j1b or CR 3j1b but which satisfy $1.0 < \Delta R(\ell, b_{top}) \leq 1.5$ are assigned to validation regions for the W +jets background. These regions are referred to as VR 2j1b and VR 3j1b and are orthogonal to both the signal regions and the control regions.

Validation regions are also defined for the $t\bar{t}$ background by selecting events with two or three jets, two of which are required to be b -tagged. Orthogonality to the signal regions is maintained by requiring $1.0 < \Delta R(\ell, b_{top}) \leq 2.4$. These regions are referred to as VR 2j2b and VR 3j2b.

A summary of the region definitions is given in table 1, while a schematic view is shown in figure 6.




Figure 6. Events in the 1-lepton channel are categorised according to the number of jets and b -tagged jets in the event. Events are assigned to signal, control or validation regions depending on the angular separation between the lepton and jet used to reconstruct the top-quark candidate (b_{top}).

The distribution of the reconstructed m_{tb} in the four signal regions is shown in figure 7 for selected simulated signal samples with a right-handed W' boson and a coupling value of $g'/g = 1$. The behaviour is similar to the 0-lepton case, with distributions peaking around the W' -boson mass and a long tail towards lower masses. The product of fiducial acceptance and selection efficiency for the same regions is shown in figure 8. After an initial rise due to threshold effects for a W' -boson mass of 500 GeV, the efficiency drops as the mass increases. This is mainly caused by b -tagging and lepton efficiency dropping as a function of p_T . As transverse momentum increases and the angular distance between the top-quark decay products is reduced, the efficiency to identify isolated leptons degrades accordingly. For high W' -boson masses the relative importance of the W' peak becomes small due to reconstruction and PDF effects, as can be seen in figure 7. When that happens the low mass tail dominates the efficiency calculation, causing it to increase slightly. The fraction of W' -boson signal events is negligible in both the validation and control regions.




Figure 7. Reconstructed m_{tb} distributions for the right-handed W' -boson signal with coupling value of $g'/g = 1$ in (a) signal region 2j1b, (b) signal region 3j1b, (c) signal region 2j2b and (d) signal region 3j2b. Distributions are normalised to unit area. The first and last bin in each distribution includes the underflow and overflow, respectively.




Figure 8. The product of fiducial acceptance and selection efficiency for the signal regions of the 1-lepton channel as a function of the mass of the W' boson, for W' bosons with (a) right-handed chirality and (b) left-handed chirality. The W' boson coupling strength is set to $g'/g = 1.0$.

7 Background estimation for the 0-lepton channel

The dominant background in the 0-lepton channel, from multi-jet production, is estimated using a data-driven method that predicts both the shape and normalisation of the multi-jet m_{tb} distribution in the signal and validation regions. The initial template for the multi-jet background in each signal region, SR j in figure 3 ($j = 1, 2, 3$), is the m_{tb} histogram in the corresponding template region, TR j in figure 3 ($j = 1, 2, 3$). TR4 is used to obtain the template in the VR. The correct normalisation for each template in the target region is obtained by multiplying each bin in the template histograms by the ratio $N_{\text{CR1a}}^{\text{obs}}/N_{\text{CR2a}}^{\text{obs}}$ obtained in the same bin of regions CR1a and CR2a or the corresponding ratio $N_{\text{CR3a}}^{\text{obs}}/N_{\text{CR4a}}^{\text{obs}}$ obtained in CR3a and CR4a. This ratio represents the number of top-candidate jets for which the b -candidate jet is b -tagged divided by the number of top-candidate jets for which the b -candidate jet is not b -tagged. It is expected to not depend strongly on the top-tagging criteria for the top-candidate. This ratio is obtained using the pairs composed of a top-proxy jet and a b -candidate jet in the control regions indicated with the letter “a” in figure 3, CR j a, which are completely dominated by multi-jet events. The pairs of jets in the control regions form the same kinematic relationship as the pairs of jets in the signal and validation regions. This equivalence allows the ratio to be used to scale the m_{tb} distribution from the template to the signal and validation regions. The ratio varies between 0.14 (0.15) at low m_{tb} , around 1 TeV, and 0.19 (0.21) at high m_{tb} , above 5 TeV, for the 0- b -tag-in-top (1- b -tag-in-top) category.

The $t\bar{t}$ background is subdominant and is estimated using the simulated event samples described in section 3. It is non-negligible in the signal and template regions because it contains two b -hadrons. The predicted $t\bar{t}$ background ($N^{t\bar{t}}$) is subtracted from data (N^{obs}) in the template regions to obtain the multi-jet background template. The small background from V +jets is similar in flavour composition to the multi-jet background and is thus accounted for by the data-driven multi-jet background estimate.

The data-driven estimate of the multi-jet background in bin i of m_{tb} in each of the signal regions and the validation region is then given by

$$N_{\text{SR1,SR2}}^{\text{data-driven background}}(i) = R_{\text{corr}}^1(i) \times \left(N_{\text{TR1,TR2}}^{\text{obs}}(i) - N_{\text{TR1,TR2}}^{t\bar{t}}(i) \right) \times \frac{N_{\text{CR1a}}^{\text{obs}}(i)}{N_{\text{CR2a}}^{\text{obs}}(i)} \quad (7.1)$$

and

$$N_{\text{SR3,VR}}^{\text{data-driven background}}(i) = R_{\text{corr}}^0(i) \times \left(N_{\text{TR3,TR4}}^{\text{obs}}(i) - N_{\text{TR3,TR4}}^{t\bar{t}}(i) \right) \times \frac{N_{\text{CR3a}}^{\text{obs}}(i)}{N_{\text{CR4a}}^{\text{obs}}(i)}. \quad (7.2)$$

The correction factors $R_{\text{corr}}^{1,0}$ take into account possible correlations between the top-tagging of the top-candidate jet and the b -tagging of the b -candidate jet in the 1- b -tag-in-top and 0- b -tag-in-top categories, respectively. The nominal value is $R_{\text{corr}} = 1$ because the correlations are small. This is verified in simulated multi-jet samples by computing the corresponding ratio of yields, for example $(N_{\text{SR1}}N_{\text{CR2a}})/(N_{\text{TR1}}N_{\text{CR1a}})$ for SR1. Deviations from unity are considered as uncertainties of the method and are described in section 9.

In order to mitigate the impact of the smaller number of data events in the tails of the m_{tb} distributions in the control regions, bins are merged from high to low m_{tb} to ensure

	SR1	SR2	SR3	VR
Data-driven	23190 ± 520	94000 ± 2000	76100 ± 500	297000 ± 1900
All-hadronic $t\bar{t}$	4700 ± 830	6400 ± 1600	1090 ± 170	1400 ± 300
Non-all-hadronic $t\bar{t}$	1220 ± 180	1730 ± 350	330 ± 46	445 ± 87
Total background	29000 ± 1000	102400 ± 2400	77560 ± 550	299000 ± 2000
W' ($m = 3.0$ TeV)	500 ± 100	390 ± 85	260 ± 58	190 ± 41
Data	29220	100383	78407	298727

Table 2. Predicted and observed event yields for the signal regions and the validation region of the 0-lepton channel before the fit to data. The uncertainty in each estimate is the sum in quadrature of the systematic uncertainties from all sources described in section 9. Signal yields correspond to the theoretical prediction for a 3 TeV right-handed W' boson with a coupling strength of $g'/g = 1.0$.

that the statistical uncertainty in each bin of the b -tagging ratio is less than 5%. The same operation is done when calculating the correction factor R_{corr} .

The expected and observed event yields in the three signal regions and the validation region are shown in table 2, together with the predicted yields for a W' boson with a mass of 3 TeV, right-handed chirality and a coupling value of $g'/g = 1.0$. The uncertainty in each estimate is the sum in quadrature of the systematic uncertainties from all sources described in section 9. The uncertainty in the $t\bar{t}$ background is large, mostly due to the uncertainty in the theory modelling. The data-driven background uncertainty is small thanks to the very large event yields in data. It is larger in regions SR1 and SR2 than in SR3 and VR due to two main factors: the larger $t\bar{t}$ background propagated through the data-driven method and the presence of b -tagged jets inside the top-candidate (top-proxy) jet, which increases the impact of the flavour composition on the correlation factor.

8 Background estimation for the 1-lepton channel

The dominant background components in the 1-lepton channel are those from W +jets and $t\bar{t}$ production. They are estimated using the simulated event samples described in section 3. The subdominant diboson, Z +jets and single-top-quark processes are also estimated using simulated event samples. The small multi-jet contamination from jets misreconstructed as isolated leptons is estimated using a data-driven method known as the template method.

The initial template for the multi-jet distribution is obtained by defining “loose” regions with exactly the same selection requirements as those described in section 6.3 except for a looser lepton selection. The lepton selection is modified as follows to obtain regions enriched in multi-jet events:

- Electrons must pass the “medium” but not the “tight” likelihood identification requirements [92].
- Muons must pass the “loose” but not the “medium” identification requirements [93].

	SR 2j1b	SR 2j2b	SR 3j1b	SR 3j2b
$t\bar{t}$	856 ± 51	3910 ± 220	8150 ± 210	7480 ± 250
W +jets	3140 ± 170	329 ± 32	3600 ± 230	204 ± 22
Z +jets	205 ± 95	100 ± 44	380 ± 160	64 ± 28
Single-top-quark	300 ± 40	1130 ± 110	1660 ± 270	990 ± 140
Diboson	69 ± 28	13 ± 6	190 ± 77	15.7 ± 7.2
Multi-jet	89 ± 11	82 ± 37	179 ± 24	11 ± 11
Total background	4670 ± 220	5560 ± 290	14160 ± 490	8760 ± 310
W' ($m = 3.0$ TeV)	15.2 ± 1.2	42.8 ± 4.9	33.4 ± 3.6	51.7 ± 5.3
Data	5081	5150	14496	8060

Table 3. Predicted and observed event yields for the signal regions of the 1-lepton channel before the fit to data. The uncertainty in each estimate is the sum in quadrature of the systematic uncertainties from all sources described in section 9. Signal yields correspond to the theoretical prediction for a 3 TeV right-handed W' boson with a coupling strength of $g'/g = 1.0$.

- Electrons and muons must fail the isolation requirements described in section 4.

Templates are obtained in each loose region for two variables, m_{tb} and m_{T}^W , by subtracting the background components described above from data.

The distribution of m_{T}^W is used to obtain correction factors from a binned maximum-likelihood fit performed independently in each signal, control and validation region. In these m_{T}^W fits, the contributions from $t\bar{t}$ and W +jets, as well as the initial multi-jet template from the corresponding loose region, are allowed to float freely. The resulting multi-jet normalisation factors, one per region, are used to scale the corresponding loose multi-jet m_{tb} template and obtain the m_{tb} multi-jet distribution to be used in the statistical analysis. The m_{T}^W distribution is chosen for this method because multi-jet contribution's shape is different from that of other backgrounds and the bin-by-bin signal significance is extremely small, even for signal regions.

The expected and observed event yields in the signal, control and validation regions are shown in tables 3, 4 and 5 respectively, together with the predicted yields for a W' boson with a mass of 3 TeV, right-handed chirality and a coupling value of $g'/g = 1.0$. The uncertainty in each estimate is the sum in quadrature of the systematic uncertainties from all sources described in section 9. The amount of multi-jet background is small in all regions, particularly in those with two b -tagged jets, where it is compatible with zero.

	CR 2j1b	CR 3j1b
$t\bar{t}$	1386 ± 58	4940 ± 160
W +jets	7720 ± 470	6780 ± 530
Z +jets	150 ± 60	160 ± 66
Single-top-quark	640 ± 160	1380 ± 360
Diboson	168 ± 68	300 ± 120
Multi-jet	236 ± 26	273 ± 38
Total background	10300 ± 520	13700 ± 800
Data	11553	14431

Table 4. Predicted and observed event yields for the control regions of the 1-lepton channel before the fit to data. The uncertainty in each estimate is the sum in quadrature of the systematic uncertainties from all sources described in section 9.

	VR 2j1b	VR 3j1b	VR 2j2b	VR 3j2b
$t\bar{t}$	677 ± 36	6420 ± 160	3010 ± 180	6500 ± 240
W +jets	3730 ± 220	4160 ± 300	1340 ± 110	620 ± 68
Z +jets	100 ± 40	153 ± 64	82 ± 34	40 ± 17
Single-top-quark	330 ± 80	1640 ± 240	1660 ± 340	1370 ± 390
Diboson	83 ± 35	203 ± 83	31 ± 13	33 ± 14
Multi-jet	83 ± 9	195 ± 27	283 ± 55	92 ± 66
Total background	5000 ± 260	12770 ± 540	6410 ± 440	8650 ± 490
Data	5398	13091	6413	8310

Table 5. Predicted and observed event yields for the validation regions of the 1-lepton channel before the fit to data. The uncertainty in each estimate is the sum in quadrature of the systematic uncertainties from all sources described in section 9.

9 Systematic uncertainties

The modelling of signal, $t\bar{t}$, V +jets, single-top-quark, and diboson events described in section 3 is affected by experimental uncertainties related to the reconstruction and calibration of the physics objects. In addition, uncertainties in the theoretical modelling of the $t\bar{t}$, single-top-quark, and V +jets backgrounds are also taken into account.

In the 0-lepton channel, these uncertainties affecting the simulated backgrounds also affect the data-driven background estimate because they are propagated through the $t\bar{t}$ subtraction in the template regions. Additional sources of uncertainty affecting the data-driven background in the 0-lepton channel are considered in order to account for possible deviations from the core assumptions of the method described in section 7. The multi-jet background in the 1-lepton channel is small, so all uncertainties in this background are expected to be covered by a single normalisation uncertainty.

9.1 Experimental uncertainties

Uncertainties related to the energy scale and resolution of small- and large- R jets are evaluated by combining information about detector reconstruction performance in simulated events with *in situ* methods using data collected with ATLAS during LHC Run 2 [73, 81]. Uncertainties related to the mass scale of large- R jets are evaluated by using a forward-folding technique combining fits to the W -boson and top-quark mass peaks in order to extract both the mass scale and resolution differences between data and simulation [94]. This approach is complemented by the R_{trk} method [73]. A constant jet mass resolution uncertainty of 20% is assigned to the mass of large- R jets [73].

Uncertainties in the correction factors for the b -tagging identification response are derived from dedicated flavour-enriched samples in data. An additional term is included to extrapolate the measured uncertainties to the high- p_{T} region with jet $p_{\text{T}} > 400$ GeV. This term is calculated from simulated events by considering variations of the quantities affecting the b -tagging performance, such as the impact parameter resolution, percentage of poorly measured tracks, description of the detector material, and track multiplicity per jet. The dominant uncertainty affecting the extrapolation to high- p_{T} is related to the interactions of high- p_{T} b -hadrons in the innermost pixel layer, which were not considered in the simulation of the samples used for this analysis [83].

Uncertainties in the correction factors for the top-tagging identification are considered [95], taking into account effects on the selection and reconstruction of jets involved in the scale factor estimation. These uncertainties are obtained by taking into account uncertainties related to the jet energy scale and b -tagging, as well as MC generator uncertainties and statistical uncertainties. Additional uncertainties related to the modelling of the samples used for the scale factor estimation are also taken into account. Uncertainties are also considered for jets with $p_{\text{T}} > 800$ GeV in the extrapolation of the measured uncertainties to the high- p_{T} region.

Uncertainties are considered on the electrons energy scale and energy resolution, the muons momentum scale and resolution and on the data-to-MC correction factors applied to the trigger, reconstruction, identification and isolation efficiencies [89, 90]. Uncertainties are also considered on the soft term used in the $E_{\text{T}}^{\text{miss}}$ calculation and on the $E_{\text{T}}^{\text{miss}}$ energy scale and resolution [91].

Variations in the reweighting applied to simulated event samples to match the mean number of pp interactions observed in each bunch crossing in data are included. They cover the uncertainty in the ratio of the predicted and measured inelastic cross-sections. A constant 1.7% [32] normalisation uncertainty is applied to all simulated event samples to account for uncertainty in the combined 2015–2018 integrated luminosity, obtained using the LUCID-2 detector [96] for the primary luminosity measurements, complemented by measurements using the inner detector and calorimeters.

9.2 Modelling uncertainties in background simulations

For the 0-lepton channel, uncertainties in modelling the $t\bar{t}$ background are included. Other possible backgrounds are only included as part of the data-driven estimation. For the 1-

lepton channel, uncertainties in modelling the $t\bar{t}$, single-top-quark, and W +jets backgrounds are included.

Several uncertainties in the theoretical modelling of the $t\bar{t}$ background samples are considered. Systematic uncertainties due to the choice of parton shower and hadronisation model are evaluated by comparing the nominal $t\bar{t}$ sample with a sample produced with the POWHEG BOX v2 generator using the NNPDF3.0NLO PDF set. Events in the latter sample were passed to HERWIG 7.04 [97, 98], which used the H7UE set of tuned parameters [98] and the MMHT2014LO PDF set [99]. To assess the uncertainty in the matching of NLO matrix elements to the parton shower, the nominal POWHEG sample is compared with a sample of events generated with MADGRAPH5_AMC@NLO 2.6.0 [33] interfaced with PYTHIA 8.230. The MADGRAPH5_AMC@NLO calculation used the NNPDF3.0NLO PDF set, and PYTHIA 8 used the A14 tune and the NNPDF2.3LO PDF set. Before the comparisons, these samples were corrected to match the NNLO predictions of the top-quark p_T distribution using the procedure outlined in section 3.2. Systematic uncertainties associated with alternative choices of renormalisation and factorisation scales, in which their nominal values are varied by factors of 0.5 and 2.0, and with the choice of PDF set, which is changed to LUXQED+PDF4LHC15 [35, 100], are included by correcting the nominal sample to dedicated alternative NNLO calculations [53]. Additional uncertainties are calculated using internal weights associated with each event for alternative MC tune choices [36] corresponding to changes in the amount of initial-state and final-state radiation and in the modelling of multiple parton interactions.

For the three production modes contributing to the single-top-quark background, the uncertainty due to the parton shower and hadronisation model is evaluated by comparing the nominal sample of events with a sample where the events generated with the POWHEG BOX v2 generator are interfaced to HERWIG 7.04, which used the H7UE tune and the MMHT2014LO PDF set. To estimate the uncertainty in the matching of NLO matrix elements to the parton shower, the nominal samples are compared with samples generated with the MADGRAPH5_AMC@NLO 2.6.2 generator at NLO in QCD using the five-flavour scheme and the NNPDF2.3NLO PDF set. The events are interfaced with PYTHIA 8.230, which used the A14 tune and the NNPDF2.3LO PDF set. Additional uncertainties are considered for the choice of PDF set, analogous to the $t\bar{t}$ uncertainties. These are included by using weights related to variations of the NNPDF3.0NLO set and alternative baseline PDF sets, namely the MMHT2014NLO set [99] and the CT14NLO set [101]. Finally, the nominal tW -channel POWHEG+PYTHIA 8 sample is also compared with an alternative sample generated using the diagram subtraction scheme [50, 66] to estimate the uncertainty arising from the interference with $t\bar{t}$ production.

For the W +jets background in the 1-lepton channel, internal weights are used to consider alternative renormalisation and factorisation scale choices. Uncertainties related to the choice of PDF set are estimated using internal weights corresponding to the NNPDF3.0NNLO set and two alternative baseline PDF sets, CT18NNLO [102] and MSHT2020NNLO [103], as well as two variations of the NNPDF3.0NNLO set with different values of α_s . Uncertainties associated with the inclusion of approximate NLO electroweak corrections are included by using internal weights corresponding to different ways of combining the QCD and

electroweak contributions: additive, multiplicative, or exponentiated [104]. Each option is compared with the nominal prediction independently.

A 6% uncertainty is assigned to the $t\bar{t}$ normalisation in the 0-lepton channel, in accord with the inclusive cross-section calculation described in section 3. No overall normalisation uncertainty is assigned to the $t\bar{t}$ or W +jet inclusive cross-section for the 1-lepton channel as the normalisation of each of these background components is controlled by free-floating parameters in the final likelihood fit described in section 10. An overall normalisation uncertainty of 5% is assigned to the single-top-quark backgrounds to account for the inclusive cross-section uncertainty. A conservative 40% normalisation uncertainty is assigned to both Z +jets and diboson production to take into account any possible mismodelling in the production of additional jets [105] and heavy-flavour jets [106] in these minor backgrounds.

9.3 Uncertainties related to the data-driven background estimation

Dedicated uncertainties in the data-driven background estimation in the 0-lepton channel are obtained by measuring the correlation between the top-tagging DNN score of the top-candidate jet and the b -tagging score of the b -candidate jet directly in data, using the control regions defined in section 7.

The assumption of $R_{\text{corr}} = 1$ in eqs. (7.1) and (7.2) is replaced by the ratio of diagonal products,

$$R_{\text{corr}}^1(i) = \frac{N_{\text{CR1a}}^{\text{obs}}(i) \times N_{\text{CR2b}}^{\text{obs}}(i)}{N_{\text{CR2a}}^{\text{obs}}(i) \times N_{\text{CR1b}}^{\text{obs}}(i)}$$

and

$$R_{\text{corr}}^0(i) = \frac{N_{\text{CR3a}}^{\text{obs}}(i) \times N_{\text{CR4b}}^{\text{obs}}(i)}{N_{\text{CR4a}}^{\text{obs}}(i) \times N_{\text{CR3b}}^{\text{obs}}(i)},$$

for the 1- b -tag-in-top and 0- b -tag-in-top categories, respectively, to obtain the varied background estimates used to define the uncertainty. The counts $N_{\text{CR}j}(i)$ represent the number of events in bin i of the m_{tb} distribution in region $\text{CR}j$. These R_{corr} values deviate from unity as the correlation between top-tagging and b -tagging increases.

The background rejection factors for top-tagging and b -tagging decrease for high- p_T large- R and small- R jets, resulting in a dependence on m_{tb} . This can cause R_{corr} to deviate from unity. In addition, the flavour composition of the partons in multi-jet events changes as a function of m_{tb} . The uncertainty in R_{corr} is separated into two uncorrelated components, one for low masses ($m_{tb} \leq 2 \text{ TeV}$) and the other for high masses ($m_{tb} > 2 \text{ TeV}$), to account for these two effects. This uncertainty ranges from 1% at low m_{tb} values to approximately 5% for large m_{tb} in SR3. It takes slightly larger values for SR2 and SR1, with values up to 13% in the high m_{tb} region.

In the 1-lepton channel, the uncertainty in the normalisation factor for the multi-jet background described in section 8 is taken into account.

9.4 Uncertainty impact

In order to estimate the importance of the different categories of systematic uncertainties, their post-fit impact is calculated for right-handed W' bosons with various masses and $g'/g = 1.0$. The result of such an estimation in the combined fit of the two channels is

Uncertainty	$m(W') = 2 \text{ TeV}$	$m(W') = 4 \text{ TeV}$
Background modelling	0.36	0.42
$t\bar{t}$	0.21	0.11
W +jets	0.19	0.11
Multi-jet and data-driven background	0.16	0.37
Single-top-quark	0.20	0.21
Other processes	0.03	0.00
Instrumental	0.40	0.21
Top-tagging	0.14	0.16
Flavour-tagging	0.35	0.11
Large- R jets	0.04	0.00
Small- R jets	0.11	0.05
Other	0.01	0.00
Total systematic uncertainty	0.61	0.47
MC and data-driven bkg. statistics	0.25	0.42
Statistical uncertainty	0.75	0.79

Table 6. Post-fit fractional contributions of different uncertainty categories to the total uncertainty in the observed signal strength, as determined in the combined fit of the 0-lepton and 1-lepton channels. Different masses of a right-handed W' boson with $g'/g = 1.0$ are considered. For each category, the fit to data is repeated with the corresponding group of nuisance parameters fixed to their best-fit values. Each category's contribution is evaluated from the difference of the squares of the uncertainty of the original fit and the modified fit, by dividing the square root of this difference by the uncertainty of the original fit. The sum in quadrature is different from unity due to correlations among nuisance parameters in the fit.

shown in table 6 as the fractional contribution of each category to the total uncertainty in the observed signal strength. For each category, the fit to data is repeated with the corresponding group of nuisance parameters fixed to their best-fit values. Each category's contribution is evaluated from the difference of the squares of the uncertainty of the original fit and the modified fit, by dividing the square root of this difference by the uncertainty of the original fit. The sum in quadrature is different from one due to correlations among nuisance parameters in the fit.

The relative importance of systematic uncertainties falls with increasing W' -boson mass, and the measurement becomes very statistically dominated at large masses. The MC and data-driven background statistics category from table 6 is dominated by the statistical uncertainty of the data-driven background in the 0-lepton channel. Among the systematic uncertainties, background modelling uncertainties dominate mostly at large masses, while for lower masses the top- and flavour-tagging uncertainty components become more important.

10 Statistical analysis and results

In order to test for the presence of a massive resonance, templates in the variable m_{tb} obtained from the simulated signal samples, background samples, and data-driven predictions are fitted to the data. The fit uses a binned maximum-likelihood approach based on the ROOSTATS framework [107]. Separate fits are performed for each signal mass and chirality hypothesis. Each fit includes the three signal regions defined in section 5 for the 0-lepton channel and the six signal and control regions defined in section 6 for the 1-lepton channel, making a total of nine regions in which the m_{tb} distribution is fitted simultaneously.

The systematic uncertainties described in section 9 can change the acceptance, normalisation and shape of the m_{tb} distribution for the signal and the background processes. They are incorporated into the fit as nuisance parameters with a log-normal or Gaussian constraint. The signal and background expectations in each bin are functions of these nuisance parameters.

As already indicated in section 9, the normalisations of the $t\bar{t}$ and W +jets background components in the 1-lepton channel are allowed to float freely in the fit. Two independent normalisation factors are used for each of the W +jets and $t\bar{t}$ background components, one for the 2-jet regions and one for the 3-jet regions, making a total of four normalisation factors. This choice is motivated by the previous search [22], where significant differences in the modelling as a function of the jet multiplicity were observed. For the same reason, all modelling uncertainties in the 1-lepton channel are kept uncorrelated between regions with different jet multiplicities. Both the normalisation and modelling uncertainties are assumed to be correlated between regions with different b -tagged jet multiplicities but the same jet multiplicity. In the 0-lepton channel, all modelling uncertainties are kept fully uncorrelated between the three signal regions.

Given the different treatment of the modelling uncertainties in the different channels, these uncertainties are considered uncorrelated between the 0-lepton and 1-lepton channels. The same is true for the normalisation uncertainties present in the 0-lepton channel and the floating normalisations present in the 1-lepton channel. Experimental uncertainties, when present in both channels, are considered correlated.

The probability that the data are compatible with the background-only hypothesis is estimated by integrating the distribution of the log-likelihood ratio, approximated using the asymptotic formulae described in ref. [108]. In the absence of any significant excess above the expected background, upper limits at the 95% confidence level (CL) on the signal production cross-section times the $W' \rightarrow tb$ decay branching ratio are derived using the CL_s method [109].

For left-handed W' hypotheses, interference with s -channel single-top-quark production is included in the fit by changing the signal template shape. If the signal is scaled by a factor μ_s , the interference contribution is scaled by $\sqrt{\mu_s}$ and the signal template is modified correspondingly by the interference contribution [110].

The m_{tb} distribution in the validation region of the 0-lepton channel before the likelihood fit is shown in figure 9(a). There is good agreement between data and prediction, and the uncertainty becomes significant only for m_{tb} above 3 TeV. Figures 9(b)–9(d) show the m_{tb} distributions in the three signal regions of the 0-lepton channel after a background-only

fit to data. The pre-fit background prediction is also shown; it is very close to the post-fit background in all three regions. The maximum value of m_{tb} observed in data is 7.8 TeV, for an event in SR3.

The distributions of m_{tb} in the 1-lepton channel are shown in figure 10 for the two control regions and the two W +jets validation regions after a background-only fit to data. Agreement is good in all 1-lepton-channel regions, with the uncertainty remaining relatively small until values of m_{tb} are higher than 3.5 TeV. Figure 11 shows the distributions for the four signal regions. The post-fit normalisation factors for the $t\bar{t}$ background component have values of 0.89 ± 0.07 and 0.92 ± 0.04 for the 2-jet and 3-jet regions respectively. The corresponding factors for the W +jets background component are 1.19 ± 0.07 and 1.21 ± 0.11 for the 2-jet and 3-jet regions respectively.

Good agreement between the background prediction and data is observed in all regions. Upper limits on the production cross-section times decay branching ratio as a function of the W' -boson mass are therefore derived and are shown in figure 12 for a right-handed W' boson and in figure 13 for a left-handed W' boson. For each chirality, three different values of g'/g are used to generate upper limits. In all cases, the expected limit in each channel is shown in addition to the combination. The observed limits and expected limits are derived by linear interpolation between those obtained for several different signal mass hypotheses. Mass-limit values are obtained from the intersection of the limit curves with the theory curve, obtained at NLO using ZTOP [13]. For a right-handed W' boson, masses below 4.6 TeV (4.2 TeV) are observed (expected) to be excluded for a g'/g value of 1.0, while for a left-handed W' boson, masses below 4.2 TeV (4.1 TeV) are observed (expected) to be excluded for the same coupling value. The observed limits are higher than expected because of statistical fluctuations of the data around $m_{tb} = 4$ TeV in the signal regions of the 0-lepton channel. The lower exclusion limits obtained for a left-handed W' boson are partially explained by the higher branching ratio to tb in the right-handed scenario, where the W' boson cannot decay leptonically. The sensitivity to high W' -boson masses is limited by statistical uncertainties. For a right-handed W' boson with $g'/g = 1.0$, the expected mass limit is more than 1 TeV higher than in the previous combination of the two channels [22]. The mass limit for a left-handed W' boson is also a more than 1 TeV improvement on the previous 0-lepton-channel-only results [21].

Figures 14(a) and 14(b) show the observed and expected exclusion contours as functions of the W' -boson mass and coupling strength for the right-handed and left-handed hypotheses respectively. The interpolation between coupling values is performed using a quadratic function. In both figures, the expected limit in each channel is shown in addition to the combination. For low W' -boson masses, the 1-lepton channel dominates the sensitivity because the large multi-jet background reduces the sensitivity in the 0-lepton channel. For high W' -boson masses, the efficiency of the signal selection in the 1-lepton channel decreases due to the lepton isolation requirement, while the 0-lepton channel remains highly efficient. For very high coupling strengths, the width of the W' boson increases and the reconstructed signal peaks become very wide or disappear completely. In this scenario, the signal distributions shift towards lower m_{tb} values, making the 1-lepton channel competitive even at high W' masses.




Figure 9. Distribution of the reconstructed m_{tb} for data and backgrounds in the 0-lepton channel in (a) the validation region before the fit to data, and the three signal regions after the background-only fit to data: (b) signal region 1, (c) signal region 2 and (d) signal region 3. The bottom panel in each plot shows the ratio of data to the background sum. For the signal regions, the dashed blue line shows the pre-fit background sum, and in the bottom panel the ratio of pre-fit to post-fit background sum. The hatched band includes all of the systematic uncertainties (a) before and (b, c, d) after the fit to data. The dashed red line shows the distribution of the W' -boson signal for a mass of 3 TeV and a coupling strength of $g'/g = 1.0$, normalised to the predicted cross-section. The last bin in each distribution includes overflow. The blue arrows in the ratio panel indicate that the data point is outside the range shown.




Figure 10. Reconstructed $m_{t\bar{b}}$ distributions for data and backgrounds in the 1-lepton channel, (a) and (b) in the control regions, and (c) and (d) in the W +jets validation regions. They are shown after the background-only fit to data. Each bottom panel shows the ratio of data to the background sum. The dashed blue line shows the pre-fit background sum, and in the bottom panel the ratio of pre-fit to post-fit background sum. The hatched band includes all of the systematic uncertainties after the fit to data. The dashed red line shows the distribution of the W' -boson signal for a mass of 3 TeV and a coupling strength of $g'/g = 1.0$, normalised to the predicted cross-section. The last bin in each distribution includes overflow. The blue arrows in the ratio panel indicate that the data point is outside the range shown.




Figure 11. Reconstructed m_{tb} distributions for data and backgrounds in the four 1-lepton signal regions after the background-only fit to data. Each bottom panel shows the ratio of data to the background sum. For the signal regions, the dashed blue line shows the pre-fit background sum, and in the bottom panel the ratio of pre-fit to post-fit background sum. The hatched band includes all of the systematic uncertainties after the fit to data. The dashed red line shows the distribution of the W' -boson signal for a mass of 3 TeV and a coupling strength of $g'/g = 1.0$, normalised to the predicted cross-section. The last bin in each distribution includes overflow. The blue arrows in the ratio panel indicate that the data point is outside the range shown.




Figure 12. Observed and expected 95% CL limits on the cross-section times branching ratio for the production of a W' boson with decay into tb and right-handed couplings as a function of the mass of the W' boson and a coupling value of (a) $g'/g = 1.0$, (b) $g'/g = 0.5$ and (c) $g'/g = 2.0$. They are obtained from the combination of the 0-lepton and 1-lepton channels. The expected limits for the individual channels are also shown. The observed limits and expected limits are derived by linear interpolation between those obtained for several different signal mass hypotheses. The uncertainty in the theory prediction includes components from the factorisation and renormalisation scales, PDFs, strong coupling constant, and top-quark mass.




Figure 13. Observed and expected 95% CL limits on the cross-section times branching ratio for the production of a W' boson with decay into tb and left-handed couplings as a function of the mass of the W' boson and a coupling value of (a) $g'/g = 1.0$, (b) $g'/g = 0.5$ and (c) $g'/g = 2.0$. They are obtained from the combination of the 0-lepton and 1-lepton channels. The expected limits for the individual channels are also shown. The observed limits and expected limits are derived by linear interpolation between those obtained for several different signal mass hypotheses. The uncertainty in the theory prediction includes components from the factorisation and renormalisation scales, PDFs, strong coupling constant, and top-quark mass.




Figure 14. Observed and expected limits as a function of the coupling value and the W' -boson mass for (a) right-handed and (b) left-handed W' -boson couplings. They are obtained from the combination of the 0-lepton and 1-lepton channels. The expected limits for the individual channels are also shown. The area above the line is excluded.

11 Conclusions

A search for $W' \rightarrow tb$ using 139 fb^{-1} of $\sqrt{s} = 13 \text{ TeV}$ pp collision data collected with the ATLAS detector at the LHC is presented. The search combines two channels, named according to the targeted decay of the top quark. The 0-lepton channel employs a DNN-based algorithm to identify large-radius jets originating from hadronically decaying top quarks. They are combined with small-radius jets selected with a b -tagging algorithm to reconstruct the W' boson. The dominant background from multi-jet production is estimated using a data-driven method. The 1-lepton channel selects events with one lepton (electron or muon), a certain amount of E_T^{miss} , and two or more jets. These objects are combined using top-quark and W -boson mass constraints to reconstruct the W' boson. The dominant backgrounds come from $t\bar{t}$ and W +jets production.

The observed distributions of the reconstructed W' -boson mass in various analysis regions are consistent with the background-only prediction, and exclusion limits at 95% CL are set on the production cross-section times branching ratio for $W' \rightarrow tb$. Several signal hypotheses are considered: W' -boson masses in the range 0.5–6 TeV, right-handed and left-handed couplings, and different coupling strengths relative to the coupling of the W boson to fermions in the SM. Effects of interference between the left-handed W' boson and the SM W boson are taken into account.

Right-handed W' bosons with masses below 4.6 TeV (4.2 TeV) are observed (expected) to be excluded for a coupling value of $g'/g = 1.0$. For the same coupling value, left-handed W' bosons with masses below 4.2 TeV (4.1 TeV) are observed (expected) to be excluded. The observed mass limits for right-handed W' bosons with $g'/g = 1.0$ are more than 1 TeV higher than in the previous 0-lepton-channel CMS search and the previous ATLAS combination of the two channels. The observed mass limits for left-handed W' bosons with the same coupling strength are approximately 0.8 TeV higher than in the previous 0-lepton-channel CMS search. The obtained limits are the most stringent to date.

Acknowledgments

We thank CERN for the very successful operation of the LHC, as well as the support staff from our institutions without whom ATLAS could not be operated efficiently.

We acknowledge the support of ANPCyT, Argentina; YerPhI, Armenia; ARC, Australia; BMWFW and FWF, Austria; ANAS, Azerbaijan; CNPq and FAPESP, Brazil; NSERC, NRC and CFI, Canada; CERN; ANID, Chile; CAS, MOST and NSFC, China; Minciencias, Colombia; MEYS CR, Czech Republic; D NRF and DNSRC, Denmark; IN2P3-CNRS and CEA-DRF/IRFU, France; SRNSFG, Georgia; BMBF, HGF and MPG, Germany; GSRI, Greece; RGC and Hong Kong SAR, China; ISF and Benozziyo Center, Israel; INFN, Italy; MEXT and JSPS, Japan; CNRST, Morocco; NWO, Netherlands; RCN, Norway; MEiN, Poland; FCT, Portugal; MNE/IFA, Romania; MESTD, Serbia; MSSR, Slovakia; ARRS and MIZŠ, Slovenia; DSI/NRF, South Africa; MICINN, Spain; SRC and Wallenberg Foundation, Sweden; SERI, SNSF and Cantons of Bern and Geneva, Switzerland; MOST, Taiwan; TENMAK, Türkiye; STFC, United Kingdom; DOE and NSF, United States of America. In addition, individual groups and members have received support from BCKDF, CANARIE, Compute Canada and CRC, Canada; PRIMUS 21/SCI/017 and UNCE SCI/013, Czech Republic; COST, ERC, ERDF, Horizon 2020, ICSC-NextGenerationEU and Marie Skłodowska-Curie Actions, European Union; Investissements d’Avenir Labex, Investissements d’Avenir Idex and ANR, France; DFG and AvH Foundation, Germany; Herakleitos, Thales and Aristeia programmes co-financed by EU-ESF and the Greek NSRF, Greece; BSF-NSF and MINERVA, Israel; Norwegian Financial Mechanism 2014-2021, Norway; NCN and NAWA, Poland; La Caixa Banking Foundation, CERCA Programme Generalitat de Catalunya and PROMETEO and GenT Programmes Generalitat Valenciana, Spain; Göran Gustafssons Stiftelse, Sweden; The Royal Society and Leverhulme Trust, United Kingdom.

The crucial computing support from all WLCG partners is acknowledged gratefully, in particular from CERN, the ATLAS Tier-1 facilities at TRIUMF (Canada), NDGF (Denmark, Norway, Sweden), CC-IN2P3 (France), KIT/GridKA (Germany), INFN-CNAF (Italy), NL-T1 (Netherlands), PIC (Spain), ASGC (Taiwan), RAL (U.K.) and BNL (U.S.A.), the Tier-2 facilities worldwide and large non-WLCG resource providers. Major contributors of computing resources are listed in ref. [111].

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] K.R. Dienes, E. Dudas and T. Gherghetta, *Grand unification at intermediate mass scales through extra dimensions*, *Nucl. Phys. B* **537** (1999) 47 [[hep-ph/9806292](#)] [[INSPIRE](#)].
- [2] S. Weinberg, *Implications of dynamical symmetry breaking*, *Phys. Rev. D* **13** (1976) 974 [[INSPIRE](#)].
- [3] L. Susskind, *Dynamics of spontaneous symmetry breaking in the Weinberg-Salam theory*, *Phys. Rev. D* **20** (1979) 2619 [[INSPIRE](#)].

- [4] S. Dimopoulos and L. Susskind, *Mass without scalars*, *Nucl. Phys. B* **155** (1979) 237 [[INSPIRE](#)].
- [5] E. Eichten and K.D. Lane, *Dynamical breaking of weak interaction symmetries*, *Phys. Lett. B* **90** (1980) 125 [[INSPIRE](#)].
- [6] D.J. Muller and S. Nandi, *Top flavor: a separate SU(2) for the third family*, *Phys. Lett. B* **383** (1996) 345 [[hep-ph/9602390](#)] [[INSPIRE](#)].
- [7] G. Burdman, B.A. Dobrescu and E. Pontón, *Resonances from two universal extra dimensions*, *Phys. Rev. D* **74** (2006) 075008 [[hep-ph/0601186](#)] [[INSPIRE](#)].
- [8] E. Malkawi, T.M.P. Tait and C.P. Yuan, *A model of strong flavor dynamics for the top quark*, *Phys. Lett. B* **385** (1996) 304 [[hep-ph/9603349](#)] [[INSPIRE](#)].
- [9] J.C. Pati and A. Salam, *Lepton number as the fourth color*, *Phys. Rev. D* **10** (1974) 275 [*Erratum ibid.* **11** (1975) 703] [[INSPIRE](#)].
- [10] C.T. Hill, *Topcolor assisted technicolor*, *Phys. Lett. B* **345** (1995) 483 [[hep-ph/9411426](#)] [[INSPIRE](#)].
- [11] G. Altarelli, B. Mele and M. Ruiz-Altaba, *Searching for new heavy vector bosons in $p\bar{p}$ colliders*, *Z. Phys. C* **45** (1989) 109 [*Erratum ibid.* **47** (1990) 676] [[INSPIRE](#)].
- [12] Z. Sullivan, *Fully differential W' production and decay at next-to-leading order in QCD*, *Phys. Rev. D* **66** (2002) 075011 [[hep-ph/0207290](#)] [[INSPIRE](#)].
- [13] D. Duffy and Z. Sullivan, *Model independent reach for W' bosons at the LHC*, *Phys. Rev. D* **86** (2012) 075018 [[arXiv:1208.4858](#)] [[INSPIRE](#)].
- [14] D0 collaboration, *Search for $W' \rightarrow tb$ resonances with left- and right-handed couplings to fermions*, *Phys. Lett. B* **699** (2011) 145 [[arXiv:1101.0806](#)] [[INSPIRE](#)].
- [15] CDF collaboration, *Search for resonances decaying to top and bottom quarks with the CDF experiment*, *Phys. Rev. Lett.* **115** (2015) 061801 [[arXiv:1504.01536](#)] [[INSPIRE](#)].
- [16] CMS collaboration, *Search for a W' boson decaying to a bottom quark and a top quark in pp collisions at $\sqrt{s} = 7$ TeV*, *Phys. Lett. B* **718** (2013) 1229 [[arXiv:1208.0956](#)] [[INSPIRE](#)].
- [17] CMS collaboration, *Search for $W' \rightarrow tb$ decays in the lepton + jets final state in pp collisions at $\sqrt{s} = 8$ TeV*, *JHEP* **05** (2014) 108 [[arXiv:1402.2176](#)] [[INSPIRE](#)].
- [18] CMS collaboration, *Search for heavy resonances decaying to a top quark and a bottom quark in the lepton+jets final state in proton-proton collisions at 13 TeV*, *Phys. Lett. B* **777** (2018) 39 [[arXiv:1708.08539](#)] [[INSPIRE](#)].
- [19] ATLAS collaboration, *Search for $W' \rightarrow t\bar{b}$ in the lepton plus jets final state in proton-proton collisions at a centre-of-mass energy of $\sqrt{s} = 8$ TeV with the ATLAS detector*, *Phys. Lett. B* **743** (2015) 235 [[arXiv:1410.4103](#)] [[INSPIRE](#)].
- [20] ATLAS collaboration, *Search for $W' \rightarrow tb \rightarrow qqbb$ decays in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector*, *Eur. Phys. J. C* **75** (2015) 165 [[arXiv:1408.0886](#)] [[INSPIRE](#)].
- [21] ATLAS collaboration, *Search for $W' \rightarrow tb$ decays in the hadronic final state using pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Lett. B* **781** (2018) 327 [[arXiv:1801.07893](#)] [[INSPIRE](#)].
- [22] ATLAS collaboration, *Search for vector-boson resonances decaying to a top quark and bottom quark in the lepton plus jets final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Lett. B* **788** (2019) 347 [[arXiv:1807.10473](#)] [[INSPIRE](#)].

- [23] CMS collaboration, *Search for W' bosons decaying to a top and a bottom quark at $\sqrt{s} = 13$ TeV in the hadronic final state*, *Phys. Lett. B* **820** (2021) 136535 [[arXiv:2104.04831](#)] [[INSPIRE](#)].
- [24] ATLAS collaboration, *The ATLAS experiment at the CERN Large Hadron Collider*, 2008 *JINST* **3** S08003 [[INSPIRE](#)].
- [25] ATLAS collaboration, *ATLAS insertable B-layer: technical design report*, ATLAS-TDR-19, CERN, Geneva, Switzerland (2010).
- [26] ATLAS IBL collaboration, *Production and integration of the ATLAS insertable B-layer*, 2018 *JINST* **13** T05008 [[arXiv:1803.00844](#)] [[INSPIRE](#)].
- [27] ATLAS collaboration, *Performance of the ATLAS trigger system in 2015*, *Eur. Phys. J. C* **77** (2017) 317 [[arXiv:1611.09661](#)] [[INSPIRE](#)].
- [28] ATLAS collaboration, *The ATLAS collaboration software and firmware*, ATL-SOFT-PUB-2021-001, CERN, Geneva, Switzerland (2021).
- [29] ATLAS collaboration, *ATLAS data quality operations and performance for 2015–2018 data-taking*, 2020 *JINST* **15** P04003 [[arXiv:1911.04632](#)] [[INSPIRE](#)].
- [30] GEANT4 collaboration, *GEANT4 — a simulation toolkit*, *Nucl. Instrum. Meth. A* **506** (2003) 250 [[INSPIRE](#)].
- [31] ATLAS collaboration, *The ATLAS simulation infrastructure*, *Eur. Phys. J. C* **70** (2010) 823 [[arXiv:1005.4568](#)] [[INSPIRE](#)].
- [32] ATLAS collaboration, *Luminosity determination in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector at the LHC*, ATLAS-CONF-2019-021, CERN, Geneva, Switzerland (2019).
- [33] J. Alwall et al., *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, *JHEP* **07** (2014) 079 [[arXiv:1405.0301](#)] [[INSPIRE](#)].
- [34] T. Sjöstrand et al., *An introduction to PYTHIA 8.2*, *Comput. Phys. Commun.* **191** (2015) 159 [[arXiv:1410.3012](#)] [[INSPIRE](#)].
- [35] J. Butterworth et al., *PDF4LHC recommendations for LHC run II*, *J. Phys. G* **43** (2016) 023001 [[arXiv:1510.03865](#)] [[INSPIRE](#)].
- [36] ATLAS collaboration, *ATLAS Pythia 8 tunes to 7 TeV data*, ATL-PHYS-PUB-2014-021, CERN, Geneva, Switzerland (2014).
- [37] E. Boos, V. Bunichev, L. Dudko and M. Perfilov, *Interference between W' and W in single-top quark production processes*, *Phys. Lett. B* **655** (2007) 245 [[hep-ph/0610080](#)] [[INSPIRE](#)].
- [38] M. Beneke, P. Falgari, S. Klein and C. Schwinn, *Hadronic top-quark pair production with NNLL threshold resummation*, *Nucl. Phys. B* **855** (2012) 695 [[arXiv:1109.1536](#)] [[INSPIRE](#)].
- [39] M. Cacciari et al., *Top-pair production at hadron colliders with next-to-next-to-leading logarithmic soft-gluon resummation*, *Phys. Lett. B* **710** (2012) 612 [[arXiv:1111.5869](#)] [[INSPIRE](#)].
- [40] P. Bärnreuther, M. Czakon and A. Mitov, *Percent level precision physics at the Tevatron: first genuine NNLO QCD corrections to $q\bar{q} \rightarrow t\bar{t} + X$* , *Phys. Rev. Lett.* **109** (2012) 132001 [[arXiv:1204.5201](#)] [[INSPIRE](#)].

- [41] M. Czakon and A. Mitov, *NNLO corrections to top-pair production at hadron colliders: the all-fermionic scattering channels*, *JHEP* **12** (2012) 054 [[arXiv:1207.0236](#)] [[INSPIRE](#)].
- [42] M. Czakon and A. Mitov, *NNLO corrections to top pair production at hadron colliders: the quark-gluon reaction*, *JHEP* **01** (2013) 080 [[arXiv:1210.6832](#)] [[INSPIRE](#)].
- [43] M. Czakon, P. Fiedler and A. Mitov, *Total top-quark pair-production cross section at hadron colliders through $O(\alpha_S^4)$* , *Phys. Rev. Lett.* **110** (2013) 252004 [[arXiv:1303.6254](#)] [[INSPIRE](#)].
- [44] M. Czakon and A. Mitov, *Top++: a program for the calculation of the top-pair cross-section at hadron colliders*, *Comput. Phys. Commun.* **185** (2014) 2930 [[arXiv:1112.5675](#)] [[INSPIRE](#)].
- [45] S. Frixione, P. Nason and G. Ridolfi, *A positive-weight next-to-leading-order Monte Carlo for heavy flavour hadroproduction*, *JHEP* **09** (2007) 126 [[arXiv:0707.3088](#)] [[INSPIRE](#)].
- [46] P. Nason, *A new method for combining NLO QCD with shower Monte Carlo algorithms*, *JHEP* **11** (2004) 040 [[hep-ph/0409146](#)] [[INSPIRE](#)].
- [47] S. Frixione, P. Nason and C. Oleari, *Matching NLO QCD computations with parton shower simulations: the POWHEG method*, *JHEP* **11** (2007) 070 [[arXiv:0709.2092](#)] [[INSPIRE](#)].
- [48] S. Alioli, P. Nason, C. Oleari and E. Re, *A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX*, *JHEP* **06** (2010) 043 [[arXiv:1002.2581](#)] [[INSPIRE](#)].
- [49] NNPDF collaboration, *Parton distributions for the LHC run II*, *JHEP* **04** (2015) 040 [[arXiv:1410.8849](#)] [[INSPIRE](#)].
- [50] ATLAS collaboration, *Studies on top-quark Monte Carlo modelling for Top2016*, [ATL-PHYS-PUB-2016-020](#), CERN, Geneva, Switzerland (2016).
- [51] R.D. Ball et al., *Parton distributions with LHC data*, *Nucl. Phys. B* **867** (2013) 244 [[arXiv:1207.1303](#)] [[INSPIRE](#)].
- [52] D.J. Lange, *The EvtGen particle decay simulation package*, *Nucl. Instrum. Meth. A* **462** (2001) 152 [[INSPIRE](#)].
- [53] M. Czakon et al., *Top-pair production at the LHC through NNLO QCD and NLO EW*, *JHEP* **10** (2017) 186 [[arXiv:1705.04105](#)] [[INSPIRE](#)].
- [54] SHERPA collaboration, *Event generation with Sherpa 2.2*, *SciPost Phys.* **7** (2019) 034 [[arXiv:1905.09127](#)] [[INSPIRE](#)].
- [55] T. Gleisberg and S. Höche, *Comix, a new matrix element generator*, *JHEP* **12** (2008) 039 [[arXiv:0808.3674](#)] [[INSPIRE](#)].
- [56] F. Buccioni et al., *OpenLoops 2*, *Eur. Phys. J. C* **79** (2019) 866 [[arXiv:1907.13071](#)] [[INSPIRE](#)].
- [57] F. Cascioli, P. Maierhöfer and S. Pozzorini, *Scattering amplitudes with OpenLoops*, *Phys. Rev. Lett.* **108** (2012) 111601 [[arXiv:1111.5206](#)] [[INSPIRE](#)].
- [58] A. Denner, S. Dittmaier and L. Hofer, *Collier: a fortran-based Complex One-Loop Library in Extended Regularizations*, *Comput. Phys. Commun.* **212** (2017) 220 [[arXiv:1604.06792](#)] [[INSPIRE](#)].
- [59] S. Schumann and F. Krauss, *A parton shower algorithm based on Catani-Seymour dipole factorisation*, *JHEP* **03** (2008) 038 [[arXiv:0709.1027](#)] [[INSPIRE](#)].

- [60] S. Höche, F. Krauss, M. Schönherr and F. Siegert, *A critical appraisal of NLO+PS matching methods*, *JHEP* **09** (2012) 049 [[arXiv:1111.1220](#)] [[INSPIRE](#)].
- [61] S. Höche, F. Krauss, M. Schönherr and F. Siegert, *QCD matrix elements + parton showers: the NLO case*, *JHEP* **04** (2013) 027 [[arXiv:1207.5030](#)] [[INSPIRE](#)].
- [62] S. Catani, F. Krauss, R. Kuhn and B.R. Webber, *QCD matrix elements + parton showers*, *JHEP* **11** (2001) 063 [[hep-ph/0109231](#)] [[INSPIRE](#)].
- [63] S. Höche, F. Krauss, S. Schumann and F. Siegert, *QCD matrix elements and truncated showers*, *JHEP* **05** (2009) 053 [[arXiv:0903.1219](#)] [[INSPIRE](#)].
- [64] C. Anastasiou, L.J. Dixon, K. Melnikov and F. Petriello, *High precision QCD at hadron colliders: electroweak gauge boson rapidity distributions at NNLO*, *Phys. Rev. D* **69** (2004) 094008 [[hep-ph/0312266](#)] [[INSPIRE](#)].
- [65] R. Frederix, E. Re and P. Torrielli, *Single-top t -channel hadroproduction in the four-flavour scheme with POWHEG and aMC@NLO*, *JHEP* **09** (2012) 130 [[arXiv:1207.5391](#)] [[INSPIRE](#)].
- [66] S. Frixione et al., *Single-top hadroproduction in association with a W boson*, *JHEP* **07** (2008) 029 [[arXiv:0805.3067](#)] [[INSPIRE](#)].
- [67] ATLAS collaboration, *Topological cell clustering in the ATLAS calorimeters and its performance in LHC run 1*, *Eur. Phys. J. C* **77** (2017) 490 [[arXiv:1603.02934](#)] [[INSPIRE](#)].
- [68] M. Cacciari, G.P. Salam and G. Soyez, *The anti- k_t jet clustering algorithm*, *JHEP* **04** (2008) 063 [[arXiv:0802.1189](#)] [[INSPIRE](#)].
- [69] M. Cacciari, G.P. Salam and G. Soyez, *FastJet user manual*, *Eur. Phys. J. C* **72** (2012) 1896 [[arXiv:1111.6097](#)] [[INSPIRE](#)].
- [70] D. Krohn, J. Thaler and L.-T. Wang, *Jet trimming*, *JHEP* **02** (2010) 084 [[arXiv:0912.1342](#)] [[INSPIRE](#)].
- [71] S. Catani, Y.L. Dokshitzer, M.H. Seymour and B.R. Webber, *Longitudinally invariant K_t clustering algorithms for hadron hadron collisions*, *Nucl. Phys. B* **406** (1993) 187 [[INSPIRE](#)].
- [72] S.D. Ellis and D.E. Soper, *Successive combination jet algorithm for hadron collisions*, *Phys. Rev. D* **48** (1993) 3160 [[hep-ph/9305266](#)] [[INSPIRE](#)].
- [73] ATLAS collaboration, *In situ calibration of large-radius jet energy and mass in 13 TeV proton-proton collisions with the ATLAS detector*, *Eur. Phys. J. C* **79** (2019) 135 [[arXiv:1807.09477](#)] [[INSPIRE](#)].
- [74] ATLAS collaboration, *Performance of top-quark and W -boson tagging with ATLAS in run 2 of the LHC*, *Eur. Phys. J. C* **79** (2019) 375 [[arXiv:1808.07858](#)] [[INSPIRE](#)].
- [75] J. Thaler and K. Van Tilburg, *Identifying boosted objects with N -subjettiness*, *JHEP* **03** (2011) 015 [[arXiv:1011.2268](#)] [[INSPIRE](#)].
- [76] J. Thaler and L.-T. Wang, *Strategies to identify boosted tops*, *JHEP* **07** (2008) 092 [[arXiv:0806.0023](#)] [[INSPIRE](#)].
- [77] A.J. Larkoski, G.P. Salam and J. Thaler, *Energy correlation functions for jet substructure*, *JHEP* **06** (2013) 108 [[arXiv:1305.0007](#)] [[INSPIRE](#)].
- [78] ATLAS collaboration, *Boosted hadronic vector boson and top quark tagging with ATLAS using run 2 data*, [ATL-PHYS-PUB-2020-017](#), CERN, Geneva, Switzerland (2020).

- [79] M. Cacciari, G.P. Salam and G. Soyez, *The catchment area of jets*, *JHEP* **04** (2008) 005 [[arXiv:0802.1188](#)] [[INSPIRE](#)].
- [80] ATLAS collaboration, *Jet reconstruction and performance using particle flow with the ATLAS detector*, *Eur. Phys. J. C* **77** (2017) 466 [[arXiv:1703.10485](#)] [[INSPIRE](#)].
- [81] ATLAS collaboration, *Jet energy scale and resolution measured in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Eur. Phys. J. C* **81** (2021) 689 [[arXiv:2007.02645](#)] [[INSPIRE](#)].
- [82] ATLAS collaboration, *Performance of pile-up mitigation techniques for jets in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector*, *Eur. Phys. J. C* **76** (2016) 581 [[arXiv:1510.03823](#)] [[INSPIRE](#)].
- [83] ATLAS collaboration, *ATLAS b-jet identification performance and efficiency measurement with $t\bar{t}$ events in pp collisions at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **79** (2019) 970 [[arXiv:1907.05120](#)] [[INSPIRE](#)].
- [84] ATLAS collaboration, *ATLAS flavour-tagging algorithms for the LHC run 2 pp collision dataset*, *Eur. Phys. J. C* **83** (2023) 681 [[arXiv:2211.16345](#)] [[INSPIRE](#)].
- [85] ATLAS collaboration, *Measurement of the c-jet mistagging efficiency in $t\bar{t}$ events using pp collision data at $\sqrt{s} = 13$ TeV collected with the ATLAS detector*, *Eur. Phys. J. C* **82** (2022) 95 [[arXiv:2109.10627](#)] [[INSPIRE](#)].
- [86] ATLAS collaboration, *Calibration of the light-flavour jet mistagging efficiency of the b-tagging algorithms with Z+jets events using 139 fb^{-1} of ATLAS proton-proton collision data at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **83** (2023) 728 [[arXiv:2301.06319](#)] [[INSPIRE](#)].
- [87] B. Nachman et al., *Jets from jets: re-clustering as a tool for large radius jet reconstruction and grooming at the LHC*, *JHEP* **02** (2015) 075 [[arXiv:1407.2922](#)] [[INSPIRE](#)].
- [88] D. Krohn, J. Thaler and L.-T. Wang, *Jets with variable R*, *JHEP* **06** (2009) 059 [[arXiv:0903.0392](#)] [[INSPIRE](#)].
- [89] ATLAS collaboration, *Electron and photon performance measurements with the ATLAS detector using the 2015–2017 LHC proton-proton collision data*, *2019 JINST* **14** P12006 [[arXiv:1908.00005](#)] [[INSPIRE](#)].
- [90] ATLAS collaboration, *Muon reconstruction and identification efficiency in ATLAS using the full run 2 pp collision data set at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **81** (2021) 578 [[arXiv:2012.00578](#)] [[INSPIRE](#)].
- [91] ATLAS collaboration, *E_T^{miss} performance in the ATLAS detector using 2015–2016 LHC pp collisions*, *ATLAS-CONF-2018-023*, CERN, Geneva, Switzerland (2018).
- [92] ATLAS collaboration, *Electron reconstruction and identification in the ATLAS experiment using the 2015 and 2016 LHC proton-proton collision data at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **79** (2019) 639 [[arXiv:1902.04655](#)] [[INSPIRE](#)].
- [93] ATLAS collaboration, *Muon reconstruction performance of the ATLAS detector in proton-proton collision data at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **76** (2016) 292 [[arXiv:1603.05598](#)] [[INSPIRE](#)].
- [94] ATLAS collaboration, *Measurement of the ATLAS detector jet mass response using forward folding with 80 fb^{-1} of $\sqrt{s} = 13$ TeV pp data*, *ATLAS-CONF-2020-022*, CERN, Geneva, Switzerland (2020).

- [95] ATLAS collaboration, *Optimisation of large-radius jet reconstruction for the ATLAS detector in 13 TeV proton-proton collisions*, *Eur. Phys. J. C* **81** (2021) 334 [[arXiv:2009.04986](#)] [[INSPIRE](#)].
- [96] G. Avoni et al., *The new LUCID-2 detector for luminosity measurement and monitoring in ATLAS*, 2018 *JINST* **13** P07017 [[INSPIRE](#)].
- [97] M. Bähr et al., *Herwig++ physics and manual*, *Eur. Phys. J. C* **58** (2008) 639 [[arXiv:0803.0883](#)] [[INSPIRE](#)].
- [98] J. Bellm et al., *Herwig 7.0/Herwig++ 3.0 release note*, *Eur. Phys. J. C* **76** (2016) 196 [[arXiv:1512.01178](#)] [[INSPIRE](#)].
- [99] L.A. Harland-Lang, A.D. Martin, P. Motylinski and R.S. Thorne, *Parton distributions in the LHC era: MMHT 2014 PDFs*, *Eur. Phys. J. C* **75** (2015) 204 [[arXiv:1412.3989](#)] [[INSPIRE](#)].
- [100] A. Manohar, P. Nason, G.P. Salam and G. Zanderighi, *How bright is the proton? A precise determination of the photon parton distribution function*, *Phys. Rev. Lett.* **117** (2016) 242002 [[arXiv:1607.04266](#)] [[INSPIRE](#)].
- [101] S. Dulat et al., *New parton distribution functions from a global analysis of quantum chromodynamics*, *Phys. Rev. D* **93** (2016) 033006 [[arXiv:1506.07443](#)] [[INSPIRE](#)].
- [102] T.-J. Hou et al., *New CTEQ global analysis of quantum chromodynamics with high-precision data from the LHC*, *Phys. Rev. D* **103** (2021) 014013 [[arXiv:1912.10053](#)] [[INSPIRE](#)].
- [103] S. Bailey et al., *Parton distributions from LHC, HERA, Tevatron and fixed target data: MSHT20 PDFs*, *Eur. Phys. J. C* **81** (2021) 341 [[arXiv:2012.04684](#)] [[INSPIRE](#)].
- [104] S. Kallweit et al., *NLO electroweak automation and precise predictions for $W +$ multijet production at the LHC*, *JHEP* **04** (2015) 012 [[arXiv:1412.5157](#)] [[INSPIRE](#)].
- [105] J. Alwall et al., *Comparative study of various algorithms for the merging of parton showers and matrix elements in hadronic collisions*, *Eur. Phys. J. C* **53** (2008) 473 [[arXiv:0706.2569](#)] [[INSPIRE](#)].
- [106] ATLAS collaboration, *Search for the standard model Higgs boson produced in association with top quarks and decaying into a $b\bar{b}$ pair in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Phys. Rev. D* **97** (2018) 072016 [[arXiv:1712.08895](#)] [[INSPIRE](#)].
- [107] M. Baak et al., *HistFitter software framework for statistical data analysis*, *Eur. Phys. J. C* **75** (2015) 153 [[arXiv:1410.1280](#)] [[INSPIRE](#)].
- [108] G. Cowan, K. Cranmer, E. Gross and O. Vitells, *Asymptotic formulae for likelihood-based tests of new physics*, *Eur. Phys. J. C* **71** (2011) 1554 [*Erratum ibid.* **73** (2013) 2501] [[arXiv:1007.1727](#)] [[INSPIRE](#)].
- [109] A.L. Read, *Presentation of search results: the CL_s technique*, *J. Phys. G* **28** (2002) 2693 [[INSPIRE](#)].
- [110] ATLAS collaboration, *Search for heavy Higgs bosons A/H decaying to a top quark pair in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector*, *Phys. Rev. Lett.* **119** (2017) 191803 [[arXiv:1707.06025](#)] [[INSPIRE](#)].
- [111] ATLAS collaboration, *ATLAS computing acknowledgements*, ATLAS-SOFT-PUB-2023-001, CERN, Geneva, Switzerland (2023).

The ATLAS collaboration

G. Aad [ID](#)¹⁰², B. Abbott [ID](#)¹²⁰, K. Abeling [ID](#)⁵⁵, N.J. Abicht [ID](#)⁴⁹, S.H. Abidi [ID](#)²⁹,
A. Aboulhorma [ID](#)^{35e}, H. Abramowicz [ID](#)¹⁵¹, H. Abreu [ID](#)¹⁵⁰, Y. Abulaiti [ID](#)¹¹⁷,
A.C. Abusleme Hoffman [ID](#)^{137a}, B.S. Acharya [ID](#)^{69a,69b,n}, C. Adam Bourdarios [ID](#)⁴, L. Adamczyk [ID](#)^{86a},
L. Adamek [ID](#)¹⁵⁵, S.V. Addepalli [ID](#)²⁶, M.J. Addison [ID](#)¹⁰¹, J. Adelman [ID](#)¹¹⁵, A. Adiguzel [ID](#)^{21c},
T. Adye [ID](#)¹³⁴, A.A. Affolder [ID](#)¹³⁶, Y. Afik [ID](#)³⁶, M.N. Agaras [ID](#)¹³, J. Agarwala [ID](#)^{73a,73b},
A. Aggarwal [ID](#)¹⁰⁰, C. Agheorghiesei [ID](#)^{27c}, A. Ahmad [ID](#)³⁶, F. Ahmadov [ID](#)^{38,z}, W.S. Ahmed [ID](#)¹⁰⁴,
S. Ahuja [ID](#)⁹⁵, X. Ai [ID](#)^{62a}, G. Aielli [ID](#)^{76a,76b}, A. Aikot [ID](#)¹⁶³, M. Ait Tamlihat [ID](#)^{35e},
B. Aitbenchikh [ID](#)^{35a}, I. Aizenberg [ID](#)¹⁶⁹, M. Akbiyik [ID](#)¹⁰⁰, T.P.A. Åkesson [ID](#)⁹⁸, A.V. Akimov [ID](#)³⁷,
D. Akiyama [ID](#)¹⁶⁸, N.N. Akolkar [ID](#)²⁴, K. Al Khoury [ID](#)⁴¹, G.L. Alberghi [ID](#)^{23b}, J. Albert [ID](#)¹⁶⁵,
P. Albicocco [ID](#)⁵³, G.L. Albouy [ID](#)⁶⁰, S. Alderweireldt [ID](#)⁵², M. Aleksa [ID](#)³⁶, I.N. Aleksandrov [ID](#)³⁸,
C. Alexa [ID](#)^{27b}, T. Alexopoulos [ID](#)¹⁰, F. Alfonsi [ID](#)^{23b}, M. Algren [ID](#)⁵⁶, M. Alhroob [ID](#)¹²⁰, B. Ali [ID](#)¹³²,
H.M.J. Ali [ID](#)⁹¹, S. Ali [ID](#)¹⁴⁸, S.W. Alibocus [ID](#)⁹², M. Aliev [ID](#)¹⁴⁵, G. Alimonti [ID](#)^{71a}, W. Alkakhki [ID](#)⁵⁵,
C. Allaire [ID](#)⁶⁶, B.M.M. Allbrooke [ID](#)¹⁴⁶, J.F. Allen [ID](#)⁵², C.A. Allendes Flores [ID](#)^{137f}, P.P. Allport [ID](#)²⁰,
A. Aloisio [ID](#)^{72a,72b}, F. Alonso [ID](#)⁹⁰, C. Alpigiani [ID](#)¹³⁸, M. Alvarez Estevez [ID](#)⁹⁹,
A. Alvarez Fernandez [ID](#)¹⁰⁰, M. Alves Cardoso [ID](#)⁵⁶, M.G. Alviggi [ID](#)^{72a,72b}, M. Aly [ID](#)¹⁰¹,
Y. Amaral Coutinho [ID](#)^{83b}, A. Ambler [ID](#)¹⁰⁴, C. Amelung [ID](#)³⁶, M. Amerl [ID](#)¹⁰¹, C.G. Ames [ID](#)¹⁰⁹,
D. Amidei [ID](#)¹⁰⁶, S.P. Amor Dos Santos [ID](#)^{130a}, K.R. Amos [ID](#)¹⁶³, V. Ananiev [ID](#)¹²⁵,
C. Anastopoulos [ID](#)¹³⁹, T. Andeen [ID](#)¹¹, J.K. Anders [ID](#)³⁶, S.Y. Andrean [ID](#)^{47a,47b},
A. Andreatza [ID](#)^{71a,71b}, S. Angelidakis [ID](#)⁹, A. Angerami [ID](#)^{41,ac}, A.V. Anisenkov [ID](#)³⁷, A. Annovi [ID](#)^{74a},
C. Antel [ID](#)⁵⁶, M.T. Anthony [ID](#)¹³⁹, E. Antipov [ID](#)¹⁴⁵, M. Antonelli [ID](#)⁵³, F. Anulli [ID](#)^{75a}, M. Aoki [ID](#)⁸⁴,
T. Aoki [ID](#)¹⁵³, J.A. Aparisi Pozo [ID](#)¹⁶³, M.A. Aparo [ID](#)¹⁴⁶, L. Aperio Bella [ID](#)⁴⁸, C. Appelt [ID](#)¹⁸,
A. Apyan [ID](#)²⁶, N. Aranzabal [ID](#)³⁶, C. Arcangeletti [ID](#)⁵³, A.T.H. Arce [ID](#)⁵¹, E. Arena [ID](#)⁹²,
J-F. Arguin [ID](#)¹⁰⁸, S. Argyropoulos [ID](#)⁵⁴, J.-H. Arling [ID](#)⁴⁸, O. Arnaez [ID](#)⁴, H. Arnold [ID](#)¹¹⁴,
G. Artoni [ID](#)^{75a,75b}, H. Asada [ID](#)¹¹¹, K. Asai [ID](#)¹¹⁸, S. Asai [ID](#)¹⁵³, N.A. Asbah [ID](#)⁶¹, J. Assahsah [ID](#)^{35d},
K. Assamagan [ID](#)²⁹, R. Astalos [ID](#)^{28a}, S. Atashi [ID](#)¹⁶⁰, R.J. Atkin [ID](#)^{33a}, M. Atkinson [ID](#)¹⁶², H. Atmani [ID](#)^{35f},
P.A. Atmasiddha [ID](#)¹⁰⁶, K. Augsten [ID](#)¹³², S. Auricchio [ID](#)^{72a,72b}, A.D. Auriol [ID](#)²⁰, V.A. Austrup [ID](#)¹⁰¹,
G. Avolio [ID](#)³⁶, K. Axiotis [ID](#)⁵⁶, G. Azuelos [ID](#)^{108,ah}, D. Babal [ID](#)^{28b}, H. Bachacou [ID](#)¹³⁵,
K. Bachas [ID](#)^{152,q}, A. Bachiu [ID](#)³⁴, F. Backman [ID](#)^{47a,47b}, A. Badea [ID](#)⁶¹, P. Bagnaia [ID](#)^{75a,75b},
M. Bahmani [ID](#)¹⁸, A.J. Bailey [ID](#)¹⁶³, V.R. Bailey [ID](#)¹⁶², J.T. Baines [ID](#)¹³⁴, L. Baines [ID](#)⁹⁴,
C. Bakalis [ID](#)¹⁰, O.K. Baker [ID](#)¹⁷², E. Bakos [ID](#)¹⁵, D. Bakshi Gupta [ID](#)⁸, V. Balakrishnan [ID](#)¹²⁰,
R. Balasubramanian [ID](#)¹¹⁴, E.M. Baldin [ID](#)³⁷, P. Balek [ID](#)^{86a}, E. Ballabene [ID](#)^{23b,23a}, F. Balli [ID](#)¹³⁵,
L.M. Baltes [ID](#)^{63a}, W.K. Balunas [ID](#)³², J. Balz [ID](#)¹⁰⁰, E. Banas [ID](#)⁸⁷, M. Bandieramonte [ID](#)¹²⁹,
A. Bandyopadhyay [ID](#)²⁴, S. Bansal [ID](#)²⁴, L. Barak [ID](#)¹⁵¹, M. Barakat [ID](#)⁴⁸, E.L. Barberio [ID](#)¹⁰⁵,
D. Barberis [ID](#)^{57b,57a}, M. Barbero [ID](#)¹⁰², M.Z. Barel [ID](#)¹¹⁴, K.N. Barends [ID](#)^{33a}, T. Barillari [ID](#)¹¹⁰,
M-S. Barisits [ID](#)³⁶, T. Barklow [ID](#)¹⁴³, P. Baron [ID](#)¹²², D.A. Baron Moreno [ID](#)¹⁰¹, A. Baroncelli [ID](#)^{62a},
G. Barone [ID](#)²⁹, A.J. Barr [ID](#)¹²⁶, J.D. Barr [ID](#)⁹⁶, L. Barranco Navarro [ID](#)^{47a,47b}, F. Barreiro [ID](#)⁹⁹,
J. Barreiro Guimarães da Costa [ID](#)^{14a}, U. Barron [ID](#)¹⁵¹, M.G. Barros Teixeira [ID](#)^{130a}, S. Barsov [ID](#)³⁷,
F. Bartels [ID](#)^{63a}, R. Bartoldus [ID](#)¹⁴³, A.E. Barton [ID](#)⁹¹, P. Bartos [ID](#)^{28a}, A. Basan [ID](#)¹⁰⁰,
M. Baselga [ID](#)⁴⁹, A. Bassalat [ID](#)^{66,b}, M.J. Basso [ID](#)^{156a}, C.R. Basson [ID](#)¹⁰¹, R.L. Bates [ID](#)⁵⁹,
S. Batlamous [ID](#)^{35e}, J.R. Batley [ID](#)³², B. Batool [ID](#)¹⁴¹, M. Battaglia [ID](#)¹³⁶, D. Battulga [ID](#)¹⁸,
M. Bauge [ID](#)^{75a,75b}, M. Bauer [ID](#)³⁶, P. Bauer [ID](#)²⁴, L.T. Bazzano Hurrell [ID](#)³⁰, J.B. Beacham [ID](#)⁵¹,

T. Beau [ID](#)¹²⁷, P.H. Beauchemin [ID](#)¹⁵⁸, F. Becherer [ID](#)⁵⁴, P. Bechtle [ID](#)²⁴, H.P. Beck [ID](#)^{19,p},
 K. Becker [ID](#)¹⁶⁷, A.J. Beddall [ID](#)⁸², V.A. Bednyakov [ID](#)³⁸, C.P. Bee [ID](#)¹⁴⁵, L.J. Beemster¹⁵,
 T.A. Beermann [ID](#)³⁶, M. Begalli [ID](#)^{83d}, M. Begel [ID](#)²⁹, A. Behera [ID](#)¹⁴⁵, J.K. Behr [ID](#)⁴⁸, J.F. Beirer [ID](#)⁵⁵,
 F. Beisiegel [ID](#)²⁴, M. Belfkir [ID](#)¹⁵⁹, G. Bella [ID](#)¹⁵¹, L. Bellagamba [ID](#)^{23b}, A. Bellerive [ID](#)³⁴, P. Bellos [ID](#)²⁰,
 K. Beloborodov [ID](#)³⁷, N.L. Belyaev [ID](#)³⁷, D. Benckekroun [ID](#)^{35a}, F. Bendebba [ID](#)^{35a},
 Y. Benhammou [ID](#)¹⁵¹, M. Benoit [ID](#)²⁹, J.R. Bensinger [ID](#)²⁶, S. Bentvelsen [ID](#)¹¹⁴, L. Beresford [ID](#)⁴⁸,
 M. Beretta [ID](#)⁵³, E. Bergeas Kuutmann [ID](#)¹⁶¹, N. Berger [ID](#)⁴, B. Bergmann [ID](#)¹³², J. Beringer [ID](#)^{17a},
 G. Bernardi [ID](#)⁵, C. Bernius [ID](#)¹⁴³, F.U. Bernlochner [ID](#)²⁴, F. Bernon [ID](#)^{36,102}, T. Berry [ID](#)⁹⁵,
 P. Berta [ID](#)¹³³, A. Berthold [ID](#)⁵⁰, I.A. Bertram [ID](#)⁹¹, S. Bethke [ID](#)¹¹⁰, A. Betti [ID](#)^{75a,75b}, A.J. Bevan [ID](#)⁹⁴,
 M. Bhamjee [ID](#)^{33c}, S. Bhatta [ID](#)¹⁴⁵, D.S. Bhattacharya [ID](#)¹⁶⁶, P. Bhattarai [ID](#)¹⁴³, V.S. Bhopatkar [ID](#)¹²¹,
 R. Bi^{29,aj}, R.M. Bianchi [ID](#)¹²⁹, G. Bianco [ID](#)^{23b,23a}, O. Biebel [ID](#)¹⁰⁹, R. Bielski [ID](#)¹²³, M. Biglietti [ID](#)^{77a},
 T.R.V. Billoud [ID](#)¹³², M. Bindi [ID](#)⁵⁵, A. Bingul [ID](#)^{21b}, C. Bini [ID](#)^{75a,75b}, A. Biondini [ID](#)⁹²,
 C.J. Birch-sykes [ID](#)¹⁰¹, G.A. Bird [ID](#)^{20,134}, M. Birman [ID](#)¹⁶⁹, M. Biros [ID](#)¹³³, S. Biryukov [ID](#)¹⁴⁶,
 T. Bisanz [ID](#)⁴⁹, E. Bisceglie [ID](#)^{43b,43a}, J.P. Biswal [ID](#)¹³⁴, D. Biswas [ID](#)¹⁴¹, A. Bitadze [ID](#)¹⁰¹,
 K. Bjørke [ID](#)¹²⁵, I. Bloch [ID](#)⁴⁸, C. Blocker [ID](#)²⁶, A. Blue [ID](#)⁵⁹, U. Blumenschein [ID](#)⁹⁴,
 J. Blumenthal [ID](#)¹⁰⁰, G.J. Bobbink [ID](#)¹¹⁴, V.S. Bobrovnikov [ID](#)³⁷, M. Boehler [ID](#)⁵⁴, B. Boehm [ID](#)¹⁶⁶,
 D. Bogavac [ID](#)³⁶, A.G. Bogdanchikov [ID](#)³⁷, C. Bohm [ID](#)^{47a}, V. Boisvert [ID](#)⁹⁵, P. Bokan [ID](#)⁴⁸,
 T. Bold [ID](#)^{86a}, M. Bomben [ID](#)⁵, M. Bona [ID](#)⁹⁴, M. Boonekamp [ID](#)¹³⁵, C.D. Booth [ID](#)⁹⁵,
 A.G. Borbély [ID](#)⁵⁹, I.S. Bordulev [ID](#)³⁷, H.M. Borecka-Bielska [ID](#)¹⁰⁸, L.S. Borgna [ID](#)⁹⁶, G. Borissov [ID](#)⁹¹,
 D. Bortoletto [ID](#)¹²⁶, D. Boscherini [ID](#)^{23b}, M. Bosman [ID](#)¹³, J.D. Bossio Sola [ID](#)³⁶, K. Bouaouda [ID](#)^{35a},
 N. Bouchhar [ID](#)¹⁶³, J. Boudreau [ID](#)¹²⁹, E.V. Bouhova-Thacker [ID](#)⁹¹, D. Boumediene [ID](#)⁴⁰,
 R. Bouquet [ID](#)⁵, A. Boveia [ID](#)¹¹⁹, J. Boyd [ID](#)³⁶, D. Boye [ID](#)²⁹, I.R. Boyko [ID](#)³⁸, J. Bracinik [ID](#)²⁰,
 N. Brahimi [ID](#)^{62d}, G. Brandt [ID](#)¹⁷¹, O. Brandt [ID](#)³², F. Braren [ID](#)⁴⁸, B. Brau [ID](#)¹⁰³, J.E. Brau [ID](#)¹²³,
 R. Brenner [ID](#)¹⁶⁹, L. Brenner [ID](#)¹¹⁴, R. Brenner [ID](#)¹⁶¹, S. Bressler [ID](#)¹⁶⁹, D. Britton [ID](#)⁵⁹,
 D. Britzger [ID](#)¹¹⁰, I. Brock [ID](#)²⁴, G. Brooijmans [ID](#)⁴¹, W.K. Brooks [ID](#)^{137f}, E. Brost [ID](#)²⁹,
 L.M. Brown [ID](#)¹⁶⁵, L.E. Bruce [ID](#)⁶¹, T.L. Bruckler [ID](#)¹²⁶, P.A. Bruckman de Renstrom [ID](#)⁸⁷,
 B. Brüers [ID](#)⁴⁸, A. Bruni [ID](#)^{23b}, G. Bruni [ID](#)^{23b}, M. Bruschi [ID](#)^{23b}, N. Bruscinò [ID](#)^{75a,75b}, T. Buanes [ID](#)¹⁶,
 Q. Buat [ID](#)¹³⁸, D. Buchin [ID](#)¹¹⁰, A.G. Buckley [ID](#)⁵⁹, O. Bulekov [ID](#)³⁷, B.A. Bullard [ID](#)¹⁴³, S. Burdin [ID](#)⁹²,
 C.D. Burgard [ID](#)⁴⁹, A.M. Burger [ID](#)⁴⁰, B. Burghgrave [ID](#)⁸, O. Burlayenko [ID](#)⁵⁴, J.T.P. Burr [ID](#)³²,
 C.D. Burton [ID](#)¹¹, J.C. Burzynski [ID](#)¹⁴², E.L. Busch [ID](#)⁴¹, V. Büscher [ID](#)¹⁰⁰, P.J. Bussey [ID](#)⁵⁹,
 J.M. Butler [ID](#)²⁵, C.M. Buttar [ID](#)⁵⁹, J.M. Butterworth [ID](#)⁹⁶, W. Buttinger [ID](#)¹³⁴,
 C.J. Buxo Vazquez [ID](#)¹⁰⁷, A.R. Buzykaev [ID](#)³⁷, S. Cabrera Urbán [ID](#)¹⁶³, L. Cadamuro [ID](#)⁶⁶,
 D. Caforio [ID](#)⁵⁸, H. Cai [ID](#)¹²⁹, Y. Cai [ID](#)^{14a,14e}, V.M.M. Cairo [ID](#)³⁶, O. Cakir [ID](#)^{3a}, N. Calace [ID](#)³⁶,
 P. Calafiura [ID](#)^{17a}, G. Calderini [ID](#)¹²⁷, P. Calfayan [ID](#)⁶⁸, G. Callea [ID](#)⁵⁹, L.P. Caloba ^{83b}, D. Calvet [ID](#)⁴⁰,
 S. Calvet [ID](#)⁴⁰, T.P. Calvet [ID](#)¹⁰², M. Calvetti [ID](#)^{74a,74b}, R. Camacho Toro [ID](#)¹²⁷, S. Camarda [ID](#)³⁶,
 D. Camarero Munoz [ID](#)²⁶, P. Camarri [ID](#)^{76a,76b}, M.T. Camerlingo [ID](#)^{72a,72b}, D. Cameron [ID](#)³⁶,
 C. Camincher [ID](#)¹⁶⁵, M. Campanelli [ID](#)⁹⁶, A. Camplani [ID](#)⁴², V. Canale [ID](#)^{72a,72b}, A. Canesse [ID](#)¹⁰⁴,
 J. Cantero [ID](#)¹⁶³, Y. Cao [ID](#)¹⁶², F. Capocasa [ID](#)²⁶, M. Capua [ID](#)^{43b,43a}, A. Carbone [ID](#)^{71a,71b},
 R. Cardarelli [ID](#)^{76a}, J.C.J. Cardenas [ID](#)⁸, F. Cardillo [ID](#)¹⁶³, T. Carli [ID](#)³⁶, G. Carlino [ID](#)^{72a},
 J.I. Carlotto [ID](#)¹³, B.T. Carlson [ID](#)^{129,r}, E.M. Carlson [ID](#)^{165,156a}, L. Carminati [ID](#)^{71a,71b},
 A. Carnelli [ID](#)¹³⁵, M. Carnesale [ID](#)^{75a,75b}, S. Caron [ID](#)¹¹³, E. Carquin [ID](#)^{137f}, S. Carrá [ID](#)^{71a,71b},
 G. Carratta [ID](#)^{23b,23a}, F. Carrio Argos [ID](#)^{33g}, J.W.S. Carter [ID](#)¹⁵⁵, T.M. Carter [ID](#)⁵²,
 M.P. Casado [ID](#)^{13,i}, M. Caspar [ID](#)⁴⁸, E.G. Castiglia [ID](#)¹⁷², F.L. Castillo [ID](#)⁴, L. Castillo Garcia [ID](#)¹³,

V. Castillo Gimenez [ID](#)¹⁶³, N.F. Castro [ID](#)^{130a,130e}, A. Catinaccio [ID](#)³⁶, J.R. Catmore [ID](#)¹²⁵,
V. Cavaliere [ID](#)²⁹, N. Cavalli [ID](#)^{23b,23a}, V. Cavasinni [ID](#)^{74a,74b}, Y.C. Cekmecelioglu [ID](#)⁴⁸, E. Celebi [ID](#)^{21a},
F. Celli [ID](#)¹²⁶, M.S. Centonze [ID](#)^{70a,70b}, V. Cepaitis [ID](#)⁵⁶, K. Cerny [ID](#)¹²², A.S. Cerqueira [ID](#)^{83a},
A. Cerri [ID](#)¹⁴⁶, L. Cerrito [ID](#)^{76a,76b}, F. Cerutti [ID](#)^{17a}, B. Cervato [ID](#)¹⁴¹, A. Cervelli [ID](#)^{23b},
G. Cesarini [ID](#)⁵³, S.A. Cetin [ID](#)⁸², Z. Chadi [ID](#)^{35a}, D. Chakraborty [ID](#)¹¹⁵, J. Chan [ID](#)¹⁷⁰,
W.Y. Chan [ID](#)¹⁵³, J.D. Chapman [ID](#)³², E. Chapon [ID](#)¹³⁵, B. Chargeishvili [ID](#)^{149b}, D.G. Charlton [ID](#)²⁰,
T.P. Charman [ID](#)⁹⁴, M. Chatterjee [ID](#)¹⁹, C. Chauhan [ID](#)¹³³, S. Chekanov [ID](#)⁶, S.V. Chekulaev [ID](#)^{156a},
G.A. Chelkov [ID](#)^{38,a}, A. Chen [ID](#)¹⁰⁶, B. Chen [ID](#)¹⁵¹, B. Chen [ID](#)¹⁶⁵, H. Chen [ID](#)^{14c}, H. Chen [ID](#)²⁹,
J. Chen [ID](#)^{62c}, J. Chen [ID](#)¹⁴², M. Chen [ID](#)¹²⁶, S. Chen [ID](#)¹⁵³, S.J. Chen [ID](#)^{14c}, X. Chen [ID](#)^{62c,135},
X. Chen [ID](#)^{14b,ag}, Y. Chen [ID](#)^{62a}, C.L. Cheng [ID](#)¹⁷⁰, H.C. Cheng [ID](#)^{64a}, S. Cheong [ID](#)¹⁴³,
A. Cheplakov [ID](#)³⁸, E. Cheremushkina [ID](#)⁴⁸, E. Cherepanova [ID](#)¹¹⁴, R. Cherkaoui El Moursli [ID](#)^{35e},
E. Cheu [ID](#)⁷, K. Cheung [ID](#)⁶⁵, L. Chevalier [ID](#)¹³⁵, V. Chiarella [ID](#)⁵³, G. Chiarelli [ID](#)^{74a}, N. Chiedde [ID](#)¹⁰²,
G. Chiodini [ID](#)^{70a}, A.S. Chisholm [ID](#)²⁰, A. Chitan [ID](#)^{27b}, M. Chitishvili [ID](#)¹⁶³, M.V. Chizhov [ID](#)³⁸,
K. Choi [ID](#)¹¹, A.R. Chomont [ID](#)^{75a,75b}, Y. Chou [ID](#)¹⁰³, E.Y.S. Chow [ID](#)¹¹⁴, T. Chowdhury [ID](#)^{33g},
K.L. Chu [ID](#)¹⁶⁹, M.C. Chu [ID](#)^{64a}, X. Chu [ID](#)^{14a,14e}, J. Chudoba [ID](#)¹³¹, J.J. Chwastowski [ID](#)⁸⁷,
D. Cieri [ID](#)¹¹⁰, K.M. Ciesla [ID](#)^{86a}, V. Cindro [ID](#)⁹³, A. Ciochio [ID](#)^{17a}, F. Cirotto [ID](#)^{72a,72b},
Z.H. Citron [ID](#)^{169,l}, M. Citterio [ID](#)^{71a}, D.A. Ciubotaru [ID](#)^{27b}, B.M. Ciungu [ID](#)¹⁵⁵, A. Clark [ID](#)⁵⁶,
P.J. Clark [ID](#)⁵², J.M. Clavijo Columbie [ID](#)⁴⁸, S.E. Clawson [ID](#)⁴⁸, C. Clement [ID](#)^{47a,47b}, J. Clercx [ID](#)⁴⁸,
L. Clissa [ID](#)^{23b,23a}, Y. Coadou [ID](#)¹⁰², M. Cobal [ID](#)^{69a,69c}, A. Coccaro [ID](#)^{57b}, R.F. Coelho Barrue [ID](#)^{130a},
R. Coelho Lopes De Sa [ID](#)¹⁰³, S. Coelli [ID](#)^{71a}, H. Cohen [ID](#)¹⁵¹, A.E.C. Coimbra [ID](#)^{71a,71b}, B. Cole [ID](#)⁴¹,
J. Collot [ID](#)⁶⁰, P. Conde Muiño [ID](#)^{130a,130g}, M.P. Connell [ID](#)^{33c}, S.H. Connell [ID](#)^{33c}, I.A. Connelly [ID](#)⁵⁹,
E.I. Conroy [ID](#)¹²⁶, F. Conventi [ID](#)^{72a,ai}, H.G. Cooke [ID](#)²⁰, A.M. Cooper-Sarkar [ID](#)¹²⁶,
A. Cordeiro Oudot Choi [ID](#)¹²⁷, F. Cormier [ID](#)¹⁶⁴, L.D. Corpe [ID](#)⁴⁰, M. Corradi [ID](#)^{75a,75b},
F. Corriveau [ID](#)^{104,x}, A. Cortes-Gonzalez [ID](#)¹⁸, M.J. Costa [ID](#)¹⁶³, F. Costanza [ID](#)⁴, D. Costanzo [ID](#)¹³⁹,
B.M. Cote [ID](#)¹¹⁹, G. Cowan [ID](#)⁹⁵, K. Cranmer [ID](#)¹⁷⁰, D. Cremonini [ID](#)^{23b,23a}, S. Crépe-Renaudin [ID](#)⁶⁰,
F. Crescioli [ID](#)¹²⁷, M. Cristinziani [ID](#)¹⁴¹, M. Cristoforetti [ID](#)^{78a,78b}, V. Croft [ID](#)¹¹⁴, J.E. Crosby [ID](#)¹²¹,
G. Crosetti [ID](#)^{43b,43a}, A. Cueto [ID](#)⁹⁹, T. Cuhadar Donszelmann [ID](#)¹⁶⁰, H. Cui [ID](#)^{14a,14e}, Z. Cui [ID](#)⁷,
W.R. Cunningham [ID](#)⁵⁹, F. Curcio [ID](#)^{43b,43a}, P. Czodrowski [ID](#)³⁶, M.M. Czurylo [ID](#)^{63b},
M.J. Da Cunha Sargeddas De Sousa [ID](#)^{57b,57a}, J.V. Da Fonseca Pinto [ID](#)^{83b}, C. Da Via [ID](#)¹⁰¹,
W. Dabrowski [ID](#)^{86a}, T. Dado [ID](#)⁴⁹, S. Dahbi [ID](#)^{33g}, T. Dai [ID](#)¹⁰⁶, D. Dal Santo [ID](#)¹⁹,
C. Dallapiccola [ID](#)¹⁰³, M. Dam [ID](#)⁴², G. D’amen [ID](#)²⁹, V. D’Amico [ID](#)¹⁰⁹, J. Damp [ID](#)¹⁰⁰,
J.R. Dandoy [ID](#)¹²⁸, M.F. Daneri [ID](#)³⁰, M. Danninger [ID](#)¹⁴², V. Dao [ID](#)³⁶, G. Darbo [ID](#)^{57b},
S. Darmora [ID](#)⁶, S.J. Das [ID](#)^{29,aj}, S. D’Auria [ID](#)^{71a,71b}, C. David [ID](#)^{156b}, T. Davidek [ID](#)¹³³,
B. Davis-Purcell [ID](#)³⁴, I. Dawson [ID](#)⁹⁴, H.A. Day-hall [ID](#)¹³², K. De [ID](#)⁸, R. De Asmundis [ID](#)^{72a},
N. De Biase [ID](#)⁴⁸, S. De Castro [ID](#)^{23b,23a}, N. De Groot [ID](#)¹¹³, P. de Jong [ID](#)¹¹⁴, H. De la Torre [ID](#)¹¹⁵,
A. De Maria [ID](#)^{14c}, A. De Salvo [ID](#)^{75a}, U. De Sanctis [ID](#)^{76a,76b}, A. De Santo [ID](#)¹⁴⁶,
J.B. De Vivie De Regie [ID](#)⁶⁰, D.V. Dedovich [ID](#)³⁸, J. Degen [ID](#)¹¹⁴, A.M. Deiana [ID](#)⁴⁴,
F. Del Corso [ID](#)^{23b,23a}, J. Del Peso [ID](#)⁹⁹, F. Del Rio [ID](#)^{63a}, F. Deliot [ID](#)¹³⁵, C.M. Delitzsch [ID](#)⁴⁹,
M. Della Pietra [ID](#)^{72a,72b}, D. Della Volpe [ID](#)⁵⁶, A. Dell’Acqua [ID](#)³⁶, L. Dell’Asta [ID](#)^{71a,71b},
M. Delmastro [ID](#)⁴, P.A. Delsart [ID](#)⁶⁰, S. Demers [ID](#)¹⁷², M. Demichev [ID](#)³⁸, S.P. Denisov [ID](#)³⁷,
L. D’Eramo [ID](#)⁴⁰, D. Derendarz [ID](#)⁸⁷, F. Derue [ID](#)¹²⁷, P. Dervan [ID](#)⁹², K. Desch [ID](#)²⁴, C. Deutsch [ID](#)²⁴,
F.A. Di Bello [ID](#)^{57b,57a}, A. Di Ciaccio [ID](#)^{76a,76b}, L. Di Ciaccio [ID](#)⁴, A. Di Domenico [ID](#)^{75a,75b},
C. Di Donato [ID](#)^{72a,72b}, A. Di Girolamo [ID](#)³⁶, G. Di Gregorio [ID](#)⁵, A. Di Luca [ID](#)^{78a,78b},

B. Di Micco [ID](#)^{77a,77b}, R. Di Nardo [ID](#)^{77a,77b}, C. Diaconu [ID](#)¹⁰², M. Diamantopoulou [ID](#)³⁴,
 F.A. Dias [ID](#)¹¹⁴, T. Dias Do Vale [ID](#)¹⁴², M.A. Diaz [ID](#)^{137a,137b}, F.G. Diaz Capriles [ID](#)²⁴,
 M. Didenko [ID](#)¹⁶³, E.B. Diehl [ID](#)¹⁰⁶, L. Diehl [ID](#)⁵⁴, S. Díez Cornell [ID](#)⁴⁸, C. Díez Pardos [ID](#)¹⁴¹,
 C. Dimitriadi [ID](#)^{161,24,161}, A. Dimitrievska [ID](#)^{17a}, J. Dingfelder [ID](#)²⁴, I-M. Dimu [ID](#)^{27b},
 S.J. Dittmeier [ID](#)^{63b}, F. Dittus [ID](#)³⁶, F. Djama [ID](#)¹⁰², T. Djobava [ID](#)^{149b}, J.I. Djuvsland [ID](#)¹⁶,
 C. Doglioni [ID](#)^{101,98}, A. Dohnalova [ID](#)^{28a}, J. Dolejsi [ID](#)¹³³, Z. Dolezal [ID](#)¹³³, K.M. Dona [ID](#)³⁹,
 M. Donadelli [ID](#)^{83c}, B. Dong [ID](#)¹⁰⁷, J. Donini [ID](#)⁴⁰, A. D’Onofrio [ID](#)^{77a,77b}, M. D’Onofrio [ID](#)⁹²,
 J. Dopke [ID](#)¹³⁴, A. Doria [ID](#)^{72a}, N. Dos Santos Fernandes [ID](#)^{130a}, P. Dougan [ID](#)¹⁰¹, M.T. Dova [ID](#)⁹⁰,
 A.T. Doyle [ID](#)⁵⁹, M.A. Draguet [ID](#)¹²⁶, E. Dreyer [ID](#)¹⁶⁹, I. Drivas-koulouris [ID](#)¹⁰, A.S. Drobac [ID](#)¹⁵⁸,
 M. Drozdova [ID](#)⁵⁶, D. Du [ID](#)^{62a}, T.A. du Pree [ID](#)¹¹⁴, F. Dubinin [ID](#)³⁷, M. Dubovsky [ID](#)^{28a},
 E. Duchovni [ID](#)¹⁶⁹, G. Duckeck [ID](#)¹⁰⁹, O.A. Ducu [ID](#)^{27b}, D. Duda [ID](#)⁵², A. Dudarev [ID](#)³⁶,
 E.R. Duden [ID](#)²⁶, M. D’uffizi [ID](#)¹⁰¹, L. DufLOT [ID](#)⁶⁶, M. Dührssen [ID](#)³⁶, C. Dülsen [ID](#)¹⁷¹,
 A.E. Dumitriu [ID](#)^{27b}, M. Dunford [ID](#)^{63a}, S. Dungs [ID](#)⁴⁹, K. Dunne [ID](#)^{47a,47b}, A. Duperrin [ID](#)¹⁰²,
 H. Duran Yildiz [ID](#)^{3a}, M. Düren [ID](#)⁵⁸, A. Durglishvili [ID](#)^{149b}, B.L. Dwyer [ID](#)¹¹⁵, G.I. Dyckes [ID](#)^{17a},
 M. Dyndal [ID](#)^{86a}, S. Dysch [ID](#)¹⁰¹, B.S. Dziedzic [ID](#)⁸⁷, Z.O. Earnshaw [ID](#)¹⁴⁶, G.H. Eberwein [ID](#)¹²⁶,
 B. Eckerova [ID](#)^{28a}, S. Eggebrecht [ID](#)⁵⁵, E. Egidio Purcino De Souza [ID](#)¹²⁷, L.F. Ehrke [ID](#)⁵⁶,
 G. Eigen [ID](#)¹⁶, K. Einsweiler [ID](#)^{17a}, T. Ekelof [ID](#)¹⁶¹, P.A. Ekman [ID](#)⁹⁸, S. El Farkh [ID](#)^{35b},
 Y. El Ghazali [ID](#)^{35b}, H. El Jarrari [ID](#)^{35e,148}, A. El Moussaouy [ID](#)^{35a}, V. Ellajosyula [ID](#)¹⁶¹,
 M. Ellert [ID](#)¹⁶¹, F. Ellinghaus [ID](#)¹⁷¹, A.A. Elliot [ID](#)⁹⁴, N. Ellis [ID](#)³⁶, J. Elmsheuser [ID](#)²⁹, M. Elsing [ID](#)³⁶,
 D. Emelianov [ID](#)¹³⁴, Y. Enari [ID](#)¹⁵³, I. Ene [ID](#)^{17a}, S. Epari [ID](#)¹³, J. Erdmann [ID](#)⁴⁹, P.A. Erland [ID](#)⁸⁷,
 M. Errenst [ID](#)¹⁷¹, M. Escalier [ID](#)⁶⁶, C. Escobar [ID](#)¹⁶³, E. Etzion [ID](#)¹⁵¹, G. Evans [ID](#)^{130a}, H. Evans [ID](#)⁶⁸,
 L.S. Evans [ID](#)⁹⁵, M.O. Evans [ID](#)¹⁴⁶, A. Ezhilov [ID](#)³⁷, S. Ezzarqtouni [ID](#)^{35a}, F. Fabbri [ID](#)⁵⁹,
 L. Fabbri [ID](#)^{23b,23a}, G. Facini [ID](#)⁹⁶, V. Fadeyev [ID](#)¹³⁶, R.M. Fakhrutdinov [ID](#)³⁷, S. Falciano [ID](#)^{75a},
 L.F. Falda Ulhoa Coelho [ID](#)³⁶, P.J. Falke [ID](#)²⁴, J. Faltova [ID](#)¹³³, C. Fan [ID](#)¹⁶², Y. Fan [ID](#)^{14a},
 Y. Fang [ID](#)^{14a,14e}, M. Fanti [ID](#)^{71a,71b}, M. Faraj [ID](#)^{69a,69b}, Z. Farazpay [ID](#)⁹⁷, A. Farbin [ID](#)⁸,
 A. Farilla [ID](#)^{77a}, T. Farooque [ID](#)¹⁰⁷, S.M. Farrington [ID](#)⁵², F. Fassi [ID](#)^{35e}, D. Fassouliotis [ID](#)⁹,
 M. Fauci Giannelli [ID](#)^{76a,76b}, W.J. Fawcett [ID](#)³², L. Fayard [ID](#)⁶⁶, P. Federic [ID](#)¹³³, P. Federicova [ID](#)¹³¹,
 O.L. Fedin [ID](#)^{37,a}, G. Fedotov [ID](#)³⁷, M. Feickert [ID](#)¹⁷⁰, L. Feligioni [ID](#)¹⁰², D.E. Fellers [ID](#)¹²³,
 C. Feng [ID](#)^{62b}, M. Feng [ID](#)^{14b}, Z. Feng [ID](#)¹¹⁴, M.J. Fenton [ID](#)¹⁶⁰, A.B. Fenyuk [ID](#)³⁷, L. Ferencz [ID](#)⁴⁸,
 R.A.M. Ferguson [ID](#)⁹¹, S.I. Fernandez Luengo [ID](#)^{137f}, M.J.V. Fernoux [ID](#)¹⁰², J. Ferrando [ID](#)⁴⁸,
 A. Ferrari [ID](#)¹⁶¹, P. Ferrari [ID](#)^{114,113}, R. Ferrari [ID](#)^{73a}, D. Ferrere [ID](#)⁵⁶, C. Ferretti [ID](#)¹⁰⁶, F. Fiedler [ID](#)¹⁰⁰,
 A. Filipčić [ID](#)⁹³, E.K. Filmer [ID](#)¹, F. Filthaut [ID](#)¹¹³, M.C.N. Fiolhais [ID](#)^{130a,130c,c}, L. Fiorini [ID](#)¹⁶³,
 W.C. Fisher [ID](#)¹⁰⁷, T. Fitschen [ID](#)¹⁰¹, P.M. Fitzhugh [ID](#)¹³⁵, I. Fleck [ID](#)¹⁴¹, P. Fleischmann [ID](#)¹⁰⁶,
 T. Flick [ID](#)¹⁷¹, M. Flores [ID](#)^{33d,ad}, L.R. Flores Castillo [ID](#)^{64a}, L. Flores Sanz De Acedo [ID](#)³⁶,
 F.M. Follega [ID](#)^{78a,78b}, N. Fomin [ID](#)¹⁶, J.H. Foo [ID](#)¹⁵⁵, B.C. Forland [ID](#)⁶⁸, A. Formica [ID](#)¹³⁵,
 A.C. Forti [ID](#)¹⁰¹, E. Fortin [ID](#)³⁶, A.W. Fortman [ID](#)⁶¹, M.G. Foti [ID](#)^{17a}, L. Fountas [ID](#)^{9,j},
 D. Fournier [ID](#)⁶⁶, H. Fox [ID](#)⁹¹, P. Francavilla [ID](#)^{74a,74b}, S. Francescato [ID](#)⁶¹, S. Franchellucci [ID](#)⁵⁶,
 M. Franchini [ID](#)^{23b,23a}, S. Franchino [ID](#)^{63a}, D. Francis [ID](#)³⁶, L. Franco [ID](#)¹¹³, L. Franconi [ID](#)⁴⁸,
 M. Franklin [ID](#)⁶¹, G. Frattari [ID](#)²⁶, A.C. Freegard [ID](#)⁹⁴, W.S. Freund [ID](#)^{83b}, Y.Y. Frid [ID](#)¹⁵¹,
 J. Friend [ID](#)⁵⁹, N. Fritzsche [ID](#)⁵⁰, A. Froch [ID](#)⁵⁴, D. Froidevaux [ID](#)³⁶, J.A. Frost [ID](#)¹²⁶, Y. Fu [ID](#)^{62a},
 M. Fujimoto [ID](#)^{118,ae}, E. Fullana Torregrosa [ID](#)^{163,*}, K.Y. Fung [ID](#)^{64a}, E. Furtado De Simas Filho [ID](#)^{83b},
 M. Furukawa [ID](#)¹⁵³, J. Fuster [ID](#)¹⁶³, A. Gabrielli [ID](#)^{23b,23a}, A. Gabrielli [ID](#)¹⁵⁵, P. Gadow [ID](#)³⁶,
 G. Gagliardi [ID](#)^{57b,57a}, L.G. Gagnon [ID](#)^{17a}, E.J. Gallas [ID](#)¹²⁶, B.J. Gallop [ID](#)¹³⁴, K.K. Gan [ID](#)¹¹⁹,

S. Ganguly [ID](#)¹⁵³, J. Gao [ID](#)^{62a}, Y. Gao [ID](#)⁵², F.M. Garay Walls [ID](#)^{137a,137b}, B. Garcia^{29,aj},
C. García [ID](#)¹⁶³, A. García Alonso [ID](#)¹¹⁴, A.G. Garcia Caffaro [ID](#)¹⁷², J.E. García Navarro [ID](#)¹⁶³,
M. Garcia-Sciveres [ID](#)^{17a}, G.L. Gardner [ID](#)¹²⁸, R.W. Gardner [ID](#)³⁹, N. Garelli [ID](#)¹⁵⁸, D. Garg [ID](#)⁸⁰,
R.B. Garg [ID](#)^{143,o}, J.M. Gargan⁵², C.A. Garner¹⁵⁵, S.J. Gasiorowski [ID](#)¹³⁸, P. Gaspar [ID](#)^{83b},
G. Gaudio [ID](#)^{73a}, V. Gautam¹³, P. Gauzzi [ID](#)^{75a,75b}, I.L. Gavrilenko [ID](#)³⁷, A. Gavriluk [ID](#)³⁷,
C. Gay [ID](#)¹⁶⁴, G. Gaycken [ID](#)⁴⁸, E.N. Gazis [ID](#)¹⁰, A.A. Geanta [ID](#)^{27b}, C.M. Gee [ID](#)¹³⁶, C. Gemme [ID](#)^{57b},
M.H. Genest [ID](#)⁶⁰, S. Gentile [ID](#)^{75a,75b}, A.D. Gentry [ID](#)¹¹², S. George [ID](#)⁹⁵, W.F. George [ID](#)²⁰,
T. Geralis [ID](#)⁴⁶, P. Gessinger-Befurt [ID](#)³⁶, M.E. Geyik [ID](#)¹⁷¹, M. Ghani [ID](#)¹⁶⁷, M. Ghneimat [ID](#)¹⁴¹,
K. Ghorbanian [ID](#)⁹⁴, A. Ghosal [ID](#)¹⁴¹, A. Ghosh [ID](#)¹⁶⁰, A. Ghosh [ID](#)⁷, B. Giacobbe [ID](#)^{23b},
S. Giagu [ID](#)^{75a,75b}, T. Giani [ID](#)¹¹⁴, P. Giannetti [ID](#)^{74a}, A. Giannini [ID](#)^{62a}, S.M. Gibson [ID](#)⁹⁵,
M. Gignac [ID](#)¹³⁶, D.T. Gil [ID](#)^{86b}, A.K. Gilbert [ID](#)^{86a}, B.J. Gilbert [ID](#)⁴¹, D. Gillberg [ID](#)³⁴, G. Gilles [ID](#)¹¹⁴,
N.E.K. Gillwald [ID](#)⁴⁸, L. Ginabat [ID](#)¹²⁷, D.M. Gingrich [ID](#)^{2,ah}, M.P. Giordani [ID](#)^{69a,69c},
P.F. Giraud [ID](#)¹³⁵, G. Giugliarelli [ID](#)^{69a,69c}, D. Giugni [ID](#)^{71a}, F. Giuli [ID](#)³⁶, I. Gkialas [ID](#)^{9,j},
L.K. Gladilin [ID](#)³⁷, C. Glasman [ID](#)⁹⁹, G.R. Gledhill [ID](#)¹²³, G. Glemža [ID](#)⁴⁸, M. Glisic¹²³,
I. Gnesi [ID](#)^{43b,f}, Y. Go [ID](#)^{29,aj}, M. Goblirsch-Kolb [ID](#)³⁶, B. Gocke [ID](#)⁴⁹, D. Godin¹⁰⁸, B. Gokturk [ID](#)^{21a},
S. Goldfarb [ID](#)¹⁰⁵, T. Golling [ID](#)⁵⁶, M.G.D. Gololo^{33g}, D. Golubkov [ID](#)³⁷, J.P. Gombas [ID](#)¹⁰⁷,
A. Gomes [ID](#)^{130a,130b}, G. Gomes Da Silva [ID](#)¹⁴¹, A.J. Gomez Delegido [ID](#)¹⁶³, R. Gonçalo [ID](#)^{130a,130c},
G. Gonella [ID](#)¹²³, L. Gonella [ID](#)²⁰, A. Gongadze [ID](#)^{149c}, F. Gonnella [ID](#)²⁰, J.L. Gonski [ID](#)⁴¹,
R.Y. González Andana [ID](#)⁵², S. González de la Hoz [ID](#)¹⁶³, S. Gonzalez Fernandez [ID](#)¹³,
R. Gonzalez Lopez [ID](#)⁹², C. Gonzalez Renteria [ID](#)^{17a}, M.V. Gonzalez Rodrigues [ID](#)⁴⁸,
R. Gonzalez Suarez [ID](#)¹⁶¹, S. Gonzalez-Sevilla [ID](#)⁵⁶, G.R. Gonzalvo Rodriguez [ID](#)¹⁶³, L. Goossens [ID](#)³⁶,
B. Gorini [ID](#)³⁶, E. Gorini [ID](#)^{70a,70b}, A. Gorišek [ID](#)⁹³, T.C. Gosart [ID](#)¹²⁸, A.T. Goshaw [ID](#)⁵¹,
M.I. Gostkin [ID](#)³⁸, S. Goswami [ID](#)¹²¹, C.A. Gottardo [ID](#)³⁶, S.A. Gotz [ID](#)¹⁰⁹, M. Gouighri [ID](#)^{35b},
V. Goumarre [ID](#)⁴⁸, A.G. Goussiou [ID](#)¹³⁸, N. Govender [ID](#)^{33c}, I. Grabowska-Bold [ID](#)^{86a}, K. Graham [ID](#)³⁴,
E. Gramstad [ID](#)¹²⁵, S. Grancagnolo [ID](#)^{70a,70b}, M. Grandi [ID](#)¹⁴⁶, C.M. Grant^{1,135}, P.M. Gravila [ID](#)^{27f},
F.G. Gravili [ID](#)^{70a,70b}, H.M. Gray [ID](#)^{17a}, M. Greco [ID](#)^{70a,70b}, C. Grefe [ID](#)²⁴, I.M. Gregor [ID](#)⁴⁸,
P. Grenier [ID](#)¹⁴³, C. Grieco [ID](#)¹³, A.A. Grillo [ID](#)¹³⁶, K. Grimm [ID](#)³¹, S. Grinstein [ID](#)^{13,t},
J.-F. Grivaz [ID](#)⁶⁶, E. Gross [ID](#)¹⁶⁹, J. Grosse-Knetter [ID](#)⁵⁵, C. Grud¹⁰⁶, J.C. Grundy [ID](#)¹²⁶,
L. Guan [ID](#)¹⁰⁶, W. Guan [ID](#)²⁹, C. Gubbels [ID](#)¹⁶⁴, J.G.R. Guerrero Rojas [ID](#)¹⁶³, G. Guerrieri [ID](#)^{69a,69c},
F. Guescini [ID](#)¹¹⁰, R. Gugel [ID](#)¹⁰⁰, J.A.M. Guhit [ID](#)¹⁰⁶, A. Guida [ID](#)¹⁸, T. Guillemin [ID](#)⁴,
E. Guilloton [ID](#)^{167,134}, S. Guindon [ID](#)³⁶, F. Guo [ID](#)^{14a,14e}, J. Guo [ID](#)^{62c}, L. Guo [ID](#)⁴⁸, Y. Guo [ID](#)¹⁰⁶,
R. Gupta [ID](#)⁴⁸, S. Gurbuz [ID](#)²⁴, S.S. Gurdasani [ID](#)⁵⁴, G. Gustavino [ID](#)³⁶, M. Guth [ID](#)⁵⁶,
P. Gutierrez [ID](#)¹²⁰, L.F. Gutierrez Zagazeta [ID](#)¹²⁸, C. Gutsche [ID](#)⁹⁶, C. Gwenlan [ID](#)¹²⁶,
C.B. Gwilliam [ID](#)⁹², E.S. Haaland [ID](#)¹²⁵, A. Haas [ID](#)¹¹⁷, M. Habedank [ID](#)⁴⁸, C. Haber [ID](#)^{17a},
H.K. Hadavand [ID](#)⁸, A. Hadeef [ID](#)¹⁰⁰, S. Hadzic [ID](#)¹¹⁰, J.J. Hahn [ID](#)¹⁴¹, E.H. Haines [ID](#)⁹⁶,
M. Haleem [ID](#)¹⁶⁶, J. Haley [ID](#)¹²¹, J.J. Hall [ID](#)¹³⁹, G.D. Hallelwell [ID](#)¹⁰², L. Halser [ID](#)¹⁹, K. Hamano [ID](#)¹⁶⁵,
M. Hamer [ID](#)²⁴, G.N. Hamity [ID](#)⁵², E.J. Hampshire [ID](#)⁹⁵, J. Han [ID](#)^{62b}, K. Han [ID](#)^{62a}, L. Han [ID](#)^{14c},
L. Han [ID](#)^{62a}, S. Han [ID](#)^{17a}, Y.F. Han [ID](#)¹⁵⁵, K. Hanagaki [ID](#)⁸⁴, M. Hance [ID](#)¹³⁶, D.A. Hangal [ID](#)^{41,ac},
H. Hanif [ID](#)¹⁴², M.D. Hank [ID](#)¹²⁸, R. Hankache [ID](#)¹⁰¹, J.B. Hansen [ID](#)⁴², J.D. Hansen [ID](#)⁴²,
P.H. Hansen [ID](#)⁴², K. Hara [ID](#)¹⁵⁷, D. Harada [ID](#)⁵⁶, T. Harenberg [ID](#)¹⁷¹, S. Harkusha [ID](#)³⁷,
M.L. Harris [ID](#)¹⁰³, Y.T. Harris [ID](#)¹²⁶, J. Harrison [ID](#)¹³, N.M. Harrison [ID](#)¹¹⁹, P.F. Harrison¹⁶⁷,
N.M. Hartman [ID](#)¹¹⁰, N.M. Hartmann [ID](#)¹⁰⁹, Y. Hasegawa [ID](#)¹⁴⁰, A. Hasib [ID](#)⁵², S. Haug [ID](#)¹⁹,
R. Hauser [ID](#)¹⁰⁷, C.M. Hawkes [ID](#)²⁰, R.J. Hawkins [ID](#)³⁶, Y. Hayashi [ID](#)¹⁵³, S. Hayashida [ID](#)¹¹¹,

D. Hayden [107](#), C. Hayes [106](#), R.L. Hayes [114](#), C.P. Hays [126](#), J.M. Hays [94](#),
H.S. Hayward [92](#), F. He [62a](#), M. He [14a,14e](#), Y. He [154](#), Y. He [48](#), N.B. Heatley [94](#),
V. Hedberg [98](#), A.L. Heggelund [125](#), N.D. Hehir [94](#), C. Heidegger [54](#), K.K. Heidegger [54](#),
W.D. Heidorn [81](#), J. Heilman [34](#), S. Heim [48](#), T. Heim [17a](#), J.G. Heinlein [128](#),
J.J. Heinrich [123](#), L. Heinrich [110,af](#), J. Hejbal [131](#), L. Helary [48](#), A. Held [170](#),
S. Hellesund [16](#), C.M. Helling [164](#), S. Hellman [47a,47b](#), R.C.W. Henderson [91](#), L. Henkelmann [32](#),
A.M. Henriques Correia [36](#), H. Herde [98](#), Y. Hernández Jiménez [145](#), L.M. Herrmann [24](#),
T. Herrmann [50](#), G. Herten [54](#), R. Hertenberger [109](#), L. Hervas [36](#), M.E. Hespings [100](#),
N.P. Hessey [156a](#), H. Hibi [85](#), S.J. Hillier [20](#), J.R. Hinds [107](#), F. Hinterkeuser [24](#),
M. Hirose [124](#), S. Hirose [157](#), D. Hirschbuehl [171](#), T.G. Hitchings [101](#), B. Hiti [93](#),
J. Hobbs [145](#), R. Hobincu [27e](#), N. Hod [169](#), M.C. Hodgkinson [139](#), B.H. Hodgkinson [32](#),
A. Hoecker [36](#), J. Hofer [48](#), T. Holm [24](#), M. Holzbock [110](#), L.B.A.H. Hommels [32](#),
B.P. Honan [101](#), J. Hong [62c](#), T.M. Hong [129](#), B.H. Hooberman [162](#), W.H. Hopkins [6](#),
Y. Horii [111](#), S. Hou [148](#), A.S. Howard [93](#), J. Howarth [59](#), J. Hoya [6](#), M. Hrabovsky [122](#),
A. Hrynevich [48](#), T. Hryn'ova [4](#), P.J. Hsu [65](#), S.-C. Hsu [138](#), Q. Hu [62a](#), Y.F. Hu [14a,14e](#),
S. Huang [64b](#), X. Huang [14c](#), Y. Huang [139](#), Y. Huang [14a](#), Z. Huang [101](#), Z. Hubacek [132](#),
M. Huebner [24](#), F. Huegging [24](#), T.B. Huffman [126](#), C.A. Hugli [48](#), M. Huhtinen [36](#),
S.K. Huiberts [16](#), R. Hulsken [104](#), N. Huseynov [12,a](#), J. Huston [107](#), J. Huth [61](#),
R. Hyneman [143](#), G. Iacobucci [56](#), G. Iakovidis [29](#), I. Ibragimov [141](#), L. Iconomidou-Fayard [66](#),
P. Iengo [72a,72b](#), R. Iguchi [153](#), T. Iizawa [126](#), Y. Ikegami [84](#), N. Ilic [155](#), H. Imam [35a](#),
M. Ince Lezki [56](#), T. Ingebretsen Carlson [47a,47b](#), G. Introzzi [73a,73b](#), M. Iodice [77a](#),
V. Ippolito [75a,75b](#), R.K. Irwin [92](#), M. Ishino [153](#), W. Islam [170](#), C. Issever [18,48](#),
S. Istin [21a,al](#), H. Ito [168](#), J.M. Iturbe Ponce [64a](#), R. Iuppa [78a,78b](#), A. Ivina [169](#), J.M. Izen [45](#),
V. Izzo [72a](#), P. Jacka [131,132](#), P. Jackson [1](#), R.M. Jacobs [48](#), B.P. Jaeger [142](#), C.S. Jagfeld [109](#),
G. Jain [156a](#), P. Jain [54](#), G. Jäkel [171](#), K. Jakobs [54](#), T. Jakoubek [169](#), J. Jamieson [59](#),
K.W. Janas [86a](#), M. Javurkova [103](#), F. Jeanneau [135](#), L. Jeanty [123](#), J. Jejelava [149a,aa](#),
P. Jenni [54,g](#), C.E. Jessiman [34](#), S. Jézéquel [4](#), C. Jia [62b](#), J. Jia [145](#), X. Jia [61](#), X. Jia [14a,14e](#),
Z. Jia [14c](#), Y. Jiang [62a](#), S. Jiggins [48](#), J. Jimenez Pena [13](#), S. Jin [14c](#), A. Jinaru [27b](#),
O. Jinnouchi [154](#), P. Johansson [139](#), K.A. Johns [7](#), J.W. Johnson [136](#), D.M. Jones [32](#),
E. Jones [48](#), P. Jones [32](#), R.W.L. Jones [91](#), T.J. Jones [92](#), H.L. Joos [55,36](#), R. Joshi [119](#),
J. Jovicevic [15](#), X. Ju [17a](#), J.J. Junggeburth [103](#), T. Junkermann [63a](#), A. Juste Rozas [13,t](#),
M.K. Juzek [87](#), S. Kabana [137e](#), A. Kaczmarek [87](#), M. Kado [110](#), H. Kagan [119](#),
M. Kagan [143](#), A. Kahn [41](#), A. Kahn [128](#), C. Kahra [100](#), T. Kaji [153](#), E. Kajomovitz [150](#),
N. Kakati [169](#), I. Kalaitzidou [54](#), C.W. Kalderon [29](#), A. Kamenshchikov [155](#), N.J. Kang [136](#),
D. Kar [33g](#), K. Karava [126](#), M.J. Kareem [156b](#), E. Karentzos [54](#), I. Karkanas [152](#),
O. Karkout [114](#), S.N. Karpov [38](#), Z.M. Karpova [38](#), V. Kartvelishvili [91](#), A.N. Karyukhin [37](#),
E. Kasimi [152](#), J. Katzy [48](#), S. Kaur [34](#), K. Kawade [140](#), M.P. Kawale [120](#), T. Kawamoto [135](#),
E.F. Kay [36](#), F.I. Kaya [158](#), S. Kazakos [107](#), V.F. Kazanin [37](#), Y. Ke [145](#), J.M. Keaveney [33a](#),
R. Keeler [165](#), G.V. Kehris [61](#), J.S. Keller [34](#), A.S. Kelly [96](#), J.J. Kempster [146](#),
K.E. Kennedy [41](#), P.D. Kennedy [100](#), O. Kepka [131](#), B.P. Kerridge [167](#), S. Kersten [171](#),
B.P. Kerševan [93](#), S. Keshri [66](#), L. Keszeghova [28a](#), S. Ketabchi Haghighat [155](#),
M. Khandoga [127](#), A. Khanov [121](#), A.G. Kharlamov [37](#), T. Kharlamova [37](#), E.E. Khoda [138](#),
T.J. Khoo [18](#), G. Khoriauli [166](#), J. Khubua [149b](#), Y.A.R. Khwaira [66](#), A. Kilgallon [123](#),

D.W. Kim [ID](#)^{47a,47b}, Y.K. Kim [ID](#)³⁹, N. Kimura [ID](#)⁹⁶, M.K. Kingston [ID](#)⁵⁵, A. Kirchhoff [ID](#)⁵⁵,
C. Kirfel [ID](#)²⁴, F. Kirfel [ID](#)²⁴, J. Kirk [ID](#)¹³⁴, A.E. Kiryunin [ID](#)¹¹⁰, C. Kitsaki [ID](#)¹⁰, O. Kivernyk [ID](#)²⁴,
M. Klassen [ID](#)^{63a}, C. Klein [ID](#)³⁴, L. Klein [ID](#)¹⁶⁶, M.H. Klein [ID](#)¹⁰⁶, M. Klein [ID](#)⁹², S.B. Klein [ID](#)⁵⁶,
U. Klein [ID](#)⁹², P. Klimek [ID](#)³⁶, A. Klimentov [ID](#)²⁹, T. Klioutchnikova [ID](#)³⁶, P. Kluit [ID](#)¹¹⁴, S. Kluth [ID](#)¹¹⁰,
E. Kneringer [ID](#)⁷⁹, T.M. Knight [ID](#)¹⁵⁵, A. Knue [ID](#)⁴⁹, R. Kobayashi [ID](#)⁸⁸, D. Kobylanski [ID](#)¹⁶⁹,
S.F. Koch [ID](#)¹²⁶, M. Kocian [ID](#)¹⁴³, P. Kodyš [ID](#)¹³³, D.M. Koeck [ID](#)¹²³, P.T. Koenig [ID](#)²⁴, T. Koffas [ID](#)³⁴,
M. Kolb [ID](#)¹³⁵, I. Koletsou [ID](#)⁴, T. Komarek [ID](#)¹²², K. Köneke [ID](#)⁵⁴, A.X.Y. Kong [ID](#)¹, T. Kono [ID](#)¹¹⁸,
N. Konstantinidis [ID](#)⁹⁶, B. Konya [ID](#)⁹⁸, R. Kopeliansky [ID](#)⁶⁸, S. Koperny [ID](#)^{86a}, K. Korcyl [ID](#)⁸⁷,
K. Kordas [ID](#)^{152,e}, G. Koren [ID](#)¹⁵¹, A. Korn [ID](#)⁹⁶, S. Korn [ID](#)⁵⁵, I. Korolkov [ID](#)¹³, N. Korotkova [ID](#)³⁷,
B. Kortman [ID](#)¹¹⁴, O. Kortner [ID](#)¹¹⁰, S. Kortner [ID](#)¹¹⁰, W.H. Kostecka [ID](#)¹¹⁵, V.V. Kostyukhin [ID](#)¹⁴¹,
A. Kotsokechagia [ID](#)¹³⁵, A. Kotwal [ID](#)⁵¹, A. Koulouris [ID](#)³⁶, A. Kourkoumeli-Charalampidi [ID](#)^{73a,73b},
C. Kourkoumelis [ID](#)⁹, E. Kourlitis [ID](#)^{110,af}, O. Kovanda [ID](#)¹⁴⁶, R. Kowalewski [ID](#)¹⁶⁵,
W. Kozanecki [ID](#)¹³⁵, A.S. Kozhin [ID](#)³⁷, V.A. Kramarenko [ID](#)³⁷, G. Kramberger [ID](#)⁹³, P. Kramer [ID](#)¹⁰⁰,
M.W. Krasny [ID](#)¹²⁷, A. Krasznahorkay [ID](#)³⁶, J.W. Kraus [ID](#)¹⁷¹, J.A. Kremer [ID](#)¹⁰⁰, T. Kresse [ID](#)⁵⁰,
J. Kretzschmar [ID](#)⁹², K. Kreul [ID](#)¹⁸, P. Krieger [ID](#)¹⁵⁵, S. Krishnamurthy [ID](#)¹⁰³, M. Krivos [ID](#)¹³³,
K. Krizka [ID](#)²⁰, K. Kroeninger [ID](#)⁴⁹, H. Kroha [ID](#)¹¹⁰, J. Kroll [ID](#)¹³¹, J. Kroll [ID](#)¹²⁸,
K.S. Krowpman [ID](#)¹⁰⁷, U. Kruchonak [ID](#)³⁸, H. Krüger [ID](#)²⁴, N. Krumnack [ID](#)⁸¹, M.C. Kruse [ID](#)⁵¹,
J.A. Krzysiak [ID](#)⁸⁷, O. Kuchinskaia [ID](#)³⁷, S. Kuday [ID](#)^{3a}, S. Kuehn [ID](#)³⁶, R. Kuesters [ID](#)⁵⁴, T. Kuhl [ID](#)⁴⁸,
V. Kukhtin [ID](#)³⁸, Y. Kulchitsky [ID](#)^{37,a}, S. Kuleshov [ID](#)^{137d,137b}, M. Kumar [ID](#)^{33g}, N. Kumari [ID](#)⁴⁸,
A. Kupco [ID](#)¹³¹, T. Kupfer [ID](#)⁴⁹, A. Kupich [ID](#)³⁷, O. Kuprash [ID](#)⁵⁴, H. Kurashige [ID](#)⁸⁵,
L.L. Kurchaninov [ID](#)^{156a}, O. Kurdysh [ID](#)⁶⁶, Y.A. Kurochkin [ID](#)³⁷, A. Kurova [ID](#)³⁷, M. Kuze [ID](#)¹⁵⁴,
A.K. Kvam [ID](#)¹⁰³, J. Kvita [ID](#)¹²², T. Kwan [ID](#)¹⁰⁴, N.G. Kyriacou [ID](#)¹⁰⁶, L.A.O. Laatu [ID](#)¹⁰²,
C. Lacasta [ID](#)¹⁶³, F. Lacava [ID](#)^{75a,75b}, H. Lacker [ID](#)¹⁸, D. Lacour [ID](#)¹²⁷, N.N. Lad [ID](#)⁹⁶, E. Ladygin [ID](#)³⁸,
B. Laforge [ID](#)¹²⁷, T. Lagouri [ID](#)^{137e}, F.Z. Lahbabi [ID](#)^{35a}, S. Lai [ID](#)⁵⁵, I.K. Lakomic [ID](#)^{86a},
N. Lalloue [ID](#)⁶⁰, J.E. Lambert [ID](#)¹⁶⁵, S. Lammers [ID](#)⁶⁸, W. Lampl [ID](#)⁷, C. Lampoudis [ID](#)^{152,e},
A.N. Lancaster [ID](#)¹¹⁵, E. Lançon [ID](#)²⁹, U. Landgraf [ID](#)⁵⁴, M.P.J. Landon [ID](#)⁹⁴, V.S. Lang [ID](#)⁵⁴,
R.J. Langenberg [ID](#)¹⁰³, O.K.B. Langrekken [ID](#)¹²⁵, A.J. Lankford [ID](#)¹⁶⁰, F. Lanni [ID](#)³⁶, K. Lantzsck [ID](#)²⁴,
A. Lanza [ID](#)^{73a}, A. Lapertosa [ID](#)^{57b,57a}, J.F. Laporte [ID](#)¹³⁵, T. Lari [ID](#)^{71a}, F. Lasagni Manghi [ID](#)^{23b},
M. Lassnig [ID](#)³⁶, V. Latonova [ID](#)¹³¹, A. Laudrain [ID](#)¹⁰⁰, A. Laurier [ID](#)¹⁵⁰, S.D. Lawlor [ID](#)⁹⁵,
Z. Lawrence [ID](#)¹⁰¹, M. Lazzaroni [ID](#)^{71a,71b}, B. Le [ID](#)¹⁰¹, E.M. Le Boulicaut [ID](#)⁵¹, B. Leban [ID](#)⁹³,
A. Lebedev [ID](#)⁸¹, M. LeBlanc [ID](#)¹⁰¹, F. Ledroit-Guillon [ID](#)⁶⁰, A.C.A. Lee [ID](#)⁹⁶, S.C. Lee [ID](#)¹⁴⁸,
S. Lee [ID](#)^{47a,47b}, T.F. Lee [ID](#)⁹², L.L. Leeuw [ID](#)^{33c}, H.P. Lefebvre [ID](#)⁹⁵, M. Lefebvre [ID](#)¹⁶⁵,
C. Leggett [ID](#)^{17a}, G. Lehmann Miotto [ID](#)³⁶, M. Leigh [ID](#)⁵⁶, W.A. Leight [ID](#)¹⁰³, W. Leinonen [ID](#)¹¹³,
A. Leisos [ID](#)^{152,s}, M.A.L. Leite [ID](#)^{83c}, C.E. Leitgeb [ID](#)⁴⁸, R. Leitner [ID](#)¹³³, K.J.C. Leney [ID](#)⁴⁴,
T. Lenz [ID](#)²⁴, S. Leone [ID](#)^{74a}, C. Leonidopoulos [ID](#)⁵², A. Leopold [ID](#)¹⁴⁴, C. Leroy [ID](#)¹⁰⁸, R. Les [ID](#)¹⁰⁷,
C.G. Lester [ID](#)³², M. Levchenko [ID](#)³⁷, J. Levêque [ID](#)⁴, D. Levin [ID](#)¹⁰⁶, L.J. Levinson [ID](#)¹⁶⁹,
M.P. Lewicki [ID](#)⁸⁷, D.J. Lewis [ID](#)⁴, A. Li [ID](#)⁵, B. Li [ID](#)^{62b}, C. Li [ID](#)^{62a}, C-Q. Li [ID](#)^{62c}, H. Li [ID](#)^{62a}, H. Li [ID](#)^{62b},
H. Li [ID](#)^{14c}, H. Li [ID](#)^{14b}, H. Li [ID](#)^{62b}, K. Li [ID](#)¹³⁸, L. Li [ID](#)^{62c}, M. Li [ID](#)^{14a,14e}, Q.Y. Li [ID](#)^{62a}, S. Li [ID](#)^{14a,14e},
S. Li [ID](#)^{62d,62c,d}, T. Li [ID](#)⁵, X. Li [ID](#)¹⁰⁴, Z. Li [ID](#)¹²⁶, Z. Li [ID](#)¹⁰⁴, Z. Li [ID](#)⁹², Z. Li [ID](#)^{14a,14e}, S. Liang [ID](#)^{14a,14e},
Z. Liang [ID](#)^{14a}, M. Liberatore [ID](#)¹³⁵, B. Liberti [ID](#)^{76a}, K. Lie [ID](#)^{64c}, J. Lieber Marin [ID](#)^{83b}, H. Lien [ID](#)⁶⁸,
K. Lin [ID](#)¹⁰⁷, R.E. Lindley [ID](#)⁷, J.H. Lindon [ID](#)², E. Lipeles [ID](#)¹²⁸, A. Lipniacka [ID](#)¹⁶, A. Lister [ID](#)¹⁶⁴,
J.D. Little [ID](#)⁴, B. Liu [ID](#)^{14a}, B.X. Liu [ID](#)¹⁴², D. Liu [ID](#)^{62d,62c}, J.B. Liu [ID](#)^{62a}, J.K.K. Liu [ID](#)³²,
K. Liu [ID](#)^{62d,62c}, M. Liu [ID](#)^{62a}, M.Y. Liu [ID](#)^{62a}, P. Liu [ID](#)^{14a}, Q. Liu [ID](#)^{62d,138,62c}, X. Liu [ID](#)^{62a},

Y. Liu [14d,14e](#), Y.L. Liu [62b](#), Y.W. Liu [62a](#), J. Llorente Merino [142](#), S.L. Lloyd [94](#),
 E.M. Lobodzinska [48](#), P. Loch [7](#), S. Loffredo [76a,76b](#), T. Lohse [18](#), K. Lohwasser [139](#),
 E. Loiacono [48](#), M. Lokajicek [131,*](#), J.D. Lomas [20](#), J.D. Long [162](#), I. Longarini [160](#),
 L. Longo [70a,70b](#), R. Longo [162](#), I. Lopez Paz [67](#), A. Lopez Solis [48](#), J. Lorenz [109](#),
 N. Lorenzo Martinez [4](#), A.M. Lory [109](#), G. Lösckce Centeno [146](#), O. Loseva [37](#),
 X. Lou [47a,47b](#), X. Lou [14a,14e](#), A. Lounis [66](#), J. Love [6](#), P.A. Love [91](#), G. Lu [14a,14e](#),
 M. Lu [80](#), S. Lu [128](#), Y.J. Lu [65](#), H.J. Lubatti [138](#), C. Luci [75a,75b](#), F.L. Lucio Alves [14c](#),
 A. Lucotte [60](#), F. Luehring [68](#), I. Luise [145](#), O. Lukianchuk [66](#), O. Lundberg [144](#),
 B. Lund-Jensen [144](#), N.A. Luongo [123](#), M.S. Lutz [151](#), D. Lynn [29](#), H. Lyons [92](#), R. Lysak [131](#),
 E. Lytken [98](#), V. Lyubushkin [38](#), T. Lyubushkina [38](#), M.M. Lyukova [145](#), H. Ma [29](#), K. Ma [62a](#),
 L.L. Ma [62b](#), Y. Ma [121](#), D.M. Mac Donell [165](#), G. Maccarrone [53](#), J.C. MacDonald [100](#),
 P.C. Machado De Abreu Farias [83b](#), R. Madar [40](#), W.F. Mader [50](#), T. Madula [96](#), J. Maeda [85](#),
 T. Maeno [29](#), M. Maerker [50](#), H. Maguire [139](#), V. Maiboroda [135](#), A. Maio [130a,130b,130d](#),
 K. Maj [86a](#), O. Majersky [48](#), S. Majewski [123](#), N. Makovec [66](#), V. Maksimovic [15](#),
 B. Malaescu [127](#), Pa. Malecki [87](#), V.P. Maleev [37](#), F. Malek [60](#), M. Mali [93](#), D. Malito [95](#),
 U. Mallik [80](#), S. Maltezos [10](#), S. Malyukov [38](#), J. Mamuzic [13](#), G. Mancini [53](#), G. Manco [73a,73b](#),
 J.P. Mandalia [94](#), I. Mandić [93](#), L. Manhaes de Andrade Filho [83a](#), I.M. Maniatis [169](#),
 J. Manjarres Ramos [102,ab](#), D.C. Mankad [169](#), A. Mann [109](#), B. Mansoulie [135](#), S. Manzoni [36](#),
 A. Marantis [152,s](#), G. Marchiori [5](#), M. Marcisovsky [131](#), C. Marcon [71a,71b](#), M. Marinescu [20](#),
 M. Marjanovic [120](#), E.J. Marshall [91](#), Z. Marshall [17a](#), S. Marti-Garcia [163](#), T.A. Martin [167](#),
 V.J. Martin [52](#), B. Martin dit Latour [16](#), L. Martinelli [75a,75b](#), M. Martinez [13,t](#),
 P. Martinez Agullo [163](#), V.I. Martinez Outschoorn [103](#), P. Martinez Suarez [13](#),
 S. Martin-Haugh [134](#), V.S. Martoiu [27b](#), A.C. Martyniuk [96](#), A. Marzin [36](#),
 D. Mascione [78a,78b](#), L. Masetti [100](#), T. Mashimo [153](#), J. Masik [101](#), A.L. Maslennikov [37](#),
 L. Massa [23b](#), P. Massarotti [72a,72b](#), P. Mastrandrea [74a,74b](#), A. Mastroberardino [43b,43a](#),
 T. Masubuchi [153](#), T. Mathisen [161](#), J. Matousek [133](#), N. Matsuzawa [153](#), J. Maurer [27b](#),
 B. Maček [93](#), D.A. Maximov [37](#), R. Mazini [148](#), I. Maznas [152](#), M. Mazza [107](#),
 S.M. Mazza [136](#), E. Mazzeo [71a,71b](#), C. Mc Ginn [29](#), J.P. Mc Gowan [104](#), S.P. Mc Kee [106](#),
 E.F. McDonald [105](#), A.E. McDougall [114](#), J.A. Mcfayden [146](#), R.P. McGovern [128](#),
 G. Mchedlidze [149b](#), R.P. McKenzie [33g](#), T.C. Mclachlan [48](#), D.J. Mclaughlin [96](#),
 K.D. McLean [165](#), S.J. McMahon [134](#), P.C. McNamara [105](#), C.M. Mcpartland [92](#),
 R.A. McPherson [165,x](#), S. Mehlhase [109](#), A. Mehta [92](#), D. Melini [150](#), B.R. Mellado Garcia [33g](#),
 A.H. Melo [55](#), F. Meloni [48](#), A.M. Mendes Jacques Da Costa [101](#), H.Y. Meng [155](#), L. Meng [91](#),
 S. Menke [110](#), M. Mentink [36](#), E. Meoni [43b,43a](#), C. Merlassino [126](#), L. Merola [72a,72b](#),
 C. Meroni [71a,71b](#), G. Merz [106](#), O. Meshkov [37](#), J. Metcalfe [6](#), A.S. Mete [6](#), C. Meyer [68](#),
 J-P. Meyer [135](#), R.P. Middleton [134](#), L. Mijović [52](#), G. Mikenberg [169](#), M. Mikestikova [131](#),
 M. Mikuž [93](#), H. Mildner [100](#), A. Milic [36](#), C.D. Milke [44](#), D.W. Miller [39](#), L.S. Miller [34](#),
 A. Milov [169](#), D.A. Milstead [47a,47b](#), T. Min [14c](#), A.A. Minaenko [37](#), I.A. Minashvili [149b](#),
 L. Mince [59](#), A.I. Mincer [117](#), B. Mindur [86a](#), M. Mineev [38](#), Y. Mino [88](#), L.M. Mir [13](#),
 M. Miralles Lopez [163](#), M. Mironova [17a](#), A. Mishima [153](#), M.C. Missio [113](#), A. Mitra [167](#),
 V.A. Mitsou [163](#), Y. Mitsumori [111](#), O. Miu [155](#), P.S. Miyagawa [94](#), T. Mkrtchyan [63a](#),
 M. Mlinarevic [96](#), T. Mlinarevic [96](#), M. Mlynarikova [36](#), S. Mobius [19](#), P. Moder [48](#),
 P. Mogg [109](#), A.F. Mohammed [14a,14e](#), S. Mohapatra [41](#), G. Mokgatitwane [33g](#), L. Moleri [169](#),































































































































B. Mondal¹⁴¹, S. Mondal¹³², K. Mönig⁴⁸, E. Monnier¹⁰², L. Monsonis Romero¹⁶³,
 J. Montejo Berlingen¹³, M. Montella¹¹⁹, F. Montekali^{77a,77b}, F. Monticelli⁹⁰,
 S. Monzani^{69a,69c}, N. Morange⁶⁶, A.L. Moreira De Carvalho^{130a}, M. Moreno Llácer¹⁶³,
 C. Moreno Martinez⁵⁶, P. Morettini^{57b}, S. Morgenstern³⁶, M. Morii⁶¹, M. Morinaga¹⁵³,
 A.K. Morley³⁶, F. Morodei^{75a,75b}, L. Morvaj³⁶, P. Moschovakos³⁶, B. Moser³⁶,
 M. Mosidze^{149b}, T. Moskalets⁵⁴, P. Moskvitina¹¹³, J. Moss^{31,m}, E.J.W. Moyses¹⁰³,
 O. Mtintsilana^{33g}, S. Muanza¹⁰², J. Mueller¹²⁹, D. Muenstermann⁹¹, R. Müller¹⁹,
 G.A. Mullier¹⁶¹, A.J. Mullin³², J.J. Mullin¹²⁸, D.P. Mungo¹⁵⁵, D. Munoz Perez¹⁶³,
 F.J. Munoz Sanchez¹⁰¹, M. Murin¹⁰¹, W.J. Murray^{167,134}, A. Murrone^{71a,71b},
 J.M. Muse¹²⁰, M. Muškinja^{17a}, C. Mwewa²⁹, A.G. Myagkov^{37,a}, A.J. Myers⁸,
 A.A. Myers¹²⁹, G. Myers⁶⁸, M. Myska¹³², B.P. Nachman^{17a}, O. Nackenhorst⁴⁹,
 A. Nag⁵⁰, K. Nagai¹²⁶, K. Nagano⁸⁴, J.L. Nagle^{29,aj}, E. Nagy¹⁰², A.M. Nairz³⁶,
 Y. Nakahama⁸⁴, K. Nakamura⁸⁴, K. Nakkalil⁵, H. Nanjo¹²⁴, R. Narayan⁴⁴,
 E.A. Narayanan¹¹², I. Naryshkin³⁷, M. Naseri³⁴, S. Nasri¹⁵⁹, C. Nass²⁴, G. Navarro^{22a},
 J. Navarro-Gonzalez¹⁶³, R. Nayak¹⁵¹, A. Nayaz¹⁸, P.Y. Nechaeva³⁷, F. Nechansky⁴⁸,
 L. Nedic¹²⁶, T.J. Neep²⁰, A. Negri^{73a,73b}, M. Negrini^{23b}, C. Nellist¹¹⁴, C. Nelson¹⁰⁴,
 K. Nelson¹⁰⁶, S. Nemecek¹³¹, M. Nessi^{36,h}, M.S. Neubauer¹⁶², F. Neuhaus¹⁰⁰,
 J. Neundorff⁴⁸, R. Newhouse¹⁶⁴, P.R. Newman²⁰, C.W. Ng¹²⁹, Y.W.Y. Ng⁴⁸,
 B. Ngair^{35e}, H.D.N. Nguyen¹⁰⁸, R.B. Nickerson¹²⁶, R. Nicolaidou¹³⁵, J. Nielsen¹³⁶,
 M. Niemeyer⁵⁵, J. Niermann^{55,36}, N. Nikiforou³⁶, V. Nikolaenko^{37,a}, I. Nikolic-Audit¹²⁷,
 K. Nikolopoulos²⁰, P. Nilsson²⁹, I. Ninca⁴⁸, H.R. Nindhito⁵⁶, G. Ninio¹⁵¹, A. Nisati^{75a},
 N. Nishu², R. Nisius¹¹⁰, J-E. Nitschke⁵⁰, E.K. Nkadimeng^{33g}, T. Nobe¹⁵³, D.L. Noel³²,
 T. Nommensen¹⁴⁷, M.B. Norfolk¹³⁹, R.R.B. Norisam⁹⁶, B.J. Norman³⁴, J. Novak⁹³,
 T. Novak⁴⁸, L. Novotny¹³², R. Novotny¹¹², L. Nozka¹²², K. Ntekas¹⁶⁰,
 N.M.J. Nunes De Moura Junior^{83b}, E. Nurse⁹⁶, J. Ocariz¹²⁷, A. Ochi⁸⁵, I. Ochoa^{130a},
 S. Oerdek⁴⁸, J.T. Offermann³⁹, A. Ogrodnik¹³³, A. Oh¹⁰¹, C.C. Ohm¹⁴⁴, H. Oide⁸⁴,
 R. Oishi¹⁵³, M.L. Ojeda⁴⁸, M.W. O’Keefe⁹², Y. Okumura¹⁵³, L.F. Oleiro Seabra^{130a},
 S.A. Olivares Pino^{137d}, D. Oliveira Damazio²⁹, D. Oliveira Goncalves^{83a}, J.L. Oliver¹⁶⁰,
 A. Olszewski⁸⁷, Ö.O. Öncel⁵⁴, A.P. O’Neill¹⁹, A. Onofre^{130a,130e}, P.U.E. Onyisi¹¹,
 M.J. Oreglia³⁹, G.E. Orellana⁹⁰, D. Orestano^{77a,77b}, N. Orlando¹³, R.S. Orr¹⁵⁵,
 V. O’Shea⁵⁹, L.M. Osojnak¹²⁸, R. Ospanov^{62a}, G. Otero y Garzon³⁰, H. Otono⁸⁹,
 P.S. Ott^{63a}, G.J. Ottino^{17a}, M. Ouchrif^{35d}, J. Ouellette²⁹, F. Ould-Saada¹²⁵,
 M. Owen⁵⁹, R.E. Owen¹³⁴, K.Y. Oyulmaz^{21a}, V.E. Ozcan^{21a}, N. Ozturk⁸, S. Ozturk⁸²,
 H.A. Pacey¹²⁶, A. Pacheco Pages¹³, C. Padilla Aranda¹³, G. Padovano^{75a,75b},
 S. Pagan Griso^{17a}, G. Palacino⁶⁸, A. Palazzo^{70a,70b}, S. Palestini³⁶, J. Pan¹⁷²,
 T. Pan^{64a}, D.K. Panchal¹¹, C.E. Pandini¹¹⁴, J.G. Panduro Vazquez⁹⁵, H.D. Pandya¹,
 H. Pang^{14b}, P. Pani⁴⁸, G. Panizzo^{69a,69c}, L. Paolozzi⁵⁶, C. Papadatos¹⁰⁸, S. Parajuli⁴⁴,
 A. Paramonov⁶, C. Paraskevopoulos¹⁰, D. Paredes Hernandez^{64b}, T.H. Park¹⁵⁵,
 M.A. Parker³², F. Parodi^{57b,57a}, E.W. Parrish¹¹⁵, V.A. Parrish⁵², J.A. Parsons⁴¹,
 U. Parzefall⁵⁴, B. Pascual Dias¹⁰⁸, L. Pascual Dominguez¹⁵¹, E. Pasqualucci^{75a},
 S. Passaggio^{57b}, F. Pastore⁹⁵, P. Pasuwan^{47a,47b}, P. Patel⁸⁷, U.M. Patel⁵¹,
 J.R. Pater¹⁰¹, T. Pauly³⁶, J. Pearkes¹⁴³, M. Pedersen¹²⁵, R. Pedro^{130a},
 S.V. Peleganchuk³⁷, O. Penc³⁶, E.A. Pender⁵², H. Peng^{62a}, K.E. Pinski¹⁰⁹,

M. Penzin [ID](#)³⁷, B.S. Peralva [ID](#)^{83d}, A.P. Pereira Peixoto [ID](#)⁶⁰, L. Pereira Sanchez [ID](#)^{47a,47b},
D.V. Perepelitsa [ID](#)^{29,aj}, E. Perez Codina [ID](#)^{156a}, M. Perganti [ID](#)¹⁰, L. Perini [ID](#)^{71a,71b,*},
H. Pernegger [ID](#)³⁶, O. Perrin [ID](#)⁴⁰, K. Peters [ID](#)⁴⁸, R.F.Y. Peters [ID](#)¹⁰¹, B.A. Petersen [ID](#)³⁶,
T.C. Petersen [ID](#)⁴², E. Petit [ID](#)¹⁰², V. Petousis [ID](#)¹³², C. Petridou [ID](#)^{152,e}, A. Petrukhin [ID](#)¹⁴¹,
M. Pettee [ID](#)^{17a}, N.E. Pettersson [ID](#)³⁶, A. Petukhov [ID](#)³⁷, K. Petukhova [ID](#)¹³³, R. Pezoa [ID](#)^{137f},
L. Pezzotti [ID](#)³⁶, G. Pezzullo [ID](#)¹⁷², T.M. Pham [ID](#)¹⁷⁰, T. Pham [ID](#)¹⁰⁵, P.W. Phillips [ID](#)¹³⁴,
G. Piacquadio [ID](#)¹⁴⁵, E. Pianori [ID](#)^{17a}, F. Piazza [ID](#)^{71a,71b}, R. Piegaia [ID](#)³⁰, D. Pietreanu [ID](#)^{27b},
A.D. Pilkington [ID](#)¹⁰¹, M. Pinamonti [ID](#)^{69a,69c}, J.L. Pinfeld [ID](#)², B.C. Pinheiro Pereira [ID](#)^{130a},
A.E. Pinto Pinoargote [ID](#)^{100,135}, L. Pintucci [ID](#)^{69a,69c}, K.M. Piper [ID](#)¹⁴⁶, A. Pirttikoski [ID](#)⁵⁶,
D.A. Pizzi [ID](#)³⁴, L. Pizzimento [ID](#)^{64b}, A. Pizzini [ID](#)¹¹⁴, M.-A. Pleier [ID](#)²⁹, V. Plesanovs⁵⁴,
V. Pleskot [ID](#)¹³³, E. Plotnikova³⁸, G. Poddar [ID](#)⁴, R. Poettgen [ID](#)⁹⁸, L. Poggioli [ID](#)¹²⁷, I. Pokharel [ID](#)⁵⁵,
S. Polacek [ID](#)¹³³, G. Polesello [ID](#)^{73a}, A. Poley [ID](#)^{142,156a}, R. Polifka [ID](#)¹³², A. Polini [ID](#)^{23b},
C.S. Pollard [ID](#)¹⁶⁷, Z.B. Pollock [ID](#)¹¹⁹, V. Polychronakos [ID](#)²⁹, E. Pompa Pacchi [ID](#)^{75a,75b},
D. Ponomarenko [ID](#)¹¹³, L. Pontecorvo [ID](#)³⁶, S. Popa [ID](#)^{27a}, G.A. Popeneciu [ID](#)^{27d}, A. Poreba [ID](#)³⁶,
D.M. Portillo Quintero [ID](#)^{156a}, S. Pospisil [ID](#)¹³², M.A. Postill [ID](#)¹³⁹, P. Postolache [ID](#)^{27c},
K. Potamianos [ID](#)¹⁶⁷, P.A. Potepa [ID](#)^{86a}, I.N. Potrap [ID](#)³⁸, C.J. Potter [ID](#)³², H. Potti [ID](#)¹,
T. Poulsen [ID](#)⁴⁸, J. Poveda [ID](#)¹⁶³, M.E. Pozo Astigarraga [ID](#)³⁶, A. Prades Ibanez [ID](#)¹⁶³, J. Pretel [ID](#)⁵⁴,
D. Price [ID](#)¹⁰¹, M. Primavera [ID](#)^{70a}, M.A. Principe Martin [ID](#)⁹⁹, R. Privara [ID](#)¹²², T. Procter [ID](#)⁵⁹,
M.L. Proffitt [ID](#)¹³⁸, N. Proklova [ID](#)¹²⁸, K. Prokofiev [ID](#)^{64c}, G. Proto [ID](#)¹¹⁰, S. Protopopescu [ID](#)²⁹,
J. Proudfoot [ID](#)⁶, M. Przybycien [ID](#)^{86a}, W.W. Przygoda [ID](#)^{86b}, J.E. Puddefoot [ID](#)¹³⁹, D. Pudzha [ID](#)³⁷,
D. Pyatiizbyantseva [ID](#)³⁷, J. Qian [ID](#)¹⁰⁶, D. Qichen [ID](#)¹⁰¹, Y. Qin [ID](#)¹⁰¹, T. Qiu [ID](#)⁵², A. Quadt [ID](#)⁵⁵,
M. Queitsch-Maitland [ID](#)¹⁰¹, G. Quetant [ID](#)⁵⁶, R.P. Quinn [ID](#)¹⁶⁴, G. Rabanal Bolanos [ID](#)⁶¹,
D. Rafanoharana [ID](#)⁵⁴, F. Ragusa [ID](#)^{71a,71b}, J.L. Rainbolt [ID](#)³⁹, J.A. Raine [ID](#)⁵⁶, S. Rajagopalan [ID](#)²⁹,
E. Ramakoti [ID](#)³⁷, K. Ran [ID](#)^{48,14e}, N.P. Rapheeha [ID](#)^{33g}, H. Rasheed [ID](#)^{27b}, V. Raskina [ID](#)¹²⁷,
D.F. Rassloff [ID](#)^{63a}, S. Rave [ID](#)¹⁰⁰, B. Ravina [ID](#)⁵⁵, I. Ravinovich [ID](#)¹⁶⁹, M. Raymond [ID](#)³⁶,
A.L. Read [ID](#)¹²⁵, N.P. Readioff [ID](#)¹³⁹, D.M. Rebuzzi [ID](#)^{73a,73b}, G. Redlinger [ID](#)²⁹, A.S. Reed [ID](#)¹¹⁰,
K. Reeves [ID](#)²⁶, J.A. Reidelsturz [ID](#)¹⁷¹, D. Reikher [ID](#)¹⁵¹, A. Rej [ID](#)¹⁴¹, C. Rembser [ID](#)³⁶,
A. Renardi [ID](#)⁴⁸, M. Renda [ID](#)^{27b}, M.B. Rendel¹¹⁰, F. Renner [ID](#)⁴⁸, A.G. Rennie [ID](#)¹⁶⁰, A.L. Rescia [ID](#)⁴⁸,
S. Resconi [ID](#)^{71a}, M. Ressegotti [ID](#)^{57b,57a}, S. Rettie [ID](#)³⁶, J.G. Reyes Rivera [ID](#)¹⁰⁷, E. Reynolds [ID](#)^{17a},
O.L. Rezanova [ID](#)³⁷, P. Reznicek [ID](#)¹³³, N. Ribaric [ID](#)⁹¹, E. Ricci [ID](#)^{78a,78b}, R. Richter [ID](#)¹¹⁰,
S. Richter [ID](#)^{47a,47b}, E. Richter-Was [ID](#)^{86b}, M. Ridel [ID](#)¹²⁷, S. Ridouani [ID](#)^{35d}, P. Rieck [ID](#)¹¹⁷,
P. Riedler [ID](#)³⁶, E.M. Riefel [ID](#)^{47a,47b}, M. Rijssenbeek [ID](#)¹⁴⁵, A. Rimoldi [ID](#)^{73a,73b}, M. Rimoldi [ID](#)⁴⁸,
L. Rinaldi [ID](#)^{23b,23a}, T.T. Rinn [ID](#)²⁹, M.P. Rinnagel [ID](#)¹⁰⁹, G. Ripellino [ID](#)¹⁶¹, I. Riu [ID](#)¹³,
P. Rivadeneira [ID](#)⁴⁸, J.C. Rivera Vergara [ID](#)¹⁶⁵, F. Rizatdinova [ID](#)¹²¹, E. Rizvi [ID](#)⁹⁴, B.A. Roberts [ID](#)¹⁶⁷,
B.R. Roberts [ID](#)^{17a}, S.H. Robertson [ID](#)^{104,x}, D. Robinson [ID](#)³², C.M. Robles Gajardo^{137f},
M. Robles Manzano [ID](#)¹⁰⁰, A. Robson [ID](#)⁵⁹, A. Rocchi [ID](#)^{76a,76b}, C. Roda [ID](#)^{74a,74b},
S. Rodriguez Bosca [ID](#)^{63a}, Y. Rodriguez Garcia [ID](#)^{22a}, A. Rodriguez Rodriguez [ID](#)⁵⁴,
A.M. Rodríguez Vera [ID](#)^{156b}, S. Roe³⁶, J.T. Roemer [ID](#)¹⁶⁰, A.R. Roepe-Gier [ID](#)¹³⁶, J. Roggel [ID](#)¹⁷¹,
O. Røhne [ID](#)¹²⁵, R.A. Rojas [ID](#)¹⁰³, C.P.A. Roland [ID](#)⁶⁸, J. Roloff [ID](#)²⁹, A. Romaniouk [ID](#)³⁷,
E. Romano [ID](#)^{73a,73b}, M. Romano [ID](#)^{23b}, A.C. Romero Hernandez [ID](#)¹⁶², N. Rompotis [ID](#)⁹²,
L. Roos [ID](#)¹²⁷, S. Rosati [ID](#)^{75a}, B.J. Rosser [ID](#)³⁹, E. Rossi [ID](#)¹²⁶, E. Rossi [ID](#)^{72a,72b}, L.P. Rossi [ID](#)^{57b},
L. Rossini [ID](#)⁵⁴, R. Rosten [ID](#)¹¹⁹, M. Rotaru [ID](#)^{27b}, B. Rottler [ID](#)⁵⁴, C. Rougier [ID](#)^{102,ab},
D. Rousseau [ID](#)⁶⁶, D. Rousso [ID](#)³², A. Roy [ID](#)¹⁶², S. Roy-Garand [ID](#)¹⁵⁵, A. Rozanov [ID](#)¹⁰²,

Y. Rozen [ID](#)¹⁵⁰, X. Ruan [ID](#)^{33g}, A. Rubio Jimenez [ID](#)¹⁶³, A.J. Ruby [ID](#)⁹², V.H. Ruelas Rivera [ID](#)¹⁸,
 T.A. Ruggeri [ID](#)¹, A. Ruggiero [ID](#)¹²⁶, A. Ruiz-Martinez [ID](#)¹⁶³, A. Rummler [ID](#)³⁶, Z. Rurikova [ID](#)⁵⁴,
 N.A. Rusakovich [ID](#)³⁸, H.L. Russell [ID](#)¹⁶⁵, G. Russo [ID](#)^{75a,75b}, J.P. Rutherford [ID](#)⁷,
 S. Rutherford Colmenares [ID](#)³², K. Rybacki⁹¹, M. Rybar [ID](#)¹³³, E.B. Rye [ID](#)¹²⁵, A. Ryzhov [ID](#)⁴⁴,
 J.A. Sabater Iglesias [ID](#)⁵⁶, P. Sabatini [ID](#)¹⁶³, L. Sabetta [ID](#)^{75a,75b}, H.F.W. Sadrozinski [ID](#)¹³⁶,
 F. Safai Tehrani [ID](#)^{75a}, B. Safarzadeh Samani [ID](#)¹⁴⁶, M. Safdari [ID](#)¹⁴³, S. Saha [ID](#)¹⁶⁵, M. Sahinsoy [ID](#)¹¹⁰,
 M. Saimpert [ID](#)¹³⁵, M. Saito [ID](#)¹⁵³, T. Saito [ID](#)¹⁵³, D. Salamani [ID](#)³⁶, A. Salnikov [ID](#)¹⁴³, J. Salt [ID](#)¹⁶³,
 A. Salvador Salas [ID](#)¹³, D. Salvatore [ID](#)^{43b,43a}, F. Salvatore [ID](#)¹⁴⁶, A. Salzburger [ID](#)³⁶, D. Sammel [ID](#)⁵⁴,
 D. Sampsonidis [ID](#)^{152,e}, D. Sampsonidou [ID](#)¹²³, J. Sánchez [ID](#)¹⁶³, A. Sanchez Pineda [ID](#)⁴,
 V. Sanchez Sebastian [ID](#)¹⁶³, H. Sandaker [ID](#)¹²⁵, C.O. Sander [ID](#)⁴⁸, J.A. Sandesara [ID](#)¹⁰³,
 M. Sandhoff [ID](#)¹⁷¹, C. Sandoval [ID](#)^{22b}, D.P.C. Sankey [ID](#)¹³⁴, T. Sano [ID](#)⁸⁸, A. Sansoni [ID](#)⁵³,
 L. Santi [ID](#)^{75a,75b}, C. Santoni [ID](#)⁴⁰, H. Santos [ID](#)^{130a,130b}, S.N. Santpur [ID](#)^{17a}, A. Santra [ID](#)¹⁶⁹,
 K.A. Saoucha [ID](#)^{116b}, J.G. Saraiva [ID](#)^{130a,130d}, J. Sardain [ID](#)⁷, O. Sasaki [ID](#)⁸⁴, K. Sato [ID](#)¹⁵⁷,
 C. Sauer^{63b}, F. Sauerburger [ID](#)⁵⁴, E. Sauvan [ID](#)⁴, P. Savard [ID](#)^{155,ah}, R. Sawada [ID](#)¹⁵³, C. Sawyer [ID](#)¹³⁴,
 L. Sawyer [ID](#)⁹⁷, I. Sayago Galvan [ID](#)¹⁶³, C. Sbarra [ID](#)^{23b}, A. Sbrizzi [ID](#)^{23b,23a}, T. Scanlon [ID](#)⁹⁶,
 J. Schaarschmidt [ID](#)¹³⁸, P. Schacht [ID](#)¹¹⁰, D. Schaefer [ID](#)³⁹, U. Schäfer [ID](#)¹⁰⁰, A.C. Schaffer [ID](#)^{66,44},
 D. Schaile [ID](#)¹⁰⁹, R.D. Schamberger [ID](#)¹⁴⁵, C. Scharf [ID](#)¹⁸, M.M. Schefer [ID](#)¹⁹, V.A. Schegelsky [ID](#)³⁷,
 D. Scheirich [ID](#)¹³³, F. Schenck [ID](#)¹⁸, M. Schernau [ID](#)¹⁶⁰, C. Scheulen [ID](#)⁵⁵, C. Schiavi [ID](#)^{57b,57a},
 E.J. Schioppa [ID](#)^{70a,70b}, M. Schioppa [ID](#)^{43b,43a}, B. Schlag [ID](#)^{143,o}, K.E. Schleicher [ID](#)⁵⁴,
 S. Schlenker [ID](#)³⁶, J. Schmeing [ID](#)¹⁷¹, M.A. Schmidt [ID](#)¹⁷¹, K. Schmieden [ID](#)¹⁰⁰, C. Schmitt [ID](#)¹⁰⁰,
 S. Schmitt [ID](#)⁴⁸, L. Schoeffel [ID](#)¹³⁵, A. Schoening [ID](#)^{63b}, P.G. Scholer [ID](#)⁵⁴, E. Schopf [ID](#)¹²⁶,
 M. Schott [ID](#)¹⁰⁰, J. Schovancova [ID](#)³⁶, S. Schramm [ID](#)⁵⁶, F. Schroeder [ID](#)¹⁷¹, T. Schroer [ID](#)⁵⁶,
 H-C. Schultz-Coulon [ID](#)^{63a}, M. Schumacher [ID](#)⁵⁴, B.A. Schumm [ID](#)¹³⁶, Ph. Schune [ID](#)¹³⁵,
 A.J. Schuy [ID](#)¹³⁸, H.R. Schwartz [ID](#)¹³⁶, A. Schwartzman [ID](#)¹⁴³, T.A. Schwarz [ID](#)¹⁰⁶,
 Ph. Schwemling [ID](#)¹³⁵, R. Schwienhorst [ID](#)¹⁰⁷, A. Sciandra [ID](#)¹³⁶, G. Sciolla [ID](#)²⁶, F. Scuri [ID](#)^{74a},
 C.D. Sebastiani [ID](#)⁹², K. Sedlaczek [ID](#)¹¹⁵, P. Seema [ID](#)¹⁸, S.C. Seidel [ID](#)¹¹², A. Seiden [ID](#)¹³⁶,
 B.D. Seidlitz [ID](#)⁴¹, C. Seitz [ID](#)⁴⁸, J.M. Seixas [ID](#)^{83b}, G. Sekhniaidze [ID](#)^{72a}, S.J. Sekula [ID](#)⁴⁴,
 L. Selem [ID](#)⁶⁰, N. Semprini-Cesari [ID](#)^{23b,23a}, D. Sengupta [ID](#)⁵⁶, V. Senthilkumar [ID](#)¹⁶³, L. Serin [ID](#)⁶⁶,
 L. Serkin [ID](#)^{69a,69b}, M. Sessa [ID](#)^{76a,76b}, H. Severini [ID](#)¹²⁰, F. Sforza [ID](#)^{57b,57a}, A. Sfyrta [ID](#)⁵⁶,
 E. Shabalina [ID](#)⁵⁵, R. Shaheen [ID](#)¹⁴⁴, J.D. Shahinian [ID](#)¹²⁸, D. Shaked Renous [ID](#)¹⁶⁹, L.Y. Shan [ID](#)^{14a},
 M. Shapiro [ID](#)^{17a}, A. Sharma [ID](#)³⁶, A.S. Sharma [ID](#)¹⁶⁴, P. Sharma [ID](#)⁸⁰, S. Sharma [ID](#)⁴⁸,
 P.B. Shatalov [ID](#)³⁷, K. Shaw [ID](#)¹⁴⁶, S.M. Shaw [ID](#)¹⁰¹, A. Shcherbakova [ID](#)³⁷, Q. Shen [ID](#)^{62c,5},
 P. Sherwood [ID](#)⁹⁶, L. Shi [ID](#)⁹⁶, X. Shi [ID](#)^{14a}, C.O. Shimmin [ID](#)¹⁷², J.D. Shinner [ID](#)⁹⁵, I.P.J. Shipsey [ID](#)¹²⁶,
 S. Shirabe [ID](#)^{56,h}, M. Shiyakova [ID](#)^{38,v}, J. Shlomi [ID](#)¹⁶⁹, M.J. Shochet [ID](#)³⁹, J. Shojaii [ID](#)¹⁰⁵,
 D.R. Shope [ID](#)¹²⁵, B. Shrestha [ID](#)¹²⁰, S. Shrestha [ID](#)^{119,ak}, E.M. Shrif [ID](#)^{33g}, M.J. Shroff [ID](#)¹⁶⁵,
 P. Sicho [ID](#)¹³¹, A.M. Sickles [ID](#)¹⁶², E. Sideras Haddad [ID](#)^{33g}, A. Sidoti [ID](#)^{23b}, F. Siegert [ID](#)⁵⁰,
 Dj. Sijacki [ID](#)¹⁵, R. Sikora [ID](#)^{86a}, F. Sili [ID](#)⁹⁰, J.M. Silva [ID](#)²⁰, M.V. Silva Oliveira [ID](#)²⁹,
 S.B. Silverstein [ID](#)^{47a}, S. Simion⁶⁶, R. Simoniello [ID](#)³⁶, E.L. Simpson [ID](#)⁵⁹, H. Simpson [ID](#)¹⁴⁶,
 L.R. Simpson [ID](#)¹⁰⁶, N.D. Simpson⁹⁸, S. Simsek [ID](#)⁸², S. Sindhu [ID](#)⁵⁵, P. Sinervo [ID](#)¹⁵⁵, S. Singh [ID](#)¹⁵⁵,
 S. Sinha [ID](#)⁴⁸, S. Sinha [ID](#)¹⁰¹, M. Sioli [ID](#)^{23b,23a}, I. Siral [ID](#)³⁶, E. Sitnikova [ID](#)⁴⁸, S.Yu. Sivoklov [ID](#)^{37,*},
 J. Sjölin [ID](#)^{47a,47b}, A. Skaf [ID](#)⁵⁵, E. Skorda [ID](#)²⁰, P. Skubic [ID](#)¹²⁰, M. Slawinska [ID](#)⁸⁷, V. Smakhtin¹⁶⁹,
 B.H. Smart [ID](#)¹³⁴, J. Smiesko [ID](#)³⁶, S.Yu. Smirnov [ID](#)³⁷, Y. Smirnov [ID](#)³⁷, L.N. Smirnova [ID](#)^{37,a},
 O. Smirnova [ID](#)⁹⁸, A.C. Smith [ID](#)⁴¹, E.A. Smith [ID](#)³⁹, H.A. Smith [ID](#)¹²⁶, J.L. Smith [ID](#)⁹², R. Smith¹⁴³,

M. Smizanska [ID](#)⁹¹, K. Smolek [ID](#)¹³², A.A. Snesev [ID](#)³⁷, S.R. Snider [ID](#)¹⁵⁵, H.L. Snoek [ID](#)¹¹⁴, S. Snyder [ID](#)²⁹, R. Sobie [ID](#)^{165,x}, A. Soffer [ID](#)¹⁵¹, C.A. Solans Sanchez [ID](#)³⁶, E.Yu. Soldatov [ID](#)³⁷, U. Soldevila [ID](#)¹⁶³, A.A. Solodkov [ID](#)³⁷, S. Solomon [ID](#)²⁶, A. Soloshenko [ID](#)³⁸, K. Solovieva [ID](#)⁵⁴, O.V. Solovyanov [ID](#)⁴⁰, V. Solovyev [ID](#)³⁷, P. Sommer [ID](#)³⁶, A. Sonay [ID](#)¹³, W.Y. Song [ID](#)^{156b}, J.M. Sonneveld [ID](#)¹¹⁴, A. Sopczak [ID](#)¹³², A.L. Soppio [ID](#)⁹⁶, F. Sopkova [ID](#)^{28b}, V. Sothilingam [ID](#)^{63a}, S. Sottocornola [ID](#)⁶⁸, R. Soualah [ID](#)^{116b}, Z. Soumami [ID](#)^{35e}, D. South [ID](#)⁴⁸, N. Soybelman [ID](#)¹⁶⁹, S. Spagnolo [ID](#)^{70a,70b}, M. Spalla [ID](#)¹¹⁰, D. Sperlich [ID](#)⁵⁴, G. Spigo [ID](#)³⁶, S. Spinali [ID](#)⁹¹, D.P. Spiteri [ID](#)⁵⁹, M. Spousta [ID](#)¹³³, E.J. Staats [ID](#)³⁴, A. Stabile [ID](#)^{71a,71b}, R. Stamen [ID](#)^{63a}, A. Stampekis [ID](#)²⁰, M. Standke [ID](#)²⁴, E. Stanecka [ID](#)⁸⁷, M.V. Stange [ID](#)⁵⁰, B. Stanislaus [ID](#)^{17a}, M.M. Stanitzki [ID](#)⁴⁸, B. Stapf [ID](#)⁴⁸, E.A. Starchenko [ID](#)³⁷, G.H. Stark [ID](#)¹³⁶, J. Stark [ID](#)^{102,ab}, D.M. Starke [ID](#)^{156b}, P. Staroba [ID](#)¹³¹, P. Starovoitov [ID](#)^{63a}, S. Starz [ID](#)¹⁰⁴, R. Staszewski [ID](#)⁸⁷, G. Stavropoulos [ID](#)⁴⁶, J. Steentoft [ID](#)¹⁶¹, P. Steinberg [ID](#)²⁹, B. Stelzer [ID](#)^{142,156a}, H.J. Stelzer [ID](#)¹²⁹, O. Stelzer-Chilton [ID](#)^{156a}, H. Stenzel [ID](#)⁵⁸, T.J. Stevenson [ID](#)¹⁴⁶, G.A. Stewart [ID](#)³⁶, J.R. Stewart [ID](#)¹²¹, M.C. Stockton [ID](#)³⁶, G. Stoicea [ID](#)^{27b}, M. Stolarski [ID](#)^{130a}, S. Stonjek [ID](#)¹¹⁰, A. Straessner [ID](#)⁵⁰, J. Strandberg [ID](#)¹⁴⁴, S. Strandberg [ID](#)^{47a,47b}, M. Stratmann [ID](#)¹⁷¹, M. Strauss [ID](#)¹²⁰, T. Strebler [ID](#)¹⁰², P. Strizenec [ID](#)^{28b}, R. Strohmer [ID](#)¹⁶⁶, D.M. Strom [ID](#)¹²³, L.R. Strom [ID](#)⁴⁸, R. Stroynowski [ID](#)⁴⁴, A. Strubig [ID](#)^{47a,47b}, S.A. Stucci [ID](#)²⁹, B. Stugu [ID](#)¹⁶, J. Stupak [ID](#)¹²⁰, N.A. Styles [ID](#)⁴⁸, D. Su [ID](#)¹⁴³, S. Su [ID](#)^{62a}, W. Su [ID](#)^{62d}, X. Su [ID](#)^{62a,66}, K. Sugizaki [ID](#)¹⁵³, V.V. Sulim [ID](#)³⁷, M.J. Sullivan [ID](#)⁹², D.M.S. Sultan [ID](#)^{78a,78b}, L. Sultanaliev [ID](#)³⁷, S. Sultansoy [ID](#)^{3b}, T. Sumida [ID](#)⁸⁸, S. Sun [ID](#)¹⁰⁶, S. Sun [ID](#)¹⁷⁰, O. Sunneborn Gudnadottir [ID](#)¹⁶¹, N. Sur [ID](#)¹⁰², M.R. Sutton [ID](#)¹⁴⁶, H. Suzuki [ID](#)¹⁵⁷, M. Svatos [ID](#)¹³¹, M. Swiatlowski [ID](#)^{156a}, T. Swirski [ID](#)¹⁶⁶, I. Sykora [ID](#)^{28a}, M. Sykora [ID](#)¹³³, T. Sykora [ID](#)¹³³, D. Ta [ID](#)¹⁰⁰, K. Tackmann [ID](#)^{48,u}, A. Taffard [ID](#)¹⁶⁰, R. Tafirout [ID](#)^{156a}, J.S. Tafoya Vargas [ID](#)⁶⁶, E.P. Takeva [ID](#)⁵², Y. Takubo [ID](#)⁸⁴, M. Talby [ID](#)¹⁰², A.A. Talyshiev [ID](#)³⁷, K.C. Tam [ID](#)^{64b}, N.M. Tamir [ID](#)¹⁵¹, A. Tanaka [ID](#)¹⁵³, J. Tanaka [ID](#)¹⁵³, R. Tanaka [ID](#)⁶⁶, M. Tanasini [ID](#)^{57b,57a}, Z. Tao [ID](#)¹⁶⁴, S. Tapia Araya [ID](#)^{137f}, S. Tapprogge [ID](#)¹⁰⁰, A. Tarek Abouelfadl Mohamed [ID](#)¹⁰⁷, S. Tarem [ID](#)¹⁵⁰, K. Tariq [ID](#)^{14a}, G. Tarna [ID](#)^{102,27b}, G.F. Tartarelli [ID](#)^{71a}, P. Tas [ID](#)¹³³, M. Tasevsky [ID](#)¹³¹, E. Tassi [ID](#)^{43b,43a}, A.C. Tate [ID](#)¹⁶², G. Tateno [ID](#)¹⁵³, Y. Tayalati [ID](#)^{35e,w}, G.N. Taylor [ID](#)¹⁰⁵, W. Taylor [ID](#)^{156b}, H. Teagle [ID](#)⁹², A.S. Tee [ID](#)¹⁷⁰, R. Teixeira De Lima [ID](#)¹⁴³, P. Teixeira-Dias [ID](#)⁹⁵, J.J. Teoh [ID](#)¹⁵⁵, K. Terashi [ID](#)¹⁵³, J. Terron [ID](#)⁹⁹, S. Terzo [ID](#)¹³, M. Testa [ID](#)⁵³, R.J. Teuscher [ID](#)^{155,x}, A. Thaler [ID](#)⁷⁹, O. Theiner [ID](#)⁵⁶, N. Themistokleous [ID](#)⁵², T. Theveneaux-Pelzer [ID](#)¹⁰², O. Thielmann [ID](#)¹⁷¹, D.W. Thomas [ID](#)⁹⁵, J.P. Thomas [ID](#)²⁰, E.A. Thompson [ID](#)^{17a}, P.D. Thompson [ID](#)²⁰, E. Thomson [ID](#)¹²⁸, Y. Tian [ID](#)⁵⁵, V. Tikhomirov [ID](#)^{37,a}, Yu.A. Tikhonov [ID](#)³⁷, S. Timoshenko [ID](#)³⁷, D. Timoshyn [ID](#)¹³³, E.X.L. Ting [ID](#)¹, P. Tipton [ID](#)¹⁷², S.H. Tlou [ID](#)^{33g}, A. Thourji [ID](#)⁴⁰, K. Todome [ID](#)¹⁵⁴, S. Todorova-Nova [ID](#)¹³³, S. Todt [ID](#)⁵⁰, M. Togawa [ID](#)⁸⁴, J. Tojo [ID](#)⁸⁹, S. Tokar [ID](#)^{28a}, K. Tokushuku [ID](#)⁸⁴, O. Toldaiev [ID](#)⁶⁸, R. Tombs [ID](#)³², M. Tomoto [ID](#)^{84,111}, L. Tompkins [ID](#)^{143,o}, K.W. Topolnicki [ID](#)^{86b}, E. Torrence [ID](#)¹²³, H. Torres [ID](#)^{102,ab}, E. Torro Pastor [ID](#)¹⁶³, M. Toscani [ID](#)³⁰, C. Tosciri [ID](#)³⁹, M. Tost [ID](#)¹¹, D.R. Tovey [ID](#)¹³⁹, A. Traeet [ID](#)¹⁶, I.S. Trandafir [ID](#)^{27b}, T. Trefzger [ID](#)¹⁶⁶, A. Tricoli [ID](#)²⁹, I.M. Trigger [ID](#)^{156a}, S. Trincaz-Duvoid [ID](#)¹²⁷, D.A. Trischuk [ID](#)²⁶, B. Trocme [ID](#)⁶⁰, C. Troncon [ID](#)^{71a}, L. Truong [ID](#)^{33c}, M. Trzebinski [ID](#)⁸⁷, A. Trzupek [ID](#)⁸⁷, F. Tsai [ID](#)¹⁴⁵, M. Tsai [ID](#)¹⁰⁶, A. Tsiamis [ID](#)^{152,e}, P.V. Tsiarshka [ID](#)³⁷, S. Tsigaridas [ID](#)^{156a}, A. Tsirigotis [ID](#)^{152,s}, V. Tsiskaridze [ID](#)¹⁵⁵, E.G. Tskhadadze [ID](#)^{149a}, M. Tsopoulou [ID](#)^{152,e}, Y. Tsujikawa [ID](#)⁸⁸, I.I. Tsukerman [ID](#)³⁷, V. Tsulaia [ID](#)^{17a}, S. Tsuno [ID](#)⁸⁴, O. Tsur [ID](#)¹⁵⁰, K. Tsurii [ID](#)¹¹⁸, D. Tsybychev [ID](#)¹⁴⁵, Y. Tu [ID](#)^{64b}, A. Tudorache [ID](#)^{27b}, V. Tudorache [ID](#)^{27b}, A.N. Tuna [ID](#)³⁶, S. Turchikhin [ID](#)^{57b,57a}, I. Turk Cakir [ID](#)^{3a}, R. Turra [ID](#)^{71a}, T. Turtuvshin [ID](#)^{38,y},

P.M. Tuts [ID](#)⁴¹, S. Tzamarias [ID](#)^{152,e}, P. Tzannis [ID](#)¹⁰, E. Tzovara [ID](#)¹⁰⁰, F. Ukegawa [ID](#)¹⁵⁷,
 P.A. Ulloa Poblete [ID](#)^{137c,137b}, E.N. Umaka [ID](#)²⁹, G. Unal [ID](#)³⁶, M. Unal [ID](#)¹¹, A. Undrus [ID](#)²⁹,
 G. Unel [ID](#)¹⁶⁰, J. Urban [ID](#)^{28b}, P. Urquijo [ID](#)¹⁰⁵, G. Usai [ID](#)⁸, R. Ushioda [ID](#)¹⁵⁴, M. Usman [ID](#)¹⁰⁸,
 Z. Uysal [ID](#)^{21b}, L. Vacavant [ID](#)¹⁰², V. Vacek [ID](#)¹³², B. Vachon [ID](#)¹⁰⁴, K.O.H. Vadla [ID](#)¹²⁵,
 T. Vafeiadis [ID](#)³⁶, A. Vaitkus [ID](#)⁹⁶, C. Valderanis [ID](#)¹⁰⁹, E. Valdes Santurio [ID](#)^{47a,47b}, M. Valente [ID](#)^{156a},
 S. Valentinetti [ID](#)^{23b,23a}, A. Valero [ID](#)¹⁶³, E. Valiente Moreno [ID](#)¹⁶³, A. Vallier [ID](#)^{102,ab},
 J.A. Valls Ferrer [ID](#)¹⁶³, D.R. Van Arneman [ID](#)¹¹⁴, T.R. Van Daalen [ID](#)¹³⁸, A. Van Der Graaf [ID](#)⁴⁹,
 P. Van Gemmeren [ID](#)⁶, M. Van Rijnbach [ID](#)^{125,36}, S. Van Stroud [ID](#)⁹⁶, I. Van Vulpen [ID](#)¹¹⁴,
 M. Vanadia [ID](#)^{76a,76b}, W. Vandelli [ID](#)³⁶, M. Vandenbroucke [ID](#)¹³⁵, E.R. Vandewall [ID](#)¹²¹,
 D. Vannicola [ID](#)¹⁵¹, L. Vannoli [ID](#)^{57b,57a}, R. Vari [ID](#)^{75a}, E.W. Varnes [ID](#)⁷, C. Varni [ID](#)^{17b}, T. Varol [ID](#)¹⁴⁸,
 D. Varouchas [ID](#)⁶⁶, L. Varriale [ID](#)¹⁶³, K.E. Varvell [ID](#)¹⁴⁷, M.E. Vasile [ID](#)^{27b}, L. Vaslin [ID](#)⁴⁰,
 G.A. Vasquez [ID](#)¹⁶⁵, A. Vasyukov [ID](#)³⁸, F. Vazeille [ID](#)⁴⁰, T. Vazquez Schroeder [ID](#)³⁶, J. Veatch [ID](#)³¹,
 V. Vecchio [ID](#)¹⁰¹, M.J. Veen [ID](#)¹⁰³, I. Veliscek [ID](#)¹²⁶, L.M. Veloce [ID](#)¹⁵⁵, F. Veloso [ID](#)^{130a,130c},
 S. Veneziano [ID](#)^{75a}, A. Ventura [ID](#)^{70a,70b}, S. Ventura Gonzalez [ID](#)¹³⁵, A. Verbitskiy [ID](#)¹¹⁰,
 M. Verducci [ID](#)^{74a,74b}, C. Vergis [ID](#)²⁴, M. Verissimo De Araujo [ID](#)^{83b}, W. Verkerke [ID](#)¹¹⁴,
 J.C. Vermeulen [ID](#)¹¹⁴, C. Vernieri [ID](#)¹⁴³, M. Vessella [ID](#)¹⁰³, M.C. Vetterli [ID](#)^{142,ah},
 A. Vgenopoulos [ID](#)^{152,e}, N. Viaux Maira [ID](#)^{137f}, T. Vickey [ID](#)¹³⁹, O.E. Vickey Boeriu [ID](#)¹³⁹,
 G.H.A. Viehhauser [ID](#)¹²⁶, L. Vigani [ID](#)^{63b}, M. Villa [ID](#)^{23b,23a}, M. Villaplana Perez [ID](#)¹⁶³,
 E.M. Villhauer [ID](#)⁵², E. Vilucchi [ID](#)⁵³, M.G. Vincter [ID](#)³⁴, G.S. Virdee [ID](#)²⁰, A. Vishwakarma [ID](#)⁵²,
 A. Visibile [ID](#)¹¹⁴, C. Vittori [ID](#)³⁶, I. Vivarelli [ID](#)¹⁴⁶, V. Vladimirov [ID](#)¹⁶⁷, E. Voevodina [ID](#)¹¹⁰, F. Vogel [ID](#)¹⁰⁹,
 P. Vokac [ID](#)¹³², Yu. Volkotrub [ID](#)^{86a}, J. Von Ahnen [ID](#)⁴⁸, E. Von Toerne [ID](#)²⁴, B. Vormwald [ID](#)³⁶,
 V. Vorobel [ID](#)¹³³, K. Vorobev [ID](#)³⁷, M. Vos [ID](#)¹⁶³, K. Voss [ID](#)¹⁴¹, J.H. Vosseveld [ID](#)⁹², M. Vozak [ID](#)¹¹⁴,
 L. Vozdecky [ID](#)⁹⁴, N. Vranjes [ID](#)¹⁵, M. Vranjes Milosavljevic [ID](#)¹⁵, M. Vreeswijk [ID](#)¹¹⁴,
 R. Vuillermet [ID](#)³⁶, O. Vujanovic [ID](#)¹⁰⁰, I. Vukotic [ID](#)³⁹, S. Wada [ID](#)¹⁵⁷, C. Wagner [ID](#)¹⁰³,
 J.M. Wagner [ID](#)^{17a}, W. Wagner [ID](#)¹⁷¹, S. Wahdan [ID](#)¹⁷¹, H. Wahlberg [ID](#)⁹⁰, M. Wakida [ID](#)¹¹¹,
 J. Walder [ID](#)¹³⁴, R. Walker [ID](#)¹⁰⁹, W. Walkowiak [ID](#)¹⁴¹, A. Wall [ID](#)¹²⁸, T. Wamorkar [ID](#)⁶,
 A.Z. Wang [ID](#)¹⁷⁰, C. Wang [ID](#)¹⁰⁰, C. Wang [ID](#)^{62c}, H. Wang [ID](#)^{17a}, J. Wang [ID](#)^{64a}, R.-J. Wang [ID](#)¹⁰⁰,
 R. Wang [ID](#)⁶¹, R. Wang [ID](#)⁶, S.M. Wang [ID](#)¹⁴⁸, S. Wang [ID](#)^{62b}, T. Wang [ID](#)^{62a}, W.T. Wang [ID](#)⁸⁰,
 W. Wang [ID](#)^{14a}, X. Wang [ID](#)^{14c}, X. Wang [ID](#)¹⁶², X. Wang [ID](#)^{62c}, Y. Wang [ID](#)^{62d}, Y. Wang [ID](#)^{14c},
 Z. Wang [ID](#)¹⁰⁶, Z. Wang [ID](#)^{62d,51,62c}, Z. Wang [ID](#)¹⁰⁶, A. Warburton [ID](#)¹⁰⁴, R.J. Ward [ID](#)²⁰,
 N. Warrack [ID](#)⁵⁹, A.T. Watson [ID](#)²⁰, H. Watson [ID](#)⁵⁹, M.F. Watson [ID](#)²⁰, E. Watton [ID](#)^{59,134},
 G. Watts [ID](#)¹³⁸, B.M. Waugh [ID](#)⁹⁶, C. Weber [ID](#)²⁹, H.A. Weber [ID](#)¹⁸, M.S. Weber [ID](#)¹⁹,
 S.M. Weber [ID](#)^{63a}, C. Wei [ID](#)^{62a}, Y. Wei [ID](#)¹²⁶, A.R. Weidberg [ID](#)¹²⁶, E.J. Weik [ID](#)¹¹⁷,
 J. Weingarten [ID](#)⁴⁹, M. Weirich [ID](#)¹⁰⁰, C. Weiser [ID](#)⁵⁴, C.J. Wells [ID](#)⁴⁸, T. Wenaus [ID](#)²⁹,
 B. Wendland [ID](#)⁴⁹, T. Wengler [ID](#)³⁶, N.S. Wenke [ID](#)¹¹⁰, N. Wermes [ID](#)²⁴, M. Wessels [ID](#)^{63a},
 A.M. Wharton [ID](#)⁹¹, A.S. White [ID](#)⁶¹, A. White [ID](#)⁸, M.J. White [ID](#)¹, D. Whiteson [ID](#)¹⁶⁰,
 L. Wickremasinghe [ID](#)¹²⁴, W. Wiedenmann [ID](#)¹⁷⁰, C. Wiel [ID](#)⁵⁰, M. Wielers [ID](#)¹³⁴, C. Wiglesworth [ID](#)⁴²,
 D.J. Wilbern [ID](#)¹²⁰, H.G. Wilkens [ID](#)³⁶, D.M. Williams [ID](#)⁴¹, H.H. Williams [ID](#)¹²⁸, S. Williams [ID](#)³²,
 S. Willocq [ID](#)¹⁰³, B.J. Wilson [ID](#)¹⁰¹, P.J. Windischhofer [ID](#)³⁹, F.I. Winkel [ID](#)³⁰, F. Winklmeier [ID](#)¹²³,
 B.T. Winter [ID](#)⁵⁴, J.K. Winter [ID](#)¹⁰¹, M. Wittgen [ID](#)¹⁴³, M. Wobisch [ID](#)⁹⁷, Z. Wolffs [ID](#)¹¹⁴, J. Wollrath [ID](#)¹⁶⁰,
 M.W. Wolter [ID](#)⁸⁷, H. Wolters [ID](#)^{130a,130c}, A.F. Wongel [ID](#)⁴⁸, S.D. Worm [ID](#)⁴⁸, B.K. Wosiek [ID](#)⁸⁷,
 K.W. Woźniak [ID](#)⁸⁷, S. Wozniowski [ID](#)⁵⁵, K. Wraight [ID](#)⁵⁹, C. Wu [ID](#)²⁰, J. Wu [ID](#)^{14a,14e}, M. Wu [ID](#)^{64a},
 M. Wu [ID](#)¹¹³, S.L. Wu [ID](#)¹⁷⁰, X. Wu [ID](#)⁵⁶, Y. Wu [ID](#)^{62a}, Z. Wu [ID](#)¹³⁵, J. Wuerzinger [ID](#)^{110,af},

T.R. Wyatt ¹⁰¹, B.M. Wynne ⁵², S. Xella ⁴², L. Xia ^{14c}, M. Xia ^{14b}, J. Xiang ^{64c},
M. Xie ^{62a}, X. Xie ^{62a}, S. Xin ^{14a,14e}, J. Xiong ^{17a}, D. Xu ^{14a}, H. Xu ^{62a}, L. Xu ^{62a},
R. Xu ¹²⁸, T. Xu ¹⁰⁶, Y. Xu ^{14b}, Z. Xu ⁵², Z. Xu ^{14a}, B. Yabsley ¹⁴⁷, S. Yacoob ^{33a},
Y. Yamaguchi ¹⁵⁴, E. Yamashita ¹⁵³, H. Yamauchi ¹⁵⁷, T. Yamazaki ^{17a}, Y. Yamazaki ⁸⁵,
J. Yan ^{62c}, S. Yan ¹²⁶, Z. Yan ²⁵, H.J. Yang ^{62c,62d}, H.T. Yang ^{62a}, S. Yang ^{62a},
T. Yang ^{64c}, X. Yang ^{62a}, X. Yang ^{14a}, Y. Yang ⁴⁴, Y. Yang ^{62a}, Z. Yang ^{62a},
W-M. Yao ^{17a}, Y.C. Yap ⁴⁸, H. Ye ^{14c}, H. Ye ⁵⁵, J. Ye ^{14a}, S. Ye ²⁹, X. Ye ^{62a},
Y. Yeh ⁹⁶, I. Yeletsikh ³⁸, B.K. Yeo ^{17b}, M.R. Yexley ⁹⁶, P. Yin ⁴¹, K. Yorita ¹⁶⁸,
S. Younas ^{27b}, C.J.S. Young ³⁶, C. Young ¹⁴³, C. Yu ^{14a,14e}, Y. Yu ^{62a}, M. Yuan ¹⁰⁶,
R. Yuan ^{62b,k}, L. Yue ⁹⁶, M. Zaazoua ^{62a}, B. Zabinski ⁸⁷, E. Zaid ⁵², T. Zakareishvili ^{149b},
N. Zakharchuk ³⁴, S. Zambito ⁵⁶, J.A. Zamora Saa ^{137d,137b}, J. Zang ¹⁵³, D. Zanzi ⁵⁴,
O. Zaplatilek ¹³², C. Zeitnitz ¹⁷¹, H. Zeng ^{14a}, J.C. Zeng ¹⁶², D.T. Zenger Jr ²⁶,
O. Zenin ³⁷, T. Ženiš ^{28a}, S. Zenz ⁹⁴, S. Zerradi ^{35a}, D. Zerwas ⁶⁶, M. Zhai ^{14a,14e},
B. Zhang ^{14c}, D.F. Zhang ¹³⁹, J. Zhang ^{62b}, J. Zhang ⁶, K. Zhang ^{14a,14e}, L. Zhang ^{14c},
P. Zhang ^{14a,14e}, R. Zhang ¹⁷⁰, S. Zhang ¹⁰⁶, T. Zhang ¹⁵³, X. Zhang ^{62c}, X. Zhang ^{62b},
Y. Zhang ^{62c,5}, Y. Zhang ⁹⁶, Z. Zhang ^{17a}, Z. Zhang ⁶⁶, H. Zhao ¹³⁸, P. Zhao ⁵¹,
T. Zhao ^{62b}, Y. Zhao ¹³⁶, Z. Zhao ^{62a}, A. Zhemchugov ³⁸, J. Zheng ^{14c}, K. Zheng ¹⁶²,
X. Zheng ^{62a}, Z. Zheng ¹⁴³, D. Zhong ¹⁶², B. Zhou ¹⁰⁶, H. Zhou ⁷, N. Zhou ^{62c}, Y. Zhou ⁷,
C.G. Zhu ^{62b}, J. Zhu ¹⁰⁶, Y. Zhu ^{62c}, Y. Zhu ^{62a}, X. Zhuang ^{14a}, K. Zhukov ³⁷,
V. Zhulanov ³⁷, N.I. Zimine ³⁸, J. Zinsser ^{63b}, M. Ziolkowski ¹⁴¹, L. Živković ¹⁵,
A. Zoccoli ^{23b,23a}, K. Zoch ⁵⁶, T.G. Zorbas ¹³⁹, O. Zormpa ⁴⁶, W. Zou ⁴¹, L. Zwalinski ³⁶

¹ Department of Physics, University of Adelaide, Adelaide; Australia

² Department of Physics, University of Alberta, Edmonton AB; Canada

³ ^(a) Department of Physics, Ankara University, Ankara; ^(b) Division of Physics, TOBB University of Economics and Technology, Ankara; Türkiye

⁴ LAPP, Université Savoie Mont Blanc, CNRS/IN2P3, Annecy; France

⁵ APC, Université Paris Cité, CNRS/IN2P3, Paris; France

⁶ High Energy Physics Division, Argonne National Laboratory, Argonne IL; United States of America

⁷ Department of Physics, University of Arizona, Tucson AZ; United States of America

⁸ Department of Physics, University of Texas at Arlington, Arlington TX; United States of America

⁹ Physics Department, National and Kapodistrian University of Athens, Athens; Greece

¹⁰ Physics Department, National Technical University of Athens, Zografou; Greece

¹¹ Department of Physics, University of Texas at Austin, Austin TX; United States of America

¹² Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan

¹³ Institut de Física d'Altes Energies (IFAE), Barcelona Institute of Science and Technology, Barcelona; Spain

¹⁴ ^(a) Institute of High Energy Physics, Chinese Academy of Sciences, Beijing; ^(b) Physics Department, Tsinghua University, Beijing; ^(c) Department of Physics, Nanjing University, Nanjing; ^(d) School of Science, Shenzhen Campus of Sun Yat-sen University; ^(e) University of Chinese Academy of Science (UCAS), Beijing; China

¹⁵ Institute of Physics, University of Belgrade, Belgrade; Serbia

¹⁶ Department for Physics and Technology, University of Bergen, Bergen; Norway

¹⁷ ^(a) Physics Division, Lawrence Berkeley National Laboratory, Berkeley CA; ^(b) University of California, Berkeley CA; United States of America

¹⁸ Institut für Physik, Humboldt Universität zu Berlin, Berlin; Germany

¹⁹ Albert Einstein Center for Fundamental Physics and Laboratory for High Energy Physics, University of Bern, Bern; Switzerland

- ²⁰ *School of Physics and Astronomy, University of Birmingham, Birmingham; United Kingdom*
- ²¹ ^(a) *Department of Physics, Bogazici University, Istanbul;* ^(b) *Department of Physics Engineering, Gaziantep University, Gaziantep;* ^(c) *Department of Physics, Istanbul University, Istanbul; Türkiye*
- ²² ^(a) *Facultad de Ciencias y Centro de Investigaciones, Universidad Antonio Nariño, Bogotá;* ^(b) *Departamento de Física, Universidad Nacional de Colombia, Bogotá; Colombia*
- ²³ ^(a) *Dipartimento di Fisica e Astronomia A. Righi, Università di Bologna, Bologna;* ^(b) *INFN Sezione di Bologna; Italy*
- ²⁴ *Physikalisches Institut, Universität Bonn, Bonn; Germany*
- ²⁵ *Department of Physics, Boston University, Boston MA; United States of America*
- ²⁶ *Department of Physics, Brandeis University, Waltham MA; United States of America*
- ²⁷ ^(a) *Transilvania University of Brasov, Brasov;* ^(b) *Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest;* ^(c) *Department of Physics, Alexandru Ioan Cuza University of Iasi, Iasi;* ^(d) *National Institute for Research and Development of Isotopic and Molecular Technologies, Physics Department, Cluj-Napoca;* ^(e) *University Politehnica Bucharest, Bucharest;* ^(f) *West University in Timisoara, Timisoara;* ^(g) *Faculty of Physics, University of Bucharest, Bucharest; Romania*
- ²⁸ ^(a) *Faculty of Mathematics, Physics and Informatics, Comenius University, Bratislava;* ^(b) *Department of Subnuclear Physics, Institute of Experimental Physics of the Slovak Academy of Sciences, Kosice; Slovak Republic*
- ²⁹ *Physics Department, Brookhaven National Laboratory, Upton NY; United States of America*
- ³⁰ *Universidad de Buenos Aires, Facultad de Ciencias Exactas y Naturales, Departamento de Física, y CONICET, Instituto de Física de Buenos Aires (IFIBA), Buenos Aires; Argentina*
- ³¹ *California State University, CA; United States of America*
- ³² *Cavendish Laboratory, University of Cambridge, Cambridge; United Kingdom*
- ³³ ^(a) *Department of Physics, University of Cape Town, Cape Town;* ^(b) *iThemba Labs, Western Cape;* ^(c) *Department of Mechanical Engineering Science, University of Johannesburg, Johannesburg;* ^(d) *National Institute of Physics, University of the Philippines Diliman (Philippines);* ^(e) *University of South Africa, Department of Physics, Pretoria;* ^(f) *University of Zululand, KwaDlangezwa;* ^(g) *School of Physics, University of the Witwatersrand, Johannesburg; South Africa*
- ³⁴ *Department of Physics, Carleton University, Ottawa ON; Canada*
- ³⁵ ^(a) *Faculté des Sciences Ain Chock, Réseau Universitaire de Physique des Hautes Energies — Université Hassan II, Casablanca;* ^(b) *Faculté des Sciences, Université Ibn-Tofail, Kénitra;* ^(c) *Faculté des Sciences Semlalia, Université Cadi Ayyad, LPHEA-Marrakech;* ^(d) *LPMR, Faculté des Sciences, Université Mohamed Premier, Oujda;* ^(e) *Faculté des sciences, Université Mohammed V, Rabat;* ^(f) *Institute of Applied Physics, Mohammed VI Polytechnic University, Ben Guerir; Morocco*
- ³⁶ *CERN, Geneva; Switzerland*
- ³⁷ *Affiliated with an institute covered by a cooperation agreement with CERN*
- ³⁸ *Affiliated with an international laboratory covered by a cooperation agreement with CERN*
- ³⁹ *Enrico Fermi Institute, University of Chicago, Chicago IL; United States of America*
- ⁴⁰ *LPC, Université Clermont Auvergne, CNRS/IN2P3, Clermont-Ferrand; France*
- ⁴¹ *Nevis Laboratory, Columbia University, Irvington NY; United States of America*
- ⁴² *Niels Bohr Institute, University of Copenhagen, Copenhagen; Denmark*
- ⁴³ ^(a) *Dipartimento di Fisica, Università della Calabria, Rende;* ^(b) *INFN Gruppo Collegato di Cosenza, Laboratori Nazionali di Frascati; Italy*
- ⁴⁴ *Physics Department, Southern Methodist University, Dallas TX; United States of America*
- ⁴⁵ *Physics Department, University of Texas at Dallas, Richardson TX; United States of America*
- ⁴⁶ *National Centre for Scientific Research “Demokritos”, Agia Paraskevi; Greece*
- ⁴⁷ ^(a) *Department of Physics, Stockholm University;* ^(b) *Oskar Klein Centre, Stockholm; Sweden*
- ⁴⁸ *Deutsches Elektronen-Synchrotron DESY, Hamburg and Zeuthen; Germany*
- ⁴⁹ *Fakultät Physik, Technische Universität Dortmund, Dortmund; Germany*
- ⁵⁰ *Institut für Kern- und Teilchenphysik, Technische Universität Dresden, Dresden; Germany*
- ⁵¹ *Department of Physics, Duke University, Durham NC; United States of America*
- ⁵² *SUPA — School of Physics and Astronomy, University of Edinburgh, Edinburgh; United Kingdom*

- ⁵³ INFN e Laboratori Nazionali di Frascati, Frascati; Italy
- ⁵⁴ Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, Freiburg; Germany
- ⁵⁵ II. Physikalisches Institut, Georg-August-Universität Göttingen, Göttingen; Germany
- ⁵⁶ Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland
- ⁵⁷ ^(a) Dipartimento di Fisica, Università di Genova, Genova; ^(b) INFN Sezione di Genova; Italy
- ⁵⁸ II. Physikalisches Institut, Justus-Liebig-Universität Giessen, Giessen; Germany
- ⁵⁹ SUPA — School of Physics and Astronomy, University of Glasgow, Glasgow; United Kingdom
- ⁶⁰ LPSC, Université Grenoble Alpes, CNRS/IN2P3, Grenoble INP, Grenoble; France
- ⁶¹ Laboratory for Particle Physics and Cosmology, Harvard University, Cambridge MA; United States of America
- ⁶² ^(a) Department of Modern Physics and State Key Laboratory of Particle Detection and Electronics, University of Science and Technology of China, Hefei; ^(b) Institute of Frontier and Interdisciplinary Science and Key Laboratory of Particle Physics and Particle Irradiation (MOE), Shandong University, Qingdao; ^(c) School of Physics and Astronomy, Shanghai Jiao Tong University, Key Laboratory for Particle Astrophysics and Cosmology (MOE), SKLPPC, Shanghai; ^(d) Tsung-Dao Lee Institute, Shanghai; China
- ⁶³ ^(a) Kirchhoff-Institut für Physik, Ruprecht-Karls-Universität Heidelberg, Heidelberg; ^(b) Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg; Germany
- ⁶⁴ ^(a) Department of Physics, Chinese University of Hong Kong, Shatin, N.T., Hong Kong; ^(b) Department of Physics, University of Hong Kong, Hong Kong; ^(c) Department of Physics and Institute for Advanced Study, Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong; China
- ⁶⁵ Department of Physics, National Tsing Hua University, Hsinchu; Taiwan
- ⁶⁶ IJCLab, Université Paris-Saclay, CNRS/IN2P3, 91105, Orsay; France
- ⁶⁷ Centro Nacional de Microelectrónica (IMB-CNM-CSIC), Barcelona; Spain
- ⁶⁸ Department of Physics, Indiana University, Bloomington IN; United States of America
- ⁶⁹ ^(a) INFN Gruppo Collegato di Udine, Sezione di Trieste, Udine; ^(b) ICTP, Trieste; ^(c) Dipartimento Politecnico di Ingegneria e Architettura, Università di Udine, Udine; Italy
- ⁷⁰ ^(a) INFN Sezione di Lecce; ^(b) Dipartimento di Matematica e Fisica, Università del Salento, Lecce; Italy
- ⁷¹ ^(a) INFN Sezione di Milano; ^(b) Dipartimento di Fisica, Università di Milano, Milano; Italy
- ⁷² ^(a) INFN Sezione di Napoli; ^(b) Dipartimento di Fisica, Università di Napoli, Napoli; Italy
- ⁷³ ^(a) INFN Sezione di Pavia; ^(b) Dipartimento di Fisica, Università di Pavia, Pavia; Italy
- ⁷⁴ ^(a) INFN Sezione di Pisa; ^(b) Dipartimento di Fisica E. Fermi, Università di Pisa, Pisa; Italy
- ⁷⁵ ^(a) INFN Sezione di Roma; ^(b) Dipartimento di Fisica, Sapienza Università di Roma, Roma; Italy
- ⁷⁶ ^(a) INFN Sezione di Roma Tor Vergata; ^(b) Dipartimento di Fisica, Università di Roma Tor Vergata, Roma; Italy
- ⁷⁷ ^(a) INFN Sezione di Roma Tre; ^(b) Dipartimento di Matematica e Fisica, Università Roma Tre, Roma; Italy
- ⁷⁸ ^(a) INFN-TIFPA; ^(b) Università degli Studi di Trento, Trento; Italy
- ⁷⁹ Universität Innsbruck, Department of Astro and Particle Physics, Innsbruck; Austria
- ⁸⁰ University of Iowa, Iowa City IA; United States of America
- ⁸¹ Department of Physics and Astronomy, Iowa State University, Ames IA; United States of America
- ⁸² Istinye University, Sariyer, Istanbul; Türkiye
- ⁸³ ^(a) Departamento de Engenharia Elétrica, Universidade Federal de Juiz de Fora (UFJF), Juiz de Fora; ^(b) Universidade Federal do Rio De Janeiro COPPE/EE/IF, Rio de Janeiro; ^(c) Instituto de Física, Universidade de São Paulo, São Paulo; ^(d) Rio de Janeiro State University, Rio de Janeiro; Brazil
- ⁸⁴ KEK, High Energy Accelerator Research Organization, Tsukuba; Japan
- ⁸⁵ Graduate School of Science, Kobe University, Kobe; Japan
- ⁸⁶ ^(a) AGH University of Krakow, Faculty of Physics and Applied Computer Science, Krakow; ^(b) Marian Smoluchowski Institute of Physics, Jagiellonian University, Krakow; Poland
- ⁸⁷ Institute of Nuclear Physics Polish Academy of Sciences, Krakow; Poland
- ⁸⁸ Faculty of Science, Kyoto University, Kyoto; Japan
- ⁸⁹ Research Center for Advanced Particle Physics and Department of Physics, Kyushu University, Fukuoka; Japan

- ⁹⁰ *Instituto de Física La Plata, Universidad Nacional de La Plata and CONICET, La Plata; Argentina*
- ⁹¹ *Physics Department, Lancaster University, Lancaster; United Kingdom*
- ⁹² *Oliver Lodge Laboratory, University of Liverpool, Liverpool; United Kingdom*
- ⁹³ *Department of Experimental Particle Physics, Jožef Stefan Institute and Department of Physics, University of Ljubljana, Ljubljana; Slovenia*
- ⁹⁴ *School of Physics and Astronomy, Queen Mary University of London, London; United Kingdom*
- ⁹⁵ *Department of Physics, Royal Holloway University of London, Egham; United Kingdom*
- ⁹⁶ *Department of Physics and Astronomy, University College London, London; United Kingdom*
- ⁹⁷ *Louisiana Tech University, Ruston LA; United States of America*
- ⁹⁸ *Fysiska institutionen, Lunds universitet, Lund; Sweden*
- ⁹⁹ *Departamento de Física Teórica C-15 and CIAFF, Universidad Autónoma de Madrid, Madrid; Spain*
- ¹⁰⁰ *Institut für Physik, Universität Mainz, Mainz; Germany*
- ¹⁰¹ *School of Physics and Astronomy, University of Manchester, Manchester; United Kingdom*
- ¹⁰² *CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille; France*
- ¹⁰³ *Department of Physics, University of Massachusetts, Amherst MA; United States of America*
- ¹⁰⁴ *Department of Physics, McGill University, Montreal QC; Canada*
- ¹⁰⁵ *School of Physics, University of Melbourne, Victoria; Australia*
- ¹⁰⁶ *Department of Physics, University of Michigan, Ann Arbor MI; United States of America*
- ¹⁰⁷ *Department of Physics and Astronomy, Michigan State University, East Lansing MI; United States of America*
- ¹⁰⁸ *Group of Particle Physics, University of Montreal, Montreal QC; Canada*
- ¹⁰⁹ *Fakultät für Physik, Ludwig-Maximilians-Universität München, München; Germany*
- ¹¹⁰ *Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), München; Germany*
- ¹¹¹ *Graduate School of Science and Kobayashi-Maskawa Institute, Nagoya University, Nagoya; Japan*
- ¹¹² *Department of Physics and Astronomy, University of New Mexico, Albuquerque NM; United States of America*
- ¹¹³ *Institute for Mathematics, Astrophysics and Particle Physics, Radboud University/Nikhef, Nijmegen; Netherlands*
- ¹¹⁴ *Nikhef National Institute for Subatomic Physics and University of Amsterdam, Amsterdam; Netherlands*
- ¹¹⁵ *Department of Physics, Northern Illinois University, DeKalb IL; United States of America*
- ¹¹⁶ ^(a) *New York University Abu Dhabi, Abu Dhabi;* ^(b) *University of Sharjah, Sharjah; United Arab Emirates*
- ¹¹⁷ *Department of Physics, New York University, New York NY; United States of America*
- ¹¹⁸ *Ochanomizu University, Otsuka, Bunkyo-ku, Tokyo; Japan*
- ¹¹⁹ *Ohio State University, Columbus OH; United States of America*
- ¹²⁰ *Homer L. Dodge Department of Physics and Astronomy, University of Oklahoma, Norman OK; United States of America*
- ¹²¹ *Department of Physics, Oklahoma State University, Stillwater OK; United States of America*
- ¹²² *Palacký University, Joint Laboratory of Optics, Olomouc; Czech Republic*
- ¹²³ *Institute for Fundamental Science, University of Oregon, Eugene, OR; United States of America*
- ¹²⁴ *Graduate School of Science, Osaka University, Osaka; Japan*
- ¹²⁵ *Department of Physics, University of Oslo, Oslo; Norway*
- ¹²⁶ *Department of Physics, Oxford University, Oxford; United Kingdom*
- ¹²⁷ *LPNHE, Sorbonne Université, Université Paris Cité, CNRS/IN2P3, Paris; France*
- ¹²⁸ *Department of Physics, University of Pennsylvania, Philadelphia PA; United States of America*
- ¹²⁹ *Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh PA; United States of America*
- ¹³⁰ ^(a) *Laboratório de Instrumentação e Física Experimental de Partículas — LIP, Lisboa;* ^(b) *Departamento de Física, Faculdade de Ciências, Universidade de Lisboa, Lisboa;* ^(c) *Departamento de Física, Universidade de Coimbra, Coimbra;* ^(d) *Centro de Física Nuclear da Universidade de Lisboa, Lisboa;* ^(e) *Departamento de Física, Universidade do Minho, Braga;* ^(f) *Departamento de Física Teórica y del Cosmos, Universidad de Granada, Granada (Spain);* ^(g) *Departamento de Física, Instituto Superior Técnico, Universidade de Lisboa, Lisboa; Portugal*

- ¹³¹ *Institute of Physics of the Czech Academy of Sciences, Prague; Czech Republic*
- ¹³² *Czech Technical University in Prague, Prague; Czech Republic*
- ¹³³ *Charles University, Faculty of Mathematics and Physics, Prague; Czech Republic*
- ¹³⁴ *Particle Physics Department, Rutherford Appleton Laboratory, Didcot; United Kingdom*
- ¹³⁵ *IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette; France*
- ¹³⁶ *Santa Cruz Institute for Particle Physics, University of California Santa Cruz, Santa Cruz CA; United States of America*
- ¹³⁷ ^(a) *Departamento de Física, Pontificia Universidad Católica de Chile, Santiago;* ^(b) *Millennium Institute for Subatomic physics at high energy frontier (SAPHIR), Santiago;* ^(c) *Instituto de Investigación Multidisciplinario en Ciencia y Tecnología, y Departamento de Física, Universidad de La Serena;* ^(d) *Universidad Andres Bello, Department of Physics, Santiago;* ^(e) *Instituto de Alta Investigación, Universidad de Tarapacá, Arica;* ^(f) *Departamento de Física, Universidad Técnica Federico Santa María, Valparaíso; Chile*
- ¹³⁸ *Department of Physics, University of Washington, Seattle WA; United States of America*
- ¹³⁹ *Department of Physics and Astronomy, University of Sheffield, Sheffield; United Kingdom*
- ¹⁴⁰ *Department of Physics, Shinshu University, Nagano; Japan*
- ¹⁴¹ *Department Physik, Universität Siegen, Siegen; Germany*
- ¹⁴² *Department of Physics, Simon Fraser University, Burnaby BC; Canada*
- ¹⁴³ *SLAC National Accelerator Laboratory, Stanford CA; United States of America*
- ¹⁴⁴ *Department of Physics, Royal Institute of Technology, Stockholm; Sweden*
- ¹⁴⁵ *Departments of Physics and Astronomy, Stony Brook University, Stony Brook NY; United States of America*
- ¹⁴⁶ *Department of Physics and Astronomy, University of Sussex, Brighton; United Kingdom*
- ¹⁴⁷ *School of Physics, University of Sydney, Sydney; Australia*
- ¹⁴⁸ *Institute of Physics, Academia Sinica, Taipei; Taiwan*
- ¹⁴⁹ ^(a) *E. Andronikashvili Institute of Physics, Iv. Javakishvili Tbilisi State University, Tbilisi;* ^(b) *High Energy Physics Institute, Tbilisi State University, Tbilisi;* ^(c) *University of Georgia, Tbilisi; Georgia*
- ¹⁵⁰ *Department of Physics, Technion, Israel Institute of Technology, Haifa; Israel*
- ¹⁵¹ *Raymond and Beverly Sackler School of Physics and Astronomy, Tel Aviv University, Tel Aviv; Israel*
- ¹⁵² *Department of Physics, Aristotle University of Thessaloniki, Thessaloniki; Greece*
- ¹⁵³ *International Center for Elementary Particle Physics and Department of Physics, University of Tokyo, Tokyo; Japan*
- ¹⁵⁴ *Department of Physics, Tokyo Institute of Technology, Tokyo; Japan*
- ¹⁵⁵ *Department of Physics, University of Toronto, Toronto ON; Canada*
- ¹⁵⁶ ^(a) *TRIUMF, Vancouver BC;* ^(b) *Department of Physics and Astronomy, York University, Toronto ON; Canada*
- ¹⁵⁷ *Division of Physics and Tomonaga Center for the History of the Universe, Faculty of Pure and Applied Sciences, University of Tsukuba, Tsukuba; Japan*
- ¹⁵⁸ *Department of Physics and Astronomy, Tufts University, Medford MA; United States of America*
- ¹⁵⁹ *United Arab Emirates University, Al Ain; United Arab Emirates*
- ¹⁶⁰ *Department of Physics and Astronomy, University of California Irvine, Irvine CA; United States of America*
- ¹⁶¹ *Department of Physics and Astronomy, University of Uppsala, Uppsala; Sweden*
- ¹⁶² *Department of Physics, University of Illinois, Urbana IL; United States of America*
- ¹⁶³ *Instituto de Física Corpuscular (IFIC), Centro Mixto Universidad de Valencia — CSIC, Valencia; Spain*
- ¹⁶⁴ *Department of Physics, University of British Columbia, Vancouver BC; Canada*
- ¹⁶⁵ *Department of Physics and Astronomy, University of Victoria, Victoria BC; Canada*
- ¹⁶⁶ *Fakultät für Physik und Astronomie, Julius-Maximilians-Universität Würzburg, Würzburg; Germany*
- ¹⁶⁷ *Department of Physics, University of Warwick, Coventry; United Kingdom*
- ¹⁶⁸ *Waseda University, Tokyo; Japan*
- ¹⁶⁹ *Department of Particle Physics and Astrophysics, Weizmann Institute of Science, Rehovot; Israel*
- ¹⁷⁰ *Department of Physics, University of Wisconsin, Madison WI; United States of America*

¹⁷¹ *Fakultät für Mathematik und Naturwissenschaften, Fachgruppe Physik, Bergische Universität Wuppertal, Wuppertal; Germany*

¹⁷² *Department of Physics, Yale University, New Haven CT; United States of America*

^a *Also Affiliated with an institute covered by a cooperation agreement with CERN*

^b *Also at An-Najah National University, Nablus; Palestine*

^c *Also at Borough of Manhattan Community College, City University of New York, New York NY; United States of America*

^d *Also at Center for High Energy Physics, Peking University; China*

^e *Also at Center for Interdisciplinary Research and Innovation (CIRI-AUTH), Thessaloniki; Greece*

^f *Also at Centro Studi e Ricerche Enrico Fermi; Italy*

^g *Also at CERN, Geneva; Switzerland*

^h *Also at Département de Physique Nucléaire et Corpusculaire, Université de Genève, Genève; Switzerland*

ⁱ *Also at Departament de Física de la Universitat Autònoma de Barcelona, Barcelona; Spain*

^j *Also at Department of Financial and Management Engineering, University of the Aegean, Chios; Greece*

^k *Also at Department of Physics and Astronomy, Michigan State University, East Lansing MI; United States of America*

^l *Also at Department of Physics, Ben Gurion University of the Negev, Beer Sheva; Israel*

^m *Also at Department of Physics, California State University, Sacramento; United States of America*

ⁿ *Also at Department of Physics, King's College London, London; United Kingdom*

^o *Also at Department of Physics, Stanford University, Stanford CA; United States of America*

^p *Also at Department of Physics, University of Fribourg, Fribourg; Switzerland*

^q *Also at Department of Physics, University of Thessaly; Greece*

^r *Also at Department of Physics, Westmont College, Santa Barbara; United States of America*

^s *Also at Hellenic Open University, Patras; Greece*

^t *Also at Institutio Catalana de Recerca i Estudis Avancats, ICREA, Barcelona; Spain*

^u *Also at Institut für Experimentalphysik, Universität Hamburg, Hamburg; Germany*

^v *Also at Institute for Nuclear Research and Nuclear Energy (INRNE) of the Bulgarian Academy of Sciences, Sofia; Bulgaria*

^w *Also at Institute of Applied Physics, Mohammed VI Polytechnic University, Ben Guerir; Morocco*

^x *Also at Institute of Particle Physics (IPP); Canada*

^y *Also at Institute of Physics and Technology, Ulaanbaatar; Mongolia*

^z *Also at Institute of Physics, Azerbaijan Academy of Sciences, Baku; Azerbaijan*

^{aa} *Also at Institute of Theoretical Physics, Iliia State University, Tbilisi; Georgia*

^{ab} *Also at L2IT, Université de Toulouse, CNRS/IN2P3, UPS, Toulouse; France*

^{ac} *Also at Lawrence Livermore National Laboratory, Livermore; United States of America*

^{ad} *Also at National Institute of Physics, University of the Philippines Diliman (Philippines); Philippines*

^{ae} *Also at Ochanomizu University, Otsuka, Bunkyo-ku, Tokyo; Japan*

^{af} *Also at Technical University of Munich, Munich; Germany*

^{ag} *Also at The Collaborative Innovation Center of Quantum Matter (CICQM), Beijing; China*

^{ah} *Also at TRIUMF, Vancouver BC; Canada*

^{ai} *Also at Università di Napoli Parthenope, Napoli; Italy*

^{aj} *Also at University of Colorado Boulder, Department of Physics, Colorado; United States of America*

^{ak} *Also at Washington College, Chestertown, MD; United States of America*

^{al} *Also at Yeditepe University, Physics Department, Istanbul; Türkiye*

* *Deceased*