

# Motor imagery, forward models and the cerebellum

Miall, R C

DOI:

[10.1007/s00426-023-01916-7](https://doi.org/10.1007/s00426-023-01916-7)

License:

Creative Commons: Attribution (CC BY)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Miall, RC 2024, 'Motor imagery, forward models and the cerebellum: a commentary on Rieger et al., 2023', *Psychological Research*. <https://doi.org/10.1007/s00426-023-01916-7>

[Link to publication on Research at Birmingham portal](#)

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.



# Motor imagery, forward models and the cerebellum: a commentary on Rieger et al., 2023

R. C. Miall<sup>1</sup>

Received: 27 June 2023 / Accepted: 14 December 2023  
© The Author(s) 2024

## Abstract

In this commentary on Rieger et al., *Psychological Research Psychologische Forschung*, 2023, I discuss possible ways to test the hypothesis that action imagery is achieved by simulations of actions through an internal forward model. These include brain imaging, perturbation through TMS, and psychophysical tests of adaptation of intended reach actions.

Rieger and colleagues (2023) make a convincing argument for the role of internal models operating in action imagery, and specifically forward models being the mechanism through which actions are simulated, and through which errors in imagery are detected. This fits well with the general concept of forward internal models in motor control (Miall & Wolpert, 1996), neural representations that are understood to receive efference copy of motor commands, as well as sensory inputs and that then generate an internal estimate of the sensory consequences of those commands (Wolpert et al., 1998). As the output is a sensory representation (or an estimate of the internal state of the motor system, that can be converted to a sensory representation), the forward model output could be available as an imagined representation of action. During imagery, no motor commands should reach the musculature, or else movement would be generated. Hence, the internal efferent copies must be either derived independently of descending commands, or the commands inhibited downstream of the internal model. Errors in action imagery are occasionally generated, with the implication that mistakes are made either in the generation of the command or efference copy, relative to the desired action, or in the forward model estimation based on this command, or possibly in the comparison process. Finally, there are possible alternatives to motor system-based simulation of actions, particularly the use of generalized knowledge, possibly gained through observation of one's own or others' actions.

These basic facts (as laid out in greater detail by Rieger et al.) suggest that their hypothesis could be tested by challenging the forward model.

First, it is likely that the cerebellum performs forward model operations and is potentially the main—if not exclusive—site of a motor-related forward model (Miall et al., 1993; Sokolov et al., 2017; Wolpert et al., 1998). It receives descending motor commands and its outputs project back to fronto-parietal cortical areas that might subserve mental imagery. Hence, one could test for cerebellar activation during mental imagery, and indeed many studies have done so. Héту et al. (2013) provide a meta-analysis of evidence from 75 brain imaging papers, and the cerebellum is consistently activated, with ipsilateral loci consistent with upper and lower limb actions. However, imaging studies alone cannot exclude a cerebellar contribution in other ways, independent of forward modeling—for example in the inverse model responsible for motor command generation.

Second, one could look for a causal relationship, by testing mental imagery during disturbances of cerebellar function. Battaglia et al. (2006) found that stroke affecting the cerebellum disrupted the changes in excitability of motor cortex that are normally induced by motor imagery. More directly, González et al. (2005) found that cerebellar stroke survivors showed slowing of finger sequences in both actual and imagined conditions, consistent with the cerebellum contributing to mental simulation. There are few reports of direct modulation of the cerebellum by transcranial magnetic or electrical stimulation, although Grami et al. (2022) and Cengiz and Boran (2016) have independently found that TDCS of the cerebellum influenced the extent that imagery of actions could modulate cerebral cortical activity, analogous to Battaglia et al.'s (2006) result. However,

✉ R. C. Miall  
r.c.miall@bham.ac.uk

<sup>1</sup> School of Psychology, University of Birmingham, Birmingham, UK

we previously reported that cerebellar TMS can selectively bias reaching movements to a visual target, and this shift in reach direction is likely to be because of the temporary blockade of cerebellar forward model output (Miall et al., 2007). This task would be well suited to test mental imagery: the final position of the hand could be reported relative to the target for imagined reaches with and without TMS. I would predict a directional bias in imagined hand end-point during stimulated trials, as is seen in active movement, and this would be strong evidence for the forward model operating during action imagery.

Third, one might explore the issue of forward model corrections and learning during imagined action errors. It is well documented that imagining and rehearsal of actions can lead to improved performance, and the assumption is that success in the imagined action leads to beneficial changes in execution. Rieger et al. (2023) discuss the converse situation and report imagined errors, albeit at a frequency lower than in actual actions. It seems plausible that errors might accumulate throughout the neural chain of command, from planning of intended actions, the generation of commands, prediction of their consequences, in the integration of the commands with brainstem and spinal circuit activity, or in muscular execution. Because there are no action outcomes in imagery, perception of errors in imagined actions is probably only possible if they occur in the first three stages, or in the comparison of intention and imagined outcome. One can ask what changes these errors lead to—is it in the intended actions or the forward modeling of their consequences? Recent papers (Morehead et al., 2017; Tsay et al., 2022) have shown that presentation of a “visually clamped” error after a reaching action, regardless of the actual hand reach direction, leads to the implicit adaptation of the reach movements, to compensate for the sensory prediction error between action and displayed feedback. These sensory prediction errors drive cerebellar-dependent learning (Tseng et al., 2007). One could perform an experiment with imagined reach actions toward a visual target with a visual “error” presented after each imagined action that is clamped to one side of the target. I would predict that the imagined reaches would gradually adapt as do actual actions (Tsay et al., 2022). After adaptation, one could test the intended action, the imagined outcome, and the direction of actual reaches, to separate out where the adaptive changes have occurred. If the forward model was adapted because of these errors in imagery, then there should be a remapping of both intended action and visual outcome, but if only the intentions are adapted, there would be no remapping, only a shift in intended reach direction.

Finally, Rieger et al. (2023) discuss the relationship between imagery through action simulation (i.e., forward models) versus propositional knowledge of actions. It is interesting to explore the experience-dependent nature of

imagery: forward models are developed through experience of actions, with a sensory prediction error between expected and actual outcome driving their improved accuracy (Tseng et al., 2007). One approach would be to test action imagery in the absence of recent experience of certain actions. Chronic or congenital loss of limbs is an obvious choice, and Malouin et al. (2009) report reduced vividness of imagined actions after amputation or disuse of a limb. Congenital or chronic loss of sensation is also likely to lead to a degraded forward model process, and IW, a man with 4 decades of profound somatosensory loss, has degraded kinaesthetic motor imagery, while still having intact or enhanced visual imagery (ter Horst et al., 2012). While not directly implicating the cerebellum, these results are consistent with a motor simulation degraded because of the chronic absence of sensory inflow to the forward model.

**Funding** Not available.

**Availability of data and materials** Not available.

## Declarations

**Conflict of interest** The author declares no competing interests.

**Ethical approval** Not available.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Battaglia, F., Quartarone, A., Ghilardi, M. F., Dattola, R., Bagnato, S., Rizzo, V., Morgante, L., & Girlanda, P. (2006). Unilateral cerebellar stroke disrupts movement preparation and motor imagery. *Clinical Neurophysiology*, *117*(5), 1009–1016. <https://doi.org/10.1016/j.clinph.2006.01.008>
- Cengiz, B., & Boran, H. E. (2016). The role of the cerebellum in motor imagery. *Neuroscience Letters*, *617*, 156–159. <https://doi.org/10.1016/j.neulet.2016.01.045>
- González, B., Rodríguez, M., Ramirez, C., & Sabaté, M. (2005). Disturbance of motor imagery after cerebellar stroke. *Behavioral Neuroscience*, *119*(2), 622–626. <https://doi.org/10.1037/0735-7044.119.2.622>
- Grami, F., de Marco, G., Bodranghien, F., Manto, M., & Habas, C. (2022). Cerebellar transcranial direct current stimulation

- reconfigures brain networks involved in motor execution and mental imagery. *Cerebellum*, 21(4), 665–680. <https://doi.org/10.1007/s12311-021-01322-y>
- Hétu, S., Grégoire, M., Saimpont, A., Coll, M.-P., Eugène, F., Michon, P.-E., & Jackson, P. L. (2013). The neural network of motor imagery: An ALE meta-analysis. *Neuroscience & Biobehavioral Reviews*, 37(5), 930–949. <https://doi.org/10.1016/j.neubiorev.2013.03.017>
- Malouin, F., Richards, C. L., Durand, A., Descent, M., Poiré, D., Frémont, P., Pelet, S., Gresset, J., & Doyon, J. (2009). Effects of practice, visual loss, limb amputation, and disuse on motor imagery vividness. *Neurorehabilitation and Neural Repair*, 23(5), 449–463. <https://doi.org/10.1177/1545968308328733>
- Miall, R. C., Christensen, L. O. D., Cain, O., & Stanley, J. (2007). Disruption of state estimation in the human lateral cerebellum. *PLoS Biology*, 5(11), 2733–2744. <https://doi.org/10.1371/journal.pbio.0050316>
- Miall, R. C., Weir, D. J., Wolpert, D. M., & Stein, J. F. (1993). Is the cerebellum a smith predictor? *Journal of Motor Behavior*, 25(3), 203–216. <https://doi.org/10.1080/00222895.1993.9942050>
- Miall, R. C., & Wolpert, D. M. (1996). Forward models for physiological motor control. *Neural Networks*, 9(8), 1265–1279.
- Morehead, J. R., Taylor, J. A., Parvin, D. E., & Ivry, R. B. (2017). Characteristics of implicit sensorimotor adaptation revealed by task-irrelevant clamped feedback. *Journal of Cognitive Neuroscience*, 29(6), 1061–1074. [https://doi.org/10.1162/jocn\\_a\\_01108](https://doi.org/10.1162/jocn_a_01108)
- Rieger, M., Boe, S. G., Ingram, T. G. J., Bart, V. K. E., & Dahm, S. F. (2023). A theoretical perspective on action consequences in action imagery: Internal prediction as an essential mechanism to detect errors. *Psychological Research Psychologische Forschung*. <https://doi.org/10.1007/s00426-023-01812-0>
- Sokolov, A. A., Miall, R. C., & Ivry, R. B. (2017). The cerebellum: Adaptive prediction for movement and cognition. *Trends in Cognitive Sciences*, 21(5), 313–332. <https://doi.org/10.1016/j.tics.2017.02.005>
- ter Horst, A. C., Cole, J., van Lier, R., & Steenbergen, B. (2012). The effect of chronic deafferentation on mental imagery: A case study. *PLoS ONE*, 7(8), e42742. <https://doi.org/10.1371/journal.pone.0042742>
- Tsay, J. S., Kim, H. E., Haith, A. M., & Ivry, R. B. (2022). Understanding implicit sensorimotor adaptation as a process of proprioceptive re-alignment. *BioRxiv*. <https://doi.org/10.1101/2021.12.21.473747>
- Tseng, Y.-W., Diedrichsen, J., Krakauer, J. W., Shadmehr, R., & Bastian, A. J. (2007). Sensory prediction errors drive cerebellum-dependent adaptation of reaching. *Journal of Neurophysiology*, 98(1), 54–62. <https://doi.org/10.1152/jn.00266.2007>
- Wolpert, D. M., Miall, R. C., & Kawato, M. (1998). Internal models in the cerebellum. *Trends in Cognitive Sciences*, 2(9), 338–347. [https://doi.org/10.1016/S1364-6613\(98\)01221-2](https://doi.org/10.1016/S1364-6613(98)01221-2)

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.