

## Using GWAS to identify genetic predisposition in hepatic autoimmunity

Webb, Gwilym; Hirschfield, G M

DOI:

[10.1016/j.jaut.2015.08.016](https://doi.org/10.1016/j.jaut.2015.08.016)

License:

Creative Commons: Attribution (CC BY)

*Document Version*

Publisher's PDF, also known as Version of record

*Citation for published version (Harvard):*

Webb, G & Hirschfield, GM 2016, 'Using GWAS to identify genetic predisposition in hepatic autoimmunity', *Journal of Autoimmunity*, vol. 66, pp. 25-39. <https://doi.org/10.1016/j.jaut.2015.08.016>

[Link to publication on Research at Birmingham portal](#)

**Publisher Rights Statement:**

Eligibility for repository : checked 16/12/2015

**General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

**Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.



Contents lists available at ScienceDirect

Journal of Autoimmunity

journal homepage: [www.elsevier.com/locate/jautimm](http://www.elsevier.com/locate/jautimm)

## Review article

## Using GWAS to identify genetic predisposition in hepatic autoimmunity

G.J. Webb, G.M. Hirschfield\*

NIHR Birmingham Liver Biomedical Research Unit, University of Birmingham, Birmingham, UK

## ARTICLE INFO

## Article history:

Received 21 August 2015

Accepted 23 August 2015

Available online xxx

## Keywords:

Primary biliary cirrhosis  
 Primary sclerosing cholangitis  
 Autoimmune hepatitis  
 Biliary autoimmunity  
 Genetic risk  
 Immune regulation  
 Epistasis  
 Epigenetics  
 Heritability

## ABSTRACT

Primary biliary cirrhosis (PBC), primary sclerosing cholangitis (PSC) and autoimmune hepatitis (AIH) represent the three major hepatic autoimmune conditions. Patient morbidity and mortality remain high across these three diseases, and an unmet need for rational therapy exists. Disease understanding has focused on combining clinical and laboratory based science to provide better insights into the joint host and environmental factors necessary for the initiation, and perpetuation, of hepato-biliary inflammation. Twin studies, family studies, population studies and an inter-relationship with other autoimmune phenomena suggest a genetic component to risk for each disease. Until recently, understanding of this genetic risk has been limited to HLA haplotypes. Associations with risk-conferring and protective HLA haplotypes are present in all three diseases. Over the last few years, genome-wide association studies (GWAS), and related genetic association studies, have greatly increased understanding of the genetic risk signature of these three diseases and autoimmunity in general. Here we consider the rationale for GWAS in general and with specific reference to hepatic autoimmunity. We consider the process of GWAS, and highlight major findings to date. Potential functional implications of key findings are discussed including the IL-12/STAT4 pathway in PBC and the CD28/IL-2 pathway in PSC. We describe the marked pleiotropy demonstrated by PBC and PSC, which is consistent with other autoimmune diseases. Further, we focus on specific gene associations including *SH2B3*, which is common to all three diseases, and *FUT2* in PSC, which represents a link between environment and genetics. We review attempts to translate GWAS findings into basic laboratory models including *in vivo* systems and highlight where clinical observations relate to genetics. Finally we describe deficiencies in GWAS to date and consider future study of genetics in hepatic autoimmunity.

© 2015 Published by Elsevier Ltd.

## Contents

1. Introduction .....	00
2. Why study genetics in hepatic autoimmunity? .....	00
2.1. Twin studies .....	00
2.2. Familial risk .....	00
2.3. Populations .....	00
2.4. Individuals developing multiple autoimmune conditions suggests shared risk .....	00
3. The genetic risk profile of hepatic autoimmunity is challenging to assess .....	00
4. The rationale behind GWAS .....	00
5. The process of GWAS .....	00
6. Major observations in hepatic autoimmunity resulting from GWAS .....	00
6.1. HLA associations .....	00
6.2. Non-HLA associations .....	00
6.2.1. The IL-2 pathway in PSC .....	00

\* Corresponding author. Centre for Liver Research, Institute of Biomedical Research, Wolfson Drive, University of Birmingham, Birmingham, B15 2TT, UK.

E-mail address: [g.hirschfield@bham.ac.uk](mailto:g.hirschfield@bham.ac.uk) (G.M. Hirschfield).

6.2.2.	The IL12-STAT4 pathway in PBC .....	00
6.2.3.	FUT2 .....	00
6.2.4.	SH2B3 – Lnk .....	00
6.2.5.	Epistasis .....	00
7.	Translating GWAS findings to clinical cohorts .....	00
7.1.	HLA .....	00
7.2.	Non-HLA .....	00
8.	Deficiencies in GWAS to date .....	00
8.1.	GWAS results do not account for all the observed risk .....	00
8.2.	Genetic information cannot yet explain key aspects of disease phenotype .....	00
8.3.	GWAS does not reflect disease risk in varied ethnicities .....	00
8.4.	Positive selection pressure for genes implicated by GWAS is not yet explained .....	00
8.5.	GWAS does not yet map to animal models nor therapies .....	00
8.6.	GWAS may not capture epigenetic modifications, including heritable epigenetic traits .....	00
9.	The future of genetic investigation of hepatic autoimmunity .....	00
10.	Conclusion .....	00
	Conflict of interest .....	00
	Acknowledgments .....	00
	References .....	00

## 1. Introduction

Autoimmunity may be considered as a “clinical syndrome caused by the activation of T cells or B cells, or both, in the absence of an ongoing infection or other discernible cause” [1] and is typically associated with autoantibodies reactive against self antigens [2]. The prevalence of autoimmunity in general is approaching 10% [3], and many of these individuals develop autoimmune phenomena affecting multiple organ systems simultaneously [4].

Hepatic autoimmunity comprises three major – and sometimes overlapping – clinico-pathological syndromes: primary biliary cirrhosis (PBC; also referred to as primary biliary cholangitis), autoimmune hepatitis (AIH) and primary sclerosing cholangitis (PSC) [5–7]. Each is associated with autoantibodies and with an increased likelihood of autoimmunity affecting other organ systems. Notably, these are uncommon conditions with the prevalence of PBC quoted as 1.91–40.2 per 100,000 and PSC as 0–16.2 per 100,000 [8]. Data for AIH are more sparse, but a recent Danish study has quoted a prevalence of 23.9 per 100,000 [9]. These diseases can impact people of all ages, persist as chronic illnesses rather than acute illnesses, and remain challenging to treat. Of the three diseases, only AIH typically responds to immunosuppression. AIH may be separated into type 1 and type 2 with the former more responsive to immunosuppression. There are few data on the genetics of type 2 AIH and the remainder of this article discusses type 1 AIH.

In each of the three major hepatic autoimmune conditions, as in most major autoimmune diseases, there is evidence for a strong genetic component to risk [10]. Genome wide association studies (GWAS), and related high-density genotyping arrays, represent a powerful method for exploring the genetic makeup of this risk. This review for a Special Edition of Journal of Autoimmunity focuses on the process of applying such studies to hepatic autoimmunity. This is apt given the life-long contribution from Professors Diego and Giorgina Vergani in the field of hepatology, particularly the presentation and management, as well as immunologic phenotype, of autoimmune liver disease in children. The studies discussed herein parallel and complement such clinical and immunologic evaluation of these diseases, collectively driving forward our efforts to better treat patients.

## 2. Why study genetics in hepatic autoimmunity?

There is a clear genetic component to the risk of developing

autoimmunity in general, and hepatic autoimmunity specifically. This inference is based on observations made on twin pairs, family studies, population-based studies, and clinical observations, and underpins the logic behind conducting GWAS. Together these observations strongly suggest a significant genetic component to the development of hepatic autoimmunity. A deeper understanding of genetic associations is hoped to guide investigation of underlying pathobiology and so translate to improved therapies, personalized medicine and more accurate prognostication [11].

### 2.1. Twin studies

Comparing disease concordance in mono- and dizygotic twins living in similar environments allows estimation of the relative contribution from environmental and genetic inputs to disease risk. Unfortunately, the low frequency of monozygotic twins (approximately 3.5 per 1000 live births) makes sufficient recruitment in rarer diseases such as hepatic autoimmunity challenging.

PBC has the highest reported concordance in monozygotic twins of any autoimmune condition, with the exception of celiac disease: 63% in one small series as compared to a risk in dizygotic twins close to the population level of <0.5% with no concordance between the 8 pairs studied [12]. Intriguingly, ages of presentation and presenting symptoms were similar among concordant twins and two of the discordant twins had other major autoimmune conditions. Similar data are not currently available for AIH or PSC, although in a recent report there was no genetic concordance in five dizygotic twin pairs with concordance in the single set of monozygotic twins from a national Dutch AIH study [13].

### 2.2. Familial risk

A particularly increased risk within a family as compared to the surrounding population tends to suggest an increased genetic risk, although unrecognized environmental risks are always also potentially relevant. Such genetic risk can be measured by the relative sibling risk (denoted as  $\lambda_s$ ) i.e. the increased risk conferred on the sibling of an individual affected with a certain trait as a multiple of the baseline population risk. For PBC  $\lambda_s$  has been calculated at  $\times 10.5$  in a large UK series [14] and  $\times 10.7$  in the USA [15]. For, PSC  $\lambda_s$  is approximately  $\times 10$  [16], whilst for AIH estimates

**Table 1**  
Reported coincidence of other autoimmune disease with PBC, PSC and AIH.

Probable or definite co-incident condition	PBC [108]	PSC [109]	AIH [110]
Sjögren syndrome	25		3
Autoimmune thyroid disease	23	4	14 <sup>a</sup>
Rheumatoid arthritis	17	1	28 <sup>a</sup>
Scleroderma	8		
Raynaud's phenomenon	24		
Systemic lupus erythematosus	1		
Autoimmune thrombocytopenic purpura	1		
Pernicious anemia	4		
Inflammatory bowel disease		62%	8
Celiac disease		1	2
Diabetes mellitus		4	9
Psoriasis		3	3
Sarcoidosis		4	1
<b>Proportion with ≥ 1 extra-hepatic autoimmune condition</b>	<b>53%</b>	<b>71%</b>	<b>49%</b>

<sup>a</sup> The definitions used when searching for associations with these conditions in this study were likely to also include non-autoimmune phenomena e.g. osteoarthritis.

are lacking. However, in AIH it is estimated that 40% of individuals have a family history of autoimmunity in general [17]. In PBC, the risk for expressing AMA and for developing disease is increased in first-degree relatives [18,19] and a disproportionate number of PBC patients have relatives with the disease at the point of diagnosis [8,20].

### 2.3. Populations

When one subset of genetically admixed populations in a shared geographic area have an increased risk of a trait this may also bely a genetic component to risk. Examples of such phenomena in hepatic autoimmunity include reports of numbers of individuals affected by PBC are also abnormally high in ethnically defined sub-populations [21,22] and within selected wider families in characterized comprehensive healthcare systems [19,23]. Data on PSC sub-populations are more scarce, and although significant variation is seen between geographically separated populations, this may also indicate varying environmental exposure [8]. Data in AIH are similarly lacking with diagnostic variability between countries a

key consideration when drawing comparisons [24].

### 2.4. Individuals developing multiple autoimmune conditions suggests shared risk


All three major hepatic autoimmune conditions are strongly associated with coincident extra-hepatic autoimmunity. The findings of three representative reports are summarized in Table 1. Such observations are consistent with autoimmune conditions in general, where development of one autoimmune condition increases the risk of another. However, this is not always the case and, for example, a diagnosis of rheumatoid arthritis is protective against a diagnosis of multiple sclerosis and *vice versa* [3].

### 3. The genetic risk profile of hepatic autoimmunity is challenging to assess

As suggested by the significant but relatively modest sibling relative risk and incomplete concordance in monozygotic twins, the major hepatic autoimmune conditions do not follow autosomal inheritance. In contrast, only a small minority of causes of hepatic autoimmunity are monogenetically inherited: Autoimmune polyendocrinopathy candidiasis ectodermal dystrophy [25], heterozygous *CTLA4* mutations [26], immunodysregulation polyendocrinopathy enteropathy X-linked syndrome [27], and *GATA2* mutation [28] may all present with AIH-like syndromes. Notably, all of these are associated with disruption of the regulatory CD4<sup>+</sup> T cell pathway.

In addition, there is strong evidence of environmental influences on PBC [15], PSC [29] and AIH [17]. Similarly to their genetic profile, no currently identified environmental risk factor is sufficient to explain a large proportion of disease risk. This combination of an uncertain relative contribution to disease risk from multiple genetic and environmental variables – as well as probable interactions between these factors – marks these conditions as multifactorial, polygenetic and complex (Fig. 1) [30].

Unraveling the genetic contribution to disease risk is therefore a challenge. Prior to GWAS, a large number of candidate gene studies had been undertaken in hepatic autoimmune diseases [31–33]. However, these studies suffered from relatively weak statistical



**Fig. 1. The make up of risk for multifactorial autoimmune diseases.** The total risk of autoimmune diseases is made up of uncertain contributions from environmental and genetic risk. Interpreting their relative contributions is made more challenging by those with similar genetics frequently being exposed to similar environments. There is also a contribution from gene–gene (epistasis) and gene–environment interactions. Heritable epigenetic modifications provide a further potential influence and may be missed by standard genotyping techniques.

power, a strong potential for type 1 statistical errors, lacked reproducibility and were criticized for their *a priori* approach based on an incomplete understanding of disease biology. Of the non-HLA candidate gene associations listed in the three pre-GWAS reviews cited above, none has subsequently shown significant association in subsequent GWAS.

#### 4. The rationale behind GWAS

GWAS offers an approach that presupposes that there is a genetic component to the risk of developing a disease trait, but seeks evidence of association across the whole genome without assuming other prior knowledge. GWAS exploits the common genetic variability that exists between individuals. When considering a particular trait, GWAS works on the assumption that the risk of developing the trait of interest is at least partly derived from the cumulative effect of multiple variants, each of which confers low risk by itself. If two groups with and without a trait of interest are compared, certain variants may be disproportionately represented. If these associations are sufficiently specific and sufficiently divergent from the distribution seen in a control group drawn from the same population, risk may be associated. Further analysis of identified risk loci can aid prediction of specific implicated genes. This hypothesis-free approach avoids much of the criticism leveled at candidate gene studies.

GWAS has been made possible by leveraging advances in chip-based genotyping technology and incremental decreases in the cost of such techniques [34]. By generating data that are amenable to combination, the power – and number of associations identified

– of individual GWAS may be increased by meta-analysis. This also permits combination of datasets to improve statistical power and facilitate replication analyses, especially where the same genotyping platforms or an overlapping selection of SNVs are used.

Considering fixed genetic information inherently focuses on the root rather than result of disease process. Potential gains from GWAS include improvements in predicting whether an individual will develop a disease, earlier prediction of disease severity or sub-phenotype, potential biomarker identification and improved diagnostics. Identifying specific genes and therefore protein targets guides the generation of pre-clinical models, informs the development of novel candidate drug treatments and may also reveal opportunities for drug re-purposing [11,35].

#### 5. The process of GWAS

Conducting GWAS requires the assembly of cohorts of individuals with and without a trait of interest. Care must be taken to assess for the presence of over-lapping conditions, which may bias genetic risk associations and misdiagnoses, which may dilute risk associations. Relevant examples include the presence or absence of ulcerative colitis in PSC or in distinguishing Type 1 from Type 2 AIH [30,36]. Although, finding sufficient well-phenotyped cases in these relatively rare diseases can be challenging, over the last decade high-throughput genetic studies have transformed our understanding of the risk architecture of autoimmune conditions in general and hepatic autoimmunity in particular (Table 2).

Once appropriate test and control cohorts have been assembled, each subject of both groups is genotyped for a wide array of

**Table 2**  
Genome-wide analyses in autoimmune liver disease.

Study	Year	Patients	Controls	Replication cohort Pat/cont <sup>a</sup>	Population	Loci implicated	Notes
<b>PBC</b>							
Hirschfield et al. [111]	2009	536	1536	526/1206	White N American	<i>HLA, IL12A, IL12RB2</i>	
Hirschfield et al. [112]	2010	494	1502	857/3198	White N American	<i>IRF-TNPO3, MMEL1, IKZF3</i>	Extension of [111]
Liu et al. [113]	2010	453	945	None	Italian and meta N American	<i>SPIB, IKZF3, IRF5-TNPO3</i>	Combination study with [111] population
Mells et al. [114]	2011	1840	5163	620/2514	N European	<i>DENND1B, STAT4, CD80, NFKB1, IL7R, CXCR5, TNFRSF1A, RAD51L1, CLEC16A, MAP3K7IP1; PLCL2, RPS6KA4, TNFAIP2</i> (in combination with Italian data [115])	
Nakamura et al. [39]	2012	487	476	787/615	Japanese	<i>HLA, TNFSF15, POU2AF1</i>	Japanese
Cordell [98]	2015	2745	9802	3616/4261	N European, Italian, N American	<i>IL1R1, CCL20, DGKQ, C5orf30, IL12B, TNFAIP3</i>	Large meta-analysis
<b>PSC</b>							
Karlsen et al. [116]	2010	285	298	766/2935	Scandinavian, German	<i>HLA</i>	
Melum et al. [117]	2011	715	2962	1025/2174	Scandinavian, German	<i>HLA, BCL2L11, MST1</i>	
Folseraas et al. [118]	2012	715	2962	1221/3508	Scandinavian and German discovery; Scandinavian, Central European and US replication	<i>MMEL1-TNFRSF14</i>	Fine-mapping study based on [117]
Srivastava et al. [119]	2012	992	5162	N/A	UK Caucasians	<i>MST1, IL2RA</i>	Replication study
Ellinghaus et al. [120]	2013	392	2977	1012/2694	German, Scandinavian	<i>GPR35, TCF4</i>	Combined analysis with ulcerative cases
Liu et al. [97]	2013	3789	25079		Pan-European, N America	<i>HLA, MMEL1-TNFRSF14, CD28, MST1, IL2/IL21, BACH2, IL2RA, SIK2, HDAC7, SH2B3, CD226, FUT2, PSMG1</i>	
<b>Type 1 AIH</b>							
de Boer et al. [36]	2014	649	15,638	466/4103	Dutch (discovery); German/Swiss (replication)	<i>HLA, SH2B3</i>	

Only loci associated at  $p < 5 \times 10^{-8}$  are included with the exception of SH2B3 ( $p = 7.7 \times 10^{-8}$ ).

<sup>a</sup> Total of all replication cohorts when several used. Where is not yet possible to distinguish between several candidate genes, a single example has been given.

individual single nucleotide variants (SNV). These SNV are pre-selected from those sufficiently prevalent in the general population, and sufficiently spaced across the genome, to power subsequent statistical association analysis. SNV selections are drawn from large haplotype mapping projects, particularly the International HapMap project. In the case of GWAS, these traits are chosen from variations identified during sequencing efforts to dissect the genome and are spread across it. In more focused studies, a targeted selection of SNVs may be used. In the case of the ImmunoChip, SNVs implicated in immune pathways and other autoimmune diseases – including some rarer than those used in most GWAS—are used to allow finer-resolution analysis of loci of interest [37].

The analysis of large-scale GWAS-derived datasets enables identification of individual variants with estimates of the risk that they confer. Notably, far more common SNVs have been described than have been assessed in any one GWAS. Therefore over-representation of a given SNV in one cohort or another is likely to represent linkage disequilibrium rather than the causative mutation itself. Exploitation of linkage disequilibrium – the phenomenon by which genes located closer together on are more frequently inherited together – allows likely risk loci to be identified and therefore candidate genes within or near such loci inferred [38]. Thus, GWAS does not typically detect specific mutations and the resolution of a particular study may make it difficult to comment on the specific gene implicated. Several GWAS ‘hits’ in hepatic autoimmunity may still relate to one of several proposed genes.

The multiple comparisons made in GWAS make both statistical correction and the use of a conservative level of statistical significance mandatory. Typically, overall values of  $p < 5 \times 10^{-8}$  are taken as indicating significance in a genome-wide study. A key potential confounder in GWAS is the potential for false positives/type 1 errors because of the multiple comparisons inherent to its analysis. Thus, careful adjustment for the number of such comparisons is mandatory. Also mandatory is confirmation of associations by replication in separate validation cohorts.

An important component of GWAS is comparing the test cohort with a suitable control: i.e. one that would be expected to have a similar distribution of common SNVs other than those associated with the disease trait of interest. Typically this means a control cohort drawn from healthy individuals in an ethnically and geographically similar area. With the notable exception of the Japanese PBC GWAS [39], all of the studies performed in hepatic GWAS to date have concentrated on populations of predominantly white European origin. Two associations from the Japanese PBC cohort (*POU2AF1* and *TNFSF15*) are unique to that study. Also, recent work in a Chinese cohort failed to replicate an association in the IL-12 pathway that has proven reproducible in European populations [40].

Following the broad model described above, risk loci associated with PBC, PSC and AIH at genome-wide significance levels ( $p < 5 \times 10^{-8}$ ) have been identified and these are summarized in Tables 3 and 4. This number will increase with larger, and higher resolution, studies e.g. there is an ongoing effort to perform a meta-analysis of PSC GWAS datasets. Notably, there is a high degree of overlap (or pleiotropy) between loci implicated in hepatic autoimmunity to date and those associated with other autoimmune conditions (Fig. 2).

## 6. Major observations in hepatic autoimmunity resulting from GWAS

### 6.1. HLA associations

GWAS has given robust support to a number of associations with HLA haplotypes with each of the major diseases. Several of these

**Table 3**  
HLA associations of hepatic autoimmune diseases.

<b>PBC</b>		
<i>Risk conferring</i>		
HLA-DQA1*04:01-DQB1*04:02-DRB1*08:01-B*39:05	[58,96]	*
HLA-DRB1*04:04-DQB1*03:02	[58,96]	*
	N. Am, UK, Ital	
HLA-DRB1*14-DPB1*03:01	[42]	*
	Ital	
HLA-DRB1*08:03-DQB1*06:01	[64,69]	
HLA-DRB1*04:05-DQB1*04:01	[69]	Jap
<i>Protective</i>		
HLA-DQB1*06:02-DRB1*15:01-DQA1*01:02-B*07:02	[58,96]	*
HLA-DQB1*03:01-DRB1*11:01-DQA1*05:01-DRB1*11:04	[58,64,96]	*
HLA-DRB1*13:02-DQB1*06:04	[69]	Jap
HLA-DRB1*11:01-DQB1*03:01	[69]	Jap
<b>PSC</b>		
<i>Risk conferring</i>		
HLA-B*08:01	[86]	N. Am inc Afr-Am
HLA-DRB1*03:01-DQA1*05:01-DQB1*02:01	[121]	N. Am, N. Euro
HLA-DRB1*13:01-DQA1*01:03-DQB1*06:03		
HLA-DRB1*15:01-DQA1*01:02-DQB1*06:02		
HLA-DRB1*01:01-DQA1*01:01		
<i>Protective</i>		
HLA-DRB4*01:03-DRB1*04:01-DQA1*03-DQB1*03:02	[122]	
HLA-DRB4*01:03-DRB1*07:01-DQA1*02:01-DQB1*03:03		
HLA-DRB4*02:02-DRB1*11:01-DQA1*05:01-DQB1*03:01		
<b>AIH (Type 1)</b>		
<i>Risk conferring</i>		
HLA-A1-HLA-B8-DRB3*01:01-DRB1*03:01-DQA1*05:01-DQB1*02:01		N Euro [123]
HLA-DR4*04:04 and HLA-DRB*04:05		Jap [124], Cent Am, China
HLA-DRB1*04:01		N Euro
DRB1*13:01-DQB1*06		Lat Am [125]
HLA-DRB1*01 and HLA-DRB1*14		W. India [126]
<i>Protective</i>		
HLA-DRB5*01:01-DRB1*15:01		N Euro [123]
HLA-DQB1*04-DQB1*03:01		Lat. Am. [125]

[126]\* Dense SNV analysis and subsequent conditional analysis of this HLA haplotype in the Italian PBC cohort has suggested that risk-conferring/protective effects are predominantly due to variants in HLA-DRB1 with associated linkage disequilibrium [42].

associations were made prior to the GWAS era. HLA associations strongly implicate the adaptive immune system: the HLA region on chromosome 6 is highly variable encodes both the class I and II major histocompatibility complexes. Associations are summarized in Table 3. All three hepatic autoimmune diseases have associations with variants in both class I (encoded by HLA-A, HLA-B and HLA-C) and class II (HLA-D) major histocompatibility complexes, with more associations in HLA-D. HLA haplotype associations are seen in all major autoimmune diseases (Fig. 2).

SNVs affecting HLA can alter the MHC antigen presentation repertoire and have been associated with disease provoking responses in response to otherwise innocuous certain agents: for example HLA-B\*57:01 and abacavir hypersensitivity [41]. In PBC much of the risk has been narrowed to HLA-DRB1 with much of these rest of the association signal attributed to linkage disequilibrium [42].

One theory that complements specific HLA associations with specific diseases alongside multiple shared other non-HLA variants is that HLA haplotypes may direct autoimmunity in an otherwise prone individual, perhaps after an environmental trigger. For

**Table 4**  
Major non-HLA associations from GWAS in hepatic autoimmunity.

Locus	PBC	PSC	AIH	Study	SNV	OR	P-value	Candidate gene(s)	Selected functional notes
1p36	x			[112]	rs3748816	1.33	3.15E-08	<i>MMEL1</i>	
1p36		x		[97]	rs3748816	1.21	7.41E-12	<i>MMEL1, TNFSFR14</i>	
1q31	x			[114]	rs12134279	1.34	2.06E-14	<i>DENND1B</i>	
2q12	x			[98]	rs12712133	1.14	5.19E-09	<i>IL1RL1, IL1RL2,</i>	Also known as ST2. Antigen-presenting cell receptor for IL-33. Promotes type 2 immune responses [127]
2q13		x		[117]	rs6720394	1.6	4.10E-08	<i>BCL2L11</i>	Also known as BIM; may promote apoptosis and inhibit autophagy [128]
2q32	x			[114]	rs10931468	1.5	2.35E-19	<i>STAT4, STAT1</i>	See Fig. 4
2q33		x		[97]	rs7426056	1.3	1.89E-20	<i>CD28</i>	Receptor that supplies co-stimulatory signals to T cells; See Fig. 3 [129]
2q36	x			[98]	rs4973341	1.22	2.34E-10	<i>CCL20</i>	Chemo-attractant to lymphocytes and neutrophils; strongly expressed in the liver [130]
2q37		x		[120]	rs4676410	1.38	2.43E-09	<i>GPR35</i>	Also known as CXCR8. Receptor for CXCL17 involved in mucosal leucocyte recruitment, including monocytes-macrophages [131]
3p21		x		[97]	rs3197999	1.33	2.45E-26	<i>MST1</i>	Involved in TNF induced cell death, apoptosis, tumor suppressing in liver [132]
3p24	x			[114]	rs1372072	1.2	2.28E-08	<i>PLCL2</i>	Transcription factor involved in B cell proliferation
3q13	x			[96]	rs2293370	1.39	6.84E-16	<i>CD80</i>	Co-stimulatory molecule – which together with CD86 – is important for T cell activation via CD28 [129]
3q25	x			[96]	rs2366643	1.35	3.92E-22	<i>IL12A</i>	See Fig. 4
4p16	x			[98]	rs11724804	1.22	9.01E-12	<i>DGKQ</i>	
4q24	x			[114]	rs7665090	1.26	8.48E-14	<i>NFKB1</i>	The transcription factor NFκB plays important roles in both the innate and adaptive immune system [133,134], influencing both B and T cell activation downstream of the antigen receptors and promoting inflammatory responses
4q27		x		[97]	rs13140464	1.3	8.87E-13	<i>IL2, IL21</i>	IL2: see figure
5p13	x			[96]	rs6871748	1.3	2.26E-13	<i>IL7R</i>	Important in lymphocyte development; loss of function mutations result in lymphopaenia and an autoimmune pre-disposition (product also known as CD127)
5q21	x			[98]	rs526231	1.15	1.14E-08	<i>C5orf30</i>	
5q33	x			[98]	rs2546890	1.15	1.06E-10	<i>IL12B, LOC285626</i>	See Fig. 4
6q15		x		[97]	rs56258221	1.23	8.36E-12	<i>BACH2</i>	Regulatory protein with effects on apoptosis and key to antibody class switching [135]; represses effector lines of CD4 <sup>+</sup> T cells [136]
6q23	x			[98]	rs6933404	1.18	1.27E-10	<i>OLIG3, TNFAIP3</i>	TNFAIP3: zinc finger protein which modulates NFκB and STAT pathways, and TNF receptors signaling [137]
7p14	x			[114]	rs6974491	1.25	4.44E-08	<i>ELMO1</i>	
7q32	x			[96]	rs35188261	1.52	6.52E-22	<i>IRF5</i>	Transcription factor mediating effects of TLR ligation
9p32	x			[39]	rs4979462	1.57	1.85E-14	<i>TNFSF15</i>	
10p15		x		[97]	rs4147359	1.24	8.19E-17	<i>IL2RA</i>	Product also known as CD25. See Fig. 3
11q23	x			[39]	rs4938534	1.38	3.27E-08	<i>POU2AF1</i>	AKA Oct binding factor 1 (OBF1).A transcription factor involved in B cell development [138].
11q13	x			[114]	rs538147	1.23	2.06E-10	<i>RPS6KA4</i>	
11q23		x		[97]	rs7937682	1.17	3.17E-09	<i>SIK2</i>	Modulation of inflammatory cytokine production by murine macrophages [139]
11q23	x			[96]	rs80065107	1.39	7.20E-16	<i>CXCR5, DDX6</i>	CXCR5: Major role in B cell and T follicular helper cell trafficking to germinal centres.
12p13	x			[96]	rs1800693	1.27	1.18E-14	<i>TNFRSF1A, LTBR</i>	TNFRSF1A encodes a member of the tumor necrosis factor family of receptors. It is predominantly expressed on antigen-presenting cells and represents a major receptor for tumor necrosis factor alpha (TNFα). Activation of this receptor can cause apoptosis through activation of NFκB and mutations leading to its constitutive activation are associated with periodic fever syndrome [140].
12q13		x		[97]	rs11168249	1.15	5.49E-09	<i>HDAC7</i>	Histone deacetylase implicated in the suppression of myeloid lineage genes in lymphoid cells [141]
12q24		x		[97]	rs3184504	1.18	5.91E-11	<i>SH2B3, ATXN2</i>	SH2B3: Also known as 'Lnk'. Adaptor protein involved in multiple cell surface signaling pathways in both hematopoietic and non hematopoietic cells. Mutations are implicated in myeloproliferative conditions including malignancies [142].
12q24	x			[96]	rs11065979	1.2	2.87E-09	<i>SH2B3</i>	
12q24			x	[36]	rs3184504	1.4	7.70E-08	<i>SH2B3*</i>	
13q14	x			[96]	rs3862738	1.33	2.18E-08	<i>TNFSF11</i>	AKA RANK-ligand or TNF-related activation-induced cytokine (TRANCE). TNFR ligand involved in control of osteoclast activity, and dendritic cell survival factor. Deficiency results in impaired lymphocyte differentiation [143].
14q24	x			[96]	rs911263	1.26	9.95E-11	<i>RAD51B</i>	
16p13	x			[96]	rs12708715	1.29	2.19E-13	<i>CLEC16A, SOCS1</i>	SOCS1: has a regulatory effect on both NFκB and JAK-STAT pathways including STAT4 [134]
16q24	x			[114]	rs11117432	1.31	4.66E-11	<i>IRF8</i>	Transcription factor mediating effects of TLR ligation.
14q32	x			[114]	rs8017161	1.22	2.61E-13	<i>TNFAIP2</i>	
17q12	x			[96]	rs17564829	1.26	6.05E-14	<i>IKZF3</i>	hematopoietic transcription factors and is involved in lymphocyte development and proliferation, especially in B cells. Dysfunction causes generalized autoimmunity [144]
17q21	x			[96]	rs17564829	1.25	2.15E-09	<i>MAPT</i>	
18q21		x		[120]	rs1452787	1.33	2.61E-08	<i>TCF4</i>	Transcription factor implicated in lymphoid development and epithelial –mesenchymal transition [145]
18q22		x		[97]	rs1788097	1.15	3.06E-08	<i>CD226</i>	Adhesion molecule expressed on NK cells, T cells, platelets and monocytes; modulator of NK cytotoxicity [146]

Table 4 (continued)

Locus	PBC	PSC	AIH	Study	SNV	OR	P-value	Candidate gene(s)	Selected functional notes
19p12	x			[96]	rs34536443	1.91	1.23E-12	TYK2	Signaling kinase in IL-12 and IL-23 receptor signaling. In systemic lupus erythematosus variants to influence IFN $\gamma$ production [147].
1p31	x			[96]	rs72678531	1.61	2.47E-38	IL12RB2	See Fig. 4
19q13		x		[97]	rs60652743	1.25	6.51E-10	FUT2, PRKD2, STRN4	FUT2: Antigen expression – see main text
19q13	x			[115]	rs3745516	1.46	7.97E-11	SPB	Transcription factor mediating B cell receptor signaling
21q22		x		[97]	rs2836883	1.28	3.19E-17	PSMG1	Embryonically lethal if functionally absent in mice; deficient hepatocytes demonstrated premature senescence [148]
22q13	x			[96]	rs2267407	1.29	1.29E-13	SYNGR1	

Lowest p value shown where different studies have implicated the same variant. All candidate genes identified here have association signals with  $p < 5 \times 10^{-8}$  with the exception of the SH2B3 signal in AIH (see text). Not all candidate genes represented for some loci.

example, in the autoimmunity-prone NOD mouse typically develops insulinitis and diabetes, but altering its HLA alleles can redirect the autoimmune focus to thyroiditis [43]. Of note however, the haplotypes associated with each of the hepatic autoimmune diseases are variable between geographic and ethnic populations (Table 3). It remains unclear as to whether the overlapping gene variants contained within these haplotypes of the highly variable HLA region represent true divergent developments resulting in a risk for a similar phenotype, or whether the risk will in due course be narrowed down to very few SNVs as it has in rheumatoid arthritis [44].

## 6.2. Non-HLA associations

Multiple non-HLA associations have now been identified in both PBC and PSC by GWAS. These are summarized with brief notes as to the function of better-understood implicated genes in Table 4. Below, we consider associations forming patterns within pathways and of associations with genes of particular interest. Notably biological studies are needed to complete the understanding of proposed gene associations.

### 6.2.1. The IL-2 pathway in PSC

The combination of loci associated with PSC risk containing CD28, IL2 and IL2RA – which encodes the alpha subunit of the high affinity IL-2 receptor – suggest that the T lymphocyte focused IL-2 pathway has a major role. The multiple effects of IL-2 are summarized as Fig. 3 and the molecule has key roles in the development and maintenance of regulatory T cells but is also important in the adaptive effector response.

The IL2RA deficient mouse develops marked lymphadenopathy with ultimately fatal autoimmunity with lymphocytic infiltrates in skin, intestine, lung and liver in the context of a complete deficiency of regulatory T cells. The liver infiltrate is peri-portal but occurs without overt fibrosis and, owing to its development of anti-mitochondrial antibodies, the mouse has in fact been described as a model of PBC [45]. Indeed, a single case of a boy with a homozygous mutation in IL2RA developing a PBC-like disorder with Treg deficiency has been described [46]. By analogy to IL2RA mutations, mice deficient in the leucocyte anti-apoptotic protein Bcl2l11 also develop multi-system autoimmunity with lymphocytic infiltration and autoantibodies, but without a hepatic focus in a pattern that has been described as more like systemic lupus erythematosus [47].

### 6.2.2. The IL12-STAT4 pathway in PBC

PBC risk has been attributed to variants in a number of genes involved in the T cell activation pathway involving IL-12 and STAT4, which leads to Th1 T cell polarization (Fig. 4). IL-12 is a heterodimeric molecule made up of the two subunits, p35 and p40, encoded by the IL12A and IL12B genes respectively. The latter

protein also heterodimerizes with IL-23p19 to form IL-23, a key signaling component in the Th-17 pathway. The IL-12 receptor is also encoded by two genes: IL12RB1, which is constitutively expressed, and IL12RB2 which is upregulated by interferon- $\gamma$  (IFN $\gamma$ ) to act as a positive feedback loop in antigenic stimulation. STAT4 and tyrosine kinase 2 (TYK2) protein are important in both IL-12 and IL-23 receptor signaling. Each of IL12A, IL12RB2, STAT4 and TYK2 have risk associations with PBC. Deficiencies in IL12 and STAT4 confer susceptibility to intracellular pathogens such as tuberculosis [48,49], whilst STAT4 is required for developing autoimmune diabetes in a mouse model [50].

Using animal models of PBC, efforts have been made to see whether interruption of the IL-12 pathway ameliorates autoimmune liver disease. Results have been variable and all depended on models of PBC created by introducing profound defects in T cell regulatory pathways: deletion of the IL-12p40 subunit worsens peri-biliary inflammation and upregulates hepatic expression of gene products associated with fibrosis in IL2R $\alpha$  deficient mice [51]; in dnTGF $\beta$ R11 mice, IL-12p40 deletion greatly reduced peri-biliary inflammation [52]; whereas IL-12p35 delayed but did not prevent disease in dnTGF $\beta$ R11 mice and was also associated with more marked fibrosis [53].


### 6.2.3. FUT2

The linkage of the gene FUT2 to PSC risk is of interest because it resonates with two other major potential etiological considerations in the disease: the gut microbiome and antigen presentation. FUT2 encodes fucosyl transferase 2 and is involved in the regulation of the expression of antigens including the ABO system and in the synthesis of the H antigen oligosaccharide. The latter acts a binding moiety for a number of bacteria and viruses including the Norwalk virus and also a source of carbon for certain bacteria. In those homozygous for the disease-associated SNV demonstrate alterations in gut microbial community composition [54], FUT2 variants appear to affect carbohydrate metabolism in the gut, so affecting the microbiome. Such alterations, and FUT2 variations, have been associated with sub-clinical colitis in both mice and man [55]. FUT2 variants are also recognized as conferring risk in inflammatory bowel disease.

### 6.2.4. SH2B3 – Lnk

SH2B3 – also known as Lnk – is notable for being associated with all three of the hepatic autoimmune diseases. The association signal in the single AIH GWAS to date falls slightly short of the generally-accepted level for genome-wide significance, but it has been argued that the high likelihood of SH2B3's involvement given its role in so many other autoimmune diseases means that it should be considered a significant association. SH2B3 encodes one a number of SH2B adaptor proteins and is within a widely shared autoimmune locus. It is involved in multiple growth factor and





**Fig. 2.** Many genetic associations of PBC, PSC and AIH are shared with other autoimmune diseases. Circos plot [101] of gene variants associated with PBC, PSC and AIH alongside other autoimmune conditions in high-throughput genetics studies and their meta-analyses. Note that significant pleiotropy exists with some loci implicated in multiple conditions and variable HLA associations a universal feature. PBC = primary biliary cirrhosis; PSC = primary sclerosing cholangitis; AIH = autoimmune hepatitis; AA = alopecia areata; AS = ankylosing spondylitis; ATD = autoimmune thyroid disease; JIA = juvenile idiopathic arthritis; MG = myasthenia gravis; MS = multiple sclerosis; Narc = narcolepsy; Pso = psoriasis; RA = rheumatoid arthritis; Scl = scleroderma; Sjö = Sjögren's syndrome; SLE = systemic lupus erythematosus; T1DM = type 1 diabetes mellitus; UC = ulcerative colitis; Vit = vitiligo. Only validated associations at  $p < 5 \times 10^{-8}$  are included (with the exception of *SH2B3* in AIH – see text); supporting citations are available at request.


cytokine signaling pathways and is a negative regulator of T cell activation, tumor necrosis factor and Janus kinase 2 and 3 (JAK2/3) signaling and is required for normal hematopoiesis. Mice deficient in *SH2B3* have greater levels of activated T cells and a tendency to autoimmunity [56].

*SH2B3* variants are also implicated in celiac disease. Investigation of the functional effect of *SH2B3* genotype in response to lipopolysaccharide and muramyl dipeptide has demonstrated that the risk that carriers of the *SH2B3* rs3184504\*A risk allele – which is the same as that implicated in PSC and AIH – show stronger activation of the NOD2 pathogen recognition pathway [57]. This suggests that *SH2B3* has a role in modulating the immune response

to intestinal pathogens.

#### 6.2.5. Epistasis

Epistasis is the process by which carrying a combination of two or more risk variants has a more than additive effect on total risk. One example of such interactions in PBC is the risk-conferring epistatic interaction between the 1p31 (*IL12RB2*) and 7q32 (*IRF5*) loci [58]. A potential risk-amplifying interaction between *CTLA4* and *TNF $\alpha$*  variants has also been reported in the pre-GWAS era, but not replicated [59]. Numbers of epistatic gene–gene interactions are likely to grow as sample sizes increase and meta-analysis is carried out.



**Fig. 3. Schematic representation of the IL-2 pathway, which is implicated in PSC by GWAS.** The combination of TCR and CD28 signals upregulate IL-2 production in activated T cells, especially CD4<sup>+</sup> T cells. T cell activation without co-stimulatory CD28 signal results in anergy with minimal IL-2 production. In general, IL-2 has pro-effector actions. However, in IL-2 deficiency or deficiency of IL-2 receptor alpha, there is widespread autoimmunity and lymphoproliferation ascribed to a failure to generate regulatory T cells. In developed immune systems however, exogenous IL-2 typically amplifies T cell responses. IL-2 receptor complex may exist as a dimer (from IL-2R $\beta$  and/or IL-2R $\gamma$ ) or trimer (with IL-2R $\alpha$ ). The addition of IL-2R $\alpha$  renders the receptor complex with much higher affinity to IL-2, and is particularly important for regulatory T cells. Cytokines secretion is shown as dashed arrows; positive effects are shown as solid arrows. Labels in red denote factors implicated by GWAS to date. Abbreviations: APC = antigen-presenting cell; CD = cluster of differentiation; IL = interleukin; AICD = activation-induced cell death; IL2R = IL-2 receptor; IFN $\gamma$  = interferon gamma; TNF $\alpha$  = tumor necrosis factor alpha; Treg = regulatory T cell [102–105]. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

## 7. Translating GWAS findings to clinical cohorts

Translating findings from GWAS studies in general to changes of clinical care has proved challenging. Notable exceptions exist however, for example the utility of IL-12 and IL-23 blockade in psoriasis [60] and the identification of the link between *IL28B* variants and sustained virological response in hepatitis C treatment [61].

### 7.1. HLA

There has been some success in predicting clinical characteristics from GWAS data, but prospective validation is required. For example, work arising from the Dutch AIH GWAS has suggested that HLA types influence clinical characteristics: HLA-DRB1\*03:01 being associated with higher serum gamma immunoglobulin levels and HLA-DRB1\*04:01 being associated with an older age of first presentation [62].

A number of pre-GWAS studies have related clinical characteristics to HLA haplotypes, although many have not been replicated. The well-established inter-population variability between HLA associations means that results must be applied between varying populations with caution. In PBC for example, HLA haplotypes correlate with anti-gp210 status in Japanese [63] and Chinese [64] patients. The presence of anti-gp210 antibodies is independently predictive of outcome. HLA-associated susceptibility to UC is different with and without the presence of concurrent PSC [65] and acute rejection after transplantation may also be associated with


HLA-B\*08, HLA-C\*07 and HLA-DRB1\*03 [66]. In AIH, HLA-DR4 may predict treatment failure [67]. Sequencing of DRB1\*03-DRB1\*04 amongst North American patients has shown a lysine-rich subset in which carriers with AIH more frequently develop cirrhosis and a requirement for liver transplantation [68]. Similarly, amongst Japanese with AIH, the DRB1\*08:03-DQB1\*06:01 haplotype was more frequent in patients who developed hepatic failure [69].

The mechanism by which given HLA variants may exert their effects in hepatic autoimmunity is unclear. However, in PBC recent important work has directly linked HLA-types to cellular phenotype. T cells with the risk-conferring HLA-DRB1\*08:01 shown a high-affinity response to specific pyruvate dehydrogenase E2 subunit peptides; an analogous response was absent in cells expressing the protective HLA-DRB1\*11:01 [70].

### 7.2. Non-HLA

The implication of the IL-2 pathway in PSC might logically suggest a role for immunosuppressants. However, no immunosuppressive regime, including those with particular activity against IL-2 has proven efficacy in PSC [71]. Similarly, despite a number of molecules implicated in activation of the adaptive immune system in PBC, no immunosuppressive therapy has been shown effective to date.

The alpha-subunit of the IL2 receptor complex (*IL2RA*; CD25) is also implicated by GWAS in PSC. This had led to particular interest in regulatory T cells, which strongly express IL2R $\alpha$ . Recent work attempting to address the role of regulatory T cells in PSC has



**Fig. 4. Schematic representation of the IL-12 pathway, which is implicated in PBC by GWAS.** Antigen activates APC through TLR. APC in turn produce IL-12 after phosphorylation of IRF5. Antigen is presented to CD4<sup>+</sup> T cells by HLA II with co-stimulation via CD80 and 86 to CD28. There is competitive inhibition of this co-stimulation by CTLA4. IL-12 activates a cascade of signaling factors including NFKB and STAT4 to promote the production of Th1-type cytokines including TNF $\alpha$  and IFN $\gamma$ ; the transcription factor IRF8 is involved. IL7R supports lymphocyte development. There is positive feedback from Th1 cytokines to APCs. Red text denotes confirmed risk associations with PBC. Arrows denote positive effects; barred lines denote negative effects. APC = antigen-presenting cell; CD = cluster of differentiation; CTLA4 = Cytotoxic T lymphocyte antigen 4; HLA II = human leucocyte antigen class II; IFN- $\gamma$  = interferon- $\gamma$ ; IFN $\gamma$ R = interferon- $\gamma$  receptor; IL-12 = interleukin-12; IL-12R $\beta$ 1/2 = IL-12 receptor  $\beta$  subunits 1 and 2; IL7R = interleukin-7 receptor; IRF5 and IRF8 = interferon response factors 5 and 8; JAK2 = Janus kinase 2; Lck = lymphocyte-specific protein tyrosine kinase; NFKB = nuclear factor kappa-light-chain-enhancer of activated B cells; PKC = protein kinase C; SOCS1 = suppressor of cytokine signaling 1; STAT4 = signal transducer and activator of transcription 4; TCR = T-cell receptor; TLR = Toll-like receptor; TNFAIP3 = tumor necrosis factor alpha-induced protein 3; TNFRSF1a = Tumor necrosis factor receptor superfamily 1a; TNF $\alpha$  = tumor necrosis factor alpha; TYK2 = Tyrosine kinase 2 [106,107]. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

suggested a reduction in their number in PSC livers as compared with PBC [72]. Further, peripheral populations were reduced and this correlated with homozygosity for the risk *IL2RA* allele.

The proportion of liver-infiltrating lymphocytes expressing CD28 in PSC is reduced and these cells have a more activated phenotype and produce more of the effector cytokines tumor necrosis factor  $\alpha$  and interferon- $\gamma$  than their CD28<sup>+</sup> counterparts [73]. Interestingly, the CD28-negative T cell compartment is also expanded in rheumatoid arthritis, in which the CD28 gene is similarly implicated by GWAS [74].

In follow-up studies of GWAS datasets, associations of *STAT4* variations with ANA status have been identified in Japanese PBC

patients [75] and *TNFSF15* polymorphisms have been linked to functional changes in the gene product. Differences in mRNA and protein expression were also seen in both PBC and healthy subjects carrying this variant, although a link to outcome or phenotype remains unclear [76].

The risk association with *CXCR5* is of particular interest because of *CXCR5*'s major role in lymphocyte trafficking, especially of B and T cells migrating to germinal centers. Subsequent to GWAS, two observations have strengthened the link to the clinical picture: firstly that numbers of *CXCR5*<sup>+</sup> T follicular helper cells are increased both peripherally and in PBC liver tissue [77,78] and secondly that the ligand for *CXCR5*, *CXCL13*, is upregulated in PBC as compared

with other liver diseases [79].

A common feature of PBC, PSC and AIH is that they all recur after transplantation. This supports a key role for the immune system rather than an intrinsic biliary epithelial cell or hepatocyte defect. In PBC, data gathered as part of the UK-PBC cohort was re-analyzed according to individuals' *IL12A* variant status and this has suggested a correlation between the AG variant and decreased time to diagnosis of PBC recurrence [80].

The *IL12* pathway is also the target of the only GWAS-derived therapeutic intervention to date in hepatic autoimmunity. This study used ustekinumab, a monoclonal antibody that blocks *IL-12* and *IL-23* signaling. However, administration to PBC patients with established cholestatic, ursodeoxycholic-unresponsive disease did not significantly alter biochemical surrogates of disease activity [81]. However intriguingly in those patients in whom there was a fall in alkaline phosphatase noted, there was evidence of modulation of biological pathways (including Th17 lymphocytes) related to ustekinumab response.

Finally, work has been published attempting to calculate risk using aggregated GWAS data and applying these to cohorts. There has been some success, with a recent risk score generating a modest AuRoC of 0.72. This is despite the same authors calculating a combinatorial disease risk from all extant PBC susceptibility alleles of only 5.3% of the observed heritable risk, suggesting the capacity for future refinement [82].

## 8. Deficiencies in GWAS to date

### 8.1. GWAS results do not account for all the observed risk

One reason for this is that these rare diseases are likely to have some of their risk conferred by rare genetic variants. The design of GWAS means that identified risk variants are, by definition, more frequent in the population than is the disease *per se* [83]. Further, rarer genes with smaller effect sizes require larger cohorts for association signals to reach the stringent levels of statistical significance: this effect has been especially apparent in more common autoimmune diseases. For example, there is an almost direct correlation between study size and the number of variants identified in inflammatory bowel disease [37] and studies of super-cohorts involving over 100,000 subjects have greatly clarified the genetic landscape of rheumatoid arthritis [84].

Importantly, it is apparent that GWAS will never account for all of the observed risk in complex diseases like those in hepatic autoimmunity: in PSC it has been commented that “more than 50% of the susceptibility to these diseases is still likely to be of an environmental origin.” [85] Equally, in PBC, which has seen the greatest amount of GWAS investigation, calculation suggests that findings to date only account for a few percent of the observed heritability [82].

### 8.2. Genetic information cannot yet explain key aspects of disease phenotype

As yet, GWAS does not explain key phenotypic features of the hepatic autoimmune diseases. Examples of these include the marked differences in risk between genders, the co-incidence of other autoimmune phenomena, treatment response in PBC and AIH, and malignancy risk in PSC. Genetic risk also does not help explain the evolution of clinical phenotype that is particularly recognized in pediatric patients presenting with autoimmune hepatitis and sclerosing cholangitis.

### 8.3. GWAS does not reflect disease risk in varied ethnicities

As mentioned above, all major GWAS in hepatic autoimmunity to date have focused on populations of European Caucasian descent, with the exception of the Japanese PBC study [39]. The Japanese GWAS identified two associations not yet replicated in other populations. Similarly, a recent replication study from China in a Han population has suggested incomplete overlap with previous studies [40]. Observations like the lower rate of concurrent ulcerative colitis in African American PSC patients suggest genetic variation may be implicated [86]. The selection of SNVs assessed by ImmunoChip are also biased to white Caucasian populations [87]. It therefore remains possible that associations in those of non-European backgrounds will yield different associations.

### 8.4. Positive selection pressure for genes implicated by GWAS is not yet explained

Several associations uncovered in GWAS have putative links to resistance to microbial infection: *IL-12/STAT4* pathway and intracellular/tubercular infection in PBC; *FUT2* and Norwalk virus in PSC; *SH2B3* and response via *NOD2* in AIH, PSC and PBC [49,54,57]. However, for the most variants associated with hepatic autoimmunity no explanation exists for the selection force that explains their presence in the population. It is likely that responses to microbial agents will be implicated: most of human variation is thought attributable to infectious disease [88] and gut pathogens are thought to have dictated the genetic signature of inflammatory bowel disease [89].

### 8.5. GWAS does not yet map to animal models nor therapies

As considered above, there is a current lack of either a direct animal model in which abnormalities in a gene identified by GWAS recapitulate the key phenotypic components of one of the major autoimmune hepatic disease. Equally, no targeted therapy based on GWAS findings has yet been successful.

### 8.6. GWAS may not capture epigenetic modifications, including heritable epigenetic traits

Epigenetic modifications are alterations to genes other than in their base-pair code that alter how, or whether, they are transcribed. They may be the result of environmental influences or a consequence of the activity of other genes. They may be inherited. However, epigenetic modifications are not captured by some current chip-based GWAS techniques nor sequencing technologies. Importantly, a high proportion of fine-mapped variants are seen in promoter regions or where regulatory proteins bind, and epigenetic variation has been identified in non-hepatic autoimmunity [90].

One epigenetic mechanism that has been proposed as explaining some of the gender differences in prevalence of PBC is biased X or Y chromosome inactivation. This process typically involves silencing by chromosome packaging in heterochromatin and has been reported as being biased in a number of autoimmune diseases [91,92]. However, it is notable that to date, no risk genes for hepatic autoimmunity have been identified on either the X or Y chromosome.

In PBC further interest a role for epigenetic mechanisms has been given further support by two recent studies. First is the observation that the promoter region of the gene encoding CD40 ligand (*CD40L*) demonstrates reduced methylation in PBC patients as compared with controls. This is of particular interest given the key role CD40-CD40 ligand interactions play in B cells activation and because this observation correlates with serum IgM [93].

Second, recent reports have demonstrated differential DNA methylation of the promoter region of the gene *CXCR3* in T lymphocytes from patients with PBC [94]. Demethylation of *CXCR3* correlated with increased protein expression of *CXCR3*. Functionally, *CXCR3* is present in many inflamed tissues and is associated with the generation of Th1 type responses and the subsequent migration of Th1 CD4<sup>+</sup> T cells. Neither *CXCR3* nor *CD40L* has yet been associated with risk for developing PBC.

## 9. The future of genetic investigation of hepatic autoimmunity

Future investigation of the genetics of hepatic autoimmunity is likely to take advantage of the increasing efficiency and reducing cost of finer-resolution SNV chips and sparse or whole genome sequencing. Such “post-GWAS” strategies are starting to reveal new risk loci in other conditions [95]. Moves towards higher-resolution assessment of the genome that stop short of full sequencing may also yield extra information as suggested by ImmunoChip studies [96,97] (with their higher resolution coverage of targeted subsections of the genome) and recent work in other autoimmune conditions [90]. In addition, improved analyses of multiple SNVs associated with risk loci have improved our ability to identify specific risk-causing variants/mutations. Combining such information with knowledge of the epigenome of individual cell types has begun to direct researchers to highlight where individual variants may have particular effects [90]. For example, in PBC candidate SNVs are enriched in CD4<sup>+</sup> T cells and B cells. Larger cohorts of patients will be valuable but may prove challenging to recruit and meta-analysis including smaller new cohorts as they become available is likely to continue to provide small incremental returns [98]. In addition, where patient – and importantly control – cohorts are well phenotyped there is the potential of linking genetic data to clinical outcomes and disease subtypes. Indeed, recent work has taken data on HLA haplotype associations from GWAS and associated them with presentation and outcome in AIH [62]. The three carefully phenotyped cohort projects known as UK-PBC, UK-PSC and UK-AIH are examples of current work that may provide insights in this direction.

Future genetic work will need to proceed hand-in-hand with closer examination of environmental influences, epigenetics and gene–environment interactions and will need to consider genomes other than in Caucasian populations.

## 10. Conclusion

In each of PBC, PSC and AIH, there is a clear genetic component to disease risk. Each also has a significant environmental input and it is uncertain as to the relative contribution of genes, environment and interactions within and between these influences. Until recently, although several risk-bearing HLA haplotypes in these diseases had been identified, these were insufficient to explain much disease heritability. GWAS and analogous focused chip-based studies have now greatly increased our knowledge of a number of variants associated with risk for PBC and PSC and begun to inform our knowledge in AIH. These variants have been mapped to a number of likely candidate genes. Of particular interest are links to the IL-12-STAT4 pathway in PBC and the CD28-IL2 pathway in PSC. However, despite the multiple genetic associations described by genetic studies to date, the observed genetic component to disease risk in hepatic autoimmunity is far from explained. Possible reasons for this include relatively small sample sizes, inherent deficiencies in GWAS methodology in detecting rare variants, and the uncertain extent of environmental and epigenetic influences. Further, GWAS does not yet explain a number of key phenotypic features of these

diseases, nor have results yet mapped to definite clinical insights as they have in other autoimmune diseases. These problems are likely to be addressed by larger, sequencing-based or higher-resolution SNV-based studies alongside investigation of gene–environment interactions. Such efforts are starting to gain leverage in more common diseases. Finally, it is a great pleasure to contribute this paper to the special issue devoted to Diego and Giorgina Vergani, both of whom have excelled in every aspect of clinical medicine, from patient care to teaching and, of course, research. This paper is part of this dedicated issue which is a component of the yearly efforts of the Journal of Autoimmunity to recognize truly outstanding figures in autoimmunology. Previous recipients have included Abul Abbas, Michael Sela, Ruth Arnon, Noel Rose and Ian Mackay [99,100]. Giorgina and Diego are fitting to include in this group.

## Conflict of interest

None declared.

## Acknowledgments

GJW is a UK Medical Research Council Clinical Research Training Fellow. GMH is a co-investigator for UK-PBC ([www.uk-pbc.com](http://www.uk-pbc.com)) supported by a Stratified Medicine Award from the UK Medical Research Council, and Principal Investigator for UK-PSC ([www.uk-psc.com](http://www.uk-psc.com)), an NIHR Rare Disease Translational Collaboration.

The views expressed are those of the authors(s) and not necessarily those of the National Health System, the National Institute for Health Research, or the Department of Health.

## References

- [1] A. Davidson, B. Diamond, Autoimmune diseases, *N. Engl. J. Med.* 345 (2001) 340–350.
- [2] A. Ileo, P. Invernizzi, B. Gao, M. Podda, M.E. Gershwin, Definition of human autoimmunity — autoantibodies versus autoimmune disease, *Autoimmun. Rev.* 9 (2010) A259–A266.
- [3] G.S. Cooper, M.L.K. Bynum, E.C. Somers, Recent insights in the epidemiology of autoimmune diseases: improved prevalence estimates and understanding of clustering of diseases, *J. Autoimmun.* 33 (2009) 197–207.
- [4] E.C. Somers, S.L. Thomas, L. Smeeth, A.J. Hall, Are individuals with an autoimmune disease at higher risk of a second autoimmune disorder? *Am. J. Epidemiol.* 169 (2009) 749–755.
- [5] G.M. Hirschfield, M.E. Gershwin, The immunobiology and Pathophysiology of Primary Biliary Cirrhosis, *Annu. Rev. Pathol. Mech. Dis.* 8 (2013) 303–330.
- [6] M. Corrigan, G.M. Hirschfield, Y.H. Oo, D.H. Adams, Autoimmune hepatitis: an approach to disease understanding and management, *Br. Med. Bull.* 114 (2015) 181–191.
- [7] G.M. Hirschfield, T.H. Karlsen, K.D. Lindor, D.H. Adams, Primary sclerosing cholangitis, *Lancet* 382 (2013) 1587–1599.
- [8] K. Boonstra, U. Beuers, C.Y. Ponsioen, Epidemiology of primary sclerosing cholangitis and primary biliary cirrhosis: A systematic review, *J. Hepatol.* 56 (2012) 1181–1188.
- [9] L. Gronbaek, H. Vilstrup, P. Jepsen, Autoimmune hepatitis in Denmark: incidence, prevalence, prognosis, and causes of death. A nationwide registry-based cohort study, *J. Hepatol.* 60 (2014) 612–617.
- [10] A. Marson, W.J. Housley, D.A. Hafler, Genetic basis of autoimmunity, *J. Clin. Invest.* 125 (2015) 2234–2241.
- [11] M.I. McCarthy, G.R. Abecasis, L.R. Cardon, D.B. Goldstein, J. Little, J.P. Ioannidis, et al., Genome-wide association studies for complex traits: consensus, uncertainty and challenges, *Nat. Rev. Genet.* 9 (2008) 356–369.
- [12] C. Selmi, M.J. Mayo, N. Bach, H. Ishibashi, P. Invernizzi, R.G. Gish, et al., Primary biliary cirrhosis in monozygotic and dizygotic twins: genetics, epigenetics, and environment, *YGASt* 127 (2004) 485–492.
- [13] N.M. van Gerven, B.J. Verwer, B.I. Witte, K.J. van Erpecum, H.R. van Buuren, I. Majers, et al., Epidemiology and clinical characteristics of autoimmune hepatitis in the Netherlands, *Scand. J. Gastroenterol.* 49 (2014) 1245–1254.
- [14] D.E. Jones, F.E. Watt, J.V. Metcalf, M.F. Bassendine, O.F. James, Familial primary biliary cirrhosis reassessed: a geographically-based population study, *J. Hepatol.* 30 (1999) 402–407.
- [15] M.E. Gershwin, C. Selmi, H.J. Worman, E.B. Gold, M. Watnik, J. Utts, et al., Risk factors and comorbidities in primary biliary cirrhosis: a controlled interview-based study of 1032 patients, *Hepatology* 42 (2005) 1194–1202.
- [16] A. Bergquist, S.M. Montgomery, S. Bahmanyar, R. Olsson, A. Danielsson,

- S. Lindgren, et al., Increased risk of primary sclerosing cholangitis and ulcerative colitis in first-degree relatives of patients with primary sclerosing cholangitis, *Clin. Gastroenterol. Hepatol. Off. Clin. Pract. J. Am. Gastroenterol. Assoc.* 6 (2008) 939–943.
- [17] D.S. Smyk, D.P. Bogdanos, E.I. Rigopoulou, Risk Factors for Autoimmune Hepatitis: From Genes and Pregnancy to Vaccinations and Pollutants, *Immunogastroenterology* 2 (2013) 29–37.
- [18] K.N. Lazaridis, B.D. Juran, G.M. Boe, J.P. Slusser, M. de Andrade, H.A. Homburger, et al., Increased prevalence of antimitochondrial antibodies in first-degree relatives of patients with primary biliary cirrhosis, *Hepatology* 46 (2007) 785–792.
- [19] A. Mantaka, M. Koulentaki, G. Chlouverakis, J.M. Enele-Melono, A. Darivianaki, M. Tzardi, et al., Primary biliary cirrhosis in a genetically homogeneous population: Disease associations and familial occurrence rates, *BMC Gastroenterol.* 12 (2012) 1.
- [20] C. Selmi, P. Invernizzi, M. Zuin, M. Podda, M.F. Seldin, M.E. Gershwin, Genes and (auto)immunity in primary biliary cirrhosis, *Genes Immun.* 6 (2005) 543–556.
- [21] L. Arbour, R. Rupps, L. Field, P. Ross, A. Erikson, H. Henderson, et al., Characteristics of primary biliary cirrhosis in British Columbia's first nations population, *Can. J. Gastroenterol.* 19 (2005) 305–310.
- [22] J. Delgado, A.D. Sperber, V. Novack, B. Delgado, L. Edelman, N. Gaspar, et al., The epidemiology of primary biliary cirrhosis in southern Israel, *Israel Med. Assoc. J.* 7 (2005) 717–721.
- [23] K. Tsuji, Y. Watanabe, J. Van de Water, T. Nakanishi, G. Kajiyama, A. Parikh-Patel, et al., Familial primary biliary cirrhosis in Hiroshima, *J. Autoimmun.* 13 (1999) 171–178.
- [24] A.W. Lohse, G. Mieli-Vergani, Autoimmune hepatitis, *J. Hepatol.* 55 (2011) 171–182.
- [25] G. Bussone, L. Mouthon, Autoimmune manifestations in primary immune deficiencies, *Autoimmun. Rev.* 8 (2009) 332–336.
- [26] D. Schubert, C. Bode, R. Kenefack, T.Z. Hou, J.B. Wing, A. Kennedy, et al., Autosomal dominant immune dysregulation syndrome in humans with CTLA4 mutations, *Nat. Med.* (2014) 1–9.
- [27] F. Barzaghi, L. Passerini, R. Bacchetta, Immune dysregulation, polyendocrinopathy, enteropathy, X-linked syndrome: a paradigm of immunodeficiency with autoimmunity, *Front. Immunol.* 3 (2012 Jul 31) 211, <http://dx.doi.org/10.3389/fimmu.2012.00211>.
- [28] G. Webb, A. Richter, M. Blahova, S. Curbishley, K.-K. Li, D. Neil, et al., P1196: Autoimmune hepatitis with T-regulatory cell deficiency in the context of GATA-2 dysfunction: a novel single gene association, *J. Hepatol.* 62 (2015) S804.
- [29] K. Boonstra, E.M. de Vries, N. van Geloven, K.J. van Erpecum, M. Spanier, A.C. Poen, et al., Risk factors for primary sclerosing cholangitis, *Liver Int. Off. J. Int. Assoc. Study Liver* (Published online 28 Aug 2015), <http://dx.doi.org/10.1111/liv.12894>.
- [30] K.J. Mitchell, What is complex about complex disorders? *Genome Biol.* 13 (2012) 237.
- [31] T.H. Karlsen, E. Schruppf, K.M. Boberg, Genetic epidemiology of primary sclerosing cholangitis, *World J. Gastroenterol.* 13 (2007) 5421–5431.
- [32] A.J. Czaja, Genetic factors affecting the occurrence, clinical phenotype, and outcome of autoimmune hepatitis, *Clin. Gastroenterol. Hepatol. Off. Clin. Pract. J. Am. Gastroenterol. Assoc.* 6 (2008) 379–388.
- [33] P. Invernizzi, C. Selmi, I.R. Mackay, M. Podda, M.E. Gershwin, From bases to basis: linking genetics to causation in primary biliary cirrhosis, *Clin. Gastroenterol. Hepatol. Off. Clin. Pract. J. Am. Gastroenterol. Assoc.* 3 (2005) 401–410.
- [34] P.M. Visscher, M.A. Brown, M.I. McCarthy, J. Yang, Five years of GWAS discovery, *Am. J. Hum. Genet.* 90 (2012) 7–24.
- [35] P. Sanseau, P. Agarwal, M.R. Barnes, T. Pastinen, J.B. Richards, L.R. Cardon, et al., Use of genome-wide association studies for drug repositioning, *Nat. Biotechnol.* 30 (2012) 317–320.
- [36] Y.S. de Boer, N.M. van Gerven, A. Zwiwers, B.J. Verwer, B. van Hoek, K.J. van Erpecum, et al., Genome-wide association study identifies variants associated with autoimmune hepatitis type 1, *Gastroenterology* 147 (2014), 443–52 e5.
- [37] M. Parkes, A. Cortes, D.A. van Heel, M.A. Brown, Genetic insights into common pathways and complex relationships among immune-mediated diseases, *Nat. Rev. Genet.* 14 (2013) 661–673.
- [38] C.J. Lessard, J.A. Ice, I. Adrianto, G.B. Wiley, J.A. Kelly, P.M. Gaffney, et al., The genomics of autoimmune disease in the era of genome-wide association studies and beyond, *Autoimmun. Rev.* 11 (2012) 267–275.
- [39] M. Nakamura, N. Nishida, M. Kawashima, Y. Aiba, A. Tanaka, M. Yasunami, et al., Genome-wide association study identifies TNFSF15 and POU2AF1 as susceptibility loci for primary biliary cirrhosis in the Japanese population, *Am. J. Hum. Genet.* 91 (2012) 721–728.
- [40] M. Dong, J. Li, R. Tang, P. Zhu, F. Qiu, C. Wang, et al., Multiple genetic variants associated with primary biliary cirrhosis in a Han Chinese population, *Clin. Rev. Allergy Immunol.* 48 (2–3) (2015 Jun) 316–321, <http://dx.doi.org/10.1007/s12016-015-8472-0>.
- [41] P.T. Illing, J.P. Vivian, N.L. Dudek, L. Kostenko, Z. Chen, M. Bharadwaj, et al., Immune self-reactivity triggered by drug-modified HLA-peptide repertoire, *Nature* 486 (2012) 554–558.
- [42] P. Invernizzi, M. Ransom, S. Raychaudhuri, R. Kosoy, A. Lleo, R. Shigeta, et al., Classical HLA-DRB1 and DPB1 alleles account for HLA associations with primary biliary cirrhosis, *Genes Immun.* 13 (2012) 461–468.
- [43] H. Braley-Mullen, G.C. Sharp, B. Medling, H. Tang, Spontaneous autoimmune thyroiditis in NOD.H-2h4 mice, *J. Autoimmun.* 12 (1999) 157–165.
- [44] S. Raychaudhuri, C. Sandor, E.A. Stahl, J. Freudenberg, H.S. Lee, X. Jia, et al., Five amino acids in three HLA proteins explain most of the association between MHC and seropositive rheumatoid arthritis, *Nat. Genet.* 44 (2012) 291–296.
- [45] K. Wakabayashi, Z.X. Lian, Y. Moritoki, R.Y. Lan, K. Tsuneyama, Y.H. Chuang, et al., IL-2 receptor alpha(-/-) mice and the development of primary biliary cirrhosis, *Hepatology* 44 (2006) 1240–1249.
- [46] C.A. Aoki, C.M. Roifman, Z.-X. Lian, C.L. Bowlus, G.L. Norman, Y. Shoenfeld, et al., IL-2 receptor alpha deficiency and features of primary biliary cirrhosis, *J. Autoimmun.* 2012 (2006) 50–53.
- [47] P. Bouillet, D. Metcalf, D.C. Huang, D.M. Tarlinton, T.W. Kay, F. Kontgen, et al., Proapoptotic Bcl-2 relative Bim required for certain apoptotic responses, leukocyte homeostasis, and to preclude autoimmunity, *Science* 286 (1999) 1735–1738.
- [48] I. Sugawara, H. Yamada, S. Mizuno, Relative importance of STAT4 in murine tuberculosis, *J. Med. Microbiol.* 52 (2003) 29–34.
- [49] C. Picard, C. Fieschi, F. Altare, S. Al-Jumaah, S. Al-Hajjar, J. Feinberg, et al., Inherited interleukin-12 deficiency: IL12B genotype and clinical phenotype of 13 patients from six kindreds, *Am. J. Hum. Genet.* 70 (2002) 336–348.
- [50] Z. Yang, M. Chen, J.D. Ellett, L.B. Fialkow, J.D. Carter, M. McDuffie, et al., Autoimmune diabetes is blocked in Stat4-deficient mice, *J. Autoimmun.* 22 (2004) 191–200.
- [51] Y. Yao, W. Yang, Y.-Q. Yang, H.-D. Ma, F.-T. Lu, L. Li, et al., Distinct from its canonical effects, deletion of IL-12p40 induces cholangitis and fibrosis in interleukin-2R $\alpha$ -/- mice, *J. Autoimmun.* 51 (2014) 99–108.
- [52] K. Yoshida, G.-X. Yang, W. Zhang, M. Tsuda, K. Tsuneyama, Y. Moritoki, et al., Deletion of interleukin-12p40 suppresses autoimmune cholangitis in dominant negative transforming growth factor  $\beta$  receptor type II mice, *Hepatology* 50 (2009) 1494–1500.
- [53] M. Tsuda, W. Zhang, G.X. Yang, K. Tsuneyama, Y. Ando, K. Kawata, et al., Deletion of interleukin (IL)-12p35 induces liver fibrosis in dominant-negative TGF $\beta$  receptor type II mice, *Hepatology* 57 (2013) 806–816.
- [54] P. Rausch, A. Rehman, S. Kunzel, R. Hasler, S.J. Ott, S. Schreiber, et al., Colonic mucosa-associated microbiota is influenced by an interaction of Crohn disease and FUT2 (Secretor) genotype, *Proc. Natl. Acad. Sci. U. S. A.* 108 (2011) 19030–19035.
- [55] M. Tong, I. McHardy, P. Ruegger, M. Goudarzi, P.C. Kashyap, T. Haritunians, et al., Reprogramming of gut microbiome energy metabolism by the FUT2 Crohn's disease risk polymorphism, *ISME J.* 8 (2014) 2193–2206.
- [56] H. Katayama, T. Mori, Y. Seki, M. Anraku, M. Iseki, M. Ikutani, et al., Lnk prevents inflammatory CD8 $^{+}$  T-cell proliferation and contributes to intestinal homeostasis, *Eur. J. Immunol.* 44 (2014) 1622–1632.
- [57] A. Zhernakova, C.C. Elbers, B. Ferwerda, J. Romanos, G. Trynka, P.C. Dubois, et al., Evolutionary and functional analysis of celiac risk loci reveals SH2B3 as a protective factor against bacterial infection, *Am. J. Hum. Genet.* 86 (2010) 970–977.
- [58] B.D. Juran, G.M. Hirschfield, P. Invernizzi, E.J. Atkinson, Y. Li, G. Xie, et al., Immunochip analyses identify a novel risk locus for primary biliary cirrhosis at 13q14, multiple independent associations at four established risk loci and epistasis between 1p31 and 7q32 risk variants, *Hum. Mol. Genet.* 21 (2012) 5209–5221.
- [59] B.D. Juran, E.J. Atkinson, J.J. Larson, E.M. Schlicht, X. Liu, E.J. Heathcote, et al., Carriage of a tumor necrosis factor polymorphism amplifies the cytotoxic T-lymphocyte antigen 4 attributed risk of primary biliary cirrhosis: evidence for a gene-gene interaction, *Hepatology* 52 (2010) 223–229.
- [60] T. Kopp, E. Riedl, C. Bangert, E.P. Bowman, E. Greisenegger, A. Horowitz, et al., Clinical improvement in psoriasis with specific targeting of interleukin-23, *Nature* 521 (2015) 222–226.
- [61] D. Ge, J. Fellay, A.J. Thompson, J.S. Simon, K.V. Shianna, T.J. Urban, et al., Genetic variation in IL28B predicts hepatitis C treatment-induced viral clearance, *Nature* 461 (2009) 399–401.
- [62] N.M.F. van Gerven, Y.S. de Boer, A. Zwiwers, B.J. Verwer, J.P.H. Drenth, B. van Hoek, et al., HLA-DRB1[ast]03:01 and HLA-DRB1[ast]04:01 modify the presentation and outcome in autoimmune hepatitis type-1, *Genes Immun.* 16 (2015) 247–252.
- [63] M. Nakamura, M. Yasunami, H. Kondo, H. Horie, Y. Aiba, A. Komori, et al., Analysis of HLA-DRB1 polymorphisms in Japanese patients with primary biliary cirrhosis (PBC): the HLA-DRB1 polymorphism determines the relative risk of antinuclear antibodies for disease progression in PBC, *Hepatol. Res. Off. J. Jpn. Soc. Hepatol.* 40 (2010) 494–504.
- [64] D.T. Zhao, H.Y. Liao, X. Zhang, Y.M. Liu, Y. Zhao, H.P. Zhang, et al., Human leukocyte antigen alleles and haplotypes and their associations with antinuclear antibodies features in Chinese patients with primary biliary cirrhosis, *Liver Int. Off. J. Int. Assoc. Study Liver* 34 (2014) 220–226.
- [65] T.H. Karlsen, K.M. Boberg, M. Vatn, A. Bergquist, J. Hampe, E. Schruppf, et al., Different HLA class II associations in ulcerative colitis patients with and without primary sclerosing cholangitis, *Genes Immun.* 8 (2007) 275–278.
- [66] B. Fosby, S. Naess, J.R. Hov, J. Traherne, K.M. Boberg, J. Trowsdale, et al., HLA variants related to primary sclerosing cholangitis influence rejection after liver transplantation, *World J. Gastroenterol.* 20 (2014) 3986–4000.
- [67] A.J. Czaja, H.A. Carpenter, S.B. Moore, Clinical and HLA phenotypes of type 1 autoimmune hepatitis in North American patients outside DR3 and DR4,

- Liver Int. Off. J. Int. Assoc. Study Liver 26 (2006) 552–558.
- [68] A.J. Montano-Loza, H.A. Carpenter, A.J. Czaja, Clinical significance of HLA DRB1\*03:01 and DRB1\*04:01 in type 1 autoimmune hepatitis, *Liver Int. Off. J. Int. Assoc. Study Liver* 26 (2006) 1201–1208.
- [69] T. Umemura, S. Joshita, T. Ichijo, K. Yoshizawa, Y. Katsuyama, E. Tanaka, et al., Human leukocyte antigen class II molecules confer both susceptibility and progression in Japanese patients with primary biliary cirrhosis, *Hepatology* 55 (2012) 506–511.
- [70] I.T. Chow, E.A. James, T.J. Gates, V. Tan, A.K. Moustakas, G.K. Papadopoulos, et al., Differential binding of pyruvate dehydrogenase complex-E2 epitopes by DRB1\*08:01 and DRB1\*11:01 is predicted by their structural motifs and correlates with disease risk, *J. Immunol.* 190 (2013) 4516–4524.
- [71] R. Chapman, J. Fevery, A. Kalloo, D.M. Nagorney, K.M. Boberg, B. Shneider, et al., Diagnosis and management of primary sclerosing cholangitis, *Hepatology* 51 (2010) 660–678.
- [72] M. Sebode, M. Peiseler, B. Franke, D. Schwinge, T. Schoknecht, F. Wortmann, et al., Reduced FOXP3(+) regulatory T cells in patients with primary sclerosing cholangitis are associated with IL2RA gene polymorphisms, *J. Hepatol.* 60 (2014) 1010–1016.
- [73] E. Liaskou, L.E. Jeffery, P.J. Trivedi, G.M. Reynolds, S. Suresh, T. Bruns, et al., Loss of CD28 expression by liver-infiltrating T cells contributes to pathogenesis of primary sclerosing cholangitis, *Gastroenterology* 147 (2014), 221–32 e7.
- [74] A. Pawlik, I. Ostanek, I. Brzosko, M. Brzosko, M. Masiuk, B. Machalinski, et al., The expansion of CD4+CD28- T cells in patients with rheumatoid arthritis, *Arthritis Res. Ther.* 5 (2003) R210–R213.
- [75] Y. Aiba, M. Nakamura, S. Joshita, T. Inamine, A. Komori, K. Yoshizawa, et al., Genetic polymorphisms in CTLA4 and SLC4A2 are differentially associated with the pathogenesis of primary biliary cirrhosis in Japanese patients, *J. Gastroenterol.* 46 (2011) 1203–1212.
- [76] Y. Hitomi, M. Kawashima, Y. Aiba, N. Nishida, M. Matsushashi, H. Okazaki, et al., Human primary biliary cirrhosis-susceptible allele of rs4979462 enhances TNFSF15 expression by binding NF-1, *Hum. Genet.* 134 (7) (2015 Jul) 737–747, <http://dx.doi.org/10.1007/s00439-015-1556-3>.
- [77] G.J. Webb, G.M. Hirschfield, Follicles, germinal centers, and immune mechanisms in primary biliary cirrhosis, *Hepatology* 61 (2015) 424–427.
- [78] L. Wang, Y. Sun, Z. Zhang, Y. Jia, Z. Zou, J. Ding, et al., CXCR5+ CD4+ T follicular helper cells participate in the pathogenesis of primary biliary cirrhosis, *Hepatology* 61 (2) (2015 Feb) 627–638, <http://dx.doi.org/10.1002/hep.27306>.
- [79] K. Kikuchi, K. Tsuneyama, H. Yamada, Y. Kajiyama, K. Matsumoto, H. Tsunashima, et al., Splenic lymph follicles generate immunoglobulin M-producing B cells in primary biliary cirrhosis, *Hepatol. Res.* 44 (10) (2014 Oct) E253–E256, <http://dx.doi.org/10.1111/hepr.12231>.
- [80] M. Carbone, G.F. Mellis, G.J. Alexander, R.H. Westbrook, M.A. Heneghan, R.N. Sandford, et al., Calcineurin inhibitors and the IL12A locus influence risk of recurrent primary biliary cirrhosis after liver transplantation, *Am. J. Transplant. Off. J. Am. Soc. Transplant. Am. Soc. Transpl. Surg.* 13 (2013) 1110–1111.
- [81] G. Hirschfield, M. Gershwin, R. Strauss, M. Mayo, C. Levy, B. Zou, et al., P367 phase 2 study evaluating the efficacy and safety of Ustekinumab in patients with primary biliary cirrhosis who had an inadequate response to ursodeoxycholic acid, *J. Hepatol.* 1 (2014) S189–S190.
- [82] R. Tang, H. Chen, Q. Miao, Z. Bian, W. Ma, X. Feng, et al., The Cumulative Effects of Known Susceptibility Variants to Predict Primary Biliary Cirrhosis Risk, 2015, pp. 1–6.
- [83] T.A. Manolio, F.S. Collins, N.J. Cox, D.B. Goldstein, L.A. Hindorf, D.J. Hunter, et al., Finding the missing heritability of complex diseases, *Nature* 461 (2009) 747–753.
- [84] Y. Okada, D. Wu, G. Trynka, T. Raj, C. Terao, K. Ikari, et al., Genetics of rheumatoid arthritis contributes to biology and drug discovery, *Nature* 506 (2014) 376–381.
- [85] T.H. Karlsen, F. Lammert, R.J. Thompson, Genetics of liver disease: from pathophysiology to clinical practice, *J. Hepatol.* 62 (2015) S6–S14.
- [86] C.L. Bowlus, C.S. Li, T.H. Karlsen, B.A. Lie, C. Selmi, Primary sclerosing cholangitis in genetically diverse populations listed for liver transplantation: unique clinical and human leukocyte antigen associations, *Liver Transplant. Off. Publ. Am. Assoc. Study Liver Dis. Int. Liver Transplant. Soc.* 16 (2010) 1324–1330.
- [87] A. Cortes, M.A. Brown, Promise and pitfalls of the immunochip, *Arthritis Res. Ther.* 13 (2011) 101.
- [88] E.K. Karlsson, D.P. Kwiatkowski, P.C. Sabeti, Natural selection and infectious disease in human populations, *Nat. Rev. Genet.* 15 (2014) 379–393.
- [89] L. Jostins, S. Ripke, R.K. Weersma, R.H. Duerr, D.P. McGovern, K.Y. Hui, et al., Host-microbe interactions have shaped the genetic architecture of inflammatory bowel disease, *Nature* 491 (2012) 119–124.
- [90] K.K.-H. Farh, A. Marson, J. Zhu, M. Kleinewietfeld, W.J. Housley, S. Beik, et al., Genetic and epigenetic fine mapping of causal autoimmune disease variants, *Nature* 518 (2015) 337–343.
- [91] P. Invernizzi, M. Miozzo, P.M. Battezzati, I. Bianchi, F.R. Grati, G. Simoni, et al., Frequency of monosomy X in women with primary biliary cirrhosis, *Lancet* 363 (2004) 533–535.
- [92] P. Invernizzi, M. Miozzo, C. Selmi, L. Persani, P.M. Battezzati, M. Zuin, et al., X chromosome monosomy: a common mechanism for autoimmune diseases, *J. Immunol.* 175 (2005) 575–578.
- [93] A. Lleo, J. Liao, P. Invernizzi, M. Zhao, F. Bernuzzi, L. Ma, et al., Immunoglobulin M levels inversely correlate with CD40 ligand promoter methylation in patients with primary biliary cirrhosis, *Hepatology* 55 (2012) 153–160.
- [94] A. Lleo, W. Zhang, M. Zhao, Y. Tan, F. Bernuzzi, B. Zhu, et al., DNA methylation profiling of the X chromosome reveals an aberrant demethylation on CXCR3 promoter in primary biliary cirrhosis, *Clin. Epigenetics* 7 (2015) 61.
- [95] Consortium C, Sparse whole-genome sequencing identifies two loci for major depressive disorder, *Nature* 523 (2015) 588–591.
- [96] J.Z. Liu, M.A. Almarri, D.J. Gaffney, G.F. Mellis, L. Jostins, H.J. Cordell, et al., Dense fine-mapping study identifies new susceptibility loci for primary biliary cirrhosis, *Nat. Publ. Group* 44 (2012) 1137–1141.
- [97] J.Z. Liu, J.R. Hov, T. Folseraas, E. Ellinghaus, S.M. Rushbrook, N.T. Doncheva, et al., Dense genotyping of immune-related disease regions identifies nine new risk loci for primary sclerosing cholangitis, *Nat. Genet.* 45 (2013) 670–675.
- [98] H.J. Cordell, Y. Han, Y. Li, G.F. Mellis, G.M. Hirschfield, International genome-wide meta-analysis in primary biliary cirrhosis discovers six new risk loci and highlights the role of JAK-STAT/IL12/IL27 signaling and cytokine-cytokine pathways in disease etiology, *Nat. Commun.* (2015) 1–13 [in press].
- [99] M.E. Gershwin, Y. Shoenfeld, Abul Abbas: an epitome of scholarship, *J. Autoimmun.* 45 (2013) 1–6.
- [100] L. Steinman, Y. Shoenfeld, From defining antigens to new therapies in multiple sclerosis: honoring the contributions of Ruth Arnon and Michael Sela, *J. Autoimmun.* 54 (2014) 1–7.
- [101] M. Krzywinski, J. Schein, I. Birol, J. Connors, R. Gascoyne, D. Horsman, et al., Circos: an information aesthetic for comparative genomics, *Genome Res.* 19 (2009) 1639–1645.
- [102] T.K. Yu, E.G. Caudell, C. Smid, E.A. Grimm, IL-2 activation of NK cells: involvement of MKK1/2/ERK but not p38 kinase pathway, *J. Immunol.* 164 (2000) 6244–6251.
- [103] M.C. Mingari, F. Gerosa, G. Carra, R.S. Accolla, A. Moretta, R.H. Zubler, et al., Human interleukin-2 promotes proliferation of activated B cells via surface receptors similar to those of activated T cells, *Nature* 312 (1984) 641–643.
- [104] T.R. Malek, The biology of interleukin-2, *Annu. Rev. Immunol.* 26 (2008) 453–479.
- [105] B.H. Nelson, IL-2, regulatory T cells, and tolerance, *J. Immunol.* 172 (2004) 3983–3988.
- [106] G. Trinchieri, Interleukin-12 and the regulation of innate resistance and adaptive immunity, *Nat. Rev. Immunol.* 3 (2003) 133–146.
- [107] A.L. Croxford, P. Kulig, B. Becher, IL-12 and IL-23 in health and disease, *Cytokine Growth Factor Rev.* 25 (2014) 415–421.
- [108] F.E. Watt, O.F.W. James, D.E.J. Jones, Patterns of autoimmunity in primary biliary cirrhosis patients and their families: a population-based cohort study, *QJM* 97 (2004) 397–406.
- [109] L.E. Lamberts, M. Janse, E.B. Haagsma, A.P. van den Berg, R.K. Weersma, Immune-mediated diseases in primary sclerosing cholangitis, *Dig. Liver Dis. Off. J. Italian Soc. Gastroenterol. Italian Assoc. Study Liver* 43 (2011) 802–806.
- [110] M. Werner, H. Prytz, B. Ohlsson, S. Almer, E. Bjornsson, A. Bergquist, et al., Epidemiology and the initial presentation of autoimmune hepatitis in Sweden: a nationwide study, *Scand. J. Gastroenterol.* 43 (2008) 1232–1240.
- [111] G.M. Hirschfield, X. Liu, C. Xu, Y. Lu, G. Xie, Y. Lu, et al., Primary biliary cirrhosis associated with HLA, IL12A, and IL12RB2 variants, *N. Engl. J. Med.* 360 (2009) 2544–2555.
- [112] G.M. Hirschfield, X. Liu, Y. Han, I.P. Gorlov, Y. Lu, C. Xu, et al., Variants at IRF5-TNPO3, 17q12-21 and MMEL1 are associated with primary biliary cirrhosis, *Nat. Publ. Group* 42 (2010) 655–657.
- [113] X. Liu, I. Pietro, Y. Lu, R. Kosoy, Y. Lu, I. Bianchi, et al., Genome-wide meta-analyses identify three loci associated with primary biliary cirrhosis, *Nat. Publ. Group* 42 (2010) 658–660.
- [114] G.F. Mellis, J.A.B. Floyd, K.I. Morley, H.J. Cordell, C.S. Franklin, S.-Y. Shin, et al., Genome-wide association study identifies 12 new susceptibility loci for primary biliary cirrhosis, *Nat. Genet.* 43 (2011) 329–332.
- [115] X. Liu, P. Invernizzi, Y. Lu, R. Kosoy, Y. Lu, I. Bianchi, et al., Genome-wide meta-analyses identify three loci associated with primary biliary cirrhosis, *Nat. Publ. Group* 42 (2010) 658–660.
- [116] T.H. Karlsen, A. Franke, E. Melum, A. Kaser, J.R. Hov, T. Balschun, et al., Genome-wide association analysis in primary sclerosing cholangitis, *Gastroenterology* 138 (2010) 1102–1111.
- [117] E. Melum, A. Franke, C. Schramm, T.J. Weismüller, D.N. Gotthardt, F.A. Offner, et al., Genome-wide association analysis in primary sclerosing cholangitis identifies two non-HLA susceptibility loci, *Nat. Publ. Group* 43 (2010) 17–19.
- [118] T. Folseraas, E. Melum, P. Rausch, B.D. Juran, E. Ellinghaus, A. Shiryayev, et al., Extended analysis of a genome-wide association study in primary sclerosing cholangitis detects multiple novel risk loci, *J. Hepatol.* 57 (2012) 366–375.
- [119] B. Srivastava, G.F. Mellis, H.J. Cordell, A. Muriithi, M. Brown, E. Ellinghaus, et al., Fine mapping and replication of genetic risk loci in primary sclerosing cholangitis, *Scand. J. Gastroenterol.* 47 (2012) 820–826.
- [120] D. Ellinghaus, T. Folseraas, K. Holm, E. Ellinghaus, E. Melum, T. Balschun, et al., Genome-wide association analysis in primary sclerosing cholangitis and ulcerative colitis identifies risk loci at GPR35 and TCF4, *Hepatology* 58 (2013) 1074–1083.
- [121] P.T. Donaldson, J.M. Farrant, M.L. Wilkinson, K. Hayllar, B.C. Portmann, R. Williams, Dual association of HLA DR2 and DR3 with primary sclerosing cholangitis, *Hepatology* 13 (1991) 129–133.
- [122] P.T. Donaldson, Genetics of liver disease: immunogenetics and disease

- pathogenesis, *Gut* 53 (2004) 599–608.
- [123] M.D. Strettell, P.T. Donaldson, L.J. Thomson, P.J. Santrach, S.B. Moore, A.J. Czaja, et al., Allelic basis for HLA-encoded susceptibility to type 1 autoimmune hepatitis, *Gastroenterology* 112 (1997) 2028–2035.
- [124] K. Yoshizawa, M. Ota, Y. Katsuyama, T. Ichijo, A. Matsumoto, E. Tanaka, et al., Genetic analysis of the HLA region of Japanese patients with type 1 autoimmune hepatitis, *J. Hepatol.* 42 (2005) 578–584.
- [125] P.L. Bittencourt, A.C. Goldberg, E.L. Cancado, G. Porta, F.J. Carrilho, A.Q. Farias, et al., Genetic heterogeneity in susceptibility to autoimmune hepatitis types 1 and 2, *Am. J. Gastroenterol.* 94 (1999) 1906–1913.
- [126] D.N. Amarapurkar, N.D. Patel, A.D. Amarapurkar, S.R. Kankonkar, HLA genotyping in type-I autoimmune hepatitis in Western India, *J. Assoc. Phys. India* 51 (2003) 967–969.
- [127] J.M. Lott, T.L. Sumpter, H.R. Turnquist, New dog and new tricks: evolving roles for IL-33 in type 2 immunity, *J. Leukoc. Biol.* 97 (2015) 1037–1048.
- [128] S. Luo, D.C. Rubinsztein, BCL2L1/BIM, *Autophagy* 9 (2012) 104–105.
- [129] C.E. Rudd, A. Taylor, H. Schneider, CD28 and CTLA-4 coreceptor expression and signal transduction, *Immunol. Rev.* 229 (2009) 12–26.
- [130] K. Hieshima, T. Imai, G. Opdenakker, J. Van Damme, J. Kusuda, H. Tei, et al., Molecular cloning of a novel human CC chemokine liver and activation-regulated chemokine (LARC) expressed in liver. Chemotactic activity for lymphocytes and gene localization on chromosome 2, *J. Biol. Chem.* 272 (1997) 5846–5853.
- [131] J.L. Maravillas-Montero, A.M. Burkhardt, P.A. Hevezi, C.D. Carnevale, M.J. Smit, A. Zlotnik, Cutting edge: GPR35/CXCR8 is the receptor of the mucosal chemokine CXCL17, *J. Immunol.* 194 (2015) 29–33.
- [132] H. Song, K.K. Mak, L. Topol, K. Yun, J. Hu, L. Garrett, et al., Mammalian Mst1 and Mst2 kinases play essential roles in organ size control and tumor suppression, *Proc. Natl. Acad. Sci. U. S. A.* 107 (2010) 1431–1436.
- [133] M.S. Hayden, S. Ghosh, NF-kappaB, the first quarter-century: remarkable progress and outstanding questions, *Genes Dev.* 26 (2012) 203–234.
- [134] J. Strebovsky, P. Walker, R. Lang, A.H. Dalpke, Suppressor of cytokine signaling 1 (SOCS1) limits NFkappaB signaling by decreasing p65 stability within the cell nucleus, *FASEB J. Off. Publ. Fed. Am. Soc. Exp. Biol.* 25 (2011) 863–874.
- [135] A. Muto, S. Tashiro, O. Nakajima, H. Hoshino, S. Takahashi, E. Sakoda, et al., The transcriptional programme of antibody class switching involves the repressor Bach2, *Nature* 429 (2004) 566–571.
- [136] R. Roychoudhuri, K. Hirahara, K. Mousavi, D. Clever, C.A. Klebanoff, M. Bonelli, et al., BACH2 represses effector programs to stabilize T(reg)-mediated immune homeostasis, *Nature* 498 (2013) 506–510.
- [137] L. Vereecke, R. Beyaert, G. van Loo, The ubiquitin-editing enzyme A20 (TNFAIP3) is a central regulator of immunopathology, *Trends Immunol.* 30 (2009) 383–391.
- [138] D.B. Schubart, A. Rolink, M.H. Kosco-Vilbois, F. Botteri, P. Matthias, B-cell-specific coactivator OBF-1/OCA-B/Bob1 required for immune response and germinal centre formation, *Nature* 383 (1996) 538–542.
- [139] M. Sanosaka, M. Fujimoto, T. Ohkawara, T. Nagatake, Y. Itoh, M. Kagawa, et al., Salt-inducible kinase 3 deficiency exacerbates lipopolysaccharide-induced endotoxin shock accompanied by increased levels of pro-inflammatory molecules in mice, *Immunology* 145 (2015) 268–278.
- [140] T. Kumpfel, R. Hohlfeld, Multiple sclerosis. TNFRSF1A, TRAPS and multiple sclerosis, *Nat. Rev. Neurol.* 5 (2009) 528–529.
- [141] B. Barneda-Zahonero, L. Roman-Gonzalez, O. Collazo, H. Rafati, A.B. Islam, L.H. Bussmann, et al., HDAC7 is a repressor of myeloid genes whose down-regulation is required for transdifferentiation of pre-B cells into macrophages, *PLoS Genet.* 9 (2013) e1003503.
- [142] J. Devalliere, B. Charreau, The adaptor Lnk (SH2B3): an emerging regulator in vascular cells and a link between immune and inflammatory signaling, *Biochem. Pharmacol.* 82 (2011) 1391–1402.
- [143] E.A. Green, R.A. Flavell, TRANCE-RANK, a new signal pathway involved in lymphocyte development and T cell activation, *J. Exp. Med.* 189 (1999) 1017–1020.
- [144] J. Sun, G. Matthias, M.J. Mihatsch, K. Georgopoulos, P. Matthias, Lack of the transcriptional coactivator OBF-1 prevents the development of systemic lupus erythematosus-like phenotypes in aiolos mutant mice, *J. Immunol.* 170 (2003) 1699–1706.
- [145] M.P. Forrest, M.J. Hill, A.J. Quantock, E. Martin-Rendon, D.J. Blake, The emerging roles of TCF4 in disease and development, *Trends Mol. Med.* 20 (2014) 322–331.
- [146] L. Martinet, M.J. Smyth, Balancing natural killer cell activation through paired receptors, *Nat. Rev. Immunol.* 15 (2015) 243–254.
- [147] M. Ishizaki, T. Akimoto, R. Muromoto, M. Yokoyama, Y. Ohshiro, Y. Sekine, et al., Involvement of tyrosine kinase-2 in both the IL-12/Th1 and IL-23/Th17 axes in vivo, *J. Immunol.* 187 (2011) 181–189.
- [148] K. Sasaki, J. Hamazaki, M. Koike, Y. Hirano, M. Komatsu, Y. Uchiyama, et al., PAC1 gene knockout reveals an essential role of chaperone-mediated 20S proteasome biogenesis and latent 20S proteasomes in cellular homeostasis, *Mol. Cell. Biol.* 30 (2010) 3864–3874.