

# Emerging halogenated flame retardants and hexabromocyclododecanes in food samples from an e-waste processing area in Vietnam

Tao, Fang; Matsukami, Hidenori; Suzuki, Go; Tue, Nguyen Minh; Viet, Pham Hung; Takigami, Hidetaka; Harrad, Stuart

DOI:

[10.1039/c5em00593k](https://doi.org/10.1039/c5em00593k)

License:

None: All rights reserved

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Tao, F, Matsukami, H, Suzuki, G, Tue, NM, Viet, PH, Takigami, H & Harrad, S 2016, 'Emerging halogenated flame retardants and hexabromocyclododecanes in food samples from an e-waste processing area in Vietnam', *Environmental Science Processes and Impacts*. <https://doi.org/10.1039/c5em00593k>

[Link to publication on Research at Birmingham portal](#)

**Publisher Rights Statement:**

Final version of record published as above.

Checked Feb 2016

**General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

**Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.



**EMERGING HALOGENATED FLAME RETARDANTS AND  
HEXABROMOCYCLODODECANES IN FOOD SAMPLES FROM  
AN E-WASTE PROCESSING AREA IN VIETNAM**

Journal:	<i>Environmental Science: Processes &amp; Impacts</i>
Manuscript ID	EM-ART-11-2015-000593.R1
Article Type:	Paper
Date Submitted by the Author:	n/a
Complete List of Authors:	Tao, Fang; University of Birmingham, Matsukami, Hidenori ; National Institute for Environmental Studies Suzuki, Go; NIES, Tue, Nguyen; Ehime University, Center for Marine Environmental Studies Viet, Pham Hung; Research Centre for Environmental Technology and Sustainable Development, Hanoi University of Science, Takigami, Hidetaka; National Institute for Environmental Studies, Res. Centre for Material Recycle + Waste Management Harrad, S; School of Chemistry, Institute of Environmental Health

1  
2  
3  
4 1 **EMERGING HALOGENATED FLAME RETARDANTS AND**  
5  
6 2 **HEXABROMOCYCLODODECANES IN FOOD SAMPLES FROM**  
7  
8  
9 3 **AN E-WASTE PROCESSING AREA IN VIETNAM**  
10  
11  
12 4

13  
14 5 Fang Tao<sup>a\*</sup>, Hidenori Matsukami<sup>b</sup>, Go Suzuki<sup>b</sup>, Nguyen Minh Tue<sup>c,d</sup>, Pham Hung  
15  
16 6 Viet<sup>c</sup>, Hidetaka Takigami<sup>b</sup>, Stuart Harrad<sup>a</sup>  
17  
18 7

19  
20  
21 8 <sup>a</sup> School of Geography, Earth and Environmental Sciences,  
22

23  
24 9 University of Birmingham,  
25

26 10 Birmingham  
27

28  
29 11 B15 2TT,  
30

31 12 UK  
32  
33

34 13  
35

36 14 <sup>b</sup> National Institute for Environmental Studies,  
37

38  
39 15 Ibaraki 305-8506,  
40

41 16 Tsukuba,  
42

43  
44 17 Japan  
45

46 18  
47

48  
49 19 <sup>c</sup> Centre for Environmental Technology and Sustainable Development  
50

51 20 Hanoi University of Science  
52

53  
54 21 334 Nguyen Trai  
55

56 22 Hanoi  
57  
58  
59  
60

1  
2  
3  
4 1 Vietnam

5  
6 2

7  
8  
9 3 <sup>d</sup>Center for Marine Environmental Studies

10  
11 4 Ehime University, 2-5 Bunkyo-cho

12  
13 5 Matsuyama

14  
15 6 Japan

16  
17 7 \* Corresponding Author

18  
19 8 E-mail address: [fangtao\\_bham@sina.com](mailto:fangtao_bham@sina.com)

20  
21 9

22  
23 10

24  
25 11

26  
27 12

28  
29 13

30  
31 14

32  
33 15

34  
35 16

36  
37 17

38  
39 18

40  
41 19

42  
43 20

44  
45 21

46  
47 22

1  
2  
3  
4 **Abstract**

5  
6 This study reports concentrations of selected emerging halogenated flame retardants  
7  
8 (HFRs) and hexabromocyclododecanes (HBCDs) in foodstuffs sourced from an e-  
9  
10 waste processing area in Vietnam and two reference sites in Vietnam and Japan.  
11  
12 Concentrations of all target HFRs in e-waste-impacted samples in this study exceed  
13  
14 significantly ( $p < 0.05$ ) those detected in the controls, suggesting e-waste processing  
15  
16 activities exert a substantial impact on local environmental contamination and human  
17  
18 dietary exposure. Significant linear positive correlations in concentrations of *syn*-  
19  
20 Dechlorane Plus (DP) and *anti*-DP were found between soils and those in co-located  
21  
22 chicken samples ( $p < 0.05$ ). This implies soil is an important exposure source of DPs in  
23  
24 chickens at our sampling sites. The estimated dietary intakes of emerging HFRs in  
25  
26 this study were 170 and 420 ng/kg bw/day, for adults and children respectively; while  
27  
28 daily ingestions of HBCDs were an estimated 480, 1500 ng/kg bw/day for adults and  
29  
30 children, respectively. Exposure at the site monitored in this study exceeds  
31  
32 substantially estimates of dietary exposure to HBCDs in e-waste processing sites and  
33  
34 non e-waste processing areas elsewhere.  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45

46 **Keywords:** Emerging HFRs; HBCDs; Food; Soil; E-Waste; Dietary exposure  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 14  
5 **1. Introduction**  
6

7 HFRs have been used widely to ensure manufactured goods such as plastics, textiles,  
8 building materials, vehicle components and electronic equipment meet fire safety  
9 regulations.<sup>1,2</sup> Some HFRs like polybrominated diphenyl ethers (PBDEs) are subject  
10 to bans and restrictions within the EU as well as elsewhere.<sup>3</sup> As a result, there is likely  
11 increasing demand for emerging HFRs, such as DP, 1,2-bis(2,4,6-  
12 tribromophenoxy)ethane (BTBPE), 2-ethylhexyl-2,3,4,5-tetrabromobenzoate (EH-  
13 TBB), bis(2-ethylhexyl)-3,4,5,6-tetrabromophthalate (BEH-TEBP),  
14 pentabromobenzene (PBBz), hexabromobenzene (HBB), and decabromodiphenyl  
15 ethane (DBDPE). Recently, such emerging HFRs have been detected in various  
16 environmental matrices, including indoor dust, air, sediment, soil, and sewage  
17 sludge.<sup>4-10</sup> Studies of the environmental behaviour of emerging HFRs are scarce, with  
18 the limited information available suggesting that, similar to regulated HFRs like  
19 PBDEs, they may be toxic, persistent, bioaccumulative, and subject to long-range  
20 transport. This information includes: the detection of DP, EH-TBB, BEH-TEBP, HBB  
21 and BTBPE in the Arctic,<sup>11,12</sup> the long half-life and biomagnification potential of  
22 BTBPE in juvenile rainbow trout,<sup>13</sup> a recent report that oral exposure to DP can  
23 induce hepatic oxidative damage and perturbations of metabolism and signal  
24 transduction in male mice,<sup>14</sup> and studies showing accumulation of some emerging  
25 HFRs in fish, birds and dolphins.<sup>15-17</sup>

26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 1 Moreover, increasing attention is being paid to environmental contamination with  
5  
6 2 HFRs arising from emissions occurring during end-of-life treatment of treated goods.  
7  
8  
9 3 Particular concern exists about situations where electrical and electronic waste (e-  
10  
11 4 waste) is dismantled under rudimentary conditions. Numerous studies have shown  
12  
13 5 elevated contamination of air, dust, soil, and sediments with PBDEs in such locations,  
14  
15 6 <sup>18–20</sup> with a smaller number of recent studies reporting elevated human dietary  
16  
17 7 exposure to local populations in such areas. In contrast, relatively few data exist about  
18  
19 8 environmental contamination with emerging HFRs in areas where such informal e-  
20  
21 9 waste processing is conducted, although a recent study reported concentrations of EH-  
22  
23 10 TBB and BEH-TEBP in foodstuffs from a Chinese e-waste impacted location to  
24  
25 11 exceed those from control locations.<sup>21</sup> In addition to PBDEs,  
26  
27 12 hexabromocyclododecanes (HBCDs) have also been frequently detected in the  
28  
29 13 environment and human milk sampled in the vicinity of rudimentary e-waste  
30  
31 14 processing sites.<sup>22–24</sup> However, studies of human exposure via consumption of food  
32  
33 15 sourced from locations where rudimentary processing of e-waste is conducted are  
34  
35 16 scarce. Zheng et al.<sup>25</sup> reported the estimated human exposure to HBCDs via ingestion  
36  
37 17 of eggs sourced from an e-waste recycling site in South China, while Labunska et al.<sup>21</sup>  
38  
39 18 reported human dietary exposure to HBCDs via consumption of a variety of  
40  
41 19 foodstuffs including fish and pork as well as chicken eggs, liver, and muscle from e-  
42  
43 20 waste recycling sites in Taizhou, China.  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 1 The available data for PBDEs and HBCDs suggest that fish, pigs, and free-range  
5  
6 2 chickens reared in e-waste impacted locations accumulate these compounds – likely  
7  
8  
9 3 through uptake from their environment (e.g. soil and sediment) and food.  
10  
11 4 Consequently, elevated dietary exposure to such HFRs is likely for people consuming  
12  
13 5 food derived from such animals reared in the vicinity of e-waste processing areas. To  
14  
15  
16 6 date however, evidence of similar exposure to emerging HFRs via consumption of  
17  
18  
19 7 fish, pork, and chicken liver, eggs, skin and muscle derived from e-waste impacted  
20  
21 8 areas is very limited. Bui Dau is a small village where e-waste processing started at  
22  
23 9 the beginning of the 21<sup>st</sup> century in Vietnam. Concentrations of polychlorinated  
24  
25  
26 10 biphenyls, and organophosphorus flame retardants, tetrabromobisphenol A, emerging  
27  
28  
29 11 HFRs, PBDEs and HBCDs in human and environment samples have already been  
30  
31 12 reported in this area.<sup>23,24,26</sup> Against this backdrop, and also as part of the study above  
32  
33  
34 13 to investigate site-specific contamination levels in Bui Dau, this study's objective is to  
35  
36  
37 14 determine concentrations of emerging HFRs and HBCDs in such foodstuffs collected  
38  
39  
40 15 from a location impacted by rudimentary e-waste processing in Bui Dau, Vietnam.  
41  
42 16 These data are compared with concentrations detected in samples of the same  
43  
44 17 matrices sourced from non-e-waste impacted control locations in Vietnam and Japan,  
45  
46  
47 18 and are combined with information on their consumption by local inhabitants to  
48  
49 19 estimate human exposure to these contaminants. Concentrations of emerging HFRs  
50  
51 20 and HBCDs are also measured in soil samples from the same e-waste impacted  
52  
53  
54 21 locations, to investigate the impact of soil contamination on chicken tissues and eggs.  
55  
56  
57 22



## 2. Methods

### 2.1 Sampling and pretreatment

Samples were collected in January 2014 from an e-waste processing area in Bui Dau (Cam Xa, Hung Yen province) in Vietnam. The locations of the sampling sites are presented in Fig. 1. The sampling area is a rural village of approximately 200 households. The main supplies of livestock products and fish for the local people in Bui Dau are from neighbouring communities,<sup>24</sup> and the livestock and fish raised in farm yards in Bui Dau are intended mainly for consumption by the families themselves with any small surpluses sold commercially. E-waste processing activities such as dismantling of electrical wires and metals, shredding of plastics into pellets, manual recycling of TVs, printers, printed circuit boards and other computer components are mainly family-based, with e-waste handled in the backyards of domestic buildings, in which livestock are raised. Fresh chicken eggs (n=3 from each site) were collected from chicken farm owners at six sampling sites (site 1-site 6) shown in Fig. 1. In addition, five chickens (one from each site) were also purchased from chicken owners at five locations (site 1-site 5), from which samples of chicken muscle (n=5), chicken liver (n=5), and chicken skin (n=5) were derived. Soil samples (one from each site) were collected at the same time from the backyards where the chickens were raised. Each soil sample analysed consisted of five sub-samples from the same backyard homogenised before analysis.

1  
2  
3  
4 1 In addition to chicken, egg, and soil samples; five tilapia (*Oreochromis mossambicus*)  
5  
6 2 samples were collected from the river (site 7), with a further five fish samples (3 rohu  
7  
8 3 (*Labeo rohita*) and 2 tilapia) collected from the fish pond located in Bui Dau.  
9  
10 4 Moreover, two samples of pork muscle were purchased from a market stall in Bui Dau  
11  
12 5 village.  
13  
14  
15  
16  
17  
18

19 7 *Control samples* One control chicken egg sample and one control chicken  
20  
21 8 muscle sample were purchased from Thanh Hoa province, situated approximately 175  
22  
23 9 km distant from the e-waste processing location. Additional control samples of  
24  
25 10 chicken egg, muscle, liver, and skin were purchased in Tsukuba, Japan, with control  
26  
27 11 samples (n=1 in each case except for chicken egg (n=3)) of pork and fish purchased in  
28  
29 12 Hanoi City.  
30  
31  
32  
33  
34  
35

36 14 *Sample treatment prior to analysis* Following collection, egg samples were boiled on  
37  
38 15 location in a large pan for 8 minutes at 100 °C and then cooled to room temperature.  
39  
40 16 Egg yolks were then separated from the egg white and wrapped in pre-cleaned  
41  
42 17 aluminum foil, prior to storage at -20 °C until shipping to the laboratory with ice  
43  
44 18 bags.  
45  
46  
47  
48  
49  
50

## 51 2.2 Sample Preparation for GC/MS Analysis

52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 1 Seven emerging HFRs (PBBz, HBB, BEH-TEBP, BTBPE, DBDPE, *syn*- and *anti*-  
5  
6 2 DP) as well as  $\alpha$ -,  $\beta$ - and  $\gamma$ -HBCDs were measured in all samples (CAS numbers and  
7  
8  
9 3 full names of emerging HFRs are provided as supporting material).

10  
11 4  
12  
13  
14 5 Following homogenization and freeze-drying, approximately 5 g of each sample was  
15  
16 6 extracted firstly using a rapid solvent extractor (SE-100; Mitsubishi Chemical  
17  
18 7 Analytec Co., Ltd.) at 35 °C for 40 min with 50% acetone in n-hexane at a flow rate  
19  
20  
21 8 of 2 mL/min, followed by secondary extraction at 80 °C for 40 min with toluene at a  
22  
23  
24 9 flow rate of 2 mL/min. Resultant extracts were evaporated and solvent exchanged to  
25  
26 10 10 mL toluene. An aliquot (1 mL) of the extract was used to determine lipids by  
27  
28  
29 11 evaporation to dryness and gravimetry. A further 2 mL aliquot of the crude solvent  
30  
31 12 extract was taken before being spiked with a known amount of  $^{13}\text{C}_{12}$ -labelled internal  
32  
33  
34 13 (or surrogate) standards ( $^{13}\text{C}_{12}$ -HBB,  $^{13}\text{C}_{12}$ -BTBPE,  $^{13}\text{C}_{12}$ -BDE209,  $^{13}\text{C}_{12}$ - $\alpha$ -,  $\beta$ -,  $\gamma$ -  
35  
36 14 HBCDs). These 2 mL aliquots were eluted through a gel permeation chromatography  
37  
38  
39 15 column packed with SX-3 Bio-Beads (Bio-Rad Laboratories, Inc.); the eluate from  
40  
41 16 which was separated into two further aliquots. The first of these aliquots was  
42  
43  
44 17 subjected to further purification by passing through a 2 g deactivated silica gel column  
45  
46 18 (eluted with 10 mL 50% diethyl ether in hexane) followed by a 8 g florisil column  
47  
48  
49 19 (eluted with 50 mL hexane and 50 mL dichloromethane). The final eluate was  
50  
51 20 concentrated and solvent exchanged to 50  $\mu\text{L}$  nonane containing  $^{13}\text{C}_{12}$ -BDE 138 as a  
52  
53  
54 21 recovery determination standard. The second aliquot was evaporated to nearly dryness  
55  
56 22 and transferred to 2 mL hexane, which was then partitioned against 3-4 mL  
57  
58  
59  
60

1  
2  
3  
4 1 concentrated sulfuric acid, prior to evaporation to incipient dryness and  
5  
6 2 resolubilisation in 100  $\mu$ L methanol. Instrumental analysis of target emerging HFRs  
7  
8  
9 3 was performed using an Agilent 6890 GC coupled to an Agilent 5973 MS operated in  
10  
11 4 electron impact (EI) mode. 2  $\mu$ L of cleaned extract was injected on a DB-1HT  
12  
13 5 capillary column (15 m  $\times$  0.25 mm i.d., 0.1  $\mu$ m film thickness; Agilent Technologies,  
14  
15 6 Inc.) via cool on-column injection. The injection temperature was set at 100  $^{\circ}$ C, hold  
16  
17 7 0.1 min, ramp 100  $^{\circ}$ C/min to 300  $^{\circ}$ C, hold 15 min. The GC temperature program was  
18  
19 8 set at 100  $^{\circ}$ C, hold 1 min, ramp 8  $^{\circ}$ C/min to 310  $^{\circ}$ C, hold 10 min.  
20  
21  
22

23  
24 9 Determination of  $\alpha$ -,  $\beta$ - and  $\gamma$ -HBCDs was achieved using an LC-MS/MS system  
25  
26 10 composed of a dual pump Shimadzu LC-20AB Prominence liquid chromatograph  
27  
28 11 equipped with a SIL-20A autosampler and a DGU-20A3 vacuum degasser, coupled to  
29  
30 12 a Sciex API 2000 triple quadrupole mass spectrometer. (Applied Biosystems, Foster  
31  
32 13 City, CA, USA) operated in electrospray negative ionization (ESI) mode equipped  
33  
34 14 with a Varian Pursuit XRS3 C18 analytical column (150 mm  $\times$  2 mm I.D., 3  $\mu$ m  
35  
36 15 particle size). Details of this instrumental method are provided elsewhere.<sup>21,27</sup>  
37  
38  
39  
40  
41  
42  
43

### 44 17 **2.3 Quality assurance/quality control**

45  
46 18 Quality assurance (QA) and control (QC) was achieved via regular analysis of  
47  
48 19 procedural blanks and matrix spike samples – the latter comprising food samples  
49  
50 20 purchased in Japan and spiked with known amounts of the target emerging HFRs and  
51  
52 21 HBCDs. A 9-point calibration curve spanning the concentration range 1.25–1,000  
53  
54 22 pg/ $\mu$ L including the  $^{13}\text{C}_{12}$ -labelled HFRs was employed.  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 1  
5  
6 2 Initial evaluation of analyte recoveries was assessed via the following matrix spiking  
7  
8  
9 3 experiment. Samples of chicken liver, egg, skin, muscle, and fish were purchased in  
10  
11 4 Japan as low-contamination controls. Aliquots (5 g) of these were spiked with known  
12  
13 5 amounts of both native and isotopically-labelled analytes included in the present study,  
14  
15  
16 6 and the samples were extracted and cleaned-up using the standard procedure. After  
17  
18  
19 7 deducting levels in the unspiked control samples, the recoveries of native compounds  
20  
21 8 were from 56-98% and the recoveries of internal standards ranged from 90-117%.  
22  
23  
24 9

#### 10 **2.4 Statistical Analysis and daily intake calculations**

11 Pearson's correlation analysis was conducted to investigate relationships in  
12  
13 concentrations of BTBPE, *syn*-DP, *anti*-DP, and HBCDs between chicken and soil  
14  
15 samples. Statistical analysis was performed using IBM SPSS Statistics 21.0 and a p  
16  
17 value <0.05 was regarded as indicating statistical significance.  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

16 Dietary intakes (DI) (ng/kg bw/day) were calculated thus:

$$DI = \sum_{i=1}^n \frac{C_i \times CR_i}{BW}$$

18 Where  $C_i$  is the concentration (ng/g ww) of HFR in a food item  $i$  and  $CR_i$  is the daily  
19  
20 consumption rate of the foodstuff  $i$  (g/day; values given in supporting material). Body  
21  
22 weight (BW; kg) values employed in this study were assumed to be 51 kg for adults  
23  
24 according to our sampling questionnaire and 14.65 kg for children based on WHO-

1  
2  
3  
4 1 growth chart data for Vietnamese children under 3 years old. No previous information  
5  
6 2 exists about consumption rates of the foodstuffs (fish, chicken liver, poultry, and meat)  
7  
8  
9 3 in Vietnam, so we used information provided in a previous study of an e-waste  
10  
11 4 processing area in Eastern China.<sup>21</sup>  
12  
13  
14  
15  
16

17 6 As we measured HFRs concentrations in egg yolk only, concentrations on a whole  
18  
19 7 egg basis were calculated and the weight of eggs was corrected for the eggshell  
20  
21 8 weight for calculating daily intakes of HFRs via chicken egg consumption according  
22  
23  
24 9 to a previous study.<sup>19</sup>  
25  
26  
27  
28  
29

### 30 11 **3. Results and discussion**

#### 31 32 12 **3.1 Detection of HFRs in food samples**

33  
34  
35 13 Table 1 shows concentrations of HFRs in food, co-located soil and co-located  
36  
37 14 sediment samples.<sup>28</sup> Of our target emerging HFRs: PBBz, HBB, BEH-TEBP, BTBPE,  
38  
39 15 DBDPE, *syn*-DP, *anti*-DP were detectable in chicken samples, with the same HFRs  
40  
41  
42 16 (except PBBz) found in soil samples. In contrast, only BTBPE was detected in river  
43  
44  
45 17 fish, only DBDPE was detectable in pork and no emerging HFRs were detected in  
46  
47 18 pond fish.  
48  
49  
50  
51

52  
53 20 Most strikingly, concentrations of all target emerging HFRs in e-waste-impacted  
54  
55 21 samples in this study exceed substantially those detected in the corresponding  
56  
57 22 controls. This demonstrates substantial impact of the e-waste processing activities on  
58  
59  
60

1  
2  
3  
4 1 the environment in Bui Dau. These findings are consistent with previous studies of  
5  
6 2 PBDEs, PBEB, HBB, EH-TBB, BEH-TEBP, BTBPE, and DBDPE in Taizhou,  
7  
8 3 eastern China.<sup>21,29</sup>  
9  
10 4  
11  
12 5 DPs were the most frequently detected emerging HFR (100% detection) in chicken  
13  
14 6 samples (muscle, liver, egg, skin), followed by BTBPE and HBB with a detection  
15  
16 7 frequency of 70% and 50%, respectively. PBBz was less frequently detected in  
17  
18 8 chicken samples while BEH-TEBP was only detected in chicken egg samples. In  
19  
20 9 contrast, the most frequently detected compounds in foods from e-waste processing  
21  
22 10 sites in Taizhou, eastern China were EH-TBB and BEH-TEBP.<sup>21</sup> This likely reflects  
23  
24 11 different waste compositions at the two locations. Interestingly, in this study, despite  
25  
26 12 the theoretical predictions that organic contaminants for which  $\text{Log } K_{\text{OW}} > 7$  become  
27  
28 13 increasingly less prone to uptake by biota,<sup>30</sup> we detected DBDPE (for which  $\text{Log } K_{\text{OW}}$   
29  
30 14 is reported to be 11.1<sup>3</sup> in chicken muscle, egg and pork samples in this study. The  
31  
32 15 most abundant emerging HFR detected in chicken was DP, with average  
33  
34 16 concentrations of  $\Sigma\text{DP}$  in chicken ranged between 520 and 5,100 ng/g lipid weight  
35  
36 17 (lw), comprising around 90% of the sum of emerging HFRs measured in this study.  
37  
38 18 This is consistent with previous findings for chicken egg samples from e-waste  
39  
40 19 recycling sites in South China.<sup>25</sup> DP was not measured in samples taken from Taizhou  
41  
42 20 in the study of Labunska et al.,<sup>21</sup> so comparison is not possible in this instance. The  
43  
44 21 predominance of DP is likely attributable to its classification as a high production  
45  
46 22 volume (HPV) compound<sup>31</sup>, a term used to describe a chemical produced in the  
47  
48 23 United States in quantities exceeding 450 t per year, and the significant  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 biomagnification potentials of both *syn*- and *anti*-DP in food samples.<sup>32,33</sup>  
2 Furthermore, DP is used in electrical cable coatings<sup>34</sup> and dismantling of electrical  
3 wires was one of the main e-waste processing activities in Bui Dau. Consequently, DP  
4 may be discharged to the environment during the handling of electrical cables. With  
5 respect to our other targeted emerging HFRs, average concentrations in chicken  
6 samples of PBBz, HBB, BEH-TEBP, BTBPE, and DBDPE were <1.5-2.0, 1.5-6.8,  
7 <1.3-2.0, 54-70, and <2.8-280 ng/g lipid weight (lw) respectively. The highest  
8 concentration of  $\Sigma$ DP (25,000 ng/g lw) was found in chicken liver, at the high end of  
9 values (<9630 ng/g lw) previously reported for  $\Sigma$ DP in food samples.<sup>25,33,35,36</sup> Only  
10 limited studies of the presence of emerging HFRs in food samples from e-waste  
11 processing sites, especially in different chicken tissues, are available. One previous  
12 study reported concentrations of  $\Sigma$ DP, DBDPE, BTBPE and HBB in chicken eggs  
13 from e-waste recycling sites in South China, to fall within the ranges 665-3,290, 5.97-  
14 37.9, 37.2-264 and 7.32-25.7 ng/g lw, respectively.<sup>25</sup> These concentrations exceed  
15 those found for  $\Sigma$ DP (4.0-2,300 ng/g lw) and HBB (<1.3-2.0 ng/g lw) in this study,  
16 but are similar to those we report here for DBDPE (<2.8-620 ng/g lw) and BTBPE  
17 (<2.8-160 ng/g lw) in eggs. Average concentrations of HBB and BTBPE in chicken  
18 livers (<0.15 and 15.0 ng/g lw, respectively), chicken muscle samples (0.41 and 1.46  
19 ng/g lw) and chicken eggs (<0.15 and 2.93 ng/g lw) from e-waste processing areas in  
20 Taizhou,<sup>21</sup> are much lower than those in this study (results shown in Table 1).  
21 Moreover, the concentrations of  $\Sigma$ DP in chicken samples in this study were  
22 comparable to those in chicken liver (4.4 ng/g ww) and muscle samples (0.92 ng/g



1  
2  
3  
4 1 ww) from e-waste processing sites in China.<sup>37</sup> Our analyses of DP concentrations in  
5  
6 2 liver, muscle and skin tissues taken from individual chickens, revealed DP  
7  
8  
9 3 concentrations were highest in chicken liver, followed by chicken muscle, similar to  
10  
11 4 data reported recently for chicken in China.<sup>37</sup> Additionally, concentrations of BTBPE  
12  
13 5 in avian (watercock) and fish samples taken from an e-waste processing area in  
14  
15 6 southern China ranged between 0.07-2.41 and <0.012-0.15 ng/g lw, respectively.<sup>16</sup>  
16  
17 7 These values are exceeded substantially in our chicken and fish samples. In contrast,  
18  
19 8 concentrations of DBDPE in avian muscle and fish samples in our study are similar to  
20  
21 9 those reported previously.<sup>16</sup>  
22  
23  
24  
25  
26  
27  
28

29 11 In samples originating from areas not impacted by e-waste processing activities,  
30  
31 12 concentrations of HBB, BTBPE, and DBDPE were determined in a selection of UK  
32  
33 13 and Irish food samples.<sup>38</sup> Only BTBPE was detectable in this earlier study, at  
34  
35 14 concentrations of 0.96, 0.75, 0.29, and 0.55 ng/g lw in fish, chicken liver, and chicken  
36  
37 15 eggs, respectively,<sup>38</sup> substantially lower than those detected in our study. Compared to  
38  
39 16 the presence of BTBPE in UK food samples, BTBPE was not detected in control  
40  
41 17 samples collected in Vietnam and Japan.  
42  
43  
44  
45  
46  
47  
48

49 19 *Concentrations in soil* In co-located soil samples, concentrations of HBB, BEH-  
50  
51 20 TEBP, BTBPE, DBDPE, *syn*-DP, and *anti*-DP were in the range 0.15-21, <0.050-0.4,  
52  
53 21 0.19-34, 0.42-64, 0.20-13, and 0.83-31 ng/g dry wt (dw) respectively. The detection  
54  
55 22 frequency was 100% except for BEH-TEBP which was detected in only 20% of  
56  
57  
58  
59  
60

1 samples. PBBz was not detected in any soil samples in this study. DBDPE was the  
2 most dominant compound in soil, accounting for around 50% of the total emerging  
3 HFRs in our study.

4  
5 Concentrations of emerging HFRs in soil in our study exceed (DBDPE and BTBPE)  
6 or are comparable with ( $\Sigma$ DP) those detected in soils taken from locations in China  
7 surrounding but not directly impacted (e.g. close to workshops) by e-waste  
8 processing.<sup>16,39</sup> Moreover, concentrations of  $\Sigma$ DP in soil in this study exceeded by 3  
9 orders of magnitude those found in soils from residential, business and industrial areas  
10 in northern China.<sup>40</sup> In contrast, the concentration of  $\Sigma$ DP in a single surface soil  
11 directly impacted (i.e. close to a workshop) by e-waste recycling in South China was  
12 3,327 ng/g dw<sup>39</sup> - 2 orders of magnitude higher than the average concentration  
13 detected in our study. Similarly elevated concentrations of DP (5,900-10,000 ng/g dw)  
14 have also been reported in soil samples directly impacted by e-waste activities in  
15 Qingyuan county.<sup>37</sup>

### 16 17 **3.2 Concentrations and diastereomer patterns of HBCDs in food samples and** 18 **co-located soil samples**

19 Concentrations of HBCDs in food samples in this study are shown in Table 1. HBCDs  
20 were detected in all chicken tissues, river fish, pork, and soil samples. In chicken and  
21 fish samples, the levels of HBCDs tend to be higher than those of emerging HFRs  
22 except DP while in pork samples the levels of HBCDs were comparable to those of

1  
2  
3  
4 1 emerging HFRs in line with reports by Labunska et al.<sup>21</sup> and Zheng et al..<sup>25</sup> While  
5  
6 2 HBCDs are known to be mainly used in polystyrene foam and fabrics, they were used  
7  
8  
9 3 to a minor extent in electrical equipment housing.<sup>41</sup> This latter minor application,  
10  
11 4 could explain the elevated levels of HBCDs in this study, as TVs, DVDs, computers  
12  
13  
14 5 and printer housings were processed on a large scale in Bui Dau. Coupled with their  
15  
16 6 environmental stability, persistence and past high production volume<sup>42</sup>, these factors  
17  
18  
19 7 may account for the levels of HBCDs in this study exceeding those for most of the  
20  
21 8 emerging HFRs monitored. In contrast, HBCDs were not detected in pond fish. This  
22  
23  
24 9 study's finding that HBCD concentrations in e-waste-impacted samples exceed those  
25  
26 10 in corresponding controls differs to that of Labunska et al.,<sup>21</sup> who reported that HBCD  
27  
28  
29 11 concentrations in some control samples exceeded those in samples derived from e-  
30  
31 12 waste-impacted locations. To our knowledge, very few data exist about concentrations  
32  
33  
34 13 of HBCDs in foods reared in locations where unregulated e-waste processing is  
35  
36 14 conducted. Average concentrations of  $\Sigma$ HBCDs in chicken liver (3,200 ng/g lw) and  
37  
38  
39 15 egg (3,600 ng/g lw) samples in our study exceeded substantially those found in  
40  
41 16 chicken liver (42.5 ng/g lw) and eggs (42.6 ng/g lw) from an e-waste processing area  
42  
43  
44 17 in Taizhou City<sup>21</sup> as well as in chicken eggs (44.2-350 ng/g lw) from another e-waste  
45  
46 18 processing site in South China (Qingyuan City).<sup>25</sup> Moreover, HBCDs concentrations  
47  
48  
49 19 (0.59-670 ng/g wet weight (ww)) in food samples in this study exceed markedly those  
50  
51 20 detected in similar foodstuffs around the world including China (<LOD to 9.2 ng/g  
52  
53  
54 21 lw),<sup>43</sup> the USA (12-616 ng/g lw),<sup>44</sup> Romania (0.04-0.25 ng/g ww),<sup>45</sup> Sweden (0.005-  
55  
56 22 0.63 ng/g ww),<sup>46</sup> Belgium (<0.01-0.35 ng/g ww),<sup>47</sup> and the UK (0.02-0.30 ng/g ww).<sup>48</sup>  
57  
58  
59  
60

1  
2  
3  
4 1 As shown in Table 1, HBCD concentrations in animal-related food sampled from the  
5  
6 2 e-waste processing site in Vietnam varied substantially between species and different  
7  
8 3 chicken tissues. The highest concentrations were found in chicken eggs, followed by  
9  
10 4 chicken liver, chicken skin and chicken muscle; with concentrations in fish and pork  
11  
12 5 samples much lower than those from chickens. Such interspecies differences indicate  
13  
14 6 that the uptake and metabolism of HBCDs is organism-dependent.  
15  
16 7

17  
18  
19  
20  
21 8 The mean  $\Sigma$ HBCDs concentration in soil in this study was 120 ng/g dw, varying from  
22  
23 9 0.030 to 580 ng/g dw, comparable to HBCD concentrations in surface soils from e-  
24  
25 10 waste processing areas in South China, ranging from 0.38-284 ng/g dw.<sup>22</sup> In contrast,  
26  
27 11 concentrations of HBCDs in soil from the vicinity of HBCD production facilities in  
28  
29 12 Sweden, Belgium, Germany and China (111-23,200 ng/g dw) exceed significantly  
30  
31 13 those in this study; while those in soils from urban Guangzhou (1.7-5.6 ng/g dw) and  
32  
33 14 from open waste dumping sites in India, Vietnam, Malaysia, Indonesia, and Cambodia  
34  
35 15 (< nd to 2.4 ng/g dw) were at the low end of the range detected in our study.<sup>22,49</sup>  
36  
37  
38  
39  
40  
41  
42  
43

44 17 The HBCD diastereomer profiles detected in foodstuffs and co-located soils in this  
45  
46 18 study are shown in Fig. 2 along with the profile reported for the HBCD commercial  
47  
48 19 formulation.<sup>50</sup> Fig. 2 shows  $\gamma$ -HBCD to be the dominant isomer in soil samples in this  
49  
50 20 study, similar to profiles observed in commercial technical products and related  
51  
52 21 abiotic environmental matrices such as sediment, soil and sewage sludge.<sup>42</sup> However,  
53  
54 22 in all food samples (whether sourced from e-waste impacted or control locations),  $\alpha$ -  
55  
56  
57  
58  
59  
60

1 HBCD predominated, in line with previous data for biota.<sup>42,51</sup> Furthermore,  $\alpha$ -HBCD  
2 was relatively more abundant in chicken egg, muscle, and skin than in liver,  
3 indicating tissue-specific variation in the relative abundance of different diastereomers,  
4 whereby  $\gamma$ -HBCD is more prevalent in liver samples than the other tissues studied. As  
5 highlighted previously,<sup>21</sup> we believe the diastereomer pattern in avian liver samples  
6 reflects more closely the pattern present in its diet and soil, as liver is the first organ  
7 exposed following the gastrointestinal tract. In contrast, other avian tissues display a  
8 pattern more influenced by metabolism post-exposure.

9  
10 Fig. 3 shows the enantiomer fraction (EF) values for  $\alpha$ -,  $\beta$ -, and  $\gamma$ -HBCD in chicken  
11 and co-located soil samples. Average EF values in our soil samples were 0.46, 0.53,  
12 and 0.54 for  $\alpha$ -,  $\beta$ -, and  $\gamma$ -HBCD, respectively, compared with those in commercial  
13 HBCD (0.514, 0.510, and 0.503).<sup>22</sup> This slight deviation from racemic suggests some  
14 edaphic enantioselective degradation of HBCDs, consistent with an earlier study that  
15 reported enantioselective biodegradation of (+)- $\alpha$ -HBCD in soils.<sup>52</sup> In contrast, in soil  
16 samples from e-waste recycling areas and industrial areas in South China, negligible  
17 enantioselective degradation was implied by mean EF values ranging from 0.503 to  
18 0.507, 0.494 to 0.506, and 0.502 to 0.511 for  $\alpha$ -,  $\beta$ -, and  $\gamma$ -HBCD, respectively.<sup>22</sup>

19  
20 With respect to biotic matrices, in this study, (+)- $\alpha$ -HBCD was clearly enriched in  
21 chicken skin and egg while (-)- $\alpha$ -HBCD dominated in chicken liver. As proposed  
22 above to explain the different diastereomer profile detected in chicken liver; the

1 different enantiomer profile observed in the liver may reflect the profile to which the  
2 bird is exposed; while that in skin and egg may reflect *in vivo* enantioselective  
3 processing post-exposure. To the best of our knowledge, no studies have been  
4 reported to investigate the enantioselectivity of  $\alpha$ -HBCD enantiomers in all three of  
5 these chicken tissue types examined here. However, in an earlier study by our group  
6 of chicken liver, eggs and muscle tissue from e-waste impacted locations in Taizhou,  
7 China<sup>21</sup>; while (-)- $\alpha$ -HBCD was enriched in all three tissue types, the enrichment was  
8 markedly greater in liver than in egg or muscle tissue – an observation not in  
9 consistent with the present study.

10  
11 Other studies have reported EFs of HBCDs in birds. He et al.<sup>52</sup> reported that spotted  
12 dove and Chinese francolin displayed EF values enriched in the (-)- $\alpha$ -enantiomer,  
13 while Chinese pond heron and its main prey (fish) displayed relatively more (+)- $\alpha$ -  
14 HBCD in an e-waste region in South China. Similarly, Janak et al.<sup>53</sup> found peregrine  
15 falcon eggs and common tern eggs were enriched in (-)- $\alpha$ -HBCD, while white-tailed  
16 sea eagle eggs were depleted in the same enantiomer. Chicken muscle, egg, and liver  
17 in our study displayed relative enrichment of (+)- $\gamma$ -HBCD, consistent with previous  
18 reports for chicken eggs and muscle from e-waste recycling sites in eastern China,<sup>21</sup>  
19 bird samples from an e-waste area in South China<sup>52</sup> and predatory birds' eggs from  
20 Sweden and the Netherlands.<sup>55</sup> In summary, our findings confirm the complex species  
21 and tissue-specific variations that exist in the enantioselective behaviour of HBCDs in  
22 birds.

1  
2  
3  
4 15  
6 2 **3.3 Relationships between concentrations of HFRs in chicken and co-located**  
7  
8  
9 3 **soil samples**10  
11 4 Significant linear positive correlations were found between concentrations of *syn*-DP  
12  
13 5 in soils and those in co-located chicken muscle, liver, and eggs ( $p < 0.05$ ,  $r^2 = 0.921$ ,  
14  
15 6  $0.925$ ,  $0.928$ , respectively for muscle, liver, and egg). Similar correlations were found  
16  
17 7 for *anti*-DP ( $p < 0.05$ ,  $r^2 = 0.876$ ,  $0.879$ ,  $0.885$ , respectively for muscle, liver, and egg).18  
19 8 Combined, these observations suggest that at our sampling sites, soil is an important  
20  
21 9 source of DPs in chickens, consistent with the findings of Zheng et al.<sup>37</sup> This is  
22  
23 10 further substantiated by comparison of values of the fraction of *anti*-DP ( $f_{anti}$  – the  
24  
25 11 fractional contribution of *anti*-DP to  $\Sigma$ DP (sum of *anti*- and *syn*-DP) in soil and co-  
26  
27 12 located chicken tissue samples. Observed  $f_{anti}$  values ranged from 0.67-0.81 in soil,  
28  
29 13 compared to 0.67-0.82 in chicken tissues and 0.65–0.80 in technical DP products.<sup>25,57</sup>30  
31 14 A slight caveat to this conclusion is the fact that significant correlation was not  
32  
33 15 observed between concentrations of either DP isomer in chicken skin samples and co-  
34  
35 16 located soil samples. With respect to other HFRs, the potential importance of soil as a  
36  
37 17 source of BTBPE contamination in chickens at our sites is indicated by the  
38  
39 18 observation of significant correlations between its concentrations in soil and in co-  
40  
41 19 located samples of chicken liver ( $r^2 = 0.985$ ,  $p < 0.01$ ) and muscle ( $r^2 = 0.909$ ,  $p < 0.05$ ).42  
43 20 Conversely, no such correlations were detected between BTBPE concentrations in soil  
44  
45 21 and those in chicken skin and eggs. Similarly, no significant correlations were found  
46  
47 22 between concentrations of any HBCD isomer in any chicken tissue and soil, which  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4 suggests substantial metabolism and/or that food and air could be more important  
5  
6 influences on the HBCD concentrations in chickens.  
7  
8

9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### 3.4 Estimated daily intake of emerging HFRs and HBCDs via consumption of foodstuffs included in this study

Very few studies have estimated human dietary exposure to emerging HFRs. Estimated dietary exposure of the sum of our target emerging HFRs for adults and children in this study were 170 and 420 ng/kg bw/day, respectively. The health effects of this exposure cannot be assessed due to the current lack of health based limit values for emerging HFRs exposure by other researchers. We have, however, compared our estimates with those reported previously. DPs account for >90% of estimated exposure for both adults and children (130 and 350 ng/kg bw/day, respectively) followed by BTBPE (31 and 61 ng/kg bw/day, respectively), while EH-TBB predominated in one study of dietary exposure of the population living in the vicinity of e-waste impacted sites in eastern China, in which DPs were not investigated.<sup>21</sup> Another study calculated average estimated daily intakes of emerging BFRs via eggs from one South China recycling area to range from 970 to 4,530 ng/day,<sup>25</sup> which is higher than our estimate of exposure via egg ingestion of 350 ng/day. Furthermore, the same study concurred with our finding that DP was the dominant contributor to emerging HFRs exposure via egg ingestion.



1  
2  
3  
4 1 Daily dietary exposure to HBCDs of individuals living in an e-waste impacted area in  
5  
6 2 this study was estimated at 480 and 1500 ng/kg bw/day for adults and children,  
7  
8  
9 3 respectively. This exceeds estimated dietary exposure to HBCDs in e-waste impacted  
10  
11 4 locations in China (10.4 and 36.1 ng/kg bw/day for adults and children),<sup>21</sup> and is  
12  
13 5 substantially in excess of estimated fish-related dietary exposure in the Netherlands  
14  
15 6 and Sweden (0.12 and 0.14 ng/kg bw/day, respectively)<sup>46,58</sup> as well as estimated  
16  
17 7 dietary exposure of non-e-waste impacted populations in Spain, Belgium and China  
18  
19 8 (2.58, 0.99, 0.432, 39.28 ng/kg bw/day, respectively).<sup>43,59,60</sup> However, as stated above,  
20  
21 9 the main food supplies for the community in Bui Dau are from neighbouring  
22  
23 10 communities. As a consequence, our assumption that Bui Dau inhabitants source all  
24  
25 11 their fish, chicken meat, liver, and eggs from e-waste impacted sites and thus our  
26  
27 12 estimates of the daily intake of emerging HFRs and HBCDs via consumption of such  
28  
29 13 foodstuffs represents a worst case – albeit not wholly unrealistic - scenario.  
30  
31  
32  
33  
34  
35  
36  
37  
38

39 15 For most of our target emerging HFRs and HBCDs, the main contributors to dietary  
40  
41 16 exposure of both adults and children in our study were chicken liver and chicken eggs  
42  
43 17 while fish was the predominant contributor to dietary exposure to BTBPE in our study.  
44  
45  
46 18 In previous studies it has been reported that fish, seafood, meat and meat products are  
47  
48 19 the principal contributors to HBCDs dietary exposure.<sup>43,46,59,60</sup> We are aware of only  
49  
50 20 one previous study of dietary exposure to emerging HFRs in e-waste impacted areas<sup>21</sup>.  
51  
52 21 In that study, consumption of pork was the principal contributor to dietary exposure of  
53  
54 22 both adults and children to EH-TBB, BEH-TEBP, and BTBPE.<sup>21</sup>  
55  
56  
57  
58  
59  
60

1

2 **4. Conclusions**

3 A number of emerging HFRs were found in chicken, fish, and pork samples from an  
4 e-waste processing site in Vietnam, as well as high levels of HBCDs. The most  
5 abundant emerging HFR detected was DP and some enantioselective enrichment of  
6 both  $\alpha$ - and  $\gamma$ -HBCD was observed in chicken samples. For most of our target  
7 compounds, the main contributors to dietary exposure of both adults and children  
8 were chicken liver and chicken eggs. Estimated daily dietary intakes of HBCDs were  
9 higher than those reported from other countries. This study provides evidence that  
10 emerging HFRs are already entering the waste stream leading to environmental  
11 contamination when such waste is treated in an unregulated fashion. The elevated  
12 concentrations of DP are of particular note. We hypothesise that over time,  
13 environmental contamination with emerging HFRs will rise as increasing numbers of  
14 products containing these chemicals reach the end of their useful life.

15

16 **5. Acknowledgements**

17 This research is supported by a Li Siguang scholarship to Fang Tao funded by the  
18 China Scholarship Council and the University of Birmingham, as well as the  
19 European Union Seventh Framework Programme FP7/2007-2013 under the  
20 INTERFLAME project # 295138. We also gratefully acknowledge the  
21 following: local people from Vietnam for help with translation and sample collection;  
22 Natsuyo Noda, Tomoko Oguri, and Akinori Hashimoto for assistance with sample

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 collection and preparation.

**Table 1. Average, median and range of concentrations of Emerging HFRs and HBCD (ng/g lw) in food and co-located soil and sediment samples (ng/g dw) from an e-waste processing (Bui Dau, Vietnam) and control sites.**


		Chicken muscle (n=5)	Chicken liver (n=5)	Chicken skin (n=5)	Chicken egg (n=15)	Soil (n=5)	River fish (n=5)	Pork (n=2)	Sediment <sup>c</sup> (n=8)	Control chicken muscle- 1 (n=1) <sup>a</sup>	Control chicken egg-1 (n=3) <sup>a</sup>	Control chicken muscle- 2 (n=1) <sup>b</sup>	Control chicken egg-2 (n=1) <sup>b</sup>	Control chicken liver (n=1) <sup>b</sup>	Control chicken skin (n=1) <sup>b</sup>	Control fish (n=1) <sup>a</sup>	Control pork (n=1) <sup>a</sup>
<b>Lipid%</b>		6.0	15	43	42	--	4.0	18	--	21	42	2.0	44	12	46	17	30
<b>PBBz</b>	average	<1.5	2.0	1.3	0.92	<0.15	<0.36	<0.36	--	<1.5	<0.31	<1.5	<0.31	<0.36	<0.31	<0.36	<0.36
	median	<1.5	<0.36	0.42	0.96	<0.15	<0.36	<0.36	--	<1.5	<0.31	<1.5	<0.31	<0.36	<0.31	<0.36	<0.36
	range	--	<0.50- 3.7	<0.50- 2.0	0.19-1.9	--	--	--	--	--	--	--	--	--	--	--	--
<b>HBB</b>	average	6.4	6.8	3.0	1.5	6.9	<0.36	<0.36	0.033	<1.5	<0.30	<1.5	<0.30	<0.36	<0.30	<0.36	<0.36
	median	<1.5	<0.36	0.49	1.4	1.3	<0.36	<0.36	--	<1.5	<0.30	<1.5	<0.30	<0.36	<0.30	<0.36	<0.36
	range	<1.5-10	<0.36- 11	<0.30- 5.8	<1.3-2.0	0.15-21	--	--	ND <sup>d</sup> -0.23	--	--	--	--	--	--	--	--
<b>BEH- TEBP</b>	average	<6.5	<1.5	<1.3	2.0	0.13	<1.5	<1.5	<0.20	<6.5	<1.3	<6.5	<1.3	<1.5	<1.3	<1.5	<1.5
	median	<6.5	<1.5	<1.3	<1.3	<0.050	<1.5	<1.5	--	<6.5	<1.3	<6.5	<1.3	<1.5	<1.3	<1.5	<1.5
	range	--	--	--	<1.3-3.0	<0.050- 0.40	--	--	ND-1.6	--	--	--	--	--	--	--	--
<b>BTBPE</b>	average	60	54	70	67	10	40	<0.70	0.89	<3.1	<0.62	<3.1	<0.62	<0.70	<0.62	<0.70	<0.70
	median	46	12	25	18	5.2	55	<0.70	--	<3.1	<0.62	<3.1	<0.62	<0.70	<0.62	<0.70	<0.70
	range	<3.1-100	<0.70- 130	<0.62- 100	<2.8- 160	0.19-34	<1.0- 57	--	ND-5.7	--	--	--	--	--	--	--	--
<b>DBDPE</b>	average	6.9	<3.0	<2.8	280	21	<3.0	2.5	3.1	<7.5	<2.5	<7.5	<2.5	<3.0	<2.5	<3.0	<3.0
	median	<7.5	<3.0	<2.8	<2.5	12	<3.0	--	--	<7.5	<2.5	<7.5	<2.5	<3.0	<2.5	<3.0	<3.0

	range	<7.5-9.9	--	--	<2.5-620	0.42-64	--	--	ND-20	--	--	--	--	--	--	--	--
<i>syn</i> -DP	average	310	1600	170	140	3.8	<0.36	<0.36	0.17	<1.6	<0.32	<1.6	<0.32	<0.36	<0.32	<0.36	<0.36
	median	76	170	90	5.0	0.60	<0.36	<0.36	--	<1.6	<0.32	<1.6	<0.32	<0.36	<0.32	<0.36	<0.36
	range	<1.6-1400	<0.36-8000	0.13-790	3.4-560	0.20-13	--	--	ND-1.8	--	--	--	--	--	--	--	--
<i>anti</i> -DP	average	1200	3500	350	450	10	<0.36	<0.36	0.45	<1.6	<0.32	<1.6	<0.32	<0.36	<0.32	<0.36	<0.36
	median	260	210	160	8.0	2.3	<0.36	<0.36	--	--	--	--	--	--	--	--	--
	range	<1.6-1600	7.5-17000	0.32-3000	5.1-1800	0.83-31	--	--	ND-4.8	--	--	--	--	--	--	--	--
$\Sigma$ DP	average	1500	5100	520	590	14	<0.36	<0.36	0.59	<1.6	<0.32	<1.6	<0.32	<0.36	<0.32	<0.36	<0.36
	median	340	380	250	13	2.9	<0.36	<0.36	--	--	--	--	--	--	--	--	--
	range	<1.6-3000	7.5-25000	0.44-3800	4.0-2300	1.3-44	--	--	ND-6.7	--	--	--	--	--	--	--	--
$\alpha$ -HBCD	average	34	1700	600	2800	6.7	2.5	0.90	--	0.10	0.10	17	1.6	<0.020	1.0	<0.020	<0.020
	median	20	1000	640	2500	5.0	2.3	--	--	--	--	--	--	--	--	--	--
	range	1.2-55	180-2500	20-850	330-3500	1.0-8.9	0.10-3.3	--	--	--	--	--	--	--	--	--	--
$\beta$ -HBCD	average	0.15	28	0.06	79	4.8	0.41	<0.020	--	<0.010	<0.010	<0.010	<0.010	<0.010	<0.010	<0.010	<0.010
	median	0.25	20	0.10	70	3.8	0.51	--	--	--	--	--	--	--	--	--	--
	range	0.10-2.0	2.5-35	0.02-0.15	15-80	0.50-8.0	0.040-0.65	--	--	--	--	--	--	--	--	--	--
$\gamma$ -HBCD	average	5.3	1500	330	700	110	0.50	0.20	--	<0.020	<0.020	0.57	0.95	<0.020	<0.020	<0.020	<0.020
	median	8.9	890	450	910	100	0.60	--	--	--	--	--	--	--	--	--	--
	range	0.10-15	160-3000	3.8-580	200-2300	56-710	0.050-0.12	--	--	--	--	--	--	--	--	--	--
$\Sigma$ HBCD	average	39	3200	930	3600	120	3.4	1.1	--	0.10	0.10	18	2.6	<0.050	1.0	<0.050	<0.050
	median	29	1900	1000	3500	110	3.4	--	--	--	--	--	--	--	--	--	--
	range	2.0-80	330-5500	25-1400	540-5800	0.030-580	0.20-4.1	--	--	--	--	--	--	--	--	--	--


<sup>a</sup> Control samples collected in Vietnam; <sup>b</sup> Control samples collected in Japan; <sup>c</sup> from Someya et al., submitted; <sup>d</sup> ND= not detected.

**Table 2. Estimated dietary exposure (ng/kg bw/day) to emerging HFRs and HBCD of adults and children living in e-waste impacted areas of Bui Dau, Vietnam. ne = not estimated**


Foodstuff	Exposed group/HFR	PBBz	HBB	BTBPE	BEH-TEBP	DBDPE	<i>syn</i> -DP	<i>anti</i> -DP	$\alpha$ -HBCD	$\beta$ -HBCD	$\gamma$ -HBCD	$\Sigma$ HBCD
Chicken Muscle	Adult	ne	0.17	1.2	ne	0.16	11	32	14	0.06	2.2	16
	Children	ne	0.28	2.1	ne	0.26	19	54	23	0.10	3.6	27
Chicken Egg	Adult	0.010	0.010	0.28	0.010	0.20	1.7	5.0	73	2.1	18	93
	Children	0.020	0.030	0.97	0.040	0.71	5.8	16	250	7.1	63	320
Chicken Liver	Adult	0.030	0.10	0.75	ne	ne	29	50	180	3.0	160	350
	Children	0.090	0.31	2.4	ne	ne	93	157	580	9.6	510	1100
Chicken Skin	Adult	0.010	0.010	0.35	ne	ne	0.76	2.9	11	ne	6.2	17
	Children	0.010	0.020	0.58	ne	ne	1.3	4.8	18	ne	10	29
Fish	Adult	ne	ne	28	ne	ne	ne	ne	1.8	0.29	0.35	2.4
	Children	ne	ne	55	ne	ne	ne	ne	3.4	0.56	0.68	4.7
Pork	Adult	ne	ne	ne	ne	4.6	ne	ne	1.7	ne	0.37	2.0
	Children	ne	ne	ne	ne	1.7	ne	ne	0.61	ne	0.14	0.75
Total	Adult	0.040	0.29	31	0.010	5.0	43	90	290	5.0	190	480
	Children	0.12	0.64	61	0.040	2.7	120	230	880	17	590	1500




**Fig. 1. Sampling sites in Bui Dau, Cam Xa, Hung Yen province, Vietnam.**



**Fig. 2. Diastereomer profiles in food and co-located surface soil samples from e-waste processing areas in Vietnam this study and commercial HBCD mixtures (Covaci et al., 2006)**




**Fig. 3: Enantiomer fractions (EF) of  $\alpha$ -HBCD,  $\gamma$ -HBCD and  $\beta$ -HBCD in chicken and co-located samples (line at EF=0.50 indicates racemic value)**



**Fig. 4: Contributions of different food groups to total dietary exposures to emerging HFRs and HBCDs of children in Bui Dau, Vietnam**



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



**Fig. 5: Contributions of different food groups to total dietary exposures to emerging HFRs and HBCDs of adults in Bui Dau, Vietnam**

## References:

- 1 N. Ali, S. Harrad, E. Goosey, H. Neels and A. Covaci, *Chemosphere*, 2011, **83**, 1360–1365.
- 2 T. Reistad, E. Mariussen and F. Fonnum, *Toxicol. Sci.*, 2005, **83**, 89–100.
- 3 A. Covaci, S. Harrad, M. A. E. Abdallah, N. Ali, R. J. Law, D. Herzke and C. A. de Wit, *Environ. Int.*, 2011, **37**, 532–556.
- 4 N. Ali, A. C. Dirtu, N. Van den Eede, E. Goosey, S. Harrad, H. Neels, A. 't Mannetje, J. Coakley, J. Douwes and A. Covaci, *Chemosphere*, 2012, **88**, 1276–1282.
- 5 M. Karlsson, A. Julander, B. van Bavel and L. Hardell, *Environ. Int.*, 2007, **33**, 62–69.
- 6 H. M. Stapleton, J. G. Allen, S. M. Kelly, A. Konstantinov, S. Klosterhaus, D. Watkins, M. D. McClean and T. F. Webster, *Environ. Sci. Technol.*, 2008, **42**, 6910–6916.
- 7 M. Venier and R. A. Hites, *Environ. Sci. Technol.*, 2008, **42**, 4745–4751.
- 8 J. Cristale, A. García Vázquez, C. Barata and S. Lacorte, *Environ. Int.*, 2013, **59**, 232–243.
- 9 L. Zeng, R. Yang, Q. Zhang, H. Zhang, K. Xiao, H. Zhang, Y. Wang, P. K. S. Lam and G. Jiang, *Environ. Sci. Technol.*, 2014, **48**, 12586–12594.
- 10 Q. Zheng, L. Nizzetto, J. Li, M. D. Mulder, O. Sanka, G. Lammel, H. Bing, X. Liu, Y. Jiang, C. Luo and G. Zhang, *Environ. Sci. Technol.*, 2015, **49**, 2904–2911.
- 11 C. de Wit, A. Kierkegaard, N. Ricklund and U. Sellström, eds. E. Eljarrat and D. Barceló, Springer Berlin / Heidelberg, 2011, vol. 16, pp. 241–286.

- 1  
2  
3 12 A. Salamova, M. H. Hermanson and R. A. Hites, *Environ. Sci. Technol.*, 2014,  
4 48, 6133–40.  
5  
6  
7  
8 13 G. T. Tomy, V. P. Palace, K. Pleskach, N. Ismail, T. Oswald, R. Danell, K.  
9 Wautier and B. Evans, *Environ. Sci. Technol.*, 2007, 41, 4913–4918.  
10  
11  
12 14 B. Wu, S. Liu, X. Guo, Y. Zhang, X. Zhang, M. Li and S. Cheng, *Environ. Sci.*  
13 *Technol.*, 2012, 46, 10758–10764.  
14  
15  
16  
17 15 E. Barón, C. Hauler, C. Gallistl, J. Giménez, P. Gauffier, J. J. Castillo, C.  
18 Fernández-Maldonado, R. de Stephanis, W. Vetter, E. Eljarrat and D. Barceló,  
19 *Environ. Sci. Technol.*, 2015.  
20  
21  
22  
23 16 T. Shi, S. J. Chen, X. J. Luo, X. L. Zhang, C. M. Tang, Y. Luo, Y. J. Ma, J. P.  
24 Wu, X. Z. Peng and B. X. Mai, *Chemosphere*, 2009, 74, 910–916.  
25  
26  
27  
28 17 Y. Sun, X. Luo, L. Mo, Q. Zhang, J. Wu, S. Chen, F. Zou and B. Mai, *Environ.*  
29 *Pollut.*, 2012, 162, 381–8.  
30  
31  
32  
33 18 D. Chen, X. Bi, J. Zhao, L. Chen, J. Tan, B. Mai, G. Sheng, J. Fu and M. Wong,  
34 *Environ. Pollut.*, 2009, 157, 1051–7.  
35  
36  
37  
38 19 I. Labunska, S. Harrad, D. Santillo, P. Johnston and L. Yun, *Environ. Sci.*  
39 *Technol.*, 2013, 47, 9258–9266.  
40  
41  
42  
43 20 H. Liu, Q. Zhou, Y. Wang, Q. Zhang, Z. Cai and G. Jiang, *Environ. Int.*, 2008,  
44 34, 67–72.  
45  
46  
47  
48 21 I. Labunska, M. A.-E. Abdallah, I. Eulaers, A. Covaci, F. Tao, M. Wang, D.  
49 Santillo, P. Johnston and S. Harrad, *Environ. Int.*, 2015, 74, 209–220.  
50  
51  
52  
53 22 S. Gao, J. Wang, Z. Yu, Q. Guo, G. Sheng and J. Fu, *Environ. Sci. Technol.*,  
54 2011, 45, 2093–2099.  
55  
56  
57  
58 23 N. M. Tue, S. Takahashi, G. Suzuki, T. Isobe, P. H. Viet, Y. Kobara, N. Seike, G.  
59 Zhang, A. Sudaryanto and S. Tanabe, *Environ. Int.*, 2013, 51, 160–167.  
60

- 1  
2  
3 24 N. M. Tue, A. Sudaryanto, T. B. Minh, T. Isobe, S. Takahashi, P. H. Viet and S.  
4 Tanabe, *Sci. Total Environ.*, 2010, **408**, 2155–2162.  
5  
6  
7  
8 25 X.-B. Zheng, J.-P. Wu, X.-J. Luo, Y.-H. Zeng, Y.-Z. She and B.-X. Mai,  
9 *Environ. Int.*, 2012, **45**, 122–128.  
10  
11  
12 26 H. Matsukami, N. M. Tue, G. Suzuki, M. Someya, L. H. Tuyen, P. H. Viet, S.  
13 Takahashi, S. Tanabe and H. Takigami, *Sci. Total Environ.*, 2015, **514**, 492–9.  
14  
15  
16  
17 27 M. A. Abdallah and S. Harrad, *Environ. Int.*, 2011, **37**, 443–448.  
18  
19  
20 28 M. Someya, G. Suzuki, A. C. Ionas, N. M. Tue, F. Xu, H. Matsukami, A.  
21 Covaci, L. H. Tuyen, P. H. Viet, S. Takahashi, S. Tanabe and H. Takigami,  
22 *Emerg. Contam.*, 2015.  
23  
24  
25  
26 29 I. Labunska, S. Harrad, M. Wang, D. Santillo and P. Johnston, *Environ. Sci.*  
27 *Technol.*, 2014, **48**, 5555–5564.  
28  
29  
30 30 R. V Thomann, *Environ. Sci. Technol.*, 1989, **23**, 699–707.  
31  
32  
33  
34 31 Q. Xian, S. Siddique, T. Li, Y. Feng, L. Takser and J. Zhu, *Environ. Int.*, 2011,  
35 **37**, 1273–1284.  
36  
37  
38 32 G. T. Tomy, C. R. Thomas, T. M. Zidane, K. E. Murison, K. Pleskach, J. Hare,  
39 G. Arsenaault, C. H. Marvin and E. Sverko, *Environ. Sci. Technol.*, 2008, **42**,  
40 5562–5567.  
41  
42  
43  
44 33 J.-P. Wu, Y. Zhang, X.-J. Luo, J. Wang, S.-J. Chen, Y.-T. Guan and B.-X. Mai,  
45 *Environ. Sci. Technol.*, 2010, **44**, 606–611.  
46  
47  
48  
49 34 E. Sverko, G. T. Tomy, E. J. Reiner, Y.-F. Li, B. E. McCarry, J. A. Arnot, R. J.  
50 Law and R. A. Hites, *Environ. Sci. Technol.*, 2011, **45**, 5088–98.  
51  
52  
53 35 J.-H. Kang, J.-C. Kim, G.-Z. Jin, H. Park, S.-Y. Baek and Y.-S. Chang,  
54 *Chemosphere*, 2010, **79**, 850–4.  
55  
56  
57  
58 36 R.-X. Sun, X.-J. Luo, X.-X. Tan, B. Tang, Z.-R. Li and B.-X. Mai, *Mar. Pollut.*  
59  
60

- 1  
2  
3 *Bull.*, 2015, **93**, 61–7.  
4  
5  
6 37 X.-B. Zheng, X.-J. Luo, Y.-H. Zeng, J.-P. Wu and B.-X. Mai, *Chemosphere*,  
7 2014, **114**, 241–246.  
8  
9  
10  
11 38 A. Fernandes, F. Smith, R. Petch, S. Panton, M. Carr, D. Mortimer, C. Tlustos  
12 and M. Rose, *Proc. BFR 2010, Kyoto, Japan*, 2010, 90028.  
13  
14  
15 39 Z. Yu, S. Lu, S. Gao, J. Wang, H. Li, X. Zeng, G. Sheng and J. Fu, *Environ.*  
16 *Pollut.*, 2010, **158**, 2920–2925.  
17  
18  
19  
20 40 W.-L. Ma, L.-Y. Liu, H. Qi, D.-Z. Sun, J.-M. Shen, D.-G. Wang and Y.-F. Li,  
21 *Environ. Int.*, 2011, **37**, 66–70.  
22  
23  
24  
25 41 M. Alaei, P. Arias, A. Sjödin and Å. Bergman, *Environ. Int.*, 2003, **29**, 683–689.  
26  
27  
28 42 A. Covaci, A. C. Gerecke, R. J. Law, S. Voorspoels, M. Kohler, N. V Heeb, H.  
29 Leslie, C. R. Allchin and J. de Boer, *Environ. Sci. Technol.*, 2006, **40**, 3679–  
30 3688.  
31  
32  
33  
34 43 Z. X. Shi, Y. N. Wu, J. G. Li, Y. F. Zhao and J. F. Feng, *Environ. Sci. Technol.*,  
35 2009, **43**, 4314–4319.  
36  
37  
38  
39 44 A. Schecter, T. R. Harris, S. Brummitt, N. Shah and O. Paepke, *Epidemiology*,  
40 2008, **19**, S76–S76.  
41  
42  
43 45 A. C. Dirtu and A. Covaci, *Environ. Sci. Technol.*, 2010, **44**, 6297–6304.  
44  
45  
46 46 A. Törnkvist, A. Glynn, M. Aune, P. O. Darnerud and E. H. Ankarberg,  
47 *Chemosphere*, 2011, **83**, 193–199.  
48  
49  
50  
51 47 L. Roosens, M. A. E. Abdallah, S. Harrad, H. Neels and A. Covaci, *Environ.*  
52 *Health Perspect.*, 2009, **117**, 1707–1712.  
53  
54  
55  
56 48 M. Driffield, N. Harmer, E. Bradley, A. R. Fernandes, M. Rose, D. Mortimer  
57 and P. Dicks, *Food Addit. Contam. Part A*, 2008, **25**, 895–903.  
58  
59  
60

- 1  
2  
3  
4  
5 49 H. Li, Q. Zhang, P. Wang, Y. Li, J. Lv, W. Chen, D. Geng, Y. Wang, T. Wang  
6 and G. Jiang, *J. Environ. Monit.*, 2012, **14**, 2591–2597.  
7  
8  
9  
10 50 K. Law, V. P. Palace, T. Halldorson, R. Danell, K. Wautier, B. Evans, M. Alaei,  
11 C. Marvin and G. T. Tomy, *Environ. Toxicol. Chem.*, 2006, **25**, 1757–1761.  
12  
13  
14 51 T. Reistad, F. Fonnum and E. Mariussen, *Arch. Toxicol.*, 2006, **80**, 785–796.  
15  
16  
17 52 M.-J. He, X.-J. Luo, L.-H. Yu, J. Liu, X.-L. Zhang, S.-J. Chen, D. Chen and B.-  
18 X. Mai, *Environ. Sci. Technol.*, 2010, **44**, 5748–54.  
19  
20  
21 53 S. Esslinger, R. Becker, R. Maul and I. Nehls, *Environ. Sci. Technol.*, 2011, **45**,  
22 3938–3944.  
23  
24  
25  
26 54 M. A.-E. Abdallah, C. Uchea, J. K. Chipman and S. Harrad, *Environ. Sci.*  
27 *Technol.*, 2014, **48**, 2732–2740.  
28  
29  
30  
31 55 K. Janák, U. Sellström, A.-K. Johansson, G. Becher, C. A. de Wit, P. Lindberg  
32 and B. Helander, *Chemosphere*, 2008, **73**, S193–S200.  
33  
34  
35  
36 56 Y. Zhang, H. Sun and Y. Ruan, *J. Hazard. Mater.*, 2014, **264**, 8–15.  
37  
38  
39 57 X. Qiu, C. H. Marvin and R. A. Hites, *Environ. Sci. Technol.*, 2007, **41**, 6014–  
40 6019.  
41  
42  
43  
44 58 S. P. J. van Leeuwen and J. de Boer, *Mol. Nutr. Food Res.*, 2008, **52**, 194–203.  
45  
46  
47 59 E. Eljarrat, M. Gorga, M. Gasser, J. Díaz-Ferrero and D. Barceló, *J. Agric.*  
48 *Food Chem.*, 2014, **62**, 2462–2468.  
49  
50  
51 60 S. Goscinny, S. Vandevijvere, M. Maleki, I. Van Overmeire, I. Windal, V.  
52 Hanot, M.-N. Blaude, C. Vleminckx and J. Van Looco, *Chemosphere*, 2011, **84**,  
53 279–288.  
54  
55  
56  
57  
58  
59  
60